

# Bayesian Optimisation for Likelihood Free Inference

Make model parameterisation go brrr

Jacob Cumming

University of Melbourne, Walter and Eliza Hall Institute

April 2024



# Notation

- ▶ Model is considered a (random) function  $f(\boldsymbol{\theta})$  that maps  $\boldsymbol{\theta}$  (a vector of parameters) to a model output, that can be transformed into  $\mathbf{X}$ , that has the same shape as:
- ▶  $\mathbf{X}_{\text{obs}}$ , a vector of outputs given to us usually in the forms of summary statistics (incidence, prevalence, hospitalisations etc).

# Parameter inference would become easy if we had...

- ▶ An explicit form for the likelihood:  $\mathcal{L}(\theta|\mathbf{X}_{\text{obs}}) := \Pr(\mathbf{X}_{\text{obs}}|\theta)$

# Parameter inference would become easy if we had...

- ▶ An explicit form for the likelihood:  $\mathcal{L}(\theta|\mathbf{X}_{\text{obs}}) := \Pr(\mathbf{X}_{\text{obs}}|\theta)$
- ▶ Or even  $\mathcal{L}(\theta|S(\mathbf{X}_{\text{obs}})) := \Pr(S(\mathbf{X}_{\text{obs}})|\theta)$ , where  $S(\mathbf{X}_{\text{obs}})$  is a (vector of) summary statistic(s)

# Parameter inference would become easy if we had...

- ▶ An explicit form for the likelihood:  $\mathcal{L}(\theta|\mathbf{X}_{\text{obs}}) := \Pr(\mathbf{X}_{\text{obs}}|\theta)$
- ▶ Or even  $\mathcal{L}(\theta|S(\mathbf{X}_{\text{obs}})) := \Pr(S(\mathbf{X}_{\text{obs}})|\theta)$ , where  $S(\mathbf{X}_{\text{obs}})$  is a (vector of) summary statistic(s)
- ▶  $\hat{\theta} = \arg \max_{\theta} \mathcal{L}(\theta|S(\mathbf{X}_{\text{obs}}))$

# Parameter inference would become easy if we had...

- ▶ An explicit form for the likelihood:  $\mathcal{L}(\theta|\mathbf{X}_{\text{obs}}) := \Pr(\mathbf{X}_{\text{obs}}|\theta)$
- ▶ Or even  $\mathcal{L}(\theta|S(\mathbf{X}_{\text{obs}})) := \Pr(S(\mathbf{X}_{\text{obs}})|\theta)$ , where  $S(\mathbf{X}_{\text{obs}})$  is a (vector of) summary statistic(s)
- ▶  $\hat{\theta} = \arg \max_{\theta} \mathcal{L}(\theta|S(\mathbf{X}_{\text{obs}}))$
- ▶  $\Pr(\theta|S(\mathbf{X}_{\text{obs}})) \propto \Pr(S(\mathbf{X}_{\text{obs}})|\theta) \Pr(\theta)$

# The Sad Truth

- ▶ As models become more complicated, explicit likelihoods don't exist (think agent based models).

# A Standard Bayesian Solution

- ▶ Approximate Bayesian Computation (ABC)
  1. Sample from prior
  2. Run model
  3. Accept or reject parameters run based on how well  $\mathbf{X}$  'matches'  $\mathbf{X}_{\text{obs}}$ .



# What is 'matches'

- ▶ Discrepancy function  $D : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$

- ▶ Can be a norm such as

$$\|S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})\|_p := (\sum_{i=1}^d |S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})|^p)^{1/p}$$


# What is 'matches'

- ▶ Discrepancy function  $D : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ 
  - ▶ Can be a norm such as
$$\|S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})\|_p := (\sum_{i=1}^d |S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})|^p)^{1/p}$$
  - ▶ Care should be taken to rescale  $S(\mathbf{X}_{\text{obs}})$  and  $S(\mathbf{X})$  appropriately (ie via a covariance matrix).


# What is 'matches'

- ▶ Discrepancy function  $D : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ 
  - ▶ Can be a norm such as
$$\|S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})\|_p := (\sum_{i=1}^d |S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})|^p)^{1/p}$$
  - ▶ Care should be taken to rescale  $S(\mathbf{X}_{\text{obs}})$  and  $S(\mathbf{X})$  appropriately (ie via a covariance matrix).
- ▶  $D(S(\mathbf{X}), S(\mathbf{X}_{\text{obs}}))$ , gives acceptance probability of  $\theta$ .

# Acceptance Probability



# Uniform Acceptance Probability



# Overall Idea of my Research

- ▶ What if we could 'predict' discrepancy values we hadn't seen before?

# Overall Idea of my Research

- ▶ What if we could 'predict' discrepancy values we hadn't seen before?
- ▶ For parameters 'close' to parameters we've already tried it should be easy.

# Gaussian Processes

- ▶ Random functions
- ▶ Common examples - Brownian motion, Ornstein Uhlenbeck process



# Gaussian Processes on $\mathbb{R}^d$

## Definition (Gaussian Process)


A collection of random variables  $\{f(\mathbf{x})\}_{\mathbf{x} \in \mathbb{R}^d}$  is a Gaussian process if all finite dimensional distributions are multivariate normal distributed. That is, there is a function  $m : \mathcal{X} \rightarrow \mathbb{R}$  and kernel  $k : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$  such that for all finite sets  $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n\}$ ,

$$\begin{bmatrix} f(\mathbf{x}_1) \\ f(\mathbf{x}_2) \\ \vdots \\ f(\mathbf{x}_n) \end{bmatrix} \sim \text{MVN} \left( \begin{bmatrix} m(\mathbf{x}_1) \\ m(\mathbf{x}_2) \\ \vdots \\ m(\mathbf{x}_n) \end{bmatrix}, \mathbf{K} \right)$$

where

$$\mathbf{K} = \begin{bmatrix} k(\mathbf{x}_1, \mathbf{x}_1) & k(\mathbf{x}_1, \mathbf{x}_2) & \dots & k(\mathbf{x}_1, \mathbf{x}_n) \\ k(\mathbf{x}_2, \mathbf{x}_1) & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ k(\mathbf{x}_n, \mathbf{x}_1) & \dots & \dots & k(\mathbf{x}_n, \mathbf{x}_n) \end{bmatrix}$$

# Gaussian Process Example Realisations



# Covariance Kernel Motivation

- ▶ Kernel determines the amount of covariance between sets of indices.
- ▶ When the distance between indices is small, covariance needs to be large

# Common Covariance Kernels

- ▶ Matern Kernel


$$k_{\nu}(x, x') = \sigma^2 \frac{2^{1-\nu}}{\Gamma(\nu)} \left( \frac{\sqrt{2\nu} \|x - x'\|}{\ell} \right)^{\nu} K_{\nu} \left( -\frac{\sqrt{2\nu} \|x - x'\|}{\ell} \right)$$

where  $K_{\nu}$  is a modified Bessel function ( $\|\cdot\|$  is the euclidean distance)


- ▶  $\lfloor \nu \rfloor$  times mean square differentiable.
- ▶ As  $\nu \rightarrow \infty$  you get squared exponential covariance function, which results in realisations that are infinitely mean square differentiable:

$$k(x, x') = \sigma^2 \exp\left(-\frac{\|x - x'\|^2}{\ell}\right)$$


# Kernel Choices - Kernel Type




# Kernel Choices - Kernel Type



# Kernel Choices - Kernel Type



# Kernel Choices - Kernel Type







## Kernel Choices - length scale

$$k(x, x') = \sigma^2 \exp\left(-\frac{\|x - x'\|^2}{\ell}\right)$$


# Kernel Choices - length scale



# Kernel Choices - length scale



# Kernel Choices - length scale




# Fitting our GP to data

GPs are 'priors'


# Fitting our GP to data

GPs are 'priors'




# Fitting our GP to data

GPs are 'priors'



# Fitting our GP to data


GPs are 'priors'






# Fitting our GP to data

GPs are 'priors'




# Fitting our GP to data

GPs are 'priors'




# Fitting our GP to data

GPs are 'priors'




# Fitting our GP to data

GPs are 'priors'




GP regression on  $x(x-1)(x+1)$


# GP regression on $x(x - 1)(x + 1)$



# GP regression on $x(x-1)(x+1)$




# GP regression on $x(x-1)(x+1)$






# GP regression on $x(x-1)(x+1)$



# GP regression on $x(x-1)(x+1)$




# What if we have noise?

Add in observation variance, so that


$$\begin{bmatrix} f(\mathbf{x}_1) \\ f(\mathbf{x}_2) \\ \vdots \\ f(\mathbf{x}_n) \end{bmatrix} \sim \text{MVN} \left( \begin{bmatrix} m(\mathbf{x}_1) \\ m(\mathbf{x}_2) \\ \vdots \\ m(\mathbf{x}_n) \end{bmatrix}, \mathbf{K} + \sigma^2 \mathbf{I}_n \right)$$

GP regression on  $x(x-1)(x+1)$


# GP regression on $x(x-1)(x+1)$




# GP regression on $x(x-1)(x+1)$



# GP regression on $x(x - 1)(x + 1)$




# GP regression on $x(x-1)(x+1)$





# GP regression on $x(x-1)(x+1)$



# Overall Idea again

- ▶ What if we could 'predict' discrepancy values we hadn't seen before?

# Overall Idea again

- ▶ What if we could 'predict' discrepancy values we hadn't seen before?
- ▶ For parameters 'close' to parameters we've already tried it should be easy

# Overall Idea again

- ▶ What if we could 'predict' discrepancy values we hadn't seen before?
- ▶ Use Gaussian process to predict discrepancy function

# About Vivax Malaria

- ▶ Has dormant liver stage on top of blood stage infection

# Champagne Model Parameters

- ▶  $\alpha$  : proportion of those infected but cleared of blood stage infections (through treatment)
- ▶  $\beta$  : a further proportion that are also cleared of liver stage parasites, given that they were also cleared of blood stage infection (radical cure)
- ▶  $\lambda$  : the rate of infection
- ▶  $\gamma_L$  : rate of clearance of liver stage disease
- ▶  $f$  : rate of relapse
- ▶  $r$  : rate of blood stage clearance
- ▶  $\delta$  : importation rate (which we assume is 0)

# Champagne ODEs

$$\begin{aligned}\frac{dI_L}{dt} = & (1 - \alpha)(\lambda I_{\text{total}} + \delta)(S_0 + S_L) + (\lambda I_{\text{total}} + \delta)I_0 \\ & + (1 - \alpha)fS_L - \gamma_L I_L - rI_L\end{aligned}$$

$$\frac{dI_0}{dt} = -(\lambda I_{\text{total}} + \delta)I_0 + \gamma_L I_L - rI_0$$

$$\begin{aligned}\frac{dS_L}{dt} = & -(1 - \alpha(1 - \beta))(\lambda I_{\text{total}} + \delta + f)S_L + \alpha(1 - \beta)(\lambda I_{\text{total}} \\ & + \delta)S_0 - \gamma_L S_L + rI_L\end{aligned}$$

$$\begin{aligned}\frac{dS_0}{dt} = & -(1 - \alpha\beta)(\lambda I_{\text{total}} + \delta)S_0 + (\lambda I_{\text{total}} + \delta)\alpha\beta S_L + \alpha\beta fS_L \\ & + \gamma_L S_L + rI_0\end{aligned}$$

# Champagne Model Diagram




Figure: *P. vivax* model described by Champagne et al. 2022



# Model Calibration Data

- ▶ Doob-Gillespie algorithm with paper parameters reported in the original paper for 'observed data', 10 initial infections, 1000 people.


# Model Calibration Data

- ▶ Doob-Gillespie algorithm with paper parameters reported in the original paper for 'observed data', 10 initial infections, 1000 people.
- ▶  $S(\mathbf{X}_{\text{obs}}) := \{w_{\text{obs}}, p_{\text{obs}}, m_{\text{obs}}\}$ 
  - ▶  $w_{\text{obs}}$  : weekly incidence around (stochastic) equilibrium
  - ▶  $p_{\text{obs}}$  : prevalence around (stochastic) equilibrium
  - ▶  $m_{\text{obs}}$  : incidence in the first month of the epidemic

# Model Calibration Data

- ▶ Doob-Gillespie algorithm with paper parameters reported in the original paper for 'observed data', 10 initial infections, 1000 people.
- ▶  $S(\mathbf{X}_{\text{obs}}) := \{w_{\text{obs}}, p_{\text{obs}}, m_{\text{obs}}\}$ 
  - ▶  $w_{\text{obs}}$  : weekly incidence around (stochastic) equilibrium
  - ▶  $p_{\text{obs}}$  : prevalence around (stochastic) equilibrium
  - ▶  $m_{\text{obs}}$  : incidence in the first month of the epidemic
- ▶  $D(S(\mathbf{X}), S(\mathbf{X}_{\text{obs}})) := \left| \frac{w_{\text{obs}} - w}{w_{\text{obs}}} \right| + \left| \frac{p_{\text{obs}} - p}{p_{\text{obs}}} \right| + \left| \frac{m_{\text{obs}} - m}{m_{\text{obs}}} \right|$ 
  - ▶  $L_1$  norm on the relative differences

# Example Simulation



# What's the Bayesian part?

- ▶ Choosing the next point to sample

# What's the Bayesian part?

- ▶ Choosing the next point to sample
- ▶  $\arg \min_{\theta} A(\theta)$  where  $A$  is an acquisition function.

# What's the Bayesian part?

- ▶ Choosing the next point to sample
- ▶  $\arg \min_{\theta} A(\theta)$  where  $A$  is an acquisition function.
- ▶ BOLFI paper uses

$$\mu(\theta) - \eta_t \sqrt{v(\theta)}$$

- ▶  $\eta_t := \sqrt{c + 2 \ln(t^{d/2+2})}$ , and  $c$  can be chosen
- ▶  $\mu(\theta)$  and  $v(\theta)$  are the posterior mean and variance

# What's the Bayesian part?

- ▶ Choosing the next point to sample
- ▶  $\arg \min_{\theta} A(\theta)$  where  $A$  is an acquisition function.
- ▶ BOLFI paper uses

$$\mu(\theta) - \eta_t \sqrt{v(\theta)}$$

- ▶  $\eta_t := \sqrt{c + 2 \ln(t^{d/2+2})}$ , and  $c$  can be chosen
  - ▶  $\mu(\theta)$  and  $v(\theta)$  are the posterior mean and variance
- ▶ Could use expected information


$$(\mu_{\min} - \mu(\theta))\Phi\left(\frac{\mu_{\min} - \mu(\theta)}{\sqrt{v(\theta)}}\right) + \sqrt{v(\theta)}\phi\left(\frac{\mu_{\min} - \mu(\theta)}{\sqrt{v(\theta)}}\right)$$

- ▶  $\mu_{\min} := \min_{\theta} \mu(\theta)$
  - ▶  $\Phi, \phi$  CDF and PDF of standard normal




# Synthetic Likelihood

- ▶  $L(\theta|\mathbf{X}_{\text{obs}}) \approx P(D_{\mathcal{GP}}(\theta) = 0)$  where  $D_{\mathcal{GP}}$  is the discrepancy modelled the Gaussian process
- ▶ This is equivalent to using the half normal acceptance criteria




# How did it go?


# How did it go?




# How did it go?



# How did it go?



# How did it go?



# Big Problem (Big Solutions?)

- ▶ Observation variance is considered constant across the GP
  - ▶ Particularly a problem at the threshold

# Big Problem (Big Solutions?)

- ▶ Observation variance is considered constant across the GP
  - ▶ Particularly a problem at the threshold
  - ▶ Fix by modelling observation variance as another GP



# Thanks to

- ▶ Eamon Conway and the Mueller lab at WEHI
- ▶ Jennifer Flegg at Unimelb
- ▶ Damon for explaining disease modelling so I don't have to