

Author's Proof

Carefully read the entire proof and mark all corrections in the appropriate place, using the Adobe Reader commenting tools ([Adobe Help](#)). Do not forget to reply to the queries.

We do not accept corrections in the form of edited manuscripts.

In order to ensure the timely publication of your article, please submit the corrections within 48 hours.

If you have any questions, please contact science.production.office@frontiersin.org.

Author Queries Form

Q1	The citation and surnames of all of the authors have been highlighted. Please check all of the names carefully and indicate if any are incorrect. Please note that this may affect the indexing of your article in repositories such as PubMed.	All are correct. Request: can we indicate joint first authorship for Mu and Chaudhuri on the title page - perhaps a footnote indicating authors contributed equally?
Q2	Confirm that the email address in your correspondence section is accurate.	Correct
Q3	Please ask the following authors to register with Frontiers (at https://www.frontiersin.org/Registration/Register.aspx) if they would like their names on the article abstract page and PDF to be linked to a Frontiers profile. Please ensure to provide us with the profile link(s) when submitting the proof corrections. Non-registered authors will have the default profile image displayed. "Kallol R. Chaudhuri."	Dr. Chaudhuri has not had time to register, so I think we are fine with publication for now.
Q4	If you decide to use previously published, copyrighted figures in your article, please keep in mind that it is your responsibility, as the author, to obtain the appropriate permissions and licenses and to follow any citation instructions requested by third-party rights holders. If obtaining the reproduction rights involves the payment of a fee, these charges are to be paid by the authors.	Acknowledged. All figures in this manuscript are original.
Q5	Ensure that all the figures, tables and captions are correct.	See below
Q6	Verify that all the equations and special characters are displayed correctly.	Typo: missing period after (ARI = 0.32) (line 452). Otherwise, yes
Q7	Ensure, if it applies to your study, the ethics statement is included in the article.	Yes, under "Standard Protocol Approvals, Registrations, and Patient Consent"
Q8	Ensure to add all grant numbers and funding information, as after publication this is no longer possible.	Yes.
Q9	Please ensure that any supplementary material is correctly published at this link: " http://journal.frontiersin.org/article/10.3389/fnagi.2017.00301/full#supplementary-material ." Please provide new files if you have any corrections.	Yes.

Q10	We have included volume number for the reference “Romenets et al., 2012”. Please confirm if this is fine.	Yes.
Q11	Please provide the meaning of superscript values in “Table 3.”	See below
Q12	We have included the numbers “10 & 11” in the last two rows of “Table 3.” Please confirm if this is fine.	See below

Q5:

Table 1:

The rows starting with “Non-Motor Symptoms Scale Total Score” and “SCOPA-Motor Total Score” are sections of the table just like “Cardinal Motor Features”. Thus:

There should be a horizontal rule above the row starting with “Non-Motor Symptoms Scale Total Score”.

There should NOT be a horizontal rule BELOW the row starting with “SCOPA-Motor Total Score”. (The horizontal rule ABOVE should be kept).

“Cardinal Motor Features” should not be boldface, and should look like “Non-Motor Symptoms Scale Total Score” and “SCOPA-Motor Total Score”.

Please add numbers 1-9 in front of the symptoms under “Non-Motor Symptoms Scale Total Score”, i.e. (1. Cardiovascular, 2. Sleep/Fatigue, ..., 9. Miscellaneous).

Please add in parentheses the abbreviation (CISI-PD) after “Clinical impression of severity Index” in the last row. Please capitalize “impression” and “severity”.

Table 3: There should be a horizontal rule below the row starting with “n (%)”. Also, please add a space in between n and (%). Also see Q11 and Q12

Table 4: Domains Clustering should be bold.

I have made comments in the manuscript.

Q11: The following table caption already in the table:

n, significant difference with cluster Sn (corrected $p < 0.05$)

indicates the meaning of the superscript values, where n is the number from 1-6. The “n”s should be italicized and the first “n” should be superscript (should look like captions in Table 2). Also, there should not be a comma after the “n”.

If this is not clear, then all 6 captions can be listed as follows, similar to Table 2, though it's a bit repetitive:

- 1 significant difference with cluster S1 (corrected $p < 0.05$)
- 2 significant difference with cluster S2 (corrected $p < 0.05$)
- 3 significant difference with cluster S3 (corrected $p < 0.05$)
- 4 significant difference with cluster S4 (corrected $p < 0.05$)
- 5 significant difference with cluster S5 (corrected $p < 0.05$)
- 6 significant difference with cluster S6 (corrected $p < 0.05$)

where the first number should be superscript.

Q12: These rows should *not* be numbered, as the numbering of the first nine rows references the numerical domains of the Non-motor symptoms scale. Could you please change “10. Motor symptoms” to “Motor symptoms” and “11. Variables not used in analysis” to “Other variables not used in analysis” (slight clarification here).



Parkinson's Disease Subtypes Identified from Cluster Analysis of Motor and Non-motor Symptoms

Jesse Mu¹, Kallol R. Chaudhuri^{2*}, Concha Bielza³, Jesus de Pedro-Cuesta⁴, Pedro Larrañaga³ and Pablo Martinez-Martin⁴

¹ Department of Computer Science, Boston College, Chestnut Hill, MA, United States, ² Department of Basic and Clinical Neuroscience, The Maurice Wohl Clinical Neuroscience Institute, King's College London, London, United Kingdom,

³ Computational Intelligence Group, Department of Artificial Intelligence, Universidad Politécnica de Madrid, Madrid, Spain,

⁴ National Center of Epidemiology, Instituto de Salud Carlos III, Madrid, Spain

OPEN ACCESS

Edited by:

Atsushi Takeda,
National Hospital Organization (NHO),
Japan

Reviewed by:

Kazushi Takahashi,
Saitama Medical University, Japan
Tien K. Khoo,
Griffith University, Australia

*Correspondence:

Kallol R. Chaudhuri
ray.chaudhuri@nhs.net

Received: 01 August 2017

Accepted: 01 September 2017

Published: xx September 2017

Citation:

Mu J, Chaudhuri KR, Bielza C, de Pedro-Cuesta J, Larrañaga P and Martinez-Martin P (2017) Parkinson's Disease Subtypes Identified from Cluster Analysis of Motor and Non-motor Symptoms. *Front. Aging Neurosci.* 9:301. doi: 10.3389/fnagi.2017.00301

Parkinson's disease is now considered a complex, multi-peptide, central, and peripheral nervous system disorder with considerable clinical heterogeneity. Non-motor symptoms play a key role in the trajectory of Parkinson's disease, from prodromal premotor to end stages. To understand the clinical heterogeneity of Parkinson's disease, this study used cluster analysis to search for subtypes from a large, multi-center, international, and well-characterized cohort of Parkinson's disease patients across all motor stages, using a combination of cardinal motor features (bradykinesia, rigidity, tremor, axial signs) and, for the first time, specific validated rater-based non-motor symptom scales. Two independent international cohort studies were used: (a) the validation study of the Non-Motor Symptoms Scale ($n = 411$) and (b) baseline data from the global Non-Motor International Longitudinal Study ($n = 540$). k -means cluster analyses were performed on the non-motor and motor domains (domains clustering) and the 30 individual non-motor symptoms alone (symptoms clustering), and hierarchical agglomerative clustering was performed to group symptoms together. Four clusters are identified from the domains clustering supporting previous studies: mild, non-motor dominant, motor-dominant, and severe. In addition, six new smaller clusters are identified from the symptoms clustering, each characterized by clinically-relevant non-motor symptoms. The clusters identified in this study present statistical confirmation of the increasingly important role of non-motor symptoms (NMS) in Parkinson's disease heterogeneity and take steps toward subtype-specific treatment packages.

Keywords: Parkinson's disease, subtypes, non-motor symptoms, motor symptoms, cluster analysis

INTRODUCTION

Parkinson's disease (PD) is classically considered a motor disorder, with resting tremor, rigidity, bradykinesia, and postural instability and gait disorder as its core features. However, the concept of PD has changed considerably in the last few years, now prompting a revision of its diagnostic criteria to include non-motor symptoms (NMS) in the core parameters (Postuma et al., 2015; Marras and Chaudhuri, 2016). There has been growing recognition that NMS in PD are caused by neurotransmitter pathway dysfunctions which involve both the central and peripheral nervous

systems (Jellinger, 2012; Gjerløff et al., 2015). The significant clinical heterogeneity of NMS in PD suggests the existence of specific non-motor subtypes (Marras and Chaudhuri, 2016; Sauerbier et al., 2016).

Previous cluster analyses have already identified motor- and non-motor-based clusters in PD patients (e.g., van Rooden et al., 2011; Erro et al., 2013; Ma et al., 2015; Pont-Sunyer et al., 2015). Recently, it has been argued that the recent concept of non-motor endophenotypes of PD provides a stronger basis for subtyping, since these relate to the central pathophysiology of specific neurotransmitter systems and are therefore likely to remain stable over time (Marras and Chaudhuri, 2016). As such, several studies have explored PD subtypes while considering motor subtypes and their association with non-motor aspects of the disease such as, psychopathology and cognition (Graham and Sagar, 1999; Reijnders et al., 2009; Selikhova et al., 2009; Burn et al., 2012; Flensburg Damholdt et al., 2012), REM sleep behavior disorder (Romenets et al., 2012), and daily visual activities (Seichepine et al., 2011). To our knowledge, however, no studies have used cluster analysis techniques to examine subtypes present in NMS only.

In this study, we used cluster analysis techniques to search for PD subtypes from a large, multi-center, international, and well-characterized cohort of patients across all stages, using a combination of motor cardinal features (bradykinesia, rigidity, tremor, axial signs) and comprehensive NMS assessed using specific validated rater-based scales. We believe this is the largest study of its size with these characteristics, and the first to focus on exclusively NMS-based phenotyping.

MATERIALS AND METHODS

Design

Data from two independent international studies were used in the analysis: the validation study of the Non-Motor Symptoms Scale (NMSS) ($n = 411$) (Martinez-Martin et al., 2009a) and baseline data from the global Non-Motor International Longitudinal Study (NILS) ($n = 540$) (Ray Chaudhuri et al., 2013). NILS has been adopted as a national study by the National Institute of Health Research in the UK (UKCRN No: 10084) and is a 5-year follow-up study addressing the range, nature, and natural history of NMS in PD across all motor stages. All data in NMSS and NILS have been anonymized and entered into a secure database at the National Center of Epidemiology, Carlos III Institute of Health, Madrid, Spain.

Patients

PD patients diagnosed according to internationally recognized criteria (Gibb and Lees, 1988; Lees et al., 2009) were included, and represented a mixed cohort of drug-naïve and treated PD across all disease stages. For the NMSS study, patients were older than 30 years, but for inclusion of NILS patients there was no age limit. Exclusion criteria were: inability to read, understand, or answer written questionnaires; comorbidity, sequelae, or any disorder interfering with the assessment of PD; and inability to give informed consent. Patient recruitment was carried out across 15 countries in America, Asia, and Europe from 2007 to 2011.

Assessments

For all patients, socio-demographic and historical data were recorded and the following assessments were applied:

1. The Scales for Outcomes in Parkinson's Disease-Motor (SCOPA-Motor), a scale with three dimensions: A. Examination (10 items); B. Activities of daily living (7 items); and C. Complications (4 items). Each item scores from 0 (normal) to 3 (severe), the total score ranging from 0 to 75. This scale was derived from the Unified Parkinson's Disease Rating Scale and showed high correlation with the original scale ($r > 0.85$) and satisfactory clinimetric attributes in validation studies (Marinus et al., 2004; Martinez-Martin et al., 2005).
2. The NMSS, a 30-item scale with nine domains: cardiovascular (2 items), sleep/fatigue (4 items), mood/apathy (6 items), perceptual problems/hallucinations (3 items), attention/memory (3 items), gastrointestinal tract (3 items), urinary function (3 items), sexual function (2 items), and miscellaneous (4 items). Each item scores from 0 to 12 (severity, 0–3, multiplied by frequency, 1–4) and the total NMSS score varies from 0 to 360, a value representing the total non-motor symptomatic burden (Chaudhuri et al., 2007; Martinez-Martin et al., 2009a).
3. The Hoehn and Yahr (HY) scale (Hoehn and Yahr, 1967).
4. The Clinical Impression of Severity Index for PD (CISI-PD), a global evaluation of motor signs, disability, motor complications, and cognitive status. Items are rated from 0 (normal) to 6 (very severe), the total score ranging from 0 to 24 (Martinez-Martin et al., 2006, 2009b).

Standard Protocol Approvals, Registrations, and Patient Consent

The NMSS validation study received ethical approval from the Carlos III Institute of Health, Madrid, Spain, and local research ethics committees (Martinez-Martin et al., 2009a). The NILS is included in the UK Department of Health portfolio of approved studies (UK CRN portfolio Nr. 10084) and has been approved at all relevant institutions and corresponding ethics committees/institutional review boards. All patients gave written informed consent before inclusion in accordance with the Declaration of Helsinki.

Statistical Analysis

SCOPA-Motor examination items were aggregated to obtain four “cardinal motor signs”: tremor (items 1 and 2), bradykinesia (item 3), rigidity (item 4), and axial signs (items 5 to 10). Additionally, an aggregate “motor complications” variable was obtained from the sum of items 18 to 21 (dyskinesias and motor fluctuations). All variables were standardized before clustering, and unstandardized afterwards for interpretation. Analyses were conducted in R version 3.2.4 (www.r-project.org) and Stata version 14 (<http://www.stata.com/>).

Cluster Analysis

k-means was used for cluster analysis. We performed two analyses on the data: the first clustering on the nine aggregate

non-motor symptom domains, the four cardinal motor signs (tremor, bradykinesia, rigidity, axial), and motor complications, henceforth the “domains clustering,” and the second on the 30 individual NMS of the NMSS only, henceforth the “symptoms clustering.” Average-linkage hierarchical agglomerative clustering on the 30 NMS, 4 motor signs, and motor complications was also performed to observe the grouping of the variables.

Various formal measures were used to determine the optimal number of clusters for the dataset. For the domains clustering, the optimal k according to the Gap Statistic and the 1-standard-error method (Tibshirani et al., 2001) was $k = 4$ (Supplementary Figure 1A). Other cluster determination methods suggested $k = 2, 3, 4$, where $k = 2, 3$ simply divided the data uninformatively into groups with varying levels of overall disease severity. Thus, $k = 4$ was selected to offer a good combination of model fit and parsimony. The same method was applied for the symptoms clustering, where the number of clusters was $k = 6$ (Supplementary Figure 1B).

Comparative Subgroup Analysis

For each variable in both clusterings, we used one-way ANOVA and χ^2 tests to, respectively, check the equality of variable means and proportions across the clusters found, using Bonferroni correction for multiple testing with corrected $p < 0.05$ considered significant. Differences among pairwise clusters were tested *post-hoc* using Tukey's range test for continuous means, or pairwise χ^2 tests for proportions, with Bonferroni correction both for the within-variable pairwise tests and the multiple variable comparisons.

To compare the domains and symptoms clusterings, we depicted cluster alignment with a contingency table, and computed the adjusted rand index (ARI) (Hubert and Arabie, 1985) to evaluate similarity between the two clusterings.

Lastly, to explore the relationship between symptom severity and disease duration, we computed the correlation of each variable with disease duration and fitted smoothed loess curves to the data both globally and for each cluster in the domains clustering.

RESULTS

Study Sample

Out of the 951 patients in the study, we used list wise deletion to exclude 47 patients due to missing measurements, resulting in 904 remaining patients. There were no significant differences between the included and excluded groups with respect to age, sex, disease duration, and HY ($\chi^2 \geq 0.19$). The characteristics of the sample included for analysis ($n = 904$) are displayed in Table 1. Patients were predominantly male (62.17%). 13.38% were in HY stage 1; 43.36% in stage 2; 29.65% in stage 3; 11.50% in stage 4; and 2.10% in stage 5.

Domains Clustering

Results from the k -means clustering on the nine non-motor domains, the four cardinal motor signs, and motor complications are reported in Table 2 along with additional variables not used

in the analysis (heatmap in Figure 1; boxplots in Supplementary Figure 2). Cluster means for all variables were found to be statistically significantly different except for age at disease onset and sex (adjusted $p < 0.05$). Specific pairwise differences are noted in the table.

Cluster D1 ($n = 428$) patients were mildly affected in all domains. This cluster was characterized by relatively lower disease durations and ages.

Cluster D2 ($n = 180$) patients were severely affected in non-motor domains but mildly affected in motor domains. This cluster had a severity of motor variables relatively similar to the cluster D1 (mild) subtype especially in tremor, but expressed significantly higher scores for non-motor domains than clusters D1 and D3, especially in the sleep/fatigue, mood/apathy, urinary, and miscellaneous domains. Except for motor complications, scores for every variable were statistically significantly different from those in cluster D3.

Cluster D3 ($n = 232$) patients were severely affected in motor domains but mildly affected in non-motor domains. Mean motor scores were greater than the means of clusters D1 and D2, with the exception of motor complications. Additionally, mean motor scores were less than D4, with the exception of

TABLE 1 | Description of the sample.

	Mean	SD	Median	Range
Age at study	64.28	9.86	65	34–89
Age at onset of Parkinson's disease (PD onset)	56.27	10.72	57	25–89
Duration of the disease (PD duration)	8.01	5.80	7	0–40
Non-Motor Symptoms Scale total score	50.45	41.72	39	0–225
Cardiovascular	1.74	3.26	0	0–24
Sleep/Fatigue	8.76	8.71	6	0–48
Mood/Apathy	8.67	11.54	4	0–60
Perceptual problems/Hallucinations	1.64	3.86	0	0–33
Attention/Memory	5.40	7.42	2	0–36
Gastrointestinal	5.53	6.78	3	0–36
Urinary	8.07	8.93	5	0–36
Sexual function	3.53	5.98	0	0–24
Miscellaneous	7.12	7.78	4	0–48
Cardinal motor features*				
Tremor	2.59	2.58	2	0–12
Bradykinesia	2.40	1.41	2	0–6
Rigidity	2.23	1.36	2	0–6
Axial	3.25	2.67	3	0–12
SCOPA-Motor total score	21.07	12.06	19	1–72
A. Examination	11.54	6.56	10	0–41
B. Activities of daily living	6.80	4.19	7	0–21
C. Complications	2.73	3.01	2	0–12
Clinical impression of severity Index	8.29	4.61	8	0–24

SD: Standard deviation. SCOPA: Scales for Outcomes in Parkinson's Disease.

*Scores derived from items of the SCOPA-Motor A. Examination.

TABLE 2 | Domains clustering summary.

Cluster		D1	D2	D3	D4
<i>n</i> (%)		428 (47%)	180 (20%)	232 (26%)	64 (7%)
Non-motor domains	1. Cardiovascular	0.7 (1.5) ^{2,4}	3.2 (3.7) ^{1,3,4}	1.1 (2.1) ^{2,4}	6.9 (6.4) ^{1,2,3}
	2. Sleep/fatigue	4.5(5.0) ^{2,3,4}	16 (8.7) ^{1,3,4}	7.5 (6.6) ^{1,2,4}	21.7 (9.7) ^{1,2,3}
	3. Mood/apathy	3.4 (4.8) ^{2,3,4}	19.2 (15.0) ^{1,3}	6.6 (8.0) ^{1,2,4}	21.7 (13.5) ^{1,3}
	4. Perception/hallucination	0.5 (1.7) ^{2,4}	2.7 (4.3) ^{1,3,4}	0.8 (1.8) ^{2,4}	9.7 (6.9) ^{1,2,3}
	5. Attention/memory	3.0 (4.5) ^{2,4}	10.5 (9.2) ^{1,3,4}	3.3 (4.4) ^{2,4}	14.5 (11.0) ^{1,2,3}
	6. Gastrointestinal	2.9 (4.1) ^{2,3,4}	8.5 (7.1) ^{1,3,4}	4.7 (5.3) ^{1,2,4}	17.4 (9.2) ^{1,2,3}
	7. Urinary	4.7 (6.2) ^{2,4}	14.0 (9.9) ^{1,3,4}	6.2 (6.7) ^{2,4}	20.3 (9.7) ^{1,2,3}
	8. Sexual function	1.7 (3.4) ^{2,3,4}	7.3 (7.8) ^{1,3}	2.4 (4.1) ^{1,2,4}	9.0 (9.9) ^{1,3}
	9. Miscellaneous	4.0 (4.8) ^{2,3,4}	13.2 (8.7) ^{1,3}	6.2 (6.8) ^{1,2,4}	14.5 (10.1) ^{1,3}
Motor symptoms	Axial	1.7 (1.5) ^{2,3,4}	3.6 (2.2) ^{1,3,4}	4.5 (2.3) ^{1,2,4}	8.2 (2.7) ^{1,2,3}
	Bradykinesia	1.6 (0.9) ^{2,3,4}	2.2 (1.1) ^{1,3,4}	3.5 (1.0) ^{1,2,4}	4.5 (1.3) ^{1,2,3}
	Rigidity	1.5 (0.9) ^{2,3,4}	1.8 (1.2) ^{1,3,4}	3.3 (1.0) ^{1,2,4}	4.2(1.2) ^{1,2,3}
	Tremor	2.0 (1.9) ^{3,4}	1.5 (1.8) ^{3,4}	4.1 (2.8) ^{1,2}	4.1 (4.1) ^{1,2}
	Motor complications	1.4 (2.1) ^{2,3,4}	3.1 (2.9) ^{1,4}	3.7 (2.9) ^{1,4}	7.0 (3.6) ^{1,2,3}
Variables not used in analysis	Sex (% male)	64	54	67	58
	CISI-PD total	5.5 (3.0) ^{2,3,4}	9.6 (3.8) ^{1,4}	10.1 (3.5) ^{1,4}	16.4 (4.6) ^{1,2,3}
	Age	62.5 (9.7) ⁴	65.2 (9.4) ⁴	64.9 (10.1) ⁴	71.1 (7.9) ^{1,2,3}
	PD onset	56 (10.5)	56.6 (10.6)	56.3 (11.3)	56.7 (10.6)
	PD duration	6.5 (4.7) ^{2,3,4}	8.6 (5.7) ^{1,4}	8.6 (5.7) ^{1,4}	14.4 (8.0) ^{1,2,3}

Unless otherwise specified, statistics are reported as mean (sd).

¹ Significant difference with cluster D1 (corrected $p < 0.05$).

² Significant difference with cluster D2 (corrected $p < 0.05$).

³ Significant difference with cluster D3 (corrected $p < 0.05$).

⁴ Significant difference with cluster D4 (corrected $p < 0.05$).

tremor, which was especially high. Importantly, CISI-PD scores of clusters D2 and D3 were not statistically significantly different, and no differences were observed in cluster D2 and cluster D3 age or disease duration.

Cluster D4 ($n = 64$) patients were severely affected in all domains, having the greatest symptom mean out of all four clusters with the exception of tremor. Consequently, patients in cluster D4 had the longest average disease duration and oldest ages, but did not have a significantly different age of disease onset.

Symptoms Clustering

k -means performed on the 30 individual NMS found six clusters ordered according to increasing CISI-PD score (Table 3, heatmap in Figure 2). Means of all symptoms were found to differ across clusters except for disease onset, sex, and tremor, with pairwise differences again noted in the table.

Cluster S1 ($n = 456$), the largest cluster representing 50% of the group, was similar to domains cluster D1, and was composed of patients relatively mildly affected in all NMS. Cluster S2 ($n = 201$) had higher mean symptom scores than cluster S1's in several cases, including restless legs syndrome (RLS), swallowing, and the miscellaneous domain, but could nonetheless be classified as a mild/moderate cluster.

Although clusters S3–S6 increased in motor and overall disease severity, they varied significantly in their non-motor

expression and each expressed a unique subset of NMS. These groups of NMS aligned well with the established non-motor domains. Cluster S3 ($n = 100$) mainly expressed domain 7 (urinary), while cluster S4 ($n = 73$) was affected severely in domain 3 (mood/apathy). Cluster S5 ($n = 54$) showed severe impact in most NMS but especially in domain 5 (attention/memory). Similarly, cluster S6 ($n = 20$) had severe scores across all NMS and motor features, but was most severely affected in the cardiovascular, perception/hallucination, and gastrointestinal NMSS domains. Overall, the symptoms clustering fragmented the domains clusters into smaller groups, as explored in the next section.

Comparison between Clusterings

Alignment of the D and S clusters is visualized in Table 4. While S1 grouped patients from D1 (mild) and D3 (motor-dominant), and D4 (severe) showed a dominant contribution from S5 (severe non-motor) and S6 (severe motor and non-motor), remaining clusters were fragmentarily distributed, as indicated by the low similarity between the clusterings ($ARI = 0.32$). For the domains clustering, patients in clusters mildly affected in non-motor domains (D1, D3) were distributed among the milder symptoms clusters (S1–S4, skewed left). Conversely, patients in clusters with severe NMS (D2, D4) were split among the various




FIGURE 1 | Heatmap of variables for each cluster in the domains clustering, separated by white lines according to nine non-motor domains, four cardinal motor features, motor complications, and four general variables not included in the analyses. Since symptoms have different scales, cluster means for each symptom are displayed as standardized scores relative to each overall symptom mean.

specific non-motor-dominant clusters (S2–S6), suggesting that the symptoms clustering is clinically more specific than the domains clustering.

Hierarchical Clustering on Variables

Hierarchical clustering on the 30 NMS and the four cardinal motor signs is depicted in **Figure 3**. Symptoms belonging to the same domain of the NMSS tended to cluster together, with some exceptions. Diplopia (domain 4) was grouped closer to domain 8 (sexual) symptoms than to symptoms in its own domain. Similarly, RLS (domain 2) was closer to domain 9 (miscellaneous) symptoms, and drowsiness (domain 2) with domain 5 (attention/memory) symptoms. Notably, tremor was the most isolated symptom, occupying a single branch at the top of the tree.

Correlation Analysis

Due to high variance, most variables had little to no correlation with disease duration (Supplementary Figure 3). In **Figure 4**, we plotted 4 variables especially relevant to the domains clustering against disease progression: CISI-PD, Tremor, Anxiety, and Depression. Notable differences in disease progression for each cluster can be seen in the scatterplots: for example, patients in NMS dominant cluster D2 actually tended to have higher scores for anxiety and depression at disease onset, decreasing with increasing disease duration.

DISCUSSION

We believe that this is the largest cluster analysis-based study of PD-related motor and NMS from a large, international, multi-center cohort. Previous cluster-analysis based studies have either focused on early/untreated Parkinson's disease (Erro et al., 2013; Pont-Sunyer et al., 2015) or lack detailed assessments based on the severity and frequency of non-motor domains and symptoms (van Rooden et al., 2011). Additionally, we believe this is the first study to perform cluster analysis exclusively on NMS to reveal NMS-specific subtypes.

The domains clustering's four clusters closely correspond with several previous studies (van Rooden et al., 2011;Erro et al., 2013; Ma et al., 2015; Pont-Sunyer et al., 2015), especially those reported by van Rooden et al. (2011). Both clusters D1 (mild) and D4 (severe) are groups which are present in most analyses, but unlike van Rooden et al., our data show that mean differences in disease duration *do* exist between mild and severe subtypes. Clusters D2 and D3 represent a divergence in symptomatic expression: D2 representing a non-motor dominant phenotype also described in many clinical phenotype-driven studies (Sauerbier et al., 2016), and D3 corresponding to the traditional motor-dominant view of PD. Due to these clusters' similar overall PD severity (CISI-PD) and duration, differences in disease progression do not explain the differences between D2 and D3. Finally, the high incidence of tremor in D3, even higher than D4, is interesting and reflects not only the motor-dominant subtype of van Rooden et al. but also the tremor-dominant/slow-progression cluster described by Ma et al. (2015).

Our correlation analysis demonstrates notable differences in disease progression among these clusters. The high initial depression and anxiety scores for cluster D2 suggests that patients susceptible to NMS-dominant PD can be identified by high NMS scores early after disease onset. Furthermore, the general improvement in depression and anxiety scores for this cluster contrasts with the relatively stable scores in clusters D1, D3, and D4.

From the symptoms clustering (**Figure 2**), six smaller clusters were identified. S1 was similar to D1. S2 to S6, while increasing in motor severity, expressed specific NMS, thus supporting the clinical concept of NMS-based subtyping. Cluster S2, with principal components including RLS, swallowing, pain, and others, may be a new finding from this study. Cluster S3, with significant urinary dysfunction, fits the descriptions by Erro et al. (2013), highlighting the relevance of this symptom as a

TABLE 3 | Symptoms clustering summary.

	Cluster	S1	S2	S3	S4	S5	S6
	<i>n</i> (%)	456 (50%)	201 (22%)	100 (11%)	73 (8%)	54 (6%)	20 (2%)
1. Cardiovascular	Lightheadedness	0.5 (1.1) ^{2,3,4,5,6}	1.7 (2.5) ^{1,5,6}	2.5 (3.4) ^{1,6}	1.9 (2.5) ^{1,6}	3.3 (3.7) ^{1,2,6}	7.7 (3.6) ^{1,2,3,4,5}
	Fainting	0.1 (0.6) ⁶	0.3 (1) ⁶	0.1 (0.5) ⁶	0.5 (1.2) ⁶	0.1 (0.5) ⁶	6.3 (2.9) ^{1,2,3,4,5}
2. Sleep/fatigue	Drowsiness	0.9 (1.8) ^{2,3,4,5,6}	2.4 (2.9) ^{1,5,6}	2.3 (2.9) ^{1,5,6}	2.2 (3) ^{1,5,6}	6.6 (4.2) ^{1,2,3,4}	5.2 (3) ^{1,2,3,4}
	Fatigue	1.2 (1.9) ^{2,3,4,5,6}	4.2 (3.7) ^{1,4,5}	4.5 (4) ^{1,5}	5.9 (4.2) ^{1,2}	8 (3.5) ^{1,2,3}	6.9 (3.1) ¹
	Insomnia	1.1 (2.2) ^{2,3,4,5,6}	3 (3.8) ^{1,4}	3.1 (4.2) ^{1,4}	5.3 (4.8) ^{1,2,3}	4.7 (4.8) ¹	5 (2.4) ¹
	RLS	0.5 (1.3) ^{2,4,5,6}	2.4 (3.5) ^{1,3,6}	1 (2.4) ^{2,5,6}	1.8 (3.2) ^{1,6}	3.2 (4.2) ^{1,3}	4.9 (2.6) ^{1,2,3,4}
3. Mood/apathy	Loss interest	0.4 (0.9) ^{2,4,5,6}	1.1 (1.8) ^{1,4,5,6}	0.6 (1.3) ^{4,5,6}	6.6 (3.9) ^{1,2,3,5}	4.5 (3.7) ^{1,2,3,4}	5.3 (2.8) ^{1,2,3}
	Loss activities	0.6 (1.3) ^{2,4,5,6}	1.9 (2.7) ^{1,4,5,6}	1 (2) ^{4,5,6}	7.8 (3.5) ^{1,2,3,5,6}	6 (4.5) ^{1,2,3,4}	4.7 (2.9) ^{1,2,3,4}
	Anxiety	0.8 (1.6) ^{2,4,5,6}	2.2 (2.9) ^{1,4,5,6}	1.6 (2.9) ^{4,5,6}	5.8 (4.3) ^{1,2,3}	6.4 (4.4) ^{1,2,3}	4.7 (3) ^{1,2,3}
	Depression	0.7 (1.4) ^{2,4,5,6}	2.7 (3.2) ^{1,3,4,5,6}	1.4 (2.3) ^{2,4,5,6}	7.4 (4) ^{1,2,3}	5.8 (4.3) ^{1,2,3}	5.3 (3.2) ^{1,2,3}
	Flat affect	0.3 (1) ^{2,4,5,6}	1.1 (2.1) ^{1,4,5}	0.8 (2.1) ^{4,5,6}	4.8 (4.3) ^{1,2,3,5}	2.9 (3.7) ^{1,2,3,4}	3 (2.1) ^{1,3}
	Loss pleasure	0.3 (1.1) ^{4,5,6}	1 (1.8) ^{4,5,6}	0.8 (1.8) ^{4,5,6}	7.3 (3.8) ^{1,2,3,5,6}	4.6 (4.2) ^{1,2,3,4}	3.6 (2.8) ^{1,2,3,4}
4. Perception/hallucination	Hallucination	0.2 (0.9) ^{5,6}	0.6 (1.7) ^{5,6}	0.7 (1.9) ^{5,6}	0.6 (1.8) ^{5,6}	2.7 (3.3) ^{1,2,3,4,6}	4.8 (3.3) ^{1,2,3,4,5}
	Delusion	0.1 (0.6) ^{4,5,6}	0.3 (1.4) ^{4,5,6}	0.1 (0.6) ^{4,5,6}	1.3 (3) ^{1,2,3,6}	2 (3.5) ^{1,2,3,6}	4.7 (3.4) ^{1,2,3,4,5}
	Diplopia	0.2 (0.8) ^{5,6}	0.6 (1.4) ^{5,6}	0.4 (1.6) ^{5,6}	0.8 (2.2) ^{5,6}	3.1 (4.4) ^{1,2,3,4}	3.6 (2.7) ^{1,2,3,4}
5. Attention/memory	Loss concentration	0.7 (1.5) ^{2,3,4,5,6}	2.6 (2.9) ^{1,5,6}	2.2 (3.4) ^{1,5,6}	2.9 (2.9) ^{1,5}	7.5 (3.9) ^{1,2,3,4}	5.1 (2.8) ^{1,2,3}
	Forget explicit	0.7 (1.3) ^{2,3,4,5,6}	2.2 (2.7) ^{1,5,6}	2.2 (3) ^{1,5,6}	2.1 (2.5) ^{1,5,6}	8.5 (3.2) ^{1,2,3,4,6}	4.8 (2.9) ^{1,2,3,4,5}
	Forget implicit	0.5 (1.2) ^{2,3,4,5,6}	1.9 (2.6) ^{1,5,6}	1.5 (2.7) ^{1,5,6}	1.8 (2.2) ^{1,5,6}	7.6 (4.1) ^{1,2,3,4,6}	5.1 (3.2) ^{1,2,3,4,5}
6. Gastrointestinal	Droping	0.6 (1.5) ^{2,3,4,5,6}	2.3 (3.3) ^{1,5,6}	3.3 (4.1) ^{1,6}	3.3 (4.2) ^{1,6}	4.2 (4.8) ^{1,2}	6.2 (3.3) ^{1,2,3,4}
	Swallowing	0.3 (0.8) ^{2,3,5,6}	2 (3) ^{1,6}	1.2 (2) ^{1,6}	1.2 (2) ⁶	2.3 (2.9) ^{1,6}	6.5 (4.2) ^{1,2,3,4,5}
	Constipation	1.5 (2.7) ^{2,3,4,5,6}	3 (3.8) ^{1,6}	3.8 (4.4) ^{1,6}	3.7 (4.4) ^{1,6}	4.5 (4.9) ^{1,6}	8.8 (4.4) ^{1,2,3,4,5}
7. Urinary	Urinary urgency	0.9 (1.7) ^{2,3,4,5,6}	1.8 (2.4) ^{1,3,5,6}	6.6 (3.8) ^{1,2,4}	2.4 (3.4) ^{1,3,5,6}	7.7 (4.3) ^{1,2,4}	6.2 (3.5) ^{1,2,4}
	Urinary frequency	0.9 (1.8) ^{3,4,5,6}	1.3 (1.9) ^{3,4,5,6}	7.7 (3.5) ^{1,2,4}	3.1 (3.6) ^{1,2,3,5,6}	6.2 (4.5) ^{1,2,4}	6.7 (3.1) ^{1,2,4}
	Nocturia	1.7 (2.3) ^{3,4,5,6}	2.6 (2.9) ^{3,4,5,6}	8.5 (3.5) ^{1,2,4}	4.6 (4.2) ^{1,2,3,5}	6.9 (4.4) ^{1,2,4}	7.1 (3.4) ^{1,2}
8. Sexual	Sex drive	0.7 (1.7) ^{2,3,4,5,6}	1.9 (3.4) ^{1,5}	3 (4.1) ^{1,5}	3.6 (4.4) ^{1,5}	6 (5.3) ^{1,2,3,4}	3.9 (5.3) ¹
	Sex dysfunction	0.7 (2) ^{3,4,5,6}	1.8 (3.3) ^{3,5}	3.4 (4.3) ^{1,2}	2.3 (4.2) ^{1,5}	5 (5.2) ^{1,2,4}	3.8 (5.6) ¹
9. Miscellaneous	Unexplained pain	0.7 (1.8) ^{2,3,4,5,6}	2.6 (3.9) ¹	2.4 (4) ¹	2.3 (3.6) ¹	4.3 (5) ¹	4.3 (2.2) ¹
	Gustation/olfaction	1.2 (2.5) ^{2,4,5,6}	4 (4.2) ¹	2.5 (3.6)	3 (3.9) ¹	4 (4.7) ¹	5.5 (4.7) ¹
	Weight change	0.8 (1.4) ^{2,4,5,6}	1.9 (3.1) ^{1,5}	1.8 (3) ⁵	2.1 (2.8) ¹	3.6 (4.3) ^{1,2,3}	4 (3.9) ¹
	Sweating	0.6 (1.6) ^{2,3,4,5,6}	2.5 (3.9) ¹	2.4 (3.9) ¹	2.1 (3.6) ¹	2.8 (3.9) ¹	3.9 (3.4) ¹
10. Motor symptoms	Axial	2.3 (2) ^{2,3,4,5,6}	3.5 (2.6) ^{1,5,6}	4 (2.8) ^{1,6}	4.4 (2.7) ^{1,6}	5.3 (3.3) ^{1,2,6}	8.5 (2.3) ^{1,2,3,4,5}
	Bradykinesia	2.1 (1.2) ^{4,5,6}	2.5 (1.4) ⁶	2.5 (1.4) ⁶	2.9 (1.4) ^{1,6}	3 (1.6) ^{1,6}	4.5 (1.7) ^{1,2,3,4,5}
	Rigidity	2 (1.2) ⁶	2.3 (1.5) ⁶	2.3 (1.4) ⁶	2.5 (1.3) ⁶	2.6 (1.5) ⁶	4.7 (1) ^{1,2,3,4,5}
	Tremor	2.6 (2.4) ⁶	2.5 (2.4) ⁶	2.1 (2.5) ⁶	2.5 (2.6) ⁶	2.4 (2.8) ⁶	5.8 (5.3) ^{1,2,3,4,5}
	Motor comp	1.9 (2.5) ^{2,4,5,6}	3.5 (3) ^{1,6}	2.7 (2.7) ⁶	3.7 (3.3) ^{1,6}	3.9 (3.7) ^{1,6}	8 (3.7) ^{1,2,3,4,5}
11. Variables not in analysis	Sex	67	57	64	49	63	50
	CISI PD total	6.4 (3.5) ^{2,3,4,5,6}	9.3 (4.1) ^{1,5,6}	9.4 (4.1) ^{1,6}	10.7 (4.6) ^{1,6}	11.8 (5.5) ^{1,2,6}	18.4 (4.2) ^{1,2,3,4,5}
	Age	62.8 (10) ^{3,5,6}	63.8 (10) ⁵	68.3 (7.5) ¹	63.1 (9.2)	69.7 (9.2) ^{1,2}	72.1 (7.7) ¹
	PD onset	56.2 (10.8)	55 (10.7)	58.2 (10)	54.8 (10.3)	58.8 (11.7)	59.2 (8.6)
	PD duration	6.6 (4.8) ^{2,3,5,6}	8.8 (5.9) ¹	10.1 (6.7) ¹	8.3 (4.9)	11 (8.5) ¹	12.8 (6.5) ¹

Unless otherwise specified, statistics are reported as mean (sd).

n, significant difference with cluster *S_n* (corrected *p* < 0.05)






TABLE 4 | Contingency table describing cross-categorization of individuals in the domains and symptoms clusters.



		Symptoms clusters						
Domains clusters		S1	S2	S3	S4	S5	S6	Total
	D1	335	64	26	3	0	0	428
	D2	0	54	46	49	31	0	180
	D3	121	77	22	12	0	0	232
	D4	0	6	6	9	23	20	64
Total		456	201	100	73	54	20	904

cluster, essentially expresses two NMS subtypes, one of them with motor symptoms. The low numbers observed in some cells do not allow consistent clinical interpretation. The hierarchical clustering (Figure 3) indicates that the symptoms grouping in the NMSS dimensions works as expected, as most items in each domain group together, with the exception of tremor.

What are the clinical implications of these clusterings? First, our analysis represents statistical conformation of NMS-dominant presentation of PD. The specific expression of several NMS domains such as, mood/anxiety, sleep/fatigue, cognition, and urinary function suggests that these subgroups may have different patterns of neurodegeneration involving the brain's various non-dopaminergic pathways, possibly in excess of dopaminergic degeneration, as suggested by several authors (Jellinger, 2012). Second, clinical recognition of subtypes using ad hoc criteria would allow for the development of truly subtype-specific treatment packages for PD (Marras and Chaudhuri, 2016). Third, clinical characterization of these groups will allow studies of natural history of specific subtypes.

The clinical non-specificity of D1, S1, D4, and S6, with extremely diverse disease durations and severities, contrasts with the precisely characterized clusters S3, S4, and S5, with dominant expression of urinary, mood/apathy and attention/memory symptoms, respectively, at intermediate stages of the disease. This pattern is in line with the notion of "phenotypic convergence" proposed by Warren et al. (2013) as a key clinical feature of the spread of neurodegenerative disorders due to abnormal protein aggregates. The identified clusters may represent distinct footprints of large-scale network disintegration which necessitates translation to clinical management. The Warren et al. concept is in line with the general etiologic hypothesis for late-life neurodegenerative diseases proposed by de Pedro-Cuesta et al. (2016).

Like any cohort-based, cluster-analysis driven study, there are several limitations of this analysis. Due to the data collection methods of the two studies used, selection due to prevalence bias, i.e., sample overrepresentation of patients with higher survival, is unlikely to explain this clustering; however, clustering at early PD stages may have been undermined by poorly recorded symptoms prior to diagnosis. Furthermore, we did not report a control group, although our intention was not to describe the symptoms as discriminant from normal subjects. Lastly, in the treated patients in our sample, NMS symptoms could be influenced



by dopaminergic therapy, including depression via pramipexole (Barone et al., 2010), sleep disorders via rotigotine (Trenkwalder et al., 2011), and others. The numbers of patients undergoing specific treatments are too small to conduct meaningful analyses of the effects of such therapies; we expect, however, that such

effects on single non-motor components do not significantly alter the trends observed in total NMSS scores across all treated and drug-naïve patients in our sample. Conversely, our study has several notable strengths: (1) the sample size, which to our knowledge is the largest international

sample in this kind of study; (2) the inclusion of patients in all disease stages; and (3) the use of detailed assessments both for motor and NMS.

In conclusion, we present statistical confirmation of the growing recognition of NMS-dominant presentation of PD and its heterogeneity. The clinical recognition of these subtypes could allow for subtype-specific treatment packages for PD, and in the future, clinical characterization of these groups will allow for studies of natural history of the various non-motor dominant clusters identified in this paper. Translating results to clinical management or experimental designs would require the identification of inclusion and exclusion criteria of patients into specific subgroups.

AUTHOR CONTRIBUTIONS

JM conducted the statistical analysis and interpretation of the data, drafted the methods and results section of the paper, and revised the manuscript for content. KC obtained the data, designed the study, and drafted and revised the introduction and conclusion sections of the paper. CB, JdP, and PL revised the manuscript for content and contributed to the analysis and interpretation of data. PM obtained the data, designed the study, and drafted and revised the introduction and conclusion sections of the paper.

REFERENCES

- Barone, P., Poewe, W., Albrecht, S., Debieve, C., Massey, D., Rascol, O., et al. (2010). Pramipexole for the treatment of depressive symptoms in patients with Parkinson's disease: a randomised, double-blind, placebo-controlled trial. *Lancet Neurol.* 9, 573–580. doi: 10.1016/S1474-4422(10)70106-X
- Burn, D. J., Landau, S., Hindle, J. V., Samuel, M., Wilson, K. C., Hurt, C. S., et al. (2012). Parkinson's disease motor subtypes and mood. *Mov. Disord.* 27, 379–386. doi: 10.1002/mds.24041
- Chaudhuri, K. R., Martinez-Martin, P., Brown, R. G., Sethi, K., Stocchi, F., Odin, P., et al. (2007). The metric properties of a novel non-motor symptoms scale for Parkinson's disease: results from an international pilot study. *Mov. Disord.* 22, 1901–1911. doi: 10.1002/mds.21596
- de Pedro-Cuesta, J., Martinez-Martin, P., Rábano, A., Ruiz-Tovar, M., Alcalde-Cabero, E., and Calero, M. (2016). Etiologic framework for the study of neurodegenerative disorders as well as vascular and metabolic comorbidities on the grounds of shared epidemiologic and biologic features. *Front. Aging Neurosci.* 8:138. doi: 10.3389/fnagi.2016.00138
- Erro, R., Vitale, C., Amboni, M., Picillo, M., Moccia, M., Longo, K., et al. (2013). The heterogeneity of early Parkinson's disease: a cluster analysis on newly diagnosed untreated patients. *PLoS ONE* 8:e70244. doi: 10.1371/journal.pone.0070244
- Flensburg Damholdt, M., Shevlin, M., Borghammer, P., Larsen, L., and Ostergaard, K. (2012). Clinical heterogeneity in Parkinson's disease revisited: a latent profile analysis. *Acta Neurol. Scand.* 125, 311–318. doi: 10.1111/j.1600-0404.2011.01561.x
- Gibb, W. R., and Lees, A. J. (1988). The relevance of the Lewy body to the pathogenesis of idiopathic Parkinson's disease. *J. Neurol. Neurosurg. Psychiatr.* 51, 745–752. doi: 10.1136/jnnp.51.6.745
- Gjerløff, T., Fedorova, T., Knudsen, K., Munk, O. L., Nahimi, A., Jacobsen, S., et al. (2015). Imaging acetylcholinesterase density in peripheral organs in Parkinson's disease with 11C-donepezil PET. *Brain* 138(Pt 3), 653–663. doi: 10.1093/brain/awu369

FUNDING

This study was funded by the Spanish Ministry of Economy and Competitiveness through the Cajal Blue Brain (C080020-09; the Spanish partner of the Blue Brain initiative from École Polytechnique Fédérale de Lausanne) and TIN2016-79684-P projects, the Regional Government of Madrid through the S2013/ICE-2845-CASI-CAM-CM project, the European Union's Horizon 2020 research and innovation programme under grant agreement No. 720270, and the National Institute of Health Research in the UK (UKCRN No: 10084).

ACKNOWLEDGMENTS

We acknowledge data collection efforts by all contributors, collaborators, and administrative staff of the NELS study and the scale development group of the NMSS. KRC also acknowledges the NIHR Biomedical Research Center, Kings College, London, and the National Parkinson Foundation, USA.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fnagi.2017.00301/full#supplementary-material>

- Graham, J. M., and Sagar, H. J. (1999). A data-driven approach to the study of heterogeneity in idiopathic Parkinson's disease: identification of three distinct subtypes. *Mov. Disord.* 14, 10–20. doi: 10.1002/1531-8257(199901)14:1<10::AID-MDS1005>3.0.CO;2-4
- Hoehn, M. M., and Yahr, M. D. (1967). Parkinsonism: onset, progression, and mortality. *Neurology* 17, 427–442. doi: 10.1212/WNL.17.5.427
- Hubert, L., and Arabie, P. (1985). Comparing partitions. *J. Classif.* 2, 193–218. doi: 10.1007/BF01908075
- Jellinger, K. A. (2012). Neuropathology of sporadic PD disease: evaluation and change of concepts. *Mov. Disord.* 27, 8–30. doi: 10.1002/mds.23795
- Lees, A. J., Hardy, J., and Revesz, T. (2009). Parkinson's disease. *Lancet* 373, 2055–2066. doi: 10.1016/S0140-6736(09)60492-X
- Ma, L. Y., Chan, P., Gu, Z. Q., Li, F. F., and Feng, T. (2015). Heterogeneity among patients with Parkinson's disease: cluster analysis and genetic association. *J. Neurol. Sci.* 351, 41–45. doi: 10.1016/j.jns.2015.02.029
- Marinus, J., Visser, M., Stiggelbout, A. M., Rabey, J. M., Martinez-Martin, P., Bonuccelli, U., et al. (2004). A short scale for the assessment of motor impairments and disabilities in Parkinson's disease: the SPES/SCOPA. *J. Neurol. Neurosurg. Psychiatr.* 75, 388–395. doi: 10.1136/jnnp.2003.017509
- Marras, C., and Chaudhuri, K. R. (2016). Non-motor features of Parkinson's disease subtypes. *Mov. Disord.* 31, 1095–1102. doi: 10.1002/mds.26510
- Martinez-Martin, P., Benito-Leon, J., Burguera, J. A., Castro, A., Linazasoro, G., Martinez-Castrillo, J. C., et al. (2005). The SCOPA-motor scale for assessment of Parkinson's disease is a consistent and valid measure. *J. Clin. Epidemiol.* 58, 674–679. doi: 10.1016/j.jclinepi.2004.09.014
- Martinez-Martin, P., Forjaz, M. J., Cubo, E., Frades, B., de Pedro Cuesta, J., and ELEG Project Members (2006). Global vs. factor-related impression of severity in Parkinson's disease: a new clinimetric index (CISI-PD). *Mov. Disord.* 21, 208–214. doi: 10.1002/mds.20697
- Martinez-Martin, P., Rodriguez-Blazquez, C., Abe, K., Bhattacharyya, K. B., Bloem, B. R., Carod-Artal, F. J., et al. (2009a). International study on the psychometric attributes of the non-motor symptoms Scale in Parkinson disease. *Neurology* 73, 1584–1591. doi: 10.1212/WNL.0b013e3181c0d416

- Martinez-Martin, P., Rodriguez-Blazquez, C., Forjaz, M. J., de Pedro, J., and Spanish-American longitudinal PD patient study group (2009b). The clinical impression of severity index for Parkinson's Disease: international validation study. *Mov. Disord.* 24, 211–217. doi: 10.1002/mds.22320
- Pont-Sunyer, C., Hotter, A., Gaig, C., Seppi, K., Compta, Y., Katzenschlager, R., et al. (2015). The onset of nonmotor symptoms in Parkinson's disease (the ONSET PD study). *Mov. Disord.* 30, 229–237. doi: 10.1002/mds.26077
- Postuma, R. B., Berg, D., Stern, M., Poewe, W., Olanow, C. W., Oertel, W., et al. (2015). MDS clinical diagnostic criteria for Parkinson's disease. *Mov. Disord.* 30, 1591–1601. doi: 10.1002/mds.26424
- Ray Chaudhuri, K., Rojo, J. M., Schapira, A. H., Brooks, D. J., Stocchi, F., Odin, P., et al. (2013). A proposal for a comprehensive grading of Parkinson's disease severity combining motor and non-motor assessments: meeting an unmet need. *PLoS ONE* 8:e57221. doi: 10.1371/journal.pone.0057221
- Reijnen, J. S., Ehrt, U., Lousberg, R., Aarsland, D., and Leentjens, A. F. (2009). The association between motor subtypes and psychopathology in Parkinson's disease. *Parkinsonism Relat. Disord.* 15, 379–382. doi: 10.1016/j.parkreldis.2008.09.003
- Romenets, S. R., Gagnon, J. F., Latreille, V., Panniset, M., Chouinard, S., Montplaisir, J., et al. (2012). Rapid eye movement sleep behavior disorder and subtypes of Parkinson's disease. *Mov. Disord.* 27, 996–1003. doi: 10.1002/mds.25086
- Sauerbier, A., Jenner, P., Todorova, A., and Chaudhuri, K. R. (2016). Non motor subtypes and Parkinson's disease. *Parkinsonism Relat. Disord.* 22(Suppl. 1), S41–S46. doi: 10.1016/j.parkreldis.2015.09.027
- Seichepine, D. R., Neargarder, S., Miller, I. N., Riedel, T. M., Gilmore, G. C., and Cronin-Golomb, A. (2011). Relation of Parkinson's disease subtypes to visual activities of daily living. *J. Int. Neuropsychol. Soc.* 17, 841–852. doi: 10.1017/S1355617711000853
- Selikhova, M., Williams, D. R., Kempster, P. A., Holton, J. L., Revesz, T., and Lees, A. J. (2009). A clinico-pathological study of subtypes in Parkinson's disease. *Brain* 132(Pt 11), 2947–2957. doi: 10.1093/brain/awp234
- Tibshirani, R., Walther, G., and Hastie, T. (2001). Estimating the number of clusters in a data set via the gap statistic. *J. R. Stat. Soc. B Stat. Methodol.* 63, 411–423. doi: 10.1111/1467-9868.00293
- Trenkwalder, C., Kies, B., Rudzinska, M., Fine, J., Nikl, J., Honczarenko, K., et al. (2011). Rotigotine effects on early morning motor function and sleep in Parkinson's disease: a double-blind, randomized, placebo-controlled study (RECOVER). *Mov. Disord.* 26, 90–99. doi: 10.1002/mds.23441
- van Rooden, S. M., Colas, F., Martinez-Martin, P., Visser, M., Verbaan, D., Marinus, J., et al. (2011). Clinical subtypes of Parkinson's disease. *Mov. Disord.* 26, 51–58. doi: 10.1002/mds.23346
- Warren, J. D., Rohrer, J. D., Schott, J. M., Fox, N. C., Hardy, J., and Rossor, M. N. (2013). Molecular nexopathies: a new paradigm of neurodegenerative disease. *Trends Neurosci.* 36, 561–569. doi: 10.1016/j.tins.2013.06.007

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Mu, Chaudhuri, Bielza, de Pedro-Cuesta, Larrañaga and Martinez-Martin. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.