

Identifying biomarkers of spatial memory with direct brain recordings in the Treasure Hunt task

Miller, J.¹, Watrous, A.¹, Novich, C.¹, Lee, S.A.¹, Sperling, M.², Sharan, A.², Worrell, G.³, Berry, B.³, Lega, B.⁴, Jobst, B.⁵, Davis, K.⁶, Sheth, S.⁷, Das, S.⁶, Stein, J.⁶, Gorniak, R.², Rizzuto, D.⁸, Jacobs, J.¹

COLUMBIA UNIVERSITY
IN THE CITY OF NEW YORK

¹Columbia University, ²Thomas Jefferson University Hospital, ³Mayo Clinic, ⁴University of Texas, Southwestern, ⁵Dartmouth University, ⁶Hospital of the University of Pennsylvania, ⁷Columbia University Medical Center, ⁸University of Pennsylvania

Introduction

- Treasure Hunt: a novel 3D adaptation of a Morris water maze, designed to investigate memory for object locations in a rich virtual environment.
- Key features: 1) Rapid acquisition of data, 2) Well defined encoding periods, 3) Multiple measures of performance, 4) Fun to play.
- Main finding: Human intracranial recordings reveal low frequency power increases related to good spatial memory at frequencies lower than traditional rodent theta.

Task Design




Trial: Subjects navigate a 3D beach opening treasure chests that may contain objects. After four chests, subjects are probed with the objects and must indicate the original locations. Gain points based on accuracy.

Session: 40 trials, 100 objects




- 43 patients
- 90 sessions
- 5060 bipolar electrodes



Behavioral results


Normalized accuracy: distance between response location and true object location, ranked by all possible response locations. Chance = 0.5.




Multivariate decoding of spatial memory

Classification method: L2 regularized logistic regression trained on memory encoding. Features: oscillatory power at 8 (right) or 50 (below) log-spaced frequencies between 1 and 200 Hz.

Successful recall: better than median response and not low confidence.




Neural signature of good spatial memory



Low frequency increases and high frequency decreases in both cortex and hippocampus.

Theta power predicts memory

Increased low frequency power in temporal cortex and hippocampus is associated with better spatial memory. The precise frequency of this effect often varies from electrode to electrode, especially in TC.



Summary

- Reliable decoding of good spatial memory for object locations in a novel 3D task.
- Prominent increases in low frequency oscillatory power are linked to successful spatial learning.

Reprints/questions: jonathan.miller@columbia.edu