



Received on 26 January, 2015; received in revised form, 21 March, 2015; accepted, 29 March, 2015; published 01 April, 2015

HYPOLIPIDAEMIC ACTIVITY OF THE SIDDHA MEDICINAL PLANTS

A. Manoharan*¹, M. Manjula², H. Mubarak³ and C.V. Chitti Babu⁴

Department of General Medicine, Government Siddha Medical College¹, Palayamkottai Tamil Nadu, India

Department of Medicine, TVMCH², Tirunelveli, Tamil Nadu, India

Senior Research Fellow(S), SCRUC³, Palayamkottai Tamil Nadu, India

Department of Botany, Presidency College⁴, Chennai Tamil Nadu, India

Keywords:

Siddha medicines,
Adhi dhoola noi (Obesity),
Hyperlipidaemia, Atherosclerosis

Correspondence to Author:

Dr. A. Manoharan

Reader, Department of Pothu
Maruthuvam, Government Siddha
Medical College, Palayamkottai,
Tirunelveli 627002 Tamil Nadu,
India

E-mail: drmanoharan25@gmail.com

ABSTRACT: Hyperlipidaemia (*Adhi dhoola noi*) is a well known metabolic disorder, prevalent in 30-40% of the people worldwide. It is one of the most important factors of morbidity and mortality related to atherosclerotic cardiovascular disease and diabetes. Research articles prove hyperlipidaemia as an important factor that determines the risk of atherosclerotic heart disease and endocrinal disorders. Siddha Medical system is one of the ancient one in which medicinal plants contributes a vital role in the making of single and compound formulations. Siddha medicinal plants can play an important role in the reduction of hyperlipidemic state with proven evidence based research. This warrants introduction of Siddha medicinal plants in the treatment of hyperlipidaemia. This review article recites various polyherbal Siddha formulations and single herbs as a hypolipidaemic drugs with scientifically proven measures which are equally effective, safe to use, affordable and easily acceptable.

INTRODUCTION: Hyperlipidaemia is a disorder of lipid metabolism. This disorder manifests as elevation of plasma cholesterol, triglycerides (or) low HDL level or all of the above. Based on CARDIA (Coronary Artery Risk Development In Young Adults) study, which was conducted among 5000 young adults of age group 18-30 years with an increased Body Mass Index (BMI) are at a high risk of developing complications due to hyperlipidaemia. Increased VLDL, LDL and total cholesterol levels and decreased HDL level renders more atherogenic blood vessels which could result in coronary insufficiency (or) ischemic heart disease.

National cholesterol Education Program (NCEP) has mentioned about Detection, types, evaluation and management of hyperlipidaemia in adults in treatment panel ATP-III.¹

Hyperlipidaemia results in metabolic syndrome which is characterized by obesity, Insulin resistance and endothelial cell dysfunction which ultimately ends in Hypertension, Diabetes (or) Stroke.² According to the World Health Organization (WHO) most of the world's population is dependent on traditional medicines. Siddha Literature has described the various usages of polyherbal formulations in hyperlipidemia. Even though there are variable Siddha formulations, this study mainly includes the use of *Venthamarai chooranam*, *Neerzhivu chooranam*, *Thiripala chooranam*, *Maruthampattai chooranam*, *Kariveppilai chooranam* in the treatment of hyperlipidaemia. It also enumerates the phyto

QUICK RESPONSE CODE 	DOI: 10.13040/IJPSR.0975-8232.IJP.2(4).145-54 Article can be accessed online on: www.ijpjournal.com
DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.2(4).145-54	

chemistry, parts to be used along with botanical and common names.^{3, 4, 5}

This study provides a list of multi herbal formulations and comparisons of their efficacy and analysis of individual chemical constituents of various herbal formulations that could have a hyperlipidaemic lowering activity. There is a warning alarm mentioned by National commission of Macro economics and health (NCMH) in India has estimated that by 2015 approximately 52 million Indian patients will develop coronary artery disease; out of this about 23 millions will be less than 40 years of age.⁶

Siddha Medicinal Plants compared with that of Current research:

Siddha medicine is widely used in various diseases like Diabetes, Atherosclerosis, Rheumatoid arthritis etc. Siddha Medicines have been prepared as multi

herbal formulations that have a hyperlipidaemic lowering activity since ages which reinforces the effectiveness of Siddha medicinal plants in hyperlipidaemia. Several medicinal plants have been proved to act through various mechanisms like Inhibition of Microsomal Triglyceride Transfer Protein (MTP), Acyl coenzyme A (COA) Cholesterolacyltransferase (ACAT), Diacyl glycerol and acyl transferase (DGAT) and farnesoid X receptors (FXR). Among various Siddha herbal formulations, *Venthamarai Chooranam* (Table 1), *Madhumega Chooranam* (Table 2), *Thribala Chooranam* (Table 3), *Maruthampattai Chooranam* (Table 4), *Karivepillai Chooranam* (Table 5) which are having multiple bioactive ingredients act as lowering the hyperlipidaemic state. Some of the interested medicinal plants are described based on the proven scientific studies.

TABLE1: INGREDIENTS OF VENTHAMRAI CHOORANAM

S. No	Botanical Name/Tamil name	Family	Part to be used	Important Alkaloids	Actions
1.	<i>Elettaria cardamomum</i> (L) Maton ELAM	Zingiberaceae	Fruit & Seed	Oleum, Cardamoni, Terpinol	Hypolipidaemia, Hypoglycemic ^{7, 8, 9}
2.	<i>Zingiber officinalis</i> Roscoe CHUKKU	Zingiberaceae	Rhizome	Zingiberneol, β -bisabolene, Gingerin, Oleoresin	Hypolipidaemia, Diabetic, Dyslipidemic, Antioxidant ^{10, 11}
3.	<i>Piper longum</i> , Linn THIPPLI	Piperaceae	Root & fruit	Piperine, Guniesine, Methyl Pluvitilo, Pipataline	Hypolipidaemia, Antioxidant, Cardiac Disease ^{13, 14}
4.	<i>Glycyrrhiza glabra</i> , Linn ATHIMADHURAM	Fabaceae	Root	Glycyrrbizin, Glycyrrhizinate, Coumarin, Vitc, α -terpinene,	Hypolipidaemia, Hypoglycemic ^{15, 16}
5.	<i>Anethum graveolens</i> Linn SADAKUPPAI	Apiaceae	Leaves	Flavanoids, Steroids -Xanthome glucoside	Hypolipidaemia ^{17, 18}
6.	<i>Cuminum cyminum</i> CHEERAKAM	Apiaceae	Seed	Cuminaldehyde	Hypolipidaemia ^{19, 20}
7.	<i>Nelumbo nucifera</i> Garten THAMARAI	Nelumbonaceae	White flowers	Lupeol, β -Sitosterol, Nuciferine, N-Norarmeparine	Lipolytic, Cardiac Disease ^{21, 22, 23}

Elettaria cardamomum (L):

Chaulhari .HS., et al.,⁸ has mentioned in review article as Aqueous extract of *Elettaria cardamomum* (100 - 200mg / kg per oral) for 30 days has significantly decreased the level of Total cholesterol, Triglycerides, LDL - C, VLDL and has increased HDL -C Levels.

Zingiber officinale:

Bhandhari, et al.,¹⁰ has reviewed that ethanolic extract of Ginger (200mg /kg) orally for 20 days produced significant reduction of blood sugar in diabetic rats and also reduction in lipid levels are noted.

Piper longum:

Jin.Z.,^{13,14} et al., has reviewed that use of ethanolic extract of piper has reduced the cholesterol levels.

Glycyrrhiza glabra:

Santhosh kumar Maurya et al.,¹⁵ reviewed that ethanolic extract of Glycyrrhiza Glabra (100mg / kg / day) for 30 days has markedly reduced the lipids level.

Antheum graveolens:

Iyer D et al.,¹⁸ has previewed that Ethanolic extract of that plant reduces the cholesterol levels.

Cuminum cyminum:

Vaibhav Srivasta et al.,¹⁹ revised that Ethanolic extract of the seeds given for 20 days has significantly lowered the lipid levels.

Nelumbo nucifera:

Subhasini et al.,²² has studied the effects of the plant in male wistar rats which has proved anti lipidaemic activity.

Terminalia chebula:

Maruthappan et al.,²⁵ 1.05mg, 1kg, 1Bw & 2.10mg / kg/ bw for 14 days given for atherogenic diet induced hyperlipidaemic rats has reduced the lipoprotein levels.

Murraya Koenigii:

Khan BA., et al.,²⁷ has reviewed tha tethanolic Petroleum and chloroform leaf extract has markedly reduced the blood sugar and lipid levels.

Phyllanthus emblica:

Arunab Bhattacharya et al.,³² has reviewed that methanol and water solvent extract (5 - 10mg / kg) for 7 days has inhibited lipid peroxidation.

TABLE 2: INGREDIENTS OF MADHUMEGA CHOORANAM (MMC)

S. No	Botanical name/ tamil name	Family	Parts to be used	Phytochemical constituents	Actions
1.	<i>Terminalia chebula</i> (Retz) KADUKKAI	Combretaceae	Fruit	Phenolic Compounds, punicalagin, Terflavin-A, Terchebulin Girin,	Antiobesity, Hypolipidaemia ^{24, 25}
2.	<i>Murraya koenigii</i> (L) KARIVEPPILAI	Rutaceae	Leaf	isomahanibine, Koenimbine, Koengicine	Hypolipidaemia, Hypoglycemic ^{26, 27}
3.	<i>Phyllanthus emblica</i> (Linn) Gaertn NELLIKAI	Euphorbiaceae	Fruit	Trigalloylglucose, phyllembic acid, terchebin	Hypolipidaemia, Antioxidants, Rich source of vit c ^{28, 29, 30}
4.	<i>Tinospora cordifolia</i> (Willd) SEENTHIL	Menispermaceae	Stem	Flavanoid, Diterpene-10 - Hydroxy columbin, Tinosporidine	Hypolipidaemia, Antidiabetic, Anti oxidant ^{31, 32}
5.	<i>Syzygium cumini</i> (L) <i>S.Jambolanum</i> (Lam) NAAVAL	Myrtaceae	Bark, seed	Betalinic acid Kaempferol Isoquerlitrin	Hypolipidaemia, Anti diabetic, Antioxidant ^{33, 34}
6.	<i>Cyperus rotundus</i> (Linn)	Cyperaceae	Tuber	Flavonoid glucoside cypernel I, II Cyptertundone	Hypolipidaemia, Antihypertensive ^{35, 36, 37}
7.	<i>Phyllanthus amarus</i> , Schume Tnonn KEEZHANELLI	Euphorbiaeae	Whole plant	Ellagitannin- Phyllanthin D	Antidiabetic, Hypolipidaemic ^{38, 39}

Tinospora cordifolia:

Veena sharma et al.,³³ has reviewed that petroleum and other plant extract in rats (200mg / kg for 28 days) produces a marked reduction in body weight.

Syzygium cumin:

Sharma SB et al.,³⁴ has reviewed that Ethanolic seed extract of Cumini in diabetic rabbits showed marked changes in lipoprotein.

Cyprus rotundus:

Bambhole. VD et al.,³⁷ has reviewed that Aqueous and Alcoholic extract for 90 days showed. lipolytic action and it has mobilised fat in rats.

Phyllanthus amarus:

A.K.K Khana.F et al.,³⁸ has reviewed tht plant extract (250mg 1kg/ weight) for 30 days in Triton Wistar Rat 1339 showed that it inhibits hepatin

cholesterol biosynthesis and increased faecal bile acid excretion and enhanced plasma lecithin (LCAT) and cholesterol acyl transferase.

***Terminalia chebula*:** Priya. F., et al.,⁴² has reviewed that 250 - 500mg / kg powder given in

Triton Wistar Rat 1339 showed that it reduces total cholesterol.

***Phyllanthus emblica*:**

Yokazawa.T., et al.,⁴³ has reviewed that (10 - 40mg/kg/bw) for 2 months reduces the hyperlipidaemia and has antioxidative actions.

TABLE 3: INGREDIENTS OF THIRIPHALA CHOORANAM (TPC)

S.no	Botanical name/ tamil name	Family	Parts to be used	Important Alkaloids	Actions
1.	<i>Terminalia chebula</i> , Retz KADUKKAI THOOL	Combretaceae	Fruit	Phenolic Compounds, punicalagin, Terflavin-A, Terchebulin	Hypolipidaemia, anti obesity ^{40,41,42}
2.	<i>Phyllanthus emblica</i> , Linn NELLIVATRAL	Euphorbiaceae	Fruit	Trigalloyl glucose, phyllembic acid, terachebin	Rich source of Vit c, antioxidant, hypolipidaemia ⁴³
3.	<i>Terminalia bellerica</i> , Gaertn THAANDRIKKAI THOOL	Combretaceae	Fruit	Belleric acid, Bellericoside, Gallic acid, ellagic acid, Mannitol	Hypolipidaemia ^{44,45,46}

***Terminalia bellerica*:**

B. Ahirwar., et al.,⁴⁵ has reviewed that Ethanolic extract of 200 - 250g in male adult albino rats for 45 days orally which showed marked reduction in lipoproteins.

***Terminalia arjuna*:**

Saravana Subramanian., et al.,⁴⁸ has reviewed that Ethanol, Diethyl ether, Ethyl acetate extract 175 - 350mg/kg BW in Female albino mice, Male wistar rats has shown that it reduces hyperlipidaemia and anti oxidant.

***Grossypium herbaceum*:**

C. Velmurugan., et al.,⁴⁹ has reviewed that ethanol extract of leaves (200mg / kg BW) has significantly reduced lipids levels in diabetes.

***Curcuma Longa*:**

P. Suresh Babu et al.,⁵² has reviewed that Aqueous and Alcoholic extract 100mg / kg/ bw one in 9 days for 2 weeks significantly reduces lipid levels in diabetes.

***Cosicinium fenestratum*:**

Shanmugam Manoharan, et al.,⁵⁴ has reviewed that Ethanolic extract of 300 mg / kg BW for 45 days / alloxan induced in wistar rats showed anti hyperlipidaemic and anti glycemic activity.

***Salacia reticulate*:**

Yoshikawa, et al.,⁵⁶ has reviewed those polyphenel constituents with lipase inhibitory and lipolytic activities in rats.

TABLE 4: INGREDIENTS OF MARUTHAM PATTAI CHOORANAM

S.no	Botanical name/ tamil name	Family	Parts to be used	Alkaloids	Actions
1.	<i>Terminalia arjuna</i> Wight & Arn MARUDU	Combretaceae	Bark	Triterpene arjunolitin, Arjunolone, arjunetin, arjunic acid, Terminolitin	Cardiac disease, Hypolipidaemia ^{47,48}

TABLE 5: INGREDIENTS OF KARIVEPPILAI CHOORANAM

S.no	Botanical name/ tamil name	Family	Parts to be used	Important Alkaloids	Actions
1.	<i>Murraya koenigii</i> (L) KARIVEPPILAI	Rutaceae	leaf	Girin, isomahanibine, Koenimbine, Koengicine	Anti- diabetic Hypolipidemia, Anti Oxidant ^{26, 27}
2.	<i>Gossypium herbaceum</i> (L) PARUTTI	Malvaceae	Seed	Gossypin	Anti diabetic Hypolipidemia ^{49, 52}

3.	<i>Curcuma longa</i> (L) MANJAL	Zingiberaceae	Rhizome	Curcumin, β - turmenone, Demthoxy Curamin	Anti-oxidant Hypolipidemia ^{51,52, 53}
4.	<i>Coscinium fenestratum</i> (Gaertn) MARAMANJAL	Menzspermaceae	Rhizome	Berlambine, Oxo- berberine, Stigmasterol	Anti-Oxidant Hypolipidemia ⁵⁴
5.	<i>Terminalia chebula</i> KADUKKAI	Combretaceae	Pericarp	Pinicalagin, Terflavin.A, Terchebuin	Anti-diabetic Hypolipidemia Cardio protective ^{40,41,42}
6.	<i>Terminalia bellirica</i> THANDRIKKAI	Combretaceae	Pericarp	Belleric acid, Mannitol	Hypolipidemia ^{44,45,46}
7.	<i>Emblica officinalis</i> NELLI MULLI	Euphorbiaceae	Pericarp	Trigalloyl glucose, Phyllembic acid	Antioxidant, Hypolipidaemia ⁴⁶
8.	<i>Salacia reticulata</i> KADAL AZHINGIL	Celestraceae	Root	Quinonemethide, Lenmbachol C, D, Pristimeria	Antidiabetic, Lipase inhibitory, Lipolytic ^{55, 56}

TABLE 6: LIST OF HYPOLIPIDAEMIC ACTIVITY OF SIDDHA MEDICINAL PLANTS

S.no	Botanical name/ Tamil name	Family	Part to be used	Important Alkaloids
1.	<i>Curcuma amada</i> (Roxb) MANGAI INJI	Zingiberaceae	Rhizome	β ,d, Curcumene, d- Pinene, d- Camphor, β - d - curcumene, Phytosterol ⁵⁷
3.	<i>Myristica fragrans</i> . (Houtt) SAATHIKAI	Myresticeae	Fruit	Eugenol, Isoleugenol, Methy Eugenol, Myristicin, Trimyriticin ⁵⁸
4.	<i>Pueraria tuberosa</i> (Dc) NILA POOSANI	Papilionaceae	Root	Puerarin, Tuberostan cardiac glycoside ⁵⁹
5.	<i>Tinospora cordifolia</i> (Wild) SEENTHIL	Menispermaceae	Stem	Tino Cardifolin, Tinosporidine, Tino sporaside, Cleodane ⁶⁰
6.	<i>Caesalpinia bonduc</i> (L) (Roxb) Kalichikai	Caesalpiniaceae	Fruit, leaf	α , β and δ Caesalpins, Pentacyclic, Tri Terpenoid (+) Ononitd, Cephane ⁶¹
8.	<i>Allium cepa</i> (Linn) VENGAYAM	Alliaceae	Bulb	Quercetin, Cycloallin, Phenolic Acid ⁶²
9.	<i>Allium sativum</i> (Linn) VALLAI - PUNDU	Alliaceae	Bulb	Ajuene, Y - glutamyl peptides, F- gitonin, degalactotigonin ⁶³
10.	<i>Capparis decidua</i> (Forsk) SENGAM	Capparaceae	Bark, fruit	Capparilline, N- triacontanol, Capparisinine ⁶⁴
11.	<i>Cinnamomum Verum</i> (J.S.Presl) LAVANGA PATTAI	Lauraceae	Leaf	Eugenol, Lianlool, Benzaldehyde, Benzyl Cinnamaldehyde, Pinene, Cymene ⁶⁵
12.	<i>Commiphora mukul</i> (Enzler) KUNGILYUM	Burseraceae	Resin	Resins, Z-Guggulusterone, E- Guggulusterone, Guggulsterol I-V ⁶⁶
13.	<i>Enicostema axillare</i> (Lam) VELLARUGU	Gentianaceae	Whole plant	Swertiamarin, Gentiamine, Swertisin 5-O, Glycosyle swertisin, Myristic acid ⁶⁷
14.	<i>Garcinia combogia</i> (G) KODUAM PUZHI	Clusiaceae	Fruit	(-) Hydroxycitric Acid, Tartaric acid. ⁶⁸
16.	<i>Saussurea lappa</i> (C.B.Clake) KOSTUM	Asteraceae	Root	Costunolide, dehydrocostus, Lactone ⁶⁹
17.	<i>Trigonella foenum & T. Gracenum</i> (Linn) VENDHAYAM	Papilionaceae	Seed	Graecunnins A-G, Trigofenosides, Gitogenin, C- trigonelline, Quercetin ⁷⁰
18.	<i>Coccinia grandis</i> (L) Voigt KOVAI	Curcubitaceae	Leaves, fruit, root	Lupeol, cucurbitacin, β -sitosterol, β -carotene, Linoleic, Oleic acids ⁷¹
19.	<i>Aloe vera</i> (L), Burm.F CHOTTU KATHALAI	Liliaceae	Leaves	Aloesone, aloesin, barbuloin, glycoside, β - barbaloin, iso-emodin ⁷²
20.	<i>Alpinia officinarum</i> (SW) PERARATHAI	Zingiberaceae	Rhizome	Methyl cinnamate, cineol, caryophyllenol I, II ⁷³

22.	<i>Embelia Ribbes</i> , Burm. F VAIVILANGAM	Myrsinaceae	Fruit, seed	Embelin, Quercitol, vilangin, christembine ⁷⁴
23.	<i>Ocimum sanctum</i> (Linn) O. emeri canum (L) NALLA THULASI	Lamiaceae	Leaves	Stigmasterol, Volatile oil, Caryophyllene ⁷⁵
24.	<i>Medicago sativa</i> KUTHIVAI MAZAL	Leguminaceae	Leaves	Saponins, Phytosterols, Vitamins, Coumarins ⁷⁶
25.	<i>Ougenia oojensis</i> (Roxb) NARIVENGAIAM	Papilionaceae	Bark	Lupeol, Betulin, Homoferreirin, Ougenin ⁷⁷
27.	<i>Sesbania grandiflora</i> (Poir) AGATI	Papilionaceae	Leaves, flower, root	Oleanolic acid, Glucuronic acid ⁷⁸
28.	<i>Pterocarpus marsupium</i> (Roxb) VENGAI	Papilionaceae	Leaves, stem	Liquiritigenin, garbauzol, Glucosides, Aurane ⁷⁹
29.	<i>Achyranthes aspera</i> , (Linn) NAIYURUVI	Amaranthaceae	Leaves	Oleanolic acid, ecdysone, ecdysterone, achyranthine ⁸⁰
31.	<i>Linum usitatissimum</i> (Linn) ALISI VIRAI	Linaceae	Seed	Phenylpropanoid, glucoside, Linnsitamarin, linseed oil ⁸¹
33.	<i>Capparis dedicua</i> , Forsk, Edgew SENGAM	Capparaceae	Fruit, bark	Capparine, Capparilline, n-petacosane, β – sitosterol ⁸²
34.	<i>Aconitum heterophyllum</i> (Wall) ADHIVIDAIYAM	Renunculaceae	Leaves, stem	3-0- β -D- glucopyranoside 7 - 0 β - D- glucopyranosyl - (1 \rightarrow 3), quercetin 3 - 0 - β -D glucopyranoside - 7 - 0- (6E - Cafferyl) β -D (1 \rightarrow 3) ⁸³
35.	<i>Dalbergia latifolia</i> (Roxb) ITTI	Papilionaceae	Bark	Methyl dalbergin, rotenoid, dalbinol ⁸⁴
37.	<i>Hibiscus Cannabinus</i> (Linn) PULICHHAI	Malvaceae	Leaves, seed	Isoquercitrin, Cannabiscitrin, myricetin, phosphonolipids ⁸⁵
38.	<i>Eclipta prostrata</i> (L) Mant KARISALAI	Asteraceae	Whole plant	Terthienyl aldehyde ecliptal, Nicotine, Triterpenoid ⁸⁶
39.	<i>Moringa oleifera</i> (Lam) KAATU MURANGAI	Moringaceae	Leaves, flower	Quercetin, 3-Rhamnoglucoside, Kaempferol ⁸⁷
40.	<i>Elaeocarpus sphaericus</i> NATTU RUTHRATCHAM	Elaeocarpaceae	Fruit, leaves	(-) - Isoleae, Carpilline, Rudrakine, Fixed oil ⁸⁸
41.	<i>Momordica charantia</i> (Linn) REVAKAI	Cucurbitaceae	Fruit, leaf	acylglucosyl sterols ⁸⁹
43.	<i>Nardostachys jatamansi</i> (Dc) JATAMANJI	Valerianaceae	Rhizome	Jatamansone, Sequitapene, Angelicin, Jatamansin ⁹⁰
44.	<i>Coriandrum sativum</i> (Linn) KOTHAMALLI	Apiaceae	Leaves, seed	Furoisocumarins, Coriandrin, Coriandrol, D.Mannitol ⁹¹
45.	<i>Syzygium cumini</i> (L), S. <i>Jambolanum</i> (Lam) C NAVAL	Myrtaceae	Fruits, seed	Myrecetin - 3 - 0 glucoside, Robinoside, Isoquercetin, Anthocyanins ⁹²
46.	<i>Nigella sativa</i> (Linn) KARUNCHEERAGAM	Ranunculaceae	Seed	Dithymoquinone, Nigilline, Aarmacenine, Trans anetholic ⁹³
47.	<i>Semecarpus anacardium</i> Linn SHENKOTTAI	Anacardiaceae	Fruit	Bhilwanol, Binaringenin, Semecarpetin, Gallufavanone ⁹⁴
48.	<i>Acorus calamus</i> . Linn VASAMBU	Araceae	Rhizome	Callaminone, Isocalamendiol, Asarone - 3, β -asarone ⁹⁵
49.	<i>Cynodon dactylon</i> (L) Pers ARUGAMPULLU	Poaceae	Whole plant	Apigenin, Luceolin, Orientin, Vitexin ⁹⁶
50.	<i>Sesbania grandiflora</i> (L) Poir AGATI	Papilionaceae	Leaves, flower	Oleanolic acid, Galactose, Glucuronic acid, Cyanidin 3 – glucoside ⁹⁷
51.	<i>Luffa cylindrica</i> (L) M.Roem <i>L.Aegyptiaca</i> Mill. PIRKANKAI	Cucurbitaceae	Tender fruits	Saponins Lucyosides A, H, Cucurbitacin B ⁹⁸

52.	<i>Bauhinia purpureae</i> . (Linn) MANDARI	Caesalpiniaceae	Flower	Tannin, Isoquercitrin, Quercitrin, Chalcone glycoside ⁹⁹
55.	<i>Scorpioides dulcis</i> . (Linn) SARKARAI VEMBU	Scrophulariaceae	Whole plant	Scoparic acid, Scopadulciol, Amellin, Root, Mannitol ¹⁰⁰
56.	<i>Hibiscus sabdariffa</i> (Linn) PULICHAI KEERAI	Malvaceae	Leaves, seed	Citric acid, d - malic, Hibiscus acid, Gossypetin, Sabdaritrin ¹⁰¹
57.	<i>Tribulus terrestris</i> NEERUNJIL	Zygophyllaceae	Whole plant	Diosgenin, giogenin, Chlorogenin, Tribuloside ¹⁰²
58.	<i>Macrotyloma uniflorum</i> KOLLU	Fabaceae	Seeds	Dolichin A and B ¹⁰³
59.	<i>Mukia maderaspatana</i> (Linn) MUSUMUSUKKAI	Curcubitaceae	Leaves, Root	Umbelliferone, Xanthotoxol, Isopimpinellin, Scopoletin ¹⁰⁴
60.	<i>Aegle marmelos</i> , (Linn). Corr VILVAM	Rutaceae	Whole plant	Psoralen, Xantho toxin, Scopoletin, Siummianine ¹⁰⁵
61.	<i>Tragia involucrata</i> Linn. CHENTHATTI	Euphorbiaceae	Root	Ethylbenzene, Limonene, Tragia Sapomine, ¹⁰⁶
62.	<i>Gymnema sylvestre</i> (Retz) P.Br.Ex. Schult SHIRUKURINJAN	Apocynaceae	Leaves	Gymnemasins A. D., Gymnimagenine, Gymnemic acid, Gypenosides II, Nonacosane. ^{107, 108}
63.	<i>Spermacoce hispida</i> (Linn) NATHAI CHURI	Rubiaceae	Seed	Isorhamnetin ¹⁰⁹
64.	<i>Ziziphus jujuba</i> (L) Gaertn ILLANTHAI	Rhamnaceae	Leaves	Jujuboside A, B., Jubanine A, B., Rutin, Ziziphin, Stephanine, Vit C

DISCUSSION AND CONCLUSION: Based on the Siddha literature and various review of the articles (**Table 6**), Siddha medicinal plants have been proven as beneficial role in the management of dyslipidaemia. It is also observed that *Madhumega Chooranam* is very much effective in Type 2 diabetes with hyperlipidaemia and *Venthamarai Chooranam* in hypertension with hyperlipidaemia. It is evident that from the above mentioned reviews, single herbal formulations have the same potency as that of the polyherbal formulations in controlling the hyperlipidaemic state.

REFERENCES:

1. National Cholesterol Education Program (NCEP) Expert panel: A high blood cholesterol in Adult (ATP III) Third report of the National Cholesterol Education Program. Final Report Circulation 2002; 106: 3143-3421.
2. The National Heart, Lung and Blood Institute Research Portfolio online Reporting Tools. www.nhlbi.nih.gov
3. C.S. Uthamarayan, Pharmacopeia of Hospital of Indian Medicine, Siddha Additional Pharmacopeia. 1956;47: 16-17
4. K.S. Murugesu Mudhaliar, Gunapadam Part I, Siddha Materia Medica 2006; 734 - 736.
5. Sarabendrar Vaithya Ratnavali, Tanjore Tamil University, Page No: 57.
6. Sawant.AM. Shetty. D., Mankeshwar.R., Asharvaid.TF., Prevalence of dyslipidemia in Young Adult Indian Population. Journal Association Physicians India 2008; 56; 99 - 102.
7. SK.Verma, Vartika Jain.et al., Blood Pressure Lowering, Fibronolysis enhancing and antioxidant activities of cardamom, 2009; IJBB. Vol 46, Page No: 503 - 506.
8. Chandhari.HS., Bhandari.U., Khanna. G., Preventive effect of exbelin from cardomomum on lipid metabolism and oxidative stress in high fat diet induced obesity in rats, plant Med 2012; 78 (7); 651 - 657.
9. Nair.S., Nagar.R., Gupta.R., Antioxidant phenolics and flavanoids in common Indian foods. J. Assoc Phys India 1998 ; 46; 708-710
10. Bhandari.U. Kanojia.R., Pillai.KK., Effect of Ethanolic Extract of Zingiber officinalis dyslipidemic in diabetic rats. Journal Ethnopharmacology 2005; 97(2); 227 - 230.
11. Al-Noory. AS., Amreen. A.N., Hymoor. S., Antihyperlipidemia effects of ginger extracts in alloxan-induced diabetes, and propylthiouracil oil-induced hypothyroidism in rats pharmacognosy Res 2013; 5(3); 157 - 161
12. Dhanley.JN. Raman.PH., Mugundar.AM. et al., inhibition of lipid peroxidative by piperine during experimental inflammation in rats. Indian J Exp Biol 1993; 31; 443-445.
13. Jin. Z., Borjihan. G., Zhaor Sunz., Antihyperlipidemic compounds from the fruit of Piper longum L. Pub. Med 2009; Aug 23(8): 1194-6.
14. Shaik Asdul Nabi, Ramesh Babu Kasetti, Swapna Sirasanagandla. et al., Anti diabetic and Anti hyperlipidemic activity of piper longum root aqueous extract in STZ, induced diabetic rats. BMC complementary and Alternative medicine, 2013; 13: 37
15. Santhosh kumar Maurya, Kenoval Raj..et al., Anti hyperlipidemic phytomedicine 2011. June 15; 18 (8-9) 795 - 801.
16. Birari.RB. Gupta.S., Mohan.CG. et al., Antiobesity and lipid lowering effects, A Glycyrrhiza Chalconesi experimental and corruptationary studies phytomedicine 2011. June 15; 18 (8-9) 795 - 801.

17. Iyer. D., Patil.UK. Effect of Choloroform and Aqueous basic fraction of Ethanolic extract from *A. Gravcolens* L, Uri experimentally - induced hyperlipidemia in Rats. *J. Complement integr, Med* 2011; doi : 10,2202/1553-3840-1529
18. Vaibhav Srivastara, Subodh dubey.et al., studies on hypolipidemia activity of seeds of *cuminum cyminum* (Linn) Indo - american Journal of Pharmaceutical Research, 04 November 2010.
19. Rohit Srivastara, Savayam Prakash.et al., Antidiabetic and antidiyslipedemic activities *Cuminum cyminum* in validated animal models - medicinal chemistry Research. 15th October - 2010.
20. Umishi .E. Yarnada. K., Yamad. T., Kaji. K., et al., Comparative effects crude drugs on serum lipids item pharm Bull (Tokyo) 1984; 32 646 - 650
21. Subasini.U., Thenmozhi.S.,Venkateswaran.V., et al., Phyto-chemical analysis; A anti hyper lipidemic activity of *Nelumbo nucifera* in male wistar rats. Internal tional Journal of Pharmacy Teaching of practices 2014; 5 (1): 935
22. Durairaj.B., Doral.A., Free radical Scarvenging potential of *Nelumbo nucifera* flowers (White and pink) International Journal of Natural Sciences Research 2014; 2(8); 133-146
23. Redd VRC. Rainana Kumari .SV. Reddy. B.M., et al., Cardiotonic activity of the fruits of *Terminalia Chebula* Fitoterapia 1990;61 517-525
24. Maruthappan.V.K., Sakthi Shree, Hypolipidemic activity A Haritaki in atherogenic diet induced hyperlypidemic rats. *J. Adv Pharm tech Res* 1: 229 - 35.
25. Tachibana.Y., Kikuzaki.H., et al., Antioxidative activity of carbazole from *Murraya koenigi* leaves. *J Agric food Chem* 2001; 49; 5589-5594
26. Khan.BA., Abrahama, Leelamm.S., Hypoglycemic action of *Murraya koenigii*:Indian *J.Biochem Biophs*; 1995; 32; 106 - 108
27. Usharani.P., Fatima.N., Muralidhar.N., effects of *phyllanthus Emblica* extract on endothelial dysfunction and biomarkers of oxidative stress in patients with Type II DM, randomized, double blind controlled studies. *Diabetes Met syndrobes* 2013; 6; 275 - 284.
28. Anila.L., Vijayalakshmi.NR., (2002), Flavanoids from *Emblica officinalis* and *mangifera indica* effectiveness for dyslipidemia. *Journal of Ethnopharmacology* 79 (1); 81 - 89.
29. Arunab Bhattacharya, Abhijit Chatterjee et al., Antioxidant activity of active tannoid principles of *Emblica officinalis*. *Indian Journal of experimental Biology*, Vol.37, July 1999; 676-680.
30. Dhingra Dinesh, Jindal vaneetha, Sharma Sunil .et al., Evaluation of anti-obesity activity of *Tinospora cordiflora* in rats. *International Journal of Research in Ayurvedha and Pharmacy* 2011; 21; 306-311.
31. Veena sharma, Rekha Gupta, Shatruhan Sharma, Hypolipidemic activity of *Tinospora cordiflora* root extract on aflatoxin, β induced toxicity in mice, *Int. J. Res Pharma, Sci* 2(4); 2011; 585-589.
32. Mallick.C., Mariti.R., Gosh.D., Comparative Study on anti hyperglycemic and anti hyper lipidemic effects of separte and composite extract of seed, *Eveginia Jambolans* and roots of *Musa paradisica* (or) streptozocin induced diabetic male albino rats. *J Phar Ther* 2006 ; 5(1) 27 - 33
33. Sharma.SB. Tamuar. RS., Nasir. A., et al., Antihyperlipidemic effect a active principle isolated from seed of *Evaginia jambolana* and alloxan induced diabetic rabbits . *J.Med food* 2011; 14(4); 353-359
34. Nagulendran. KR., Mahesh. R., Begum. VH., Preventive role of *Cyperus rotundus* rhizomes extract on age associated changes in glucose and lipids pharmacology online, 2007;2; 318 - 325
35. Fried Wald. W.T., Levy. R., et al., Estimation of Concentration of LDL-Cholesterol in Plasma without use a preparative Ultra centrifuge clinical chemistry 1972; 18; 499 - 502
36. Bambhole. V.D., Effect of Some medicinea plant preparation adipose tissue metabolism, *Ancient sci Life* 1988; 8, 117 - 124
37. Akkhana. F., Rizvi Lipid Lowering activity of *Phyllanthus niruri* in hyperlipidemic rats; 2002; *Journal of Ethropharmacology*.
38. R.Harish.T., Shivanandappa.A., Antioxidant activity and hepato protective potential of *phyllanthus niruri*.
39. Redd. URC. Rainana Kumari S.V., Reddy B.M. et al., cardio protective activity of the fruits of the *Terminalia chebula*, *Fitoterpia* 1990; 61; 517-525.
40. Maruthapan V.K., Sakthi Shree. Hypolipidaemic activity a Haritaki in atherogenic diet induced hyperlipidemic rats, *J. Adv Phar Tech Research*, Volume I, 229 - 35.
41. Priya F. Velpandian et al. Hypolipidaemic activity of Kadukai Chooranam in Triton WR - 1339 induced Hyperlipidemia.
42. Yokozawa. T., et al., Amla prevents dyslipidemic and oxidative stress in the ageing process *Br. J. Mutr Jun*; 97 (6): 1187 - 95
43. Patel .S.S., Goyal R. K., Cardio protective effects of gallic acid in induced myocardial dysfunction in rats, *Pharmacognosy Res* 2011; Oct 3(4) 239-40
44. B. Ahirwar, A.K. Singhai., et al., Effects of *Terminania bellerica* fruits on lipid profiles of rats, *Journals of Natural remedies*, 2003; Vol 31, 31 - 35
45. Shaila. H.P., Udupa.S.L., Udupa. A.L., Hypolipidemic activity of three indogenous drugs in experimentally induced atherosclerosis, *International journal of Cardiology*, 1998; 67 (2); 119 - 124.
46. Patil.R.H., Prakash.K., V.L. Maheshwari.V.L., Hypolipidemic effect of *Terminalia arjuna* (L) in experimeintally induced Hypercholestremic rats. *Acta Bio Sz eged* 55 (2) ; 2011; 289 - 293
47. Saravanan Subiramanian, Anti Hyperlipidaemic and antioxidant potential of different Fraction of *Terminalia arjuna*. bark agains px. 407 induced hyperlipidaemic, *Indian Journal of experimental Biology* ; Vol . 49 April 2011, Page No: 282 - 288
48. C. Velmurugan., Anurag Bhargara et al., Anti Hyperglycemic and Hypolipidemic activity of *Gossypium herbaceum* by dexamethasone induced diabetic rat *Int. J. of Research in Pharmaceutical and Nano Science* 2013; 2(2) 169 - 170
49. C. Velmurugan., A Bharagana., Anti diabetic activity of *Gossypium herbaceum* by alloxan induced model in Rats, *Journal of Pharmatutors* 2014; Vol. 2 pp. 126 - 132
50. P.Suresh Babu., K. Srinivasan., Hypolipidemic action of curcumin active principle of turmeric in STZ induced diabetic rats *Journal a molecular and cellular Biochemistry* 1997; Vol . 166 - Vol I; Page No: 169 - 174.
51. Sarah Nuzooluwatosin, Adarange, oral administration of extract *curcuma longa* blood glucose in alloxan induced Hyperlipidaemia in diabetic rabbits.
52. Kempaiah.RK., Srinivasan.K., Influence of dietary cucumin, Capaicin and garlic on the anti oxidant status of RBS and the liver in HFD in rats. *Animals and nutrition and metabolism* 48 (5); 314 - 320.

53. Shanmugam Manoharan.S., Umadevi et al., Anti hyperglycemic effect of coscinium fenestratum and Catharanthus roseus in Alloxan induced diabetic rats Int. J. of Nutrition Pharmacology, Neurological disease Vol. V, Issue II, 2011;Page No: 189 - 193
54. Kirshino.E., Itol et al., A mixture of the salacia reticulata, (Kotala himbutu) Aqueous extract and Cyclodextrin reduces the accumulation of visceral fat mass in mice and rats with HFD - induced obesity, J. Nutr in;136 (2); 2006;433-9;
55. Yoshikawa.M.,Shimodel.H.,et al.,Polyphenolic constituent with lipase inhibitory and lipolytic activities have mild antiobesity effect in rats . J. Nutr 13; 2002; 1819, 24.
56. Babu.P.S., Srinivasan K., Hypolipidaemia action of curcumin. The active principle of turmeric (Curcumin) in STZ miduced diabetic rats. Molecular Bio chemistry 1997; Jan 166 (1-2); 169 - 175.
57. R.Dhanalakshmi., K. Balamurugan et al., Hypolipidaemia and anti atherosclerotic activity of some medical plants. A Review world Journal of Pharmacy and Parmaceutical sciences ; Vol III issue; 328 - 340 ; 2014
58. Verma S.K., Jain .V. Uyas.A., Fibrolysis enhancement by pueraria tuberosa in patients with CAD; A placeso controlled study. Journal of Cell tissue Research 2009;Vol 9(1), 1649 - 1654
59. Veen a sharma, Rekha gupta, Shatruhan sharma, Hypolipidaemia actiity of Tinospora cordifolia root extract on an aflatoxin β - induced toxicity in mice Int. J. Res Pharma Sciences; 2011; 2 (4) ; 585 - 589,
60. Sharma. S.R., Divivedi.SR. Swarup.D., Hypoglycaemic and Hypolipidaemia activities of Caesalpinia bunduc seeds in rats. Journal of Ethnopharmacology 1997; Sep ; 58 (1); 39 -44
61. Singh Ayodhya, Singh Kusum etal., Hypoglycemic activity of different extracts of various herbal plants IJRAPI (1), 2014;212 - 224,.
62. Atul Shrivastava, Upma chatured et al., A mechanism based pharmacological evaluation of efficacy of Allium sativum in regulation of dyslipidaemia and oxidative stress in hyperlipidaemic rats, Asian Journal of Pharmaceutical and clinical Research, Vol III 2010; issue 2 123 - 126
63. Neelkamal chabla., Evaluation of Hypolipidaemic activity of capparis decidua in Ethanolic extracts IJBS, 3/2009; 5(1); 70 - 73
64. Gaber, E., El oesoky etal., Anti diabetic and Hypolipidaemic activity of cinnamom miallanan - diabetic rats, Journal of medicinal plants Research Vol.VI (9); 9th March 2012; PP1685 - 1691.
65. Comparative study of Hypolipidaemia profile of resinoids of commiphora mukul / commiphora weight, from different geographical locations - Int. J. a pharmaceutical sciences. Volume 74 5; 2010; pp 422 – 427.
66. Rhopal, Gnammani et al., Enicostema Litorale, A potential Hypolipidemic plant, Natural product Radiance Vol III (6) Nov. December 2004; 401 - 403
67. Subhasini.N., Nagarajan.G., et al., Invitro antioxidant and Anti cholinestrase activities of Garcinia Combogia, Inter Jrl of Pharmacy and Pharmaceutical Science, Vol III, 2011; issue 3, 129 - 132.
68. Anbuy. J., Ashwini Anjana K., et al., Evaluation of Anti hyperlipidaemic activity of Ethanolic extract saussurea lappa in Rats. Int Journal of Pharma and Bio science Vol II / issue 4/ 2011; Oct - Dec 1550 - 555.
69. Prasanna.M., Hypolipidaemic effect a Fenugreek, a clinical study, Indian Journal of Pharmacology, 2000; 34 - 26,
70. Pekamwar S.S, Kalyankar T.M., et al., Pharmacological activity of coccinia grandis- Journal of applied pharmaceutical science Vol III (05) March 2013 ; PP 114 - 119 .
71. Kwanghee kirm Hyunyulkim et al., Hypoglycemic and Hypolipidaemic effects processed. Aloe vera gel to a mouse model NIDDM, Journal of Phytomedicine, 2009; 02 - 14.
72. Shin.J.E., Joo Han.M., Kim.DH., 3 -Methyl Lether - galarugin isolated from Alpinia officinarum Inhibits pancreatic lipase, Biological and Pharmaceutical Bulletin 26(2); 2003;854 - 857.
73. Chandhari.HS., Bhandari.U., Khanna.G., Preventive effect Embelin from Embelia ribes lipid metabolism and oxidative stress in HFD induced obesity in rats, Planta med 2012; 78 (7); 651-657.
74. Gupta.S., Mediratta P.K., et al., Anti diabetic, Anti hypocholesterolaemic and Antioxidant effect of Ocimum sanctum (Linn) Seed oil . Indian .Jl. exp piol. 44(4) ; 300-304; 2006.
75. Baxv D.B., Singh P.K., Medicago Sativa Leaf extract Supplementation corrects diabetes induced dyslipidemia, oxidative stress and hepatic, Renal functions, Annals of Biological Research I(3); 107 - 109 ; (2010).
76. Velmurugan.C.T., Sundram et al., Anti diabetic and hypolipidaemic activity of Bark of Ethanolic extract Eugenia Oojeinesis , Med J Malaysia, Vol.66(1); 2011; 22 - 26 ;
77. Saravanakumar, S. Vanitha, et al., Hypolipidaemic activity of Sesbania grandiflora in Triton Wr - 1339 induced hyper lipidaemic rats, Int. J.A. Pyto medicine 2(210) 52 - 58.
78. Maruthupandian.A and R.Mohan, Antidiabetic, Anti-hyperlipidaemic and antioxidant activity of pterocarpus, marsupium (Roxb) in alloxan induced diabetic rats. Int J. Pharm Tech Research, Vol III No.3 July - Sep 2011; 1681-1687
79. Venkatalakshmi.P., Vedhavalli.N., Sangeetha.S., Hypo-lipidaemic, Effect of Achyraonthe aspra on HFD induved atherogenic rats. Research Journal of Pharmaeuctical Biological and chemical Sciences. July - Sept, RJPBCS, Vol III, Issue 3. PP 75 - 83
80. Mani (or) Mani et al., An open label study on the effect of flax seed powder - supplementation in the management of DM, J Diet Supl . Sep 7 (3) 273 - 82; 2010
81. Neel kamal, Chablia, Preliminary study into the Hypolipidaemic activity of various pars capparis dedicuas, Ethanobotanical leaflets 13; 332 - 27 (2009)
82. Arun Koorapally, Subash Anu, Augustine, Hypolipidemic effect of Methanol fraction of Aconitum hetrophyllum wall. ex Royre and the mechanism of action in diet induced obese rats J. Adv. Pharm, Tech Res Vol VIII; (issue 4) Oct - Dec 2012.
83. Mohammad, Khalid.H.H., Siddique Lipid howering and hypoglycemic potential of dried Dalbergia Latifolia Roxb, Bark extract in spraque - Dwley rats induced HFD. Ind. Jr. of Naural products Research 1(4) 2011; 49 - 54.
84. Shivali.N., Mahadevan, et al., Anti Hypolipidaemic effect of Hydro alcholic extract of Hibiscus cannabinus (L) leaves in the HFD in rats . Annals of Biologicals Research 1 (3); 174 - 181 (2010)
85. R.Dhandapani, Hypolipidemia activity of Eclipta Prostrata (L) Leaf extract atherogenic diet induuced hyperlipidemic rats, Indian Journal of experimental Biology ; vol 45, 2007; 617 - 619
86. Pankaj.G.Jain., Savita.D., et al., Hypolipidaemic actiity of Moringa olifera Lam. Moringaceae on HFD induced

- Hyperlipidaemic in albino rats, Brazillan. Journal of Pharmacognosy 20 (6); 969-973 (2010).
87. Kalpana garg, Komal goswami et al., A Pharmacognostical review on *Elaeocarpus sphaericus* Int.N. Journal of Pharmacy and Pharmaceutical sciences 2013; vol.5 issue 1
88. Abd. El Sattar., El., Batan.S El Gengaini .SC et al., Toxicological studies of *Momordica charantia* on albino rats in normal and alloxan diabetic rats. Journal of Ethnopharmacology 2006; 108 (2) 236 - 242.
89. Subashini et al., Biochemical study on the protective potential of *Nardostachys jatamansi* extract on lipid profile and lipid metabolizing enzyme pharmazie 62 (5); 2007; 382-387.
90. Lal A.A., Kumar.I., et al., Hypolipidaemic effect of *Coriandrum sativum* in triton - induced hyperlipidaemic rats. Indian Journal of experimental Biology 42 (9); 2004; 909 - 912.
91. Ravi. K., Rajasekaran. S., Subiramanian .S, Anti hyperlipidaemic effect of *Eugenia jambolana* seed kernel on STZ induced diabetes in rats food and chemical toxicology 43 (9); 2005; 1433 - 1439.
92. Morikawa.I., F.Winomiya et al., Nigellamines A₃, A₄, A₅ and C . New dolabellane - type of peno alkaloids with lipid metabolism - promoting activities chemical and pharmaceutical bulletin 2004; 52(4); 494 - 497.
93. Tripathi Y. B and Pandery R.S. *Semecarpus anacardium* nuts inhibit lipopoly saccharide induced nitrus oxide production in rat, macrophages along with its hypolipidemic property. Ind. J. of experimental Biology 2004; 42(4); 432 - 432.
94. Parab.R.S. Mengi, Hypolipidemic actiity of *Acorus calamus* Linn rats. *Atoterpia* 73 (6) 451 - 455.
95. Sighn.SK. Kesari.AN., et al., assessment of antidiabetic potential of *cynodon dactylon* extract in STZ diabetic rats Journal of ethnopharmacology; 114 (2); 2002; 174 - 179.
96. Saravana Kumar.A., Vanitha. S., Ganesh. M., et al., Hypolipidaemic activity of *Sesbania grandiflora* intriton WR - 1339 induced, Hyper lipidaemic rats. In Jr. of Phytomedicine 2; 2010; 52 - 58.
97. Abdul Hameed Thayyil. M.K.M., Surulivel et al., Hypolipidaemic activity of *Luffa aegyptiaca* fruits. Int. J of Pharmaceutical application 2011; Vol.2 (issue) 81 - 88.
98. Lakshmi BVS. N., Neelima N., Kasthuri et al., Anti hyper lipidaemic activity of *Bauhinia purpurea* extracts in hyper cholestremic albino rats. Int. J. Pharm. Tech Res Vol- III, 2011; No.3; 126 - 1272.
99. Pari. L Latha. M., Anti hyperlipedaemic effect of *scorpiadulcis* in STZ induced diabetic rats. Journal of Medicinal food 9(1) 2006; 102 - 107.
100. Hirunpanich.V., Vtaipat.A., et al., Hypocholesterolemic and antioxidant effects of aqueous extracts from the dried calyx of *Hibiscus sabdorriffa* (L) in Hypercholestrenic the dried calyx of *Hibsus Sabdariffa* (L) in Hyper cholestrenic rats. Journal of Ethnopharmacology 103 (2) 252 - 260.
101. Hypoglycemic and Hypolipidaemic effects of Alcholoic extract of *Tribulus* in STZ induced diabetic rats in comparative study with *Tribulus terrestris*. Indian Jr. of experimental Biology 45 sep 2007 pp 785 - 770.
102. Sathikumar.G. Prashanth, Antihypercholesterol- -enic effect is *Macrotyloma uniflorum* (lam) Vende (Fabaceae) extract on HFD induced. Hyper Cholestremia in spraque - Dauley Fates, Journal of diet supply 10 (2); 2013; 116 - 128.
103. A Ja Petrus, Ethonobotanical and Pharmacological profile with propagation strategies of *Muka maderaspatana* (L) M. poem. A Corcise; Review Indian Journal of Natural products and Resources Vol IV (1) March 2013; pp 9,26.
104. Mani Satya . KL, Bairu... et al., In fluence of combination of crude extract of *Aegle marmelos* leaves and *Tamarinnus indica* seeds on sugar and lipid level in normal STZ induced Diabetic rats. Resh. Journl of Pharmaceutical Biological and Chemical Science. June 2013; Vol IV; issue 2 Page 107.
105. S. Mohamed Farook. Element Atlee, Antidiabetic and Hypolipidaemic potential of *Tragia involucrata* Linn. is STZ Nicotinamide, induced, Type II, Diabetic Rats, Int Jour of Pharmacy and Pharmaceutical Science, Vol III, Suppl 4 2011.
106. Mohan Reddy. et al., Therapeutic plants and their Anti obesity properties Global J. Res Md plants, Indigen. Medl Vol. II; Issue 9; Sept 2013; PP 648 - 655.
107. Shrivastara.R, Solankis.. et al., Comparative evaluation of polypherbal combination of hypolipidaemic activity 2009; 1(1); 9-12.
108. G.Sivaelango, P.Senthil Kumaran.. et al., Anti hyperlipidaemic activity of spermacoce hispida Ethonolic extract in Triton WR.1339 induced Hyperlipidaemic Rats. Journal, applied, Pharmaceutical Science 02 (02) 2012, 95 - 98.
109. Anbarasi, Balakrishnan, Brindha Pamalah, Hypo-glycemia and Hypolipidemic effects of *Ziziphus Jujuba*. Linn. In STZ induced diabetic rats. Res. Journal A pharmaceutical and chemical science, vol IV; Ap. June 2013. PP - 611.

How to cite this article:

Manoharan A, Manjula M, Mubarak H and Chitti Babu CV: Hypolipidaemic Activity of the Siddha Medicinal Plants. *Int J Pharmacognosy* 2015; 2(4): 145-54:doi link: [http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.2\(4\).145-54](http://dx.doi.org/10.13040/IJPSR.0975-8232.IJP.2(4).145-54).

This Journal licensed under a Creative Commons Attribution-Non-commercial-Share Alike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)