

Potential Mechanisms by Which Polyphenol-Rich Grapes Prevent Obesity-Mediated Inflammation and Metabolic Diseases

Chia-Chi Chuang and Michael K. McIntosh

Department of Nutrition, University of North Carolina at Greensboro, Greensboro, North Carolina 27402; email: c_chuan2@uncg.edu, mkmcinto@uncg.edu

Annu. Rev. Nutr. 2011.31:155–76

First published online as a Review in Advance on May 6, 2011

The *Annual Review of Nutrition* is online at nutr.annualreviews.org

This article's doi:
10.1146/annurev-nutr-072610-145149

Copyright © 2011 by Annual Reviews.
All rights reserved

0199-9885/11/0821-0155\$20.00

Keywords

phenolic phytochemicals, flavonols, anthocyanins, antioxidant, anti-inflammatory, insulin resistance

Abstract

Obesity and metabolic disease-related health problems (e.g., type 2 diabetes, atherosclerosis, and hypertension) are the most prevalent nutrition-related issues in the United States. An emerging feature of obesity and type 2 diabetes is their linkage with chronic inflammation that begins in white adipose tissue and eventually becomes systemic. One potential strategy to reduce inflammation and insulin resistance is consumption of polyphenol-rich foods like grapes or their by-products, which have anti-inflammatory properties. Polyphenols commonly found in grape products have been reported to reduce inflammation by (*a*) acting as an antioxidant or increasing antioxidant gene or protein expression, (*b*) attenuating endoplasmic reticulum stress signaling, (*c*) blocking proinflammatory cytokines or endotoxin-mediated kinases and transcription factors involved in metabolic disease, (*d*) suppressing inflammatory- or inducing metabolic-gene expression via increasing histone deacetylase activity, or (*e*) activating transcription factors that antagonize chronic inflammation. Thus, polyphenol-rich grape products may reduce obesity-mediated chronic inflammation by multiple mechanisms, thereby preventing metabolic diseases.

Contents

OBESITY, INFLAMMATION, AND METABOLIC DISEASES	156
Factors Influencing White Adipose Tissue Growth, Development, and Overexpansion that Lead to Obesity	156
Metabolic Consequences of Expanding White Adipose Tissue and Its Impact on Liver, Muscle, and Pancreas	157
Proinflammatory Role of Free Fatty Acids and Adipokines	158
TYPES AND ABUNDANCE OF PHENOLIC PHYTOCHEMICALS IN GRAPES	159
Types of Phenolic Phytochemicals in Grapes and Grape Products	159
Relatively High Abundance of Anthocyanidins/ Anthocyanins and Flavonols in Grapes	160
MECHANISMS BY WHICH ANTHOCYANIDINS/ ANTHOCYANINS AND FLAVONOLS, MAJOR PHENOLIC	

PHYTOCHEMICALS IN GRAPES, REDUCE CHRONIC INFLAMMATION	160
Acting as an Antioxidant or Increasing Antioxidant Gene or Protein Expression	160
Attenuating Endoplasmic Reticulum Stress Signaling	163
Blocking Pro-Inflammatory Cytokines or Endotoxin-Mediated Kinases and Transcription Factors Involved in Metabolic Diseases	163
Suppressing Inflammatory- or Inducing Metabolic-Gene Expression via Increasing Histone Deacetylase Activity	165
Activating Transcription Factors that Antagonize Chronic Inflammation	166
CONCLUSIONS AND IMPLICATIONS	167
Potential Health Benefits of Consuming Grapes or Grape Products	167
Potential Risks of Excess Consumption of Grapes or Grape Products	169
Future Research	169

OBESITY, INFLAMMATION, AND METABOLIC DISEASES

Factors Influencing White Adipose Tissue Growth, Development, and Overexpansion that Lead to Obesity

Obesity is a global health issue, with more than 500 million people classified as obese and 1.5 billion overweight, including 43 million

children under the age of five (<http://www.who.int/mediacentre/factsheets/fs311/en/index.html>). In the United States, obesity is rapidly increasing among all age groups. Currently, in all but two states at least 20% of the population is classified as obese (<http://www.cdc.gov/obesity/data/trends.html>). Diets high in calories, especially from sugars, saturated fatty acids (SFAs), and long-chain

omega-6 polyunsaturated fatty acids (PUFAs), and sedentary lifestyles lacking physical activity contribute significantly to this obesity epidemic. Endogenous and exogenous factors that enhance preadipocyte proliferation or adipocyte hypertrophy have the capacity to increase white adipose tissue (WAT) mass and the development of obesity (reviewed in 66, 116).

Metabolic Consequences of Expanding White Adipose Tissue and Its Impact on Liver, Muscle, and Pancreas

The rapid rise in obesity is accompanied by a similar increase in cardiovascular disease (CVD), hypertension, and insulin resistance or type 2 diabetes (<http://www.who.int/mediacentre/factsheets/fs311/en/index.html>). For instance, ~80% of people with type 2 diabetes are overweight or obese (13), suggesting a strong positive relationship between the two diseases (<http://www.cdc.gov/nccdphp/dnpa/obesity/consequences.htm>). Obese patients with type 2 diabetes have elevated levels of tumor necrosis factor alpha (TNF α) in their blood (133), WAT (50), and muscle (97). Furthermore, impaired glucose disposal is positively correlated to

TNF α expression (97, 104, 133). Metabolic endotoxemia [e.g., elevated lipopolysaccharide (LPS) level in circulation] is also associated with obesity (14).

This cluster of obesity-related, metabolic diseases is known as the metabolic syndrome. One emerging feature of the metabolic syndrome is its linkage with chronic inflammation in WAT that becomes systemic. It is characterized by engorged adipocyte death (107), increased cytokine/chemokine production and inflammatory signaling, and recruitment of leukocytes (reviewed in 48). Thus, excess WAT is an overactive endocrine organ secreting an array of inflammatory adipokines that contribute to the metabolic syndrome (reviewed in 44). Furthermore, as WAT mass increases, its lipid- and glucose-buffering capacity decreases. This results in elevated blood levels of free fatty acids (FFAs), very-low-density lipoprotein (VLDL) (hyperlipidemia), glucose (hyperglycemia), and insulin (hyperinsulinemia), as well as ectopic lipid accumulation in skeletal and cardiac muscle, liver (steatosis), and pancreas (**Figure 1**). Understanding the mediators of this inflammation and their mechanisms of action is essential in order to develop effective strategies to prevent chronic inflammatory signaling from WAT.

WAT: white adipose tissue
TNF: tumor necrosis factor




Figure 1

Decreased lipid- and glucose-buffering capacity of adipocytes leads to hyperglycemia, hyperlipidemia, and hyperinsulinemia. Nutrient overload in white adipose tissue (WAT) leads to insulin resistance [IR; i.e., decreased free fatty acids (FFAs) and glucose (GLU) uptake] coupled with increased secretion of FFAs and proinflammatory adipokines such as tumor necrosis factor alpha (TNF α). Elevated FFAs are deposited ectopically in muscle, liver, and pancreas, leading to hyperglycemia, hyperlipidemia, and hyperinsulinemia, respectively. TG, triglyceride; PPAR γ , peroxisome proliferator-activated receptor gamma; VLDL, very-low-density lipoprotein.

ROS: reactive oxygen species

MAPKs: mitogen-activated protein kinases

ERK: extracellular signal-related kinase

JNK: c-Jun-NH₂-terminal kinase

AP: activating protein

PPAR: peroxisome proliferator-activated receptor

MCP: monocyte chemoattractant protein

MΦ: macrophages

Proinflammatory Role of Free Fatty Acids and Adipokines

Free fatty acids, especially SFAs, contribute significantly to chronic inflammation (reviewed in 58). For example, palmitate activates protein kinase C (PKC) signaling, generates ceramide and reactive oxygen species (ROS), and increases oxidative or endoplasmic reticulum (ER) stress, all of which are associated with chronic inflammation or insulin resistance. Dietary constituents like SFA may also alter gut microflora, leading to elevated blood levels of LPS or TNF α that cause endotoxemia (14).

Tumor necrosis factor alpha, LPS, SFA, or FFA instigate inflammation and insulin resistance by triggering ROS-mediated oxidative or ER stress and serine/threonine kinase phosphorylation signaling that active inflammatory mitogen-activated protein kinases (MAPKs),

including extracellular signal-related kinase (ERK) and c-Jun-NH₂-terminal kinase (JNK), I κ B α kinase (IKK), and transcription factors nuclear factor-kappa B (NF- κ B) and activating protein (AP)-1 (reviewed in 45). Collectively, these proteins induce inflammatory gene transcription, antagonize peroxisome proliferator-activated receptor gamma (PPAR γ) activity, or directly impair insulin receptor substrate (IRS)-1 signaling, leading to insulin resistance. Furthermore, engorged adipocytes release chemokines such as monocyte chemoattractant protein (MCP)-1 that attract monocytes and stimulate their recruitment and differentiation into macrophages (M Φ) via chemotaxis (reviewed in 77). Monocyte chemoattractant protein-1 has also been reported to convert alternatively activated M Φ (M2) into classically activated M Φ (M1), further increasing inflammation and insulin resistance (**Figure 2**). Such an inflammatory scenario in perivascular WAT can lead to plaque formation and smooth muscle migration and proliferation in the vascular endothelium and wall, thereby increasing the development of CVD (reviewed in 95).

Thus, a positive energy balance expands WAT, generating adipokines such as TNF α and MCP-1 as well as FFA that recruit and activate M Φ and activate inflammatory cascades. These responses antagonize PPAR γ activity, leading to decreased glucose and FA uptake and metabolism, causing ectopic lipid accumulation (lipodystrophy), hyperglycemia, and hyperlipidemia. Proposed mechanisms by which these inflammatory signals reduce PPAR γ activity include (a) decreased PPAR γ mRNA levels, (b) increased phosphorylation of serine residue 112 of PPAR γ by ERK, leading to PPAR γ ubiquitination and proteasome degradation, (c) decreased PPAR γ DNA binding, and (d) decreased PPAR γ transcriptional activity (**Figure 3**) (reviewed in 129). Discovery of dietary strategies, including the consumption of polyphenol-rich grapes or grape products, to reduce obesity-related chronic inflammation could potentially attenuate metabolic diseases.




Figure 2

Mechanisms by which tumor necrosis factor alpha (TNF α), lipopolysaccharide (LPS), saturated fatty acids (SFAs), and free fatty acids (FFAs) promote inflammation and insulin resistance. Elevated levels of TNF α , LPS, SFA, or FFA activate their cognate cell surface receptors or diffuse into the cell, thereby activating protein kinases such as protein kinase C (PKC) and mitogen-activated protein kinases (MAPKs) or enzymes such as nicotinamide adenine dinucleotide phosphate (NADPH) oxidase that produce reactive oxygen species (ROS), thereby triggering oxidative or endoplasmic reticulum (ER) stress signals or activating inflammatory transcription factors such as nuclear factor-kappa B (NF- κ B) and activating protein (AP)-1. Together, these inflammatory signals induce inflammatory gene expression, enhance leukocyte recruitment and differentiation including activating monocytes into classically activated M1-macrophages (M Φ), antagonize peroxisome proliferator-activated receptor gamma (PPAR γ) activity, and impair insulin receptor substrate (IRS)-1 signaling, leading to insulin resistance. MCP-1, monocyte chemoattractant protein-1; TLR, toll-like receptor; TNFR, TNF α receptor.

TYPES AND ABUNDANCE OF PHENOLIC PHYTOCHEMICALS IN GRAPES

Types of Phenolic Phytochemicals in Grapes and Grape Products

Grapes and their by-products are consumed worldwide. There are more than 50 varieties of seeded and seedless grapes in black, blue, blue-black, golden, red, green, purple, and white colors. Common grape products include table grapes, wine, raisins, juices, and preservatives. The average annual consumption of fresh grapes in the United States is about eight pounds per person (<http://www.ers.usda.gov/Data/FoodConsumption/>). Most of the grapes consumed in the United States are grown in California. Table grapes contain essential nutrients such as water, carbohydrates, proteins, fats, vitamins, minerals, and fiber (reviewed in 127) and nonessential compounds including phytochemicals. Phenolic phytochemicals in grapes (**Table 1**) possess biological activities that have been reported to promote health (reviewed in 125). Some of the early health benefits of consuming red wine (e.g., decreased the risk of CVD), referred to as the “French Paradox” (92, 103), have been attributed to its phenolic phytochemicals including resveratrol. Ironically, the resveratrol content of grapes and red wine is relatively low compared to other polyphenols (**Table 1**).

The predominant phenolic phytochemicals in grapes are flavonoids such as flavonols, flavan-3-ols (monomers, oligomers, or polymers), and anthocyanidins/anthocyanins, and to a much lesser extent nonflavonoids such as stilbenes, phenolic acids/hydroxybenzoates, and hydroxycinnamates (**Tables 1 and 2**; 15, 17, 18, 29, 30, 41, 43, 56, 71, 78, 84, 85, 88, 90). Quercetin is the major flavonol in grapes (**Figure 4A**) and usually occurs as *O*-glucosides in the D-glucose isoform such as quercetin 3-*O*-glucoside (**Figure 4B**). Grape flavan-3-ols are mainly represented by monomeric catechins such as (+)-catechin (**Figure 4C**) and oligomeric procyanidins, also known as proanthocyanidins or condensed tannins, such as pro-




Figure 3

Mechanisms by which tumor necrosis factor alpha (TNF α) or lipopolysaccharide (LPS) antagonize peroxisome proliferator-activated receptor gamma (PPAR γ) activity. Tumor necrosis factor α and LPS or their signals antagonize PPAR γ activity by (a) decreasing PPAR γ mRNA levels, (b) phosphorylation (P) of PPAR γ on serine 112 by extracellular signal-related kinase (ERK), leading to PPAR γ ubiquitination and proteasomal degradation, (c) activating nuclear factor-kappa B (NF- κ B), which antagonizes PPAR γ activity and its DNA binding capacity, or (d) activating nuclear corepressors, which decrease PPAR γ transcriptional activity and target gene expression. Collectively, this leads to insulin resistance in white adipose tissue, thereby elevating circulating free fatty acids (lipodystrophy) and glucose (hyperglycemia) that promote ectopic lipid deposition in liver, skeletal muscle, heart, and pancreas. 9-*cis* RA, 9-*cis* retinoic acid; α P2, adipocyte fatty acid-binding protein; GLUT4, insulin-dependent glucose transporter 4; LPL, lipoprotein lipase; PPRE, peroxisome proliferator response element; RXR, retinoid X receptor, TG, triglyceride.

cyanidin B2 (**Figure 4D**). Anthocyanins (i.e., anthocyanidins with sugar groups) in grapes are primarily 3-*O*-glucosides such as malvidin (**Figure 4E**) and malvidin 3-*O*-glucoside (**Figure 4F**), which are found in red grapes. *Trans*-resveratrol (**Figure 4G**) is probably the most well-known and studied stilbene found in grapes and red wine. Phenolic acids, also known as hydroxybenzoates, are commonly represented by gallic acid (**Figure 4H**) in grapes. In grapes or grape products, hydroxycinnamates usually undergo esterification with tartaric acid such as caffeic acids (**Figure 4I**) and its tartaric acid esters such as caftaric acid (**Figure 4J**).

The phenolic phytochemical content of grapes varies due to grape type, color, and ripeness, and climatic, geographical, and cultural factors. In general, flavonols and anthocyanidins/anthocyanins represent colored

Table 1 The content of polyphenols analyzed in freeze-dried table grape powder (GP)^a

Polyphenols	Compounds	Content (mg/kg GP)
Anthocyanins	Malvidin	145.2
	Cyanidin	125.0
	Peonidin	31.7
Flavonols	Quercetin	32.6
	Isorhamnetin	6.8
	Kaempferol	5.6
Flavan-3-ols	Catechin	19.7
	Epicatechin	12.6
Stilbenes	Resveratrol	1.75

^aData are from California Table Grape Commission Information Sheet (2005), http://agbioresearch.msu.edu/rfp/ca_grape2009.pdf.

phenols and flavan-3-ols, phenolic acids/hydroxybenzoates, and hydroxycinnamates represent noncolored phenols in grapes or grape products (Table 2). These polyphenols are present mainly in skins and seeds and contribute to the astringency, bitterness, and color of the grapes and wine. Polyphenols also impact the quality of the wine produced and potential health benefits associated with wine consumption.

Relatively High Abundance of Anthocyanidins/Anthocyanins and Flavonols in Grapes

Grapes are particularly rich in anthocyanidins/anthocyanins (e.g., malvidin, cyanidin, and peonidin) and to a lesser extent flavonols (e.g., quercetin) (Tables 1 and 2). The average of daily anthocyanidin/anthocyanin and flavonol intake from all foods in the United States is approximately 12.5 mg (124) and 20.0 mg (68, 99) per person, respectively. Average adult plasma levels of flavonols such as quercetin and isorhamnetin, a metabolite of quercetin, are approximately 53.9 nM and 3.0 nM, respectively (34). However, anthocyanidins/anthocyanins such as malvidin 3-*O*-glucoside were undetectable in the plasma of adult subjects (12, 40). Also, variation in plasma levels of polyphenols or their metabolites between individuals is high at baseline and after intervention (34, 75).

The bioavailability of these anthocyanidins/anthocyanins and flavonols has been reported to be poor based on in vitro (11, 30, 37, 130) and in vivo studies (reviewed in 32, 96). However, phenolic phytochemicals may be rapidly metabolized by cells and therefore difficult to detect within the circulation or cells following consumption or supplementation (105). Also, despite having relatively low abundance and poor bioavailability, some polyphenols have potent biological actions (i.e., resveratrol; 5, 6, 62). Because of the relative high abundance and biological activity of grape anthocyanidins/anthocyanins and flavonols, the following sections focus on potential mechanisms by which they reduce chronic inflammation associated with obesity and related metabolic diseases.

MECHANISMS BY WHICH ANTHOCYANIDINS/ ANTHOCYANINS AND FLAVONOLS, MAJOR PHENOLIC PHYTOCHEMICALS IN GRAPES, REDUCE CHRONIC INFLAMMATION

Acting as an Antioxidant or Increasing Antioxidant Gene or Protein Expression

Reactive oxygen species and reactive nitrogen species (RNS) can be generated by enzyme systems such as nicotinamide adenine dinucleotide phosphate (NADPH) oxidase and nitric oxide synthase (NOS), respectively, or transition metals, and are notorious mediators of oxidative stress and inflammation. Reactive oxygen species trigger redox-sensitive kinases such as apoptosis signal-regulating kinase 1 (ASK1) that activate downstream MAPKs, NF-κB, and AP-1, which in turn induce inflammatory gene expression. Phenolic phytochemicals have a strong antioxidant potential owing to the abundance of hydroxyl groups associated with their aromatic rings. Phenolic phytochemicals also have the capacity to increase the levels of anti-inflammatory

Table 2 Phenolic phytochemicals found in grapes and grape products

Resource	Phenolic phytochemicals	References
Whole grapes	Flavonol glycosides (quercetin, kaempferol, myricetin, laricitrin, isorhamnetin, syringetin), anthocyanins ^a (malvidin, peonidin, petunidin, cyanidin, delphinidin, pelargonidin), flavan-3-ols (catechin, epicatechin), phenolic acids (protocatechuic acid, gallic acid), hydroxycinnammates (caftaric acid, coutaric acid, fertaric acid), stilbenes (<i>trans</i> -resveratrol)	(15, 17, 78)
Grape seed	Flavan-3-ols (catechin, catechin gallate, epicatechin, epicatechin gallate, procyanidin gallate, B1, B2, other dimers, trimers, and tetramers)	(18, 84)
Grape skin	Flavonol glycosides (quercetin, kaempferol, myricetin, laricitrin, isorhamnetin, syringetin), anthocyanins ^a (malvidin, peonidin, petunidin, cyanidin, delphinidin, pelargonidin), flavan-3-ols (catechin, epicatechin, gallo catechin, procyanidin B1, B2, B4, other dimers, C1, other trimers, and tetramers), phenolic acids (protocatechuic acid, gallic acid), hydroxycinnammates (caftaric acid, coutaric acid, fertaric acid), stilbenes (<i>trans</i> -resveratrol, <i>cis</i> -resveratrol, resveratrol dimmers and tetramers), flavanonol glycosides (taxifolin)	(15, 17, 18, 84)
Grape stem	Flavonols (quercetin glucoside), flavan-3-ols (catechin, catechin gallate, epicatechin, procyanidin B1 and B2), hydroxycinnammates (caffeic acid, coumaric acid), stilbenes (<i>trans</i> -resveratrol, <i>trans</i> -resveratrol glucoside, resveratrol dimers, trimers, and tetramers), flavanonols (taxifolin glucoside)	(71, 90)
Grape leaf	Flavonols (quercetin, rutin, kaempferol, myricetin), flavan-3-ols (catechin), phenolic acids (gallic acid, ellagic acid), stilbenes (resveratrol), flavanones (naringin)	(29, 84)
Red grape juice	Flavonols (quercetin glucoside, rutin, myricetin), anthocyanins ^a (malvidin, peonidin, petunidin, cyanidin, delphinidin), flavan-3-ols (catechin, procyanidin B2)	(30)
Red wine	Flavonol glycosides (quercetin, kaempferol, myricetin, isorhamnetin), anthocyanins ^a (malvidin, peonidin, petunidin, cyanidin, delphinidin), flavan-3-ols (catechin, catechin gallate, epicatechin, epicatechin gallate, procyanidin B1, B2, B4, and trimers), phenolic acids (protocatechuic acid, gallic acid, ellagic acid, vanillic acid, syringic acid), hydroxycinnammates (caffeic acid, caftaric acid, coutaric acid, ferulic acid, fertaric acid, coumaric acid, sinapic acid), stilbenes (<i>trans</i> -resveratrol, <i>trans</i> -resveratrol glucoside)	(41, 43, 85, 88)
Raisin	Flavonol glycosides (quercetin, kaempferol), phenolic acids (protocatechuic acid), hydroxycinnammates (caftaric acid, coutaric acid)	(56)

^aAnthocyanidins/anthocyanins are detected only in red grapes.

genes such as superoxide dismutase (SOD), glutathione peroxidase (GPx), and heme oxygenase (HO)-1 via activation of the transcription factor nuclear factor-erythroid 2 (NF-E2)-related factor 2 (Nrf2) (reviewed in 91). Thus, polyphenols have an inherent capacity to reduce ROS and other free radicals, thereby preventing their activation of oxidative stress and inflammation. Consistent with this antioxidant potential, Sprague-Dawley rats consuming a high-fructose diet supplemented with a grape skin extract (21 mg/kg body weight for six weeks) were protected against ROS production, possibly due to reduced levels of NADPH oxidase, thereby preventing cardiac hypertrophy and hypertension induced by a high-fructose diet (2).

In vitro, pretreatment with freeze-dried grape powder (300 µg/ml) restored glutathione (GSH) content in human hepatoma cells (Huh7 cells) and primary mouse hepatocytes treated with hydrogen peroxide (H₂O₂) (126). The flavonols quercetin and kaempferol (5–50 µM) decreased the levels of oxidized GSH, peroxides, superoxide anions, and nitric oxide in parenchymal liver cells (Chang liver cells) treated with a mixture of inflammatory cytokines (28). Quercetin pretreatment (25 µM) of primary cultures of rat neurons exposed to H₂O₂ increased GSH levels, the nuclear translocation of Nrf2, and the expression of γ-glutamate-cysteine ligase catalytic subunit (GCLC), the rate-limiting enzyme for GSH synthesis, compared with those treated with




Figure 4

Structures of polyphenols commonly found in grape products.

H₂O₂ only (3). However, cells treated with doses of quercetin higher than 25 μ M for 24 hours showed signs of cytotoxicity.

Quercetin and isorhamnetin (10 μ M) pretreatment of murine macrophages (RAW264.7 cells) treated with LPS decreased the expression of inflammatory genes and inducible NOS and increased the protein levels of HO-1 (10). However, cells treated with doses of these flavonols equal to or greater than 25 μ M for 24 hours showed signs of cytotoxicity. Quercetin pretreatment (10–50 μ M) of immortalized human keratinocytes (HaCaT cells) treated with H₂O₂ increased mitochondrial membrane potential and cell viability and reduced the percentage of apoptotic cells compared with controls (122). Quercetin pretreatment (10 and 100 μ M) of human hepatoma cells treated with H₂O₂ or CuSO₄ (to initiate the Fenton reaction for the production of hydroxyl radicals) acutely decreased ROS production (60). However, long-term treatment with high levels of quercetin (100 μ M) decreased cell viability, suggesting pro-oxidant actions of quercetin at high levels. Pretreatment with the anthocyanin cyanidin 3-glucoside (10–40 μ M) of murine adipocytes (3T3-L1 cells) exposed to H₂O₂ or TNF α decreased ROS production and insulin resistance compared with controls (46). Quercetin and rutin (1–25 μ M), a quercetin metabolite, blocked oxidized low-density lipoprotein (LDL)-mediated apoptosis of human umbilical vein endothelial cells via modulation of Janus kinase/signal transducers and activators of the transcription signaling pathway (23). Rutin, but not quercetin, inhibited JNK and p38 signaling by decreasing the activation of ASK1, a redox-sensitive kinase that triggers inflammatory MAPK signaling.

Collectively, these data suggest that grape polyphenols or their metabolites have the capacity to protect cells against oxidative damage by (a) neutralizing ROS and RNS or transition metals that produce ROS, RNS, and oxidize GSH, (b) decreasing the activity of enzymes such as NADPH oxidase and NOS that produce ROS and RNS, respectively, (c) suppressing inflammatory signaling cascades

including ASK1 and downstream MAPK, and (d) activating transcription factors such as Nrf2 that induce the transcription of antioxidant enzymes such as GPx, SOD, HO-1, and GCLC (**Figure 5**). However, high levels of these polyphenols may be cytotoxic.

Attenuating Endoplasmic Reticulum Stress Signaling

The development of obesity and the metabolic syndrome has been linked to ER stress and inflammation (reviewed in 49). Endoplasmic reticulum stress activates the unfolded protein response (UPR), which involves activation of three ER membrane-associated proteins: (a) double-stranded RNA-activated protein kinase (PKR)-like eukaryotic initiation factor 2 α (eIF2 α) kinase (PERK), (b) inositol-requiring enzyme 1 (IRE1), and (c) activating transcription factor 6 (ATF6). Studies on flavonols suggest that kaempferol (10 μ M) or quercetin (25–150 μ M) attenuate ER stress in a rat cardiac muscle cell line (H9c2 cells) and isolated rat hearts (59) or in human colon cancer cell lines (Caco-2 and LS180 cells) (76), respectively, by blocking PERK-mediated eIF2 α phosphorylation, IRE1-mediated X-box-binding protein 1 activation, and ATF6 expression (**Figure 5**). In contrast, freeze-dried grape powder (300 μ g/ml) did not prevent ER stress-mediated apoptosis of human hepatoma cell line (Huh7 cell) and primary mouse hepatocytes (126). Research investigating the inhibitory effects of anthocyanidins/anthocyanins on ER stress is lacking, and the preventive effects of grape anthocyanidins/anthocyanins and flavonols on ER stress and chronic inflammation are unknown.

Blocking Pro-Inflammatory Cytokines or Endotoxin-Mediated Kinases and Transcription Factors Involved in Metabolic Diseases

Anthocyanins and flavonols modulate inflammation and insulin resistance associated with obesity. For example, Zucker fatty rats




Figure 5

Potential mechanisms by which grape polyphenols prevent chronic inflammation associated with obesity. Grape polyphenols may attenuate oxidative or endoplasmic reticulum (ER) stress-mediated inflammation associated with obesity by (a) inhibiting the activity of nicotinamide adenine dinucleotide phosphate (NADPH) oxidase or nitric oxide synthase (NOS) that generate reactive oxygen species (ROS) or reactive nitrogen species (RNS), respectively, (b) acting as a classic antioxidant by directly scavenging ROS and RNS, (c) preventing ROS-mediated activation of apoptosis signal-regulated kinase 1 (ASK1), a redox-sensitive, inflammatory signaling pathway, (d) increasing the expression of antioxidant genes such as glutathione peroxidase (GPx), superoxide dismutase (SOD), heme oxygenase (HO)-1, and γ -glutamate-cysteine ligase catalytic subunit (GCLC) via activation of transcription factor nuclear factor-erythroid 2 (NF-E2)-related factor 2 (Nrf2), or (e) blocking ER stress-mediated unfolded protein response including double-stranded RNA-activated protein kinase (PKR)-like eukaryotic initiation factor 2 α (eIF2 α) kinase (PERK)-mediated eIF2 α , inositol requiring enzyme 1 (IRE1)-mediated X-box binding protein 1 (XBP1), and activating transcription factor 6 (ATF6) signaling pathways that trigger proinflammatory gene expression. AP-1, activating protein-1; GSH, reduced glutathione; GSSG, oxidized glutathione; JNK, c-Jun-NH2 terminal kinase; NF- κ B, nuclear factor-kappa B; P, phosphorylated; Trx1, thioredoxin-1.

fed a high-fat diet supplemented with an anthocyanin-rich, 1% tart cherry powder (w/w) for 12 weeks had lower WAT and plasma levels of TNF α and interleukin (IL)-6 and WAT NF- κ B activity compared with controls (101). Dahl salt-sensitive, hypertensive rats supplemented with 3% grape powder in the diet (w/w) for 18 weeks had lower plasma levels of TNF α and IL-6, decreased cardiac tissue oxidative damage (102), and reduced expression of inflammatory genes and NF- κ B DNA binding activity in cardiac tissue (100) compared with high-sugar controls (3% fructose:glucose; 1:1). C57BL/6J mice fed a high-fat diet supplemented for eight weeks with an anthocyanin-rich, 4% blueberry

powder (w/w) had lower indices of inflammation (e.g., decreased gene expression of markers of WAT M Φ and inflammation) and oxidative stress and greater insulin sensitivity compared with control mice (33). C57BL/6J mice fed a high-fat diet supplemented with 0.8% quercetin in the diet (w/w) for eight weeks had lower plasma levels of interferon γ , IL-1 α , and IL-4 compared with controls (106). Finally, Zucker fatty rats receiving a daily dose of quercetin (2 or 10 mg/kg body weight, o.p.) for 10 weeks had lower inflammatory markers, improved insulin sensitivity and blood lipid profiles, and decreased blood pressure compared with placebo controls (94).

IL: interleukin

In vitro, cyanidin 3-glucoside (10–40 μ M) prevented TNF α -mediated JNK activation, IRS-1 phosphorylation at serine residue 307, and insulin resistance in 3T3-L1 adipocytes (46). Notably, grape seed flavan-3-ols/procyanidins (50 and 100 mg/liter) modulated LPS- and TNF α -induced inflammatory signal and gene expression in human M Φ (differentiated THP-1 monocytes) and Simpson-Golabi-Behmel Syndrome adipocytes, respectively (19). Moreover, oligomerized grape seed polyphenols (10 and 20 μ g/ml) attenuated activation of inflammatory signaling and production of inflammatory cytokines in a murine cell line of adipocytes (HW mouse white adipocytes) cocultured with a murine M Φ cell line (RAW264 cells) (98).

Consistent with these data, our lab demonstrated that grape powder extract (GPE; 30–100 μ g/ml), made from grape powder provided by the California Table Grape Commission, attenuated inflammatory gene expression (i.e., IL-6, IL-1 β , IL-8, and MCP-1) in primary human adipocytes induced by conditioned media collected from LPS-treated M Φ (differentiated U937 monocytes) (81). Furthermore, quercetin (3–30 μ M), the most abundant polyphenol in GPE (81), prevented inflammation in human M Φ and primary human adipocytes treated with M Φ conditioned media (82). We showed that pretreatment of human M Φ with quercetin (3–30 μ M) prevented M Φ -mediated insulin resistance in primary human adipocytes (82). We also demonstrated that GPE (10–60 μ g/ml) and quercetin (3–60 μ M) attenuated TNF α -mediated inflammation and insulin resistance in primary human adipocytes by blocking activation of ERK, JNK, NF- κ B, and AP-1 signaling and negative regulators of insulin signaling (e.g., protein tyrosine phosphatase (PTP)-1B and phosphorylation of serine residue 307 on IRS) (25, 26). Furthermore, the quercetin content of the cells increased within one hour of treatment with 30 μ M quercetin, suggesting that quercetin is rapidly taken up by human adipocytes (26). No signs of cytotoxicity were observed for these doses of GPE or quercetin.

Quercetin (1–5 μ M) suppressed carcinogen-mediated inflammation by directly binding to MEK, which prevents activation of inflammatory MEK/MAPK signaling in JB6 P+ cells, a JB6 promotion-sensitive mouse skin epidermal cell line (65). Moreover, kaempferol (1 and 3 μ M) inhibited TNF α -mediated inflammation by directly blocking nuclear translocation and DNA binding of NF- κ B and AP-1 in A549 cells, an alveolar epithelial cell carcinoma cell line (21). Taken together, these data suggest that grape polyphenols, especially anthocyanins and flavonols, inhibit proinflammatory cytokine- and metabolic endotoxin-triggered activation of inflammatory MEK/MAPK, NF- κ B, and AP-1 signaling, which increase inflammatory gene expression (e.g., TNF α , IL-6, IL-8, IL-1 β , and MCP-1) and negative regulators of insulin signaling (**Figure 6**). Therefore, grape polyphenols may be useful in preventing inflammatory-mediated insulin resistance and other related metabolic diseases.

Suppressing Inflammatory- or Inducing Metabolic-Gene Expression via Increasing Histone Deacetylase Activity

Sirtuins (SIRT), class III histone deacetylases, are nicotinamide adenine dinucleotide (NAD⁺)-dependent deacetylases. Sirtuins deacetylate not only histones, but also non-histone proteins including transcription factors by transferring an acetyl group from the targeted protein to NAD⁺, generating deacetylated histones or nonhistone proteins, nicotinamide, and O-acetyl-ADP-ribose. This NAD⁺ dependency contributes to the role that SIRT play in the regulation of chromatin structure and gene expression, cell survival, and energy homeostasis. In a SIRT1 in vitro screening assay (51), several polyphenols such as *trans*-resveratrol and quercetin (100 μ M) were shown to activate SIRT1.

In vitro (0.01–10 μ M) and in vivo (2.5–400 mg/kg body weight/day) studies have reported that resveratrol activated SIRT1, thereby (*a*) decreasing inflammatory gene




Figure 6

Grape polyphenols may prevent inflammation and insulin resistance associated with obesity by blocking activation of mitogen-activated protein kinases (MAPKs), nuclear factor-kappa B (NF- κ B), and activating protein (AP)-1. Grape polyphenols may prevent inflammatory tumor necrosis factor alpha (TNF α)/TNF receptor (TNFR) or lipopolysaccharide (LPS)/toll-like receptor (TLR) signaling to MAPK kinases (MEK) and their downstream, extracellular signal-related kinase (ERK) and c-Jun-NH2 terminal kinase (JNK). This would attenuate the activation of inflammatory transcription factors NF- κ B and AP-1, which are potent inducers of inflammatory gene expression [e.g., TNF α , interleukin (IL)-6, IL-8, IL-1 β , and monocyte chemoattractant protein (MCP)-1] and negative regulators of insulin signaling [e.g., phosphorylation of serine residue 307 on insulin receptor substrate (IRS) and protein tyrosine phosphatase (PTP)-1B], and which suppress insulin signaling necessary for insulin-dependent glucose transporter 4 (GLUT4) translocation to the plasma membrane. AKT, AKT/protein kinase B; GLU, glucose; IKK, I κ B α kinase; PI3 K, phosphatidylinositol 3-kinases; P, phosphorylated; PPAR, peroxisome proliferator-activated receptor; Ser, serine; Tyr, tyrosine; Ub, ubiquitinated degradation.

expression by deacetylation/inactivation of NF- κ B (39, 79, 128, 132) and (b) improving insulin sensitivity by deacetylation/activation of PPAR γ coactivator-1 α (PGC-1 α) (62, 111), a regulator of PPAR activity (reviewed in 20). Activation of PPAR α or PPAR β/δ by PGC-1 α also enhances energy expenditure (reviewed in 38, 110). Consistent with these data, quercetin supplementation (12.5 and 25 mg/kg body weight/day) for seven days increased SIRT1 and PGC-1 α expression and mitochondrial biogenesis in male ICR mice (31). These findings suggest that resveratrol and quercetin prevent inflammation by activating SIRT1, which deacetylates NF- κ B (inactive form) and PGC-1 α (active form), thereby suppressing inflammatory- and inducing metabolic-gene expression, mitochondrial

biogenesis, and oxidative phosphorylation via activation of PPAR (Figure 7). However, more research is needed on the regulation of SIRT1 by grape polyphenols to determine which specific anthocyanidins and flavonols are involved.

Activating Transcription Factors that Antagonize Chronic Inflammation

The PPARs (i.e., PPAR α , PPAR β/δ , and PPAR γ) are ligand-dependent, nuclear transcription factors that regulate energy homeostasis, glucose and lipid metabolism, and immune response (reviewed in 7). Activation of PPARs has been reported to suppress inflammatory gene expression by directly interfering with transcriptional activation of NF- κ B or AP-1 (reviewed in 93). For example, upon

activation by ligand binding, PPAR γ can be SUMOylated by binding to SUMO1, a small ubiquitin-like modifier. The SUMOylated PPAR γ subsequently binds to nuclear receptor corepressors, which interferes with clearance of the corepressor complex of NF- κ B, thereby transrepressing LPS-mediated NF- κ B activation (83).

Several studies reported that anthocyanin-rich cherries or berries increase the level of PPAR γ gene, protein, or activity. For example, the supplementation with 1% tart cherry in the diet (w/w) for 12 weeks or 3% grape powder in the diet (w/w) for 18 weeks suppressed inflammation and improved metabolic diseases by increasing PPAR γ gene expression in Zucker fatty rats (101) or by increasing PPAR α / γ mRNA levels and protein activity in Dahl salt-sensitive, hypertensive rats (100), respectively. Consistent with these data, our lab (26) and others (36) showed that by increasing PPAR γ activation, quercetin or kaempferol (3–60 μ M) attenuated inflammation or insulin resistance in adipocytes. These findings suggest that grape polyphenols increase the expression or activation of PPARs that antagonize inflammatory transcription factors, thereby blocking inflammation and the development of metabolic diseases (Figures 6 and 7).

CONCLUSIONS AND IMPLICATIONS

Potential Health Benefits of Consuming Grapes or Grape Products

Cardiovascular disease. Inflammatory mediators released from excess WAT cause endothelial dysfunction, plaque initiation and progression, and plaque rupture, leading to CVD (reviewed in 63). Supplementation with quercetin (64 mg/kg body weight/day for 10 and 20 weeks) attenuated inflammation and endothelial dysfunction in a mouse model of atherosclerosis (ApoE $^{-/-}$ knockout mice) (69). Randomized, clinical trials with proanthocyanidin-rich cocoa powder (40 g cocoa powder in 500 ml skim milk/day for four




Figure 7

Grape polyphenols may activate sirtuin 1 (SIRT1) and peroxisome proliferator-activated receptor (PPAR) that antagonize inflammation and improve the metabolic diseases associated with obesity. Grape polyphenols may prevent inflammation and improve metabolic diseases associated with obesity by activating SIRT1. SIRT1 has been reported to (a) deacetylate/inactivate nuclear factor-kappa B (NF- κ B), resulting in the suppression of inflammatory gene expression, and (b) deacetylate/activate PPAR γ coactivator-1 alpha (PGC-1 α), resulting in the activation of PPARs. Activation of PPAR α and PPAR β / δ increases oxidative phosphorylation, metabolic gene expression, and mitochondrial (Mito) biogenesis. Activation of PPAR γ may antagonize NF- κ B transcriptional activation of inflammatory genes, thereby attenuating inflammation associated with obesity.

weeks) (73) and anthocyanin- and flavonol-rich bilberry juice (330 ml/day for four weeks) (57) showed decreased plasma markers of inflammation in adults with high CVD risk who consumed these polyphenol-rich products.

Consumption of red grape juice (50 ml concentrate twice per day for two weeks) reduced serum markers of inflammation and oxidative stress/oxidized-LDL and improved dyslipidemia in hemodialysis patients (16). Consumption of raisins (one cup per day for six weeks) reduced systolic blood pressure and inflammatory cytokines and improved lipid profiles in men and postmenopausal women (89). Furthermore, consumption of grape seed flavan-3-ols procyanidins (0.32 mg/g diet/day for 19 weeks) attenuated inflammatory markers in liver, WAT, and circulation in high-fat-fed, obese male Zucker fatty rats (114). Also, consumption of whole table grape powder (3% in the diet, w/w, for 18 weeks) attenuated systolic blood pressure, cardiac hypertrophy, and diastolic dysfunction in Dahl salt-sensitive hypertensive rats (102). Finally, Pérez-Jiménez & Saura-Calixto (86) reviewed 75 trials including human and animal studies that investigated the relationship between consumption of grapes or

grape products and their effects on lowering risk factors for CVD including (*a*) inflammation, (*b*) oxidative stress/LDL oxidation, (*c*) endothelial dysfunction, (*d*) platelet aggregation, (*e*) dyslipidemia, and (*f*) hypertension. They concluded from these studies that grapes and grape products have antihypertensive, antihyperlipidemic, antiatherosclerotic, and antioxidant effects. Therefore, supplementation with grapes or grape products rich in polyphenols may be a useful dietary strategy for the attenuation of CVD associated with obesity and inflammation.

Insulin resistance or diabetes. Obesity-associated, low-grade inflammation causes insulin resistance (**Figures 1 and 2**). Insulin resistance is a prediabetic state characterized by decreased tissue sensitivity to insulin and hyperinsulinemia. These chronic perturbations in glucose disposal lead to the development of noninsulin-dependent diabetes (type 2) and eventually pancreatic beta cell failure, resulting in insulin-dependent diabetes (type 1) (reviewed in 108). Elevated inflammatory TNF α , LPS, SFA, or FFA contribute to obesity-associated insulin resistance as shown in **Figure 2**.

Grape polyphenols such as flavan-3-ols/procyanidins or stilbenes/resveratrol improved glucose intolerance in type 1 diabetic rat models (1, 22, 35, 87, 109). Resveratrol (0.04% in the diet w/w for 12–48 weeks, or 2.5–400 mg/kg body weight/day for 15–16 weeks) improved glucose and lipid homeostasis in high-fat-fed C57BL/6J obese mice (6, 62, 111). Consumption of 150 ml of muscadine grape wine or dealcoholized grape wine per day for 28 days improved fasting blood glucose and insulin levels in subjects with type 2 diabetes (4). Overall, results suggest that grapes or grape products are good candidates for dietetic management of type 2 diabetes because of their abundant polyphenol content and low glycemic index (134). However, clinical studies investigating the effect of grape consumption on insulin resistance or type 2 diabetes are limited.

Cancer. Obesity-associated inflammation contributes to cancer initiation or progression (reviewed in 27, 52, 80). García-Lafuente et al. (42) summarized the potential anti-inflammatory mechanisms of flavonoids found in fruits and vegetables that are linked to the prevention or treatment of cancer. For example, resveratrol prevents certain cancers due to its ability to inactivate inflammatory signaling, including protein kinases such as MEK/MAPK and transcription factors such as NF- κ B and AP-1, and to modulate carcinogenic signaling by down-regulation of cyclooxygenase (COX)-2 and iNOS (47, 54, 72, 113; reviewed in 8, 112, 117).

Interestingly, the anticarcinogenic effects of red wine extract may be due to not only resveratrol but to flavonols as well (65). For example, flavonol/quercetin (9, 24, 64, 67, 123) and anthocyanidin/delphinidin (53, 61, 120, 131) exert anti-inflammatory and anticancer activity. However, most studies used pharmacological doses of these polyphenols. Future studies are needed to investigate the relationship between physiologically relevant doses of polyphenols from grapes and grape products and their ability to prevent or treat certain types of cancer.

Neurodegenerative diseases linked to obesity. Grapes and grape products have been reported to have antiaging and antineurodegenerative effects due to their antioxidant properties (reviewed in 55). Moreover, rats fed 6.5 ml/kg body weight per day white or red wine or 2.5 mg/kg body weight per day resveratrol for two weeks had higher expression levels of genes and proteins related to longevity (74). Consumption of red wine containing 0.2 mg/liter of resveratrol for seven months attenuated the development of Alzheimer's disease (AD) in Tg2576 AD transgenic mice due to reduced aggregation of brain amyloid beta-protein (119). Grape powder or extract also has been reported to modulate the pathogenesis of neurodegenerative disease (70, 115, 118, 121). However, the extent to which polyphenols in grapes other than resveratrol extend lifespan

or prevent neurodegenerative diseases is unknown, as are their mechanisms of action.

Potential Risks of Excess Consumption of Grapes or Grape Products

Overconsumption of grapes or grape products could lead to (a) excess weight gain due to the high sugar content of grapes, (b) immunosuppression due to the anti-inflammatory actions of polyphenols, (c) impairment of micronutrient absorption or metabolism, and (d) alcohol toxicity associated with excess wine consumption. Of these four, the deleterious effects of excess alcohol consumption (e.g., gastrointestinal cancers, cirrhosis, pancreatitis, malabsorption, nonsteroidal anti-inflammatory drug interactions, impaired judgment, alcoholism, mania) are well-documented risks. Thus, moderation is the key with regard to consuming grapes or their by-products, especially alcoholic beverages made from grapes. Detrimental effects of consuming high levels of supplements made from grapes or their by-products are not well reported in the literature.

Future Research

Potential mechanisms by which polyphenol-rich grape products reduce chronic inflammation and metabolic diseases associated with




Figure 8

Summary of potential mechanisms by which polyphenol-rich grape products reduce chronic inflammation and metabolic diseases associated with obesity. AP-1, activating protein-1; ER, endoplasmic reticulum; MAPK, mitogen-activated protein kinase; NF-κB, nuclear factor-kappa B; PPAR, peroxisome proliferator-activated receptor; PGC-1, PPARγ coactivator-1; ROS, reactive oxygen species; SIRT, sirtuins.

obesity are summarized in **Figure 8**. The following research foci on polyphenol-rich grape products warrant further investigation: (a) identify the most bioactive components in grapes or grape products and their mechanisms of action; (b) investigate the bioavailability and efficacy of individual grape bioactive components versus whole foods, beverages, extracts, or supplements; (c) examine additive or synergistic effects of grape polyphenols in vitro and in vivo; (d) design and carry out well-controlled cell, animal, and human studies with purified single and mixed grape compounds known to be taken up by specific tissues; and (e) determine potential side effects of consuming high levels of grapes or grape products, grape supplements, or individual or mixtures of grape polyphenols.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

We thank our lab group members Kristina Martinez, Wan Shen, and Robin Hopkins for their thorough review of this manuscript. We also thank our funding sources, the NC Agricultural Research Service Project #02288 and the W.B. Kellogg Institute for Food and Marketing, for our research with grape products. Finally, we thank the California Table Grape Commission for providing us with table grape powder.

LITERATURE CITED

1. Al-Awwadi N, Azay J, Poucheret P, Cassanas G, Krosniak M, et al. 2004. Antidiabetic activity of red wine polyphenolic extract, ethanol, or both in streptozotocin-treated rats. *J. Agric. Food Chem.* 52:1008–16
2. Al-Awwadi NA, Araiz C, Bornet A, Delbosc S, Cristol JP, et al. 2005. Extracts enriched in different polyphenolic families normalize increased cardiac NADPH oxidase expression while having differential effects on insulin resistance, hypertension, and cardiac hypertrophy in high-fructose-fed rats. *J. Agric. Food Chem.* 53:151–57
3. Arredondo F, Echeverry C, Abin-Carriquiry JA, Blasina F, Antúnez K, et al. 2010. After cellular internalization, quercetin causes Nrf2 nuclear translocation, increases glutathione levels, and prevents neuronal death against an oxidative insult. *Free Radic. Biol. Med.* 49:738–47
4. Banini AE, Boyd LC, Allen JC, Allen HG, Sauls DL. 2006. Muscadine grape products intake, diet and blood constituents of non-diabetic and type 2 diabetic subjects. *Nutrition* 22:1137–45
5. Barger JL, Kayo T, Vann JM, Arias EB, Wang J, et al. 2008. A low dose of dietary resveratrol partially mimics caloric restriction and retards aging parameters in mice. *PLoS One* 3:e2264
6. Baur JA, Pearson KJ, Price NL, Jamieson HA, Lerin C, et al. 2006. Resveratrol improves health and survival of mice on a high-calorie diet. *Nature* 444:337–42
7. Berger JP, Akiyama TE, Meinke PT. 2005. PPARs: therapeutic targets for metabolic disease. *Trends Pharmacol. Sci.* 26:244–51
8. Bishayee A. 2009. Cancer prevention and treatment with resveratrol: from rodent studies to clinical trials. *Cancer Prev. Res.* 2:409–18
9. Bobe G, Albert PS, Sansbury LB, Lanza E, Schatzkin A, et al. 2010. Interleukin-6 as a potential indicator for prevention of high-risk adenoma recurrence by dietary flavonols in the polyp prevention trial. *Cancer Prev. Res.* 3:764–75
10. Boesch-Saadatmandi C, Loboda A, Wagner AE, Stachurska A, Jozkowicz A, et al. 2011. Effect of quercetin and its metabolites isorhamnetin and quercetin-3-glucuronide on inflammatory gene expression: role of miR-155. *J. Nutr. Biochem.* 22:293–99
11. Boyer J, Brown D, Liu RH. 2004. Uptake of quercetin and quercetin 3-glucoside from whole onion and apple peel extracts by Caco-2 cell monolayers. *J. Agric. Food Chem.* 52:7172–79
12. Bub A, Watzl B, Heeb D, Rechkemmer G, Briviba K. 2001. Malvidin-3-glucoside bioavailability in humans after ingestion of red wine, dealcoholized red wine and red grape juice. *Eur. J. Nutr.* 40:113–20
13. Cañete R, Gil-Campos M, Aguilera CM, Gil A. 2007. Development of insulin resistance and its relation to diet in the obese child. *Eur. J. Nutr.* 46:181–87
14. Cani PD, Bibiloni R, Knauf C, Waget A, Neyrinck A, et al. 2008. Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat fed mice-induced obesity and diabetes in mice. *Diabetes* 57:1470–81
15. Cantos E, Espín JC, Tomás-Barberán FA. 2002. Varietal differences among the polyphenol profiles of seven table grape cultivars studied by LC-DAD-MS-MS. *J. Agric. Food Chem.* 50:5691–96
16. Castilla P, Dávalos A, Teruel JL, Cerrato F, Fernández-Lucas M, et al. 2008. Comparative effects of dietary supplementation with red grape juice and vitamin E on production of superoxide by circulating neutrophil NADPH oxidase in hemodialysis patients. *Am. J. Clin. Nutr.* 87:1053–61
17. Castillo-Muñoz N, Fernández-González M, Gómez-Alonso S, García-Romero E, Hermosín-Gutiérrez I. 2009. Red-color related phenolic composition of Garnacha Tintorera (*Vitis vinifera* L.) grapes and red wines. *J. Agric. Food Chem.* 57:7883–91
18. Cavaliere C, Foglia P, Gubbiotti R, Sacchetti P, Samperi R, et al. 2008. Rapid-resolution liquid chromatography/mass spectrometry for determination and quantitation of polyphenols in grape berries. *Rapid Commun. Mass Spectrom.* 22:3089–99
19. Chacón MR, Ceperuelo-Mallafre V, Maymó-Masip E, Mateo-Sanz JM, Arola L, et al. 2009. Grape-seed procyanidins modulate inflammation on human differentiated adipocytes in vitro. *Cytokine* 47:137–42
20. Chaudhary N, Pfluger PT. 2009. Metabolic benefits from Sirt1 and Sirt1 activators. *Curr. Opin. Clin. Nutr. Metab. Care* 12:431–37
21. Chen CC, Chow MP, Huang WC, Lin YC, Chang YJ. 2004. Flavonoids inhibit tumor necrosis factor- α -induced up-regulation of intercellular adhesion molecule-1 (ICAM-1) in respiratory epithelial cells

- through activator protein-1 and nuclear factor-kappaB: structure-activity relationships. *Mol. Pharmacol.* 66:683–93
22. Chi TC, Chen WP, Chi TL, Kuo TF, Lee SS, et al. 2007. Phosphatidylinositol-3-kinase is involved in the antihyperglycemic effect induced by resveratrol in streptozotocin-induced diabetic rats. *Life Sci.* 80:1713–20
23. Choi JS, Kang SW, Li J, Kim JL, Bae JY, et al. 2009. Blockade of oxidized LDL-triggered endothelial apoptosis by quercetin and rutin through differential signaling pathways involving JAK2. *J. Agric. Food Chem.* 57:2079–86
24. Choi SY, Park JH, Kim JS, Kim MK, Aruoma OI, et al. 2006. Effects of quercetin and beta-carotene supplementation on azoxymethane-induced colon carcinogenesis and inflammatory responses in rats fed with high-fat diet rich in omega-6 fatty acids. *Biofactors* 27:137–46
25. Chuang CC, Bumrungpert A, Kennedy A, Overman A, West T, et al. 2011. Grape powder extract attenuates tumor necrosis factor alpha-mediated inflammation and insulin resistance in primary cultures of human adipocytes. *J. Nutr. Biochem.* 22:89–94
26. Chuang CC, Martínez K, Xie G, Kennedy A, Bumrungpert A, et al. 2010. Quercetin is equally or more effective than resveratrol in attenuating tumor necrosis factor- α -mediated inflammation and insulin resistance in primary human adipocytes. *Am. J. Clin. Nutr.* 92:1511–21
27. Coussens LM, Werb Z. 2002. Inflammation and cancer. *Nature* 420:860–67
28. Crespo I, García-Mediavilla MV, Almar M, González P, Tuñón MJ, et al. 2008. Differential effects of dietary flavonoids on reactive oxygen and nitrogen species generation and changes in antioxidant enzyme expression induced by proinflammatory cytokines in Chang liver cells. *Food Chem. Toxicol.* 46:1555–69
29. Dani C, Oliboni LS, Agostini F, Funchal C, Serafini L, et al. 2010. Phenolic content of grapevine leaves (*Vitis labrusca* var. Bordo) and its neuroprotective effect against peroxide damage. *Toxicol. In Vitro* 24:148–53
30. Dávalos A, Fernández-Hernando C, Cerrato F, Martínez-Botas J, Gómez-Coronado D, et al. 2006. Red grape juice polyphenols alter cholesterol homeostasis and increase LDL-receptor activity in human cells in vitro. *J. Nutr.* 136:1766–73
31. Davis JM, Murphy EA, Carmichael MD, Davis B. 2009. Quercetin increases brain and muscle mitochondrial biogenesis and exercise tolerance. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 296:R1071–77
32. de Pascual-Teresa S, Moreno DA, García-Viguera C. 2010. Flavanols and anthocyanins in cardiovascular health: a review of current evidence. *Int. J. Mol. Sci.* 11:1679–703
33. DeFuria J, Bennett G, Strissel KJ, Perfield JW 2nd, Milbury PE, et al. 2009. Dietary blueberry attenuates whole-body insulin resistance in high fat-fed mice by reducing adipocyte death and its inflammatory sequelae. *J. Nutr.* 139:1510–16
34. Egert S, Wolfram S, Bosy-Westphal A, Boesch-Saadatmandi C, Wagner AE, et al. 2008. Daily quercetin supplementation dose-dependently increases plasma quercetin concentrations in healthy humans. *J. Nutr.* 138:1615–21
35. El-Alfy AT, Ahmed AA, Fatani AJ. 2005. Protective effect of red grape seeds proanthocyanidins against induction of diabetes by alloxan in rats. *Pharmacol. Res.* 52:264–70
36. Fang XK, Gao J, Zhu DN. 2008. Kaempferol and quercetin isolated from *Euonymus alatus* improve glucose uptake of 3T3-L1 cells without adipogenesis activity. *Life Sci.* 82:615–22
37. Faria A, Pestana D, Azevedo J, Martel F, de Freitas V, et al. 2009. Absorption of anthocyanins through intestinal epithelial cells—putative involvement of GLUT2. *Mol. Nutr. Food Res.* 53:1430–37
38. Feige JN, Auwerx J. 2007. Transcriptional coregulators in the control of energy homeostasis. *Trends Cell Biol.* 17:292–301
39. Fischer-Posovszky P, Kukulus V, Tews D, Unterkircher T, Debatin KM, et al. 2010. Resveratrol regulates human adipocyte number and function in a Sirt1-dependent manner. *Am. J. Clin. Nutr.* 92:5–15
40. Garcia-Alonso M, Miniñane AM, Rimbach G, Rivas-Gonzalo JC, de Pascual-Teresa S. 2009. Red wine anthocyanins are rapidly absorbed in humans and affect monocyte chemoattractant protein 1 levels and antioxidant capacity of plasma. *J. Nutr. Biochem.* 20:521–29
41. García-Falcón MS, Pérez-Lamela C, Martínez-Carballo E, Simal-Gándara J. 2007. Determination of phenolic compounds in wines: influence of bottle storage of young red wines on their evolution. *Food Chem.* 105:248–59

42. García-Lafuente A, Guillaumon E, Villares A, Rostagno MA, Martínez JA. 2009. Flavonoids as anti-inflammatory agents: implications in cancer and cardiovascular disease. *Inflamm. Res.* 58:537–52
43. Gómez-Alonso S, García-Romero E, Hermosín-Gutiérrez I. 2007. HPLC analysis of diverse grape and wine phenolics using direct injection and multidetection by DAD and fluorescence. *J. Food Compos. Anal.* 20:618–26
44. Grundy S, Brewer H, Cleeman J, Smith S, Lenfant C. 2004. Definition of metabolic syndrome: Report of the National Heart, Lung, and Blood Institute/American Heart Association conference on scientific issues related to definition. *Circulation* 109:433–38
45. Guilherme A, Virbasius JV, Puri V, Czech MP. 2008. Adipocyte dysfunctions linking obesity to insulin resistance and type 2 diabetes. *Nat. Rev. Mol. Cell Biol.* 9:367–77
46. Guo H, Ling W, Wang Q, Liu C, Hu Y, Xia M. 2008. Cyanidin 3-glucoside protects 3T3-L1 adipocytes against H₂O₂- or TNF- α -induced insulin resistance by inhibiting c-Jun NH₂-terminal kinase activation. *Biochem. Pharmacol.* 75:1393–401
47. Holmes-McNary M, Baldwin AS Jr. 2000. Chemopreventive properties of trans-resveratrol are associated with inhibition of activation of the I κ B kinase. *Cancer Res.* 60:3477–83
48. Hotamisligil GS. 2006. Inflammation and metabolic disorders. *Nature* 444:860–67
49. Hotamisligil GS. 2010. Endoplasmic reticulum stress and the inflammatory basis of metabolic disease. *Cell* 140:900–17
50. Hotamisligil GS, Arner P, Caro JF, Atkinson RL, Spiegelman BM. 1995. Increased adipose expression of tumor necrosis factor- α in human obesity and insulin resistance. *J. Clin. Invest.* 95:2409–15
51. Howitz KT, Bitterman KJ, Cohen HY, Lamming DW, Lavu S, et al. 2003. Small molecule activators of sirtuins extend *Saccharomyces cerevisiae* lifespan. *Nature* 425:191–96
52. Hursting SD, Nunez NP, Varticovski L, Vinson C. 2007. The obesity-cancer link: lessons learned from a fatless mouse. *Cancer Res.* 67:2391–93
53. Hwang MK, Kang NJ, Heo YS, Lee KW, Lee HJ. 2009. Fyn kinase is a direct molecular target of delphinidin for the inhibition of cyclooxygenase-2 expression induced by tumor necrosis factor- α . *Biochem. Pharmacol.* 77:1213–22
54. Jang M, Cai L, Udeani GO, Slowing KV, Thomas CF, et al. 1997. Cancer chemopreventive activity of resveratrol, a natural product derived from grapes. *Science* 275:218–20
55. Joseph JA, Shukitt-Hale B, Willis LM. 2009. Grape juice, berries, and walnuts affect brain aging and behavior. *J. Nutr.* 139:1813–17S
56. Karadeniz F, Durst RW, Wrolstad RE. 2000. Polyphenolic composition of raisins. *J. Agric. Food Chem.* 48:5343–50
57. Karlsen A, Paur I, Bohn SK, Sakhi AK, Borge GI, et al. 2010. Bilberry juice modulates plasma concentration of NF- κ B related inflammatory markers in subjects at increased risk of CVD. *Eur. J. Nutr.* 49:345–55
58. Kennedy A, Martinez K, Chuang CC, LaPoint K, McIntosh M. 2008. Saturated fatty acid-mediated inflammation and insulin resistance in adipose tissue: mechanisms of action and implications. *J. Nutr.* 139:1–4
59. Kim DS, Ha KC, Kwon DY, Kim MS, Kim HR, et al. 2008. Kaempferol protects ischemia/reperfusion-induced cardiac damage through the regulation of endoplasmic reticulum stress. *Immunopharmacol. Immunotoxicol.* 30:257–70
60. Kim GN, Jang HD. 2009. Protective mechanism of quercetin and rutin using glutathione metabolism on HO-induced oxidative stress in HepG2 cells. *Ann. N. Y. Acad. Sci.* 1171:530–37
61. Kwon JY, Lee KW, Kim JE, Jung SK, Kang NJ, et al. 2009. Delphinidin suppresses ultraviolet B-induced cyclooxygenase-2 expression through inhibition of MAPK4 and PI-3 kinase. *Carcinogenesis* 30:1932–40
62. Lagouge M, Argmann C, Gerhart-Hines Z, Meziane H, Lerin C, et al. 2006. Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1 α . *Cell* 127:1109–22
63. Lau DC, Dhillon B, Yan H, Szmítko PE, Verma S. 2005. Adipokines: molecular links between obesity and atherosclerosis. *Am. J. Physiol. Heart Circ. Physiol.* 288:H2031–41
64. Lee KM, Hwang MK, Lee DE, Lee KW, Lee HJ. 2010. Protective effect of quercetin against arsenite-induced COX-2 expression by targeting PI3K in rat liver epithelial cells. *J. Agric. Food Chem.* 58:5815–20

65. Lee KW, Kang NJ, Heo YS, Rogozin EA, Pugliese A, et al. 2008. Raf and MEK protein kinases are direct molecular targets for the chemopreventive effect of quercetin, a major flavonol in red wine. *Cancer Res.* 68:946–55
66. Lefferova MI, Lazar MA. 2009. New developments in adipogenesis. *Trends Endocrinol. Metab.* 20:107–14
67. Lin CW, Hou WC, Shen SC, Juan SH, Ko CH, et al. 2008. Quercetin inhibition of tumor invasion via suppressing PKC delta/ERK/AP-1-dependent matrix metalloproteinase-9 activation in breast carcinoma cells. *Carcinogenesis* 29:1807–15
68. Lin J, Rexrode KM, Hu F, Albert CM, Chae CU, et al. 2007. Dietary intakes of flavonols and flavones and coronary heart disease in US women. *Am. J. Epidemiol.* 165:1305–13
69. Loke WM, Proudfoot JM, Hodgson JM, McKinley AJ, Hime N, et al. 2010. Specific dietary polyphenols attenuate atherosclerosis in apolipoprotein E-knockout mice by alleviating inflammation and endothelial dysfunction. *Arterioscler. Thromb. Vasc. Biol.* 30:749–57
70. Long J, Gao H, Sun L, Liu J, Zhao-Wilson X. 2009. Grape extract protects mitochondria from oxidative damage and improves locomotor dysfunction and extends lifespan in a *Drosophila* Parkinson's disease model. *Rejuvenation Res.* 12:321–31
71. Makris DP, Boskou G, Andrikopoulos NK, Kefalas P. 2008. Characterisation of certain major polyphenolic antioxidants in grape (*Vitis vinifera* cv. Roditis) stems by liquid chromatography-mass spectrometry. *Eur. Food Res. Technol.* 226:1075–79
72. Manna SK, Mukhopadhyay A, Aggarwal BB. 2000. Resveratrol suppresses TNF-induced activation of nuclear transcription factors NF-kappa B, activator protein-1, and apoptosis: potential role of reactive oxygen intermediates and lipid peroxidation. *J. Immunol.* 164:6509–19
73. Monagas M, Khan N, Andres-Lacueva C, Casas R, Urpí-Sardà M, et al. 2009. Effect of cocoa powder on the modulation of inflammatory biomarkers in patients at high risk of cardiovascular disease. *Am. J. Clin. Nutr.* 90:1144–50
74. Mukherjee S, Lekli I, Gurusamy N, Bertelli AA, Das DK. 2009. Expression of the longevity proteins by both red and white wines and their cardioprotective components, resveratrol, tyrosol, and hydroxytyrosol. *Free Radic. Biol. Med.* 1:573–78
75. Mullen W, Edwards CA, Crozier A. 2006. Absorption, excretion and metabolite profiling of methyl-, glucuronyl-, glucosyl- and sulpho-conjugates of quercetin in human plasma and urine after ingestion of onions. *Br. J. Nutr.* 96:107–16
76. Natsume Y, Ito S, Satsu H, Shimizu M. 2009. Protective effect of quercetin on ER stress caused by calcium dynamics dysregulation in intestinal epithelial cells. *Toxicology* 258:164–75
77. Neels JG, Olefsky JM. 2006. Inflamed fat: What starts the fire? *J. Clin. Invest.* 116:33–35
78. Nicoletti I, Bello C, De Rossi A, Corradini D. 2008. Identification and quantification of phenolic compounds in grapes by HPLC-PDA-ESI-MS on a semimicro separation scale. *J. Agric. Food Chem.* 56:8801–8
79. Olholm J, Paulsen SK, Cullberg KB, Richelsen B, Pedersen SB. 2010. Anti-inflammatory effect of resveratrol on adipokine expression and secretion in human adipose tissue explants. *Int. J. Obes.* 34:1546–53
80. Osório-Costa F, Rocha GZ, Dias MM, Carvalheira JB. 2009. Epidemiological and molecular mechanisms aspects linking obesity and cancer. *Arq. Bras. Endocrinol. Metabol.* 53:213–26
81. Overman A, Bumrungpert A, Kennedy A, Martinez K, Chuang CC, et al. 2010. Polyphenol-rich grape powder extract (GPE) attenuates inflammation in human macrophages and in human adipocytes exposed to macrophage-conditioned media. *Int. J. Obes. (Lond.)* 34:800–8
82. Overman A, Chuang CC, McIntosh MK. 2011. Quercetin attenuates inflammation in human macrophages and in human adipocytes exposed to macrophage-conditioned media. *Int. J. Obesity (Lond.)* In press. doi:10.1038/ijo.2010.272
83. Pascual G, Fong AL, Ogawa S, Gamliel A, Li AC, et al. 2005. A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR-gamma. *Nature* 437:759–63
84. Pastrana-Bonilla E, Akoh CC, Sellappan S, Krewer G. 2003. Phenolic content and antioxidant capacity of muscadine grapes. *J. Agric. Food Chem.* 51:5497–503
85. Pereira V, Câmara JS, Cacho J, Marques JC. 2010. HPLC-DAD methodology for the quantification of organic acids, furans and polyphenols by direct injection of wine samples. *J. Sep. Sci.* 33:1204–15

86. Pérez-Jiménez J, Saura-Calixto F. 2008. Grape products and cardiovascular disease risk factors. *Nutr. Res. Rev.* 21:158–73
87. Pinent M, Blay M, Bladé MC, Salvadó MJ, Arola L, et al. 2004. Grape seed-derived procyanidins have an antihyperglycemic effect in streptozotocin-induced diabetic rats and insulinomimetic activity in insulin-sensitive cell lines. *Endocrinology* 145:4985–90
88. Preys S, Mazerolles G, Courcoux P, Samson A, Fischer U, et al. 2006. Relationship between polyphenolic composition and some sensory properties in red wines using multiway analyses. *Anal. Chim. Acta* 563:126–36
89. Puglisi MJ, Vaishnav U, Shrestha S, Torres-Gonzalez M, Wood RJ, et al. 2008. Raisins and additional walking have distinct effects on plasma lipids and inflammatory cytokines. *Lipids Health Dis.* 7:14
90. Püssa T, Floren J, Kuldkepp P, Raal A. 2006. Survey of grapevine *Vitis vinifera* stem polyphenols by liquid chromatography-diode array detection-tandem mass spectrometry. *J. Agric. Food Chem.* 54:7488–94
91. Rahman I, Biswas SK, Kirkham PA. 2006. Regulation of inflammation and redox signaling by dietary polyphenols. *Biochem. Pharmacol.* 72:1439–52
92. Renaud S, de Lorgeril M. 1992. Wine, alcohol, platelets, and the French paradox for coronary heart disease. *Lancet* 339:1523–26
93. Ricote M, Glass CK. 2007. PPARs and molecular mechanisms of transrepression. *Biochim. Biophys. Acta* 1771:926–35
94. Rivera L, Morón R, Sánchez M, Zarzuelo A, Galisteo M. 2008. Quercetin ameliorates metabolic syndrome and improves the inflammatory status in obese Zucker rats. *Obesity* 16:2081–87
95. Rocha VZ, Libby P. 2009. Obesity, inflammation, and atherosclerosis. *Nat. Rev. Cardiol.* 6:399–409
96. Ross JA, Kasum CM. 2002. Dietary flavonoids: bioavailability, metabolic effects, and safety. *Annu. Rev. Nutr.* 22:19–34
97. Saghizadeh M, Ong JM, Garvey WT, Henry RR, Kern PA. 1996. The expression of TNF alpha by human muscle cells. Relationship to insulin resistance. *J. Clin. Invest.* 97:1111–16
98. Sakurai T, Kitadate K, Nishioka H, Fujii H, Kizaki T, et al. 2010. Oligomerized grape seed polyphenols attenuate inflammatory changes due to antioxidative properties in coculture of adipocytes and macrophages. *J. Nutr. Biochem.* 21:47–54
99. Sesso HD, Gaziano JM, Liu S, Buring JE. 2003. Flavonoid intake and the risk of cardiovascular disease in women. *Am. J. Clin. Nutr.* 77:1400–8
100. Seymour EM, Bennink MR, Watts SW, Bolling SF. 2010. Whole grape intake impacts cardiac peroxisome proliferator-activated receptor and nuclear factor kappaB activity and cytokine expression in rats with diastolic dysfunction. *Hypertension* 55:1179–85
101. Seymour EM, Lewis SK, Urcuyo-Llanes DE, Tanone II, Kirakosyan A, et al. 2009. Regular tart cherry intake alters abdominal adiposity, adipose gene transcription, and inflammation in obesity-prone rats fed a high fat diet. *J. Med. Food* 12:935–42
102. Seymour EM, Singer AA, Bennink MR, Parikh RV, Kirakosyan A, et al. 2008. Chronic intake of a phytochemical-enriched diet reduces cardiac fibrosis and diastolic dysfunction caused by prolonged salt-sensitive hypertension. *J. Gerontol. A Biol. Sci. Med. Sci.* 63:1034–42
103. St. Leger AS, Cochrane AL, Moore F. 1979. Factors associated with cardiac mortality in developed countries with particular reference to the consumption of wine. *Lancet* 1:1017–20
104. Steinberg GR, Michell BJ, Van Denderen BJ, Watt MJ, Carey AL, et al. 2006. Tumor necrosis factor alpha-induced skeletal muscle insulin resistance involves suppression of AMP-kinase signaling. *Cell Metab.* 4:465–74
105. Steinert RE, Ditscheid B, Netzel M, Jahreis G. 2008. Absorption of black currant anthocyanins by monolayers of human intestinal epithelial Caco-2 cells mounted in ussing type chambers. *J. Agric. Food Chem.* 56:4995–5001
106. Stewart LK, Soileau JL, Ribnick D, Wang ZQ, Raskin I, et al. 2008. Quercetin transiently increases energy expenditure but persistently decreases circulating markers of inflammation in C57BL/6J mice fed a high-fat diet. *Metabolism* 57:S39–46
107. Strissel KJ, Stancheva Z, Miyoshi H, Perfield JW 2nd, DeFuria J, et al. 2007. Adipocyte death, adipose tissue remodeling, and obesity complications. *Diabetes* 56:2910–18

108. Stumvoll M, Goldstein BJ, van Haeften TW. 2005. Type 2 diabetes: principles of pathogenesis and therapy. *Lancet* 365:1333–46
109. Su HC, Hung LM, Chen JK. 2006. Resveratrol, a red wine antioxidant, possesses an insulin-like effect in streptozotocin-induced diabetic rats. *Am. J. Physiol. Endocrinol. Metab.* 290:E1339–46
110. Sugden MC, Caton PW, Holness MJ. 2010. PPAR control: It's SIRTainly as easy as PGC. *J. Endocrinol.* 204:93–104
111. Sun C, Zhang F, Ge X, Yan T, Chen X, et al. 2007. SIRT1 improves insulin sensitivity under insulin-resistant conditions by repressing PTP1B. *Cell Metab.* 6:307–19
112. Surh YJ, Chun KS, Cha HH, Han SS, Keum YS, et al. 2001. Molecular mechanisms underlying chemopreventive activities of anti-inflammatory phytochemicals: down-regulation of COX-2 and iNOS through suppression of NF-kappa B activation. *Mutat. Res.* 480–481:243–68
113. Tao HY, Wu CF, Zhou Y, Gong WH, Zhang X, et al. 2004. The grape component resveratrol interferes with the function of chemoattractant receptors on phagocytic leukocytes. *Cell Mol. Immunol.* 1:50–56
114. Terra X, Montagut G, Bustos M, Llopiz N, Ardèvol A, et al. 2009. Grape-seed procyanidins prevent low-grade inflammation by modulating cytokine expression in rats fed a high-fat diet. *J. Nutr. Biochem.* 20:210–18
115. Thomas P, Wang YJ, Zhong JH, Kosaraju S, O'Callaghan NJ, et al. 2009. Grape seed polyphenols and curcumin reduce genomic instability events in a transgenic mouse model for Alzheimer's disease. *Mutat. Res.* 661:25–34
116. Tontonoz P, Spiegelman BM. 2008. Fat and beyond: the diverse biology of PPARgamma. *Annu. Rev. Biochem.* 77:289–312
117. Udenigwe CC, Ramprasath VR, Aluko RE, Jones PJ. 2008. Potential of resveratrol in anticancer and anti-inflammatory therapy. *Nutr. Rev.* 66:445–54
118. Wang J, Ho L, Zhao W, Ono K, Rosensweig C, et al. 2008. Grape-derived polyphenolics prevent Abeta oligomerization and attenuate cognitive deterioration in a mouse model of Alzheimer's disease. *J. Neurosci.* 28:6388–92
119. Wang J, Ho L, Zhao Z, Seror I, Humala N, et al. 2006. Moderate consumption of cabernet sauvignon attenuates Abeta neuropathology in a mouse model of Alzheimer's disease. *FASEB J.* 20:2313–20
120. Wang LS, Hecht SS, Carmella SG, Yu N, Larue B, et al. 2009. Anthocyanins in black raspberries prevent esophageal tumors in rats. *Cancer Prev. Res.* 2:84–93
121. Wang Q, Simonyi A, Li W, Sisk BA, Miller RL, et al. 2005. Dietary grape supplement ameliorates cerebral ischemia-induced neuronal death in gerbils. *Mol. Nutr. Food Res.* 49:443–51
122. Wang XY, He PY, Du J, Zhang JZ. 2010. Quercetin in combating H2O2 induced early cell apoptosis and mitochondrial damage to normal human keratinocytes. *Chin. Med. J. (Engl.)* 123:532–36
123. Warren CA, Paulhill KJ, Davidson LA, Lupton JR, Taddeo SS, et al. 2009. Quercetin may suppress rat aberrant crypt foci formation by suppressing inflammatory mediators that influence proliferation and apoptosis. *J. Nutr.* 139:101–5, erratum *J. Nutr.* 139:792
124. Wu X, Beecher GR, Holden JM, Haytowitz DB, Gebhardt SE, et al. 2006. Concentrations of anthocyanins in common foods in the United States and estimation of normal consumption. *J. Agric. Food Chem.* 54:4069–75
125. Xia EQ, Deng GF, Guo YJ, Li HB. 2010. Biological activities of polyphenols from grapes. *Int. J. Mol. Sci.* 11:622–46
126. Xu Y, Khaoustov VI, Wang H, Yu J, Tabassam F, Yoffe B. 2009. Freeze-dried grape powder attenuates mitochondria- and oxidative stress-mediated apoptosis in liver cells. *J. Agric. Food Chem.* 57:9324–31
127. Yadav M, Jain S, Bhardwaj A, Nagpal R, Puniya M, et al. 2009. Biological and medicinal properties of grapes and their bioactive constituents: an update. *J. Med. Food* 12:473–84
128. Yang Z, Kahn BB, Shi H, Xue BZ. 2010. Macrophage alpha1 AMP-activated protein kinase (alpha1AMPK) antagonizes fatty acid-induced inflammation through SIRT1. *J. Biol. Chem.* 285:19051–59
129. Ye J. 2008. Regulation of PPARgamma function by TNFalpha. *Biochem. Biophys. Res. Commun.* 374:405–8
130. Yi W, Akoh CC, Fischer J, Krewer G. 2006. Absorption of anthocyanins from blueberry extracts by caco-2 human intestinal cell monolayers. *J. Agric. Food Chem.* 54:5651–58

131. Yun JM, Afaq F, Khan N, Mukhtar H. 2009. Delphinidin, an anthocyanidin in pigmented fruits and vegetables, induces apoptosis and cell cycle arrest in human colon cancer HCT116 cells. *Mol. Carcinog.* 48:260–70
132. Zhu J, Yong W, Wu X, Yu Y, Lv J, et al. 2008. Anti-inflammatory effect of resveratrol on TNF-alpha-induced MCP-1 expression in adipocytes. *Biochem. Biophys. Res. Commun.* 369:471–77
133. Zinman B, Hanley AJ, Harris SB, Kwan J, Fantus IG. 1999. Circulating tumor necrosis factor-alpha concentrations in native Canadian population with high rates of type 2 diabetes mellitus. *J. Clin. Endocrinol. Metab.* 84:272–78
134. Zunino S. 2009. Type 2 diabetes and glycemic response to grapes or grape products. *J. Nutr.* 139:1794–800S



Contents

Nutritional Scientist or Biochemist? <i>J.W. Suttie</i>	1
Interaction Between Obesity and the Gut Microbiota: Relevance in Nutrition <i>Nathalie M. Delzenne and Patrice D. Cani</i>	15
The Implication of Brown Adipose Tissue for Humans <i>Eric Ravussin and José E. Galgani</i>	33
The Role of MicroRNAs in Cholesterol Efflux and Hepatic Lipid Metabolism <i>Kathryn J. Moore, Katey J. Rayner, Yajaira Suárez, and Carlos Fernández-Hernando</i>	49
Cytochrome P450s in the Regulation of Cellular Retinoic Acid Metabolism <i>A. Catharine Ross and Reza Zolfaghari</i>	65
Vitamin D in Pregnancy and Lactation in Humans <i>Patsy M. Brannon and Mary Frances Picciano</i>	89
Knockout Mouse Models of Iron Homeostasis <i>Robert E. Fleming, Qi Feng, and Robert S. Britton</i>	117
Zinc in Neurotransmission <i>Katalin Tóth</i>	139
Potential Mechanisms by Which Polyphenol-Rich Grapes Prevent Obesity-Mediated Inflammation and Metabolic Diseases <i>Chia-Chi Chuang and Michael K. McIntosh</i>	155
Mechanisms of Membrane Transport of Folate into Cells and Across Epithelia <i>Rongbao Zhao, Ndeye Diop-Bove, Michele Visentin, and I. David Goldman</i>	177
The Impact of Common Gene Variants on the Response of Biomarkers of Cardiovascular Disease (CVD) Risk to Increased Fish Oil Fatty Acids Intakes <i>Jacqueline Madden, Christine M. Williams, Philip C. Calder, Georg Lietz, Elizabeth A. Miles, Heather Cordell, John C. Mathers, and Anne Marie Minihane</i>	203

How Is Maternal Nutrition Related to Preterm Birth? <i>Frank H. Bloomfield</i>	235
How Many People Are Malnourished? <i>Peter Svedberg</i>	263
What Are the Risks and Benefits to Increasing Dietary Bone Minerals and Vitamin D Intake in Infants and Small Children? <i>Steven A. Abrams</i>	285
<u>Nutrigenomics, Rumen-Derived Bioactive Fatty Acids, and the Regulation of Milk Fat Synthesis</u> <u><i>Dale E. Bauman, Kevin J. Harvatine, and Adam L. Lock</i></u>	299
Docosahexaenoic Acid Signalolipidomics in Nutrition: Significance in Aging, Neuroinflammation, Macular Degeneration, Alzheimer's, and Other Neurodegenerative Diseases <i>Nicolas G. Bazan, Miguel F. Molina, and William C. Gordon</i>	321
Energy Intake and Response to Infection with Influenza <i>Elizabeth M. Gardner, Eleni Beli, Jonathan F. Clinthorne, and David M. Duriancik</i>	353

Indexes

Cumulative Index of Contributing Authors, Volumes 27–31	369
Cumulative Index of Chapter Titles, Volumes 27–31	372

Errata

An online log of corrections to *Annual Review of Nutrition* articles may be found at
<http://nutr.annualreviews.org/errata.shtml>