

WJD 5<sup>th</sup> Anniversary Special Issues (2): Type 2 diabetes

## Functional foods-based diet as a novel dietary approach for management of type 2 diabetes and its complications: A review

Parvin Mirmiran, Zahra Bahadoran, Fereidoun Azizi

Parvin Mirmiran, Department of Clinical Nutrition and Dietetics, Faculty of Nutrition Sciences and Food Technology, National Nutrition and Food Technology Research Institute, Shahid Beheshti University of Medical Sciences, Tehran 19395-4763, Iran  
Zahra Bahadoran, Nutrition and Endocrine Research Center, and Obesity Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran 19395-4763, Iran

Fereidoun Azizi, Endocrine Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran 19395-4763, Iran

Author contributions: All authors contributed equally to the manuscript.

Supported by Research Institute of Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Correspondence to: Fereidoun Azizi, MD, Endocrine Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, No 24 Parvaneh St, Yemen St, Chamran Exp, Tehran 19395-4763, Iran. [azizi@endocrine.ac.ir](mailto:azizi@endocrine.ac.ir)

Telephone: +98-21-22432500 Fax: +98-21-22416264

Received: November 24, 2013 Revised: January 11, 2014

Accepted: April 11, 2014

Published online: June 15, 2014

carbohydrate and lipid metabolism. Functional foods may also improve dyslipidemia and insulin resistance, and attenuate oxidative stress and inflammatory processes and subsequently could prevent the development of long-term diabetes complications including cardiovascular disease, neuropathy, nephropathy and retinopathy. In conclusion available data indicate that a functional foods-based diet may be a novel and comprehensive dietary approach for management of type 2 diabetes.

© 2014 Baishideng Publishing Group Inc. All rights reserved.

**Key words:** Type 2 diabetes; Insulin resistance; Functional foods; Whole grain; Legumes; Nuts; Fruits; Herbs or spices; Vegetables; Prebiotics; Probiotics

**Core tip:** Medical nutrition therapy (MNT) is a main part of type 2 diabetes management. Apparently the therapeutic and medicinal properties of foods maybe a missing step during MNT process, and could enhance the effectiveness of dietary management of type 2 diabetes.

### Abstract

Type 2 diabetes is a complicated metabolic disorder with both short- and long-term undesirable complications. In recent years, there has been growing evidence that functional foods and their bioactive compounds, due to their biological properties, may be used as complementary treatment for type 2 diabetes mellitus. In this review, we have highlighted various functional foods as missing part of medical nutrition therapy in diabetic patients. Several *in vitro*, animal models and some human studies, have demonstrated that functional foods and nutraceuticals may improve postprandial hyperglycemia and adipose tissue metabolism modulate

Mirmiran P, Bahadoran Z, Azizi F. Functional foods-based diet as a novel dietary approach for management of type 2 diabetes and its complications: A review. *World J Diabetes* 2014; 5(3): 267-281 Available from: URL: <http://www.wjgnet.com/1948-9358/full/v5/i3/267.htm> DOI: <http://dx.doi.org/10.4239/wjd.v5.i3.267>

### INTRODUCTION

Type 2 diabetes is a metabolic disorder characterized by hyperglycemia, developing insulin resistance,  $\beta$ -cell dysfunction and impaired insulin secretion<sup>[1,2]</sup>. Multiple metabolic disorders including impaired lipid and lipoprotein metabolism, oxidative stress (over production of free

radicals and defect in endogenous antioxidant defense system), sub-clinical inflammation, vascular endothelial dysfunction and hypertension are commonly accompanied by type 2 diabetes<sup>[3-5]</sup>; these metabolic disorders lead to long-term pathogenic conditions such as micro- and macro-vascular complications including neuropathy, retinopathy, nephropathy, and a decreased quality of life and an increased mortality rate<sup>[6,7]</sup>.

Despite availability of many pharmacological interventions including oral hypoglycemic agents and insulin therapy for diabetes management, current evidence shows an alarming rising trend in the occurrence of undesirable complications among these patients<sup>[1]</sup>.

Medical nutrition therapy (MNT) is also a main part of type 2 diabetes management; estimation of energy and nutrients requirements, carbohydrate counting as well as glycemic index and glycemic load, recommendation for dietary fats and cholesterol and protein intakes, explanation the foods exchange list for patients and common important recommendations for a healthy diet are the main components of diet planning in type 2 diabetic patients<sup>[8,9]</sup>; however it is not clear whether this approach *per se* is sufficiently adequate for prevention of long-term complications of diabetes. Administration of various supplements, including antioxidant vitamins, fibers,  $\omega$ 3 fatty acids, numerous nutraceuticals, and herbs has also been proposed for glycemic control but data available supporting these recommendations for diabetic patients are insufficient<sup>[10-14]</sup>. Apparently the therapeutic and medicinal properties of foods maybe a missing step during MNT process, and could enhance the effectiveness of dietary management of type 2 diabetes.

During the past two decades, the concept of functional food is fast expanding; functional foods beyond the basic nutritional functions have potential benefits to promote health and reduce the risk of chronic diseases and have hence been given much attention<sup>[15,16]</sup>. In recent years, researchers have focused on properties of the bioactive compounds of functional foods in the control of various aspects of diabetes mellitus; some protective effects of these compounds and food sources have been investigated *in vitro* and *in vivo*, and several clinical trials have even confirmed these advantages in diabetic patients<sup>[17-19]</sup>.

Here, based on the multiple biological properties of functional foods and their bioactive compounds, a functional foods-based diet has been hypothesized as a novel and comprehensive dietary approach for management of type 2 diabetes and prevention of long-term complications.

## RESEARCH

The evidence cited in this review was obtained through searches in PubMed, Scopus, and Google scholar using the following key words: "Type 2 diabetes or hyperglycemia", "insulin resistance", "cardiovascular disease", "obesity", "metabolic syndrome", "oxidative stress", "inflammation", "long-term diabetic complications" in combination with "functional foods", "nutraceuticals", "bio-

active food compounds", "fiber", "polyphenols", "whole grain", "legumes", "nuts", "fruits", "herbs or spices" "vegetables", "prebiotics", "probiotics", and "bioactive peptides". Relevant articles of acceptable quality were used. Briefly, in this article we tried to highlight some of the following important functional foods including whole grains, phytochemical-rich fruits and vegetables, legumes, nuts, dairy products, green tea and some spices, as required components of a health-promoting diet for diabetic patients.


### Whole grains

Grains and cereal-based products are the basic sources providing energy and carbohydrate in human diets. Since the dietary carbohydrate sources in type 2 diabetic patients play a determining role in glycemic and insulin secretory response, the use of functional grains including whole grain cereals, and bakery products prepared using whole wheat, rye, oat, and barley is the first step in planning of a functional foods-based diet.

Some previous studies report that dietary carbohydrate modification in patients with metabolic syndrome resulted in favorable metabolic consequences especially increased insulin sensitivity, decreased adipocyte cell size, and modulated expression of adipose tissue genes involved in insulin signaling pathways (insulin-like-growth-factor binding protein-5, insulin receptors, hormone-sensitive lipase<sup>[20,21]</sup>).

Compared to refined grains, whole grains (WGs) have more non-digestible complex polysaccharides including soluble and insoluble fibers, inulin,  $\beta$ -glucan, and resistant starches, as well as non-carbohydrate functional components including carotenoids, phytates and phytoestrogens, phenolic acids (ferulic acid, vanilic acid, caffeic acid, syringic acid, *P*-cumaric acid), and tocopherols. The most well-known protective effects of whole grain-based products against obesity, type 2 diabetes, cardiovascular diseases, hypertension, metabolic syndrome and various types of cancer, have been attributed to these bioactive compounds<sup>[22-25]</sup>. Among the several mechanisms available in current data regarding the beneficial effects of WGs and cereal-based products in diabetic patients, some of the more important are that bioactive compounds of WGs could effectively regulate glycemic response, increase insulin sensitivity, improve pancreatic  $\beta$ -cell functions and increase insulin secretion<sup>[26,27]</sup>. High contents of inulin and  $\beta$ -glucan, main soluble and fermentable fibers in WGs, in addition to their hypolipidemic and hypoglycemic effects, act as prebiotics in the gut and modulate gut microbiota *via* stimulation of growth and activity of bifidobacteria and lactic acid bacteria<sup>[28,29]</sup>, effects leading to more metabolic responses (Figure 1).

Long-term follow-ups of diabetic patients indicate that higher consumption of whole grain, cereal fiber, bran, and germ were associated with decreased all-cause and cardiovascular disease-cause mortality<sup>[30]</sup>. Epidemiological studies also confirmed that regular consumption of WGs products could modify the main risk factors of atherosclerotic diseases including triglyceride and LDL-C



**Figure 1** Role of prebiotic compounds of whole grains and cereal-based products in modulation of gut microbiota and consequent metabolic effects could lead to better glycemic control.

levels, blood pressure and serum homocysteine levels, as well as vascular functions, and oxidative and inflammatory status<sup>[31]</sup>.

Rye, a widely used grain especially in Northern and Eastern Europe, is considered a functional grain. The high fiber content of rye products decreases digestion and absorption of dietary carbohydrates, and increase metabolites derived from colonic fermentation of the soluble fiber of rye products, including propionic and butyric acids which effectively stimulate secretion of insulin from  $\beta$ -cells; studies have indicated that the bioactive compounds of rye (phenolic acids, tannins, benzoic acid, phenylalanine) derivatives have a similar efficacy with anti-diabetic drugs in insulin secretion<sup>[26,32]</sup>. In one study, the consumption of rye products in the breakfast meal increased colonic fermentation, decreased ghrelin levels and satiety rating in the late postprandial phase after breakfast as well as energy intake at a subsequent lunch meal, and improved acute glucose and insulin responses<sup>[32]</sup>.

Oat meal products have also been investigated as healthy carbohydrate sources for diabetic patients; they are rich sources of soluble fiber especially  $\beta$ -glucan, antioxidants and bioactive compounds including carotenoids, phytic acid, phenolic acids (hydroxycinnamic acids, caffeic acid, ferulic acid), flavonoids and phytosterols<sup>[33]</sup>. Studies show that consumption of oat products improves glycemic, insulinemic, and lipidemic responses in diabetic patients, and act as active ingredient reducing postprandial glycemia<sup>[34,35]</sup>. In diabetic animal models, oat products attenuated hyperglycemia-induced retinal oxidative stress, increased glycogen content of liver, decreased plasma free fatty acids and succinate dehydrogenase activity and inhibited pancreatic  $\beta$ -cell apoptosis as well<sup>[36]</sup>.

The beneficial effects of barley and its by products for diabetic patients are mainly attributed to its high content of  $\beta$ -glucan; Administration of barley  $\beta$ -glucan extract in pre-diabetic subjects improved glucose tolerance and insulin resistance index<sup>[27]</sup>. In addition, barley may use as base of a meal; the use of barley combined

with refined grains such as white rice maybe a practical way to attenuate their undesirable effects on glycemic control; in a randomized crossover study, combination of cooked barley with white rice dose-dependently reduced the area under the curves of plasma glucose and insulin concentrations, suppressed postprandial decrease of plasma desacyl ghrelin levels and consequently increased satiety<sup>[37]</sup>. The hypolipidemic properties, antioxidant and anti-inflammatory activities of barley products have also been investigated<sup>[38,39]</sup>. In animal diabetic models, barley improved some features of fatty liver, decreased lipid content of the liver, increased fatty acid oxidation and adiponectin levels<sup>[40]</sup>.

Several positive effects of whole wheat and its byproducts on carbohydrate and insulin metabolism have also been reported; wheat bran and whole wheat products are rich sources of dietary fiber, magnesium (main cofactor of enzymes involved in glucose metabolism and insulin secretion), potassium, phenolic acids,  $\alpha$ -tocopherols, carotenoids and antioxidants<sup>[41]</sup>. It is believed that the majority of beneficial effects of whole wheat grain are related to bran and germ fractions; wheat bran is a main source of fiber, lignans, phenolic acid and alkylresorcinol, and beyond the health promotion of gastrointestinal tract and weight management, could improve postprandial glycemic response, glycosylated hemoglobin, lipid disorders and other cardiovascular risk factors in diabetic patients<sup>[42]</sup>. Studies showed that alkylresorcinol of wheat bran inhibited platelet activity and aggregation, decreased triglyceride de novo synthesis, and decreased cardiovascular disease risk factors<sup>[43]</sup>. Wheat germ is rich in non-digestible oligosaccharides, phytosterols, benzoquinone and flavonoids that play a potent role in induction of antioxidant and anti-inflammatory properties and modulation of immunity responses<sup>[44]</sup>. Avemar, fermented wheat germ extract, had interesting properties in the treatment of cardiovascular disease, and improved metabolic abnormalities including hyperglycemia, lipid peroxidation and abdominal fat gain<sup>[45]</sup>.

Brown rice and its byproducts is another grain investigated as a functional food. Compared to white rice, brown rice has lower glycemic load and glycemic index, and higher content of fiber, vitamins and minerals, phytic acids, polyphenols, tocopherols, tocotrienols, and other bioactive compounds<sup>[46]</sup>; consumption of brown rice has benefits on glycemic control, dyslipidemia, endothelial function, abdominal obesity and liver functions in type 2 diabetic patients<sup>[47]</sup>. Studies show that  $\gamma$ -orizanol found in brown rice modulates high-fat diet induces oxidative stress, improves  $\beta$ -cell function, enhances glucose-stimulated insulin secretion and prevents the development of type 2 diabetes<sup>[48]</sup>. Germinated and pre-germinated brown rice, as more interesting functional foods, have unique components including  $\gamma$ -amino butyric acid, and bioactive acylated sterol glucosides with potent anti-diabetic properties; these bioactive components attenuate oxidative-induced peripheral nervous system, prevent diabetic neuropathy, inhibit oxidative-induced pancreatic  $\beta$ -cell apoptosis and enhance insulin secretion<sup>[49-51]</sup>. Bran rice, a byproduct of brown rice, contains within 31% fiber (mainly insoluble fiber),  $\beta$ -glucan, pectin, tocopherols, orizanol, ferulic acid, lutein, xanthine, vitamin K, thiamin, niacin, pantothenic acid,  $\alpha$ -lipoic acid, coenzyme Q<sub>10</sub> and other nutraceuticals; administration of bran rice in diabetic patients reduced glycosylated hemoglobin, LDL-C and total cholesterol as well as increased HDL-C<sup>[52]</sup>.

In conclusion, replacement of whole grain and cereal-based products with refined grains in diet planning may be an effective and practical strategy for MNT in type 2 diabetic patients; this approach beyond the improvement of glycemic control, leads to more benefits for management of other aspects of diabetes, attenuation of diabetes-induced metabolic disorders, and prevents long-term complications especially atherosclerosis and cardiovascular disease.

## PHYTOCHEMICAL-RICH FRUITS AND VEGETABLES

Fruits and vegetables are rich sources of dietary fiber (soluble and insoluble fiber), vitamins, and various phytochemicals and play a vital role in health promotion and prevention of chronic disease<sup>[53]</sup>. Dietary modification based on fruits and vegetables certainly is a definitely important strategy for management of type 2 diabetes and prevention of its complications; several studies indicate that regular consumption of various fruits and vegetables in diabetic patients can lead to an improved glycemic control, reduced HbA<sub>1c</sub> and triglyceride levels, enhanced antioxidant defense system, attenuated oxidative stress and inflammatory markers, decreased risk of diabetic retinopathy, and a lower burden of carotid atherosclerosis<sup>[54-57]</sup>. Since various fruits and vegetables provide many different micronutrients and bioactive compounds, consumption of varied fruits and vegetables is mainly recommended; it should be noted that the color of fruits and vegetables reflects predominant pigmented phytochemi-

cals, and considering the colors in selection of these food groups provide a wide range of nutraceuticals. In Table 1, some phytochemical-rich fruits and vegetables, their bioactive compounds and favorable effects on diabetic related conditions are reviewed. Studies showed that tomato and its by products, as main sources of lycopene,  $\beta$ -carotene, flavonoids and other bioactive components, could attenuate blood pressure and dyslipidemia, decrease cardiovascular risk factors and enhance antioxidant defense system; other sources of lycopene and carotenoids such as grapefruit and watermelon have also beneficial properties to regulate lipid and lipoprotein metabolism, blood pressure and vascular function. Anthocyanins-rich fruits including red apple, berries family, grapes, cherries, red cabbage, and pomegranate have mainly hypoglycemic effects ( $\downarrow$  digestion and absorption of dietary carbohydrates,  $\downarrow$  postprandial glycemic response and  $\downarrow$  glycosylated hemoglobin) as well as protective properties against oxidative damages (Table 1).

## LEGUMES

Legumes (peas, beans, lentils, peanuts) are valuable sources of dietary protein, non-digestible carbohydrates including dietary fiber, resistance starches, oligosaccharides, and bioactive compounds such as functional fatty acids (linoleic acid,  $\alpha$ -linolenic acid), isoflavones (daidzein, genistein, glycitein), phenolic acids, saponins, and phytic acid; some polyphenols including pelargonidin, cyanidin, delphinidin, and malvidin are also found in legumes<sup>[134,135]</sup>. Legumes are considered a component of a healthy diet and there is much evidence showing that regular consumption of legumes has protective effects against obesity, type 2 diabetes, and cardiovascular disease<sup>[136]</sup>. Legumes may be considered as an important component of a functional-foods based diet for management of type 2 diabetes.  $\alpha$ -amylase inhibitory peptides are one of the bioactive compounds in legumes and beans that reduce digestion and absorption of dietary carbohydrates, and modulate postprandial glycemic response; other bioactive peptides of grain legumes including the 7S globulin  $\alpha$  chain and conglutin  $\gamma$  have unique properties to regulate lipid metabolism and normalize lipid and lipoprotein levels<sup>[137]</sup>. Low glycemic index, high fiber and phytochemical content of legumes have made them functional food for diabetic patients.

Lentils (*Lens culinaris*), the most consumed legume grains, are rich sources of dietary fiber, slowly digestible starch and resistant starch, tannins,  $\beta$ -glucan, functional antioxidant ingredients, a wide range of phenolic acids including gallic acid, proanthocyanidins, prodelphinidin, procyanidins, catechins, epicatechin, kampferol, quercetin, cinapic acid and apigenin<sup>[138]</sup>. Studies show that bioactive proteins of lentil reduce plasma levels of LDL-C, triglyceride content of the liver, and adipose tissue lipoprotein lipase activity; moreover, polyphenols of lentil could prevent angiotensin II-induced hypertension, and pathological changes including vascular remodeling and



**Table 1 Bioactive compounds and functional properties of some of favorable fruits and vegetables**

Ref.	Possible functional properties in diabetes	Main bioactive components and phytochemicals	Fruits and vegetables
[58-62]	↓ Systolic and diastolic blood pressure ↑ apolipoprotein a1 and HDL-C ↓ LDL oxidation, improve diabetes-induced lipid disorders ↓ cardiovascular risk factors ↓ aldose reductase activity and cataract ↑ antioxidative enzymes activity	Lycopene, β-carotene, flavonoids, anthocyanins, phytoan, phyto flava, quercetin, kampferol	Tomato and its by products
[63-65]	↓ Triglyceride levels, enhance endogenous antioxidant defense system, regulation of appetite	Lycopene, pectin, naringin, hesperidin	Grapefruit
[66-69]	↑ Nitric oxide biosynthesis, improve endothelial function ↓ blood pressure ↑ plasma arginine levels and consequently ↓ insulin resistance and adipocyte size	Lycopene, carotenoids, cytolin	Watermelon
[70-73]	↓ Absorption of dietary carbohydrate ↓ postprandial glycemia, improve pancreatic β-cell function ↓ free radical generation ↓ lipid peroxidation ↑ plasma total antioxidant capacity, prevent vascular damage, improve dyslipidemia	Soluble fiber, quercetin, catechins, epicatechin, P-cumaric acid, chlorogenic acid, gallic acid, phloridizin, procyanidins	Red apple, apple peel, apple and its by products
[74-81]	Glycemic control, inhibit α-glucosidase and α-amylase activity ↓ digestion and absorption of dietary carbohydrates ↓ insulin resistance, improve dyslipidemia ↓ postprandial oxidative stress ↓ lipid peroxidation ↑ plasma total antioxidant capacity ↓ systolic blood pressure ↑ antioxidative enzymes activity ↑ adipocytes lipolysis ↓ inflammatory processes, modulation of peroxisome proliferator-activated receptors	Anthocyanins, tannins, ellagitanins, α-carotene, β-carotene, lutein, delphinidins, pelargonidins, cyanidins, catechins, hydroxy-cinnamic acid	Berries; cranberry, blackberry, black raspberry, blueberry, red raspberry, strawberries
[82-86]	Protective effects on vascular system ↓ platelet hyperactivity and aggregation ↓ cardiovascular diseases ↓ oxidative damage ↓ rennin-angiotensin activity ↑ production of nitric oxide ↓ blood pressure ↑ bone-marrow-derived endothelial progenitor cells	Anthocyanins, resveratrol	Grapes, grape by products
[87-91]	↓ Hyperglycemia ↓ HbA1c, improve lipid disorders, anti-inflammatory properties (inhibit cyclooxygenase) ↓ abdominal fat ↓ microalbuminuria, improve metabolic syndrome and fatty liver features ↓ oxidative stress ↓ production of cytokines, induction of PPARγ ↓ diabetic neuropathy	Anthocyanins, quercetin, hydroxy-cinnamic acid, carotenoids, melatonin, phenolic acids, gallic acid, lutein, xanthine, β-carotene	Cherries
[92-95]	↓ Hyperglycemia, attenuate hyperglycemia-induced metabolic disorders ↓ lipid peroxidation, induction of glutathione reductase, glutathione peroxidase, superoxide dismutase, delay progression of nephropathy ↓ inflammatory processes, improve dyslipidemia	Isothiocyanates, anthocyanins (red cabbage), carotenoids, lutein, β-carotene	Cabbage, Cauliflower
[96-100]	↓ Hyperglycemia ↑ endothelial nitric oxide synthase activity, inhibit angiotensin converting enzyme ↓ blood pressure, improve vascular function ↓ cholesterol and atherogenic lipids ↓ lipid peroxidation ↓ progression of atherosclerosis ↑ plasma total antioxidant capacity, modulate activation of PPARγ and nuclear factor κB ↑ activity of paraxonase 1 and HDL-C levels ↓ serum resistin levels and ameliorate obesity-induced insulin resistance	Anthocyanins, tannins, catechins, gallic acid, punicalagin acid, ellagic acid, gallic acid, oleanolic acid, ursolic acid, uallic acid	Pomegranate and its by products, pomegranate peel and seeds

[101-105]	↓ Hyperglycemia, induce insulin secretion from $\beta$ -cell ↓ blood pressure, inhibit enzyme involved in cholesterol biosynthesis, improve dyslipidemia, prevent atherosclerosis ↓ lipid peroxidation ↓ platelet hyperactivity and aggregation, regulate glycolysis, gluconeogenesis and carbohydrate metabolism pathways ↑ insulin sensitivity	Allyl sulfors, flavonoids, quercetin, dihydroflavonols, anthocyanins (red onion)	Garlic, onions
[106-111]	↓ Endothelial macrophage activation ↓ hyperactivity and aggregation of platelet, improve vascular function ↓ oxidative stress and inhibit stress-sensitive signaling pathways ↓ digestion of dietary lipids, improve dyslipidemia ↓ pro-inflammatory cytokines ↓ lipid peroxidation	Lutein, xanthine, $\alpha$ -cryptoxanthin, $\beta$ -cryptoxanthin, naringenin, hesperidin, $\beta$ -carotene, phytosterols	Citrus fruits
[112-113]	↓ Free radical generation and lipid peroxidation, binding to bile acids ↑ cholesterol excretion, improve lipid profile ↑ plasma total antioxidant capacity	Lutein, betaine, violaxanthine, opioid peptides (rubisculins), <i>P</i> -cumaric acid, ferulic acid	Spinach
[114-115]	Improve glycemic and insulinemic response ↓ systemic inflammation ↓ cardiovascular disease risk factors	Carotenoids, pectin, oleic and linolenic acids	Pumpkin
[116]	Improve hyperglycemia and dyslipidemia ↑ adiponectin, antioxidant and anti-inflammatory effect	Fiber, polyphenols, chlorogenic acid, flavonoids, anthocyanins	Plums
[117-119]	Improve dyslipidemia, anti-inflammatory properties ↓ lipid peroxidation ↑ plasma total antioxidant capacity	Soluble fiber (pectin), $\alpha$ -carotene, $\beta$ -carotene, lutein, phenolic acids, stilbenes	Carrots
[120-122]	Inhibit $\alpha$ -amylase ↓ postprandial glycemia ↑ glycogen synthesis, improve dyslipidemia ↓ lipid peroxidation, protective effect against diabetic nephropathy	Carotenoids, quercetin, kampferol, gallic acid, caffeic acid, catechins, tannins, mangiferin	Mango
[123-127]	Regulate carbohydrate metabolism (↑ glucokinase and glucose-6-phosphate dehydrogenase activity ↓ glucose-6-phosphatase activity) ↓ lipid peroxidation ↓ protein carbonylation ↑ antioxidant enzyme activity, improve metabolic syndrome features ↑ insulin sensitivity ↓ carbohydrate absorption ↓ plasma free fatty acid	Anthocyanins, alkaloid compounds (berberine, oxycontin)	Barberry
[128-131]	Protective effects against diabetic neuropathy ↓ lipid peroxidation, induce antioxidant enzymes, protect liver and kidney against oxidative damage	Dietary fiber, polyphenols, acid cinnamic, melatonin	Date fruit
[132-133]	Improve lipid and lipoprotein metabolism ↑ insulin sensitivity ↓ blood pressure	Dietary fiber, pectin, flavonoids, gallic acid, chlorogenic acid, catechins, anthocyanins	Figs

PPAR $\gamma$ : Peroxisome proliferator-activated receptor  $\gamma$ .

vascular fibrosis<sup>[139,140]</sup>.

Beans are also other important legume grains in the human diet with high content of fiber, phytate,  $\omega_3$  fatty acids, antioxidants, phenolic compounds. The hypoglycemic effect of beans (*via* inhibition of  $\alpha$ -amylase and  $\beta$ -glucosidase activity) has been reported as being similar to those of anti-diabetic drugs<sup>[141-143]</sup>. Including beans (pinto, dark red kidney, black beans) in diet planning for type 2 diabetic patients effectively helps weight management, attenuates postprandial glycemic response, and improves dyslipidemia<sup>[144-146]</sup>.

Soybean, a rich source of unique phytoestrogens (genistein, daidzein, glycitein), is another important functional food which has been considered in diabetes; the isoflavones and bioactive peptides of soybean have

favorable effects on glycemic control and insulin sensitivity, dyslipidemia, and kidney function<sup>[147-149]</sup>. It seems that the anti-diabetic effects of soybean mainly occur through interaction with estrogen receptors (ERs); studies show that soy isoflavones selectively bind to both  $\alpha$  and  $\beta$  estrogen receptors; ER $\alpha$  is considered as key modulator of glucose and lipid metabolism, and regulate insulin biosynthesis and secretion as well as pancreatic  $\beta$ -cell survival<sup>[150]</sup>. Soy protein could induce insulin sensitivity and improve lipid homeostasis *via* activation of peroxisome proliferator-activated receptor and liver X receptors, and inhibition of the sterol regulatory element binding protein-1c<sup>[151]</sup>. Regular consumption of soy products could help diabetic patients in the management of dyslipidemia<sup>[152]</sup>. Soy protein and isoflavones decrease

production of atherogenic apolipoproteins such as apo B, increase biosynthesis of HDL-C, induce LDL-C receptors, increase biosynthesis and excretion of bile acids, decrease gastrointestinal absorption of steroids, induce favorable changes in hormonal status, including the insulin to glucagon ratio, and thyroid hormones which lead to improvement of dyslipidemia<sup>[153,154]</sup>. Recently two bioactive peptides, identified in glycinin (a main soy protein), have unique hypolipidemic properties. These peptides inhibit 3-hydroxy-3-methyl glutaryl CoA reductase, key enzyme involved in cholesterol biosynthesis.  $\beta$ -conglycinin, another main soy bioactive protein with anti-atherogenic properties *via* regulation of lipogenesis, decrease liver lipogenic enzyme activity, inhibits fatty acid biosynthesis in liver, and facilitates fatty acid  $\beta$ -oxidation; other biological activities of soy peptides include antioxidant, anti-inflammatory, and hypotensive effect<sup>[155]</sup>.

Another feature of soybean and soy products as well as other legumes which may highlight them as main part of a functional foods-based diet, is their established effectiveness in weight management; since the overweight and obesity are the common problems in diabetic patients and main contributors in development of insulin resistance, benefit from anti-obesity properties of legumes is considered another key approach in these patients. Thermogenic effects, induction of satiety through some important appetite regulatory gut peptides, mediation in gene expression and secretion of key adipocytokines such as leptin and adiponectin, as well as inhibitory effects on proliferation and differentiation of adipocytes are some of the mechanisms that could explain the role of legumes on weight management<sup>[140,156-159]</sup>. In conclusion, considering the potential benefits of legumes and its by products, regular consumption of these functional foods may be an effective strategy for management of various aspects of type 2 diabetes.

## NUTS

Based on current evidence, nuts may play a protective effect against cardiovascular disease risk factors. Almonds, pistachios, walnuts and hazelnuts are commonly used nuts; these functional foods are considered as rich sources of high-biological value proteins, bioactive peptides, functional fatty acids (mono and poly unsaturated fatty acids), fiber, phytosterols, polyphenols, tocopherols and other antioxidant vitamins; the antioxidative effect of nuts mainly is related to a high content of  $\alpha$  and  $\gamma$  tocopherol, phenolic acids, melatonin, oleic acid and selenium, while the anti-inflammatory effect is related to ellagic acid,  $\alpha$ -linolenic acid and magnesium<sup>[160,161]</sup>.

Most current evidence reveals that consumption of nuts in type 2 diabetic patients other than improving the overall diet quality also has beneficial effects on postprandial glycemic response following high-carbohydrate meals, attenuates postprandial oxidative stress and inflammatory processes, normalizes lipid and lipoprotein levels and decreases lipid atherogenicity, and improves insulin resistance<sup>[162,163]</sup>. Moreover, habitual intake of nuts could help

to effectively manage weight especially in diabetic patients; the anti-obesity effects of nuts investigated in some studies may be attributed to thermogenic effects, induction of satiety, decreased dietary fat absorption, and increased fat excretion; bioactive components of nuts also modulate regulatory appetite neurotransmitters and adipose tissue metabolism, as well as decrease proliferation and differentiation of adipocytes, inhibit lipogenesis and induce fatty acid  $\beta$ -oxidation<sup>[164,165]</sup>. Studies show that consumption of nuts effectively decreases serum levels of high-sensitivity C-reactive protein; a well measure of systemic low-grade inflammation, interleukin 6 (a potent pro-inflammatory cytokine) and fibrinogen while increase plasma concentration of adiponectin, a potent anti-inflammatory cytokine released from adipose tissue; dietary patterns, high in nuts, were also related to lower levels of soluble inflammatory and cardiovascular risk markers including intercellular adhesion molecule 1 and vascular cell adhesion molecule 1<sup>[166,167]</sup>. Another beneficial effect of nuts which is important especially in diabetic patients is favorably influence on endothelial function; high content of L-arginine, a main precursor of nitric oxide, as well as antioxidants and polyphenols could contribute to this effect<sup>[161]</sup>.

In conclusion, it seems that a diet enriched with nuts may be an effective strategy to improve glycemic control and prevent cardiovascular disease in type 2 diabetic patients.

## OTHER BENEFICIAL FUNCTIONAL FOODS AND BIOACTIVE COMPONENTS FOR DIABETIC PATIENTS

Although there are a large number of natural foods, nutraceuticals or bioactive components that could be considered as functional ingredients and have beneficial effects for diabetes management, addressing all these issues is beyond the scope of this article. Table 2 shows some of these potential functional foods including dairy products and probiotics, fish meat, green tea, spices are presented.

## CONCLUSION

Type 2 diabetes is a complicated metabolic disorder with both short- and long-term undesirable complications as well as various pathogenic conditions including dyslipidemia, vascular dysfunction, oxidative stress, sub-clinical inflammation, and altered signaling pathways. Ineffectiveness of the current medical treatments in management of long-term diabetes complications confirms that other complementary approaches are required; the use of functional foods and bioactive compounds is one of these new approaches. Functional foods and their bioactive compounds could attenuate carbohydrate metabolism and hyperglycemia, improve pancreatic  $\beta$ -cell function and insulin secretion as well as insulin resistance, regulate lipid and lipoprotein metabolism and adipose tissue metabolism, modulate oxidative/antioxidative balance and

**Table 2 Bioactive compounds and functional properties of some of favorable functional foods**

Ref.	Possible functional properties in diabetes	Main bioactive components and nutraceuticals	Functional foods
[168-179]	Improve the features of metabolic syndrome, modulate gut microbiota, regulate satiety and food intake ↑ adiponectin, modulate adipocytokines, induce thermogenesis, lipolysis and $\beta$ -oxidation ↑ dietary fat excretion ↓ adiposity and body weight ↓ oxidative stress and inflammatory markers, hypo-lipidemic and anti-thrombotic effects ↑ insulin sensitivity, modulate immune responses in diabetic patients ↑ total antioxidant capacity ↓ lipid peroxidation ↓ HbA1c	Calcium, vitamin B, bioactive proteins such as casein and whey, immunoglobulines, bioactive peptides ( $\alpha$ - and $\beta$ -lactorphanes, lactoferrin, lactoferricin, $\alpha$ -lactalbumin, $\beta$ -lactoglobulin, growth factors), conjugated linoleic acids, lactic acid bacteria and bifidobacteria	Dairy products and probiotics
[180-185]	Improve hypertriglyceridemia and hypertension ↓ cardiovascular disease ↓ insulin resistance and inflammation, improve glycemic management ↓ proteinuria ↓ oxidative stress, inhibit lipogenesis and induce lipolysis, induce PPAR $\alpha$ and PPAR $\beta$ ↓ adiposity and weight management ↑ thermogenesis and energy expenditure, inhibit angiotensin converting enzyme and modulate blood pressure	Bioactive peptides, antioxidant compounds, $\omega$ 3 fatty acids (docosahexaenoic acid, eicosapentaenoic acid), selenium, taurine	Fish and seafood
[186-189]	Regulate cholesterol metabolism ↓ LDL oxidation, protect vascular endothelium against atherogenesis, inhibit platelet aggregation ↓ atherosclerosis development ↓ pro-inflammatory cytokines, activate PPAR $\gamma$ , improve sub-clinical inflammation	Oleic acid, $\omega$ 3 fatty acids, Flavonoids, cinnamic acid, benzoic acid, lignans, cumaric acid, ferulic acid, tocopherols, carotenoids, oleuropein, oleocanthal	Olive oil
[190-193]	Promote endogenous antioxidant defense system, induce superoxide dismutase and catalase ↓ lipid peroxidation, improve glycemic control ↑ insulin sensitivity ↓ gluconeogenesis ↑ glycogen content ↓ glycation of collagen and fibrosis, protect cardiac muscle, regulate lipid metabolism as well as adipose tissue metabolism, inhibit lipogenic enzymes ↓ satiety ↑ thermogenesis ↓ proliferation and differentiation of adipocytes ↓ pro-inflammatory cytokines ↓ monocyte chemotactic protein-1	Polyphenols, phenolic acids, catechins, epigallocatechin-3-gallat, chlorophyll, carotenoids, pectin, plant sterols	Green tea
[194-196]	↑ Insulin sensitivity, improve peripheral uptake of glucose, increase glycolysis and gluconeogenesis, hypoglycemic and hypolipidemic effects, antioxidant and anti-inflammatory properties	Cinnamaldehyde, cinnamic acid, coumarin, catechins, epicatechin, procyanidins B-2	Cinnamon
[197-199]	Inhibit enzymes involved in inflammation including cyclooxygenase-2, lipoxygenase, and nuclear factor $\kappa$ B, inhibit $\alpha$ -glucosidase and $\alpha$ -amylase activity ↓ postprandial glycemic response ↓ proteinuria, activate PPAR $\gamma$ and regulate carbohydrate and lipid metabolism, prevent diabetic cataract	Curcuminoids, stigmasterol, $\beta$ -sitosterol, 2-hydroxy methyl anthraquinone, bioactive peptide turmerin	Turmeric
[200-203]	Attenuate oxidative stress, protective effects against oxidative damage ↓ serum creatinine and urea, improve dyslipidemia ↓ atherogenic lipoprotein levels ↓ lipid peroxidation in renal tissue, inhibit $\alpha$ -glucosidase activity ↓ carbohydrate digestion and absorption, protect liver against diabetes-induced oxidative damage	Tannins, flavonoids, anthocyanins, phenolic acid, gallic acid	Sumac

PPAR: Peroxisome proliferator-activated receptor.

inflammatory processes, improve weight management and prevent micro and macro vascular complications.

Considering the beneficial properties of functional foods, it seems that diet planning based on these healthy foods may be considered an effective strategy for management of various aspects of diabetes and promotion

of health in diabetic patients.

## ACKNOWLEDGMENTS

The authors express to acknowledge the assistance given by Ms Niloofar Shiva for critical editing of English gram-



mar and syntax. None of the authors had any personal or financial conflicts of interest.

## REFERENCES

- 1 **Santaguida PL**, Balion C, Hunt D, Morrison K, Gerstein H, Raina P, Booker L, Yazdi H. Diagnosis, prognosis, and treatment of impaired glucose tolerance and impaired fasting glucose. *Evid Rep Technol Assess (Summ)* 2005; **(128)**: 1-11 [PMID: 16194123]
- 2 **Evans JL**, Goldfine ID, Maddux BA, Grodsky GM. Oxidative stress and stress-activated signaling pathways: a unifying hypothesis of type 2 diabetes. *Endocr Rev* 2002; **23**: 599-622 [PMID: 12372842 DOI: 10.1210/er.2001-0039]
- 3 **Spranger J**, Kroke A, Möhlig M, Hoffmann K, Bergmann MM, Ristow M, Boeing H, Pfeiffer AF. Inflammatory cytokines and the risk to develop type 2 diabetes: results of the prospective population-based European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam Study. *Diabetes* 2003; **52**: 812-817 [PMID: 12606524 DOI: 10.2337/diabetes.52.3.812]
- 4 **Bekyarova GY**, Ivanova DG, Madjova VH. Molecular mechanisms associating oxidative stress with endothelial dysfunction in the development of various vascular complications in diabetes mellitus. *Folia Med (Plovdiv)* 2007; **49**: 13-19 [PMID: 18504928]
- 5 **Gadi R**, Samaha FF. Dyslipidemia in type 2 diabetes mellitus. *Curr Diab Rep* 2007; **7**: 228-234 [PMID: 17547840 DOI: 10.1007/s11892-007-0036-0]
- 6 **Constantino MI**, Molyneaux L, Limacher-Gisler F, Al-Saeed A, Luo C, Wu T, Twigg SM, Yue DK, Wong J. Long-term complications and mortality in young-onset diabetes: type 2 diabetes is more hazardous and lethal than type 1 diabetes. *Diabetes Care* 2013; **36**: 3863-3869 [PMID: 23846814]
- 7 **Lloyd A**, Sawyer W, Hopkinson P. Impact of long-term complications on quality of life in patients with type 2 diabetes not using insulin. *Value Health* 2001; **4**: 392-400 [PMID: 11705130 DOI: 10.1046/j.1524-4733.2001.45029.x]
- 8 **Bantle JP**, Wylie-Rosett J, Albright AL, Apovian CM, Clark NG, Franz MJ, Hoogwerf BJ, Lichtenstein AH, Mayer-Davis E, Mooradian AD, Wheeler ML. Nutrition recommendations and interventions for diabetes: a position statement of the American Diabetes Association. *Diabetes Care* 2008; **31** Suppl 1: S61-S78 [PMID: 18165339 DOI: 10.2337/dc08-S061]
- 9 **Dämon S**, Schätzer M, Höfler J, Tomasec G, Hoppichler F. Nutrition and diabetes mellitus: an overview of the current evidence. *Wien Med Wochenschr* 2011; **161**: 282-288 [PMID: 21442215 DOI: 10.1007/s10354-011-0888-4]
- 10 **Perera PK**, Li Y. Functional herbal food ingredients used in type 2 diabetes mellitus. *Pharmacogn Rev* 2012; **6**: 37-45 [PMID: 22654403 DOI: 10.4103/0973-7847.95863]
- 11 **Bahadoran Z**, Mirmiran P, Hosseinpour F, Hedayati M, Hosseinpour-Niazi S, Azizi F. Broccoli sprouts reduce oxidative stress in type 2 diabetes: a randomized double-blind clinical trial. *Eur J Clin Nutr* 2011; **65**: 972-977 [PMID: 21559038 DOI: 10.1038/ejcn.2011.59]
- 12 **Bahadoran Z**, Mirmiran P, Hosseinpour F, Rajab A, Asghari G, Azizi F. Broccoli sprouts powder could improve serum triglyceride and oxidized LDL/LDL-cholesterol ratio in type 2 diabetic patients: a randomized double-blind placebo-controlled clinical trial. *Diabetes Res Clin Pract* 2012; **96**: 348-354 [PMID: 22325157 DOI: 10.1016/j.diabres.2012.01.009]
- 13 **Bahadoran Z**, Tohidi M, Nazeri P, Mehran M, Azizi F, Mirmiran P. Effect of broccoli sprouts on insulin resistance in type 2 diabetic patients: a randomized double-blind clinical trial. *Int J Food Sci Nutr* 2012; **63**: 767-771 [PMID: 22537070 DOI: 10.3109/09637486.2012.665043]
- 14 **Kris-Etherton PM**, Lichtenstein AH, Howard BV, Steinberg D, Witztum JL. Antioxidant vitamin supplements and cardiovascular disease. *Circulation* 2004; **110**: 637-641 [PMID: 15289389 DOI: 10.1161/01.CIR.0000137822.39831.F1]
- 15 **Ballali S**, Lanci F. Functional food and diabetes: a natural way in diabetes prevention? *Int J Food Sci Nutr* 2012; **63** Suppl 1: 51-61 [PMID: 22107597 DOI: 10.3109/09637486.2011.637487]
- 16 **Gregori D**, Gafare CE. Multifunctional food: medical evidence and methodological notes on substantiating health claims. *Int J Food Sci Nutr* 2012; **63** Suppl 1: 29-36 [PMID: 22289058 DOI: 10.3109/09637486.2011.653553]
- 17 **Rudkowska I**. Functional foods for health: focus on diabetes. *Maturitas* 2009; **62**: 263-269 [PMID: 19233576 DOI: 10.1016/j.maturitas.2009.01.011]
- 18 **Bahadoran Z**, Mirmiran P, Azizi F. Dietary polyphenols as potential nutraceuticals in management of diabetes: a review. *J Diabetes Metab Disord* 2013; **12**: 43 [PMID: 23938049 DOI: 10.1186/2251-6581-12-43]
- 19 **Bahadoran Z**, Mirmiran P, Azizi F. Potential efficacy of broccoli sprouts as a unique supplement for management of type 2 diabetes and its complications. *J Med Food* 2013; **16**: 375-382 [PMID: 23631497 DOI: 10.1089/jmf.2012.2559]
- 20 **Laaksonen DE**, Toppinen LK, Juntunen KS, Autio K, Liukkonen KH, Poutanen KS, Niskanen L, Mykkanen HM. Dietary carbohydrate modification enhances insulin secretion in persons with the metabolic syndrome. *Am J Clin Nutr* 2005; **82**: 1218-1227 [PMID: 16332654]
- 21 **Kallio P**, Kolehmainen M, Laaksonen DE, Kekäläinen J, Salopuro T, Sivenius K, Pulkkinen L, Mykkanen HM, Niskanen L, Uusitupa M, Poutanen KS. Dietary carbohydrate modification induces alterations in gene expression in abdominal subcutaneous adipose tissue in persons with the metabolic syndrome: the FUNGENT Study. *Am J Clin Nutr* 2007; **85**: 1417-1427 [PMID: 17490981]
- 22 **Borneo R**, León AE. Whole grain cereals: functional components and health benefits. *Food Funct* 2012; **3**: 110-119 [PMID: 22134555 DOI: 10.1039/c1fo10165j]
- 23 **Ye EQ**, Chacko SA, Chou EL, Kugizaki M, Liu S. Greater whole-grain intake is associated with lower risk of type 2 diabetes, cardiovascular disease, and weight gain. *J Nutr* 2012; **142**: 1304-1313 [PMID: 22649266 DOI: 10.3945/jn.111.155325]
- 24 **Venn BJ**, Mann JI. Cereal grains, legumes and diabetes. *Eur J Clin Nutr* 2004; **58**: 1443-1461 [PMID: 15162131 DOI: 10.1038/sj.ejcn.1601995]
- 25 **Okarter N**, Liu RH. Health benefits of whole grain phytochemicals. *Crit Rev Food Sci Nutr* 2010; **50**: 193-208 [PMID: 20301011 DOI: 10.1080/10408390802248734]
- 26 **Rosén LA**, Silva LO, Andersson UK, Holm C, Ostman EM, Björck IM. Endosperm and whole grain rye breads are characterized by low post-prandial insulin response and a beneficial blood glucose profile. *Nutr J* 2009; **8**: 42 [PMID: 19781071 DOI: 10.1186/1475-2891-8-42]
- 27 **Shen RL**, Cai FL, Dong JL, Hu XZ. Hypoglycemic effects and biochemical mechanisms of oat products on streptozotocin-induced diabetic mice. *J Agric Food Chem* 2011; **59**: 8895-8900 [PMID: 21749041 DOI: 10.1021/jf200678q]
- 28 **Martínez I**, Lattimer JM, Hubach KL, Case JA, Yang J, Weber CG, Louk JA, Rose DJ, Kyureghian G, Peterson DA, Haub MD, Walter J. Gut microbiome composition is linked to whole grain-induced immunological improvements. *ISME J* 2013; **7**: 269-280 [PMID: 23038174 DOI: 10.1038/ismej.2012.104]
- 29 **Andersson A**, Tengblad S, Karlström B, Kamal-Eldin A, Landberg R, Basu S, Aman P, Vessby B. Whole-grain foods do not affect insulin sensitivity or markers of lipid peroxidation and inflammation in healthy, moderately overweight subjects. *J Nutr* 2007; **137**: 1401-1407 [PMID: 17513398]
- 30 **He M**, van Dam RM, Rimm E, Hu FB, Qi L. Whole-grain, cereal fiber, bran, and germ intake and the risks of all-cause and cardiovascular disease-specific mortality among women with type 2 diabetes mellitus. *Circulation* 2010;

- 121: 2162-2168 [PMID: 20458012 DOI: 10.1161/CIRCULATIONAHA.109.907360]
- 31 **Anderson JW.** Whole grains protect against atherosclerotic cardiovascular disease. *Proc Nutr Soc* 2003; **62**: 135-142 [PMID: 12740068 DOI: 10.1079/PNS2002222]
  - 32 **Rosén LA, Ostman EM, Björck IM.** Effects of cereal breakfasts on postprandial glucose, appetite regulation and voluntary energy intake at a subsequent standardized lunch; focusing on rye products. *Nutr J* 2011; **10**: 7 [PMID: 21247415 DOI: 10.1186/1475-2891-10-7]
  - 33 **Sadiq Butt M, Tahir-Nadeem M, Khan MK, Shabir R, Butt MS.** Oat: unique among the cereals. *Eur J Nutr* 2008; **47**: 68-79 [PMID: 18301937 DOI: 10.1007/s00394-008-0698-7]
  - 34 **VandenLangenberg GM, Brady WE, Nebeling LC, Block G, Forman M, Bowen PE, Stacewicz-Sapuntzakis M, Mares-Perlman JA.** Influence of using different sources of carotenoid data in epidemiologic studies. *J Am Diet Assoc* 1996; **96**: 1271-1275 [PMID: 8948389 DOI: 10.1016/S0002-8223(96)00329-X]
  - 35 **Tapola N, Karvonen H, Niskanen L, Mikola M, Sarkkinen E.** Glycemic responses of oat bran products in type 2 diabetic patients. *Nutr Metab Cardiovasc Dis* 2005; **15**: 255-261 [PMID: 16054549 DOI: 10.1016/j.numecd.2004.09.003]
  - 36 **Bays H, Frestedt JL, Bell M, Williams C, Kolberg L, Schmelzer W, Anderson JW.** Reduced viscosity Barley  $\beta$ -Glucan versus placebo: a randomized controlled trial of the effects on insulin sensitivity for individuals at risk for diabetes mellitus. *Nutr Metab (Lond)* 2011; **8**: 58 [PMID: 21846371 DOI: 10.1186/1743-7075-8-58]
  - 37 **Sakuma M, Yamanaka-Okumura H, Naniwa Y, Matsumoto D, Tsunematsu M, Yamamoto H, Taketani Y, Takeda E.** Dose-dependent effects of barley cooked with white rice on postprandial glucose and desacyl ghrelin levels. *J Clin Biochem Nutr* 2009; **44**: 151-159 [PMID: 19308269 DOI: 10.3164/jcbn.08-232]
  - 38 **Gremmel H, Konrad RM.** [Postoperative results in alloplastic closure of diaphragmatic defects]. *Fortschr Geb Rontgenstr Nuklearmed* 1964; **100**: 703-710 [PMID: 14252728]
  - 39 **Kofuji K, Aoki A, Tsubaki K, Konishi M, Isobe T, Murata Y.** Antioxidant Activity of  $\beta$ -Glucan. *ISRN Pharm* 2012; **2012**: 125864 [PMID: 22500243]
  - 40 **Brockman DA, Chen X, Gallaher DD.** Consumption of a high  $\beta$ -glucan barley flour improves glucose control and fatty liver and increases muscle acylcarnitines in the Zucker diabetic fatty rat. *Eur J Nutr* 2013; **52**: 1743-1753 [PMID: 23229409]
  - 41 **Whent M, Huang H, Xie Z, Lutterodt H, Yu L, Fuerst EP, Morris CF, Yu LL, Luthria D.** Phytochemical composition, anti-inflammatory, and antiproliferative activity of whole wheat flour. *J Agric Food Chem* 2012; **60**: 2129-2135 [PMID: 22321109 DOI: 10.1021/jf203807w]
  - 42 **Stevenson L, Phillips F, O'Sullivan K, Walton J.** Wheat bran: its composition and benefits to health, a European perspective. *Int J Food Sci Nutr* 2012; **63**: 1001-1013 [PMID: 22716911 DOI: 10.3109/09637486.2012.687366]
  - 43 **Ross AB, Kamal-Eldin A, Aman P.** Dietary alkylresorcinols: absorption, bioactivities, and possible use as biomarkers of whole-grain wheat- and rye-rich foods. *Nutr Rev* 2004; **62**: 81-95 [PMID: 15098855 DOI: 10.1111/j.1753-4887.2004.tb00029.x]
  - 44 **Brandolini A, Hidalgo A.** Wheat germ: not only a by-product. *Int J Food Sci Nutr* 2012; **63** Suppl 1: 71-74 [PMID: 22077851 DOI: 10.3109/09637486.2011.633898]
  - 45 **Iyer A, Brown L.** Fermented wheat germ extract (avemar) in the treatment of cardiac remodeling and metabolic symptoms in rats. *Evid Based Complement Alternat Med* 2011; **2011**: 508957 [PMID: 19622599]
  - 46 **Callegaro Mda D, Tirapegui J.** [Comparison of the nutritional value between brown rice and white rice]. *Arq Gastroenterol* 1996; **33**: 225-231 [PMID: 9302338]
  - 47 **Shimabukuro M, Higa M, Kinjo R, Yamakawa K, Tanaka H, Kozuka C, Yabiku K, Taira S, Sata M, Masuzaki H.** Effects of the brown rice diet on visceral obesity and endothelial function: the BRAVO study. *Br J Nutr* 2014; **111**: 310-320 [PMID: 23930929]
  - 48 **Zhang P, Tang H, Chen K, Chen Y, Xu D.** Biological variations of hematologic parameters determined by UniCel DxH 800 hematology analyzer. *Arch Pathol Lab Med* 2013; **137**: 1106-1110 [PMID: 23899069 DOI: 10.1186/1472-6882-13-198]
  - 49 **Kozuka C, Yabiku K, Takayama C, Matsushita M, Shimabukuro M.** Natural food science based novel approach toward prevention and treatment of obesity and type 2 diabetes: recent studies on brown rice and  $\gamma$ -oryzanol. *Obes Res Clin Pract* 2013; **7**: e165-e172 [PMID: 23697584 DOI: 10.1016/j.orcp.2013.02.003]
  - 50 **Torimitsu M, Nagase R, Yanagi M, Homma M, Sasai Y, Ito Y, Hayamizu K, Nonaka S, Hosono T, Kise M, Seki T, Ariga T.** Replacing white rice with pre-germinated brown rice mildly ameliorates hyperglycemia and imbalance of adipocytokine levels in type 2 diabetes model rats. *J Nutr Sci Vitaminol (Tokyo)* 2010; **56**: 287-292 [PMID: 21228498 DOI: 10.3177/jnsv.56.287]
  - 51 **Usuki S, Tsai YY, Morikawa K, Nonaka S, Okuhara Y, Kise M, Yu RK.** IGF-1 induction by acylated steryl  $\beta$ -glucosides found in a pre-germinated brown rice diet reduces oxidative stress in streptozotocin-induced diabetes. *PLoS One* 2011; **6**: e28693 [PMID: 22194889 DOI: 10.1371/journal.pone.0028693]
  - 52 **Cheng HH, Huang HY, Chen YY, Huang CL, Chang CJ, Chen HL, Lai MH.** Ameliorative effects of stabilized rice bran on type 2 diabetes patients. *Ann Nutr Metab* 2010; **56**: 45-51 [PMID: 20016147 DOI: 10.1159/000265850]
  - 53 **Heber D.** Vegetables, fruits and phytoestrogens in the prevention of diseases. *J Postgrad Med* 2004; **50**: 145-149 [PMID: 15235216]
  - 54 **Hegde SV, Adhikari P, M N, D'Souza V.** Effect of daily supplementation of fruits on oxidative stress indices and glycaemic status in type 2 diabetes mellitus. *Complement Ther Clin Pract* 2013; **19**: 97-100 [PMID: 23561067 DOI: 10.1016/j.ctcp.2012.12.002]
  - 55 **Tanaka S, Yoshimura Y, Kawasaki R, Kamada C, Tanaka S, Horikawa C, Ohashi Y, Araki A, Ito H, Akanuma Y, Yamada N, Yamashita H, Sone H.** Fruit intake and incident diabetic retinopathy with type 2 diabetes. *Epidemiology* 2013; **24**: 204-211 [PMID: 23348071 DOI: 10.1097/EDE.0b013e318281725e]
  - 56 **Takahashi K, Kamada C, Yoshimura H, Okumura R, Iimuro S, Ohashi Y, Araki A, Umegaki H, Sakurai T, Yoshimura Y, Ito H.** Effects of total and green vegetable intakes on glycated hemoglobin A1c and triglycerides in elderly patients with type 2 diabetes mellitus: the Japanese Elderly Intervention Trial. *Geriatr Gerontol Int* 2012; **12** Suppl 1: 50-58 [PMID: 22435940 DOI: 10.1111/j.1447-0594.2011.00812.x]
  - 57 **Chan HT, Yiu KH, Wong CY, Li SW, Tam S, Tse HF.** Increased dietary fruit intake was associated with lower burden of carotid atherosclerosis in Chinese patients with Type 2 diabetes mellitus. *Diabet Med* 2013; **30**: 100-108 [PMID: 22913415 DOI: 10.1111/j.1464-5491.2012.03764.x]
  - 58 **Ali MM, Agha FG.** Amelioration of streptozotocin-induced diabetes mellitus, oxidative stress and dyslipidemia in rats by tomato extract lycopene. *Scand J Clin Lab Invest* 2009; **69**: 371-379 [PMID: 19148834 DOI: 10.1080/00365510802658473]
  - 59 **Canene-Adams K, Campbell JK, Zaripheh S, Jeffery EH, Erdman JW.** The tomato as a functional food. *J Nutr* 2005; **135**: 1226-1230 [PMID: 15867308]
  - 60 **Pollack A, Oren P, Stark AH, Eisner Z, Nyska A, Madar Z.** Cataract development in sand and galactosemic rats fed a natural tomato extract. *J Agric Food Chem* 1999; **47**: 5122-5126 [PMID: 10606583 DOI: 10.1021/jf9900231]
  - 61 **Shidfar F, Froghifar N, Vafa M, Rajab A, Hosseini S, Shidfar S, Gohari M.** The effects of tomato consumption on serum

- glucose, apolipoprotein B, apolipoprotein A-I, homocysteine and blood pressure in type 2 diabetic patients. *Int J Food Sci Nutr* 2011; **62**: 289-294 [PMID: 21138408 DOI: 10.3109/09637486.2010.529072]
- 62 **Upritchard JE**, Sutherland WH, Mann JI. Effect of supplementation with tomato juice, vitamin E, and vitamin C on LDL oxidation and products of inflammatory activity in type 2 diabetes. *Diabetes Care* 2000; **23**: 733-738 [PMID: 10840987 DOI: 10.2337/diacare.23.6.733]
  - 63 **Goldwasser J**, Cohen PY, Yang E, Balaguer P, Yarmush ML, Nahmias Y. Transcriptional regulation of human and rat hepatic lipid metabolism by the grapefruit flavonoid naringenin: role of PPARalpha, PPARgamma and LXRalpha. *PLoS One* 2010; **5**: e12399 [PMID: 20811644 DOI: 10.1371/journal.pone.0012399]
  - 64 **Gorinstein S**, Caspi A, Libman I, Lerner HT, Huang D, Leontowicz H, Leontowicz M, Tashma Z, Katrich E, Feng S, Trakhtenberg S. Red grapefruit positively influences serum triglyceride level in patients suffering from coronary atherosclerosis: studies in vitro and in humans. *J Agric Food Chem* 2006; **54**: 1887-1892 [PMID: 16506849 DOI: 10.1021/jf058171g]
  - 65 **Owira PM**, Ojewole JA. The grapefruit: an old wine in a new glass? Metabolic and cardiovascular perspectives. *Cardiovasc J Afr* 2010; **21**: 280-285 [PMID: 20972517]
  - 66 **Wu G**, Collins JK, Perkins-Veazie P, Siddiq M, Dolan KD, Kelly KA, Heaps CL, Meininger CJ. Dietary supplementation with watermelon pomace juice enhances arginine availability and ameliorates the metabolic syndrome in Zucker diabetic fatty rats. *J Nutr* 2007; **137**: 2680-2685 [PMID: 18029483]
  - 67 **Figueroa A**, Sanchez-Gonzalez MA, Perkins-Veazie PM, Arjmandi BH. Effects of watermelon supplementation on aortic blood pressure and wave reflection in individuals with prehypertension: a pilot study. *Am J Hypertens* 2011; **24**: 40-44 [PMID: 20616787 DOI: 10.1038/ajh.2010.142]
  - 68 **Fatema K**, Habib B, Afza N, Ali L. Glycemic, non-esterified fatty acid (NEFA) and insulinemic responses to watermelon and apple in type 2 diabetic subjects. *Asia Pac J Clin Nutr* 2003; **12** Suppl: S53 [PMID: 15023673]
  - 69 **Martins MJ**, Negrão MR, Azevedo I. Watermelon: the value of higher plasma arginine concentrations. *Nutrition* 2007; **23**: 517 [PMID: 17467236 DOI: 10.1016/j.nut.2007.03.007]
  - 70 **Setorki M**, Asgary S, Eidi A, Rohani AH, Esmail N. Effects of apple juice on risk factors of lipid profile, inflammation and coagulation, endothelial markers and atherosclerotic lesions in high cholesterolemic rabbits. *Lipids Health Dis* 2009; **8**: 39 [PMID: 19804641 DOI: 10.1186/1476-511X-8-39]
  - 71 **Boyer J**, Liu RH. Apple phytochemicals and their health benefits. *Nutr J* 2004; **3**: 5 [PMID: 15140261]
  - 72 **Chai SC**, Hooshmand S, Saadat RL, Payton ME, Brummel-Smith K, Arjmandi BH. Daily apple versus dried plum: impact on cardiovascular disease risk factors in postmenopausal women. *J Acad Nutr Diet* 2012; **112**: 1158-1168 [PMID: 22818725 DOI: 10.1016/j.jand.2012.05.005]
  - 73 **Hyson DA**. A comprehensive review of apples and apple components and their relationship to human health. *Adv Nutr* 2011; **2**: 408-420 [PMID: 22332082 DOI: 10.3945/an.111.000513]
  - 74 **Kaume L**, Howard LR, Devareddy L. The blackberry fruit: a review on its composition and chemistry, metabolism and bioavailability, and health benefits. *J Agric Food Chem* 2012; **60**: 5716-5727 [PMID: 22082199]
  - 75 **Seymour EM**, Tanone II, Urcuyo-Llanes DE, Lewis SK, Kirakosyan A, Kondoleon MG, Kaufman PB, Bolling SF. Blueberry intake alters skeletal muscle and adipose tissue peroxisome proliferator-activated receptor activity and reduces insulin resistance in obese rats. *J Med Food* 2011; **14**: 1511-1518 [PMID: 21861718 DOI: 10.1089/jmf.2010.0292]
  - 76 **Suh JH**, Romain C, González-Barrio R, Cristol JP, Teissèdre PL, Crozier A, Rouanet JM. Raspberry juice consumption, oxidative stress and reduction of atherosclerosis risk factors in hypercholesterolemic golden Syrian hamsters. *Food Funct* 2011; **2**: 400-405 [PMID: 21894327 DOI: 10.1039/c1fo10047e]
  - 77 **Apostolidis E**, Kwon YI, Shetty K. Potential of cranberry-based herbal synergies for diabetes and hypertension management. *Asia Pac J Clin Nutr* 2006; **15**: 433-441 [PMID: 16837438]
  - 78 **Basu A**, Fu DX, Wilkinson M, Simmons B, Wu M, Betts NM, Du M, Lyons TJ. Strawberries decrease atherosclerotic markers in subjects with metabolic syndrome. *Nutr Res* 2010; **30**: 462-469 [PMID: 20797478 DOI: 10.1016/j.nutres.2010.06.016]
  - 79 **Chen CF**, Li YD, Xu Z. [Chemical principles and bioactivities of blueberry]. *Yaoxue Xuebao* 2010; **45**: 422-429 [PMID: 21355205]
  - 80 **Morimoto C**, Satoh Y, Hara M, Inoue S, Tsujita T, Okuda H. Anti-obese action of raspberry ketone. *Life Sci* 2005; **77**: 194-204 [PMID: 15862604 DOI: 10.1016/j.lfs.2004.12.029]
  - 81 **Basu A**, Rhone M, Lyons TJ. Berries: emerging impact on cardiovascular health. *Nutr Rev* 2010; **68**: 168-177 [PMID: 20384847 DOI: 10.1111/j.1753-4887.2010.00273.x]
  - 82 **Castilla P**, Echarri R, Dávalos A, Cerrato F, Ortega H, Teruel JL, Lucas MF, Gómez-Coronado D, Ortuño J, Lasunción MA. Concentrated red grape juice exerts antioxidant, hypolipidemic, and antiinflammatory effects in both hemodialysis patients and healthy subjects. *Am J Clin Nutr* 2006; **84**: 252-262 [PMID: 16825703]
  - 83 **Felice F**, Zambito Y, Di Colo G, D'Onofrio C, Fausto C, Balbarini A, Di Stefano R. Red grape skin and seeds polyphenols: Evidence of their protective effects on endothelial progenitor cells and improvement of their intestinal absorption. *Eur J Pharm Biopharm* 2012; **80**: 176-184 [PMID: 21971368 DOI: 10.1016/j.ejpb.2011.09.002]
  - 84 **Gollücke AP**, Ribeiro DA. Use of grape polyphenols for promoting human health: a review of patents. *Recent Pat Food Nutr Agric* 2012; **4**: 26-30 [PMID: 22316271 DOI: 10.2174/1876142911204010026]
  - 85 **Leifert WR**, Abeywardena MY. Cardioprotective actions of grape polyphenols. *Nutr Res* 2008; **28**: 729-737 [PMID: 19083481 DOI: 10.1016/j.nutres.2008.08.007]
  - 86 **Vislocky LM**, Fernandez ML. Biomedical effects of grape products. *Nutr Rev* 2010; **68**: 656-670 [PMID: 20961296 DOI: 10.1111/j.1753-4887.2010.00335.x]
  - 87 **Kim DO**, Heo HJ, Kim YJ, Yang HS, Lee CY. Sweet and sour cherry phenolics and their protective effects on neuronal cells. *J Agric Food Chem* 2005; **53**: 9921-9927 [PMID: 16366675 DOI: 10.1021/jf0518599]
  - 88 **Seymour EM**, Lewis SK, Urcuyo-Llanes DE, Tanone II, Kirakosyan A, Kaufman PB, Bolling SF. Regular tart cherry intake alters abdominal adiposity, adipose gene transcription, and inflammation in obesity-prone rats fed a high fat diet. *J Med Food* 2009; **12**: 935-942 [PMID: 19857054 DOI: 10.1089/jmf.2008.0270]
  - 89 **Traustadóttir T**, Davies SS, Stock AA, Su Y, Heward CB, Roberts LJ, Harman SM. Tart cherry juice decreases oxidative stress in healthy older men and women. *J Nutr* 2009; **139**: 1896-1900 [PMID: 19692530 DOI: 10.3945/jn.109.111716]
  - 90 **Ataie-Jafari A**, Hosseini S, Karimi F. Effects of sour cherry juice on blood glucose and some cardiovascular risk factors improvements in diabetic women: A pilot study. *Nutr Food Sci* 2008; **38**: 355-360 [DOI: 10.1108/00346650810891414]
  - 91 **Zhou Z**, Nair MG, Claycombe KJ. Synergistic inhibition of interleukin-6 production in adipose stem cells by tart cherry anthocyanins and atorvastatin. *Phytomedicine* 2012; **19**: 878-881 [PMID: 22703874 DOI: 10.1016/j.phymed.2012.05.001]
  - 92 **Islam MS**, Choi H. Antidiabetic effect of Korean traditional Baechu (Chinese cabbage) kimchi in a type 2 diabetes model of rats. *J Med Food* 2009; **12**: 292-297 [PMID: 19459728 DOI: 10.1089/jmf.2008.0181]
  - 93 **Kataya HA**, Hamza AA. Red Cabbage (*Brassica oleracea*) Ameliorates Diabetic Nephropathy in Rats. *Evid Based Complement Alternat Med* 2008; **5**: 281-287 [PMID: 18830445 DOI: 10.1093/ecam/nem029]



- 94 **Sankhari JM**, Thounaojam MC, Jadeja RN, Devkar RV, Ramachandran AV. Anthocyanin-rich red cabbage (*Brassica oleracea* L.) extract attenuates cardiac and hepatic oxidative stress in rats fed an atherogenic diet. *J Sci Food Agric* 2012; **92**: 1688-1693 [PMID: 22228433 DOI: 10.1002/jsfa.5532]
- 95 **Dinkova-Kostova AT**, Kostov RV. Glucosinolates and isothiocyanates in health and disease. *Trends Mol Med* 2012; **18**: 337-347 [PMID: 22578879 DOI: 10.1016/j.molmed.2012.04.003]
- 96 **Aviram M**, Dornfeld L. Pomegranate juice consumption inhibits serum angiotensin converting enzyme activity and reduces systolic blood pressure. *Atherosclerosis* 2001; **158**: 195-198 [PMID: 11500191 DOI: 10.1016/S0021-9150(01)00412-9]
- 97 **Basu A**, Penugonda K. Pomegranate juice: a heart-healthy fruit juice. *Nutr Rev* 2009; **67**: 49-56 [PMID: 19146506 DOI: 10.1111/j.1753-4887.2008.00133.x]
- 98 **Banihani S**, Swedan S, Alguraan Z. Pomegranate and type 2 diabetes. *Nutr Res* 2013; **33**: 341-348 [DOI: 10.1016/j.nutres.2013.03.003]
- 99 **Mathew AS**, Capel-Williams GM, Berry SE, Hall WL. Acute effects of pomegranate extract on postprandial lipaemia, vascular function and blood pressure. *Plant Foods Hum Nutr* 2012; **67**: 351-357 [PMID: 23093401 DOI: 10.1007/s11130-012-0318-9]
- 100 **Medjakovic S**, Jungbauer A. Pomegranate: a fruit that ameliorates metabolic syndrome. *Food Funct* 2013; **4**: 19-39 [PMID: 23060097 DOI: 10.1039/c2fo30034f]
- 101 **Slimestad R**, Fossen T, Vågen IM. Onions: a source of unique dietary flavonoids. *J Agric Food Chem* 2007; **55**: 10067-10080 [PMID: 17997520 DOI: 10.1021/jf0712503]
- 102 **Taj Eldin IM**, Ahmed EM, Elwahab H M A. Preliminary Study of the Clinical Hypoglycemic Effects of *Allium cepa* (Red Onion) in Type 1 and Type 2 Diabetic Patients. *Environ Health Insights* 2010; **4**: 71-77 [PMID: 21079693]
- 103 **Gabler NK**, Osrowska E, Imsic M, Eagling DR, Jois M, Tatham BG, Dunshea FR. Dietary onion intake as part of a typical high fat diet improves indices of cardiovascular health using the mixed sex pig model. *Plant Foods Hum Nutr* 2006; **61**: 179-185 [PMID: 17123162 DOI: 10.1007/s11130-006-0030-8]
- 104 **Ashraf R**, Aamir K, Shaikh AR, Ahmed T. Effects of garlic on dyslipidemia in patients with type 2 diabetes mellitus. *J Ayub Med Coll Abbottabad* 2005; **17**: 60-64 [PMID: 16320801]
- 105 **Ashraf R**, Khan RA, Ashraf I. Garlic (*Allium sativum*) supplementation with standard antidiabetic agent provides better diabetic control in type 2 diabetes patients. *Pak J Pharm Sci* 2011; **24**: 565-570 [PMID: 21959822]
- 106 **Yamada T**, Hayasaka S, Shibata Y, Ojima T, Saegusa T, Gotoh T, Ishikawa S, Nakamura Y, Kayaba K. Frequency of citrus fruit intake is associated with the incidence of cardiovascular disease: the Jichi Medical School cohort study. *J Epidemiol* 2011; **21**: 169-175 [PMID: 21389640 DOI: 10.2188/jea.JE20100084]
- 107 **Ramful D**, Tarnus E, Rondeau P, Robert Da Silva C, Bahr-run T, Bourdon E. Citrus Fruit Extracts Reduce Advanced Glycation End Products (AGEs)- and H<sub>2</sub>O<sub>2</sub>-Induced Oxidative Stress in Human Adipocytes. *J Agric Food Chem* 2010; **58**: 11119-11129 [PMID: 20882960]
- 108 **Mulvihill EE**, Huff MW. Citrus flavonoids and the prevention of atherosclerosis. *Cardiovasc Hematol Disord Drug Targets* 2012; **12**: 84-91 [PMID: 23030447 DOI: 10.2174/1871529X11202020084]
- 109 **Buscemi S**, Rosafio G, Arcoleo G, Mattina A, Canino B, Montana M, Verga S, Rini G. Effects of red orange juice intake on endothelial function and inflammatory markers in adult subjects with increased cardiovascular risk. *Am J Clin Nutr* 2012; **95**: 1089-1095 [PMID: 22492368 DOI: 10.3945/ajcn.111.031088]
- 110 **Chanet A**, Milenkovic D, Manach C, Mazur A, Morand C. Citrus flavanones: what is their role in cardiovascular protection? *J Agric Food Chem* 2012; **60**: 8809-8822 [PMID: 22574825 DOI: 10.1021/jf300669s]
- 111 **Assini JM**, Mulvihill EE, Huff MW. Citrus flavonoids and lipid metabolism. *Curr Opin Lipidol* 2013; **24**: 34-40 [PMID: 23254473 DOI: 10.1097/MOL.0b013e32835c07df]
- 112 **Bergman M**, Varshavsky L, Gottlieb HE, Grossman S. The antioxidant activity of aqueous spinach extract: chemical identification of active fractions. *Phytochemistry* 2001; **58**: 143-152 [PMID: 11524124 DOI: 10.1016/S0031-9422(01)00137-6]
- 113 **Moser B**, Szekeres T, Bieglmayer C, Wagner KH, Mišák M, Kundi M, Zakerska O, Nersesyan A, Kager N, Zahrl J, Hoelzl C, Ehrlich V, Knasmueller S. Impact of spinach consumption on DNA stability in peripheral lymphocytes and on biochemical blood parameters: results of a human intervention trial. *Eur J Nutr* 2011; **50**: 587-594 [PMID: 21384253 DOI: 10.1007/s00394-011-0167-6]
- 114 **Yadav M**, Jain S, Tomar R, Prasad GB, Yadav H. Medicinal and biological potential of pumpkin: an updated review. *Nutr Res Rev* 2010; **23**: 184-190 [PMID: 21110905 DOI: 10.1017/S0954422410000107]
- 115 **Yoshinari O**, Sato H, Igarashi K. Anti-diabetic effects of pumpkin and its components, trigonelline and nicotinic acid, on Goto-Kakizaki rats. *Biosci Biotechnol Biochem* 2009; **73**: 1033-1041 [PMID: 19420712 DOI: 10.1271/bbb.80805]
- 116 **Utsunomiya H**, Yamakawa T, Kamei J, Kadonosono K, Tanaka S. Anti-hyperglycemic effects of plum in a rat model of obesity and type 2 diabetes, Wistar fatty rat. *Biomed Res* 2005; **26**: 193-200 [PMID: 16295695 DOI: 10.2220/biomed-res.26.193]
- 117 **Potter AS**, Foroudi S, Stamatikos A, Patil BS, Deyhim F. Drinking carrot juice increases total antioxidant status and decreases lipid peroxidation in adults. *Nutr J* 2011; **10**: 96 [PMID: 21943297 DOI: 10.1186/1475-2891-10-96]
- 118 **Ramezani A**, Tahbaz F, Rasooli Sh, Rashidkhani B, Gharavi Noori A, Moslemi N, Hedayati M. Effects of  $\beta$ -carotene-fortified carrot juice on the lipid profile in type-2 diabetes patients. *Iranian Journal of Nutrition Sciences and Food Technology* 2010; **5**: 57-66
- 119 **Metzger BT**, Barnes DM, Reed JD. Purple carrot (*Daucus carota* L.) polyacetylenes decrease lipopolysaccharide-induced expression of inflammatory proteins in macrophage and endothelial cells. *J Agric Food Chem* 2008; **56**: 3554-3560 [PMID: 18433135 DOI: 10.1021/jf073494t]
- 120 **Lucas EA**, Li W, Peterson SK, Brown A, Kuvibidila S, Perkins-Veazie P, Clarke SL, Smith BJ. Mango modulates body fat and plasma glucose and lipids in mice fed a high-fat diet. *Br J Nutr* 2011; **106**: 1495-1505 [PMID: 21733317 DOI: 10.1017/S0007114511002066]
- 121 **Masibo M**, He Q. Mango Bioactive Compounds and Related Nutraceutical Properties: A Review. *Food Rev* 2009; **25**: 346-370 [DOI: 10.1080/87559120903153524]
- 122 **Li X**, Cui X, Sun X, Li X, Zhu Q, Li W. Mangiferin prevents diabetic nephropathy progression in streptozotocin-induced diabetic rats. *Phytother Res* 2010; **24**: 893-899 [PMID: 19960420]
- 123 **Zhou J**, Zhou S, Tang J, Zhang K, Guang L, Huang Y, Xu Y, Ying Y, Zhang L, Li D. Protective effect of berberine on beta cells in streptozotocin- and high-carbohydrate/high-fat diet-induced diabetic rats. *Eur J Pharmacol* 2009; **606**: 262-268 [PMID: 19374872]
- 124 **Fatehi M**, Saleh TM, Fatehi-Hassanabad Z, Farrokhfal K, Jafarzadeh M, Davodi S. A pharmacological study on *Berberis vulgaris* fruit extract. *J Ethnopharmacol* 2005; **102**: 46-52 [PMID: 15993555 DOI: 10.1016/j.jep.2005.05.019]
- 125 **Fatehi-Hassanabad Z**, Jafarzadeh M, Tarhini A, Fatehi M. The antihypertensive and vasodilator effects of aqueous extract from *Berberis vulgaris* fruit on hypertensive rats. *Phytother Res* 2005; **19**: 222-225 [PMID: 15934023 DOI: 10.1002/ptr.1661]
- 126 **Gao CR**, Zhang JQ, Huang QL. [Experimental study on berberin raised insulin sensitivity in insulin resistance rat mod-

- els]. *Zhongguo Zhongxiyi Jiehe Zazhi* 1997; **17**: 162-164 [PMID: 9863084]
- 127 Yin J, Hu R, Chen M, Tang J, Li F, Yang Y, Chen J. Effects of berberine on glucose metabolism in vitro. *Metabolism* 2002; **51**: 1439-1443 [PMID: 12404195 DOI: 10.1053/meta.2002.34715]
  - 128 Al-Farsi MA, Lee CY. Nutritional and functional properties of dates: a review. *Crit Rev Food Sci Nutr* 2008; **48**: 877-887 [PMID: 18949591 DOI: 10.1080/10408390701724264]
  - 129 Saafi EB, Louedi M, Elfeki A, Zakhama A, Najjar MF, Hammami M, Achour L. Protective effect of date palm fruit extract (*Phoenix dactylifera* L.) on dimethoate induced-oxidative stress in rat liver. *Exp Toxicol Pathol* 2011; **63**: 433-441 [PMID: 20359872 DOI: 10.1016/j.etp.2010.03.002]
  - 130 Vayalil PK. Date fruits (*Phoenix dactylifera* Linn): an emerging medicinal food. *Crit Rev Food Sci Nutr* 2012; **52**: 249-271 [PMID: 22214443 DOI: 10.1080/10408398.2010.499824]
  - 131 Zangiabadi N, Asadi-Shekaari M, Sheibani V, Jafari M, Shabani M, Asadi AR, Tajadini H, Jarahi M. Date fruit extract is a neuroprotective agent in diabetic peripheral neuropathy in streptozotocin-induced diabetic rats: a multimodal analysis. *Oxid Med Cell Longev* 2011; **2011**: 976948 [PMID: 22191015]
  - 132 Slatnar A, Klancar U, Stampar F, Veberic R. Effect of drying of figs (*Ficus carica* L.) on the contents of sugars, organic acids, and phenolic compounds. *J Agric Food Chem* 2011; **59**: 11696-11702 [PMID: 21958361 DOI: 10.1021/jf202707y]
  - 133 Josef B, Raj S. Pharmacognostic and phytochemical properties of *Ficus carica* Linn: An overview. *Int J Pharma Tech Res* 2011; **3**: 8-12
  - 134 Duranti M. Grain legume proteins and nutraceutical properties. *Fitoterapia* 2006; **77**: 67-82 [PMID: 16406359 DOI: 10.1016/j.fitote.2005.11.008]
  - 135 Madar Z, Stark AH. New legume sources as therapeutic agents. *Br J Nutr* 2002; **88** Suppl 3: S287-S292 [PMID: 12498629 DOI: 10.1079/BJN2002719]
  - 136 Flight I, Clifton P. Cereal grains and legumes in the prevention of coronary heart disease and stroke: a review of the literature. *Eur J Clin Nutr* 2006; **60**: 1145-1159 [PMID: 16670693 DOI: 10.1038/sj.ejcn.1602435]
  - 137 López-Amorós ML, Hernández T, Estrella I. Estrella I. Effect of germination on legume phenolic compounds and their antioxidant activity. *J Food Comp Anal* 2006; **19**: 277-283 [DOI: 10.1016/j.jfca.2004.06.012]
  - 138 Zou Y, Chang SK, Gu Y, Qian SY. Antioxidant activity and phenolic compositions of lentil (*Lens culinaris* var. Morton) extract and its fractions. *J Agric Food Chem* 2011; **59**: 2268-2276 [PMID: 21332205 DOI: 10.1021/jf104640k]
  - 139 Boualga A, Prost J, Taleb-Senouci D, Krouf D, Kharoubi O, Lamri-Senhadj M, Belleville J, Bouchenak M. Purified chickpea or lentil proteins impair VLDL metabolism and lipoprotein lipase activity in epididymal fat, but not in muscle, compared to casein, in growing rats. *Eur J Nutr* 2009; **48**: 162-169 [PMID: 19165521 DOI: 10.1007/s00394-009-0777-4]
  - 140 Yao F, Sun C, Chang SK. Lentil polyphenol extract prevents angiotensin II-induced hypertension, vascular remodelling and perivascular fibrosis. *Food Funct* 2012; **3**: 127-133 [PMID: 22159297 DOI: 10.1039/c1fo10142k]
  - 141 Barrett ML, Udani JK. A proprietary alpha-amylase inhibitor from white bean (*Phaseolus vulgaris*): a review of clinical studies on weight loss and glycemic control. *Nutr J* 2011; **10**: 24 [PMID: 21414227 DOI: 10.1186/1475-2891-10-24]
  - 142 Yao Y, Cheng XZ, Wang LX, Wang SH, Ren G. Major Phenolic Compounds, Antioxidant Capacity and Antidiabetic Potential of Rice Bean (*Vigna umbellata* L.) in China. *Int J Mol Sci* 2012; **13**: 2707-2716 [PMID: 22489119 DOI: 10.3390/ijms13032707]
  - 143 Helmstädter A. Beans and diabetes: *Phaseolus vulgaris* preparations as antihyperglycemic agents. *J Med Food* 2010; **13**: 251-254 [PMID: 20132042 DOI: 10.1089/jmf.2009.0002]
  - 144 Geil PB, Anderson JW. Nutrition and health implications of dry beans: a review. *J Am Coll Nutr* 1994; **13**: 549-558 [PMID: 7706585 DOI: 10.1080/07315724.1994.10718446]
  - 145 Preuss HG. Bean amylase inhibitor and other carbohydrate absorption blockers: effects on diabetes and general health. *J Am Coll Nutr* 2009; **28**: 266-276 [PMID: 20150600 DOI: 10.1080/07315724.2009.10719781]
  - 146 Thompson SV, Winham DM, Hutchins AM. Bean and rice meals reduce postprandial glycemic response in adults with type 2 diabetes: a cross-over study. *Nutr J* 2012; **11**: 23 [PMID: 22494488 DOI: 10.1186/1475-2891-11-23]
  - 147 Anderson JW, Bush HM. Soy protein effects on serum lipoproteins: a quality assessment and meta-analysis of randomized, controlled studies. *J Am Coll Nutr* 2011; **30**: 79-91 [PMID: 21730216 DOI: 10.1080/07315724.2011.10719947]
  - 148 Azadbakht L, Shakerhosseini R, Atabak S, Jamshidian M, Mehrabi Y, Esmail-Zadeh A. Beneficiary effect of dietary soy protein on lowering plasma levels of lipid and improving kidney function in type II diabetes with nephropathy. *Eur J Clin Nutr* 2003; **57**: 1292-1294 [PMID: 14506491 DOI: 10.1038/sj.ejcn.1601688]
  - 149 Bhathena SJ, Velasquez MT. Beneficial role of dietary phytoestrogens in obesity and diabetes. *Am J Clin Nutr* 2002; **76**: 1191-1201 [PMID: 12450882]
  - 150 Gilbert ER, Liu D. Anti-diabetic functions of soy isoflavone genistein: mechanisms underlying its effects on pancreatic  $\beta$ -cell function. *Food Funct* 2013; **4**: 200-212 [PMID: 23160185 DOI: 10.1039/c2fo30199g]
  - 151 Ronis MJ, Chen Y, Badeaux J, Badger TM. Dietary soy protein isolate attenuates metabolic syndrome in rats via effects on PPAR, LXR, and SREBP signaling. *J Nutr* 2009; **139**: 1431-1438 [PMID: 19515742 DOI: 10.3945/jn.109.107029]
  - 152 Yang B, Chen Y, Xu T, Yu Y, Huang T, Hu X, Li D. Systematic review and meta-analysis of soy products consumption in patients with type 2 diabetes mellitus. *Asia Pac J Clin Nutr* 2011; **20**: 593-602 [PMID: 22094845]
  - 153 Borradaile NM, de Dreu LE, Wilcox LJ, Edwards JY, Huff MW. Soya phytoestrogens, genistein and daidzein, decrease apolipoprotein B secretion from HepG2 cells through multiple mechanisms. *Biochem J* 2002; **366**: 531-539 [PMID: 12030847 DOI: 10.1042/BJ20020046]
  - 154 Potter SM. Overview of proposed mechanisms for the hypocholesterolemic effect of soy. *J Nutr* 1995; **125**: 606S-611S [PMID: 7884541]
  - 155 Liu CF, Pan TM. Beneficial Effects of Bioactive Peptides Derived from Soybean on Human Health and Their Production by Genetic Engineering. *Soybean and health* 2011: 311-329
  - 156 Frigolet ME, Torres N, Uribe-Figueroa L, Rangel C, Jimenez-Sanchez G, Tovar AR. White adipose tissue genome wide-expression profiling and adipocyte metabolic functions after soy protein consumption in rats. *J Nutr Biochem* 2011; **22**: 118-129 [PMID: 20471815 DOI: 10.1016/j.jnutbio.2009.12.006]
  - 157 Cheik NC, Rossi EA, Guerra RL, Tenório NM, Oller do Nascimento CM, Viana FP, Manzoni MS, Carlos IZ, Leão da Silva P, Vendramini RC, Dâmaso AR. Effects of a ferment soy product on the adipocyte area reduction and dyslipidemia control in hypercholesterolemic adult male rats. *Lipids Health Dis* 2008; **7**: 50 [PMID: 19087298 DOI: 10.1186/1476-511X-7-50]
  - 158 Zhang Z, Lanza E, Ross AC, Albert PS, Colburn NH, Rovine MJ, Bagshaw D, Ulbrecht JS, Hartman TJ. A high-legume low-glycemic index diet reduces fasting plasma leptin in middle-aged insulin-resistant and -sensitive men. *Eur J Clin Nutr* 2011; **65**: 415-418 [PMID: 21206508 DOI: 10.1038/ejcn.2010.273]
  - 159 Segura R, Javierre C, Lizarraga MA, Ros E. Other relevant components of nuts: phytosterols, folate and minerals. *Br J Nutr* 2006; **96** Suppl 2: S36-S44 [PMID: 17125532 DOI: 10.1017/BJN20061862]
  - 160 Kendall CW, Esfahani A, Truan J, Srichaikul K, Jenkins DJ. Health benefits of nuts in prevention and management of diabetes. *Asia Pac J Clin Nutr* 2010; **19**: 110-116 [PMID: 20150600 DOI: 10.1080/07315724.2009.10719781]



- 20199995]
- 161 **Ros E.** Nuts and novel biomarkers of cardiovascular disease. *Am J Clin Nutr* 2009; **89**: 1649S-1656S [PMID: 19321561 DOI: 10.3945/ajcn.2009.26736R]
  - 162 **Jenkins DJ,** Hu FB, Tapsell LC, Josse AR, Kendall CW. Possible benefit of nuts in type 2 diabetes. *J Nutr* 2008; **138**: 1752S-1756S [PMID: 18716181]
  - 163 **Li TY,** Brennan AM, Wedick NM, Mantzoros C, Rifai N, Hu FB. Regular consumption of nuts is associated with a lower risk of cardiovascular disease in women with type 2 diabetes. *J Nutr* 2009; **139**: 1333-1338 [PMID: 19420347 DOI: 10.3945/jn.108.103622]
  - 164 **Tey SL,** Brown R, Gray A, Chisholm A, Delahunty C. Nuts improve diet quality compared to other energy-dense snacks while maintaining body weight. *J Nutr Metab* 2011; **2011**: 357350 [PMID: 21845219]
  - 165 **Mattes RD,** Dreher ML. Nuts and healthy body weight maintenance mechanisms. *Asia Pac J Clin Nutr* 2010; **19**: 137-141 [PMID: 20199999]
  - 166 **Mantzoros CS,** Williams CJ, Manson JE, Meigs JB, Hu FB. Adherence to the Mediterranean dietary pattern is positively associated with plasma adiponectin concentrations in diabetic women. *Am J Clin Nutr* 2006; **84**: 328-335 [PMID: 16895879]
  - 167 **Jiang R,** Jacobs DR, Mayer-Davis E, Szklo M, Herrington D, Jenny NS, Kronmal R, Barr RG. Nut and seed consumption and inflammatory markers in the multi-ethnic study of atherosclerosis. *Am J Epidemiol* 2006; **163**: 222-231 [PMID: 16357111 DOI: 10.1093/aje/kwj033]
  - 168 **Al-Salami H,** Butt G, Fawcett JP, Tucker IG, Golocorbin-Kon S, Mikov M. Probiotic treatment reduces blood glucose levels and increases systemic absorption of gliclazide in diabetic rats. *Eur J Drug Metab Pharmacokinet* 2008; **33**: 101-106 [PMID: 18777945 DOI: 10.1007/BF03191026]
  - 169 **Ataie-Jafari A,** Larijani B, Alavi Majd H, Tahbaz F. Cholesterol-lowering effect of probiotic yogurt in comparison with ordinary yogurt in mildly to moderately hypercholesterolemic subjects. *Ann Nutr Metab* 2009; **54**: 22-27 [PMID: 19229114 DOI: 10.1159/000203284]
  - 170 **Chang BJ,** Park SU, Jang YS, Ko SH, Joo NM, Kim SI, Kim CH, Chang DK. Effect of functional yogurt NY-YP901 in improving the trait of metabolic syndrome. *Eur J Clin Nutr* 2011; **65**: 1250-1255 [PMID: 21697819 DOI: 10.1038/ejcn.2011.115]
  - 171 **Guo C,** Zhang L. [Cholesterol-lowering effects of probiotics-a review]. *Weishengwu Xuebao* 2010; **50**: 1590-1599 [PMID: 21365911]
  - 172 **Guzel-Seydim ZB,** Kok-Tas T, Greene AK, Seydim AC. Review: functional properties of kefir. *Crit Rev Food Sci Nutr* 2011; **51**: 261-268 [PMID: 21390946 DOI: 10.1080/10408390903579029]
  - 173 **Lin MY,** Yen CL. Reactive oxygen species and lipid peroxidation product-scavenging ability of yogurt organisms. *J Dairy Sci* 1999; **82**: 1629-1634 [PMID: 10480088 DOI: 10.3168/jds.S0022-0302(99)75391-9]
  - 174 **Liu S,** Choi HK, Ford E, Song Y, Klevak A, Buring JE, Manson JE. A prospective study of dairy intake and the risk of type 2 diabetes in women. *Diabetes Care* 2006; **29**: 1579-1584 [PMID: 16801582 DOI: 10.2337/dc06-0256]
  - 175 **Lopitz-Otsoa F,** Rementeria A, Elguezal N, Garaizar J. Kefir: a symbiotic yeasts-bacteria community with alleged healthy capabilities. *Rev Iberoam Micol* 2006; **23**: 67-74 [PMID: 16854180 DOI: 10.1016/S1130-1406(06)70016-X]
  - 176 **Marteau PR.** Probiotics in clinical conditions. *Clin Rev Allergy Immunol* 2002; **22**: 255-273 [PMID: 12043384 DOI: 10.1007/s12016-002-0011-0]
  - 177 **Pfeuffer M,** Schrenzenmeier J. Milk and the metabolic syndrome. *Obes Rev* 2007; **8**: 109-118 [PMID: 17300277 DOI: 10.1111/j.1467-789X.2006.00265.x]
  - 178 **Shah-Bidar S,** Neyestani TR, Djazayeri A. Efficacy of vitamin D3-fortified-yogurt drink on anthropometric, metabolic, inflammatory and oxidative stress biomarkers according to vitamin D receptor gene polymorphisms in type 2 diabetic patients: a study protocol for a randomized controlled clinical trial. *BMC Endocr Disord* 2011; **11**: 12 [PMID: 21696575 DOI: 10.1186/1472-6823-11-12]
  - 179 **Teruya K,** Yamashita M, Tominaga R, Nagira T, Shim SY, Katakura Y, Tokumaru S, Tokumaru K, Barnes D, Shirahata S. Fermented milk, Kefram-Kefir enhances glucose uptake into insulin-responsive muscle cells. *Cytotechnology* 2002; **40**: 107-116 [PMID: 19003111 DOI: 10.1023/A:1023926407877]
  - 180 **Boukourt FO,** Girard A, Prost JL, Ait-Yahia D, Bouchenak M, Belleville J. Fish protein improves the total antioxidant status of streptozotocin-induced diabetes in spontaneously hypertensive rat. *Med Sci Monit* 2004; **10**: BR397-BR404 [PMID: 15507843]
  - 181 **Farmer A,** Montori V, Dinneen S, Clar C. Fish oil in people with type 2 diabetes mellitus. *Cochrane Database Syst Rev* 2001; (3): CD003205 [PMID: 11687050]
  - 182 **Jacques H,** Gascon A, Bergeron N, Lavigne C, Hurley C, Deshaies Y, Moorjani S, Julien P. Role of dietary fish protein in the regulation of plasma lipids. *Can J Cardiol* 1995; **11** Suppl G: 63G-71G [PMID: 7585295]
  - 183 **Lankinen M,** Schwab U, Kolehmainen M, Paananen J, Poutanen K, Mykkänen H, Seppänen-Laakso T, Gylling H, Uusitupa M, Orešič M. Whole grain products, fish and bilberries alter glucose and lipid metabolism in a randomized, controlled trial: the Sysdimet study. *PLoS One* 2011; **6**: e22646 [PMID: 21901116 DOI: 10.1371/journal.pone.0022646]
  - 184 **McEwen B,** Morel-Kopp MC, Tofler G, Ward C. Effect of omega-3 fish oil on cardiovascular risk in diabetes. *Diabetes Educ* 2010; **36**: 565-584 [PMID: 20534874 DOI: 10.1177/0145721710372675]
  - 185 **Pecis M,** de Azevedo MJ, Gross JL. Chicken and fish diet reduces glomerular hyperfiltration in IDDM patients. *Diabetes Care* 1994; **17**: 665-672 [PMID: 7924775 DOI: 10.2337/diacare.17.7.665]
  - 186 **Cicerale S,** Conlan XA, Sinclair AJ, Keast RS. Chemistry and health of olive oil phenolics. *Crit Rev Food Sci Nutr* 2009; **49**: 218-236 [PMID: 19093267 DOI: 10.1080/10408390701856223]
  - 187 **Lucas L,** Russell A, Keast R. Molecular mechanisms of inflammation. Anti-inflammatory benefits of virgin olive oil and the phenolic compound oleocanthal. *Curr Pharm Des* 2011; **17**: 754-768 [PMID: 21443487 DOI: 10.2174/138161211795428911]
  - 188 **Madigan C,** Ryan M, Owens D, Collins P, Tomkin GH. Dietary unsaturated fatty acids in type 2 diabetes: higher levels of postprandial lipoprotein on a linoleic acid-rich sunflower oil diet compared with an oleic acid-rich olive oil diet. *Diabetes Care* 2000; **23**: 1472-1477 [PMID: 11023139 DOI: 10.2337/diacare.23.10.1472]
  - 189 **Wardhana ES,** Datau EA. The role of omega-3 fatty acids contained in olive oil on chronic inflammation. *Acta Med Indones* 2011; **43**: 138-143 [PMID: 21785178]
  - 190 **Babu PV,** Sabitha KE, Srinivasan P, Shyamaladevi CS. Green tea attenuates diabetes induced Maillard-type fluorescence and collagen cross-linking in the heart of streptozotocin diabetic rats. *Pharmacol Res* 2007; **55**: 433-440 [PMID: 17336542 DOI: 10.1016/j.phrs.2007.01.019]
  - 191 **Fiorino P,** Evangelista FS, Santos F, Motter Magri FM, Delorenzi JC, Ginoza M, Farah V. The effects of green tea consumption on cardiometabolic alterations induced by experimental diabetes. *Exp Diabetes Res* 2012; **2012**: 309231 [PMID: 22474420]
  - 192 **Kumar B,** Gupta SK, Nag TC, Srivastava S, Saxena R. Green tea prevents hyperglycemia-induced retinal oxidative stress and inflammation in streptozotocin-induced diabetic rats. *Ophthalmic Res* 2012; **47**: 103-108 [PMID: 21997135]
  - 193 **Kim HM,** Kim J. The effects of green tea on obesity and type 2 diabetes. *Diabetes Metab J* 2013; **37**: 173-175 [PMID: 23807919 DOI: 10.4093/dmj.2013.37.3.173]
  - 194 **Anderson RA.** Chromium and polyphenols from cinnamon

- improve insulin sensitivity. *Proc Nutr Soc* 2008; **67**: 48-53 [PMID: 18234131 DOI: 10.1017/S0029665108006010]
- 195 **Gruenwald J**, Freder J, Armbruester N. Cinnamon and health. *Crit Rev Food Sci Nutr* 2010; **50**: 822-834 [PMID: 20924865 DOI: 10.1080/10408390902773052]
  - 196 **Kirkham S**, Akilen R, Sharma S, Tsiami A. The potential of cinnamon to reduce blood glucose levels in patients with type 2 diabetes and insulin resistance. *Diabetes Obes Metab* 2009; **11**: 1100-1113 [PMID: 19930003 DOI: 10.1111/j.1463-1326.2009.01094.x]
  - 197 **Honda S**, Aoki F, Tanaka H, Kishida H, Nishiyama T, Okada S, Matsumoto I, Abe K, Mae T. Effects of ingested turmeric oleoresin on glucose and lipid metabolisms in obese diabetic mice: a DNA microarray study. *J Agric Food Chem* 2006; **54**: 9055-9062 [PMID: 17117790 DOI: 10.1021/jf061788t]
  - 198 **Khajehdehi P**, Pakfetrat M, Javidnia K, Azad F, Malekmakan L, Nasab MH, Dehghanzadeh G. Oral supplementation of turmeric attenuates proteinuria, transforming growth factor- $\beta$  and interleukin-8 levels in patients with overt type 2 diabetic nephropathy: a randomized, double-blind and placebo-controlled study. *Scand J Urol Nephrol* 2011; **45**: 365-370 [PMID: 21627399 DOI: 10.3109/00365599.2011.585622]
  - 199 **Lekshmi PC**, Arimboor R, Raghu KG, Menon AN. Turmeric, the antioxidant protein from turmeric (*Curcuma longa*) exhibits antihyperglycaemic effects. *Nat Prod Res* 2012; **26**: 1654-1658 [PMID: 21972920]
  - 200 **Rayne S**, Mazza G. Biological activities of extracts from sumac (*Rhus spp.*): a review. *Plant Foods Hum Nutr* 2007; **62**: 165-175 [PMID: 17909971 DOI: 10.1007/s11130-007-0058-4]
  - 201 **Shafiei M**, Nobakht M, Moazzam AA. Lipid-lowering effect of *Rhus coriaria* L. (sumac) fruit extract in hypercholesterolemic rats. *Pharmazie* 2011; **66**: 988-992 [PMID: 22312707]
  - 202 **Giancarlo S**, Rosa LM, Nadjafi F, Francesco M. Hypoglycaemic activity of two spices extracts: *Rhus coriaria* L. and *Bunium persicum* Boiss. *Nat Prod Res* 2006; **20**: 882-886 [PMID: 16753927 DOI: 10.1080/14786410500520186]
  - 203 **Pourahmad J**, Eskandari MR, Shakibaei R, Kamalinejad M. A search for hepatoprotective activity of aqueous extract of *Rhus coriaria* L. against oxidative stress cytotoxicity. *Food Chem Toxicol* 2010; **48**: 854-858 [PMID: 20036300]

**P- Reviewers:** Hiroyuki T, Karoly R, Nahid P, Semir O

**S- Editor:** Qi Y **L- Editor:** A **E- Editor:** Liu SQ





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>

