



ISSN NO. 2320-5407

Journal homepage: <http://www.journalijar.com>

INTERNATIONAL JOURNAL
OF ADVANCED RESEARCH

REVIEW ARTICLE

A Quick Review on Anti-diabetic Plants and Action of Phytochemicals

Md. Abdul Mannan¹, Beauty Akter Rupa², Md. Nur Kabidul Azam¹, *Md. Nasir Ahmed^{1,2},
Md. Nazmul Hasan³

1. TechB Herbal Solution, Kushtia-7040, Bangladesh.

2. Department of Biotechnology & Genetic Engineering, University of Development Alternative, Dhanmondi, Dhaka-1205, Bangladesh.

3. Department of Genetic Engineering and Biotechnology, Jessore Science & Technology University, Jessore-7408, Bangladesh.

Manuscript Info

Manuscript History:

Received: 15 March 2014

Final Accepted: 25 April 2014

Published Online: May 2014

Key words:

Medicinal plants, diabetes mellitus, antidiabetic activity, hypoglycemic, phytochemical

*Corresponding Author

Md. Nasir Ahmed

Abstract

Diabetes is an endocrinological disorder arising from insulin deficiency or due to effectiveness of the insulin produced by the body. Around 200 million people of the world are currently suffering from diabetes and the figure is projected to rise to 300 million within 2025 by the World Health Organization. The disease is caused by the inability of pancreas to produce insulin or inability of the body metabolic system to properly use the insulin produced. In Asia, India and China are the leading countries in herbal plants research, and there has been an increase in medicinal research on plants extract for diabetes treatment since 1995 in these regions. The objective of the present study was to document the medicinal plants used for the diabetes treatment, so that future work directed towards the identification of active principles from these medicinal plants may provide the opportunity for the development of a novel class of agents for the treatment of diabetes. In this review, we have attempted to compile a list of total 141 plant species belongs to 82 families, along with their type of used extracts, botanical name, Plant part used and active principles are included based on available scientific literature those has been reported to be effective in diabetes. Different mechanism of action of anti-hyperglycemic phytochemicals are also shortly discussed. The information collected shows that plant leaves are about 51% more favorable for storing active ingredients, as compared to other parts of the medicinal plants. The documentation of anti-diabetic medicinal plants stimulates the researchers for further research on the potential use of medicinal plants having anti-diabetic potential.

Copy Right, IJAR, 2014., All rights reserved.

1. Introduction

Diabetes mellitus is a common and very prevalent disease affecting the citizens of both developed and developing countries. 6% population of human beings affected by Diabetes mellitus, which is an endocrine disorder and this will affect 5 times more people in next 10 years (WHO/Acadia, 1992; ADA, 1997). Diabetes mellitus is not disease but a syndrome with combination of hereditary and environmental cause that results in high blood sugar levels may know as Hyperglycemia. Our body always maintain sugar level by chemicals and hormones especially beta cells of pancreas that produce insulin and defects in either insulin secretion or insulin action leads to diabetes mellitus (Tierney 2002). And this disease results in high urine production, thirst and blurred vision, lethargy and changes in energy metabolism (Patel et al., 2012).

Plants, can provide biologically active molecules and lead structures for the development of modified derivatives with enhanced activity and reduced toxicity. The small fraction of flowering plants that have so far been investigated have yielded about 120 therapeutic agents of known structure from about 90 species of plants. (Joy et al., 1998). Indigenous medicinal practices of various communities throughout the world have always proved to be an excellent route to discovery of many important modern drugs (Balick and Cox, 1996; Gilani and Rahman, 2005). In some cases, the crude extract of medicinal plants may be used as medicaments. About 121 (45 tropical and 76 subtropical) major plant drugs have been identified for which no synthetic one is currently available (Kumar et al., 1997). It has been estimated that more than 400 traditional plants or plant-derived products have been used for the management of type 2 diabetes across geographically (Bailey and Day, 1989) and about 800 plants have anti-diabetic potential (Alarcon-Aguilara et al., 1998). Plants have their chemical compounds which demonstrate alternative and safe effects on diabetes mellitus. Most of plants contain glycosides, alkaloids, terpenoids, flavonoids, carotenoids, etc., that are frequently implicated as having antidiabetic effect (Malviya et al., 2010). *Galega officinalis* is plant from which hypoglycemic drugs was obtained traditionally. insulin, biguanides, sulfonylurease and thiazolidinediones are modern pharmacotherapeutics, but still except glycemic control with insulin, need to look new drugs for more efficacious agents with less side effects is needed (Gover et al., 2002). Herbal medicine also known as phytomedicine, refers to usage of roots, leaves, bark, flowers, seeds and berries for medicines. Herbalism is becoming focusing point due to research on herbs and their use in treatment of diabetes and its prevention (Ang-Lee et al., 2001). The scientific study of traditional medicines, concerned medicinal plants are thus of great importance.

The anti-diabetic plant species from seventeen families have earlier been recorded by Sidhu and Sharma (2013a; 2013b). In the present review an attempt has been made to investigate the anti-diabetic medicinal plants and may be useful to the health professionals, scientists and scholars working in the field of pharmacology and therapeutics to develop anti-diabetic drugs.

2. Methodology:

The authors first carried out a computer-aided search of the literature stored in various medical data banks. In addition, relevant information was also searched on the Internet. All publications containing original data and an adequate detailed description of methodology were considered in the present review. The papers reviewed in this article are selected from the medicinal journals as tabulated in Table 1 due to reliable reputation in medicinal research on herbal plant extract. The research papers were extracted from the selected 84 journals under the key words of "Anti-diabetic activities of plants". Similar observations have also been obtained in other anti-diabetic plants review study (Chang et al., 2012). Plants which did not show any significant hypoglycemic effect were not included in the document.

3. Discussions

The paper presented a total of 141 plant species, have been enlisted in table 2 that have anti-diabetic properties. These plant species belongs to 82 families (Table 3). Four species belongs to each of families Combretaceae, Solanaceae and Verbenaceae. The families like Aizoaceae, Nyctaginaceae, Nymphaeaceae, Burseraceae, Cecropiaceae, Chenopodiaceae, Rosaceae, Rubiaceae, Ebenaceae, Urticaceae, Lauraceae, Liliaceae, Loganiaceae have three species each, followed by Annonaceae, Berberidaceae, Bignoniaceae, Bombacaceae, Boraginaceae, Convolvulaceae, Costaceae, Hippocrateaceae, Lythraceae, Mimosaceae, Oleaceae, Pandanaceae, Piperaceae, Polygonaceae, Portulacaceae, Ranunculaceae, Sapindaceae, Sapotaceae, Scrophulariaceae, Sterculiaceae, Zingiberaceae and Zygophyllaceae credited with two species. In addition to this, one species have been reported from the families like Acoraceae, Agavaceae, Alangiaceae, Apiaceae, Araceae, Arecaceae, Araliaceae, Balanitiaceae, Basellaceae, Capparidaceae, Caricaceae, Caryophyllaceae, Moringaceae, Musaceae, Myricaceae, Onagraceae, Orchidaceae, Oxalidaceae, Palmaceae, Passifloraceae, Polypodiaceae, Primulaceae, Rhamnaceae, Crassulaceae, Elaeagnaceae, Equisetaceae, Eucommiaceae, Flacourtiaceae, Fomitopsidaceae, Hericiaceae, Hypericaceae, Irvingiaceae, Juglandaceae, Melastomataceae, Melianthaceae, Salvadoraceae, Samydaceae, Sonneratiaceae, Symplocaceae, Thymelaeaceae, Tiliaceae, Ulmaceae, Violaceae and Vitaceae having antidiabetic activity.

Table 1: The 84 selected journals for the study

African Journal of Traditional, Complementary and Alternative Medicines	European Journal of Scientific Research	International Journal of Phytomedicine	Journal of Diabetology
African Journal of Plant Science	Fitoterapia	International Journal of Applied Biology and Pharmaceutical Technology	Journal of Advanced Pharmaceutical Technology and Research
African Journal of Biotechnology	Global Journal of Biotechnology and Biochemistry	International Journal of Research in Pharmaceutical and Biomedical Sciences	Journal of Advanced Pharmacy Education and Research
Applied Biochemistry and Biotechnology	Global Journal of Pharmacology	International Journal of Molecular Sciences	Journal of Pharmacognosy and Phytotherapy
Asian Journal of Medical Sciences	Indian Journal of Pharmacology	International Journal of Pharma Research and Development	Journal of Applied Biomedicine
Asian Journal of Pharmaceutical and Clinical Research	Indian Journal of Experimental Biology	International Journal of Applied Research in Natural Products	Journal of Endocrinology and Metabolism
Asian Pacific Journal of Tropical Medicine	Indian Journal of Pharmaceutical Science and Research	International Conference on Asia Agriculture and Animal	Libyan Agriculture Research Centre Journal
Asian Pacific Journal of Tropical Biomedicine	Indian Journal of Pharmaceutical Sciences	International Journal of Phytomedicine	International Methods and Findings in Experimental and Clinical Pharmacology
Biological and Pharmaceutical Bulletin	Indian Journal of Pharmaceutical Education and Research	International Journal of Molecular Sciences	Pakistan Journal of Pharmaceutical Sciences
Biological and Pharmaceutical Bulletin	International Journal of Pharmaceutical and Biological Archives	International Journal of Applied Biology and Pharmaceutical Technology	Pacific Journal of Medical Sciences
Biology and Medicine	International Journal of Pharmacy and Pharmaceutical Sciences	International Research Journal of Pharmacy	Pharmacologyonline
Bioscience Biotechnology and Biochemistry	International Journal of Applied Research in Natural Products	International Journal of Pharma Sciences and Research	Pharmacognosy Magazine
BMC Complementary and Alternative Medicine	International Journal of Pharmaceutical Sciences and Nanotechnology	International Journal of PharmTech Research	Phytotherapy Research
BMC Pharmacology	International Journal of Pharmaceutical Research and Development (IJPRD)	Iranian Journal of Pharmaceutical Research	Pharmacy and Pharmacology Communications
British Journal of Pharmaceutical Research	Journal of Medical Sciences	Journal of Applied Pharmaceutical Science	Records of Natural Products
British Journal of Nutrition	Journal of Applied Pharmaceutical Science	Journal of Ethnopharmacology	Singapore Medical Journal
Current Science	Journal of Pharmaceutical Research and Clinical Practice	Journal of Medicinal Plants Research	South African Journal of Botany

Der Pharmacia Lettre	Journal of Herbs, Spices and Medicinal Plants	Journal of Life Sciences	The Egyptian Journal of Hospital Medicine
Diabetes Research	Journal of Chemical and Pharmaceutical Research	Journal of Pharmacy and Bio allied Sciences	
Diabetologia Croatica	Journal of Herbal Medicine and Toxicology	Journal of Pharmacognosy and Phytotherapy	
Diabetes Research and Clinical Practice	Journal of Medicinal Plants	Journal of Medicinal Food	
Drug Invention Today	Journal of Medicinal Plants	Journal of the Science of Food and Agriculture	

Table 2: Medicinal plants with anti-diabetic active principles and their reported effect on experimental models

SN	Botanical Name	Family Name	Used extract	Active Principles	References
Leaves					
1	<i>Aegiceras corniculatum</i> (L.) Blanco	Primulaceae	E	Leaf extract	Gurudeeban et al., 2012
2	<i>Aframomum melegueta</i> (Rosc.) K. Schum.	Zingiberaceae	Aq.	Leaf extract	Mojekwu et al., 2011
3	<i>Alangium lamarckii</i> Thw.	Alangiaceae	Al.	Leaf extract	Kumar et al., 2011
4	<i>Aloe vera</i> (L.) Burm. Fil.	Liliaceae	Aq.	Leaf extract	Rehman et al., 2011
5	<i>Anabasis articulata</i> (Forssk) Moq.	Chenopodiaceae	Aq.	Saponins	Kambouche et al., 2009
6	<i>Annona squamosa</i> L.	Annonaceae	Aq.	Leaf extract	Gupta et al., 2005
7	<i>Areca catechu</i> L.	Palmaceae	P.E., Chlf. & M	Triterpenoids	Mondal et al., 2012
8	<i>Averrhoa bilimbi</i> L.	Oxalidaceae	E	Leaf extract	Pushparaj et al., 2000
9	<i>Basella rubra</i> L.	Basellaceae	Aq.	Leaf pulp	Nirmala et al., 2009
10	<i>Bersama engleriana</i> Gurke	Melanthaceae	Aq. & M	Leaf extract	Njike et al., 2005
11	<i>Boerhaavia diffusa</i> L.	Nyctaginaceae	Aq.	Alkaloids, sterols or triterpenoids	Koffi et al., 2011
12	<i>Bougainvillea glabra</i> L.	Nyctaginaceae	Aq.	Alkaloids, flavonoids, saponins & cardiac glycosides	Adebayo et al., 2009
13	<i>Bryophyllum pinnatum</i> (Lam.) Kurz	Crassulaceae	Aq.	Flavonoids, polyphenols, triterpenoids	Ojewole, 2005
14	<i>Cardiospermum halicacabum</i> L.	Sapindaceae	E	Leaf extract	Veeramani et al., 2008
15	<i>Cecropia obtusifolia</i> Bertol.	Cecropiaceae	Aq. & B	Flavone, isoorientin & 3-caffeoylquinic	Andrade-Cetto and Wiedenfeld (2001)
16	<i>Cecropia pachystachya</i> Mart.	Cecropiaceae	M	Leaf extract	Aragao et al., 2010
17	<i>Chamaerops humilis</i> L.	Arecaceae	Aq.	Leaf extract	Gaamoussi et al., 2010
18	<i>Cinnamomum tamala</i> Fr. Nees.	Lauraceae	Aq.	Leaf extract	Chakraborty and Das (2010)
19	<i>Cissus sicyoides</i> L.	Vitaceae	Aq.	Flavonoids & tannins	Viana et al., 2004
20	<i>Clerodendrum capitatum</i> (Willd)	Verbenaceae	Aq.	Saponins, flavonoids,	Adeneye et al., 2008

	Schumach et. Thonn.			alkaloids, tannins, glycosides	
21	<i>Combretum micranthum</i> G. Don	Combretaceae	Aq.	Leaf extract	Chika and Bello (2010)
22	<i>Costus afer</i> Ker Gawl.	Costaceae	M	Alkaloids, flavonoids, tannins, saponins	Momoh et al., 2011
23	<i>Dolichandrone falcata</i> Seem.	Bignoniaceae	Aq.	Steroidal compounds, flavonoids, tannins & sugars	Mungle et al., 2012
24	<i>Eucommia ulmoides</i> Oliv.	Eucommiaceae	Aq.	Powdered leaf extract	Lee et al., 2005
25	<i>Flacourtia jangomas</i> Raeusch.	Flacourtiaceae	M	Flavonoids, saponins, carbohydrates, steroids, tannins & phenolic compounds	Singh and Singh (2010)
26	<i>Gardenia taitensis</i> A. P. de Candolle	Rubiaceae	E	Alkaloids, phytosterols, carbohydrates & saponins	Maheswari and Gandhimathi (2011)
27	<i>Hedera helix</i> L.	Araliaceae	E	Leaf extract	Zafar et al., 2002
28	<i>Holoptelea integrifolia</i> (Roxb.)	Ulmaceae	M & P.E	Steroids & glycosides	Sharma et al., 2010
29	<i>Hypericum perforatum</i> L.	Hypericaceae	Ethl.ac.	Leaf extract	Arokiyaraj et al., 2011
30	<i>Juglans regia</i> L.	Juglandaceae	M	Leaf extract	Teimori et al., 2010
31	<i>Lagerstroemia speciosa</i> (L.) Pers.	Lythraceae	Aq.	Leaf powder or decoction	Tanquilut et al., 2009
32	<i>Memecylon umbellatum</i> Burm.F.	Melastomataceae	Al.	Leaf extract	Amalraj and Ignacimuthu (1998)
33	<i>Mimosa pudica</i> L.	Mimosaceae	E	Leaf extract	Sutar et al., 2009
34	<i>Mimusops elengi</i> L.	Sapotaceae	M	Leaf extract	Zahid et al., 2012
35	<i>Moringa oleifera</i> Lam.	Moringaceae	Aq.	Leaf extract	Manohar et al., 2012
36	<i>Myrcia uniflora</i> Barb. Rodr.	Myricaceae	Aq.	Leaf extract	Pepato et al., 1993
37	<i>Nauclea latifolia</i> Sm.	Rubiaceae	Aq.	Leaf extract	Gidado et al., 2005
38	<i>Nelumbo nucifera</i> Gaertn.	Nymphaeaceae	E	Flavonoids	Zhou et al., 2009
39	<i>Neolamarckia cadamba</i> (Roxb.) Bossier	Rubiaceae	M	Leaf extract	Ahmed et al., 2011
40	<i>Olea europaea</i> L.	Oleaceae	Al.	Leaf extract	Eidi et al., 2009
41	<i>Passiflora mollissima</i> Bailey	Passifloraceae	E	Alkaloids, tannins	Edwin et al., 2007

				& flavonoids	
42	<i>Piper betle</i> L.	Piperaceae	Aq. & E	Leaf extract	Arambewela et al., 2005
43	<i>Pisonia alba</i> Span.	Nyctaginaceae	E	Vitamin A, alkaloids, proteins & fats	Sunil et al., 2009
44	<i>Premna corymbosa</i> (Burm. F.) Rottl	Verbenaceae	E & Aq.	Alkaloids, flavonoids, glycosides, saponins, terpenes & steroids	Thiruvengkatasubramaniam and Jayakar (2010)
45	<i>Salacia fruticosa</i> Heyne exLawson	Hippocrateaceae	M	Alkaloids, carbohydrates, phytosterols, glycosides, saponins & phenolic compounds	Venkateshwarlu et al., 2009
46	<i>Salacia reticulata</i> Wight	Hippocrateaceae	Aq.	Leaf extract	Yoshino et al., 2009
47	<i>Solanum nigrum</i> L.	Solanaceae	Aq. & hydro- alcoholic	Alkaloids, flavonoids, phenolics	Meonah et al., 2012
48	<i>Solanum trilobatum</i> L.	Solanaceae	Aq.	Leaf extract	Doss et al., 2009
49	<i>Sonneratia alba</i> Sm.	Sonneratiaceae	M	Leaf extract	Morada et al., 2011
50	<i>Spinacia oleracea</i> L.	Chenopodiaceae	E	Leaf extract	Kumar and Loganathan (2010)
51	<i>Symplocos cochinchinensis</i> (Lour.) S. Moore.	Symplocaceae	H	Leaf extract	Sunil et al., 2011
52	<i>Talinum portulacifolium</i> Forssk.	Portulacaceae	M	Steroids, triterpenoids & flavonoids	Rao et al., 2007
53	<i>Urtica dioica</i> L.	Urticaceae	Aq.	Leaf extract	Das et al., 2011
54	<i>Urtica parviflora</i> roxb.	Urticaceae	Aq.	Alkaloids, reducing sugars, polysaccharides, tannins, saponins, glycosides & flavonoids	Sah et al., 2010
55	<i>Withania somnifera</i> (L.) Dunal	Solanaceae	E	Flavonoids	Udayakumar et al., 2009
56	<i>Zizyphus sativa</i> Gaertn.	Rhamnaceae	Al.	Leaf extract	Anand et al., 1989
Barks					

57	<i>Adansonia digitata</i> L.	Bombacaceae	M	Tannins, carbohydrates, terpenes, saponins, flavonoids & alkaloids	Tanko et al., 2008
58	<i>Albizia odoratissima</i> Benth.	Mimosaceae	M	Bark extract	Kumar et al., 2011
59	<i>Cinnamomum verum</i> J. S. Presl	Lauraceae	Aq.	Bark extract	El-Desoky et al., 2012
60	<i>Commiphora africana</i> (A.Rich.) Engl.	Burseraceae	Aq. & E	Alkaloids, tannins, flavonoids, steroids & saponins	Goji et al., 2009
61	<i>Crataeva nurvala</i> Buch. Ham.	Capparidaceae	P.E, Aq., Chlf. & Al.	Triterpenoids & flavonoids	Sikarwar and Patil (2010)
62	<i>Diospyros melanoxylon</i> Roxb.	Ebenaceae	E	Triterpenoids, steroids, alkaloids, flavonoids & tannins	Gupta et al., 2009
63	<i>Helicteres isora</i> L.	Sterculiaceae	Aq.	Bark extract	Kumar et al., 2009
64	<i>Madhuca indica</i> J. F. Gmel.	Sapotaceae	M	Bark extract	Choudhary et al., 2011
65	<i>Musanga cecropioides</i> R. Br. Ex Bennet	Cecropiaceae	E & Aq.	Bark extract	Adeneye et al., 2007
66	<i>Parinari excelsa</i> Sougue	Rosaceae	Aq.	Bark extract	Ndiaye et al., 2008
67	<i>Polyalthia longifolia</i> var. <i>angustifolia</i> Thw.	Annonaceae	M	Alkaloids, triterpenoids, flavonoids, steroids, saponins, glycosides & tannins	Gosh et al., 2010
68	<i>Strychnos henningsii</i> Gilg.	Loganiaceae	Aq.	Flavonoids, tannins & saponins	Oyedemi et al., 2012
69	<i>Triplochiton scleroxylon</i> Schumann	Sterculiaceae	Aq.	Bark extract	Prohp and Onoagbe (2011)
Roots					
70	<i>Anthocleista djalensis</i> A. Chev	Loganiaceae	E	Flavonoids, saponins, tannins, cardiac glycosides & anthraquinones	Okokon et al., 2012
71	<i>Asparagus racemosus</i> Willd.	Liliaceae	E	Root extract	Hannan et al., 2012
72	<i>Berberis lyceum</i> Royle	Berberidaceae	E & Aq.	Root extract	Gulfraz et al., 2007
73	<i>Casearia esculenta</i> Roxb.	Samydaceae	E	Alkaloids, glycosides, saponins, phytosterols, tannins & amino acids	Arul et al., 2006
74	<i>Costus speciosus</i> (Koenex.Retz.)	Costaceae	E	Root extract	Bavarva et al., 2008

75	<i>Lantana aculeata</i> L.	Verbenaceae	E	Root extract	Kumar et al., 2010
76	<i>Merremia tridentata</i> (L.) Hall. f.	Convolvulaceae	Aq.	Root extract	Arunachalam and Parimelazhagan (2012)
77	<i>Nyctanthes arbor-tristis</i> L.	Oleaceae	M	Root extract	Sharma et al., 2011
78	<i>Pandanus fascicularis</i> Lam.	Pandanaceae	Aq & E	Carbohydrates, proteins, amino acids, saponins, tannins, phenolic compounds, alkaloids & flavonoids	Rajeswari et al., 2012
79	<i>Pandanus odoratus</i> RIDL.	Pandanaceae	Aq.	Root extract	Peunqvicha
80	<i>Potentilla fulgens</i> L.	Rosaceae	CM	Root extract	Syiem et al., 2002
81	<i>Tectona grandis</i> L.	Verbenaceae	M	Root extract	Pooja et al., 2011
82	<i>Withania somnifera</i> (L.) Dunal	Solanaceae	E	Flavonoids	Udayakumar et al., 2009
83	<i>Zaleya decandra</i> L. N. Burm. f.	Aizoaceae	E	Root extract	Meenakshi et al., 2010
Fruits					
84	<i>Balanites aegyptiaca</i> (L.) Delile	Balanitiaceae	Aq. & E	Fruit flesh extract	Zaahkhouk et al., 2003
85	<i>Diospyros lotus</i> L.	Ebenaceae	Aq.	Fruit extract	Azadbakhta et al., 2010
86	<i>Fomitopsis pinicola</i> (Swartz. Fries) Karst.	Fomitopsidaceae	Aq. & Alkali	Fruit body extract	Lee et al., 2008
87	<i>Hericium erinaceus</i> (Bull.) Pers.	Hericiaceae	M	Fruiting bodies extract	Wang et al., 2005
88	<i>Musa paradisiaca</i> L.	Musaceae	M	Fruit extract	Ojewole and Adewunmi (2003)
89	<i>Phaleria macrocarpa</i> (Scheff.) Boerl (Pm)	Thymelaeaceae	M	Flavonoids, terpenoids & tannins	Ali et al., 2012
90	<i>Rosa canina</i> L.	Rosaceae	E	Fruit extract	Orhan et al., 2009
91	<i>Solanum nigrum</i> L.	Solanaceae	Aq. & hydro alcoholic	Alkaloids, flavonoids, phenolics & micronutrients	Meonah et al., 2012
92	<i>Terminalia belerica</i> Roxb.	Combretaceae	M	Triterpenoids-arjungenin, bellericagenins & belleric acid	Sabu and Kuttan (2009)
93	<i>Terminalia pallida</i> Brandis	Combretaceae	E	Flavonoids, phenolic acids, sterols/	Kameswara et al., 2003

				triterpenoid, alkaloids, tannins & anthocyanins	
94	<i>Trapa natans</i> L.	Lythraceae	M	Fruit peel extract	Das et al., 2011
Seeds					
95	<i>Brassica juncea</i> L.	Brassicaceae	Aq.	Seed extract	Thirumalai et al., 2011
96	<i>Carica papaya</i> L.	Caricaceae	Aq.	Alkaloids, flavonoids, saponins, tannins, anthraquinones, anthocyanosides & reducing sugars	Adeneye and Olagunju (2009)
97	<i>Carum carvi</i> L.	Apiaceae	E	Carvone, limonene, carveol, dihydrocarveol & thymol	Eidi et al., 2010
98	<i>Hippophae rhamnoides</i> L.	Elaegnaceae	Aq.	Seed extract	Zhang et al., 2010
99	<i>Irvingia gabonensis</i> (Aubry- Lecomte) Baill.	Irvingiaceae	Aq.	Seed extract	Ozolua et al., 2006
100	<i>Lepidium sativum</i> L.	Brassicaceae	Aq.	Seed extract	Eddouks et al., 2005
101	<i>Nigella sativa</i> L.	Ranunculaceae	Aq.	Essential oils, proteins, alkaloids & saponins	Mathur et al., 2011
102	<i>Persea americana</i> Mill.	Lauraceae	E	Seed extract	Edem (2009)
103	<i>Spergularia purpurea</i> (Pers.) G.Don. Fil	Caryophyllaceae	Aq.	Saponins & flavonoids	Jouad et al., 2000
104	<i>Strychnus potatorum</i> L.	Loganiaceae	E & Aq.	Seed extract	Raghu et al., 2011
105	<i>Terminalia chebula</i> Retz.	Combretaceae	Chlf.	Seed extract	Rao et al., 2006
Aerial Parts					
106	<i>Bacopa monnieri</i> L.	Scrophulariaceae	E	Aerial parts extract	Ghosh et al., 2008
107	<i>Equisetum myriochaetum</i> Schlecht and Cham	Equisetaceae	Aq. & B	Flavonol glycosides & one Caffeoylglycoside	Choudhary et al., 2011
108	<i>Laportea ovalifolia</i> Schamand Thonn.	Urticaceae	Aq.	Aerial part extract	Momo et al., 2006
109	<i>Lycium shawii</i> Roem and Schult	Solanaceae	E	Aerial part extract	Sher et al., 2011
110	<i>Salvadora oleoides</i> Decne	Salvadoraceae	E	Aerial part extract	Yadav et al., 2008
111	<i>Scoparia dulcis</i> L.	Scrophulariaceae	E	Alkaloids, carbohydrates, glycosides, flavonoids &	Zulfiker et al., 2010

				tannins	
112	<i>Suaeda fruticosa</i> Forssk. ex J. F. Gmel.	Chenopodiaceae	Aq.	Aerial part extract	Benwahhoud et al., 2001
113	<i>Zygophyllum gaetulum</i> Emb. and Maire	Zygophyllaceae	Aq.	Aerial part extract	Jaouhari et al., 2000
114	<i>Zygophyllum geslini</i> Coss.	Zygophyllaceae	Aq.	Aerial parts extract	Medjdoub et al., 2012
Flower					
115	<i>Antigonon leptopus</i> Hook & Arn.	Polygonaceae	M	Flower extract	Sujatha et al., 2012
116	<i>Kigelia pinnata</i> Jacq.	Bignoniaceae	M	Flower extract	Kumar et al., 2012
117	<i>Mimusops elengi</i> L.	Sapotaceae	M	Flower extract	Sidhu et al., 2013
118	<i>Nymphaea stellata</i> Willd.	Nymphaeaceae	E	Flower extract	Rajagopal and Sasikala (2008)
Whole Plant					
119	<i>Heliotropium zeylanicum</i> (Burm.F) Lamk	Boraginaceae	M	Plant extract	Muruges et al., 2006
120	<i>Hemionitis arifolia</i> (Burm.) Moore.	Polypodiaceae	Aq.	Plant extract	Nair et al., 2006
121	<i>Hybanthus enneaspermus</i> (L.) F. Muell	Violaceae	E	Flavonoids, terpenes, phenols, anthraquinones, glycosides, polyoses, alkaloids, saponins & tannins	Patel et al., 2011
122	<i>Jussiaea suffruticosa</i> L.	Onagraceae	M	Plant extract	Murugesan et al., 2000
123	<i>Merremia emarginata</i> Burm. F.	Convolvulaceae	M	Plant extract	Gandhi and Sasikumar (2012)
124	<i>Mollugo nudicaulis</i> Lam.	Aizoaceae	E	Plant extract	Sindhu et al., 2010
125	<i>Piper sarmentosum</i> Roxb.	Piperaceae	Aq.	Plant extract	Peunqvicha et al., 1998
126	<i>Polygala javana</i> DC.	Polygalaceae	E	Alkaloids, catechism, tannins, saponins, steroids, flavonoids, phenols, sugar, glycosides & xanthoprotein	Alagammal et al., 2012
127	<i>Portulaca oleracea</i> L.	Portulacaceae	Chlf, M	Plant extract	Syiem et al., 2002
128	<i>Trianthema portulacastrum</i> L.	Aizoaceae	M	Alkaloids, flavonoids, saponins, phenolic compounds & terpenoids	Sunder et al., 2009

129	<i>Triumfetta pilosa</i> Roth	Tiliaceae	E	Plant extract	Ramakrishna et al., 2011
Rhizome					
130	<i>Acorus calamus</i> L.	Acoraceae	M	Rhizome extract	Prisilla et al., 2012
131	<i>Alpinia galanga</i> Willd.	Zingiberaceae	M & Aq.	Rhizome extract	Akhtar et al., 2002
132	<i>Sansevieria roxburghiana</i> Schult. and Schult. f.	Agavaceae	Aq. & E	Alkaloids, triterpenes, steroids, flavonoids & saponins	Haldar et al., 2010
Root bark					
133	<i>Blighia sapida</i> K. Kong	Sapindaceae	Aq.	Terpenoids, phenol, alkaloids, tannins	Saidu et al., 2012
134	<i>Ceiba pentandra</i> (L.) Gaertner	Bombacaceae	Mthl. Cl. & M	Root bark extract	Djomeni et al., 2006
135	<i>Euclea undulata</i> Thunb. var myrtina	Ebenaceae	Act.	Root bark extract	Deutschlander et al., 2012
Stem					
136	<i>Berberis aristata</i> DC.	Berberidaceae	M	Alkaloids, glycosides, carbohydrates, bitter principles & saponins	Upwar et al., 2010
137	<i>Flacourtia jangomas</i> Raeusch.	Flacourtiaceae	M	Flavonoids, saponins, carbohydrates, steroids, tannins & phenolic compounds	Singh and Singh 2010
138	<i>Nervilia plicata</i> (Andrews) Schltr.	Orchidaceae	Al.	Stem extract	Kumar and Janardhana (2011)
139	<i>Tournefortia hirsutissima</i> L.	Boraginaceae	Aq. & B	Stem extract	Andrade-Cetto et al., 2007
Tubers					
140	<i>Anaphyllum wightii</i> Schott.	Araceae	E	Tuber extract	Mathew et al., 2013
141	<i>Nymphaea pubescens</i> Willd.	Nymphaeaceae	E	Alkaloids, flavonoids, glycosides, terpenoids, tannins, phenols, saponins & steroids	Shajeela et al., 2012

Abbreviations: Aq.- Aqueous, Acetone- Act., Al.- Alcohol, B- Buthanol, E- Ethanol, H- Hexane, M.- Methanol, Chlf.- Chloroform, CM- Crude Methanol, Ethl.ac.-Ethyl acetate, Methl. Cl.- Methyl Chloride, PE- Petroleum Ether.

Table 3: Number of plant species in different families


Acoraceae	1	Eucommiaceae	1	Piperaceae	2
Agavaceae	1	Flacourtiaceae	1	Polygonaceae	2
Aizoaceae	3	Fomitopsidaceae	1	Polypodiaceae	1
Alangiaceae	1	Hericiaceae	1	Portulacaceae	2
Annonaceae	2	Hippocrateaceae	2	Ranunculaceae	1
Apiaceae	1	Hypericaceae	1	Rhamnaceae	1
Araceae	1	Irvingiaceae	1	Rosaceae	3
Araliaceae	1	Juglandaceae	1	Rubiaceae	3
Arecaceae	1	Lauraceae	3	Salvadoraceae	1
Balanitiaceae	1	Liliaceae	2	Samydaceae	1
Basellaceae	1	Loganiaceae	3	Sapindaceae	2
Berberidaceae	2	Lythraceae	2	Sapotaceae	2
Bignoniaceae	2	Melastomataceae	1	Scrophulariaceae	2
Bombacaceae	2	Melanthaceae	1	Solanaceae	4
Boraginaceae	2	Mimosaceae	2	Sonneratiaceae	1
Burseraceae	3	Moringaceae	1	Sterculiaceae	2
Capparidaceae	1	Musaceae	1	Symplocaceae	1
Caricaceae	1	Myricaceae	1	Thymelaeaceae	1
Caryophyllaceae	1	Nyctaginaceae	3	Tiliaceae	1
Cecropiaceae	3	Nymphaeaceae	3	Ulmaceae	1
Chenopodiaceae	3	Oleaceae	2	Urticaceae	3
Combretaceae	4	Onagraceae	1	Verbenaceae	4
Convolvulaceae	2	Orchidaceae	1	Violaceae	1
Costaceae	2	Oxalidaceae	1	Vitaceae	1
Crassulaceae	1	Palmaceae	1	Zingiberaceae	2
Ebenaceae	3	Pandanaceae	2	Zygophyllaceae	2
Elaegnaceae	1	Passifloraceae	1		
Equisetaceae	1	Primulaceae	1		

Plants have also been categorized on the basis of plant part used (Table 4). Leaves were found to be the most frequently used plant part (56) in the management of diabetes, followed by roots (14), bark (13), fruits (11), seeds (11), whole plants (11), and aerial parts (9) have shown activity against diabetes. Rhizome (3), flower (4), root bark (3), stem (4), tubers (2) also possess anti-diabetic activity.

Table 4: Various plant parts used & number of their plant species

Plant Parts Used	Number of Species
Leaf	56
Root	14
Bark	13
Fruit	11
Seed	11
Whole plant	11
Aerial Part	9
Flower	4
Steam	4
Rhizome	3
Root bark	3
Tuber	2

Extraction is the first crucial step in preparation of plant formulations. Considerable effort has been made by researchers to find efficient extraction methods in order to get high efficiency and efficacy. It should be noted that choice of appropriate solvent is of essential importance along with application of a compatible extraction method (Gupta et al. 2012). In this review of 141 anti-diabetic plants, the most popular used solvent to make plant extract was aqueous solvent 39.7% among the 141 species, after then ethanol 25.5%, methanol 20.5%, alcohol 3.7%, and chloroform 2.5%. The rest of the used solvents were buthanol, petroleum ether 1.9% each, hydro alcoholic 1.2% and acetone, crude methanol, hexane, methyl chloride & ethyl acetate 0.62 % respectively (fig 1).



4. Hypoglycemic Constituents and Mechanisms of Action of Phytochemicals

It has been studied that bioactive plant, secondary metabolites are important in chemical defense mechanisms (Ehrlich and Raven 1964; Berenbaum 1983). Glucose is metabolic energy source and this undergoes storage and

mobilization under hormonal control. Plant growth regulators such as indole-3-acetic acid and similarly indole-3-butyric acid, L-tryptophan, indole-3-propionic acid and p-chlorophenoxyacetic acid, inhibits insulinase in vitro and are hypoglycemic in vivo (Mirsky et al. 1956). In the same way, insulinase and potentiate administered insulin with the help of anthranilic and nicotinic acid. An efficient hypoglycemic activity was observed when an inhibitor of indole-3-acetic acid oxidase from *Phaseolus vulgaris* (fruit exocarps) was applied for diabetic treatment, similarly hypoglycemic alkaloid trigonelline from *Trigonella foenum-graecum*, is a growth inhibitor helps in hypoglycemic activity. In the same way, another plant growth inhibitor is "Salicylic Acid" (Oliver- Bever and Zahnd, 1979). So plant metabolism regulating constituents can also be anti-diabetic, animal metabolism-regulatory agents.

Cinnamaldehyde, a phytoconstituent extracts have been reported to exhibit significant antihyperglycemic effect resulting in the lowering of both total cholesterol and triglyceride levels and, at the same time, increasing HDL-cholesterol in STZ-induced diabetic rats. This investigation reveals the potential of cinnamaldehyde for use as a natural oral agent, with both hypoglycemic and hypolipidemic effects. Recent reports indicate that Cinnamon extract and polyphenols with procyanidin type-A polymers exhibit the potential to increase the amount of TTP (Thrombotic Thrombocytopenic Purpura), IR (Insulin Resistance), and GLUT4 (Glucose Transporter-4) in 3T3-L1 Adipocytes. It was suggested that the mechanism of Cinnamon's insulin-like activity may be in part due to increase in the amounts of TTP, IR (Insulin Resistance), and GLUT4 and that Cinnamon polyphenols may have additional roles as anti-inflammatory and/or anti-angiogenesis agents (Jakhelia et al., 2010).

Summary

In Present world, the word "Diabetes" has got striking significance because this disease with its causes and effects in different ages. One of the important thing is impediment in the cure of this syndrome, which provides chief importance and critical analysis of it, to provide vital treatments of diabetes so that this disease will not affect succeeding generations. For a long time, remedy of it has been under surveillance, and for this medicinal plants played their vital role, as nature has sovereignty in a particular way to overcome harmful effects of diabetes. So this document is an endeavor to assemble those patent medicinal plants that are with benevolent use in cure of syndrome diabetes. As 141 plant species with all their particulars and data are being mentioned in this review article, especially with their action of mechanism to help in apprehension role of anti-diabetic plants. All contents of this paper, especially "tables" with their brief information about anti-diabetic medicinal plants are cognition source with systemize details. This review paper with all its particulars is compile work to elevate importance of plants in treatment of threatening syndrome diabetes so that we might be able of secure world from its vulnerable risks in near future.

Conflict of interest statement

We declare that we have no conflict of interest.

Authors' contribution

The authors are thankful to Beauty Akter Rupa for her gracious contribution in this work. All the authors edited the paper, read and approved the final paper.

References:

- ADA, (1997): Clinical practice recommendation 1997, Screening for diabetes. *Diabetes care* 20(1), 22-24.
- Adebayo GI., Alabi OT., Owoyele BV., Soladoye. (2009): Antidiabetic properties of the aqueous leaf extract of *Bougainvillea glabra* (Glory of the garden) on alloxan- induced diabetic rats. *Rec. Nat. Prod.* 3(4): 187-192.
- Adeneye AA, Olagunju, JA. (2009): Preliminary hypoglycemic and hypolipidemic activity of the aqueous seed extract of *Carica papaya* Linn in wistar rats. *Biology and Medicine.* 1(1): 1-10.
- Adeneye AA, Adeleke TI, Adeneye AK. (2008): Hypoglycemic and hypolipidemic effects of the aqueous fresh leaves extract of *Clerodendrum capitatum* in wistar rats. *J. Ethnopharmacol.* 116(1): 7-10.
- Adeneye AA, Ajagbonna OP, Ayodele OW, (2007): Hypoglycemic and antidiabetic activities on the stem bark aqueous and ethanol extracts of *Musanga cecropioides* in normal and alloxan-induced diabetic rats. *Fitoterapia*, 78: 502-505.

- Ahmed F., Rahman S., Ahmed N., Hossain M., Biswas A., et al., (2011): Evaluation of *Neolamarckia cadamba* (Roxb.) Bosser leaf extract on glucose tolerance in glucose-induced hyperglycaemic mice. *Afr. J. Tradit. Complement. Alter. Med.* 8(1): 79-81.
- Akhtar MS, Khan Ma, Malik MT, (2002): Hypoglycemic activity of *Alpinia galanga* rhizome and its extracts in rabbits. *Fitoterapia*, 73(7-8): 623-628.
- Alagammal M, Agnel RA, Mohan VR, (2012): Antidiabetic and antihyperlipidemic effect of *Polygala javana* DC on alloxan induced diabetic rats. *Int. Res. J.Pharm.* 3(9): 231-234.
- Alarcon-Aguilara FJ, Roman-Ramos R, Perez-Gutierrez S, Aguilar-Contreras A, Contreras-Weber CC, Flores-Saenz JL. (1998): Study of the anti-hyperglycemic effect of plants used as antidiabetics. *J Ethnopharmacol.* 61: 101-110.
- Ali RB et al. (2012): Hypoglycemic and anti-hyperglycemic study of *Phaleriana crocarpa* fruits pericarp. *J Med. Plant. Res.* 6(10): 1982-1990.
- Amalraj T, Ignacimuthu S. (1998): Evaluation of the hypoglycemic effect of *Memecylon umbellatum* in normal and alloxan diabetic mice. *J. Ethnopharmacol.* 62(3): 247-250.
- Anand KK, Singh B, Chand D, Chandan BK, Gupta VN. (1989): Effect of *Zizyphus sativa* leaves on blood glucose levels in normal and alloxan-induced diabetic rats. *J. Ethnopharmacol.* 27(1-2): 121-127.
- Andrade-Cetto A, Wiedenfeld H. (2001): Hypoglycemic effect of *Cecropia obtusifolia* on streptozotocin diabetic rats. *J. Ethnopharmacol.* 78: 145-149.
- Andrade-Cetto A, Revilla-Monsalve C, Wiedenfeld H. (2007): Hypoglycemic effect of *Tournefortia hirsutissima* L. on n-streptozotocin diabetic rats. *J. Ethnopharmacol.* 112(1): 96-100.
- Ang-Lee MK, Moss J, Yuan CS. (2001): Herbal medicines and perioperative care. *JAMA.* 286(2): 208-16.
- Aragao DM, Guarize L, Lanini J, Costa JC, Garcia RM, Scio E, (2010): Hypoglycemic effect of *Cecropia pachystachya* in normal and alloxan-induced diabetic rats. *J. Ethnopharmacol.* 128(3): 629-633.
- Arambewela LS, Arawawala LD, Ratnasooriya WD, (2005): Antidiabetic activities of aqueous and ethanolic extracts of *Piper betle* leaves in rats. *J. Ethnopharmacol.* 102: 239-245.
- Arokiyaraj S, Balamurugan R, Augustian P. (2011): Antihyperglycemic effect of *Hypericum perforatum* ethyl acetate extract on streptozotocin-induced diabetic rats. *Asian Pac. J Trop.Biomed.* 1(5): 386-390.
- Arul B, Kothai R, Christina AMJ. (2006): Hypoglycemic activity of *Casearia esculenta* in normal and diabetic albino rats. *Iran J Pharm Res.* 1: 47-51.
- Arunachalam K, Parimelazhagan T. (2012): Anti-diabetic activity of aqueous root extract of *Merremia tridentata* (L.) Hall. f. in streptozotocin-induced-diabetic rats. *AsianPac. J Trop. Med.* 5(3): 175-179.
- Azadbakhta M, Safapour S, Ahmadi A, Ghasemi M, Shokrzadeh M. (2010): Anti-diabetic effects of aqueous fruits extract of *Diospyros lotus* L. on streptozotocin-induced diabetic rats and the possible morphologic in the liver, kidney, and heart. *Journal of Pharmacognosy and Phytotherapy*, 2(2): 010-016.
- Balick MJ, Cox PAR. (1996): Plants people and culture. The science of ethnobotany. Scientific American Library, New York, USA. pp: 219.
- Bailey CJ and Day C. (1989): Traditional plant medicines as treatments for diabetes. *Diabetes Care*, 12, American diabetes association, 553-564.

- Bavarva JH, Narasimhacharya AV. (2008): Antihyperglycemic and hypolipidemic effects of *Costus speciosus* in alloxan induced diabetic rats. *Phytother Res.* 22(5): 620-626.
- Benwahhoud M, Jouad H, Eddouks M, Lyoussi B. (2001): Hypoglycemic effect of *Suaeda fruticosa* in streptozotocin-induced diabetic rats. *J Ethnopharmacol.* 76(1): 35-38.
- Berenbaum M. (1983): Coumarins and caterpillars: a case for co-evolution. *Evolution*, 37: 163-179.
- Cetto AA, Wiedenfeld H, Revilla MC, Sergio IA. (2000): Hypoglycemic effect of *Equisetum myriochaetum* aerial parts on streptozotocin diabetic rats. *J Ethnopharmacol.* 72 (1-2): 129-133.
- Chakraborty U, Das H. (2010): Antidiabetic and antioxidant activities of *Cinnamomum tamala* leaf extracts in STZ-treated diabetic rats. *Glob. J Biotech.Biochem.* 5(1): 12-18.
- Chang CH, Ngoh GC, Yusoff R. (2012): A brief review on anti diabetic plants: Global distribution, active ingredients, extraction techniques and acting mechanisms. *Pharmacogn Rev.* 6(11): 22-8.
- Chika A, Bello SO. (2010): Antihyperglycemic activity of aqueous leaf extract of *Combretum micranthum* (Combretaceae) in normal and alloxan-induced diabetic rats. *J Ethnopharmacol* 129(1): 34-37.
- Choudhary A et al. (2011): Anti-diabetic activity of the methanolic extract of *Madhuca indica* on normal and streptozotocin induced diabetic rats. *Int. J Pharm. Res. Dev.*, 3(4): 13-18.
- Cunningham AB. An Investigation of the herbal medicine trade in Natal/KwaZulu. Pietermaritzburg: Investigational Report No 29 Institute of Natural Resources, 1988.
- Das PK, Bhattacharya S, Pandey JN, Biswas M. (2011): Antidiabetic activity of *Trapa natans* fruit peel extract against streptozotocin induced diabetic rats. *Glob. J Pharm.* 5(3): 186-190.
- Deutschlander MS, Lall N, Venter MV de, Dewanjee S. (2012): The hypoglycemic activity of *Euclea undulata* Thunb. var. *myrtina* (Ebenaceae) root bark evaluated in a streptozotocin-nicotinamide induced type-2 diabetes rat model. *South African Journal of Botany*, 80(5): 9-12.
- Djomeni PDD et al. (2006): Hypoglycemic and antidiabetic effect of root extracts of *Ceiba pentandra* in normal and diabetic rats. *Afr. J Trad. Complemen. Altern. Med.* 3(1): 129-136.
- Doss A, Palaniswamy M, Angayarkanni J, Dhanabalan R. (2009): Antidiabetic activity of water extract of *Solanum trilobatum* (L.) in alloxan-induced diabetes in rats. *Afr J Biotech.* 8(20): 5562-5564.
- Eddouks M, Maghrani M, Zeggwagh NA, Michel JB. (2005): Study of the hypoglycemic activity of *Lepidium sativum* L. aqueous extract in normal and diabetic rats. *J Ethnopharmacol.* 97(2): 391-395.
- Edem DO. (2009): Hypoglycemic effects of ethanolic extracts of Alligator Pear seed (*Persea americana* Mill.) in rats. *European Journal of Scientific Research.* 33(4): 669-678.
- Edwin E, Sheeja E, Dhanabal S, Suresh B. (2007): Antihyperglycemic activity of *Passiflora mollissima* Bailey. *Indian Journal of Pharmaceutical Sciences.* 69(4): 570-771.
- Ehrlich PR., Raven P.H. (1964): Butterflies and plants: A study in coevolution. *Evolution* 18(4): 586-608.
- Eidi A, Eidi M, Darzi R. (2009): Antidiabetic effect of *Olea europaea* L. in normal and diabetic rats. *Phytother. Res.* 23(3): 347-350.

Eidi A, Eidi M, Haeri RA, Basati F. (2010): Hypoglycemic effect of ethanolic extract of *Carum carvi* L. seeds in normal and streptozotocin-induced diabetic rats. J Med. Plant. 9(35): 106-113.

El-Desoky GE, Aboul-Soud MAM, AL-Numair KS. (2012): Antidiabetic and hypolipidemic effects of *Ceylon cinnamon* (*Cinnamomum verum*) in alloxan-diabetic rats. J Med.Plant Res. 6(9): 1685-1691.

Gaamoussi F, Israili ZH, Lyoussi B. (2010): Hypoglycemic and hypolipidemic effects of an aqueous extract of *Chamaerops humilis* leaves in obese, hyperglycemic and hyperlipidemic meriones shawi rats. Pak J Pharm. Sci. 23(2): 212-219.

Gandhi GR, Sasikumar P. (2012): Antidiabetic effect of *Merremia emarginata* Burm. F. in streptozotocin induced diabetic rats. Asian Pac. J Trop. Biomed. 2(4): 281-286.

Ghosh T, Maity TK, Sengupta P, Dash DK, Bose A. (2008): Antidiabetic and invivoantioxidant activity of ethanolic extract of *Bacopa monnieri* Linn. aerial parts: a possible mechanism of action. Iran J Pharm.Res. 7(1): 61-68.

Gidado A., Ameh D.A., and Atawodi S.E. (2005): Effect of *Nauclea latifolia* leaves aqueous extracts on blood glucose levels of normal and alloxan-induced diabetic rats. African Journal of Biotechnology, 4(1): 91-93.

Gilani AH, Rahman AU. (2005): Trends in ethnopharmacology. J Ethnopharmacol. 100:43-49.

Goji A.D.T., Dikko A.A.AU., Bakari A.G., Mohammed A., and Tanko Y. (2009): Evaluation of the effect of aqueous-ethanolic stem bark extract of *Commiphora africana* on blood glucose levels of alloxan induced diabetic rats. AsianJournal of Medical Sciences, 1(2): 18-21.

Gosh G., M. Kar D., B. Subudhi B. and K. Mishra S. (2010): Anti-hyperglycemic and antioxidant activity of stem bark of *Polyalthia longifoliavar. angustifolia*. Der Pharmacia Lettre, 2(2): 206-216.

Gover JK, Yadav S, Vats V. (2002): Medicinal Plants of India with anti-diabetic Potential. J EthnoPharmacol. 81(1):81-100.

Gulfranz M., Qadir G., Nosheen F., Parveen Z. (2007): Antihyperglycemic effects of *Berberis lyceum* Royle in alloxan-induced diabetic rats. DiabetologiaCroatica, 36(3): 49-54.

Gupta A, Naraniwal M and Kothari V. (2012): Modern extraction methods for preparation of bioactive plant extracts. International Journal of Applied and Natural Sciences, 1(1): 8-26.

Gupta RK, Kesari AN, Watal G, Murthy P.S., et al., (2005): Hypoglycemic and antidiabetic effect of aqueous extract of leaves of *Annona squamosa* (L.) in experimental animal. Current Science, 88(8): 1244-1254.

Gurudeeban S, Satyavani K, Ramanathan T and Balasubramanian T. (2012): Antidiabetic effect of a black mangrove species *Aegiceras corniculatum* in alloxan-induced diabetic rats. Journal of Advanced Pharmaceutical Technology and Research, 3(1): 52-56.

Haldar P.K., Kar B., Bhattacharya S., Bala A., Kumar S.R.B. (2010): Antidiabetic activity and modulation of antioxidant status by *Sansevieria roxburghiana* rhizome in streptozotocin-induced diabetic rats. Diabetologia Croatica, 39(4): 115-123.

Hannan JM, Ali L, Khaleque J, Akhter M, Flatt PR, Abdel-Wahab YH. (2012): Antihyperglycemic activity of *Asparagus racemosus* roots is partly mediated by inhibition of carbohydrate digestion and absorption and enhancement of cellular insulin action. Br J Nutr. 107: 1316-1323.

Hill AF. A textbook of useful plants and plant products, 2nd ed.; Economic Botany, Mc Garw- Hill Book Company Inc, New York, 1952.

- Jakhetia V, Patel R, Khatri P, Pahuja N, Garg S, Pandey A and Sharma SA. (2010): *Cinnamon*: a pharmacological review. *Journal of Advanced Scientific Research*, 1(2): 19-23.
- Jaouhari J.T., Lazrek H.B., Jana M. (2000): The Hypoglycaemic activity of *Zygophyllum gaetulum* extracts in alloxan induced hyperglycemic rats. *J Ethnopharmacol.* 69: 17-20.
- Jouad H, Eddouks M, Lacaille-Dubois MA, Lyoussi B. (2000): Hypoglycaemic effects of *Spergularia purpurea* in normal and streptozotocin-induced diabetic rats. *J Ethnopharmacol.* 71: 169-177.
- Joy PP, Thomas J, Mathew S, Skaria BP. (1998): Medicinal plants, Kerala Agricultural University, Aromatic and Medicinal Plants Research Station.
- Kambouche N., Merah B., Derdour A., Bellahouel S., et al. (2009): Hypoglycemic and antihyperglycemic effects of *Anabasis articulata* (Forssk) Moq. (Chenopodiaceae), an Algerian Medicinal plant. *African Journal of Biotechnology*, 8(20): 5589-5594.
- Kameswara RB, Renuka SP, Rajasekhar MD, Nagaraju N, Appa RC. (2003): Antidiabetic activity of *Terminalia pallida* fruit in alloxan induced diabetic rats. *J Ethnopharmacol.* 85: 169-172.
- Koffi NG, Yvette FNGB, Henri KK. (2011): et al. Effect of aqueous extract of *Boerhaavia diffusa* leaves in the glycaemia of rabbits. *International Journal of Applied Biology and Pharmaceutical Technology*, 2(3): 330-338.
- Kumar S, Shukla YN, Lavania UC, Sharma A and Singh AK. (1997): Medicinal and Aromatic Plants: Prospects for India. *J. Med. Arom. Pl. Sc.* 19 (2):361-365.
- Kumar D, Kumar S, Kohli S, Arya R, Gupta J. (2011): Antidiabetic activity of methanolic bark extract of *Albizia odoratissima* Benth. in alloxan induced diabetic albino mice. *Asian Pac J Trop Med.* 4(11): 900-903.
- Kumar EK, Janardhana GR. (2011): Antidiabetic activity of alcoholic stem extract of *Nervilia plicata* in streptozotocin-nicotinamide induced type 2 diabetic rats. *J of Ethnopharmacol.* 133(2): 480-483.
- Kumar G et al. (2009): Anti-diabetic activity of *Helicteres isora* L. bark extracts on streptozotocin-induced diabetic rats. *International Journal of Pharmaceutical Sciences and Nanotechnology*, 1(4): 379-382.
- Kumar KV et al. (2010): Antidiabetic potential of *Lantana aculeata* root extract in alloxan-induced diabetic rats. *International Journal of Phytomedicine*, 2: 299-303.
- Kumar NJ, Loganathan P. (2010): Hypoglycaemic effect of *Spinacia oleracea* in alloxan induced diabetic rat. *Global Journal of Biotechnology and Biochemistry*, 5(2): 87-91.
- Kumar R, Pate DK, Prasad SK, Siram k, Hemalatha S. (2011): Antidiabetic activity of alcoholic leaves extract of *Alangium lamarckii* Thwaites on streptozotocin-nicotinamide induced type 2 diabetic rats. *Asian Pac JTrop Med.* 4(11): 904-909.
- Kumar S, Kumar V, Prakash OM. (2012): Antidiabetic and hypolipidemic activities of *Kigelia pinnata* flowers extract in streptozotocin induced diabetic rats. *Asian Pac J Trop Biomed.* 2(7): 543-546.
- Lee MK, Kim MJ, Cho SY, Park SA, Park KK, Jung UJ, Park HM, Choi MS. (2005): Hypoglycemic effect of *Du-zhong* (*Eucommia ulmoides* Oliv.) leaves in streptozotocin-induced diabetic rats. *Diabetes Res Clin Pract.* 67(1): 22-28.

- Lee SI, Kim JS, Oh SH, Lee HG, Kim SD. (2008): Antihyperglycemic effect of *Fomitopsis pinicola* extracts in streptozotocin-induced diabetic rats. *J Med Food*. 11(3): 518-524.
- Li F, Li Q, Gao D, Peng Y and Feng C. (2009): Preparation and antidiabetic activity of polysaccharide from *Portulaca oleracea* L. *African Journal of Biotechnology*, 8(4): 569-573.
- Lotliker MM, Rao MRR. (1996): Pharmacology of a hypoglycemic principle isolated from fruits of *Momordica charantia* Linn. *The Indian Journal of Pharmacy*, 28: 129-133.
- Maheswari JU, Gandhimathi R. (2011): Hypoglycemic and hypolipidemic activity of leaves of *Gardenia taitensis* on streptozotocin induced diabetic rats. *Indian Journal of Pharmaceutical Science and Research*, 1(1): 10-14.
- Malviya N, Jain S, Malviya S. (2010): Antidiabetic potential of medicinal plants. *Acta Pol Pharm*. 67(2):113–118.
- Manohar VS, Jayasree T., Kishore K.K., Rupa M.L., et al. (2012): Evaluation of hypoglycemic and antihyperglycemic effect of freshly prepared aqueous extract of *Moringa oleifera* leaves in normal and diabetic rabbits. *Journal of Chemical and Pharmaceutical Research*, 4(1): 249-253.
- Mathur ML, Gaur J, Sharma R, Haldiya KR. (2011): Antidiabetic properties of a spice plant *Nigella sativa*. *Journal of Endocrinology and Metabolism*, 1(1): 1-8.
- Medjdoub H, Tabti B, Baatouche M., et al. (2012): Antihyperglycemic effect of *Zygophyllum geslini* aqueous extract in streptozotocin-induced diabetic wistar rats. *Journal of Life Sciences*, 6: 652-656.
- Meenakshi P, Bhuvaneshwari r, Rath MA, Thirumoorthi L, Guravaiah DC, Jiji MJ, Gopalakrishnan VK. (2010): Antidiabetic activity of ethanolic extract of *Zaleya decandra* in alloxan-induced diabetic rats. *Appl Biochem Biotechnol*. 162(4): 1153-1159.
- Meonah STS, Palaniswamy M, et al. (2012): Pharmacognostical and hypoglycemic activity of *Solanum nigrum* Linn. *Plant. International Journal of Pharmacy and Pharmaceutical Sciences*, 4(1): 221-224.
- Mirsky LA., Diengott D., Perisutti G. (1956): The hypoglycemic and in-sulinase-inhibitory action of some plant growth regulators. *Endocrinology* 59: 715-718.
- Mojekwu TO, Yama OE, Ojokuku SA, Oyebadejo SA. (2011): Hypoglycemic effects of aqueous extract of *Aframomum melegueta* leaf on alloxan-induced diabetic male albino rats. *Pac. J. Med. Sci*. 8(1): 28-36.
- Momo CEN, Oben JE, Tazoo D, Dongo E. (2006): Antidiabetic and hypolipidemic effects of *Laportea ovalifolia* (Urticaceae) in alloxan induced diabetic rats. *Afr J Trad. Complement. Altern. Med*. 3(1): 36-43.
- Momoh S, Yusuf OW, Adamu MM, Agwu COC and Atanu FO. (2011): Evaluation of the phytochemical composition and hypoglycemic activity of methanolic leaves extract of *Costus afer* in albino rats. *British J Pharm Res*. 1(1): 1-8.
- Mondal S, Bhattacharya and Biswas M. (2012): Antidiabetic activity of *Areca catechu* leaf extracts against streptozotocin induced diabetic rats. *Journal of Advanced Pharmacy Education and Research* 2(1): 10-17.
- Morada NJ, Metillo EB, Uy MM, and Oclarit M. (2011): Anti-diabetic polysaccharide from mangrove plant, *Sonneratia alba* Sm. *International Conference on Asia Agriculture and Animal*, 13: 197-200.
- Mungle AN, Bodhankar MM, and Chandak KK. (2012): Antidiabetic potential of *Dolichandrone falcata* leaves in alloxan induced diabetic rats. *International Journal of Research in Pharmaceutical and Biomedical Sciences*, 3(1): 319-324.

- Murugesan T, Rao B, Sinha S, Biswas S, Pal M. and Saha BP. (2000): Anti-diabetic activity of *Jussiaea suffruticosa* extract in rats. *Pharmacy and Pharmacology Communications*, 6(10): 451-453.
- Murugesan K, Yeligar V, Dash DK, Sengupta P, Maiti BC, Maity TK. (2006): Antidiabetic, antioxidant and antihyperlipidemic status of *Heliotropium zeylanicum* extract on streptozotocin-induced diabetes in rats. *Bio Pharm Bull.* 29(11): 2202-2205.
- Nair SA, Shylesh BS, Gopakumar B, Subramoniam A. (2006): Antidiabetes and hypoglycemic properties of *Hemionitis arifolia* (Burm.) Moore in rats. *J Ethnopharmacol.* 106: 192-197.
- Ndiaye M, Diatta W, Sy AN, Dieye AM, Faye B, Bassene E. (2008): Antidiabetic properties of aqueous barks extract of *Parinari excelsa* in alloxan-induced diabetic rats. *Fitoterapia*, 79(4): 267-270.
- Nirmala A, Saroja S, Vasanthi HR, Lalitha G. (2009): Hypoglycemic effect of *Basella rubra* in streptozotocin-induced diabetic albino rats. *Journal of Pharmacognosy and Phytotherapy*, 1(2): 025-030.
- Njike GN, Watcho P, Nguetefack TB, Kamanyi A. (2005): Hypoglycemic activity of the leaves extracts of *Bersama engleriana* in rats. *Afr J Trad.Complement. Altern. Med.* 2(3): 215-221.
- Ojewole JA. (2005). Antinociceptive, anti-inflammatory and antidiabetic effects of *Bryophyllum pinnatum* (Crassulaceae) leaf aqueous extract. *J Ethnopharmacol.* 99(1): 13-19.
- Ojewole JA, Adewunmi CO. (2003): Hypoglycemic effect of methanolic extract of *Musa paradisiaca* (Musaceae) green fruits in normal and diabetic mice. *MethodsFind Exp Clin Pharmacol.* 25(6): 453-456.
- Okokon JE, Bassey SA and John AU. (2012): Antidiabetic activities of ethanolic extract and fraction of *Anthocleista djalensis*. *Asian Pac J Trop Biomed.* 2(6): 461-464.
- Oliver-Bever B., Zahnd G.R. (1979): Plants with oral hypoglycaemic action. *Quart. J. Crude Drug Res.* 17: 139-196.
- Orhan N, Aslan M, Hosbas S, Deliorman OD. (2009): Antidiabetic effect and antioxidant potential of *Rosa canina* fruits. *Pharmacognosy Magazine*, 5(20): 309-315.
- Oyedemi S, Bradley G, Afolayan A. (2012): Antidiabetic activities of aqueous stem bark extract of *Strychnos henningsii* Gilg. in streptozotocin-nicotinamide type 2 diabetic rats. *Iran J Pharm Res.* 11(1): 221-228.
- Ozolua RI, Eriyamremu GE, Okene EO and Ochei U. (2006): Hypoglycemic effects of viscous preparation of *Irvingia gabonensis* (Dikanut) seeds in streptozotocin induced diabetic wistar rats. *Journal of Herbs, Spicesand Medicinal Plants*, 12(4): 1-9.
- Patel D, Kumar R, Prasad S, Sairam K, Hemalatha S. (2011): Antidiabetic and in vitro antioxidant potential of *Hybanthus enneaspermus* (Linn) F. Muell in streptozotocin-induced diabetic rats. *Asian PacJ Trop Biomed.* 1(4): 316-322.
- Patel P, Harde P, Pillai J, Darji N and Patel B. (2012): Antidiabetic herbal drugs a review. *Pharmacophore*, 3(1): 18-29.
- Pepato MT, Oliveira JR, Kettelhut IC, Migliorini RH. (1993): Assessment of the antidiabetic activity of *Myrcia uniflora* extracts in streptozotocin diabetic rats. *Diabetes Res.* 22(2): 49-57.
- Peunqvicha P, Thirawarapan SS, Watanabe H. (1996): Hypoglycemic effect of water extract of the root of *Pandanus odoratus* RIDL. *Biol Pharm Bull.* 19(3): 364-366.

- Peunqvicha P, Thirawarapan SS, Temsiririrkkul R, Watanabe H, Kumar PJ, Kadota S. (1998): Hypoglycemic effect of the water extract of *Piper sarmentosum* in rats. J Ethnopharmacol. 60(1): 27-32.
- Pooja, Sharma V and Samanta KC. (2011): Hypoglycemic activity of methanolic extract of *Tectona grandis* Linn. root in alloxan induced diabetic rats. Journal of Applied Pharmaceutical Science, 1(4): 106-109.
- Prisilla DH, Balamurugan R, Shah HR. (2012): Antidiabetic activity of methanol extract of *Acorus calamus* in STZ induced diabetic rats. Asian Pac J Trop Biomed. 2(2): S941-S946.
- Prohp TA, Onoagbe IO. (2011): Anti-diabetic studies of aqueous extract of *Triplochiton scleroxylon* on platelets and associated parameters in alloxan-induced diabetic rabbits. African Journal of Plant Science 2011; 5(12): 697-701.
- Pushparaj P, Tan CH, Tan BK. (2000): Effects of *Averrhoa bilimbi* leaf extract on blood glucose and lipids in streptozotocin-diabetic rats. J Ethnopharmacol. 72: 69-76.
- Raghu S et al. (2011): Antidiabetic activity of *Strychnus potatorum* Linn. seeds against alloxan induced diabetes mellitus in rats. Journal of Pharmaceutical Research and Clinical Practice, 1(1): 18-26.
- Rajagopal K, Sasikala K. (2008): Antihyperglycemic and antihyperlipidemic effects of *Nymphaea stellata* in alloxan-induced diabetic rats. Singapore Med J. 49(2): 137-141.
- Rajeswari J, Kesavan K, Jayakar B. (2012): Antidiabetic activity of chemical characterization of aqueous/ethanol prop roots extracts of *Pandanus fascicularis* Lam in streptozotocin-induced diabetic rats. Asian Pac J Trop Biomed. 2: S170-S174.
- Ramakrishna D, Vidyasagar G, Kumar KP., et al. (2011): Evaluation of antidiabetic activity of *Triumfetta pilosa* Roth in streptozotocin-induced diabetic rats. International Journal of Pharma Sciences and Research, 2(3): 98-103.
- Rao NK, Nammi S. (2006): Antidiabetic and renoprotective effects of the chloroform extract of *Terminalia chebula* Retz. seeds in streptozotocin-induced diabetic rats. BMC Complementry and Alternative Medicine, 6(17): 1-6.
- Rao TN, Kumarappan CT, Lakshmi SM, Mandal SC. (2007): Antidiabetic activity of leaves of *Talinum portulacifolium* (Forssk) in alloxan-induced diabetic rats. Pharmacologyonline, 2: 407-417.
- Rehman SU, Jafri SA, Hassan S, Ahmed I and Naim M. (2011): Study on Antidiabetic effect of *Aloe vera* extract on alloxan induced diabetic rats. Libyan Agriculture Research Centre Journal International, 2(1): 29-32.
- Sabu MC, Kuttan R. (2009): Antidiabetic and antioxidant activity of *Terminalia belerica* Roxb. Indian J Exp Biol. 47(4): 270-275.
- Sah SP et al. (2010): Hypoglycemic activity of aqueous extract of *Urtica parviflora* Roxb. in normoglycaemic rats. International Journal of Phytomedicine, 2: 47-51.
- Saidu AN, Mann A and Onuegbu CD. (2012): Phytochemical screening and hypoglycemic effect of aqueous *Blighia sapida* root bark extract on normoglycaemic albino rats. British Journal of Pharmaceutical Research 2012; 2(2): 89-97.
- Shajeela PS, Kalpana DV, V.R. Mohan. (2012): Potential antidiabetic, hypolipidemic and antioxidant effects of *Nymphaea pubescens* extract in alloxan induced diabetic rats. Journal of Applied Pharmaceutical Science, 2(2): 83-88.
- Sharma S et al. (2010): Antidiabetic screening leaves extract of *Holoptelea integrifolia* (Roxb.). International Journal of Pharma Research and Development 2010; 2(10): 66-71.

- Sharma V, Pooja, Marwaha A. (2011): Hypoglycemic activity of methanolic extracts of *Nyctanthes arbor-tristis* Linn. root in alloxan induced diabetic rats. International Journal of Pharmacy and Pharmaceutical Sciences, 3(3): 210-212.
- Sher H, Alyemeni MN. (2011): Evaluation of anti-diabetic activity and toxic potential of *Lycium shawii* in animal models. J Med Plant Res. 5(15): 3387-3395.
- Sidhu MC, Sharma T. (2013a): A database of antidiabetic plant species of family asteraceae, euphorbiaceae, fabaceae, lamiaceae and moraceae. International Journal of Herbal Medicine, 1(2): 152-164.
- Sidhu MC, Sharma T. (2013b): Medicinal plants from twelve families having antidiabetic activity: a review. American Journal of PharmTech Research, 3(5): 37-52.
- Sikarwar MS, Patil MB. (2010): Antidiabetic activity of *Crataeva nurvala* stem bark extracts in alloxan induced diabetic rats. Journal of Pharmacy and Bioallied Sciences, 2(1): 18-21.
- Sindhu T et al. (2010): Antidiabetic activity of *Mollugo nudicaulis* against alloxan-induced diabetic rats. International Journal of Applied Biology and Pharmaceutical Technology, 1: 511-519.
- Singh AK, Singh J. (2010): Evaluation of antidiabetic potential of leaves and stem of *Flacourtia jangomas* in streptozotocin-induced diabetic rats. Indian J Pharm. 42(5): 301-305.
- Sujatha S, Swaroopa RV and Ravikumar B. (2012): Antidiabetic effect of flower extract of *Antigonon leptopus* Hook and Arn in alloxan-induced diabetic rats. Indian Journal of Pharmaceutical Education and Research, 46(1): 9-16.
- Sunder AS, Rajyalakshmi G, Bharath A and Rajeshwar Y. (2009): Antihyperglycemic activity of *Trianthema portulacastrum* plant in streptozotocin induced diabetic rats. Pharmacologyonline, 1: 1006-1011.
- Sunil C, Ignacimuthu S, Agastian P. (2011): Antidiabetic effect of *Symplocos cochinchinensis* (Lour.) S Moore in type 2 diabetic rats. J Ethnopharmacol. 134(2): 298-304.
- Sunil C et al. (2009): Effect of ethanolic extracts of *Pisonia alba* Span leaves on blood glucose levels and histological changes in tissues of alloxan induced diabetic rats. International Journal of Applied Research in Natural Products, 2(2): 4-11.
- Sutar NG, Sutar UN, Behera BC. (2009): Antidiabetic activity of the leaves of *Mimosa pudica* Linn in albino rats. Journal of Herbal Medicine and Toxicology, 3(1): 123-126.
- Syiem D, Syngai G, Khup PZ, Khongwir BS, Kharbuli B, Kayang H. (2002): Hypoglycemic effects of *Potentilla fulgens* L in normal and alloxan induced diabetic mice. J Ethnopharmacol. 83(1-2): 55-61.
- Tanko Y, Yerima m, Mahdi MA, yaro AH, Musa KY, Mohammed A. (2008): Hypoglycemic activity of methanolic stem bark of *Adansonia digitata* extract on blood glucose level of streptozotocin-induced diabetic wistar rats. International Journal of Applied Research in Natural Products, 1(2): 32-36.
- Tanquilut NC, Tanquilut MRC, Estacio MAC, Torres EB et al. (2009): Hypoglycemic effects of *Lagerstroemia speciosa* (L.) Pers on alloxan induced diabetic mice. Journal of Medicinal Plants Research, 3(12): 1066-1071.
- Teimori M, Montasser KS, Ghafarzadegan R, Hajiaghaee R. (2010): Study of hypoglycemic effect of *Juglans regia* leaves and its mechanism. Journal of Medicinal Plants, 9(6): 57-65.
- Thirumalai T, Therasa SV, Elumalai E, David E. (2011): Hypoglycemic effect of *Brassica juncea* (seeds) on streptozotocin induced diabetic male albino rat. Asian Pac J Trop Biomed. 1(4): 323-325.

Thiruvengkatasubramaniam R, Jayakar B. (2010): Anti-hyperglycemic and anti-hyperlipidemic activities of *Premna corymbosa* (Burm. F.) Rottl on streptozotocin induced diabetic rats. *Der Pharmacia Lettre*, 2(1): 505-509.

Tierney LM. (2002): *Current Medical Diagnosis and Treatment*, 2002. New York: Lange Medical Books/McGraw-Hill.

Udayakumar R et al. (2009): Hypoglycemic and hypolipidemic effects of *Withania somnifera* root and leaf extracts on alloxan-induced diabetic rats. *Int J Mol Sci*. 10(5): 2367-2382.

Upwar N, Patel R, Waseem N. (2010): Hypoglycemic effect of methanolic extract of *Berberis aristata* DC stem on normal and streptozotocin induced diabetic rats. *International Journal of Pharmacy and Pharmaceutical Sciences*, 3(1): 222-224.

Veeramani C, Pushpavalli G, Pagalendi KV. (2008): Antihyperglycemic effect of *Cardiospermum halicacabum* Linn leaf extract on STZ-induced diabetic rats. *J. Appl. Biomed*. 6: 19-26.

Venkateshwarlu E, Reddy ARN, Sunder AS, Kiran G, Rao JV and Madhusudhan S. (2009): Anti-hyperglycemic activity of methanolic extract of *Salacia fruticosa* leaves in alloxan induced diabetic rats. *Drug Invention Today*, 1(2): 95-97.

Viana GS, Medeiros AC, Lacerda AM, Leal LK, Vale TG, Matos FJ. (2004): Hypoglycemic and anti-lipemic effects of the aqueous extract from *Cissus sicyoides*. *BMC Pharmacology* 2004; 4(9): 1-7.

Wang JC, hu SH, Wang JT, Chen KS and Chia YC. (2005): Hypoglycemic effect of extract of *Hericium erinaceus*. *J Sci Food Agric*. 85: 641-646.

WHO/Acadia, (1992): *Rapport de la Journe Internationale de, diabetes* 14 octobre.

Yadav JP, Saini S, Kalia AN, Dangi AS. (2008): Hypoglycemic and hypolipidemic activity of ethanolic extract of *Salvadora oleoides* in normal and alloxan-induced diabetic rats. *Indian J Pharmacol*. 40(1): 23-27.

Yoshino K, Miyauchi Y, Kanetaka T, Takagi Y, Koga K. (2009): Anti-diabetic activity of a leaf extract prepared from *Salacia reticulata* in mice. *Biosci Biotechnol Biochem*. 73(5): 1096-1104.

Zaahkouk SAM, Rashid SZA and Mattar AF. (2003): Anti-diabetic properties of water and ethanolic extracts of *Balanites aegyptiaca* fruits flesh in senile diabetic rats. *The Egyptian Journal of Hospital Medicine*, 10: 90-108.

Zafar I, Mohammed P, Asadullah et al. (2002): Study of the hypoglycemic activity of *Hedera helix* L in alloxan-induced diabetic rabbits. *J. Med. Sci*. 2(4): 206-208.

Zahid H, Rizwani GH, Shareef H, Mahmud S and Ali T. (2012): Hypoglycaemic and hypolipidaemic effects of *Mimusops elengi* Linn extracts on normoglycaemic and alloxan-induced diabetic rats. *International Journal of Pharmaceutical and Biological Archives*, 3(1): 56-62.

Zhang W, Zhao J, Wang J, Pang X, Zhuang X, Zhu X, Qu W. (2010): Hypoglycemic effect of aqueous extract of seabuckthorn (*Hippophae rhamnoides* L.) seed residues in streptozotocin-induced diabetic rats. *Phytother Res*. 24(2): 228-232.

Zhou T, Luo D, Li X, Luo y. (2009): Hypoglycemic and hypolipidemic effects of flavonoids from Lotus (*Nelumbo nucifera* Gaertn) leaf in diabetic mice. *Journal of Medicinal Plants Research*, 3(4): 290-293.

Zulfiker AHM, Ripa FA, Rahman MM, Ullah MO, Hamid K, Khan MMR, Rana MS. (2010): Antidiabetic and antioxidant effect of *Scoparia dulcis* in alloxan induced albino mice. *International Journal of PharmTech Research*, 2(4): 2527-2534.