

[Open in app](#)

Hritika Agarwal

[Follow](#)

5 Followers

[About](#)

Cost function Intuition 2




Hritika Agarwal Jul 18 · 2 min read

Hypothesis: $h_{\theta}(x) = \theta_0 + \theta_1 x$


Parameters: θ_0, θ_1


Cost Function: $J(\theta_0, \theta_1) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$

Goal: minimize $J(\theta_0, \theta_1)$
 θ_0, θ_1




$J(\theta_0, \theta_1)$
(function of the parameters θ_0, θ_1)







A contour plot is a graph that contains many contour lines. A contour line of a two-variable function has a constant value at all points of the same line. An example of such a graph is the one to the right below.




Taking any color and going along the 'circle', one would expect to get the same value of the cost function. For example, the three green points found on the green line above have the same value for $J(\theta_0, \theta_1)$ and as a result, they are found along the same line. The circled x displays the value of the cost function for the graph on the left when $\theta_0 = 800$ and $\theta_1 = -0.15$. Taking another $h(x)$ and plotting its contour plot, one gets the following graphs:







When $\theta_0 = 360$ and $\theta_1 = 0$, the value of $J(\theta_0, \theta_1)$ in the contour plot gets closer to the center thus reducing the cost function error. Now giving our hypothesis function a slightly positive slope results in a better fit of the data.



The graph above minimizes the cost function as much as possible and consequently, the result of θ_1 and θ_0 tend to be around 0.12 and 250 respectively. Plotting those values on our graph to the right seems to put our point in the center of the innermost 'circle'.

3 dimensionally contour plot looks like this





Read Next — [Gradient Descent](#)

Machine Learning Cost Function Linear Regression

[About](#) [Help](#) [Legal](#)

Get the Medium app

