

Lindblad Equation

Jie Ren

Contents

I. Lindblad Master Equation	1
A. General Markovian Form	1
B. First Principal Deduction	2
C. Stochastic Schrödinger Equation	4
1. Poisson SSE	4
2. Gaussian SSE	4
II. Quadratic Lindbladian	5
A. Third Quantization	5
B. Quadratic Jump Operators	6
1. Majorana Case	6
2. Dirac Fermion Case	6
III. Fermionic Gaussian States	7
A. Grassmann Representation	7
1. Gaussian States	8
2. Gaussian Operators	8
3. Gaussian Linear Maps	10
B. Operator Form	10
1. Dirac Fermion Case	10
2. Majorana Case	11
References	11

I. LINDBLAD MASTER EQUATION

A. General Markovian Form

For general open quantum evolution, suppose the system and environment are separable initially: $\rho_T = \rho \otimes \rho_B$, where we assume $\rho_B = \sum_{\alpha} \lambda_{\alpha} |\phi_{\alpha}\rangle\langle\phi_{\alpha}|$. Then the evolution of system-bath is unitary: $\rho_T(t) = U(t)\rho_T U^{\dagger}(t)$. Trace out the environment's degrees of freedom, we have the quantum channel expression:

$$\rho(t) = \sum_{\alpha\beta} W_{\alpha\beta} \rho W_{\alpha\beta}^{\dagger}, \quad W_{\alpha\beta} = \sqrt{\lambda_{\beta}} \langle\phi_{\alpha}|U(t)|\phi_{\beta}\rangle. \quad (1)$$

In general, the evolution of an open quantum system has the form $\rho(t) = \mathcal{L}_t[\rho]$. The Lindblad equation assumes a semi-group relation: $\mathcal{L}_t = \lim_{N \rightarrow \infty} \mathcal{L}_{t/N} \cdot \mathcal{L}_{t/N} \cdots \mathcal{L}_{t/N}$. Such time decimation implies that the evolution is Markovian. We will show that Markovian approximation leads directly to the Lindblad equation. First, we choose a complete operator basis $\{F_i\}$ in N -dimensional Hilbert space, satisfying $\text{Tr}[F_i^{\dagger} F_j] = \delta_{ij}$, where we choose $F_0 = N^{-1/2} \cdot \mathbb{I}$. For a quantum channel, the channel operator K_{μ} can be expanded as $K_{\mu} = \sum_i \text{Tr}[F_i^{\dagger} K_{\mu}] F_i$. In general, we have:

$$\mathcal{L}_t[\rho] = \sum_{ij} c_{ij}(t) F_i \rho F_j^{\dagger},$$

where the Hermitian coefficient $c_{ij}(t)$ is $c_{ij}(t) = \sum_{\mu} \text{Tr}[F_i^{\dagger} K_{\mu}] \cdot \text{Tr}[F_j^{\dagger} K_{\mu}]^*$. Our target is to compute the limit

$$\frac{d}{dt}\rho \equiv \lim_{t \rightarrow 0} \frac{1}{t}(\mathcal{L}_t[\rho] - \rho).$$

For this purpose, we define the (Hermitian) coefficient a_{ij} as:

$$a_{00} = \lim_{t \rightarrow 0} \frac{c_{00}(t) - N}{t}, \quad a_{ij} = \lim_{t \rightarrow 0} \frac{c_{ij}(t)}{t}.$$

The limit is then

$$\frac{d}{dt}\rho = \frac{a_{00}}{N}\rho + \frac{1}{\sqrt{N}} \sum_{i>0} \left(a_{i0} F_i \rho + a_{i0}^* \rho F_i^{\dagger} \right) + \sum_{i,j>0} a_{ij} F_i \rho F_j^{\dagger}.$$

To further simplify the expression, we define

$$F = \frac{1}{\sqrt{N}} \sum_{i=1}^{N^2-1} a_{i0} F_i, \quad G = \frac{1}{2N} a_{00} \mathbb{I} + \frac{1}{2} (F^{\dagger} + F), \quad H = \frac{1}{2i} (F^{\dagger} - F).$$

The limit can be expressed by G, H in a compact form:

$$\frac{d\rho}{dt} = -i[H, \rho] + \{G, \rho\} + \sum_{i,j=1}^{N^2-1} a_{ij} F_i \rho F_j^{\dagger}. \quad (2)$$

Note the $[H, \rho]$ part is the traceless part and the $\{G, \rho\}$ is the trace part. Since the quantum channel preserves the trace (for any ρ):

$$\text{Tr} \left[\frac{d\rho}{dt} \right] = \text{Tr} \left[\left(2G + \sum_{i,j=1}^{N^2-1} a_{ij} F_j^{\dagger} F_i \right) \rho \right] = 0.$$

Therefore $G = -\frac{1}{2} \sum_{i,j=1}^{N^2-1} a_{ij} F_j^{\dagger} F_i$. We thus obtain the Lindblad form:

$$\frac{d\rho}{dt} = -i[H, \rho] + \sum_{i,j=1}^{N^2-1} a_{ij} \left(F_i \rho F_j^{\dagger} - \frac{1}{2} \{F_j^{\dagger} F_i, \rho\} \right).$$

We can further simplify the form by diagonalizing the matrix a_{ij} . It is a convention to take the norm of a_{ij} out to indicate the strength of the system-bath coupling, and the diagonalized Lindblad equation is

$$\frac{d\rho}{dt} = -i[H, \rho] + \gamma \sum_m \left(L_m \rho L_m^{\dagger} - \frac{1}{2} \{L_m^{\dagger} L_m, \rho\} \right). \quad (3)$$

B. First Principal Deduction

In this section, we consider a general system-bath coupling:¹

$$H_T = H + H_B + V, \quad V = \sum_k A_k \otimes B_k. \quad (4)$$

We will show under certain condition, the dynamics of the system is well approximated by the Lindblad equation. We first assume that initially, the total system is a product state

$$\rho_T(0) = \rho(0) \otimes \rho_B.$$

¹ Without loss of generality, we can also assume $\|A_k\| = 1$, $\text{Tr}[\rho_B B_k] = 0$.

In the following, we will adopt the interacting picture, where the density operator evolves as

$$\partial_t \rho_T(t) = -i[V(t), \rho_T(t)] \equiv -i\mathcal{V}(t)|\rho_T(t)\rangle.$$

Note that in the last equality, ρ_T is expressed as a ket in the Hilbert space of linear operator, and the commutator with V is expressed as a superoperator \mathcal{V} . This notation can simplify the expression. For example, the inner product in the operator space is the trace, so the partial trace operation can be denoted as $|\rho\rangle = \langle \mathbb{I}_B | \rho_T \rangle$. The evolution of the system is then

$$\begin{aligned} \frac{d}{dt}|\rho(t)\rangle &= -i\langle \mathbb{I}_B | \mathcal{V}(t) | \rho_T(t) \rangle = -i\langle \mathbb{I}_B | \mathcal{V}(t) | \rho_T(0) \rangle - \int_0^t \langle \mathbb{I}_B | \mathcal{V}(t) \mathcal{V}(\tau) | \rho_T(\tau) \rangle d\tau \\ &= - \int_0^t \langle \mathbb{I}_B | \mathcal{V}(t) \mathcal{V}(\tau) | \rho_T(\tau) \rangle d\tau. \end{aligned}$$

Now we are taking the **Born approximation**, which states when the coupling is weak enough compared with the energy scale of the system and the bath, the total density matrix is approximated by the product state $|\rho_T(t)\rangle \approx |\rho(t)\rangle \otimes |\rho_B\rangle$. The evolution is now

$$\begin{aligned} \frac{d}{dt}\rho(t) &\approx \int_0^t \text{Tr}_B [V(t)\rho_T(\tau)V(\tau) - \rho_T(\tau)V(\tau)V(t)] d\tau + h.c. \\ &= \sum_{kl} \int_0^t d\tau C_{lk}(\tau - t) [A_k(t)\rho(\tau)A_l(\tau) - \rho(\tau)A_l(\tau)A_k(t)] + h.c., \end{aligned}$$

where $C_{kl}(t) \equiv \text{Tr}_B[\rho_B B_k(t)B_l]$ is the correlation function of B_k 's. We then take the **Markovian approximation** which assumes that the correlations of the bath decay fast in time. We can thus make the substitution $\rho(\tau) \rightarrow \rho(t)$, the result equation of motion is Markovian:

$$\begin{aligned} \frac{d}{dt}\rho(t) &\approx \sum_{kl} \int_0^t dt' C_{lk}(-t') [A_k(t)\rho(t)A_l(t-t') - \rho(t)A_l(t-t')A_k(t)] + h.c. \\ &= \sum_k \int_0^t dt [A_k \rho B_k - \rho B_k A_k + h.c.], \end{aligned}$$

where we have defined $B_k(t) = \sum_l \int_0^\infty dt' A_l(t-t')C_{lk}(-t')$. Now we switch to the frequency domain,

$$A_k(t) = \sum_\omega A_k(\omega)e^{-i\omega t}, \quad B_k(t) = \sum_{l,\omega} e^{-i\omega t} A_l(\omega)\Gamma_{lk}(\omega), \quad \Gamma_{kl}(\omega) = \int_0^\infty dt e^{i\omega t} C_{kl}(t).$$

We then take the **rotating wave approximation**, where we only keep the contributions from canceling frequency of operator A and B ,

$$\begin{aligned} \frac{d}{dt}\rho(t) &= \sum_\omega [\Gamma_{lk}(\omega)A_k(\omega)\rho A_l(\omega) - \Gamma_{lk}(\omega)\rho A_l(\omega)A_k(\omega) + h.c.] \\ &= \sum_\omega \gamma_{kl}(\omega)(A_{l,\omega}\rho A_{k,\omega}^\dagger - \frac{1}{2}\{\rho, A_{k,\omega}^\dagger A_{l,\omega}\}) - i \left[\sum_\omega S_{kl}(\omega)A_{k,\omega}^\dagger A_{l,\omega}, \rho \right], \end{aligned} \tag{5}$$

where we defined

$$\gamma_{kl}(\omega) = \Gamma_{kl}(\omega) + \Gamma_{lk}^*(\omega), \quad S_{kl}(\omega) = \frac{\Gamma_{kl}(\omega) - \Gamma_{lk}^*(\omega)}{2i}.$$

The matrices $\gamma(\omega)$ are positive, we can then take the square root of them. The jump operator is then

$$L_{i,\omega} = \sum_j \sqrt{\gamma_{ij}(\omega)} A_{j,\omega}.$$

The evolution is then in the Lindblad form.

C. Stochastic Schrödinger Equation

The Lindblad form Eq. (3) is equivalent to the stochastic Schrödinger equation (SSE):

$$d|\psi\rangle = -iH|\psi\rangle + A[\psi]dt + B[\psi]dW, \quad (6)$$

where dW is a stochastic infinitesimal element. The expectation value is then the average over all possible evolution path (trajectory): $\langle O(t) \rangle = \overline{\langle \psi(t) | O | \psi(t) \rangle}$.

1. Poisson SSE

Consider a small time interval Δt , the Lindblad equation is equivalent to the quantum channel $\rho(t + \Delta t) = M_0 \rho(t) M_0^\dagger + \sum_m M_m \rho(t) M_m^\dagger$, where

$$M_0 = 1 - i \left(H - i \frac{\gamma}{2} \sum_m L_m^\dagger L_m \right) \Delta t, \quad M_m = \sqrt{\gamma \Delta t} L_m.$$

A quantum channel can be simulated by a stochastic evolution of pure states:

$$|\psi(t + \Delta t)\rangle \propto \begin{cases} L_m |\psi(t)\rangle & p = p_m(t) \gamma \Delta t \\ \exp(-iH_{\text{eff}} \Delta t) |\psi(t)\rangle & p = 1 - \sum_m p_m(t) \end{cases}, \quad \text{where } p_m(t) = \langle \psi(t) | L_m^\dagger L_m | \psi(t) \rangle. \quad (7)$$

Here the effective (non-Hermitian) Hamiltonian is

$$H_{\text{eff}} = H - i \frac{\gamma}{2} \sum_m L_m^\dagger L_m. \quad (8)$$

We can introduce a Poisson variable dW_m satisfying

$$dW_m dW_n = \delta_{mn} dW_m, \quad \overline{dW_m} = \langle L_m^\dagger L_m \rangle \gamma dt,$$

and the evolution can be cast into the stochastic differential equation

$$d|\psi\rangle = -iHdt|\psi\rangle + \sum_m \left[\left(\frac{L_m}{\langle L_m^\dagger L_m \rangle^{\frac{1}{2}}} - 1 \right) dW_m - \frac{\gamma}{2} (L_m^\dagger L_m - \langle L_m^\dagger L_m \rangle) dt \right] |\psi\rangle. \quad (9)$$

Note that the $-\langle L_m^\dagger L_m \rangle dt |\psi\rangle$ comes from the renormalization. For numerical simulation, we can ignore it.

2. Gaussian SSE

We can also use the Wiener processes dW_m satisfying

$$\overline{dW_m} = 0, \quad \overline{dW_m dW_n} = \delta_{mn} \gamma dt.$$

The Gaussian SSE is

$$d|\psi\rangle = -iHdt|\psi\rangle + \sum_m \left[(L_m - \langle L_m \rangle) dW_m - \frac{\gamma}{2} (L_m^\dagger - \langle L_m^\dagger \rangle) (L_m - \langle L_m \rangle) dt \right] |\psi\rangle. \quad (10)$$

To retain the Lindblad, note that $d\rho = \overline{|d\psi\rangle\langle\psi|} + \overline{|\psi\rangle\langle d\psi|} + \overline{|d\psi\rangle\langle d\psi|}$. Without going into the detail, we note that $L_m dW_m$ term in $\overline{|d\psi\rangle\langle d\psi|}$ will contribute a term $\gamma L_m \rho L_m^\dagger dt$; $-\frac{\gamma}{2} L_m^\dagger L_m dt$ term in $\overline{|d\psi\rangle\langle\psi|} + \overline{|\psi\rangle\langle d\psi|}$ contribute a term $-\frac{\gamma}{2} \{L_m^\dagger L_m, \rho\}$ term. All terms involving expectation value can be regarded as coming from the renormalization.

II. QUADRATIC LINDBLADIAN

Consider the Lindblad in the Heisenberg picture:

$$\frac{d}{dt}\hat{O} = i[\hat{H}, \hat{O}] + \sum_{\mu} \hat{L}_{\mu}^{\dagger} \hat{O} \hat{L}_{\mu} - \frac{1}{2} \sum_{\mu} \{\hat{L}_{\mu}^{\dagger} \hat{L}_{\mu}, \hat{O}\}, \quad (11)$$

where we choose $\hat{O}_{ij} = \omega_i \omega_j$ satisfying the relation $\hat{O}^T = 2\mathbb{I} - \hat{O}$. The covariance matrix is then $\Gamma_{ij} = i\langle \hat{O} \rangle - i\delta_{ij}$.

We assume that the jump operator has up to quadratic Majorana terms. In particular, we denote the linear terms and the Hermitian quadratic terms as

$$\hat{L}_r = \sum_{j=1}^{2N} L_j^r \omega_j, \quad \hat{L}_s = \sum_{j,k=1}^{2N} M_{jk}^s \omega_j \omega_k.$$

When the **jump operator** \hat{L}_{μ} contains only the linear Majorana operator, the Lindblad equation preserves Gaussianity. For jump operators containing up to quadratic Majorana terms, the evolution will break the Gaussian form, however, the $2n$ -point correlation is still solvable for free fermion systems.

A. Third Quantization

Assume only linear terms in jump operators,

$$\partial_t \hat{O} = [i\hat{H}, \hat{O}] + \mathcal{D}_r[\hat{O}] = \left[i\hat{H} - \frac{1}{2} \sum_r \hat{L}_r^{\dagger} L_r, \hat{O} \right] + \sum_r [\hat{L}_r^{\dagger}, \hat{O}] \hat{L}_r.$$

Define $B \equiv \sum_r L_i^r L_j^{r*}$, the first term of EOM is:²

$$\begin{aligned} \left[i\hat{H} - \frac{1}{2} \sum_r \hat{L}_r^{\dagger} L_r, \hat{O}_{ij} \right] &= \sum_{kl} \left(\frac{1}{4} H - \frac{1}{2} B \right)_{kl} [\omega_k \omega_l, \omega_i \omega_j] \\ &= \sum_{kl} \left(\frac{1}{2} H - B \right)_{kl} (\delta_{ki} \omega_j \omega_l - \delta_{kj} \omega_i \omega_l + \delta_{li} \omega_k \omega_j - \delta_{lj} \omega_k \omega_i) \\ &= \left[\left(\frac{1}{2} H - B \right) \cdot \hat{O}^T + \left(\frac{1}{2} H - B \right)^T \cdot \hat{O} - \hat{O} \cdot \left(\frac{1}{2} H - B \right)^T - \hat{O}^T \cdot \left(\frac{1}{2} H - B \right) \right]_{ij} \\ &= \left[(-H + 2B^T) \cdot \hat{O} + \hat{O} \cdot (H - 2B^T) \right]_{ij} \end{aligned}$$

The second term is

$$\begin{aligned} \sum_r [\hat{L}_r^{\dagger}, \hat{O}_{ij}] \hat{L}_r &= \sum_{kl} B_{kl} [\omega_k, \omega_i \omega_j] \omega_l = 2 \sum_{kl} B_{kl} (\delta_{ki} \omega_j \omega_l - \delta_{kj} \omega_i \omega_l) \\ &= \left[2B \cdot \hat{O}^T - 2\hat{O} \cdot B^T \right]_{ij} = \left[-2B \cdot \hat{O} - 2\hat{O} \cdot B^* + 4B \right]_{ij} \end{aligned}$$

Therefore

$$\partial_t \hat{O}_{ij} = \left[(-H - 2B^R) \cdot \hat{O} + \hat{O} \cdot (H - 2B^R) + 4B \right]_{ij}$$

The EOM of the covariance matrix is then

$$\partial_t \Gamma = X^T \cdot \Gamma + \Gamma \cdot X + Y, \quad (12)$$

² Use the commutation relation $\{\omega_i, \omega_j\} = 2\delta_{ij}$, we have the relation $[\omega_k, \omega_i \omega_j] = 2(\delta_{ki} \omega_j - \delta_{kj} \omega_i)$ and $[\omega_k \omega_l, \omega_i \omega_j] = 2(\delta_{ki} \omega_j \omega_l - \delta_{kj} \omega_i \omega_l + \delta_{li} \omega_k \omega_j - \delta_{lj} \omega_k \omega_i)$.

where $X = H - 2B^R$, $Y = 4B^I$. Note that the constant part is replaced by its anti-symmetric part.

The steady state of the system is solved by the Lyapunov equation

$$X^T \cdot \Gamma + \Gamma \cdot X = -Y. \quad (13)$$

B. Quadratic Jump Operators

Now include the Hermitian quadratic quantum jumps:

$$\begin{aligned} \partial_t \hat{O} &= i[\hat{H}, \hat{O}] + \mathcal{D}_r[\hat{O}] + \mathcal{D}_s[\hat{O}], \\ \mathcal{D}_s[\hat{O}] &= \sum_s \hat{L}_s \hat{O} \hat{L}_s - \frac{1}{2} \sum_r \{\hat{L}_s^2, \hat{O}\} = -\frac{1}{2} \sum_s [\hat{L}_s, [\hat{L}_s, \hat{O}]]. \end{aligned} \quad (14)$$

1. Majorana Case

A direct calculation gives

$$\begin{aligned} D_s[\hat{O}] &= -\frac{1}{2} \sum_s \sum_{kl} M_{kl}^s \langle [\hat{L}_s, [\omega_k \omega_l, \omega_i \omega_j]] \rangle \\ &= 2 \sum_s \sum_k \left\{ M_{ik}^s [\hat{L}_s, \omega_k \omega_j] - [\hat{L}_s, \omega_i \omega_k] M_{kj}^s \right\} \\ &= 8 \sum_{s,kl} \left[M_{ik}^s (-M_{kl}^s \omega_l \omega_j + \omega_k \omega_l M_{lj}^s) + (M_{il}^s \omega_l \omega_k - \omega_i \omega_l M_{lk}^s) M_{kj}^s \right] \\ &= 8 \sum_s \left[2M^s \cdot \hat{O} \cdot M^s - (M^s)^2 \cdot \hat{O} - \hat{O} \cdot (M^s)^2 \right]_{ij}. \end{aligned}$$

Together, we get the EOM of the variance matrix Γ_{ij} :

$$\partial_t \Gamma = X^T \cdot \Gamma + \Gamma \cdot X + \sum_s (Z^s)^T \cdot \Gamma \cdot Z^s + Y, \quad (15)$$

where

$$X = H - 2B^R + 8 \sum_s (\text{Im} M^s)^2, \quad Y = 4B^I, \quad Z = 4M^s. \quad (16)$$

2. Dirac Fermion Case

In this section, we consider the free fermion system preserving the U(1) charge. The jump operators are assumed to be quadratic: $\hat{L}_s = \sum_{jk} M_{jk}^s c_j^\dagger c_k$ where $\{M^s\}$ are Hermitian matrices.

For the fermion case, we choose $\hat{O}_{ij} = c_i^\dagger c_j$, and consider the Lindbladian

$$\partial_t \hat{O} = i[\hat{H}, \hat{O}] + \mathcal{D}_s[\hat{O}] = i[\hat{H}, \hat{O}] - \frac{1}{2} \sum_s [\hat{L}_s, [\hat{L}_s, \hat{O}]],$$

where each $\hat{L}_s = M_{ij}^s c_i^\dagger c_j$ is a Hermitian fermion bilinear.

The Hamiltonian part is:³

$$i \sum_{kl} H_{kl} [c_k^\dagger c_l, c_i^\dagger c_j] = i \sum_{kl} H_{kl} (\delta_{il} c_k^\dagger c_j - \delta_{jk} c_i^\dagger c_l) = i[H^T \cdot \hat{O} - \hat{O} \cdot H^T]_{ij}.$$

³ Using the fact $[c_k^\dagger c_l, c_i^\dagger c_j] = c_k^\dagger [c_l, c_i^\dagger c_j] + [c_k^\dagger, c_i^\dagger c_j] c_l = \delta_{il} c_k^\dagger c_j - \delta_{jk} c_i^\dagger c_l$, we know that for a quadratic form $\hat{A} = \sum_{ij} A_{ij} c_i^\dagger c_j$, $[\hat{A}, \hat{O}_{ij}] = [A^T, \hat{O}]_{ij}$.

Similarly, the double commutation in the second term is:

$$\mathcal{D}_s[\hat{O}] = -\frac{1}{2} \sum_s [(M^{s*})^2 \cdot \hat{O} + \hat{O} \cdot (M^{s*})^2 - 2M^{s*} \cdot \hat{O} \cdot M^{s*}].$$

Together, the EOM of correlation $G_{ij} = \langle c_i^\dagger c_j \rangle$ is

$$\partial_t G = X^\dagger \cdot G + G \cdot X + \sum_s M^{s*} \cdot G \cdot M^{s*}, \quad (17)$$

where $X = -iH^* - \frac{1}{2} \sum_s (M^{s*})^2$.

III. FERMIONIC GAUSSIAN STATES

In this section, we discuss the general fermionic Gaussian state, in the framework of the Grassmann representation. We will closely follow Ref. [1].

A. Grassmann Representation

The Majorana operators are defined as $\hat{\omega}_j^a = \hat{c}_i + \hat{c}_i^\dagger$, $\hat{\omega}_j^b = i(\hat{c}_i - \hat{c}_i^\dagger)$. A general operator in Fermionic Fock space can be expanded on the Majorana basis:

$$\hat{X} = \alpha \hat{I} + \sum_{p=1}^{2n} \sum_{1 \leq a_1 < \dots < a_p \leq 2n} \alpha_{a_1 \dots a_p} \hat{\omega}_{a_1} \dots \hat{\omega}_{a_p}. \quad (18)$$

Define a linear map from Fermionic operator space to Grassmann algebra:

$$\hat{X} \mapsto X(\theta) = \alpha + \sum_{1 \leq a_1 < \dots < a_p \leq 2n} \alpha_{a_1 \dots a_p} \theta_{a_1} \dots \theta_{a_p}. \quad (19)$$

This mapping is called the Grassmann representation of \hat{X} .

One can formally define calculus on Grassmann algebra:

$$\frac{\partial}{\partial \theta_i} \theta_j = \int d\theta_i \theta_j = \delta_{ij}, \quad \frac{\partial}{\partial \theta_i} 1 = \int d\theta_i 1 = 0. \quad (20)$$

The Gaussian integral of Grassmann algebra is

$$\int D\theta \exp \left(\eta^T \theta + \frac{i}{2} \theta^T M \theta \right) = i^n \text{Pf}(M) \exp \left(-\frac{i}{2} \eta^T M^{-1} \eta \right). \quad (21)$$

One useful result concerning the expectation value is

Theorem 1. For two operator \hat{X} and \hat{Y} , we have the following identity

$$\text{Tr}(\hat{X}\hat{Y}) = (-2)^n \int D[\theta, \mu] e^{\theta^T \cdot \mu} X(\theta) Y(\mu).$$

where $\int D\theta = \int d\theta_{2n} \dots \int d\theta_1$, $\int D\mu = \int d\mu_{2n} \dots \int d\mu_1$.

Proof. We prove the statement by considering only m -th order monomial. On the one hand

$$\text{LHS} = \text{Tr}[\hat{\omega}_1 \dots \hat{\omega}_m \hat{\omega}_1 \dots \hat{\omega}_m] = 2^n (-1)^{m(m-1)/2}.$$

On the other hand,

$$\begin{aligned} \text{RHS} &= (-2)^n \int D[\theta, \mu] \theta_1 \dots \theta_m (\theta_{m+1} \mu_{m+1} \dots \theta_{2n} \mu_{2n}) \mu_1 \dots \mu_m \\ &= (-2)^n (-1)^{(4n-m)m + (m+1+2n)(2n-m)/2} \\ &= 2^n (-1)^{-m(m+3)/2} = 2^n (-1)^{m(m-1)/2}. \end{aligned}$$

We therefore proved the statement. \square

1. Gaussian States

Definition 1. A quantum state $\hat{\rho}$ is Gaussian if it has Gaussian Grassmann representation:

$$\rho(\theta) = \frac{1}{2^n} \exp\left(\frac{i}{2} \theta^T M \theta\right),$$

where the antisymmetric matrix $M_{ab} = \frac{i}{2} \text{Tr}(\hat{\rho}[\hat{\omega}_a, \hat{\omega}_b])$ is the **covariance matrix**.

All higher correlations of a Gaussian state are determined by the Wick theorem, namely

$$\text{Tr}(i^p \hat{\rho} \hat{\omega}_{a_1} \cdots \hat{\omega}_{a_p}) = \text{Pf}(M|_{a_1, \dots, a_p}).$$

The canonical form of antisymmetric matrix M is:

$$M = R \begin{bmatrix} 0 & \text{diag}(\lambda_1, \dots, \lambda_n) \\ -\text{diag}(\lambda_1, \dots, \lambda_n) & 0 \end{bmatrix} R^T, \quad R \in \text{SO}(2n).$$

Under the new Grassmann variance $\mu = R\theta$, ρ has the form

$$\rho(\mu) = \frac{1}{2^n} \prod_j \exp(i\lambda_j \mu_j \mu_{j+n}) = \frac{1}{2^n} \prod_j (1 + i\lambda_j \mu_j \mu_{j+n}). \quad (22)$$

We can then obtain the operator form:

$$\hat{\rho} = 2^{-n} \prod_{j=1}^n (1 + i\lambda_j \hat{\gamma}_j \hat{\gamma}_{j+n}) \quad (23)$$

where $\hat{\gamma}$'s are a new set of Majorana operators. In the fermion basis

$$\hat{d}_j = \frac{\hat{\gamma}_j - i\hat{\gamma}_{j+n}}{2}, \quad \hat{d}_j^\dagger = \frac{\hat{\gamma}_j + i\hat{\gamma}_{j+n}}{2}, \quad (24)$$

the density matrix has the form

$$\hat{\rho} = \prod_j \left(\frac{1+\lambda_j}{2} - \lambda_j \hat{d}_j^\dagger \hat{d}_j \right) = \bigotimes_j \begin{bmatrix} \frac{1+\lambda_j}{2} & 0 \\ 0 & \frac{1-\lambda_j}{2} \end{bmatrix}_j. \quad (25)$$

Without loss of generality, we assume $\lambda_i \geq 0$. For pure state, $\lambda_i = 1, \forall i$. For mixed state, the entropy of ρ is just

$$S(\hat{\rho}) = \sum_j H\left(\frac{1+\lambda_j}{2}\right) = - \sum_j \left[\left(\frac{1+\lambda_j}{2}\right) \log\left(\frac{1+\lambda_j}{2}\right) + \left(\frac{1-\lambda_j}{2}\right) \log\left(\frac{1-\lambda_j}{2}\right) \right]. \quad (26)$$

2. Gaussian Operators

Definition 2. An operator \hat{X} (with nonzero trace) is Gaussian if

$$X(\theta) = C \exp\left(\frac{i}{2} \theta^T M \theta\right)$$

for some complex number C and some **complex antisymmetric** matrix M . M is called a correlation matrix of \hat{X} . If \hat{X} is traceless, it should be thought of as a limit $\hat{X} = \lim_{m \rightarrow \infty} \hat{X}_m$ for some converging sequence of Gaussian operators with nonzero trace.

Note that for traceless \hat{X} , the explicit form of $X(\theta)$ is

$$X(\theta) = C \left(\prod_{a=1}^{2k} \mu_a \right) \exp\left(\frac{i}{2} \sum_{a,b=2k+1}^{2n} M_{ab} \mu_a \mu_b\right), \quad (27)$$

where $\mu_a = \sum_b T_{ab} \theta_b$ for some invertible complex matrix T . The factor is a limiting point of the sequence:

$$\prod_{a=1}^{2k} \mu_a = \lim_{t \rightarrow \infty} \prod_{a=1}^k \left(\mu_{2a-1} \mu_{2a} + \frac{1}{t} \right) = \lim_{t \rightarrow \infty} \frac{1}{t^k} \exp \left(t \sum_{a=1}^k \mu_{2a-1} \mu_{2a} \right).$$

Introducing the operator $\hat{\Lambda} \equiv \sum_{a=1}^{2n} \hat{\omega}_a \otimes \hat{\omega}_a$, we have the following theorem:

Theorem 2. *An operator \hat{X} is Gaussian iff \hat{X} is even and satisfies*

$$[\hat{\Lambda}, \hat{X} \otimes \hat{X}] = 0.$$

Proof. The adjoint action of $\hat{\Lambda}$ in the Grassmann representation has the form:

$$\Lambda_{\text{ad}} = 2 \sum_a \left(\theta_a \otimes \frac{\partial}{\partial \theta_a} + \frac{\partial}{\partial \theta_a} \otimes \theta_a \right) \equiv \sum_a \Delta_a. \quad (28)$$

That is, $[\hat{\omega}_a \otimes \hat{\omega}_a, Y \otimes Z](\theta) = \Delta_a \cdot Y(\theta) \otimes Z(\theta)$ for any operators Y, Z having the same parity. Without loss of generality, both Y and Z are monomials in $\hat{\omega}$'s. In this case each of them either commutes or anticommutes with $\hat{\omega}_a$. Consider two cases:

1. Both Y and Z contain $\hat{\omega}_a$, or both Y and Z do not contain $\hat{\omega}_a$. Then the commutator $[\hat{\omega}_a \otimes \hat{\omega}_a, Y \otimes Z]$ is zero since both factors yield the same sign. The right-hand side is also zero, since either θ_a or $\partial/\partial \theta_a$ annihilates both Y and Z .
2. Y contains $\hat{\omega}_a$ while Z does not contain $\hat{\omega}_a$ (or vice versa). In this case $\hat{\omega}_a \otimes \hat{\omega}_a$ anticommutes with $Y \otimes Z$. Let us write $Y = \hat{\omega}_a \tilde{Y}$, where \tilde{Y} is a monomial which does not contain $\hat{\omega}_a$. We have:

$$[\hat{\omega}_a \otimes \hat{\omega}_a, Y \otimes Z] = 2(\hat{\omega}_a \otimes \hat{\omega}_a)(Y \otimes Z) = 2\tilde{Y} \otimes (\hat{\omega}_a Z).$$

On the other hand,

$$\theta_a \otimes \frac{\partial}{\partial \theta_a} \cdot Y \otimes Z = 0, \quad \frac{\partial}{\partial \theta_a} \otimes \theta_a \cdot Y \otimes Z = \tilde{Y} \otimes \theta_a Z.$$

We again get equality.

Necessity: Note that Λ_{ad} is invariant under change of variables since

$$\mu_a = \sum_b T_{ab} \theta_b, \quad \frac{\partial}{\partial \mu_a} = \sum_b (T^{-1})_{ab} \frac{\partial}{\partial \theta_b} \implies \sum_a \theta_a \otimes \frac{\partial}{\partial \theta_a} = \sum_a \mu_a \otimes \frac{\partial}{\partial \mu_a}.$$

Direct application of the operator to the general Gaussian form will prove the necessity.

Sufficiency: Denote $C = 2^{-n} \text{tr}(X) \equiv X(0)$ and represent $X(\theta)$ as

$$X(\theta) = C \cdot 1 + \frac{iC}{2} \sum_{a,b=1}^{2n} M_{ab} \theta_a \theta_b + \text{higher order terms}.$$

Applying a differential operator $1 \otimes \frac{\partial}{\partial \theta_b}$ to both sides:

$$\sum_{a=1}^{2n} \left(\theta_a X \otimes \frac{\partial^2}{\partial \theta_b \partial \theta_a} X - \frac{\partial}{\partial \theta_a} X \otimes \theta_a \frac{\partial}{\partial \theta_b} X \right) + \frac{\partial}{\partial \theta_b} X \otimes X = 0.$$

Now let us put $\theta \equiv 0$ in the second factor:

$$\frac{\partial}{\partial \theta_b} X = i \sum_{a=1}^{2n} M_{ba} \theta_a X.$$

This differential equation can be easily solved by $X(\theta) = C \exp \left(\frac{i}{2} \theta^T M \theta \right)$.

For general cases, we denote $\mathcal{K} \subseteq \mathcal{M}_1$ a subspace spanned by linear functions which annihilate \hat{X} , i.e.

$$\mathcal{K} = \{f \in \mathcal{M}_1 : f(\theta)X(\theta) = 0\}.$$

Let us perform a linear change of variables $\mu_a = \sum_b T_{ab}\theta_b$, with T being an invertible complex matrix chosen such that the first k variables μ span the subspace \mathcal{K} , i.e. $\mathcal{K} = \text{span}[\mu_1, \dots, \mu_{2k}]$. From equalities $\mu_j X = 0$, $j \in [1, 2k]$ it follows that

$$X(\theta(\mu)) = \left(\prod_a \mu_a \right) \tilde{X}(\mu),$$

where $\tilde{X}(\mu)$ depends only upon $\mu_{2k+1}, \dots, \mu_{2n}$. The function $\tilde{X}(\mu)$ satisfies the equation

$$\sum_{a=2k+1}^{2n} \left(\mu_a \otimes \frac{\partial}{\partial \mu_a} + \frac{\partial}{\partial \mu_a} \otimes \mu_a \right) \tilde{X} \otimes \tilde{X} = 0.$$

Therefore we get the general Gaussian form. □

3. Gaussian Linear Maps

We define linear maps that preserve Gaussian states as the following:

Definition 3. A linear map \mathcal{E} is Gaussian iff it admits an integral representation

$$\mathcal{E}(X)(\theta) = C \int D[\eta, \mu] \exp [S(\theta, \eta) + i\eta^T \mu] X(\mu), \quad (29)$$

where

$$S(\theta, \eta) = \frac{i}{2} (\theta^T, \eta^T) \begin{pmatrix} A & B \\ -B^T & D \end{pmatrix} \begin{pmatrix} \theta \\ \eta \end{pmatrix} \quad (30)$$

for some complex $2n \times 2n$ matrices A, B, D , and some complex number C .

Consider a Gaussian operator \hat{X} which can be described by a correlation matrix M and a Gaussian map \mathcal{E} . Applying the Gaussian integration, one can show that $\mathcal{E}(X)$ has a correlation matrix

$$\mathcal{E}(M) = A + B (M^{-1} + D)^{-1} B^T = A + B (I + MD)^{-1} M B^T,$$

while a pre-exponential factor of the operator $\mathcal{E}(X)$ can be found from an identity

$$\text{tr}(\mathcal{E}(X)) = C(-1)^n \text{Pf}(M) \text{Pf}(M^{-1} + D) \text{tr}(X).$$

The value of $\text{tr}(\mathcal{E}(X))$ can be found up to a factor ± 1 using a regularized version:

$$\text{tr}(\mathcal{E}(X))^2 = C^2 \det(I + MD) \text{tr}(X)^2.$$

B. Operator Form

1. Dirac Fermion Case

For particle number conserving systems, the Gaussian state can be represented as a matrix:

$$|B\rangle \equiv \prod_{j=1}^N \sum_i B_{ij} c_i^\dagger |0\rangle \equiv \bigotimes_{j=1}^N |B_j\rangle. \quad (31)$$

Note that the matrix B representing the Gaussian state has the unitary degree of freedom

$$|B\rangle = |B'\rangle, \quad B'_{ij} = \sum_k B_{ik} U_{kj},$$

where U_{kj} is an $N \times N$ unitary matrix. It means that the Gaussian state is determined by the linear subspace that columns of B span. The columns of B need not to be orthogonal, while the canonical form can be obtained by the QR decomposition: $B_{L \times N} = Q_{L \times N} \cdot R_{N \times N}$, where the Q matrix is orthonormal and we can set $B' = Q$.

A free fermion state maintains its structure when applied to a quasi-particle creation/annihilation operator. Consider a general quasi-particle $b^\dagger = \sum_i b_i c_i^\dagger$, creating a quasiparticle is simply adding a column to B , since

$$b^\dagger |B\rangle = \sum_k b_k c_k^\dagger \prod_{j=1}^N \sum_i c_i^\dagger B_{ij} |0\rangle = \prod_{j=1}^{N+1} \sum_i c_i^\dagger [b|B]_{ij} |0\rangle \quad (32)$$

In general, the new column b is not orthogonal to linear space B , therefore orthogonalization procedure is needed to obtain canonical form.

Using the Baker-Campbell-Hausdorff formula $e^X Y e^{-X} = \exp(\text{ad } X) Y$,

$$e^{-iHt} c_j^\dagger e^{iHt} = c_k^\dagger [e^{-iHt}]_{kj} \implies e^{-iHt} |B\rangle = \prod_{j=1}^N \sum_i [e^{-iHt}]_{ki} B_{ij} c_k^\dagger |0\rangle = |e^{-iHt} \cdot B\rangle. \quad (33)$$

For the quasiparticle annihilation operator b ,

$$b |B\rangle = \sum_k b_k^* c_k \prod_j \sum_i c_i^\dagger B_{ij} |0\rangle = \sum_j \langle b|B_j\rangle \bigotimes_{l \neq j} |B_l\rangle. \quad (34)$$

We can use the gauge freedom to restrict $\langle b|B'_j\rangle = 0$ for $j > 1$. Such matrix B' always exists since we can always find a column j that $\langle b|B_j\rangle \neq 0$ (otherwise $p_m = 0$ and the jump is impossible). We then move the column to the first and define the column as

$$|B'_j\rangle = |B_j\rangle - \frac{\langle a|B_j\rangle}{\langle a|B_1\rangle} |B_1\rangle, \quad j > 1. \quad (35)$$

Such column transformations do not alter the linear space B spans, while the orthogonality and the normalization might be affected.

2. Majorana Case

For the Majorana case, the canonical form (25) for a Gaussian pure state $|\psi\rangle$ can be reformulated as

$$|\psi\rangle\langle\psi| = \prod_{j=1}^n \hat{d}_j^\dagger \hat{d}_j, \quad \hat{d}_j^\dagger = \frac{\hat{\gamma}_j + i\hat{\gamma}_{j+n}}{2} = \sum_{i=1}^n \left(\frac{R_{i,j} + iR_{i,j+n}}{2} \right) \hat{c}_i + \left(\frac{R_{i+n,j} + iR_{i+n,j+n}}{2} \right) \hat{c}_i^\dagger. \quad (36)$$

Note that the state is annihilated by $\{\hat{d}_j^\dagger\}$. We can store the information of $|\psi\rangle$ into a $2n \times n$ complex matrix

$$|\psi\rangle \iff B = \frac{1}{2} \begin{bmatrix} R_{11} + iR_{12} \\ R_{21} + iR_{22} \end{bmatrix}. \quad (37)$$

The rest of the procedures are parallel to those of the Dirac fermion case.

[1] S. Bravyi, *Lagrangian representation for fermionic linear optics* (2004), quant-ph/0404180.