27496번 - 발머의 피크 이론 성공

☆

실버 III

시간 제한	메모리 제한	제출	정답	맞힌 사람	정답 비율
1 초 (추가 시간 없음)	1024 MB	2347	977	845	42.893%

문제

발머의 피크 이론이란 혈중 알코올 농도 C가 $(0.129 \le C \le 0.138)$일때 초인적인 프로그래밍 능력을 가지게 된다는 이론이다. 기령이는 발머의 피크 이론 신봉자이기 때문에 지금부터 코딩테스트를 볼 때까지 혈중 알코올 농도를 최대한 지키려고 한다. 기령이는 매 시간마다 정해진 술을 섭취하며, 섭취한 알코올은 일정 시간이 지나면 분해된다. 기령이가 지금부터 코딩테스트를 볼 때까지 얼마나 혈중 알코올 농도를 0.129와 0.138 사이로 지킬 수 있는지 알아내보자. 단, 혈중 알코올 농도 증가량은 술에 포함된 알코올의 양 $A \times 0.001$로 계산하며 최초의 혈중 알코올 농도는 0이다.

입력

첫째 줄에 코딩테스트를 볼 때까지 남은 시간 $N(1 \le N \le 1\,000\,000)$, 섭취한 알코올의 지속시간 $L(1 \le L \le 10\,000, L \le N)$이 공백으로 구분되어 주어진다.

둘째 줄에 술에 포함된 알코올의 양 정수 $a_i(0 \le a_i \le 200)$가 공백으로 구분되어 주어진다.

출력

혈중 알코올 농도 C를 $(0.129 \le C \le 0.138)$로 유지한 시간을 출력한다.

예제 입력 1 복사

```
5 3
45 44 44 43 40
```

예제 출력 1 복사

```
2
```

각 시간에 따른 혈중 알코올 농도는 {0.045, 0.089, **0.133**, **0.131**, 0.127}이다. 따라서 지금으로부터 2시간 후와 3시간 후, 총 두 시간 동안 혈중 알코올 농도를 유지할 수 있다.

예제 입력 2 복사

```
2 1
129 138
```

예제 출력 2 복사

```
2
```

출처

Camp (/category/220) > ICPC Sinchon Algorithm Camp (/category/499) > 2023 ICPC Sinchon Winter Algorithm Camp Contest (/category/801) > 초급 (/category/detail/3494) C번

- 문제를 만든 사람: lickelon (/user/lickelon)
- 문제를 검수한 사람: eaststar (/user/eaststar), man_of_learning (/user/man_of_learning), pinebananais (/user/pinebananais), starbow (/user/starbow), tae826 (/user/tae826), tolelom (/user/tolelom), ystaeyoon113 (/user/ystaeyoon113)

알고리즘 분류

- 누적 합 (/problem/tag/139)
- 슬라이딩 윈도우 (/problem/tag/68)

메모