1

Assignment 1 Probability and Random Processes

Padala Sai Kowshik* EE22BTECH11038

Question 1.4.2

Find the intersection **O** of the perpendicular bisectors of **AB** and **AC**.

Solution:

Given,

$$\mathbf{A} = \begin{pmatrix} 1 \\ -1 \end{pmatrix}, \ \mathbf{B} = \begin{pmatrix} -4 \\ 6 \end{pmatrix}, \ \mathbf{C} = \begin{pmatrix} -3 \\ -5 \end{pmatrix}$$

Vector equation of perpendicular bisector of $\mathbf{A} - \mathbf{B}$ is

$$(\mathbf{A} - \mathbf{B})^{\mathsf{T}} \left(\mathbf{x} - \frac{\mathbf{A} + \mathbf{B}}{2} \right) = 0 \tag{1}$$

where,

$$\mathbf{A} + \mathbf{B} = \begin{pmatrix} 1 \\ -1 \end{pmatrix} + \begin{pmatrix} -4 \\ 6 \end{pmatrix} \tag{2}$$

$$= \begin{pmatrix} -3\\5 \end{pmatrix} \tag{3}$$

$$\mathbf{A} - \mathbf{B} = \begin{pmatrix} 1 \\ -1 \end{pmatrix} - \begin{pmatrix} -4 \\ 6 \end{pmatrix} \tag{4}$$

$$= \begin{pmatrix} 5 \\ -7 \end{pmatrix} \tag{5}$$

(6)

$$\implies (\mathbf{A} - \mathbf{B})^{\mathsf{T}} = \begin{pmatrix} 5 & -7 \end{pmatrix}$$

 \therefore The vector equation of $\mathbf{O} - \mathbf{F}$ is

$$\left(5 \quad -7\right)\left(\mathbf{x} - \frac{1}{2}\begin{pmatrix} -3\\ 5 \end{pmatrix}\right) = 0\tag{7}$$

$$\implies (5 \quad -7)\mathbf{x} = \frac{1}{2} (5 \quad -7) \begin{pmatrix} -3\\ 5 \end{pmatrix} \qquad (8)$$

Performing matrix multiplication yields

Vector equation of perpendicular bisector of A - C is

$$(\mathbf{A} - \mathbf{C})^{\mathsf{T}} \left(\mathbf{x} - \frac{\mathbf{A} + \mathbf{C}}{2} \right) = 0 \tag{10}$$

where,

$$\mathbf{A} + \mathbf{C} = \begin{pmatrix} 1 \\ -1 \end{pmatrix} + \begin{pmatrix} -3 \\ -5 \end{pmatrix} \tag{11}$$

$$= \begin{pmatrix} -2\\ -6 \end{pmatrix} \tag{12}$$

$$\mathbf{A} - \mathbf{C} = \begin{pmatrix} 1 \\ -1 \end{pmatrix} - \begin{pmatrix} -3 \\ -5 \end{pmatrix} \tag{13}$$

$$= \begin{pmatrix} 4 \\ 4 \end{pmatrix} \tag{14}$$

$$\implies (\mathbf{A} - \mathbf{C})^{\mathsf{T}} = \begin{pmatrix} 4 & 4 \end{pmatrix} \tag{15}$$

 \therefore The vector equatio of $\mathbf{O} - \mathbf{E}$ is

$$\begin{pmatrix} 4 & 4 \end{pmatrix} \begin{pmatrix} \mathbf{x} - \frac{1}{2} \begin{pmatrix} -2 \\ -6 \end{pmatrix} \end{pmatrix} = 0 \tag{16}$$

$$\implies \begin{pmatrix} 4 & 4 \end{pmatrix} \mathbf{x} = \frac{1}{2} \begin{pmatrix} 4 & 4 \end{pmatrix} \begin{pmatrix} -2 \\ -6 \end{pmatrix} \tag{17}$$

Performing matrix multiplication yields

$$\begin{pmatrix} 4 & 4 \end{pmatrix} \mathbf{x} = -16 \tag{18}$$

$$\implies (1 \quad 1)\mathbf{x} = -4 \tag{19}$$

$$\begin{pmatrix} 5 & -7 & -25 \\ 1 & 1 & -4 \end{pmatrix} \xrightarrow{R_2 \leftarrow 5R_2 - R_1} \begin{pmatrix} 5 & -7 & -25 \\ 0 & 12 & 5 \end{pmatrix} \tag{20}$$

(8)
$$\begin{pmatrix} 3 & -7 & -23 \\ 0 & 12 & 5 \end{pmatrix} & \stackrel{R_1 \leftarrow \frac{7}{7}R_1 + R_2}{\longleftrightarrow} & \begin{pmatrix} \frac{27}{7} & 0 & \frac{-7}{7} \\ 0 & 12 & 5 \end{pmatrix} & (21)$$

$$\begin{pmatrix} \frac{60}{7} & 0 & \frac{-265}{7} \\ 0 & 12 & 5 \end{pmatrix} & \stackrel{R_2 \leftarrow \frac{1}{12}R_2}{\longleftrightarrow} & \begin{pmatrix} 1 & 0 & \frac{-53}{12} \\ 0 & 1 & \frac{5}{12} \end{pmatrix} & (22)$$

$$\begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} \frac{-53}{12} \\ \frac{-53}{12} \end{pmatrix}$$

$$\therefore \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \mathbf{x} = \begin{pmatrix} \frac{-53}{12} \\ \frac{5}{12} \end{pmatrix} \tag{23}$$

$$\implies \mathbf{x} = \begin{pmatrix} \frac{-53}{12} \\ \frac{5}{12} \end{pmatrix} \tag{24}$$

Therefore, the point of intersection of perpendicular bisectors of $\mathbf{A} - \mathbf{B}$ and $\mathbf{A} - \mathbf{C}$ is $\mathbf{O} = \frac{1}{12} \begin{pmatrix} -53 \\ 5 \end{pmatrix}$




Fig. 0. $\mathbf{O} - \mathbf{E}$ and $\mathbf{O} - \mathbf{F}$ are perpendicular bisectors of $\mathbf{A} - \mathbf{C}$ and $\mathbf{A} - \mathbf{B}$ respectively