### **Academic Presentations**

### A LaTeX Template Using the beamer Class

Adam Smith<sup>a, c</sup> Janet Smith<sup>b, c</sup> Jeremiah Smith<sup>a</sup>

<sup>a</sup> University of Bonn, Germany

<sup>b</sup> University of Cologne, Germany

<sup>c</sup> Collaborative Research Center Transregio 224

### Name of the Inviting Institution/Seminar Series

December 17, 2023

### **Outline**

- 1 Introduction
- 2 Study Design
- 3 Results
- **4** Discussion
- **5** Math "Torture" Test
- **6** References

### **Introduction 1: Choice of a Reasonable Aspect Ratio**

When preparing a presentation, we often do not know whether the native aspect ratio of the projector in the seminar room/lecture hall will be 4:3 or 16:9 (or 16:10).

In this case, it may be a good idea to choose an **intermediate aspect ratio**, see https://github.com/josephwright/beamer/issues/497. The idea behind this recommendation is that it minimizes the average loss of available space.

Hence, these templates include a presentation in the **14:9** aspect ratio (see https://en.wikipedia.org/wiki/14:9\_aspect\_ratio): while it is imperfect for probably every projector that you will encounter, it is good on average for all of them.

(Please note that 14:9  $\doteq$  1.556, which is pretty close to the "officially" recommended 20:13  $\doteq$  1.5385.)

Great Minds Discuss Ideas.

Average Minds Discuss Events.

Small Minds Discuss People.

—https://quoteinvestigator.com/2014/11/18/great-minds/

### **Background**

- Temporal discounting is key concept in economics.
- Normative model: exponential discounting. However, observed decisions are hard to explain (e.g., Dohmen et al., 2012).
- One alternative: the "focusing model" by Kőszegi and Szeidl (2013).
- See also Lisi (1995).

### **Research Question**

- The composition of latex and of typical rubbers is given below.
- Is it true that trees are regularly tapped and the coagulated latex which exudes is collected and worked up into rubber?

### **Research Question**

- The composition of latex and of typical rubbers is given below.
- Is it true that trees are regularly tapped and the coagulated latex which exudes is collected and worked up into rubber?

### **Preview of the Results**

- There is no feasible method at present known of preventing the inclusion of the resin of the latex with the rubber during coagulation.
- ⇒ Although the separation of the resin from the solid caoutchouc by means of solvents is possible, it is not practicable or profitable commercially.

### Introduction 4: Beamer block Environments

### Block title example: 0123456789 äöüß ÄÖÜ €. Often finding flowers in official fjords

The block environment. Block title example: 0123456789 äöüß ÄÖÜ €. Often finding flowers in official fjords.

### An exemplary example

I am the exampleblock environment. Use me for examples.

### **Summary: Things to remember**

The alertblock environment. Use this environment for really important stuff. The alertblock environment.

### Introduction 5: Beamer block Environment with Different Colors

### A block in the default color

The block environment. The block environment. The block environment. The block environment. A block in the default color.

### A block in yellow

The block environment. The block environment. The block environment. The block environment. A block in yellow.

### A block in the default color

The block environment. The block environment. The block environment. The block environment. A block in the default color.

### Introduction 6: Beamer definition and theorem Environments

# Definition (A Very, Very, Very, Very, Very, Very Long Name of a Concept that Spans Two Lines)

The definition environment. Upright.

### **Theorem (Theorem's Name)**

The theorem environment. Italic.

### Lemma (Lemma's Name)

The lemma environment. Italic.

### **Corollary (Corollary's Name)**

The corollary environment. Italic.

### **Proof of Theorem's Name**

The proof environment. Upright.

# Study Design

### **Study Design 1: Design of the Study**

- The latex of the best rubber plants furnishes from 20% to 50% of rubber.
- As the removal of the impurities of the latex is one of the essential points to be aimed at, it was thought that the use of a centrifugal machine to separate the caoutchouc as a cream from the watery part of the latex would prove to be a satisfactory process.

### **Study Design 2: Design of the Study**

The watery portion of the latex soaks into the trunk, and the soft spongy rubber which remains is kneaded and pressed into lumps or balls:

 $BAL_{1:1}^{I}$ ,  $BAL_{1:1}^{II}$ : Each payment transferred on single day.

**UNBAL** $_{1:n}^{1}$ : Earlier payoff concentrated, while later payoff dispersed over n = 2, 4, or 8 dates.

UNBAL $_{n:1}^{II}$ : Earlier payoff dispersed over n = 2, 4, or 8 dates, while later payoff concentrated.

### **Study Design 3: Control Experiment**

- · Control for alternative explanations.
- Many of the example sentences were taken from http://sentence.yourdictionary. com/latex.

### Study Design 4: An Example enumerate List

- 1. First itemtext
  - a. First itemtext
    - i. First itemtext
    - ii. Second itemtext
    - iii. Last itemtext
    - iv. First itemtext
  - b. Second itemtext
  - c. Last itemtext
  - d. First itemtext
- 2. Second itemtext
- 3. Last itemtext
- 4. First itemtext

### Study Design 5: An Example itemize List

- First itemtext
  - First itemtext
    - First itemtext
    - Second itemtext
    - Last itemtext
    - First itemtext
  - Second itemtext
  - Last itemtext
  - First itemtext
- Second itemtext
- Last itemtext
- First itemtext

### **Study Design 6: Some Example Text**

### Let's include some Greek letters: $\alpha$ , $\beta$ , $\sigma$

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.  $(a, \alpha), \beta, \sigma, (v, v), (v, v)$ 

Test: χρστ χρς τ χρσ τ. Math mode, upright: σ

### **Study Design 7: Some Example Formulas**

Let's include some additional Greek letters:  $\gamma$ ,  $\phi$ ,  $\sigma_{\varepsilon}$ ,  $c^{\alpha}$ 

$$p(R,\phi) \sim \int_{-\infty}^{\infty} \frac{\tilde{W}_n(\gamma) \exp\left[ \iota R/a \left( \sqrt{k^2 a^2 - \gamma^2} \cos \phi \right) \right]}{(k^2 a^2 - \gamma^2)^{3/4} H_n^{\prime (1)} \left( \sqrt{k^2 a^2 - \gamma^2} \right)} d\gamma$$

### **Study Design 7: Some Example Formulas**

Let's include some additional Greek letters:  $\gamma$ ,  $\phi$ ,  $\sigma_{\varepsilon}$ ,  $c^{\alpha}$ 

$$p(R,\phi) \sim \int_{-\infty}^{\infty} \frac{\tilde{W}_n(\gamma) \exp\left[ \iota R/a \left( \sqrt{k^2 a^2 - \gamma^2} \cos \phi \right) \right]}{(k^2 a^2 - \gamma^2)^{3/4} H_n^{\prime (1)} \left( \sqrt{k^2 a^2 - \gamma^2} \right)} d\gamma$$

Let's include upright Latin letters in math mode: d, e (next slide)

$$\int_a^b f(x) \, \mathrm{d}x = F(b) - F(a)$$

### **Study Design 7: Some Example Formulas**

Let's include some additional Greek letters:  $\gamma$ ,  $\phi$ ,  $\sigma_{\varepsilon}$ ,  $c^{\alpha}$ 

$$p(R,\phi) \sim \int_{-\infty}^{\infty} \frac{\tilde{W}_n(\gamma) \exp\left[ \iota R/a \left( \sqrt{k^2 a^2 - \gamma^2} \cos \phi \right) \right]}{(k^2 a^2 - \gamma^2)^{3/4} H_n^{\prime (1)} \left( \sqrt{k^2 a^2 - \gamma^2} \right)} d\gamma$$

Let's include upright Latin letters in math mode: d, e (next slide)

$$\int_a^b f(x) \, \mathrm{d}x = F(b) - F(a)$$

Let's test the math bold styles (both upright and italic)

$$\mathbf{\Sigma} := \mathbf{Cov}(\mathbf{X}) = \begin{bmatrix} \operatorname{Var}(X_1) & \cdots & \operatorname{Cov}(X_1, X_n) \\ \vdots & \ddots & \vdots \\ \operatorname{Cov}(X_n, X_1) & \cdots & \operatorname{Var}(X_n) \end{bmatrix}$$

### Study Design 8: Additional Example Formulas (with upright $\pi$ )

Only variables are set in italics according to ISO style—hence, we use upright "d," "e," and " $\pi$ " (\mathup{d}, \mathup{e}, and \mathup{\pi}, respectively).

### Theorem (simplest form of the Central Limit Theorem)

Let  $X_1, X_2, \ldots, X_n$  be a sequence of i.i.d. random variables with mean 0 and variance 1 on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . Then

$$\mathbb{P}\left(\frac{X_1+\cdots+X_n}{\sqrt{n}}\leq y\right)\to \mathfrak{N}(y):=\int_{-\infty}^y\frac{\mathrm{e}^{-v^2/2}}{\sqrt{2\pi}}\,\mathrm{d}v\quad as\quad n\to\infty,$$

or, equivalently, letting  $S_n := \sum_{1}^{n} X_k$ ,

$$\mathbb{E}f(\mathsf{S}_n/\sqrt{n})\to \int_{-\infty}^{\infty}f(\mathsf{v})\frac{\mathrm{e}^{-\mathsf{v}^2/2}}{\sqrt{2\pi}}\,\mathsf{d}\mathsf{v}\quad \text{as }n\to\infty\text{, for every }f\in\mathsf{b}\mathscr{C}(\mathbb{R}).$$

### Study Design 9: An siunitx Example Table

**Table 1.** Overview of the choice lists presented to subjects (adapted from Gerhardt, Schildberg-Hörisch, and Willrodt, 2017).

		Alternative <b>A</b>			Alternative <b>B</b>			
	CA,1	PA,1	C <sub>A,2</sub>	PA,2	CB,1	P <sub>B,1</sub>	CB,2	р <sub>в,2</sub>
Choice List I: risky/risky (x = €	22.00, r = €7.50, k	= €11.5	0; 25 rows)					
Top row	€ 3.00	50%	€22.00	50%	€ 3.00	50%	€ 7.00	509
Center row	€ 3.00	50%	€22.00	50%	€ 9.00	50%	€13.00	509
Row with $m = 0$	€ 3.00	50%	€22.00	50%	€10.50	50%	€14.50	509
Bottom row	€ 3.00	50%	€22.00	50%	€15.00	50%	€19.00	509
Choice List II: safe/risky (x = 0	16.00, r = €5.00, k	= €5.00	; 19 rows)					
Top row	€11.00	100%			€11.00	50%	€21.00	509
Center row	€11.00	100%			€ 6.50	50%	€16.50	509
Row with $m = 0$	€11.00	100%			€ 6.00	50%	€16.00	509
Bottom row	€11.00	100%			€ 2.00	50%	€12.00	509
Choice List III: "long shot" (x =	€14.00, r = -€36.0	00, k = €	7.00; 21 ro	ws)				
Top row	€ 7.00	90%	€50.00	10%	€ 7.00	90%	€10.00	109
Row with $m = 0$	€ 7.00	90%	€50.00	10%	€11.00	90%	€14.00	109
Center row	€ 7.00	90%	€50.00	10%	€12.00	90%	€15.00	109
Bottom row	€ 7.00	90%	€50.00	10%	€17.00	90%	€20.00	109
Choice List IV: delayed payoffs	(x = €18.00, r = €6	5.00, k =	€8.50, paid	d in one v	veek; 20 row	vs)		
Top row	€ 9.50	50%	€12.00	50%	€ 9.50	50%	€24.00	509
Above-center row	€ 9.50	50%	€12.00	50%	€ 5.00	50%	€19.50	509
Below-center row	€ 9.50	50%	€12.00	50%	€ 4.50	50%	€19.00	509
Row with $m = 0$	€ 9.50	50%	€12.00	50%	€ 3.50	50%	€18.00	509
Bottom row	€ 9.50	50%	€12.00	50%	€ 0.00	50%	€14.50	509

## **Results**

### **Results 1: Overview**

**1.** As a secondary function we may recognize the power of closing wounds, which results from the rapid coagulation of exuded latex in contact with the air:

### **Results 1: Overview**

- **1.** As a secondary function we may recognize the power of closing wounds, which results from the rapid coagulation of exuded latex in contact with the air:
  - **a.** In some cases (Allium, Convolvulaceae, etc.) rows of cells with latex-like contents occur.
  - b. However, the walls separating the individual cells do not break down.

### **Results 1: Overview**

- **1.** As a secondary function we may recognize the power of closing wounds, which results from the rapid coagulation of exuded latex in contact with the air:
  - **a.** In some cases (Allium, Convolvulaceae, etc.) rows of cells with latex-like contents occur.
  - **b.** However, the walls separating the individual cells do not break down.
- 2. The rows of cells from which the laticiferous vessels are formed can be distinguished (6.3 p.p. vs. 2.6 p.p.; p < 0.01).

### **Results 2: Our Main Results**

The charts are taken from Dertwinkel-Kalt et al. (2017).





Figure 1. (A) Difference between treatment and control condition. (B) Heterogeneity.

### **Results 3: Main vs. Control Experiment**

Rule out some alternative explanations (Dertwinkel-Kalt et al., 2017).



### Results 4: Another siunitx Example Table

Table 2. Example of a regression table (adapted from Gerhardt, Schildberg-Hörisch, and Willrodt, 2017). Never forget to mention the dependent variable (here,  $m_{\sim}$ )!

	(1)	(2)	(3)	(4)	(5)
Treatment	-0.390	-0.228	-0.729*	-0.449*	-0.453**
	(+0.352)	(-0.205)	[+0.377]	[-0.245]	{+0.204}
Female	0.948***	0.061	0.188	0.305	0.385*
	(0.354)	(0.233)	(0.372)	-0.449* [-0.245]	(0.222)
Female × Treatment	0.169	0.251	0.892*	0.454	0.439
	(0.514)	(0.325)	(0.533)	(0.341)	(0.307)
Final high school grade	-0.101	0.013	0.076	0.117	0.039
	(0.198)	(0.144)	(0.224)	(0.146)	(0.133)
Trait self-control	-0.016	0.002	-0.016	-0.000	-0.007
rait self-control	(0.016)	(0.010)	(0.015)	(0.010)	(0.009)
Constant	2.357***	1.512***	-0.322	2.158***	1.437**
	(0.239)	(0.144)	(0.265)	[-0.245] 0.305 (0.226) 0.454 (0.341) 0.117 (0.146) -0.000 (0.010) 2.158*** (0.161) 304 0.043	(0.152)
Observations	303	289	295	304	1191
$R^2$	0.057	0.008	0.039	0.043	0.024
Treatment × (1 + Female)	-0.221	0.023	0.163	0.004	-0.014
$p_F[Treatment \times (1 + Female) = 0]$	0.327	0.008	0.192	0.000	0.003

Notes: Dependent variable:  $m_-$ . Robust standard errors (cluster-corrected for column 5) in parentheses. \*\*\* p < 0.01. Missing observations (N < 308) due to exclusion of trials in which subjects behaved irrationally (i.e., chose a dominated option). The regressors Final high school grade and Trait self-control are mean-centered.

Table 3. Revealing a table column-by-column makes it easier to comprehend.

Treatment	
Female	
Female × Treatment	
Final high school grade	
Trait self-control	
Constant	
Observations $R^2$	
$\begin{aligned} & \text{Treatment} \times (1 + \text{Female}) \\ & p_f[\text{Treatment} \times \\ & (1 + \text{Female}) = 0] \end{aligned}$	

Table 3. Revealing a table column-by-column makes it easier to comprehend.

	(1)	
Treatment	-0.390	
	(+0.352)	
Female	0.948***	
	(0.354)	
Female × Treatment	0.169	
	(0.514)	
Final high school grade	-0.101	
	(0.198)	
Trait self-control	-0.016	
	(0.016)	
Constant	2.357***	
	(0.239)	
Observations	303	
$R^2$	0.057	
Treatment × (1 + Female)	-0.221	
$p_F$ [Treatment $\times$	0.327	
(1 + Female) = 0		

Table 3. Revealing a table column-by-column makes it easier to comprehend.

	(1)	(2)	
Treatment	-0.390	-0.228	
	(+0.352)	(-0.205)	
Female	0.948***	0.061	
	(0.354)	(0.233)	
Female × Treatment	0.169	0.251	
	(0.514)	(0.325)	
Final high school grade	-0.101	0.013	
	(0.198)	(0.144)	
Trait self-control	-0.016	0.002	
	(0.016)	(0.010)	
Constant	2.357***	1.512***	
	(0.239)	(0.144)	
Observations	303	289	
$R^2$	0.057	0.008	
Treatment × (1 + Female)	-0.221	0.023	
$p_F$ [Treatment $\times$	0.327	0.008	
(1 + Female) = 0]			

Table 3. Revealing a table column-by-column makes it easier to comprehend.

	(1)	(2)	(3)	
Treatment	-0.390	-0.228	-0.729*	
	(+0.352)	(-0.205)	[+0.377]	
Female	0.948***	0.061	0.188	
	(0.354)	(0.233)	(0.372)	
Female × Treatment	0.169	0.251	0.892*	
	(0.514)	(0.325)	(0.533)	
Final high school grade	-0.101	0.013	0.076	
	(0.198)	(0.144)	(0.224)	
Trait self-control	-0.016	0.002	-0.016	
	(0.016)	(0.010)	(0.015)	
Constant	2.357***	1.512***	-0.322	
	(0.239)	(0.144)	(0.265)	
Observations	303	289	295	
R <sup>2</sup>	0.057	0.008	0.039	
Treatment × (1 + Female)	-0.221	0.023	0.163	
$p_F$ [Treatment $\times$	0.327	0.008	0.192	
(1 + Female) = 0				

Table 3. Revealing a table column-by-column makes it easier to comprehend.

	(1)	(2)	(3)	(4)
Treatment	-0.390	-0.228	-0.729*	-0.449*
	(+0.352)	(-0.205)	[+0.377]	[-0.245]
Female	0.948***	0.061	0.188	0.305
	(0.354)	(0.233)	(0.372)	(0.226)
emale × Treatment	0.169	0.251	0.892*	0.454
	(0.514)	(0.325)	(0.533)	(0.341)
Final high school grade	-0.101	0.013	0.076	0.117
	(0.198)	(0.144)	(0.224)	(0.146)
Trait self-control	-0.016	0.002	-0.016	-0.000
	(0.016)	(0.010)	(0.015)	(0.010)
Constant	2.357***	1.512***	-0.322	2.158***
	(0.239)	(0.144)	(0.265)	(0.161)
Observations	303	289	295	304
R <sup>2</sup>	0.057	0.008	0.039	0.043
reatment × (1 + Female)	-0.221	0.023	0.163	0.004
$p_F$ [Treatment $ imes$	0.327	0.008	0.192	0.000
(1 + Female) = 0]				

Table 3. Revealing a table column-by-column makes it easier to comprehend.

	(1)	(2)	(3)	(4)	(5)
Treatment	-0.390	-0.228	-0.729*	-0.449*	-0.453**
	(+0.352)	(-0.205)	[+0.377]	[-0.245]	{+0.204}
Female	0.948***	0.061	0.188	0.305	0.385*
	(0.354)	(0.233)	(0.372)	-0.449* [-0.245]	(0.222)
Female × Treatment	0.169	0.251	0.892*	0.454	0.439
Female × Treatment	(0.514)	(0.325)	(0.533)	(0.341)	(0.307)
Final high school grade	-0.101	0.013	0.076	0.117	0.039
	(0.198)	(0.144)	(0.224)	(0.146)	(0.133)
Trait self-control	-0.016	0.002	-0.016	-0.000	-0.007
	(0.016)	(0.010)	(0.015)	(0.010)	(0.009)
Constant	2.357***	1.512***	-0.322	2.158***	1.437***
	(0.239)	(0.144)	(0.265)	(0.161)	(0.152)
Observations	303	289	295	304	1191
$R^2$	0.057	0.008	0.039	0.043	0.024
Treatment × (1 + Female)	-0.221	0.023	0.163	0.004	-0.014
$p_F$ [Treatment $\times$	0.327	0.008	0.192	0.000	0.003
(1 + Female) = 0					

### Results 6: Highlighting Cells in a Table

Table 4. There are various ways to highlight cells, columns, and rows in a table.

	(1)	(2)	(3)	(4)	(5)
Treatment	-0.390	-0.228	-0.729*	-0.449*	-0.453**
	(+0.352)	(-0.205)	[+0.377]	[-0.245]	{+0.204}
Female	0.948***	0.061	0.188	0.305	0.385*
	(0.354)	(0.233)	(0.372)	-0.449* [-0.245]	(0.222)
Female × Treatment	0.169	0.251	0.892*	0.454	0.439
	(0.514)	(0.325)	(0.533)	(0.341)	(0.307)
Final high school grade	-0.101	0.013	0.076	0.117	0.039
	(0.198)	(0.144)	(0.224)	(0.146)	(0.133)
Trait self-control	-0.016	0.002	-0.016	-0.000	-0.007
	(0.016)	(0.010)	(0.015)	(0.010)	(0.009)
Constant	2.357***	1.512***	-0.322	2.158***	1.437***
	(0.239)	(0.144)	(0.265)	(0.161)	(0.152)
Observations	303	289	295	304	1191
$R^2$	0.057	0.008	0.039	0.043	0.024
Treatment × (1 + Female)	-0.221	0.023	0.163	0.004	-0.014
$p_F[\text{Treatment} \times (1 + \text{Female}) = 0]$	0.327	0.008	0.192	0.000	0.003

Table 4. There are various ways to highlight cells, columns, and rows in a table.

	(1)	(2)	(3)	(4)	(5)
Treatment	-0.390	-0.228	-0.729*	-0.449*	-0.453**
	(+0.352)	(-0.205)	[+0.377]	[-0.245]	{+0.204}
Female	0.948***	0.061	0.188	0.305	0.385*
	(0.354)	(0.233)	(0.372)	(0.226)	(0.222)
Female × Treatment	0.169	0.251	0.892*	0.454	0.439
	(0.514)	(0.325)	(0.533)	(0.341)	(0.307)
Final high school grade	-0.101	0.013	0.076	0.117	0.039
	(0.198)	(0.144)	(0.224)	(0.146)	(0.133)
Trait self-control	-0.016	0.002	-0.016	-0.000	-0.007
	(0.016)	(0.010)	(0.015)	(0.010)	(0.009)
Constant	2.357***	1.512***	-0.322	2.158***	1.437***
	(0.239)	(0.144)	(0.265)	(0.161)	(0.152)
Observations	303	289	295	304	1191
$R^2$	0.057	0.008	0.039	0.043	0.024
Treatment × (1 + Female)	-0.221	0.023	0.163	0.004	-0.014
$p_F[Treatment \times (1 + Female) = 0]$	0.327	0.008	0.192	0.000	0.003

Table 4. There are various ways to highlight cells, columns, and rows in a table.

	(1)	(2)	(3)	(4)	(5)
Treatment	-0.390	-0.228	-0.729*	-0.449*	-0.453**
	(+0.352)	(-0.205)	[+0.377]	[-0.245]	{+0.204}
Female	0.948***	0.061	0.188	0.305	0.385*
	(0.354)	(0.233)	(0.372)	(0.226)	(0.222)
Female × Treatment	0.169	0.251	0.892*	0.454	0.439
	(0.514)	(0.325)	(0.533)	(0.341)	(0.307)
Final high school grade	-0.101	0.013	0.076	0.117	0.039
	(0.198)	(0.144)	(0.224)	(0.146)	(0.133)
Trait self-control	-0.016	0.002	-0.016	-0.000	-0.007
	(0.016)	(0.010)	(0.015)	(0.010)	(0.009)
Constant	2.357***	1.512***	-0.322	2.158***	1.437**
	(0.239)	(0.144)	(0.265)	(0.161)	(0.152)
Observations	303	289	295	304	1191
$R^2$	0.057	0.008	0.039	0.043	0.024
Treatment × (1 + Female)	-0.221	0.023	0.163	0.004	-0.014
$p_F$ [Treatment $\times$	0.327	0.008	0.192	0.000	0.003
(1 + Female) = 0					

Table 4. There are various ways to highlight cells, columns, and rows in a table.

	(1)	(2)	(3)	(4)	(5)
Treatment	-0.390	-0.228	-0.729*	-0.449*	-0.453**
	(+0.352)	(-0.205)	[+0.377]	[-0.245]	{+0.204}
Female	0.948***	0.061	0.188	0.305	0.385*
	(0.354)	(0.233)	(0.372)	(0.226)	(0.222)
Female × Treatment	0.169	0.251	0.892*	0.454	0.439
	(0.514)	(0.325)	(0.533)	(0.341)	(0.307)
Final high school grade	-0.101	0.013	0.076	0.117	0.039
	(0.198)	(0.144)	(0.224)	(0.146)	(0.133)
Trait self-control	-0.016	0.002	-0.016	-0.000	-0.007
	(0.016)	(0.010)	(0.015)	(0.010)	(0.009)
Constant	2.357***	1.512***	-0.322	2.158***	1.437**
	(0.239)	(0.144)	(0.265)	(0.161)	(0.152)
Observations	303	289	295	304	1191
$R^2$	0.057	0.008	0.039	0.043	0.024
Treatment × (1 + Female)	-0.221	0.023	0.163	0.004	-0.014
$p_F[Treatment \times (1 + Female) = 0]$	0.327	0.008	0.192	0.000	0.003

Table 4. There are various ways to highlight cells, columns, and rows in a table.

	(1)	(2)	(3)	(4)	(5)
Treatment	-0.390	-0.228	-0.729*	-0.449*	-0.453**
	(+0.352)	(-0.205)	[+0.377]	[-0.245]	{+0.204}
Female	0.948***	0.061	0.188	0.305	0.385*
	(0.354)	(0.233)	(0.372)	(0.226)	(0.222)
Female × Treatment	0.169	0.251	0.892*	0.454	0.439
	(0.514)	(0.325)	(0.533)	(0.341)	(0.307)
Final high school grade	-0.101	0.013	0.076	0.117	0.039
	(0.198)	(0.144)	(0.224)	(0.146)	(0.133)
Trait self-control	-0.016	0.002	-0.016	-0.000	-0.007
	(0.016)	(0.010)	(0.015)	(0.010)	(0.009)
Constant	2.357***	1.512***	-0.322	2.158***	1.437**
	(0.239)	(0.144)	(0.265)	(0.161)	(0.152)
Observations	303	289	295	304	1191
R <sup>2</sup>	0.057	0.008	0.039	0.043	0.024
Treatment × (1 + Female)	-0.221	0.023	0.163	0.004	-0.014
$p_F[\text{Treatment} \times (1 + \text{Female}) = 0]$	0.327	0.008	0.192	0.000	0.003

## Results 7: Yet Another siunitx Example Table

Table 5. Figure grouping via siunitx in a table.

(1)	(2)	(3)
-0.100*	-0.100 01*	-123456.444***
(2.871)	(2.871 23)	[+50000.123]

• The latex exhibits a neutral, acid, or alkaline reaction, depending on the plant from which it was obtained.

- The latex exhibits a neutral, acid, or alkaline reaction, depending on the plant from which it was obtained.
- The latex is therefore usually allowed to coagulate on the tree (Kőszegi and Szeidl, 2013).
  - ⇒ The latex, which is usually coagulated by standing or by heating, is obtained from incisions.

- The latex exhibits a neutral, acid, or alkaline reaction, depending on the plant from which it was obtained.
- The latex is therefore usually allowed to coagulate on the tree (Kőszegi and Szeidl, 2013).
  - ⇒ The latex, which is usually coagulated by standing or by heating, is obtained from incisions.
- See also Dohmen et al. (2012) and Bordalo, Gennaioli, and Shleifer (2013).

#### **Discussion 2: Conclusion**

A paragraph before a list.

- When exposed to air, the latex gradually undergoes putrefactive changes accompanied by coagulation.
  - A paragraph within a list item.
- The addition of a small quantity of ammonia or of formalin to some latices has the effect of preserving them.
- There is, however, reason to believe the following.
- The coagulation of latex into rubber is not mainly of this character.

A paragraph after a list.

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview,
   Skim, and SumatraPDF.



Figure 2. Step 1—Angle: 30.0°

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview, Skim, and SumatraPDF.



Figure 2. Step 2—Angle: 60.0°

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview, Skim, and SumatraPDF.



Figure 2. Step 3—Angle: 90.0°

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview, Skim, and SumatraPDF.



Figure 2. Step 4—Angle: 120.0°

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview, Skim, and SumatraPDF.



Figure 2. Step 5—Angle: 150.0°

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview, Skim, and SumatraPDF.



Figure 2. Step 6—Angle: 180.0°

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview, Skim, and SumatraPDF.



Figure 2. Step 7—Angle: 210.0°

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview, Skim, and SumatraPDF.



Figure 2. Step 8—Angle: 240.0°

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview, Skim, and SumatraPDF.



Figure 2. Step 9—Angle: 270.0°

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview, Skim, and SumatraPDF.



Figure 2. Step 10—Angle: 300.0°

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview, Skim, and SumatraPDF.



Figure 2. Step 11—Angle: 330.0°

- The feature is available in Adobe Acrobat and Acrobat Reader.
- Unfortunately, it is (currently, December 17, 2023) not available in macOS Preview, Skim, and SumatraPDF.



Figure 2. Step 12—Angle: 360.0°



## Discussion 4: Testing the allowframebreaks option

Let's test automatic numbering with the allowframebreaks option.

On this slide, **no** number should be included in the frame title.

## **Discussion 5: Testing the allowframebreaks Option** (1/3)

Let's test automatic numbering with the allowframebreaks option.

On this slide, "(1/3)" should appear in the frame title.

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

## **Discussion 6: Testing the allowframebreaks Option** (2/3)

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

## **Discussion 7: Testing the allowframebreaks Option** (3/3)

Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetuer adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

Math "Torture" Test

## Math "Torture" Test (1/13)

 $\alpha,\beta,\gamma,\delta,\epsilon,\varepsilon,\zeta,\eta,\theta,\vartheta,\iota,\kappa,\lambda,\mu,\nu,\xi,o,\pi,\varpi,\rho,\varrho,\sigma,\varsigma,\tau,\upsilon,\phi,\varphi,\chi,\psi,\omega,\\ \varepsilon,A,B,\Gamma,\Delta,E,Z,H,\Theta,I,K,\Lambda,M,N,\Xi,O,\Pi,P,\Sigma,T,\Upsilon,\Phi,X,\Psi,\Omega,F,-\\ g,\gamma,\Gamma,g,\gamma,\Gamma,g,\gamma,\Gamma,g,\gamma,`$ 

 $\alpha, \beta, \gamma, \delta, \epsilon, \varepsilon, \zeta, \eta, \theta, \vartheta, \iota, \kappa, \lambda, \mu, \nu, \xi, o, \pi, \varpi, \rho, \varrho, \sigma, \zeta, \tau, \upsilon, \phi, \varphi, \chi, \psi, \omega, \varepsilon, A, B, \Gamma, \Delta, E, Z, H, \Theta, I, K, A, M, N, \Xi, O, \Pi, P, \Sigma, T, \Upsilon, \Phi, X, \Psi, \Omega, F, — g, \gamma, \Gamma, g, \gamma, \Gamma, g, \gamma, \Gamma, g, \gamma, \cdot$ 

## Math "Torture" Test (2/13)

- α, β, γ, δ, ε, ε, ζ, η, θ, θ, ι, κ, λ, μ, ν, ξ, ο, π, π, ρ, ρ, σ, ς, τ, υ, φ, φ, χ, ψ, ω, ϝ, Α, Β, Γ, Δ, Ε, Ζ, Η, Θ, Ι, Κ, Λ, Μ, Ν, Ξ, Ο, Π, Ρ, Σ, Τ, Υ, Φ, Χ, Ψ, Ω, Ϝ, <math>- g, γ, Γ, g, γ, Γ, g, γ, Γ, g, γ, Γ, g, γ, Γ
- $\alpha,\,\beta,\,\gamma,\,\delta,\,\epsilon,\,\epsilon,\,\zeta,\,\eta,\,\theta,\,\theta,\,\iota,\,\kappa,\,\lambda,\,\mu,\,\nu,\,\xi,\,o,\,\pi,\,\pi,\,\rho,\,\rho,\,\sigma,\,\varsigma,\,\tau,\,\upsilon,\,\phi,\,\phi,\,\chi,\,\psi,\,\omega,\,\varsigma,\,A,\,B,\,\Gamma,\,\Delta,\,E,\,Z,\,H,\,\theta,\,I,\,K,\,\Lambda,\,M,\,N,\,\Xi,\,O,\,\Pi,\,P,\,\Sigma,\,T,\,Y,\,\Phi,\,X,\,\Psi,\,\Omega,\,F,\,-\,g,\,\gamma,\,\Gamma,\,g,\,\gamma,\,\Gamma,\,g,\,\gamma,\,\Gamma,\,g,\,\gamma,\,\Gamma$
- α, β, γ, δ, ε, ε, ζ, η, θ, θ, ι, κ, λ, μ, ν, ξ, ο, π, π, ρ, ρ, σ, ς, τ, υ, φ, φ, χ, ψ, ω, ϝ, Α, Β, Γ, Δ, Ε, Ζ, Η, Θ, Ι, Κ, Λ, Μ, Ν, Ξ, Ο, Π, Ρ, Σ, Τ, Υ, Φ, Χ, Ψ, Ω, Ϝ, g, γ, Γ, g, γ, Γ, g, γ, Γ, g, γ, Γ

## Math "Torture" Test (3/13)

Most of the following examples are taken from *The T<sub>E</sub>Xbook* (Knuth, 1984, see https://ctan.org/pkg/texbook) and were adapted for 上X from Karl Berry's torture test for plain T<sub>E</sub>X math fonts.

$$x+y-z$$
,  $x+y*z$ ,  $z*y/z$ ,  $(x+y)(x-y)=x^2-y^2$ ,  $x\times y\cdot z=[x\,y\,z]$ ,  $x\circ y\bullet z$ ,  $x\cup y\cap z$ ,  $x\sqcup y\sqcap z$ ,  $x\vee y\wedge z$ ,  $x\pm y\mp z$ ,  $x=y/z$ ,  $x:=y$ ,  $x\leq y\neq z$ ,  $x\sim y\simeq z$   $x\equiv y\not\equiv z$ ,  $x\subset y\subseteq z$   $\sin 2\theta=2\sin\theta\cos\theta$ ,  $O(n\log n\log n)$ ,  $\Pr(X>x)=\exp(-x/\mu)$ ,  $(x\in A(n)\mid x\in B(n))$ ,  $\bigcup_n X_n\mid \bigcap_n Y_n$  In-text matrices  $\binom{11}{01}$  and  $\binom{a\ b\ c}{1\ m\ n}$ .

## Math "Torture" Test (4/13)

Disambiguation: 0 O O, 1 I I | lI/, ij, rnm,  $\theta \Theta$ ,  $\phi \psi$ , --

Latin vs. Greek:  $a \alpha, d \delta, e \epsilon, i \iota, k \kappa, n \eta, o \sigma, p \rho, \beta \beta, u \upsilon, v v, w \omega, x \chi, y \gamma, A \Delta \Lambda, O \Theta \Omega, T \Gamma, Y \Upsilon.$ 

Disambiguation: 0 O O, 1 | I | I |, i j, rn m,  $\theta \Theta$ ,  $\phi \psi$ , --

Latin vs. Greek:  $\alpha \alpha$ ,  $d \delta$ ,  $e \varepsilon$ ,  $i \iota$ ,  $k \kappa$ ,  $n \eta$ ,  $o \sigma$ ,  $p \rho$ ,  $\beta \beta$ ,  $u \upsilon$ , v v,  $w \omega$ ,  $x \chi$ ,  $y \gamma$ ,  $A \Delta \Lambda$ ,  $O \Theta \Omega$ ,  $T \Gamma$ , Y Y.

## Math "Torture" Test (5/13)

$$a_{0} + \frac{1}{a_{1} + \frac{1}{a_{2} + \frac{1}{a_{3} + \frac{1}{a_{4}}}}}$$

$$\binom{p}{2}x^{2}y^{p-2} - \frac{1}{1-x}\frac{1}{1-x^{2}} = \frac{a+1}{b} / \frac{c+1}{d}.$$

$$\sqrt{1 + \sqrt{1 + \sqrt{1 + \sqrt{1 + \sqrt{1 + x}}}}}$$

$$\sqrt{1 + \sqrt[k]{1 + \sqrt[k]{1 + \sqrt[k]{1 + \sqrt[k]{1 + x}}}}}$$

## Math "Torture" Test (6/13)

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right) \left| \phi(x + iy) \right|^2 = 0$$

$$\pi(n) = \sum_{m=2}^n \left[ \left( \sum_{k=1}^{m-1} \lfloor (m/k) / \lceil m/k \rceil \rfloor \right)^{-1} \right].$$

$$\int_0^\infty \frac{t - ib}{t^2 + b^2} e^{iat} dt = e^{ab} E_1(ab), \quad a, b > 0.$$

$$\mathbf{A} := \begin{pmatrix} x - \lambda & 1 & 0 \\ 0 & x - \lambda & 1 \\ 0 & 0 & x - \lambda \end{pmatrix}.$$

## Math "Torture" Test (7/13)

$$\mathbf{A} = \begin{pmatrix} a & b & c \\ d & e & f \end{pmatrix} \begin{pmatrix} u & x \\ v & y \\ w & z \end{pmatrix}$$

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$$

$$\mathbf{C} \qquad I \qquad C'$$

$$\mathbf{M} = \begin{matrix} C & I & C' \\ C & 1 & 0 & 0 \\ b & 1 - b & 0 \\ C' & 0 & a & 1 - a \end{matrix}$$

## Math "Torture" Test (8/13)

$$\sum_{n=0}^{\infty} a_n z^n \quad \text{converges if} \quad |z| < \left( \limsup_{n \to \infty} \sqrt[n]{|a_n|} \right)^{-1}.$$

$$\frac{f(x + \Delta x) - f(x)}{\Delta x} \to f'(x) \quad \text{as } \Delta x \to 0.$$

$$||u_i|| = 1, \quad u_i \cdot u_j = 0 \quad \text{if } i \neq j.$$

The confluent image of  $\left\{ \begin{array}{l} \text{an arc} \\ \text{a circle} \\ \text{a fan} \end{array} \right\} \quad \text{is} \quad \left\{ \begin{array}{l} \text{an arc} \\ \text{an arc or a circle} \\ \text{a fan or an arc} \end{array} \right\}.$ 

## Math "Torture" Test (9/13)

$$\sum_{n=0}^{\infty} a_n z^n \quad \text{converges if} \quad |z| < \left( \limsup_{n \to \infty} \sqrt[n]{|a_n|} \right)^{-1}.$$

$$\frac{f(x + \Delta x) - f(x)}{\Delta x} \to f'(x) \quad \text{as } \Delta x \to 0.$$

$$||u_i|| = 1, \quad u_i \cdot u_j = 0 \quad \text{if } i \neq j.$$

The confluent image of  $\left\{ \begin{array}{l} \text{an arc} \\ \text{a circle} \\ \text{a fan} \end{array} \right\} \quad \left\{ \begin{array}{l} \text{an arc} \\ \text{an arc or a circle} \\ \text{a fan or an arc} \end{array} \right\}.$ 

## Math "Torture" Test (10/13)

$$T(n) \le T(2^{\lceil \lg n \rceil}) \le c(3^{\lceil \lg n \rceil} - 2^{\lceil \lg n \rceil})$$

$$< 3c \cdot 3^{\lg n}$$

$$= 3c n^{\lg 3}.$$

$$(x+y)(x-y) = x^2 - xy + yx - y^2$$

$$= x^2 - y^2$$

$$(x+y)^2 = x^2 + 2xy + y^2.$$

## Math "Torture" Test (11/13)

$$\left(\int_{-\infty}^{\infty} e^{-x^2} dx\right)^2 = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)} dx dy$$

$$= \int_{0}^{2\pi} \int_{0}^{\infty} e^{-r^2} dr d\theta$$

$$= \int_{0}^{2\pi} \left(e^{-\frac{r^2}{2}}\Big|_{r=0}^{r=\infty}\right) d\theta$$

$$= \pi.$$

## Math "Torture" Test (12/13)

$$\prod_{k\geq 0} \frac{1}{(1-q^k z)} = \sum_{n\geq 0} z^n / \prod_{1\leq k\leq n} (1-q^k).$$

$$\sum_{\substack{0 < i \le m \\ 0 < j \le n}} p(i,j) \neq \sum_{i=1}^p \sum_{j=1}^q \sum_{k=1}^r a_{ij} b_{jk} c_{ki} \neq \sum_{\substack{1 \le i \le p \\ 1 \le j \le q \\ 1 < k < r}} a_{ij} b_{jk} c_{ki}$$

## Math "Torture" Test (13/13)

$$\max_{1 \le n \le m} \log_2 P_n \quad \text{and} \quad \lim_{x \to 0} \frac{\sin x}{x} = 1$$

Inline math:  $\max_{1 \le n \le m} \log_2 P_n$  and  $\lim_{x \to 0} \frac{\sin x}{x} = 1$ 

$$p_1(n) = \lim_{m \to \infty} \sum_{v=0}^{\infty} \left( 1 - \cos^{2m} (v!^n \pi/n) \right)$$

Inline math:  $p_1(n) = \lim_{m \to \infty} \sum_{v=0}^{\infty} (1 - \cos^{2m}(v!^n \pi/n))$ 

## References

#### References

- Bordalo, Pedro, Nicola Gennaioli, and Andrei Shleifer. 2013. "Salience and Consumer Choice." Journal of Political Economy 121 (5): 803–43. https://doi.org/10.1086/673885. [PDF pp. 43–45]
- Dertwinkel-Kalt, Markus, Holger Gerhardt, Gerhard Riener, Frederik Schwerter, and Louis Strang. 2017. "Concentration Bias in Intertemporal Choice." Working paper. Bonn, Germany, et al.: University of Bonn et al. https://www.dropbox.com/s/dv20mcu0gkygmiz/Concentration Bias in Intertemporal Choice.pdf. [PDF pp. 27, 28]
- Dohmen, Thomas, Armin Falk, David Huffman, and Uwe Sunde. 2012. "Interpreting Time Horizon Effects in Inter-Temporal Choice."
  IZA Discussion Paper, IZA Discussion Paper Series 6385. Maastricht University et al. http://ftp.iza.org/dp6385.pdf. [PDF pp. 5, 43–45]
- Gerhardt, Holger, Hannah Schildberg-Hörisch, and Jana Willrodt. 2017. "Does self-control depletion affect risk attitudes?" European Economic Review 100: 463–87. https://doi.org/10.1016/j.euroecorev.2017.09.004. [PDF pp. 22, 29]
- Knuth, Donald E. 1984. The TeXbook. 483. Reading, MA, USA: Addison-Wesley. [PDF p. 66]
- Kőszegi, Botond, and Adam Szeidl. 2013. "A Model of Focusing in Economic Choice." Quarterly Journal of Economics 128 (1): 53–104. https://doi.org/10.1093/qje/qjs049. [PDF pp. 5, 43–45, 80]
- Lisi, A. Garrett. 1995. "A solitary wave solution of the Maxwell-Dirac equations." Journal of Physics A: Mathematical and General 28 (18): 5385-92. https://doi.org/10.1088/0305-4470/28/18/026. arXiv: hep-th/9410244. [PDF p. 5]

# Appendix

## **Appendix: Modeling Concentration Bias**

Subjects consider a sequences of consequences c from choice set C.

Standard discounted utility: Suppose that the instantaneous utility function u satisfies u' > 0 and u" ≤ 0, and that earlier consequences are preferred over later consequences of the same magnitude, i.e., D(t) ≤ 1:

$$U(\mathbf{c}) := \sum_{t=1}^{T} D(t) u(c_t)$$
, where, e.g.,  $D(t) = \delta^t$  or  $D(t) = \frac{1}{1+kt}$ .

Focusing model (Kőszegi and Szeidl, 2013):

$$\begin{split} \tilde{U}(\boldsymbol{c},C) &:= \sum_{t=1}^{T} g_t \, D(t) \, u(c_t), \quad \text{where} \\ g_t &\equiv g[\max_{c' \in C} D(t) \, u(c'_t) - \min_{c' \in C} D(t) \, u(c'_t)] \end{split}$$

- Weighting function  $g[\cdot]$  increases in difference of maximum and minimum possible utility at a point in time.
- Subjects overweight intertemporal consequences with a greater range.