


Theories of Generalized Change

Joseph J. Babčanec



Theories of Generalized Change

Joseph J. Babčanec

First Edition

© 2025

*To my wife Emily and my sons Andy and Jamie
To my former math teacher Heather Elfen*

Theories of Generalized Change
First Edition

Copyright © 2025 by Joseph J. Babčanec

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means—electronic, mechanical, photocopying, recording, or otherwise—without the prior written permission of the author, except in the case of brief quotations used in reviews or scholarly analysis.

Some illustrations in this book are original works by the author; others are sourced from public domain or free-to-use archives, and are credited where appropriate.

ISBN 978-1-23456-789-0

Cover design and interior layout by Joseph J. Babčanec

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1
First Printing, 2025

Set in L^AT_EX using the Computer Modern typeface.

Contents

Introduction	iv
1 Single-Variable Calculus, Vectors, and the Geometry of Space	1
Lecture 1.1: Review of Differential and Integral Calculus	2
Lecture 1.2: Three-Dimensional Coordinate Systems	25
Lecture 1.3: Vectors	38
Lecture 1.4: Lines and Planes in Space	59
Lecture 1.5: Quadric Surfaces	80
Formulas and Definitions	92
Practice Problems	94
2 Vector-Valued Functions and Motion in Space	98
Lecture 2.1: Vector-Valued Functions	99
Lecture 2.2: Motion Along a Curve	110
Lecture 2.3: The TNB Frame, Curvature and Torsion	120
Lecture 2.4: Arc Length and Reparameterization	133
Formulas and Definitions	145
Practice Problems	147
3 Partial Derivatives	151
Lecture 3.1: Functions of Several Variables	152
Lecture 3.2: Limits and Continuity	169
Lecture 3.3: Partial Derivatives	184
Lecture 3.4: The Chain Rule	194
Lecture 3.5: Directional Derivatives and Gradient Vectors	203
Lecture 3.6: Tangent Planes and Linear Approximations	217
Lecture 3.7: Extrema and Critical Points	226
Lecture 3.8: Optimization	248
Formulas and Definitions	268
Practice Problems	271
4 Multiple Integrals	275
Lecture 4.1: Double Integrals over Rectangular Regions	276
Lecture 4.2: Double Integrals over General Regions	290
Lecture 4.3: Applications of Double Integrals	310
Lecture 4.4: Triple Integrals	322

Lecture 4.5: Change of Variables in Multiple Integrals	336
Lecture 4.6: Applications of Triple Integrals	346
Formulas and Definitions	359
Practice Problems	362
5 Vector Calculus	370
Lecture 5.1: Vector Fields	371
Lecture 5.2: Line Integrals	387
Lecture 5.3: Green's Theorem	406
Lecture 5.4: Curl and Divergence	417
Lecture 5.5: Surface Integrals	434
Lecture 5.6: Stokes' Theorem	450
Lecture 5.7: Divergence Theorem	468
Formulas and Definitions	480
Practice Problems	482
6 Differential Equations	490
Lecture 6.1: Introduction to Differential Equations	491
Lecture 6.2: First-Order Ordinary Differential Equations	502
Lecture 6.3: Higher-Order Ordinary Differential Equations I	515
Lecture 6.4: Higher-Order Ordinary Differential Equations II	528
Lecture 6.5: Methods of Solving Partial Differential Equations I	539
Lecture 6.6: Methods of Solving Partial Differential Equations II	553
Formulas and Definitions	569
Practice Problems	572
7 Complex Variables	577
Lecture 7.1: The Complex Number System	578
Lecture 7.2: Complex Functions	594
Lecture 7.3: Basic Limits in Complex Analysis	611
Lecture 7.4: Complex Differentiation	623
Lecture 7.5: Elementary Complex Integration	637
Lecture 7.6: Cauchy's Integral Theorem and Residues	651
Formulas and Definitions	672
Practice Problems	675
Appendix A: Derivative and Integral Identities	683
Appendix B: Vector Properties and Formulas	685
Appendix C: Calculus Theorems	687
Appendix D: Quick Reference for First-Order ODEs	689
Appendix E: Useful Tables for Differential Equations	690

Appendix F: Complex Analysis Identities	692
References	694
Index	696

Preface

I wrote this textbook primarily to provide students of Multivariate Calculus (also known as Vector Calculus or Calculus III) with the option of having a concise reference to assist them or a springboard into deeper mathematical topics for the sake of interest. When I was a freshman engineering major in college, I reached this crossroads and, in a way, chose both paths—I added a mathematics major, thereby embracing both perspectives. These notes, compiled into textbook form and titled “Theories of Generalized Change,” are designed to bridge theory with application, emphasizing problem-solving techniques and the essential skills needed in applied fields.

This course, like most others, requires a solid foundation in Calculus I and Calculus II, including an understanding of limits, derivatives, integrals, and series. Reviewing the concepts from Calculus I will be particularly helpful before diving in, and knowledge from Calculus II will provide additional clarity and depth. While familiarity with introductory multivariable concepts is beneficial, we will introduce and review key topics as needed to ensure all students have the necessary tools to succeed. This course will feel like we are revisiting basic Differential Calculus, but with a new spirit—one of generalization. We will constantly explore how to further generalize and extend our understanding of Calculus to higher dimensions.

The course topics cover vectors, vector-valued functions, partial derivatives, multiple integrals, vector calculus theorems, introductory topics in Partial Differential Equations (PDEs), and a bit of Complex Variables, each presented with examples relevant to engineering, physics, and other applied sciences. PDEs and, less frequently, Complex Variables are seldom discussed in such a course. However, they are intrinsically tied together and embody the spirit of generalizing calculus; not to mention, they are quite useful in their own right. Through these applications, students will develop an appreciation for the power of calculus in modeling and solving practical problems, as well as gain a taste of higher mathematics.

Each chapter is divided into lectures, roughly structured to fit a 90- to 120-minute session. This format allows flexibility for instructors to adapt the material to the pace and needs of the class. The sections within each lecture focus on specific concepts, promoting a gradual mastery of each topic before progressing to more advanced material. This structure also provides a convenient framework for creating assignments and assessments. In my opinion, Chapters 1–5 are absolutely crucial for a Multivariate Calculus class, while Chapters 6 and 7 are additional topics to include if there is extra time.

One of the most challenging aspects for students of mathematics is handling notation and the over-assumptions made by some texts. Extra care is taken to ensure that notation is clearly defined and labeled—mathematics is like learning a new language. I also make sure to write out problems in full and use examples that cover a broad range of use cases, allowing students to more easily make connections to further topics.

My goal as an educator with this text is to empower students to apply multivariable calculus concepts beyond the classroom, equipping them with a robust mathematical foundation for their future studies and careers. I hope this course serves as a valuable resource for both students and instructors, providing the clarity, depth, and insight needed to master Multivariate Calculus...I also crack jokes often. Enjoy!

Joseph J. Babčanec

Adjunct Professor of Mathematics, Benedict College
Lead Data Scientist, Anika Systems Inc.

joseph.babcanec@benedict.edu.

Introduction

Purpose of This Book

This book is structured to provide comprehensive coverage of Calculus III concepts, with applications relevant to fields such as engineering, physics, and applied sciences. The aim is to build on students' foundation in single-variable calculus, introducing the complexities of multivariable calculus and vector calculus. By focusing on practical and theoretical elements, the book equips students to understand multidimensional space, vector fields, integrals over curves and surfaces, and more advanced mathematical modeling techniques.

This text is designed as a primary resource, integrating lecture material with detailed explanations, examples, and practice problems to foster a strong grasp of the subject.

How to Use This Book

This book should be used as a guide and practice tool alongside lectures. Each section breaks down complex ideas, followed by examples and exercises. Students should:

- **Preview Sections:** Familiarize yourself with topics before lectures for easier engagement.
- **Engage During Lectures:** Participate in discussions to solidify your understanding.
- **Work Through Examples:** Each example is chosen to illustrate key techniques.
- **Attempt Problems Independently:** Practice problems build problem-solving skills.
- **Use as Reference:** Refer back to specific sections when tackling homework or exams.

Overview of Chapters

This book is organized into nine chapters, progressively covering essential topics in multivariable calculus and applications:

1. **Vectors and the Geometry of Space:** Covers 3D coordinate systems, vector operations, lines, planes, and quadric surfaces, including applications in engineering.

2. **Vector-Valued Functions and Motion in Space:** Explores vector functions, derivatives, motion along curves, curvature, arc length, and engineering applications.
3. **Partial Derivatives:** Examines functions of several variables, limits, partial derivatives, chain rule, tangent planes, and optimization, crucial for multidimensional analysis.
4. **Multiple Integrals:** Discusses double and triple integrals in various coordinates, with applications in area, volume, mass, and center of mass.
5. **Vector Calculus:** Focuses on vector fields, line and surface integrals, and major theorems like Green's, Stokes', and the Divergence Theorem.
6. **Differential Equations:** Introduces ODEs and PDEs with applications.
7. **Complex Variables:** Covers basic complex analysis, including complex functions, integration, and applications in physics.

Prerequisites

This course requires a solid foundation in Calculus I and II. Students should be comfortable with:

- **Differentiation and Integration:** Mastery of techniques for single-variable calculus.
- **Series and Sequences:** Familiarity with Taylor and power series.
- **Basic Linear Algebra:** Matrix operations and systems of linear equations is very helpful.
- **Mathematical Reasoning:** Ability to follow and construct rigorous proofs will ensure you get the very most out of this book.

Acknowledgments

Special thanks to my colleagues and the mathematics department for their support and input. I also must of course mention my family who puts up with me when I write such textbooks (this is my third), and go bat-shit crazy for a few months.

Basic Terminology

Below is an extensive list of terms essential to Calculus III and related subject matter to be covered in this text.

3D Coordinate System A system for describing positions in three-dimensional space using three coordinates (e.g., (x, y, z)) relative to mutually perpendicular axes.

Cartesian Coordinates A coordinate system in which points are described by their distances along perpendicular axes, typically labeled x , y , and z .

Cylindrical Coordinates A three-dimensional coordinate system using (r, θ, z) , where r is the radius in the xy -plane, θ is the angle around the z -axis, and z is the vertical coordinate.

Spherical Coordinates A three-dimensional coordinate system using (ρ, θ, ϕ) , where ρ is the distance from the origin, θ is the angle in the xy -plane, and ϕ is the angle from the positive z -axis.

Vectors Quantities with both magnitude and direction, often represented as $\langle x, y, z \rangle$ in three dimensions.

Magnitude (Norm) of a Vector The length of a vector, computed as $\sqrt{x^2 + y^2 + z^2}$ in 3D.

Unit Vector A vector of length 1 in a particular direction.

Dot Product An operation that takes two vectors and produces a scalar, measuring the extent to which they point in the same or opposite directions.

Cross Product An operation that takes two vectors in 3D and produces a vector perpendicular to both, with magnitude related to the area of the parallelogram formed by the two vectors.

Vector-Valued Functions Functions whose outputs are vectors, commonly written as $\mathbf{r}(t) = \langle x(t), y(t), z(t) \rangle$.

Parametric Equations of Curves Expressions for x , y , and z in terms of a parameter t , used to describe curves in three dimensions.

Tangent Vector The derivative of a vector-valued function, indicating the instantaneous direction of the curve.

Arc Length The length of a curve over an interval, computed by integrating the magnitude of the derivative of the vector-valued function.

Curvature A measure of how sharply a curve bends at a given point.

T, N, B (Frenet-Serret Frame) The unit tangent (**T**), the principal normal (**N**), and the binormal (**B**) vectors associated with a curve in space.

Level Surfaces Sets of points in 3D where a function of three variables is constant.

Partial Derivatives Derivatives of a function of several variables with respect to one variable, treating the others as constants.

Gradient A vector of all partial derivatives of a function, pointing in the direction of greatest increase.

Directional Derivative The rate of change of a function in the direction of a specified vector.

Chain Rule (Multivariable) A rule for computing the derivative of a composite function where the variables themselves depend on other parameters.

Jacobian A determinant representing how a function of several variables transforms infinitesimal elements during a change of variables.

Local Extrema (Maxima/Minima) Points where a function takes on locally highest or lowest values, found using partial derivatives and tests involving second derivatives.

Saddle Point A point where a function is neither a local maximum nor a local minimum, but partial derivatives are zero.

Multiple Integrals Integrals taken over regions in higher dimensions (e.g., double integrals over areas or triple integrals over volumes).

Double Integral An integral over a two-dimensional region, typically written as $\iint_D f(x, y) dA$.

Triple Integral An integral over a three-dimensional region, typically written as $\iiint_E f(x, y, z) dV$.

Iterated Integrals The process of evaluating a multiple integral by integrating one variable at a time.

Line Integrals Integrals that sum a function along a curve, with either scalar or vector integrands.

Surface Integrals Integrals that sum a function over a surface in 3D, used for scalar fields or vector fields.

Vector Fields Functions that assign a vector to each point in space, often denoted $\mathbf{F}(x, y, z)$.

Divergence A scalar measure of how much a vector field spreads out or converges at a given point.

Curl A vector measure of the rotation or swirling strength of a vector field at a point.

Laplacian The sum of second partial derivatives of a function (in 3D, $\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}$).

Green's Theorem Relates a line integral around a simple closed curve in the plane to a double integral over the region it encloses.

Stokes' Theorem Relates a surface integral of the curl of a vector field to a line integral of the field around the boundary of the surface.

Divergence Theorem (Gauss's Theorem) Relates the flux of a vector field through a closed surface to the volume integral of the divergence of the field over the region it encloses.

Ordinary Differential Equations (ODEs) Equations involving derivatives of functions of one variable.

Order of an ODE The highest derivative present in the equation (e.g., first-order, second-order).

Linear ODE An ODE where the function and its derivatives appear to the first power and there are no products of the function and/or its derivatives.

Nonlinear ODE An ODE that does not meet the criteria for linearity; may include products of the function and its derivatives or other nonlinear terms.

Initial Value Problem (IVP) An ODE with specified initial conditions at a particular point.

Partial Differential Equations (PDEs) Equations involving partial derivatives of functions of multiple variables (e.g., the heat equation or wave equation).

Boundary Conditions Constraints specifying the values (or behavior) of a function along boundaries of a domain for PDEs.

Complex Numbers Numbers of the form $z = x + iy$, where i is the imaginary unit and x, y are real.

Complex Plane A two-dimensional plane representing the real part on the horizontal axis and the imaginary part on the vertical axis.

Polar Form of a Complex Number Writing a complex number as $re^{i\theta}$, where r is the magnitude and θ is the argument (angle).

Complex Function A function with complex input and output, often written $f(z)$.

Holomorphic (Analytic) Function A complex function that is differentiable everywhere in its domain, satisfying the Cauchy-Riemann equations.

Contour A path or curve in the complex plane along which a complex integral can be computed.

Residue A coefficient representing the behavior of a complex function near its singularity, used for evaluating contour integrals via the residue theorem.

Contour Integral An integral of a complex function along a specified path in the complex plane.

Euler's Formula A fundamental relationship $e^{i\theta} = \cos(\theta) + i\sin(\theta)$, connecting complex exponentials with trigonometric functions.

Mathematical Symbols and Notation

A comprehensive list of symbols used throughout Calculus III:

$\vec{r}$, $\mathbf{r}$ Position vector.

$|\vec{v}|$, $|\mathbf{v}|$, $\|\mathbf{v}\|$, $\|\mathbf{v}\|_2$ Magnitude of vector $\vec{v}$.

Δ Finite Difference.

$\int$ Integral sign.

$\sum$ Summation.

$\prod$ Product.

∇ Del operator (used in gradient, divergence, curl).

∂ Partial derivative symbol. Also used to indicate boundary, ie, ∂C

dA , dV Area and volume elements.

ρ , θ , ϕ Spherical coordinates.

$\vec{F}$, $\mathbf{F}$ Vector field.

$\int_C$ Line integral.

$\iint_S$ Surface integral.

$\iiint_V$ Triple integral.

$\oint_C$ Closed Contour Line integral.

$\oiint_S$ Closed Contour Surface integral.

O Big O notation.

$\approx$ Approximately equal.

$\equiv$ Identically equal.

$\perp$ Perpendicular.

$\parallel$ Parallel.

$\Rightarrow$ Implies.

This book aims to provide a comprehensive understanding of these and other core concepts, supporting students in developing the mathematical maturity required for advanced studies and professional applications.

Chapter 1

Single-Variable Calculus, Vectors, and the Geometry of Space

Introduction

In this chapter, we begin our journey into multivariable calculus by revisiting core concepts from Calculus I and Calculus II. These reviews will serve as stepping stones as we enter into the world of “high-dimensional thinking.” Our goal is to build a solid foundation for understanding functions that involve several variables, expanding beyond the single-variable functions studied previously. Functions of multiple variables naturally arise in physics and engineering. This chapter will introduce functions like these that depend on multiple variables, setting the stage for exploring complex systems where outcomes are influenced by more than one parameter.

To engage with these new concepts, we will introduce vectors, essential tools for representing quantities with both magnitude and direction. Vectors are critical for understanding spatial relationships and navigating the geometry of three-dimensional space. We will cover the fundamental operations on vectors, including addition, scalar multiplication, dot product, and cross product, each with significant applications in describing physical forces, directions, and motions. Furthermore, this chapter will introduce the geometry of space, specifically three-dimensional coordinate systems. By exploring Cartesian, cylindrical, and spherical coordinates, we will establish methods to describe points, lines, planes, and surfaces in space. These systems provide multiple ways of visualizing and solving problems, especially in fields like engineering and physics, where spatial reasoning and high-dimensional analysis are vital.

In summary, this chapter lays the groundwork for high-dimensional thinking, equipping students with the skills to analyze functions of several variables and understand the fundamental geometry of space. With a thorough review of Calculus I and II concepts, we’ll be ready to explore multivariable calculus, ultimately deepening our ability to model and interpret complex systems.

Lecture 1.1: Review of Differential and Integral Calculus

Review of Single-Variable Calculus

Before we embark on multivariable calculus, it's essential to have a strong command of foundational topics in single-variable calculus, along with proficiency in algebra. Mastery of differentiation, integration, and limits forms the backbone of the more advanced concepts we'll explore in this course. Additionally, skill in handling complex algebraic expressions will be invaluable, as calculus problems frequently require careful algebraic manipulation. These topics should already be second nature:

1. Limits and Continuity:

A precise understanding of limits is fundamental. You should be comfortable calculating limits both algebraically and conceptually, as they underpin differentiation, integration, and continuity. Key ideas include the ε - δ definition of a limit, one-sided limits, and limits approaching infinity.

2. Differentiation:

Differentiation techniques such as the power rule, product rule, quotient rule, and chain rule must be second nature. In multivariate calculus, you will extend these rules to partial derivatives and gradients. Familiarity with applications of derivatives—such as optimization, related rates, and curve sketching—is also critical, as these concepts generalize to higher dimensions.

3. Integration:

Integration techniques such as substitution, integration by parts, and partial fraction decomposition will reappear in multivariable settings. Definite and indefinite integrals, as well as their geometric interpretations (e.g., areas under curves), are fundamental. In multivariate calculus, we'll extend these concepts to double and triple integrals over various regions.

4. Fundamental Theorem of Calculus:

A firm grasp of the Fundamental Theorem of Calculus is crucial. This theorem connects differentiation and integration, and its higher-dimensional analogs—like Green's Theorem, Stokes' Theorem, and the Divergence Theorem—are central topics in multivariate calculus.

5. Algebraic Skills:

Proficiency in manipulating algebraic expressions is indispensable. Be comfortable simplifying complex fractions, solving linear and quadratic equations, and working with exponential and logarithmic functions. Multivariable calculus often involves tedious algebraic steps, and fluency here will prevent errors.

As we review these topics, consider how they might extend into three dimensions and beyond. For example, differentiation in higher dimensions involves calculating partial derivatives, gradients, and directional derivatives. Similarly, integration evolves to include volume and surface calculations. These extensions will require you to adapt and expand the skills you've already mastered in single-variable calculus. **Think of this review not as a simple refresher, but as a foundation to support the complexities of three-dimensional calculus.**

Before we review, let's knock out some elementary examples that should be second nature to you. If they are not easy to you, then it is **critical** that you ensure you are at the level to where they are before you proceed past this lecture.

▲ **Example 1.1.1** Evaluate the derivative of $f(x) = x^2e^x$.

Solution We use the product rule, which states:

$$\frac{d}{dx}f(x) = u(x)v'(x) + v(x)u'(x).$$

Here, $u(x) = x^2$ and $v(x) = e^x$. Compute:

$$\begin{aligned}\frac{d}{dx}(x^2e^x) &= x^2 \cdot \frac{d}{dx}(e^x) + e^x \cdot \frac{d}{dx}(x^2) \\ &= x^2e^x + 2xe^x = e^x(x^2 + 2x).\end{aligned}$$

Thus, the derivative is $f'(x) = e^x(x^2 + 2x)$.

▲ **Example 1.1.2** Find the general solution to $\int \frac{1}{1-x} dx$.

Solution We use substitution to evaluate the integral. Let $u = 1 - x$, so $du = -dx$. Rewrite the integral:

$$\int \frac{1}{1-x} dx = \int \frac{-1}{u} du.$$

The integral of $\frac{1}{u}$ is $\ln |u|$. Substituting back $u = 1 - x$, we obtain:

$$\int \frac{1}{1-x} dx = -\ln |1-x| + C,$$

where C is the constant of integration.

▲ **Example 1.1.3** Solve the system of equations:

$$\begin{aligned} 2x + y &= 5, \\ x - y &= 1. \end{aligned}$$

Solution Add the equations to eliminate y :

$$2x + y + x - y = 5 + 1 \implies 3x = 6 \implies x = 2.$$

Substitute $x = 2$ into $2x + y = 5$:

$$2(2) + y = 5 \implies 4 + y = 5 \implies y = 1.$$

Thus, the solution is $(x, y) = (2, 1)$.

Single-Variable Limits

Limits are fundamental in calculus, allowing us to understand the behavior of functions as they approach specific values. They provide a rigorous framework for defining instantaneous rates of change, which are crucial for differentiation and integration. Limits help us analyze functions near points of interest, including points where the function may not be defined, making them indispensable in fields like physics and engineering. You may not have encountered the formal epsilon-delta definition, but it forms the mathematical foundation for how limits are defined precisely.

- **Definition:** $\lim_{x \rightarrow a} f(x) = L$ means that as x approaches a , $f(x)$ gets arbitrarily close to L .

- **Properties:**

– **Sum Rule:**

$$\lim_{x \rightarrow a} [f(x) + g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x)$$

– **Product Rule:**

$$\lim_{x \rightarrow a} [f(x)g(x)] = \lim_{x \rightarrow a} f(x) \cdot \lim_{x \rightarrow a} g(x)$$

– **Quotient Rule:**

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \frac{\lim_{x \rightarrow a} f(x)}{\lim_{x \rightarrow a} g(x)}, \text{ if } \lim_{x \rightarrow a} g(x) \neq 0$$

- **Single-Sided Limits:** Sometimes we are interested in the behavior of $f(x)$ as x approaches a from only one side, either from the left or the right.

- **Left-Hand Limit:** $\lim_{x \rightarrow a^-} f(x) = L$ means $f(x)$ approaches L as x approaches a from values less than a .
- **Right-Hand Limit:** $\lim_{x \rightarrow a^+} f(x) = L$ means $f(x)$ approaches L as x approaches a from values greater than a .
- For the (two-sided) limit $\lim_{x \rightarrow a} f(x) = L$ to exist, both the left-hand limit and right-hand limit must exist and be equal.
- **Limits at Infinity and Infinite Limits:** In addition to limits at finite values, we often need to consider what happens to $f(x)$ as x grows arbitrarily large or small, or as $f(x)$ approaches infinity.
 - **Limit at Infinity:** $\lim_{x \rightarrow \infty} f(x) = L$ means that as x becomes very large, $f(x)$ approaches L .
 - **Infinite Limit:** $\lim_{x \rightarrow a} f(x) = \infty$ means that as x approaches a , $f(x)$ grows without bound.
- **Formal Definition of a Limit (Epsilon-Delta Definition):** To define limits rigorously, we use the epsilon-delta definition, which formalizes the concept as follows: We say that $\lim_{x \rightarrow a} f(x) = L$ if for every $\epsilon > 0$, there exists a $\delta > 0$ such that:

$$0 < |x - a| < \delta \implies |f(x) - L| < \epsilon$$

This definition means we can make $f(x)$ as close to L as desired (within ϵ) by taking x sufficiently close to a (within δ), without x being exactly a .




Figure 1.1: A limit being approached notated with $\epsilon - \delta$ notation.

- **L'Hôpital's Rule:** When calculating limits, we sometimes encounter indeterminate forms, such as $\frac{0}{0}$ or $\frac{\infty}{\infty}$. L'Hôpital's Rule is a powerful tool for handling these cases.

If $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x) = 0$ or $\pm\infty$, and both f and g are differentiable near a (except possibly at a), then:

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

Provided the limit on the right exists or is infinite. L'Hôpital's Rule can also be applied iteratively. If an indeterminate form persists after applying the rule once, we may apply the rule again to the resulting expression until the limit can be evaluated.

- **Interpreting the Epsilon-Delta Definition:**

- ϵ represents the desired closeness of $f(x)$ to L .
 - δ is how close x needs to be to a to ensure $f(x)$ is within ϵ of L .
 - The limit $\lim_{x \rightarrow a} f(x) = L$ exists if, for any chosen ϵ , there is always a corresponding δ that keeps $f(x)$ within the desired range.
-

▲ **Example 1.1.4** Use L'Hôpital's Rule to evaluate $\lim_{x \rightarrow 0} \frac{e^x - 1 - x - \frac{x^2}{2}}{x^3}$.

Solution Since both the numerator and denominator approach 0 as $x \rightarrow 0$, this is an indeterminate form of $\frac{0}{0}$. We can apply L'Hôpital's Rule by differentiating the numerator and the denominator.

1. Differentiate the numerator and denominator:

$$\lim_{x \rightarrow 0} \frac{e^x - 1 - x - \frac{x^2}{2}}{x^3} = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(e^x - 1 - x - \frac{x^2}{2})}{\frac{d}{dx}(x^3)} = \lim_{x \rightarrow 0} \frac{e^x - 1 - x}{3x^2}$$

2. This still gives an indeterminate form $\frac{0}{0}$, so we apply L'Hôpital's Rule again:

$$\lim_{x \rightarrow 0} \frac{e^x - 1 - x}{3x^2} = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(e^x - 1 - x)}{\frac{d}{dx}(3x^2)} = \lim_{x \rightarrow 0} \frac{e^x - 1}{6x}$$

3. Apply L'Hôpital's Rule one more time:

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{6x} = \lim_{x \rightarrow 0} \frac{\frac{d}{dx}(e^x - 1)}{\frac{d}{dx}(6x)} = \lim_{x \rightarrow 0} \frac{e^x}{6}$$

4. Evaluate the limit:

$$\frac{e^0}{6} = \frac{1}{6}$$

▲ **Example 1.1.5** Prove using the epsilon-delta definition that $\lim_{x \rightarrow 3}(2x + 5) = 11$.

Solution To prove this limit using the epsilon-delta definition, we need to show that for every $\epsilon > 0$, there exists a $\delta > 0$ such that whenever $0 < |x - 3| < \delta$, we have $|(2x + 5) - 11| < \epsilon$.

1. Start by simplifying $|(2x + 5) - 11|$:

$$\begin{aligned} |(2x + 5) - 11| &= |2x - 6| \\ &= |2(x - 3)| \\ &= 2|x - 3| \end{aligned}$$

2. We want $|2(x - 3)| < \epsilon$. This can be achieved by ensuring that:

$$2|x - 3| < \epsilon$$

or equivalently,

$$|x - 3| < \frac{\epsilon}{2}$$

3. Therefore, we can choose $\delta = \frac{\epsilon}{2}$. Now, if $0 < |x - 3| < \delta$, then:

$$|x - 3| < \frac{\epsilon}{2}$$

which implies that:

$$2|x - 3| < 2 \cdot \frac{\epsilon}{2} = \epsilon$$

4. Thus, $|(2x + 5) - 11| < \epsilon$, as required.

Single-Variable Derivatives

Calculus is fundamentally the theory of change, providing tools to understand and quantify how quantities evolve in relation to one another. In single-variable calculus, derivatives play a central role in this theory, allowing us to determine the rate at which a function changes at any given point. The derivative of a function represents the slope of the tangent line to the

function's graph, encapsulating the concept of instantaneous change, or how the function behaves at an exact point in time or space. Derivatives have widespread applications across various fields, enabling us to calculate velocity and acceleration in physics, model growth rates in biology, and analyze marginal changes in economics, among others.

As we progress in our study of calculus, we aim to extend this powerful framework by adding more variables, moving from single-variable functions to functions of several variables. This generalization will enable us to explore more complex systems where multiple factors interact, paving the way for multivariable calculus. Hence, this book is titled *Theories of Generalized Change*, as we will develop the tools to analyze and model change not just in one direction or along one axis, but across multiple dimensions and variables, broadening the scope and applicability of calculus.

To start with, derivatives are grounded in the concept of the **Instantaneous Rate of Change** (IROC), which refines the **Average Rate of Change** (AROC) by focusing on how a function behaves “at a single point.” While AROC provides the rate of change over a finite interval, IROC captures the behavior as this interval shrinks to zero, giving us a precise measure of change at an infinitesimal level. This approach of taking limits is foundational in calculus and will serve as the basis for more advanced studies in differentiation and integration.

- **Average Rate of Change (AROC):** The AROC of a function $f(x)$ over an interval $[a, b]$ is given by:

$$\text{AROC} = \frac{f(b) - f(a)}{b - a}$$

This is the slope of the secant line between the points $(a, f(a))$ and $(b, f(b))$ on the graph of f . The AROC gives an overall measure of change across the interval, providing insights but not specifics about any single point within that interval.




Figure 1.2: A secant line threading two points along a curve $f(x)$.

- **Instantaneous Rate of Change (IROC):** While AROC provides an average, we often seek to know how fast the function is changing at a specific point. The IROC at a point $x = a$ is the derivative of the function, calculated as the limit of the AROC as the interval becomes infinitesimally small.
- **Definition of the Derivative (Single Variable):** For a function $f(x)$, the derivative at a point $x = a$ is defined by the limit:

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

This expression represents the slope of the tangent line to f at the point $(a, f(a))$, capturing the instantaneous rate of change of $f(x)$ with respect to x at a .




Figure 1.3: A tangent line threading a point along a curve $f(x)$.

- **Basic Derivative Rules** The following rules simplify the process of finding derivatives for common functions.

– **Power Rule:** If $f(x) = x^n$, then

$$f'(x) = nx^{n-1}$$

– **Exponential Rule:** If $f(x) = e^x$, then

$$f'(x) = e^x$$

– **Logarithmic Rule:** If $f(x) = \ln(x)$, then

$$f'(x) = \frac{1}{x}$$

- **Sine Rule:** If $f(x) = \sin(x)$, then

$$f'(x) = \cos(x)$$

- **Cosine Rule:** If $f(x) = \cos(x)$, then

$$f'(x) = -\sin(x)$$

- **Product and Quotient Rules** When dealing with the derivatives of products or quotients of functions, we use these rules:

- **Product Rule:** If $f(x) = u(x)v(x)$, then

$$f'(x) = u'(x)v(x) + u(x)v'(x)$$

- **Quotient Rule:** If $f(x) = \frac{u(x)}{v(x)}$, then

$$f'(x) = \frac{u'(x)v(x) - u(x)v'(x)}{[v(x)]^2}$$

- **Chain Rule** For composite functions, we use the Chain Rule:

If $f(x) = g(h(x))$, then

$$f'(x) = g'(h(x)) \cdot h'(x)$$

The Chain Rule is essential when differentiating functions nested within other functions, allowing us to find the derivative of more complex expressions.

▲ **Example 1.1.6** Use the limit definition of the derivative to find the derivative of the function $f(x) = x^2$.

Solution The derivative of $f(x) = x^2$ at a point x can be found using the limit definition of the derivative:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

1. Substitute $f(x) = x^2$ into the formula:

$$f'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h}$$

2. Expand $(x+h)^2$:

$$f'(x) = \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - x^2}{h}$$

3. Simplify by canceling x^2 terms:

$$f'(x) = \lim_{h \rightarrow 0} \frac{2xh + h^2}{h}$$

4. Factor out h from the numerator:

$$f'(x) = \lim_{h \rightarrow 0} \frac{h(2x + h)}{h}$$

5. Cancel h in the numerator and denominator:

$$f'(x) = \lim_{h \rightarrow 0} (2x + h)$$

6. Take the limit as $h \rightarrow 0$:

$$f'(x) = 2x$$

Integrals

Integrals form a foundational concept in calculus, representing the accumulation of quantities and the area under a curve. Integrals have applications across mathematics and science, such as calculating area, volume, work, and in modeling the behavior of physical systems. The process of integration is often considered the reverse operation of differentiation, and it provides insights into quantities that are accumulated or summed continuously over intervals. In this section, we will explore the integral as a limit of finite sums (Riemann Sum), develop the concept of definite and indefinite integrals, and introduce several integration techniques.

- **Indefinite Integral:** The indefinite integral of a function $f(x)$, also known as the antiderivative, is defined as:

$$\int f(x) dx = F(x) + C$$

where $F'(x) = f(x)$ and C is the constant of integration. This represents the family of all antiderivatives of $f(x)$, capturing the essence of the function's accumulated effect.

- **Definite Integral:** The definite integral of $f(x)$ over the interval $[a, b]$ is defined as:

$$\int_a^b f(x) dx = F(b) - F(a)$$

where $F(x)$ is any antiderivative of $f(x)$. The definite integral represents the net area between the curve $f(x)$ and the x -axis from $x = a$ to $x = b$, taking into account areas above and below the axis.

- **Riemann Sum and the Integral as a Limit of Finite Sums:** The concept of integration can be understood through the Riemann Sum, which approximates the area under a curve by summing the areas of rectangles. As the number of rectangles increases (and their width decreases), the sum approaches the exact area under the curve.

For a function $f(x)$ over $[a, b]$, we divide the interval into n subintervals, each of width $\Delta x = \frac{b-a}{n}$. The Riemann sum is:

$$\sum_{i=1}^n f(x_i^*) \Delta x$$

where x_i^* is a sample point within each subinterval. The definite integral is defined as the limit of this sum as $n \rightarrow \infty$:

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$$




Figure 1.4: A Riemann Sum approximation for an integral of a function.

- **Approximations Using Left, Right, and Center Points; Maxima, Minima, Supremum, and Infimum:** Riemann sums allow for different choices of points within each subinterval to approximate the integral. These choices influence the accuracy and convergence of the sum, providing different approximations based on the choice of evaluation points within each subinterval.
 - **Left Endpoint Approximation:** The function is evaluated at the left endpoint of each subinterval x_i , producing a sum that may underestimate the area for increasing functions and overestimate for decreasing functions. The Riemann sum for n subintervals of width $\Delta x = \frac{b-a}{n}$ is:

$$\sum_{i=0}^{n-1} f(x_i) \Delta x$$

where $x_i = a + i\Delta x$ represents the left endpoint of each subinterval.

- **Right Endpoint Approximation:** The function is evaluated at the right endpoint of each subinterval x_{i+1} . This sum tends to overestimate for increasing functions and underestimate for decreasing functions:

$$\sum_{i=1}^n f(x_{i+1}) \Delta x$$

where $x_i = a + i\Delta x$ represents the right endpoint of each subinterval.

- **Midpoint (Center) Approximation:** The function is evaluated at the midpoint $x_i^* = \frac{x_i + x_{i+1}}{2}$ of each subinterval. This approach often yields a more accurate approximation by balancing overestimates and underestimates:

$$\sum_{i=0}^{n-1} f(x_i^*) \Delta x$$

where $x_i^* = a + (i + \frac{1}{2}) \Delta x$ is the midpoint.

- **Maximal Approximation:** For each subinterval $[x_i, x_{i+1}]$, choose the maximum function value, yielding an upper bound for the area:

$$\sum_{i=0}^{n-1} \max_{x \in [x_i, x_{i+1}]} f(x) \Delta x$$

This approximation gives an upper bound for the integral if $f(x)$ is continuous over $[a, b]$.

- **Minimal Approximation:** Similarly, choosing the minimum function value within each subinterval provides a lower bound:

$$\sum_{i=0}^{n-1} \min_{x \in [x_i, x_{i+1}]} f(x) \Delta x$$

This approximation serves as a lower bound if $f(x)$ is continuous over $[a, b]$.

- **Supremum and Infimum:** For theoretical purposes and cases with countably many discontinuous points, we can define Riemann sums using the supremum (least upper bound) and infimum (greatest lower bound) of $f(x)$ on each subinterval, providing bounds as $n \rightarrow \infty$. This approach is foundational in defining integrability:

$$\sum_{i=0}^{n-1} \sup_{x \in [x_i, x_{i+1}]} f(x) \Delta x \quad \text{and} \quad \sum_{i=0}^{n-1} \inf_{x \in [x_i, x_{i+1}]} f(x) \Delta x$$

These sums approximate the true integral more closely as $n \rightarrow \infty$.

- **Numerical Integration Techniques** When exact integration is challenging, numerical integration methods approximate the value of definite integrals by evaluating the function at specific points. Key methods include:

- **Trapezoidal Rule:** Approximates the area under the curve by dividing it into trapezoids rather than rectangles, averaging the left and right endpoint evaluations:

$$\int_a^b f(x) dx \approx \frac{\Delta x}{2} \left[f(x_0) + 2 \sum_{i=1}^{n-1} f(x_i) + f(x_n) \right]$$

where $\Delta x = \frac{b-a}{n}$ and $x_0, x_1, \dots, x_n$ are equally spaced points.

- **Simpson's Rule:** Uses parabolic segments to approximate the curve. It is more accurate for functions that are reasonably smooth and is defined as:

$$\int_a^b f(x) dx \approx \frac{\Delta x}{3} \left[f(x_0) + 4 \sum_{\text{odd } i} f(x_i) + 2 \sum_{\text{even } i} f(x_i) + f(x_n) \right]$$

Simpson's Rule provides excellent accuracy for a modest number of intervals, especially for polynomials and smooth functions.

▲ **Example 1.1.7** Use Simpson's Rule with $n = 4$ subintervals to approximate the integral $\int_0^2 \frac{1}{1+x^2} dx$. Compare the result with the exact value of the integral.

Solution Since Simpson's Rule requires knowing the exact value to compare, we start by calculating the exact value of the integral:

$$\int_0^2 \frac{1}{1+x^2} dx = [\arctan x]_0^2 = \arctan 2 - \arctan 0 = \arctan 2.$$

1. Calculate the width of each subinterval:

$$h = \frac{b-a}{n} = \frac{2-0}{4} = 0.5.$$

2. Determine the x -values:

$$x_0 = 0, \quad x_1 = 0.5, \quad x_2 = 1.0, \quad x_3 = 1.5, \quad x_4 = 2.0.$$

3. Compute the corresponding y -values:

$$\begin{aligned} y_0 &= \frac{1}{1+(0)^2} = 1, \\ y_1 &= \frac{1}{1+(0.5)^2} = \frac{1}{1.25} = 0.8, \\ y_2 &= \frac{1}{1+(1)^2} = \frac{1}{2} = 0.5, \end{aligned}$$

$$y_3 = \frac{1}{1 + (1.5)^2} = \frac{1}{3.25} \approx 0.3077,$$

$$y_4 = \frac{1}{1 + (2)^2} = \frac{1}{5} = 0.2.$$

4. Apply Simpson's Rule:

$$\int_0^2 \frac{1}{1+x^2} dx \approx \frac{h}{3} (y_0 + 4y_1 + 2y_2 + 4y_3 + y_4).$$

Substitute the computed values:

$$\int_0^2 \frac{1}{1+x^2} dx \approx \frac{0.5}{3} (1 + 4(0.8) + 2(0.5) + 4(0.3077) + 0.2).$$

Calculate the expression inside the parentheses:

$$\begin{aligned} &1 + 4(0.8) + 2(0.5) + 4(0.3077) + 0.2 \\ &= 1 + 3.2 + 1.0 + 1.2308 + 0.2 \\ &= 6.6308. \end{aligned}$$

Compute the approximation:

$$\int_0^2 \frac{1}{1+x^2} dx \approx \frac{0.5}{3} \times 6.6308 = \frac{3.3154}{3} \approx 1.1051.$$

5. Compare with the exact value:

$$\arctan 2 \approx 1.1071.$$

The Simpson's Rule approximation is 1.1051, while the exact value is approximately 1.1071. The error is approximately 0.0020.

- **Basic Integrals:** Several integral rules and formulas are essential for computing integrals efficiently. Here are some fundamental examples:

– **Power Rule:** For $n \neq -1$,

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C$$

– **Logarithmic Rule:**

$$\int \frac{1}{x} dx = \ln |x| + C$$

– **Exponential Rule:**

$$\int e^x dx = e^x + C$$

– **Sine Rule:**

$$\int \sin(x) dx = -\cos(x) + C$$

– **Cosine Rule:**

$$\int \cos(x) dx = \sin(x) + C$$

- **Integration by Parts:** This technique is useful when integrating products of functions. It is derived from the product rule for differentiation and is given by:

$$\int u dv = uv - \int v du$$

- **DI Method (Tabular Method):** To evaluate integrals using the DI method, construct a table with three columns: Sign, D , and I . Differentiate the function that simplifies upon differentiation (usually u) and integrate the function that does not change much upon integration (usually dv). This method is particularly useful in solving Partial Differential Equations (PDEs) because it systematically reduces the complexity of integrals involving products of functions, making it easier to handle boundary conditions and find solutions.

The objective is to continue differentiating in the D column until reaching 0. For each row, you multiply diagonally from the D term to the I term, following the specified sign. You can choose to stop early and integrate the last row separately if a pattern allows. Once the D column reaches zero, any further terms contribute nothing to the integral. We have color coded and marked with asterisks the terms in the table to add together.

Sign	D	I
+	u (*)	dv
–	u' (**)	$\int dv$ (*)
+	u'' (***)	$\int (\int dv) dv$ (**)
–	u'''	$\int (\int (\int dv) dv) dv$ (***)

Note: Integrate here for the last term.

The integral is computed by summing the diagonal products, following the pattern until the final term reaches zero.

- **Substitution Method:** Also known as u -substitution, this method is helpful for integrals involving compositions of functions. If $u = g(x)$, then

$$\int f(g(x))g'(x) dx = \int f(u) du$$

- **Definite Integral Evaluation Using the Fundamental Theorem of Calculus:** The Fundamental Theorem of Calculus links differentiation and integration, stating that if F is an antiderivative of f on $[a, b]$, then:

$$\int_a^b f(x) dx = F(b) - F(a)$$

This theorem provides a way to evaluate the exact area under a curve using antiderivatives.

▲ **Example 1.1.8** Evaluate $\int x^3 e^x dx$ using the DI (Tabular) Method.

Solution We'll apply the DI method, choosing x^3 to differentiate and e^x to integrate. Construct the table as follows:

Sign	D	I
+	x^3	e^x
-	$3x^2$	e^x
+	$6x$	e^x
-	6	e^x
+	0	e^x

Now, we combine terms according to the DI method by multiplying entries across each row (taking the signs into account) and summing the results:

$$\int x^3 e^x dx = x^3 e^x - 3x^2 e^x + 6x e^x - 6e^x + C$$

Other Useful Integration Rules

In addition to basic integration techniques, there are several advanced methods that help solve more complex integrals. These methods expand the range of integrals we can compute by breaking down complex functions into simpler parts, leveraging trigonometric identities, or applying specific substitutions.

- **Integration by Partial Fractions:** Partial fraction decomposition is useful for integrating rational functions (ratios of polynomials). This technique involves breaking a complicated rational function into a sum of simpler fractions that can be integrated individually.

– **Steps for Partial Fraction Decomposition:**

1. Factor the denominator, if possible.
2. Write the integrand as a sum of partial fractions. For example, if the denominator factors as $(x - a)(x - b)$, write:

$$\frac{P(x)}{(x - a)(x - b)} = \frac{A}{x - a} + \frac{B}{x - b}$$

3. Solve for the constants (A , B , etc.) by equating coefficients or substituting convenient values for x .
4. Integrate each term separately.

- **Trigonometric Integrals:** Trigonometric integrals involve expressions with trigonometric functions, such as $\sin^n(x)$, $\cos^m(x)$, or combinations like $\sin(x) \cos(x)$. Simplifying these integrals often requires trigonometric identities.

– **Key Identities for Simplification:**

$$\begin{aligned} * \sin^2(x) &= \frac{1 - \cos(2x)}{2} \\ * \cos^2(x) &= \frac{1 + \cos(2x)}{2} \\ * \sin(x) \cos(x) &= \frac{\sin(2x)}{2} \end{aligned}$$

- **Trigonometric Substitution:** Trigonometric substitution is useful for integrating expressions containing square roots, such as $\sqrt{a^2 - x^2}$, $\sqrt{a^2 + x^2}$, or $\sqrt{x^2 - a^2}$. By making a substitution involving trigonometric functions, we can transform these integrals into simpler forms.

– **Common Substitutions:**

Expression	Substitution	Resulting Identity
$\sqrt{a^2 - x^2}$	$x = a \sin(\theta)$	$\sqrt{a^2 - x^2} = a \cos(\theta)$
$\sqrt{a^2 + x^2}$	$x = a \tan(\theta)$	$\sqrt{a^2 + x^2} = a \sec(\theta)$
$\sqrt{x^2 - a^2}$	$x = a \sec(\theta)$	$\sqrt{x^2 - a^2} = a \tan(\theta)$

- **Reduction Formulas:** Reduction formulas provide a recursive way to evaluate integrals of certain functions, often involving powers of trigonometric functions or products of functions. These formulas allow us to reduce the power of the integrand step-by-step, eventually reaching a simpler integral.

– **Example of a Reduction Formula for Powers of Sine:**

$$\int \sin^n(x) dx = -\frac{\sin^{n-1}(x) \cos(x)}{n} + \frac{n-1}{n} \int \sin^{n-2}(x) dx$$

Taylor Series

Taylor Series provide a way to approximate functions using polynomials, which can make complex functions easier to work with in calculus, physics, engineering, and beyond. This series expansion allows us to approximate functions locally around a point by using the derivatives of the function at that point.

- **Definition:** For a function $f(x)$ that is infinitely differentiable at a , the Taylor series expansion of $f(x)$ around $x = a$ is:

$$f(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots$$

or, in summation notation:

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$$

This expansion allows us to represent $f(x)$ as an infinite series of terms involving powers of $(x - a)$, with each term weighted by the function's n -th derivative evaluated at a . The closer x is to a , the better the approximation.

- **Maclaurin Series:** A Maclaurin series is a special case of the Taylor series centered at $a = 0$:

$$f(x) = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!}x^n$$

Maclaurin series are commonly used for standard functions such as e^x , $\sin(x)$, and $\cos(x)$.

- **Common Maclaurin Series Expansions:**

These series expansions provide useful approximations for frequently encountered functions:

- **Exponential Function e^x :**

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

The exponential function e^x is unique in that it is its own derivative and its own Taylor series, making it fundamental in calculus.

- **Sine Function $\sin(x)$:**

$$\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$

This series shows the odd power expansion of $\sin(x)$, alternating in sign. It is especially useful in approximations for small x , where only a few terms may be needed.

- **Cosine Function** $\cos(x)$:

$$\cos(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \cdots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

Similar to the sine series, the cosine function's Maclaurin series consists of even powers of x , with alternating signs. This series provides an accurate approximation for $\cos(x)$ near $x = 0$.

- **Logarithmic Function** $\ln(1 + x)$ (valid for $-1 < x \leq 1$):

$$\ln(1 + x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \cdots = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n}$$

This series is useful in approximating logarithmic functions around $x = 0$.

- **Arc Tangent Function** $\arctan(x)$ (valid for $-1 \leq x \leq 1$):

$$\arctan(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \cdots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}$$

This series provides an approximation for the inverse tangent function near $x = 0$, often useful in trigonometric applications.

- **Remainder Term and Approximation Accuracy:** When using Taylor or Maclaurin series to approximate functions, it is important to understand the error or remainder involved in truncating the series. For a Taylor series approximation up to the n -th term, the remainder $R_n(x)$ gives the difference between the actual function and the Taylor polynomial.

$$R_n(x) = \frac{f^{(n+1)}(c)}{(n+1)!} (x - a)^{n+1} \text{ for some } c \text{ between } a \text{ and } x$$

This remainder term shows how the accuracy of the approximation depends on the higher derivatives of $f(x)$, the point x , and the number of terms included in the approximation.

▲ **Example 1.1.9** Find the Taylor series of $f(x) = \ln(1 + x)$ centered at $a = 0$ up to the third term.

Solution To find the Taylor series, we calculate the derivatives of $f(x)$ at $x = 0$ and use them to construct the series.

1. Calculate $f(x)$ and its derivatives:

$$f(x) = \ln(1 + x)$$

$$f'(x) = \frac{1}{1+x}$$

$$f''(x) = -\frac{1}{(1+x)^2}$$

$$f'''(x) = \frac{2}{(1+x)^3}$$

2. Evaluate these derivatives at $x = 0$:

$$f(0) = \ln(1) = 0$$

$$f'(0) = 1$$

$$f''(0) = -1$$

$$f'''(0) = 2$$

3. Use the Taylor series formula up to the third term:

$$f(x) \approx f(0) + f'(0)(x) + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3$$

4. Substitute the values:

$$f(x) \approx 0 + x - \frac{1}{2}x^2 + \frac{1}{3}x^3$$

Important Theorems and Results in Calculus

Understanding key theorems in calculus is crucial for solving ordinary differential equations (ODEs), as these results often provide the foundation for methods of integration, approximation, and analysis. They help in determining the behavior of solutions, establishing existence and uniqueness, and transforming complex problems into more manageable forms. Here are some of the most significant theorems:

- **Mean Value Theorem (MVT):** If $f(x)$ is continuous on $[a, b]$ and differentiable on (a, b) , then there exists a point c in (a, b) such that:

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

This theorem asserts that for a smooth curve between a and b , there is at least one point where the tangent (instantaneous rate of change) is parallel to the secant line connecting $(a, f(a))$ and $(b, f(b))$.

- **Rolle's Theorem:** If $f(x)$ is continuous on $[a, b]$, differentiable on (a, b) , and $f(a) = f(b)$, then there exists a point c in (a, b) such that $f'(c) = 0$. This theorem is a special

case of the MVT, asserting that if a function returns to the same value at two points, there must be at least one point in between where the derivative (slope of the tangent) is zero.

- **Intermediate Value Theorem (IVT):** If $f(x)$ is continuous on $[a, b]$ and k is any value between $f(a)$ and $f(b)$, then there exists a point c in $[a, b]$ such that $f(c) = k$. The IVT guarantees that a continuous function takes on every intermediate value between $f(a)$ and $f(b)$, a property crucial for establishing the existence of solutions.
- **Extreme Value Theorem (EVT):** If $f(x)$ is continuous on a closed interval $[a, b]$, then f attains both a maximum and a minimum value on $[a, b]$. This theorem ensures that a continuous function over a closed interval has both an absolute maximum and minimum, critical for optimization problems and boundary analyses.
- **L'Hôpital's Rule:** If $\lim_{x \rightarrow a} \frac{f(x)}{g(x)}$ results in the indeterminate form $\frac{0}{0}$ or $\frac{\infty}{\infty}$, then:

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

provided the limit on the right exists or is infinite. This rule is essential for evaluating limits that yield indeterminate forms, often encountered in calculus and differential equations.

- **First Fundamental Theorem of Calculus (FTOC I):** If f is continuous on $[a, b]$ and F is an antiderivative of f on $[a, b]$, then:

$$\int_a^b f(x) dx = F(b) - F(a)$$

This theorem links the process of differentiation and integration, showing that the definite integral of a function over an interval can be computed using any of its antiderivatives.

- **Second Fundamental Theorem of Calculus (FTOC II):** If f is continuous on an interval containing a , then for any x in that interval:

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

This theorem states that the derivative of the integral function (accumulated area) is the original function, demonstrating how integration can "undo" differentiation.

- **Leibniz Rule for Differentiation under the Integral Sign:** If $f(x, t)$ is continuous on x and t , and $\partial f / \partial t$ exists and is continuous, then:

$$\frac{d}{dx} \int_{a(x)}^{b(x)} f(x, t) dt = f(x, b(x)) \cdot \frac{db}{dx} - f(x, a(x)) \cdot \frac{da}{dx} + \int_{a(x)}^{b(x)} \frac{\partial}{\partial x} f(x, t) dt$$

This rule is crucial in scenarios where integration limits depend on x , enabling differentiation within the integral sign.

- **Taylor's Theorem** For a function $f(x)$ that is $n + 1$ -times differentiable at a , Taylor's Theorem gives an approximation for $f(x)$ near a :

$$f(x) = f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \cdots + \frac{f^{(n)}(a)}{n!}(x - a)^n + R_n(x)$$

where $R_n(x)$ is the remainder term. In the limit as $n \rightarrow \infty$, if $R_n(x) \rightarrow 0$, we obtain the Taylor Series:

$$f(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(a)}{k!}(x - a)^k$$

Taylor's Theorem is fundamental in approximating functions locally using polynomials and is widely applied in numerical analysis and differential equations.

▲ **Example 1.1.10** Verify that the function $f(x) = x^2 - 3x + 2$ satisfies the Mean Value Theorem on the interval $[1, 4]$, and find the value of c that satisfies the theorem.

Solution To apply the Mean Value Theorem (MVT), we need to confirm that $f(x)$ is continuous on $[1, 4]$ and differentiable on $(1, 4)$. Since $f(x) = x^2 - 3x + 2$ is a polynomial, it is continuous and differentiable everywhere, so the MVT applies on this interval. According to the Mean Value Theorem, there exists a point $c \in (1, 4)$ such that:

$$f'(c) = \frac{f(4) - f(1)}{4 - 1}$$

1. Calculate $f(4)$ and $f(1)$:

$$\begin{aligned} f(4) &= 4^2 - 3 \cdot 4 + 2 = 16 - 12 + 2 = 6 \\ f(1) &= 1^2 - 3 \cdot 1 + 2 = 1 - 3 + 2 = 0 \end{aligned}$$

2. Compute the average rate of change:

$$\frac{f(4) - f(1)}{4 - 1} = \frac{6 - 0}{3} = 2$$

3. Differentiate $f(x)$ and set $f'(c) = 2$:

$$f'(x) = 2x - 3$$

Setting $f'(c) = 2$, we get:

$$2c - 3 = 2$$

Solving for c :

$$2c = 5 \Rightarrow c = \frac{5}{2} = 2.5$$

4. Verify $c \in (1, 4)$: Since $c = 2.5$ lies within the interval $(1, 4)$, it satisfies the conditions of the Mean Value Theorem.

Lecture 1.2: Three-Dimensional Coordinate Systems

Transitioning from two-dimensional to three-dimensional coordinate systems marks a significant leap in our capacity to represent and analyze objects and interactions spatially. This shift extends our geometric understanding, equipping us with a powerful framework for calculating points, distances, and angles within three-dimensional space—a foundational requirement for advanced calculus, physics, engineering, and numerous other fields.

Rectangular Coordinates in Two Dimensions

In two dimensions, every point on a plane is specified by a coordinate pair, (x, y) , where x and y represent distances along two mutually perpendicular axes: the x -axis (horizontal) and y -axis (vertical). This two-dimensional system is essential for understanding how we map points in a flat plane and serves as a stepping stone for more complex spatial reasoning. In a 2D Cartesian system, an ordered pair (x, y) not only specifies location but also allows us to measure distances, areas, and angular relationships on a flat plane.




Figure 1.5: 2D Cartesian Coordinate System showing points such as $(2, 3)$ and $(-3, 2)$

Introduction to Three-Dimensional Space

By introducing a third axis, the z -axis, we can move into three-dimensional space, where each point is described by an ordered triple (x, y, z) . The addition of the z -axis—perpendicular to both x and y —allows us to locate points and measure distances not only horizontally and vertically but also in terms of depth or height. This extension from two to three dimensions mirrors our physical world and enables us to mathematically describe real objects, from simple geometric shapes to complex structures.

In three-dimensional space:

- x : Measures the position along the x -axis (typically horizontal).
- y : Measures the position along the y -axis (perpendicular to the x -axis).
- z : Measures the position along the z -axis (perpendicular to both x and y axes).

This system allows us to define planes formed by pairs of axes:

- *The xy -plane*, where $z = 0$, represents a flat surface in the horizontal and vertical dimensions. This is the familiar coordinate plane from previous mathematics courses.
- *The xz -plane*, where $y = 0$, represents a plane incorporating depth along z and position along x . This plane would appear as a vertical wall if you were looking directly at it.
- *The yz -plane*, where $x = 0$, allows us to represent depth along z and height along y . This too appears as a vertical wall, but perpendicular to the xz -plane.

These three coordinate planes divide space into eight regions called octants, with the first octant being where all coordinates are positive. This organization helps us understand spatial relationships and serves as the foundation for our study of multivariable calculus. Throughout this course, we will learn to analyze functions, curves, and surfaces in this three-dimensional setting, building upon our knowledge of calculus in one and two dimensions.

When visualizing three-dimensional objects, we often use perspective drawings where parallel lines appear to meet at a distance. This technique helps us represent three-dimensional concepts on a two-dimensional page while maintaining our intuitive understanding of spatial relationships.




Figure 1.6: 3D Cartesian Coordinate System with sample points in space connected by lines.

The intersection of these planes at the origin divides space into eight distinct regions, called octants. We locate any point (x, y, z) by moving x units along the x -axis, y units along

the y -axis, and z units along the z -axis. This spatial representation is fundamental in fields like navigation, robotics, and physics, where understanding three dimensions is critical to applications ranging from satellite positioning to structural analysis.




Figure 1.7: *The eight octants of a three-dimensional Cartesian plot.*

Toward Higher Dimensions: Imagining Four-Dimensional Space

Extending to a fourth spatial dimension—though challenging to visualize—invites us to think beyond our physical intuition. In a four-dimensional space, each point is represented by an ordered quadruple (x, y, z, w) , where w denotes the fourth dimension. Though difficult to imagine, one can think of w as adding an additional degree of freedom beyond our typical three-dimensional experience. Mathematicians and physicists work in four-dimensional (and higher-dimensional) spaces when dealing with complex phenomena, such as in the study of spacetime in general relativity.

A helpful analogy is to consider how three-dimensional objects can cast two-dimensional shadows. In the same way, a four-dimensional object, known as a tesseract or hypercube, could project a three-dimensional “shadow.” This shadow could show us a trace of four-dimensional interactions, allowing us to infer properties of the higher dimension. While our intuition may falter beyond three dimensions, mathematical models provide structure. Calculations in higher-dimensional spaces can help solve problems in fields such as quantum mechanics and data science, where multi-dimensional datasets reveal patterns not visible in lower dimensions.

Distance and Midpoint Formulas in Three Dimensions

The distance and midpoint formulas in three-dimensional space are extensions of the two-dimensional formulas, derived from the Pythagorean Theorem. By understanding these

derivations, we gain insight into the geometry of space and the relationships between points in three dimensions.

Derivation of the Distance Formula

The distance formula in geometry is a direct extension of the Pythagorean Theorem, which states that for a right triangle with legs a and b , and hypotenuse c , the relationship between the sides is $a^2 + b^2 = c^2$. We rearrange this equation for the hypotenuse to get $c = \sqrt{a^2 + b^2}$. This formula gives the length of the hypotenuse when the lengths of the other two sides are known. Now, consider finding the distance between two points $P_1(x_1, y_1)$ and $P_2(x_2, y_2)$ in a two-dimensional plane. To derive this distance, we construct a right triangle by taking the horizontal and vertical distances between these points:

$$a = x_2 - x_1 \quad \text{and} \quad b = y_2 - y_1.$$

Substituting a and b into the Pythagorean Theorem, we get:

$$(x_2 - x_1)^2 + (y_2 - y_1)^2 = d^2,$$

where d represents the distance between P_1 and P_2 . Solving for d gives us the two-dimensional distance formula:

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}.$$

Extending the Distance Formula to Three Dimensions

To generalize this result to three-dimensional space, we need to consider an additional dimension, the z -axis. Given two points $P_1(x_1, y_1, z_1)$ and $P_2(x_2, y_2, z_2)$, the change in the z -direction introduces an additional “side” in the Pythagorean relationship. Now, we have three differences:

$$a = x_2 - x_1, \quad b = y_2 - y_1, \quad \text{and} \quad c = z_2 - z_1.$$

We follow the same logic as before to obtain

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}.$$

This formula allows us to calculate the straight-line distance between any two points in three-dimensional space by considering the differences along each of the three axes. Notice, that because of the square terms, the distance will always be positive or zero. Since we saw that for the Pythagorean theorem all we had to do to increase the dimensionality was include one more square term, let's do that n -many times to go to any n th dimension we desire. Thus we obtain for any n -dimension we have

$$d = \left(\sum_{j=1}^n (y_j - x_j)^2 \right)^{1/2} = ((y_1 - x_1)^2 + \cdots + (y_n - x_n)^2)^{1/2}$$

Midpoint Formula

Similarly, the midpoint formula in three dimensions is derived from averaging the coordinates of the endpoints. For points $P_1(x_1, y_1, z_1)$ and $P_2(x_2, y_2, z_2)$, the midpoint M of the line segment joining P_1 and P_2 is given by:

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2} \right).$$

This midpoint divides the line segment into two equal halves, allowing us to locate the center point between P_1 and P_2 in three-dimensional space.

▲ Example 1.2.1 A drone starts at point P_1 with coordinates $(2, 3, 5)$ and needs to reach a target at point P_2 located at $(7, 1, 10)$. Find the straight-line distance the drone needs to travel to reach the target.

Solution To calculate the straight-line distance between points $P_1(2, 3, 5)$ and $P_2(7, 1, 10)$ in three-dimensional space, we apply the distance formula:

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

1. Identify the coordinates:

$$P_1(x_1, y_1, z_1) = (2, 3, 5), \quad P_2(x_2, y_2, z_2) = (7, 1, 10)$$

2. Substitute the coordinates into the distance formula:

$$d = \sqrt{(7 - 2)^2 + (1 - 3)^2 + (10 - 5)^2}$$

3. Calculate each component:

$$\begin{aligned} (x_2 - x_1)^2 &= (7 - 2)^2 = 5^2 = 25 \\ (y_2 - y_1)^2 &= (1 - 3)^2 = (-2)^2 = 4 \\ (z_2 - z_1)^2 &= (10 - 5)^2 = 5^2 = 25 \end{aligned}$$

4. Sum the components and take the square root:

$$d = \sqrt{25 + 4 + 25} = \sqrt{54} = 3\sqrt{6} \approx 7.35$$

Thus, the drone must travel approximately 7.35 units to reach its target.

Introduction to Polar Coordinates

Polar coordinates provide an alternative way to describe points in a plane using a distance and an angle relative to a central point, typically the origin. The term “polar” is derived from the Latin word *polus*, meaning “axis” or “pivot point,” indicating how this coordinate system revolves around a single fixed point as its central reference. In polar coordinates, each point is represented by an ordered pair (r, θ) , where:

- r : The radial distance from the origin (the straight-line distance from the point to the central origin).
- θ : The angular direction, measured from the positive x -axis, typically in degrees or radians, representing the angle at which the point is located relative to the origin.

Polar coordinates are particularly useful in situations involving circular or radial motion, such as describing points that orbit around a center. For instance, when an airplane travels in a curved path from one location to another, rectangular coordinates (i.e., x and y) may be inefficient, as they do not inherently account for direction and distance from a central point. Instead, polar coordinates allow the airplane’s position to be easily expressed in terms of distance and angle relative to the origin or any central reference, simplifying navigation, particularly over long distances along curved paths, such as on the Earth’s surface.

Conversion Between Rectangular and Polar Coordinates

The conversion between rectangular and polar coordinates relies on the Pythagorean Theorem and trigonometric relationships. Given a point (x, y) in rectangular coordinates:

- The distance r , from the origin to the point, can be derived using the Pythagorean Theorem:

$$r = \sqrt{x^2 + y^2}$$

- The angle θ , representing the direction from the positive x -axis, is found using trigonometry:

$$\theta = \tan^{-1} \left(\frac{y}{x} \right)$$

To convert back from polar to rectangular coordinates:

- The x -coordinate can be found using:

$$x = r \cos \theta$$

- The y -coordinate can be found using:

$$y = r \sin \theta$$

To illustrate the difference between rectangular and polar coordinates, let's examine the point $(3, 4)$ in both systems. In rectangular coordinates, the point $(3, 4)$ represents a location 3 units along the x -axis and 4 units along the y -axis. To express this point in polar coordinates, we calculate r and θ as follows:

$$r = \sqrt{3^2 + 4^2} = \sqrt{9 + 16} = \sqrt{25} = 5$$

$$\theta = \tan^{-1}\left(\frac{4}{3}\right) \approx 53.13^\circ$$

Thus, the point $(3, 4)$ in rectangular coordinates is equivalent to $(5, 53.13^\circ)$ in polar coordinates. The following plot demonstrates both coordinate systems for this point. Rectangular coordinates (x, y) are shown on the standard Cartesian grid, while polar coordinates (r, θ) use circles centered at the origin and rays extending out at different angles.




Figure 1.8: Plot showing both Rectangular and Polar representations of the point $(3, 4)$ with a legend for coordinate paths.

Cylindrical Coordinates

The cylindrical coordinate system is an extension of polar coordinates, adding a vertical z -dimension to describe points in three-dimensional space. It is particularly useful for representing objects with circular symmetry around an axis, such as cylinders, tubes, and circular regions extending along a vertical axis. In cylindrical coordinates, each point is represented by (r, θ, z) , where:

- r : The radial distance from the z -axis, which is the distance from the point to the z -axis in the xy -plane.

- θ : The angle measured from the positive x -axis within the xy -plane, in radians, describing the angular position around the z -axis.
- z : The vertical distance from the xy -plane, equivalent to the z -coordinate in rectangular coordinates, indicating height or depth.

In essence, cylindrical coordinates generalize polar coordinates by incorporating an additional z -coordinate, allowing us to describe points in three-dimensional space while maintaining the circular symmetry of polar coordinates around a central axis. If we set $z = 0$, cylindrical coordinates reduce to polar coordinates in the xy -plane, where a point is solely described by its distance r from the origin and the angle θ .




Figure 1.9: A point in three-dimensional space represented in cylindrical coordinates (r, θ, z) . The radius r is the distance from the z -axis, θ is the angle measured from the positive x -axis, and z is the vertical coordinate.

Cylindrical coordinates are widely used in fields like physics and engineering where problems often exhibit cylindrical or rotational symmetry. For instance, in analyzing magnetic fields around a wire, fluid flow in pipes, or any phenomenon with a circular base extending along a vertical dimension, cylindrical coordinates provide a natural and efficient way to describe the system.

Converting Between Cylindrical and Rectangular Coordinates:

To switch between cylindrical and rectangular coordinates, we use trigonometric relationships similar to those in polar coordinates, with the z -coordinate remaining consistent between both systems.

- To convert from cylindrical (r, θ, z) to rectangular (x, y, z) :

$$x = r \cos \theta, \quad y = r \sin \theta, \quad z = z$$

This transformation projects the point (r, θ) in the xy -plane into x - and y -coordinates using trigonometry, while z remains unchanged.

- To convert from rectangular (x, y, z) to cylindrical (r, θ, z) :

$$r = \sqrt{x^2 + y^2}, \quad \theta = \tan^{-1} \left(\frac{y}{x} \right), \quad z = z$$

Here, r is calculated as the distance from the point to the z -axis (similar to the Pythagorean distance in polar coordinates), and θ is the angle around the z -axis. The z -coordinate remains the same in both representations.

This conversion framework allows easy switching between cylindrical and rectangular coordinates depending on the symmetry and nature of the problem at hand. In problems with circular symmetry along an axis, cylindrical coordinates simplify analysis and calculation by leveraging the radial and angular structure inherent to the system.

Spherical Coordinates

Spherical coordinates are ideal for problems with radial symmetry, such as those involving spheres or radial fields. Each point in spherical coordinates is represented by (ρ, θ, ϕ) :

- ρ : The radial distance from the origin (distance from the point to the origin).
- θ : The angle in the xy -plane from the positive x -axis, measured counterclockwise.
- ϕ : The angle between the point and the positive z -axis (measured from the z -axis downward).

Spherical coordinates are particularly useful for scenarios where an object moves in a circular arc around a central point and has varying elevation, such as a planet orbiting a star, a camera with full rotation, or a baseball bat's swing through the air. When a softball player swings a bat to hit a ball, the bat moves in a curved, sweeping motion around the player. The motion of the bat can be best described in spherical coordinates:

- Radial Distance (ρ): The distance of the bat's tip from the player's body (the origin) varies, allowing for control over how close or far the bat reaches.
- Horizontal Angle (θ): The angle θ in the xy -plane represents the circular path of the bat around the player as they swing.
- Vertical Angle (ϕ): The vertical angle ϕ describes how high or low the bat is swung, which varies depending on whether the player adjusts to hit a high or low pitch.




Figure 1.10: A representation of the spherical coordinate system.

Unlike cylindrical coordinates, which would only capture the horizontal rotation around a central vertical axis (the player's body), spherical coordinates provide a complete representation of the bat's path, including its height and depth relative to the player's position. Thus, spherical coordinates offer a more accurate and flexible framework for capturing the full 3D motion of a softball swing.



Figure 1.11: A softball swing is best represented in spherical coordinates, capturing the radial distance, horizontal angle, and vertical angle of the bat's position relative to the player.

Converting Between Spherical and Rectangular Coordinates:

- To convert from spherical (ρ, θ, ϕ) to rectangular (x, y, z) :

$$x = \rho \sin \phi \cos \theta, \quad y = \rho \sin \phi \sin \theta, \quad z = \rho \cos \phi$$

- To convert from rectangular (x, y, z) to spherical (ρ, θ, ϕ) :

$$\rho = \sqrt{x^2 + y^2 + z^2}, \quad \theta = \tan^{-1} \left(\frac{y}{x} \right), \quad \phi = \cos^{-1} \left(\frac{z}{\rho} \right)$$

▲ **Example 1.2.2** A bird is flying along a circular path around a radio tower, while also increasing its altitude. At a certain moment, the bird's position relative to the tower is given by the cylindrical coordinates $(r, \theta, z) = (100, \frac{\pi}{4}, 50)$, where r is in meters and z represents height. Convert this position into rectangular coordinates to determine the bird's (x, y, z) coordinates in the 3D plane.

Solution To convert from cylindrical coordinates (r, θ, z) to rectangular coordinates (x, y, z) , we use the following conversion formulas:

$$x = r \cos \theta, \quad y = r \sin \theta, \quad z = z$$

Given:

$$r = 100, \quad \theta = \frac{\pi}{4}, \quad z = 50$$

1. Calculate x :

$$x = 100 \cos \frac{\pi}{4} = 100 \cdot \frac{\sqrt{2}}{2} = 50\sqrt{2}$$

2. Calculate y :

$$y = 100 \sin \frac{\pi}{4} = 100 \cdot \frac{\sqrt{2}}{2} = 50\sqrt{2}$$

3. The z -coordinate remains the same:

$$z = 50$$

4. Therefore, the rectangular coordinates of the bird's position are:

$$(x, y, z) = (50\sqrt{2}, 50\sqrt{2}, 50)$$

▲ **Example 1.2.3** In an automated car assembly plant, a spherical robotic arm is used to position car parts in 3D space. The arm operates by extending its length, rotating horizontally, and pivoting vertically. At a certain moment, the robotic arm performs the following actions:

- Extends to a length of 2 meters.
- Rotates 60° counterclockwise from the positive x -axis.
- Raises the arm 45° above the horizontal plane.

Identify which spherical coordinate (ρ , θ , or ϕ) corresponds to each action, and write down the spherical coordinates of the arm's end effector after this movement.

Solution In spherical coordinates:

- ρ represents the distance from the origin to the point (length of the arm).
- θ is the angle measured in the xy -plane from the positive x -axis (horizontal rotation).
- ϕ is the angle measured from the positive z -axis down to the point (vertical angle).

Given the actions:

1. Extension to a length of 2 meters:

- This corresponds to $\rho = 2$ m.

2. Rotation 60° counterclockwise from the positive x -axis:

- This corresponds to $\theta = 60^\circ$ or $\theta = \frac{\pi}{3}$ radians.

3. Raising the arm 45° above the horizontal plane:

- Since ϕ is measured from the positive z -axis downwards, we need to calculate ϕ .
- The horizontal plane corresponds to $\phi = 90^\circ$ (as it is 90° down from the positive z -axis).
- Raising the arm 45° above the horizontal plane means the angle from the positive z -axis is:

$$\phi = 90^\circ - 45^\circ = 45^\circ \quad \text{or} \quad \phi = \frac{\pi}{4} \text{ radians.}$$

Choice of Coordinates

The choice of coordinate system is essential in simplifying calculations and providing clarity in solving problems. Each coordinate system has advantages depending on the problem's geometry, symmetry, and underlying physical properties. Below is a comparison of when each system is most useful.

- **Rectangular (Cartesian) Coordinates:** In rectangular coordinates, each point is specified by its perpendicular distances from the x -, y -, and z -axes. This system is intuitive and ideal for objects and problems aligned along the three principal axes, such as rectangular prisms, planes, and many engineering structures.
- **Polar Coordinates:** Polar coordinates are well-suited for two-dimensional problems involving circular or rotational symmetry around a central point. Problems that naturally involve rotation or radial symmetry—such as the motion of a particle in a circular path or objects positioned relative to a central point—benefit from polar coordinates.
- **Cylindrical Coordinates:** Cylindrical coordinates are an extension of polar coordinates into three dimensions, making them ideal for problems with circular symmetry around a central vertical axis, such as cylindrical objects or fields surrounding a wire. For example, fluid flow in a pipe, magnetic fields around a wire, or the structure of a cylindrical tank are best analyzed using cylindrical coordinates.
- **Spherical Coordinates:** Spherical coordinates extend the idea of radial symmetry into three dimensions and are best suited for objects and fields with a spherical or radial symmetry, such as planets, stars, and electric fields radiating from a point charge.

Lecture 1.3: Vectors

In progressing from two-dimensional space to three-dimensional space, vectors become an essential concept for describing both position and direction. Vectors allow us to represent not only points but also the direction and magnitude of movement or force in space, forming the basis for advanced topics in calculus.

Definition of Vectors

As we move from two-dimensional space into three dimensions, vectors become crucial for describing both position and direction. Vectors aren't just about marking a location—they let us describe how something moves or acts, combining both its size (magnitude) and direction. This combination of information forms the basis of advanced ideas in calculus, physics, and engineering. For instance, knowing the strength of a wind gust without knowing the direction isn't very helpful. But by representing it as a vector, we gain both its intensity and the path it follows.

We represent a vector in three-dimensional space with an ordered triple $\mathbf{v} = \langle v_x, v_y, v_z \rangle$, where:

- v_x : Shows the vector's reach along the x -axis.
- v_y : Shows its reach along the y -axis.
- v_z : Shows its reach along the z -axis.

Each component captures the vector's effect along one axis, allowing it to describe anything with direction and magnitude. In addition to using $\mathbf{v}$, we may also write $\vec{v}$. This is typically what students write in their notebooks when indicating a vector. For listing components of a vector we may write either (v_x, v_y, v_z) or $\langle v_x, v_y, v_z \rangle$. To eliminate ambiguity from describing points, we will use the latter notation.

To make vectors more tangible, imagine them as arrows in space. Each arrow's direction and length tell us the direction and strength of an action or movement—like the current in a river. At each point along the river, an arrow (or vector) can show how fast and in which direction the water flows. Vectors make it possible to describe these continuous, directed motions, making them ideal for representing physical forces or speeds in space.

Vectors also have special properties that make them versatile tools. For example, they can be moved around in space without changing their meaning, a property called *translational invariance*. This means a vector maintains its effect regardless of its starting point, allowing us to describe forces or velocities that act consistently no matter where they occur. They can also be “stacked” or added together, which is essential in physics. For example, if two people push a box from different directions, the net effect on the box's movement can be represented by the sum of their vectors. Vectors can also be scaled up or down, describing changes in strength or intensity. For example, if a car speeds up, its

velocity vector is scaled up to reflect its increased speed while keeping the same direction. This operation, called *scalar multiplication*, lets us adjust vectors without altering their paths.




Figure 1.12: A vector in three-dimensional space represented by components along x , y , and z axes.

Vector Magnitude and Direction

The *magnitude* or *length* of a vector $\mathbf{v} = \langle v_x, v_y, v_z \rangle$ is denoted by $|\mathbf{v}|$ (and in some advanced texts as $\|\mathbf{v}\|$) and is calculated by using the Pythagorean theorem extended to three dimensions. This is because each component of the vector v_x , v_y , and v_z represents the vector's “reach” along the x -, y -, and z -axes. The magnitude formula is directly related to the distance formula covered previously:

$$|\mathbf{v}| = \sqrt{v_x^2 + v_y^2 + v_z^2}.$$

For example, if $\mathbf{v} = (3, -4, 12)$, then:

$$|\mathbf{v}| = \sqrt{3^2 + (-4)^2 + 12^2} = \sqrt{9 + 16 + 144} = \sqrt{169} = 13.$$

The magnitude provides a sense of “how much” or “how far” the vector reaches in space, without giving any indication of direction. With magnitude, we gain a crucial part of the vector’s identity, as it represents the scale or intensity of whatever the vector is describing—such as the speed of a moving object or the strength of a force.

Unit Vectors and Direction Cosines

A *unit vector* is a vector that has been scaled to have a magnitude of exactly 1 while keeping its direction unchanged. Unit vectors serve as pure directional indicators, providing only

the direction without any size or scale. They are especially useful for defining the principal directions along each axis in three-dimensional space. In three-dimensional space, we define the standard unit vectors along the x -, y -, and z -axes as:

$$\mathbf{i} = \langle 1, 0, 0 \rangle, \quad \mathbf{j} = \langle 0, 1, 0 \rangle, \quad \mathbf{k} = \langle 0, 0, 1 \rangle.$$

These vectors— $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ —represent the directions along each principal axis with a magnitude of 1, making them ideal for constructing and breaking down other vectors. In other texts you may also see them written as $\hat{\mathbf{i}}$, $\hat{\mathbf{j}}$, and $\hat{\mathbf{k}}$. Typically important unit vectors are also written this way with a hat. Any vector $\mathbf{v} = \langle v_x, v_y, v_z \rangle$ can be written in terms of $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ as:

$$\mathbf{v} = v_x \mathbf{i} + v_y \mathbf{j} + v_z \mathbf{k}.$$

Expressing vectors in this form allows us to separate and analyze their individual components along each axis, which simplifies many operations in vector algebra and physics. To create a unit vector $\mathbf{u}$ that points in the same direction as a given vector $\mathbf{v}$, we divide $\mathbf{v}$ by its magnitude $|\mathbf{v}|$. For a vector $\mathbf{v} = \langle v_x, v_y, v_z \rangle$, the unit vector $\mathbf{u}$ in the direction of $\mathbf{v}$ is:

$$\mathbf{u} = \frac{\mathbf{v}}{|\mathbf{v}|} = \left\langle \frac{v_x}{|\mathbf{v}|}, \frac{v_y}{|\mathbf{v}|}, \frac{v_z}{|\mathbf{v}|} \right\rangle.$$

This process of scaling a vector to make it a unit vector is called *normalization*. Normalization allows us to create a standardized version of any vector that preserves its direction while adjusting its magnitude to exactly 1. We will see normal vectors all over this entire study of multi-variable Calculus—more to be addressed about this soon.

The concept of unit vectors is valuable in many applications because they give us a standardized way to indicate direction independently of magnitude. For instance, if we only care about the direction an object is moving (such as “due north”) and not its speed, we can use a unit vector to represent that direction without indicating scale.

Direction cosines are another way to describe a vector’s direction, using trigonometric relationships to find the angles the vector makes with each coordinate axis. For a vector $\mathbf{v} = \langle v_x, v_y, v_z \rangle$, direction cosines are the cosines of the angles α , β , and γ that $\mathbf{v}$ makes with the x -, y -, and z -axes, respectively. They are calculated by dividing each component of the vector by its magnitude, which corresponds to taking the cosine of each directional angle:

$$\cos \alpha = \frac{v_x}{|\mathbf{v}|}, \quad \cos \beta = \frac{v_y}{|\mathbf{v}|}, \quad \cos \gamma = \frac{v_z}{|\mathbf{v}|}.$$

The use of cosines here is rooted in trigonometry: when a vector is projected onto an axis, the cosine of the angle gives the ratio between the length of the projection (the component) and the length of the vector itself. This relationship provides a way to understand how the vector is oriented relative to each axis, which is essential in fields like physics and engineering, where precise direction often matters as much as magnitude.




Figure 1.13: A 2D vector $\mathbf{v}$ showing the angle θ with the x -axis. The direction cosine, $\cos \theta$, is the ratio $\frac{v_x}{|\mathbf{v}|}$, which describes the vector's orientation relative to the x -axis.

Normal Vectors

While direction cosines help us understand how a vector is oriented relative to our coordinate axes, we often need to describe vectors that are perpendicular to other vectors or surfaces. This leads us to the concept of normal vectors.

A *normal vector* (or simply a *normal*) is a vector $\mathbf{n}$ that is perpendicular to something else, like a line or a surface. Normal vectors are especially useful because many problems in physics and geometry involve perpendicular directions. For example, when light hits a mirror, we need to know the direction perpendicular to the mirror's surface to determine how the light will reflect.

For a line in two dimensions, if we know a vector along the line, we can find a normal vector by simply swapping its x and y components and making one of them negative. For example, if a vector along a line is $\mathbf{v} = \langle 3, 4 \rangle$, then $\mathbf{n} = \langle -4, 3 \rangle$ is a normal vector to that line. We can verify this is true because perpendicular vectors have components that, when multiplied and summed, equal zero: $3(-4) + 4(3) = -12 + 12 = 0$.


In polar coordinates, we can find normal vectors in a similar way. For instance, at any point on the circle where $r = 1$, a vector pointing outward from the circle (normal to the circle) has components $\langle \cos \theta, \sin \theta \rangle$. You can visualize this by imagining the normal vector as pointing straight out from the circle at each point, like spokes on a wheel.

Just like any other vector, we can turn a normal vector into a unit normal vector by dividing by its length. Sometimes we denote this like our unit vectors $\hat{\mathbf{n}}$. For instance, in our first example:


$$\hat{\mathbf{n}} = \frac{\langle -4, 3 \rangle}{\sqrt{(-4)^2 + 3^2}} = \frac{\langle -4, 3 \rangle}{5} = \left\langle -\frac{4}{5}, \frac{3}{5} \right\rangle.$$

Unit normal vectors as we have seen are useful because they give us a standard way to describe perpendicular directions, regardless of the size of the original vectors we started

with. We'll see many applications of normal vectors as we continue our study of vectors and their uses in geometry and physics.



(a) Normal vector in Cartesian coordinates



(b) Normal vector on a circle (polar)

Figure 1.14: Normal vectors shown in different coordinate systems. In (a), the normal vector $\mathbf{n}$ is perpendicular to vector $\mathbf{v}$. In (b), the normal vector $\mathbf{n}$ is perpendicular to the radial vector $\mathbf{r}$ at a point on the circle.

▲ **Example 1.3.1** A heavy box is being pushed along a factory floor with a force vector $\mathbf{F} = \langle 9, 12, 15 \rangle$ (measured in newtons) representing the components of the force applied along the x -, y -, and z -axes. Find the magnitude of $\mathbf{F}$, determine the unit vector in the direction of $\mathbf{F}$, calculate the direction cosines of $\mathbf{F}$ with respect to each axis, and find a normal vector to $\mathbf{F}$ that lies in the xy -plane.

Solution To find the unit vector and direction cosines, we first need the magnitude of $\mathbf{F}$.

1. Calculate the magnitude $|\mathbf{F}|$ using the formula:

$$|\mathbf{F}| = \sqrt{F_x^2 + F_y^2 + F_z^2}$$

Substitute $F_x = 9$, $F_y = 12$, and $F_z = 15$:

$$|\mathbf{F}| = \sqrt{9^2 + 12^2 + 15^2} = \sqrt{81 + 144 + 225} = \sqrt{450} = 15\sqrt{2}$$

2. Now, calculate the unit vector $\mathbf{u}$ in the direction of $\mathbf{F}$:

$$\mathbf{u} = \frac{\mathbf{F}}{|\mathbf{F}|} = \left\langle \frac{9}{15\sqrt{2}}, \frac{12}{15\sqrt{2}}, \frac{15}{15\sqrt{2}} \right\rangle$$

Simplify each component:

$$\mathbf{u} = \left\langle \frac{3}{5\sqrt{2}}, \frac{4}{5\sqrt{2}}, \frac{1}{\sqrt{2}} \right\rangle$$

3. Next, we calculate the direction cosines, which are the cosines of the angles that $\mathbf{F}$ makes with each coordinate axis. These are given by:

$$\cos \alpha = \frac{F_x}{|\mathbf{F}|}, \quad \cos \beta = \frac{F_y}{|\mathbf{F}|}, \quad \cos \gamma = \frac{F_z}{|\mathbf{F}|}$$

Substitute the values of F_x , F_y , F_z , and $|\mathbf{F}| = 15\sqrt{2}$:

$$\begin{aligned} \cos \alpha &= \frac{9}{15\sqrt{2}} = \frac{3}{5\sqrt{2}} \\ \cos \beta &= \frac{12}{15\sqrt{2}} = \frac{4}{5\sqrt{2}} \\ \cos \gamma &= \frac{15}{15\sqrt{2}} = \frac{1}{\sqrt{2}} \end{aligned}$$

4. Finally, let's find a normal vector to $\mathbf{F}$ that lies in the xy -plane. We can do this by:

- First, considering only the x and y components of $\mathbf{F}$: $\langle 9, 12 \rangle$
- To find a normal vector in 2D, we can swap these components and make one negative: $\langle -12, 9 \rangle$
- Since we want this normal vector in 3D space lying in the xy -plane, we add a z -component of 0

Therefore, one possible normal vector is $\mathbf{n} = \langle -12, 9, 0 \rangle$.

We can verify this is indeed perpendicular to $\mathbf{F}$ by checking if their dot product is zero:

$$\mathbf{F} \cdot \mathbf{n} = 9(-12) + 12(9) + 15(0) = -108 + 108 + 0 = 0$$

5. Therefore, our complete results are:

- The magnitude of $\mathbf{F}$ is $15\sqrt{2}$ newtons.
- The unit vector in the direction of $\mathbf{F}$ is $\mathbf{u} = \left\langle \frac{3}{5\sqrt{2}}, \frac{4}{5\sqrt{2}}, \frac{1}{\sqrt{2}} \right\rangle$.

- The direction cosines are:

$$\cos \alpha = \frac{3}{5\sqrt{2}}, \quad \cos \beta = \frac{4}{5\sqrt{2}}, \quad \cos \gamma = \frac{1}{\sqrt{2}}$$

- A normal vector to $\mathbf{F}$ in the xy -plane is $\mathbf{n} = \langle -12, 9, 0 \rangle$.
-

Vector Operations

Vectors can be added, subtracted, and scaled, enabling a wide range of applications. Understanding vector operations involves grasping both their mathematical properties and the intuitive idea of combining directions and magnitudes.

Vector Addition and Subtraction

Vector addition is done by adding corresponding components. Given two vectors $\mathbf{u} = \langle u_x, u_y, u_z \rangle$ and $\mathbf{v} = \langle v_x, v_y, v_z \rangle$, their sum $\mathbf{u} + \mathbf{v}$ is calculated by adding each component individually:

$$\mathbf{u} + \mathbf{v} = \langle u_x + v_x, u_y + v_y, u_z + v_z \rangle.$$

Similarly, the difference $\mathbf{u} - \mathbf{v}$ is given by:

$$\mathbf{u} - \mathbf{v} = \langle u_x - v_x, u_y - v_y, u_z - v_z \rangle.$$

The reason we can add vector components directly is that each component describes how far the vector reaches along a particular axis. When we add the components, we're combining these "reaches" along each axis into a single, unified reach, effectively determining where we'd end up if we followed each vector sequentially.

An intuitive way to think about vector addition is with the *tip-to-tail* method. In this method, we place the tail (starting point) of $\mathbf{v}$ at the tip (ending point) of $\mathbf{u}$, without changing either vector's length or direction. This use of *translational invariance* that we previously talked about means the vectors can be moved around in space without changing their meaning, as their direction and magnitude stay the same. The resulting vector, drawn from the starting point of $\mathbf{u}$ to the tip of $\mathbf{v}$, represents the sum $\mathbf{u} + \mathbf{v}$. This tip-to-tail method visually demonstrates how two directions and magnitudes combine to create a new direction and magnitude.




Figure 1.15: *Tip-to-tail method of vector addition: placing $\mathbf{v}$ at the tip of $\mathbf{u}$ results in $\mathbf{u} + \mathbf{v}$.*

Analogy: Tug of War. Consider a classic game of tug-of-war: two teams pull on opposite sides of a rope that has a flag tied in the middle. Each team's force is represented by a vector. If two people on one team each pull with forces represented by vectors $\mathbf{u}$ and $\mathbf{v}$, their combined effort is represented by the sum $\mathbf{u} + \mathbf{v}$, a vector with a magnitude and direction determined by the tip-to-tail addition. If the teams pull at equal effort, the vectors which are the same length, can be combined tip to tail which would go out and then right back to the flag in the middle, meaning the flag won't move.

Now, we can extend this idea: is a three-dimensional tug-of-war possible? Imagine a 3D tug-of-war where three teams pull the rope in different directions. Each team's force is represented by a vector, and the resulting motion of the rope would depend on the sum of all these vectors. In 3D, vector addition still combines each vector's pull, and the rope would move in the direction of the resulting vector. However, the outcome could be complex, with the rope moving in a new diagonal direction rather than towards one "winning" team. This idea illustrates how vector addition generalizes even when multiple forces act in various directions.

We can see where we would place our $\mathbf{u}$ and $\mathbf{v}$ vectors in the below picture:



Figure 1.16: *Tug-of-war as a visual analogy for vector addition.*

We may also consider by itself, a negative vector $-\mathbf{v}$ with the same magnitude but an opposite direction. For a vector $\mathbf{v}$, the negative vector $-\mathbf{v}$ reverses all components:

$$-\mathbf{v} = \langle -v_x, -v_y, -v_z \rangle.$$

Negative vectors are useful in subtraction, as subtracting a vector is the same as adding its negative: $\mathbf{u} - \mathbf{v} = \mathbf{u} + (-\mathbf{v})$.

Scalar Multiplication

Scalar multiplication scales a vector by a constant factor. If c is a scalar and $\mathbf{v} = \langle v_x, v_y, v_z \rangle$, then:

$$c\mathbf{v} = \langle cv_x, cv_y, cv_z \rangle.$$

This means that each component of $\mathbf{v}$ is multiplied by c , effectively stretching or shrinking the vector by a factor of c . If $c > 1$, the vector's magnitude increases; if $0 < c < 1$, it decreases.

For example, doubling the force in a tug-of-war scenario would double each component of the vector representing that force, effectively making the pull twice as strong but in the same direction. Together, vector addition and scalar multiplication allow us to construct complex combinations of vectors and to adjust their magnitudes while maintaining direction, providing a powerful foundation for vector operations in physics and engineering. We may visualize the negative vector and scalar multiplication on the graph below:




Figure 1.17: A vector $\mathbf{v}$ and its negative counterpart $-\mathbf{v}$, along with scalar multiples of $\mathbf{v}$ for various constants.

▲ **Example 1.3.2** An engineer is designing a crane to lift materials by applying forces from multiple cables to achieve the required lift and stability. At a particular moment, two cables exert forces represented by the vectors $\mathbf{F}_1 = \langle 120, 0, 80 \rangle$ and $\mathbf{F}_2 = \langle -50, 75, 60 \rangle$, measured in newtons. To achieve a stable lift, determine the combined force applied by both cables, calculate the unit vector in the direction of the combined force, and find a third vector $\mathbf{G}$ that would need to be added to triple the resultant force from the two cables. Additionally, if we need to fully stabilize the system, calculate the force vector for a third cable $\mathbf{F}_3$ that would completely counterbalance the combined force.

Solution To solve this problem, we'll use vector addition, scalar multiplication, unit vector concepts, and explore how to achieve both triple force and full stability.

1. Combine the Forces by Vector Addition: The combined force vector $\mathbf{F} = \mathbf{F}_1 + \mathbf{F}_2$ is found by adding the components of $\mathbf{F}_1$ and $\mathbf{F}_2$:

$$\mathbf{F} = \langle 120 + (-50), 0 + 75, 80 + 60 \rangle = \langle 70, 75, 140 \rangle$$

2. Calculate the Magnitude of the Combined Force: The magnitude $|\mathbf{F}|$ of the combined force vector $\mathbf{F}$ is given by:

$$|\mathbf{F}| = \sqrt{70^2 + 75^2 + 140^2} = \sqrt{4900 + 5625 + 19600} = \sqrt{30125} \approx 173.6 \text{ newtons}$$

3. Find the Unit Vector in the Direction of the Combined Force: The unit vector $\mathbf{u}$ in the direction of $\mathbf{F}$ is calculated by dividing each component by $|\mathbf{F}|$:

$$\mathbf{u} = \frac{\mathbf{F}}{|\mathbf{F}|} = \left\langle \frac{70}{173.6}, \frac{75}{173.6}, \frac{140}{173.6} \right\rangle \approx \langle 0.403, 0.432, 0.806 \rangle$$

4. Determine the Vector $\mathbf{G}$ to Triple the Combined Force: To triple the resultant force, we need a new vector $\mathbf{G}$ such that the total force becomes $3\mathbf{F}$. This means:

$$\mathbf{G} = 3\mathbf{F} - \mathbf{F} = 2\mathbf{F} = 2 \times \langle 70, 75, 140 \rangle = \langle 140, 150, 280 \rangle$$

Therefore, the vector $\mathbf{G}$ that would need to be applied to triple the combined force is:

$$\mathbf{G} = \langle 140, 150, 280 \rangle$$

5. Determine the Counterbalancing Force Vector $\mathbf{F}_3$: To fully stabilize the system, we need a third cable with force vector $\mathbf{F}_3$ that exactly counterbalances the combined force $\mathbf{F}$. This means $\mathbf{F}_3$ must be equal in magnitude but opposite in direction to $\mathbf{F}$, so:

$$\mathbf{F}_3 = -\mathbf{F} = \langle -70, -75, -140 \rangle$$

Applying this force would bring the net force to zero, stabilizing the system entirely.

Thus, the combined force vector of the two cables is $\langle 70, 75, 140 \rangle$ newtons, with a unit direction vector $\langle 0.403, 0.432, 0.806 \rangle$. To triple this force, a third vector $\mathbf{G} = \langle 140, 150, 280 \rangle$ would be added. Finally, a fourth force vector, cable three, $\mathbf{F}_3 = \langle -70, -75, -140 \rangle$ would stabilize the system by perfectly counterbalancing the combined force.

We've covered addition and subtraction of vectors, which have intuitive meanings in terms of combining directions and magnitudes. However, multiplication raises an important question: how do we interpret multiplying vectors? Here, the dot and cross products offer two distinct perspectives. The dot product provides a scalar, measuring alignment between vectors, while the cross product results in a new vector, perpendicular to both. Simply multiplying components wouldn't capture these meaningful outcomes, so dot and cross products arise naturally as solutions to vector multiplication.

Dot Product

The *dot product* (or scalar product) of two vectors $\mathbf{u} = \langle u_x, u_y, u_z \rangle$ and $\mathbf{v} = \langle v_x, v_y, v_z \rangle$ measures how much one vector “projects” onto another. This projection quantifies how much of one vector's length is aligned in the direction of the other. Mathematically, the dot product is defined as:

$$\mathbf{u} \cdot \mathbf{v} = u_x v_x + u_y v_y + u_z v_z.$$

While this formula looks straightforward, it can be re-expressed to reveal a deeper interpretation by relating it to the angle θ between $\mathbf{u}$ and $\mathbf{v}$:

$$\mathbf{u} \cdot \mathbf{v} = |\mathbf{u}| |\mathbf{v}| \cos \theta.$$

This alternate form shows that the dot product depends on the magnitudes of $\mathbf{u}$ and $\mathbf{v}$ and their alignment. To understand where $\cos \theta$ comes into play, recall that we previously introduced direction cosines to describe a vector's orientation. In the dot product, $\cos \theta$ arises because the product measures how well-aligned two vectors are. When two vectors point in the same direction, $\theta = 0$ (cosine of 0° is 1), and the dot product is simply $|\mathbf{u}| |\mathbf{v}|$. If the vectors are perpendicular (at 90° , or $\pi/2$ radians), $\cos \theta = 0$, and the dot product is zero, meaning there's no overlap in direction.




Figure 1.18: The dot product as the projection of $\mathbf{v}$ onto $\mathbf{u}$, scaled by $\cos \theta$.

Real-World Example: Work Done by a Force. A common example of the dot product is the calculation of work done by a force acting over a displacement. If a force $\mathbf{F}$ is applied to move an object by a displacement $\mathbf{d}$, then the work done W is:

$$W = \mathbf{F} \cdot \mathbf{d} = |\mathbf{F}||\mathbf{d}| \cos \theta.$$

In this expression, only the component of the force that aligns with the displacement contributes to work. To make this intuitive, think about closing a door: if you push on the door in a direction perfectly perpendicular to its plane (where $\theta = 90^\circ$ or $\pi/2$), the force is wasted because $\cos(90^\circ) = 0$, and no work is done to close the door. However, when you push directly along the direction of the door's swing (at $\theta = 0^\circ$), the cosine is maximized, allowing the full force to contribute to moving the door.

Thus, the dot product is valuable because it captures not just magnitude but the idea of alignment, making it essential in scenarios where only the aligned portion of a force (or vector) is relevant to an outcome.

▲ Example 1.3.3 Given vectors $\mathbf{a} = \langle 3, 0, 4 \rangle$ and $\mathbf{b} = \langle 1, 2, -1 \rangle$, calculate their dot product and interpret the result.

Solution The dot product $\mathbf{a} \cdot \mathbf{b}$ is computed as:

$$\mathbf{a} \cdot \mathbf{b} = (3)(1) + (0)(2) + (4)(-1).$$

Calculating each term:

$$\begin{aligned} 3 \cdot 1 &= 3, \\ 0 \cdot 2 &= 0, \\ 4 \cdot (-1) &= -4. \end{aligned}$$

Adding these results, we get:

$$\mathbf{a} \cdot \mathbf{b} = 3 + 0 - 4 = -1.$$

The dot product -1 indicates that $\mathbf{a}$ and $\mathbf{b}$ are not aligned, as a positive result would imply some degree of alignment in the same direction. The negative value suggests that $\mathbf{a}$ and $\mathbf{b}$ point in somewhat opposite directions along their components.

Cross Product

The *cross product* (or vector product) of two vectors $\mathbf{u} = \langle u_x, u_y, u_z \rangle$ and $\mathbf{v} = \langle v_x, v_y, v_z \rangle$ gives a new vector that is perpendicular, or *normal* to both $\mathbf{u}$ and $\mathbf{v}$. The cross product

is defined such that the resulting vector points in a direction determined by the *right-hand rule*: if you point your right hand's index finger along $\mathbf{u}$ and your middle finger along $\mathbf{v}$, then your thumb points in the direction of $\mathbf{u} \times \mathbf{v}$. The magnitude of this vector corresponds to the area of the parallelogram spanned by $\mathbf{u}$ and $\mathbf{v}$, given by:

$$\mathbf{n} = \mathbf{u} \times \mathbf{v} = \langle u_y v_z - u_z v_y, u_z v_x - u_x v_z, u_x v_y - u_y v_x \rangle.$$

An alternative way to calculate the cross product is by using the *determinant* of a 3×3 matrix (maybe you have heard of this in linear algebra before). In this method, we set up a matrix with the unit vectors $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ in the first row, the components of $\mathbf{u}$ in the second row, and the components of $\mathbf{v}$ in the third row:

$$\mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_x & u_y & u_z \\ v_x & v_y & v_z \end{vmatrix}.$$

To compute this determinant, we expand along the first row:

$$\mathbf{u} \times \mathbf{v} = \mathbf{i} \begin{vmatrix} u_y & u_z \\ v_y & v_z \end{vmatrix} - \mathbf{j} \begin{vmatrix} u_x & u_z \\ v_x & v_z \end{vmatrix} + \mathbf{k} \begin{vmatrix} u_x & u_y \\ v_x & v_y \end{vmatrix}.$$

Each 2×2 determinant is calculated as follows:

$$\begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc.$$

Applying this to our cross product:

$$\mathbf{u} \times \mathbf{v} = \mathbf{i}(u_y v_z - u_z v_y) - \mathbf{j}(u_x v_z - u_z v_x) + \mathbf{k}(u_x v_y - u_y v_x).$$

Thus, we arrive at the same result as before:

$$\mathbf{u} \times \mathbf{v} = \langle u_y v_z - u_z v_y, u_z v_x - u_x v_z, u_x v_y - u_y v_x \rangle.$$

Using the determinant method provides a structured way to calculate the cross product and reinforces the concept of orientation in three-dimensional space.

The magnitude of the cross product is given by:

$$|\mathbf{u} \times \mathbf{v}| = |\mathbf{u}||\mathbf{v}|\sin \theta,$$

where θ is the angle between $\mathbf{u}$ and $\mathbf{v}$. This value is maximized when $\mathbf{u}$ and $\mathbf{v}$ are perpendicular (where $\sin \theta = 1$) and zero when the vectors are parallel, indicating no area spanned. The magnitude of the cross product is significant because it represents the area of the parallelogram formed by $\mathbf{u}$ and $\mathbf{v}$.

Geometrically, this area provides a measure of how “spread out” the two vectors are relative to each other. If the vectors are aligned (parallel), there's no area between them, so the magnitude is zero. When the vectors are perpendicular, they span the maximum

possible area, with the magnitude equal to $|\mathbf{u}||\mathbf{v}|$. This concept is especially useful in physics and engineering, where the cross product's magnitude can indicate quantities like torque or rotational force, which depend on both the strength and the orientation of the applied forces. We observe a visualization below:




Figure 1.19: Visualization of the cross product $\mathbf{a} \times \mathbf{b}$, showing the area spanned by vectors $\mathbf{a}$ and $\mathbf{b}$, as well as the normal vector direction $\mathbf{n}$.

▲ **Example 1.3.4** Compute the following cross products:

1. $\langle 1, 5, 7 \rangle \times \langle -4, 3, 3.2 \rangle$
2. $\langle xy^2, z^2, x \rangle \times \langle y, x, z \rangle$

Solution

1. Using the determinant method:

$$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 5 & 7 \\ -4 & 3 & 3.2 \end{vmatrix}$$

$$i = \begin{vmatrix} 5 & 7 \\ 3 & 3.2 \end{vmatrix} = 5 \cdot 3.2 - 7 \cdot 3 = 16 - 21 = -5$$

$$j = - \begin{vmatrix} 1 & 7 \\ -4 & 3.2 \end{vmatrix} = -(1 \cdot 3.2 - 7 \cdot (-4)) = -(3.2 + 28) = -31.2$$

$$k = \begin{vmatrix} 1 & 5 \\ -4 & 3 \end{vmatrix} = 1 \cdot 3 - 5 \cdot (-4) = 3 + 20 = 23$$

$$\langle 1, 5, 7 \rangle \times \langle -4, 3, 3.2 \rangle = \langle -5, -31.2, 23 \rangle$$

2. Using the same method:

$$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ xy^2 & z^2 & x \\ y & x & z \end{vmatrix}$$

$$i = \begin{vmatrix} z^2 & x \\ y & z \end{vmatrix} = z^2 \cdot z - x \cdot y = z^3 - xy$$

$$j = - \begin{vmatrix} xy^2 & x \\ y & z \end{vmatrix} = -(xy^2 \cdot z - x \cdot y) = -(xyz - xy) = xy(1 - z)$$

$$k = \begin{vmatrix} xy^2 & z^2 \\ y & x \end{vmatrix} = xy^2 \cdot x - z^2 \cdot y = x^2y^2 - yz^2$$

$$\langle xy^2, z^2, x \rangle \times \langle y, x, z \rangle = \langle z^3 - xy, xy(1 - z), x^2y^2 - yz^2 \rangle$$

The Right-Hand Rule

The right-hand rule provides an intuitive way to determine the orientation of the cross product vector. As you point your right hand's index finger along the direction of $\mathbf{u}$ and your middle finger along $\mathbf{v}$, your thumb will naturally point in the direction of $\mathbf{u} \times \mathbf{v}$. This orientation represents a unique perpendicular direction relative to both vectors.

Uniqueness and Ambiguity of the Normal Vector. The normal vector $\mathbf{u} \times \mathbf{v}$ is unique in that it provides a consistent perpendicular direction, yet there is also an inherent ambiguity: for any two vectors, two possible perpendicular directions exist. The right-hand rule helps us choose one of these consistently, giving us a convention for which perpendicular direction to use. However, if we were to reverse the order, calculating $\mathbf{v} \times \mathbf{u}$ instead, the resulting vector would point in the opposite direction.

Real-World Example: Torque. In engineering, the cross product is essential in calculating torque, the rotational effect of a force around an axis. If a force $\mathbf{F}$ is applied at a point represented by the position vector $\mathbf{r}$, the torque $\boldsymbol{\tau}$ around the pivot point is:

$$\boldsymbol{\tau} = \mathbf{r} \times \mathbf{F}.$$

Here, the torque vector $\boldsymbol{\tau}$ is perpendicular to the plane formed by $\mathbf{r}$ and $\mathbf{F}$, and its magnitude gives the rotational effect or leverage created by $\mathbf{F}$. This example shows how cross products are useful when interactions involve rotation or orientation.

Geometric Interpretation of Cross Product: The cross product also provides a way to measure the area spanned by two vectors. When we construct a parallelogram with $\mathbf{u}$ and $\mathbf{v}$ as adjacent sides, the magnitude of $\mathbf{u} \times \mathbf{v}$ represents the area of this parallelogram. This interpretation is valuable in physics and engineering applications where the area or spatial orientation between vectors is critical.




Figure 1.20: Completing the parallelogram using the cross product $\mathbf{u} \times \mathbf{v}$, representing the area spanned by vectors $\mathbf{u}$ and $\mathbf{v}$.

Parallelepipeds

A *parallelepiped* is a three-dimensional figure formed by six parallelograms. It is defined by three vectors $\mathbf{u}$, $\mathbf{v}$, and $\mathbf{w}$ that emanate from the same point. The volume of a parallelepiped provides a geometric interpretation of the scalar triple product, which combines the concepts of dot and cross products.

The volume V of a parallelepiped defined by $\mathbf{u}$, $\mathbf{v}$, $\mathbf{w}$ is given by:

$$V = |\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})|.$$

Why this formula works:

- The cross product $\mathbf{v} \times \mathbf{w}$ gives a vector perpendicular to the parallelogram formed by $\mathbf{v}$ and $\mathbf{w}$, with a magnitude equal to the area of this parallelogram.
- Taking the dot product $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})$ projects $\mathbf{u}$ onto the direction of $\mathbf{v} \times \mathbf{w}$, yielding the “height” of the parallelepiped.
- Multiplying the height by the base area (given by the cross product) gives the volume.

Below is a plot of a parallelepiped:




Figure 1.21: A parallelepiped spanned by vectors $\mathbf{u}$, $\mathbf{v}$, $\mathbf{w}$ originating at the origin.

▲ **Example 1.3.5** Given vectors $\mathbf{a} = \langle 3, 0, 4 \rangle$ and $\mathbf{b} = \langle 1, 2, -1 \rangle$, calculate their cross product.

Solution The cross product $\mathbf{a} \times \mathbf{b}$ is computed as:

$$\mathbf{a} \times \mathbf{b} = \langle 0 \cdot (-1) - 4 \cdot 2, 4 \cdot 1 - 3 \cdot (-1), 3 \cdot 2 - 0 \cdot 1 \rangle.$$

Calculating each component:

$$\begin{aligned} x\text{-component} &= 0 \cdot (-1) - 4 \cdot 2 = -8, \\ y\text{-component} &= 4 \cdot 1 - 3 \cdot (-1) = 4 + 3 = 7, \\ z\text{-component} &= 3 \cdot 2 - 0 \cdot 1 = 6. \end{aligned}$$

Thus, $\mathbf{a} \times \mathbf{b} = \langle -8, 7, 6 \rangle$.

Applications of Vectors in Three Dimensions and Higher

Vectors are essential tools across various fields due to their ability to represent both magnitude and direction. Some of the most common applications include:

- In **physics**, vectors describe physical quantities like forces, velocities, and accelerations in three-dimensional space, which helps analyze motion and interactions. For instance, Newton's Second Law of Motion can be written with vectors as:

$$\mathbf{F} = m\mathbf{a},$$

where $\mathbf{F}$ is the force vector, m is the mass, and $\mathbf{a}$ is the acceleration vector. Vectors also appear in equations for gravitational force. For two masses m_1 and m_2 separated

by a distance vector $\mathbf{r}$, the gravitational force is:

$$\mathbf{F}_{\text{gravity}} = -G \frac{m_1 m_2}{|\mathbf{r}|^2} \hat{\mathbf{r}},$$

where G is the gravitational constant, and $\hat{\mathbf{r}}$ is the unit vector in the direction of $\mathbf{r}$. These vector equations help us break down forces into components, making it easier to understand the effects in each direction.

- In **engineering**, vectors represent stresses, displacements, and rotations within structures, which aids in designing stable structures. For example, vectors help engineers analyze how much force a bridge can handle by representing forces and movements in each direction. Stress and strain can also be represented with multi-dimensional vectors or matrices, providing insights into material behavior under different loads.
- In **computer graphics**, vectors are crucial for simulating lighting, shading, and movement in three-dimensional environments. For example, when calculating the lighting on a surface, we use the *dot product* between the light direction vector $\mathbf{L}$ and the surface normal vector $\mathbf{N}$ to find the light's intensity on that surface:

$$I = I_{\text{light}} \cdot \max(0, \mathbf{L} \cdot \mathbf{N}),$$

where I is the resulting intensity, and I_{light} is the light source intensity. This helps create realistic shading by adjusting brightness based on the angle between the light and the surface.

Vectors are also powerful for thinking beyond physical three-dimensional space. In fields like machine learning and quantum mechanics, vectors allow us to work in higher-dimensional, abstract spaces:

- In **machine learning**, vectors represent data points with multiple features. For example, if each data point has 100 features, we can think of it as a vector in a 100-dimensional space. By measuring distances between these vectors, we can find similarities between data points. For instance, the Euclidean distance between two data points $\mathbf{x}$ and $\mathbf{y}$ is:

$$\text{Distance}(\mathbf{x}, \mathbf{y}) = \sqrt{\sum_{i=1}^n (x_i - y_i)^2},$$

where $\mathbf{x}$ and $\mathbf{y}$ are vectors representing two data points. The dot product is also used in machine learning algorithms like support vector machines to classify data by finding directions that separate categories. Neural networks rely heavily on matrix-vector multiplication, which is an extension of vector operations, to process and transform data through layers.

- In **quantum mechanics**, vectors represent quantum states in complex, high-dimensional spaces. For instance, a basic quantum unit, a qubit, can be represented by a vector in a two-dimensional complex space:

$$|\psi\rangle = \alpha|0\rangle + \beta|1\rangle,$$

where α and β are complex numbers satisfying $|\alpha|^2 + |\beta|^2 = 1$. The dot product (or inner product) between two quantum state vectors gives probabilities for measurement outcomes, which is central to quantum calculations.

These examples show that vectors are more than just arrows in space. They allow us to work with quantities that have both size and direction, whether we're studying forces in physics or organizing data in machine learning. By using vectors, we can extend our understanding into higher-dimensional spaces, making them indispensable tools in both theoretical and applied fields.

▲ **Example 1.3.6** Suppose we have two products represented by feature vectors:

$$\mathbf{p} = \langle 8, 2, 15 \rangle \quad \text{and} \quad \mathbf{q} = \langle 5, 3, 12 \rangle,$$

where each vector component represents the popularity, weight (in kg), and price (in dollars) of each product, respectively. In machine learning, we often use vector operations like the dot product and cross product to analyze relationships between data points (in this case, the products). Let's calculate:

1. The dot product $\mathbf{p} \cdot \mathbf{q}$ to measure the similarity between the products in terms of their features.
2. The cross product $\mathbf{p} \times \mathbf{q}$ to analyze how the products differ across the feature space.

Solution We'll go through each calculation step-by-step.

1. Dot Product:

The dot product is useful for assessing similarity between vectors. When comparing products, a higher dot product value suggests that the products share similar feature characteristics. Here, we calculate the dot product as:

$$\mathbf{p} \cdot \mathbf{q} = (8)(5) + (2)(3) + (15)(12) = 40 + 6 + 180 = 226.$$

In this case, a dot product of 226 implies moderate alignment between the products' popularity, weight, and price. In a recommendation system, for example, a higher dot product might indicate that two products could appeal to similar customers.

2. Cross Product:

The cross product results in a new vector perpendicular to both $\mathbf{p}$ and $\mathbf{q}$. In machine learning, this can help identify unique directions in the feature space where

the products differ significantly, which can be useful for clustering or classification tasks to understand how data points vary in multiple dimensions. The cross product is calculated as:

$$\mathbf{p} \times \mathbf{q} = \langle 2 \cdot 12 - 15 \cdot 3, 15 \cdot 5 - 8 \cdot 12, 8 \cdot 3 - 2 \cdot 5 \rangle.$$

Calculating each component:

$$x\text{-component} = (2)(12) - (15)(3) = 24 - 45 = -21,$$

$$y\text{-component} = (15)(5) - (8)(12) = 75 - 96 = -21,$$

$$z\text{-component} = (8)(3) - (2)(5) = 24 - 10 = 14.$$

Thus, the cross product $\mathbf{p} \times \mathbf{q} = \langle -21, -21, 14 \rangle$.

This cross product vector, $\langle -21, -21, 14 \rangle$, highlights the directional differences between the products. In some machine learning applications, such as Principal Component Analysis (PCA), understanding these directions helps in finding dimensions that best separate or distinguish data points.

Summary: In this example:

- The dot product of 226 suggests that the products have moderate similarity in their features. This measure could help a recommendation algorithm group these items as similar.
- The cross product $\langle -21, -21, 14 \rangle$ represents the direction of variance between these products, useful for exploring the distinctiveness of feature combinations in high-dimensional analysis.

The following Python programming example demonstrates how dot and cross products can be applied in machine learning to measure similarity and analyze differences between feature vectors of products. Dot products are used here as a basic similarity score, while cross products provide insight into directional differences across features – much like what we have done in the previous example. Students can run the following Python code to experiment with these concepts in a real-world context:

```
1 import numpy as np
2
3 # Feature vectors for two products: [popularity, weight (kg), price (dollars)]
4 p = np.array([8, 2, 15])
5 q = np.array([5, 3, 12])
6
7 # Calculate the dot product
8 dot_product = np.dot(p, q)
```

```
9 print(f"Dot Product (Similarity Score): {dot_product}")
10
11 # Calculate the cross product
12 cross_product = np.cross(p, q)
13 print(f"Cross Product (Directional Difference): {cross_product}")
14
15 # Use dot product as a basic similarity measure for recommendation
16 similarity_threshold = 200 # Set an arbitrary threshold
17 if dot_product > similarity_threshold:
18     print("Products are similar enough to recommend together.")
19 else:
20     print("Products are distinct enough to consider separately.")
21
22 # Analyze uniqueness with cross product
23 if np.linalg.norm(cross_product) > 0:
24     print("Products have unique directional attributes across feature space.")
25 else:
26     print("Products are very similar across all feature dimensions.")
```

Lecture 1.4: Lines and Planes in Space

As we extend our understanding of vectors and coordinate systems from two dimensions to three, we encounter new challenges and possibilities. In two-dimensional space, we could represent points, lines, and directions relatively simply. But as we move into three dimensions, we need to adapt our definitions to handle the additional complexity of this expanded space. Naturally, we begin asking questions like: *How do we represent a line in 3D? How do we define a flat surface, or plane, that extends in multiple directions in this space?*

Using what we know about vectors and coordinate systems, we can build definitions for lines and planes in 3D that feel like natural extensions of the concepts we used in 2D. In 2D, a line could be defined by a point and a direction (often represented by a slope). In 3D, however, a line requires both a point for positioning and a vector to set its direction in this more complex space. Similarly, defining a plane in 3D requires a point and a vector that establishes its orientation. Developing these definitions is key to understanding and navigating three-dimensional space.

Equations of Lines and Line Segments

In mathematics, we often represent points and vectors with similar symbols, but it is crucial to distinguish between them as they serve different purposes in geometry. A point, denoted as P , represents a specific location in space, typically described by its coordinates (x, y, z) . For instance, when we write $P = (x_0, y_0, z_0)$, we are explicitly identifying a unique position in space.

On the other hand, when we use bold notation such as $\mathbf{P}$ (e.g., $\mathbf{P}_0$ or $\mathbf{P}_1$), we are explicitly treating it as a **position vector**. A position vector originates from the origin and “points” to the coordinates of P . In this case, $\mathbf{P} = \langle x, y, z \rangle$ represents a vector that extends from the origin to the location (x, y, z) . When a point (x, y, z) is to be used in vector calculations, it is written as $\mathbf{P}$. This explicit use of bold notation highlights that the point is being treated as a vector, enabling operations such as addition, subtraction, and scaling to describe directions, magnitudes, and displacements in space.

To summarize:

- P refers to a specific point at coordinates (x, y, z) , which identifies a location.
- $\mathbf{P} = \langle x, y, z \rangle$ explicitly represents the position vector pointing from the origin to P .
- When a point is to be used as a vector in calculations, we write it as $\mathbf{P}$ to make its vectorial role explicit.

In this context, $\mathbf{P}$ will always be used when working with vectors in calculations, ensuring clarity and consistency, while P is reserved to identify specific points independent of vectors. By maintaining this distinction, we make the roles of points and vectors explicit in both notation and interpretation.

Defining a Line in 3D Space

To describe a line in three-dimensional space, we need:

- A specific **point** through which the line passes, denoted $\mathbf{P}_0 = (x_0, y_0, z_0)$.
- A **direction vector** that defines the orientation of the line, represented by $\mathbf{d} = \langle a, b, c \rangle$.

The point $\mathbf{P}_0$ anchors the line at a specific position, while $\mathbf{d}$ describes how the line extends in space. By combining these, we can reach any point along the line.

The Parametric Equation of a Line

To generate all points on a line that passes through $\mathbf{P}_0$ with direction vector $\mathbf{d}$, we introduce a **parameter** t that scales $\mathbf{d}$. The parametric equation of the line becomes:

$$\mathbf{r}(t) = \mathbf{P}_0 + t\mathbf{d} = \langle x_0 + at, y_0 + bt, z_0 + ct \rangle,$$

where $\mathbf{r}(t)$ represents the position vector of any point on the line. Breaking down this equation:

- When $t = 0$, $\mathbf{r}(0) = \mathbf{P}_0$, locating us at the point $\mathbf{P}_0$.
- As t varies, we move along the line in the direction of $\mathbf{d}$.

Alternatively, we can express the parametric form as a set of equations:

$$\begin{cases} x = x_0 + at, \\ y = y_0 + bt, \\ z = z_0 + ct, \end{cases} \quad t \in I,$$

where I is the interval for t that defines the line or line segment.

Line Segments

If we want only a segment of the line connecting two points $\mathbf{P}_1 = (x_1, y_1, z_1)$ and $\mathbf{P}_2 = (x_2, y_2, z_2)$, we define the line segment using t in the range $[0, 1]$.

For $t \in [0, 1]$, the position vector $\mathbf{r}(t)$ of any point on the line segment is:

$$\mathbf{r}(t) = (1 - t)\mathbf{P}_1 + t\mathbf{P}_2 = (1 - t)\langle x_1, y_1, z_1 \rangle + t\langle x_2, y_2, z_2 \rangle.$$

Expanding this:

$$\mathbf{r}(t) = \langle (1 - t)x_1 + tx_2, (1 - t)y_1 + ty_2, (1 - t)z_1 + tz_2 \rangle.$$

This form describes points on the segment:

- When $t = 0$, $\mathbf{r}(0) = \mathbf{P}_1 = \langle x_1, y_1, z_1 \rangle$.

- When $t = 1$, $\mathbf{r}(1) = \mathbf{P}_2 = \langle x_2, y_2, z_2 \rangle$.
- For $t \in (0, 1)$, $\mathbf{r}(t)$ lies between $\mathbf{P}_1$ and $\mathbf{P}_2$, tracing the line segment.

This approach uses vector notation to generalize points along the line segment, treating each location as a vector in 3D space.




Figure 1.22: The line segment connecting points $\mathbf{P}_1$ and $\mathbf{P}_2$ in three-dimensional space, represented by $\mathbf{r}(t) = (1-t)\mathbf{P}_1 + t\mathbf{P}_2$ for $0 \leq t \leq 1$. Here, $\mathbf{P}_1$ and $\mathbf{P}_2$ represent the positions of the endpoints, while $\mathbf{P}_1$ and $\mathbf{P}_2$ denote their position vectors.

Why Use a Line Segment? Line segments are useful for representing finite parts of lines, such as beams in a structure or paths between points. In many applications, we don't need the entire line, just a specific portion. In summary, by combining a point with a direction vector, we can describe a line in 3D space. The parametric form gives us flexibility in describing all points on the line and any specific segment of it. We will revisit parametric equations many more times in our study of multi-variate functions.

▲ **Example 1.4.1** Suppose we want to find the equation of a line that passes through a point $\mathbf{P}_0 = (1, -2, 3)$ and points in the direction of a vector $\mathbf{d} = \langle 4, -1, 2 \rangle$. Please also describe the line segment between two points, $\mathbf{P}_1 = (1, -2, 3)$ and $\mathbf{P}_2 = (5, -3, 5)$.

Solution

1. Equation of the Line:

To represent the line passing through $\mathbf{P}_0$ in the direction of $\mathbf{d}$, we use the parametric

form:

$$\mathbf{r}(t) = \mathbf{P}_0 + t\mathbf{d} = \langle 1, -2, 3 \rangle + t\langle 4, -1, 2 \rangle.$$

Expanding this, we get:

$$\mathbf{r}(t) = \langle 1 + 4t, -2 - t, 3 + 2t \rangle.$$

In curly bracket notation, this can be represented as:

$$\begin{cases} x = 1 + 4t, \\ y = -2 - t, \\ z = 3 + 2t \end{cases} \quad \text{for } t \in \mathbb{R}.$$

These equations describe any point on the line by varying t .

2. Equation of the Line Segment between $\mathbf{P}_1$ and $\mathbf{P}_2$:

For a line segment connecting $\mathbf{P}_1 = (1, -2, 3)$ and $\mathbf{P}_2 = (5, -3, 5)$, we restrict t to the interval $0 \leq t \leq 1$. We can describe this line segment with the following equation:

$$\mathbf{r}(t) = (1 - t)\mathbf{P}_1 + t\mathbf{P}_2 = (1 - t)\langle 1, -2, 3 \rangle + t\langle 5, -3, 5 \rangle.$$

Expanding this, we get:

$$\mathbf{r}(t) = \langle (1 - t) \cdot 1 + t \cdot 5, (1 - t) \cdot (-2) + t \cdot (-3), (1 - t) \cdot 3 + t \cdot 5 \rangle = \langle 1 + 4t, -2 - t, 3 + 2t \rangle.$$

In curly bracket notation, this line segment can be written as:

$$\begin{cases} x = 1 + 4t, \\ y = -2 - t, \\ z = 3 + 2t \end{cases} \quad \text{for } t \in [0, 1].$$

This form of $\mathbf{r}(t)$ represents points on the line segment:

- When $t = 0$, $\mathbf{r}(0) = \mathbf{P}_1 = \langle 1, -2, 3 \rangle$.
- When $t = 1$, $\mathbf{r}(1) = \mathbf{P}_2 = \langle 5, -3, 5 \rangle$.
- For values of t between 0 and 1, $\mathbf{r}(t)$ gives intermediate points along the line segment from $\mathbf{P}_1$ to $\mathbf{P}_2$.

Summary: The parametric form $\mathbf{r}(t) = \langle 1 + 4t, -2 - t, 3 + 2t \rangle$ defines a line through $P_0 = (1, -2, 3)$ in the direction $\mathbf{d} = \langle 4, -1, 2 \rangle$. Restricting t to $[0, 1]$ describes the specific line segment between $P_1 = (1, -2, 3)$ and $P_2 = (5, -3, 5)$.

Equations of Planes

After defining lines, a natural next question arises: *How can we define a flat surface in three-dimensional space?* While a line gives us a one-dimensional path in space, a plane is a natural extension of this concept, representing a two-dimensional surface that can extend infinitely in all directions within that surface. Just as a line in 3D requires a direction vector to establish its orientation, a plane requires a way to “anchor” its orientation, allowing it to stretch out in multiple directions while still remaining flat.

To define a plane, we need:

- A point on the plane, $P_0 = (x_0, y_0, z_0)$.
- A **normal vector** $\mathbf{n} = \langle A, B, C \rangle$ that is perpendicular to every direction within the plane.

Why a Normal Vector? The normal vector is crucial because it generalizes the idea of direction from lines to planes. While a line has a single direction vector that guides its path, a plane requires a unique direction that can “lock” its orientation in space. Let’s take a look:




Figure 1.23: *Illustration of a plane in 3D space defined by point P_0 , with the normal vector $\mathbf{n}$ extending from P_0 in the perpendicular direction.*

The normal vector provides this by pointing perpendicularly to the plane, creating a stable orientation despite the plane extending infinitely in two directions. With these components, the equation of the plane can be written as:

$$A(x - x_0) + B(y - y_0) + C(z - z_0) = 0.$$

Expanding this, we get:

$$Ax + By + Cz = D,$$

where $D = Ax_0 + By_0 + Cz_0$. This form represents all points (x, y, z) on the plane, as they satisfy this equation by remaining perpendicular to $\mathbf{n}$.

▲ **Example 1.4.2** Suppose we want to find the equation of a plane that passes through the point $P_0 = (2, -1, 3)$ and is oriented in space by a normal vector $\mathbf{n} = \langle 4, 5, -2 \rangle$.

Solution

To define the equation of the plane, we use the general form:

$$A(x - x_0) + B(y - y_0) + C(z - z_0) = 0,$$

where $P_0 = (x_0, y_0, z_0)$ is a point on the plane and $\mathbf{n} = \langle A, B, C \rangle$ is the normal vector.

1. Substitute Point and Normal Vector Values:

For this example, $P_0 = (2, -1, 3)$ and $\mathbf{n} = \langle 4, 5, -2 \rangle$, so we substitute $A = 4$, $B = 5$, and $C = -2$ as well as $x_0 = 2$, $y_0 = -1$, and $z_0 = 3$:

$$4(x - 2) + 5(y + 1) - 2(z - 3) = 0.$$

2. Expand and Simplify the Equation:

Expanding each term:

$$4x - 8 + 5y + 5 - 2z + 6 = 0.$$

Combining like terms, we obtain:

$$4x + 5y - 2z = -3.$$

Interpretation: This equation describes a plane passing through the point $(2, -1, 3)$ and oriented by the normal vector $\langle 4, 5, -2 \rangle$. The plane extends infinitely in two directions, and every point (x, y, z) on this plane maintains a perpendicular relationship to the normal vector $\mathbf{n}$, aligning with the orientation given by $\mathbf{n}$.

Intersections and Angles between Lines and Planes

Now that we have equations for lines and planes, it's natural to explore how these objects interact in space. Specifically, we are interested in points of intersection and the angles between lines and planes.

Intersection of a Line and a Plane

To determine if and where a line intersects a plane, we substitute the parametric form of the line into the plane equation. If there's a solution for t , the line intersects the plane at that point; otherwise, the line is parallel to the plane. Observe the below visual:




Figure 1.24: A line intersecting a tilted plane at a single point in three-dimensional space.

Consider a line with parametric equation:

$$\mathbf{r}(t) = \mathbf{P}_0 + t\mathbf{d} = \langle x_0, y_0, z_0 \rangle + t\langle a, b, c \rangle,$$

and a plane with equation:

$$A(x - x_1) + B(y - y_1) + C(z - z_1) = 0.$$

By substituting the coordinates of $\mathbf{r}(t)$ into the plane equation, we get:

$$A(x_0 + at - x_1) + B(y_0 + bt - y_1) + C(z_0 + ct - z_1) = 0.$$

This equation simplifies to a linear equation in t . If a solution exists, it gives the value of t at the intersection point. If no solution exists, the line is parallel to the plane.

▲ **Example 1.4.3** Suppose we have a line passing through the point $\mathbf{P}_0 = (1, 2, 3)$ in the direction of the vector $\mathbf{d} = \langle 2, -1, 1 \rangle$, and a plane given by the equation $3x - y + 2z = 5$. We want to find the point of intersection between this line and the plane, if it exists.

Solution

1. Set Up the Line Equation:

The parametric form of the line is:

$$\mathbf{r}(t) = \mathbf{P}_0 + t\mathbf{d} = (1, 2, 3) + t\langle 2, -1, 1 \rangle.$$

Expanding this, we get:

$$\mathbf{r}(t) = \langle 1 + 2t, 2 - t, 3 + t \rangle.$$

So the coordinates of any point on the line are:

$$x = 1 + 2t, \quad y = 2 - t, \quad z = 3 + t.$$

2. Substitute into the Plane Equation:

The plane equation is:

$$3x - y + 2z = 5.$$

Substitute $x = 1 + 2t$, $y = 2 - t$, and $z = 3 + t$ into this equation:

$$3(1 + 2t) - (2 - t) + 2(3 + t) = 5.$$

Expanding and simplifying:

$$3 + 6t - 2 + t + 6 + 2t = 5.$$

Combine like terms:

$$9t + 7 = 5.$$

Solving for t :

$$t = -\frac{2}{9}.$$

3. Find the Intersection Point:

Substitute $t = -\frac{2}{9}$ back into the parametric equations of the line:

$$x = 1 + 2\left(-\frac{2}{9}\right) = 1 - \frac{4}{9} = \frac{5}{9},$$

$$y = 2 - \left(-\frac{2}{9}\right) = 2 + \frac{2}{9} = \frac{20}{9},$$

$$z = 3 + \left(-\frac{2}{9}\right) = 3 - \frac{2}{9} = \frac{25}{9}.$$

Thus, the intersection point is:

$$\left(\frac{5}{9}, \frac{20}{9}, \frac{25}{9}\right).$$

Conclusion: The line intersects the plane at the point $\left(\frac{5}{9}, \frac{20}{9}, \frac{25}{9}\right)$.

Line of Intersection Between Two Planes

When two planes intersect in three-dimensional space, they do so along a line. However, there is a specific condition for this to occur: *the planes must not be parallel*. When two planes are parallel, their normal vectors are either equal or scalar multiples of each other, meaning they have no intersection or they coincide completely.




Figure 1.25: The intersection of two planes forming a line in three-dimensional space. The red and blue shaded regions represent the two planes, with their line of intersection shown as the thick black line.

For two non-parallel planes, we can determine the line of intersection by using the following systematic approach:

1. **Represent each plane with an equation:** The general equation of a plane is

$$A_1x + B_1y + C_1z + D_1 = 0$$

for the first plane, and

$$A_2x + B_2y + C_2z + D_2 = 0$$

for the second plane. Here, A_1, B_1, C_1 and A_2, B_2, C_2 are the components of the normal vectors of the planes, $\mathbf{n}_1 = \langle A_1, B_1, C_1 \rangle$ and $\mathbf{n}_2 = \langle A_2, B_2, C_2 \rangle$, respectively.

2. **Understand why these planes intersect along a line:** Since $\mathbf{n}_1$ and $\mathbf{n}_2$ are not scalar multiples of each other, the planes are not parallel, meaning they must intersect. Geometrically, if two planes intersect, they do so along a line. Intuitively, this is because two planes cutting across space create a shared set of points, forming a one-dimensional path—the line of intersection.
3. **Find the direction vector of the line of intersection:** The line of intersection lies along a direction that is perpendicular to both normal vectors $\mathbf{n}_1$ and $\mathbf{n}_2$. To find this direction, we compute the cross product of the normal vectors:

$$\mathbf{d} = \mathbf{n}_1 \times \mathbf{n}_2.$$

Since the cross product of two vectors is perpendicular to both, $\mathbf{d}$ serves as the direction vector of the intersection line.

4. **Find a specific point on the line:** To fully define the line, we need a point through which it passes. To find such a point, we solve the system of equations given by the two plane equations:

$$A_1x + B_1y + C_1z = -D_1,$$

$$A_2x + B_2y + C_2z = -D_2.$$

We can solve this system by eliminating one of the variables by setting it equal to 0 and finding values for two of x , y , and z that satisfy both equations. This gives us a point $\mathbf{P}_0 = (x_0, y_0, z_0)$ on the line.

5. **Write the parametric equations of the line:** With a direction vector $\mathbf{d} = \langle d_x, d_y, d_z \rangle$ and a point $\mathbf{P}_0 = (x_0, y_0, z_0)$ on the line, the parametric form of the line of intersection can be expressed as:

$$\mathbf{r}(t) = \mathbf{P}_0 + t\mathbf{d} = \langle x_0, y_0, z_0 \rangle + t\langle d_x, d_y, d_z \rangle.$$

Expanding, we get:

$$\mathbf{r}(t) = \langle x_0 + td_x, y_0 + td_y, z_0 + td_z \rangle.$$

In curly bracket notation, this is:

$$\begin{cases} x = x_0 + td_x, \\ y = y_0 + td_y, \\ z = z_0 + td_z \end{cases} \quad \text{for } t \in \mathbb{R}.$$

This derivation shows that the intersection of two planes, when they are not parallel, is always a line. By solving the system of equations to find a point on the intersection and calculating the cross product of the planes' normal vectors, we can fully define the line of intersection in space.

Angles Between Planes

To understand the angle between two planes, we look at the angle between their normal vectors. Given two planes with normal vectors $\mathbf{n}_1$ and $\mathbf{n}_2$, we can think of the angle θ between the planes as the angle by which their normal vectors are “tilted” relative to each other. This makes sense because if the planes are tilted, so are their normal vectors.

To find this angle, we use the dot product of the normal vectors, which is related to the cosine of the angle between them:

$$\mathbf{n}_1 \cdot \mathbf{n}_2 = |\mathbf{n}_1||\mathbf{n}_2| \cos \theta.$$

Rearranging, we solve for $\cos \theta$ to find the angle between the planes:

$$\cos \theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1||\mathbf{n}_2|}.$$

This expression gives us a clear and direct way to find θ .

Special cases arise in two situations:

- If $\cos \theta = 0$, then $\theta = 90^\circ$, meaning the planes are **perpendicular**.
- If $\cos \theta = \pm 1$, then $\theta = 0^\circ$ or $\theta = 180^\circ$, meaning the planes are **parallel**.

Angle Between a Line and a Plane

For a line intersecting a plane, the angle ϕ measures how “steeply” the line intersects the plane. Intuitively, we define this angle using the direction vector $\mathbf{d}$ of the line and the normal vector $\mathbf{n}$ of the plane. To find this angle, we consider the following:

- The angle θ between $\mathbf{d}$ and $\mathbf{n}$ represents how much the line “tilts” away from being perpendicular to the plane.
- Since the angle between the line and the plane is complementary to θ , we have $\phi = 90^\circ - \theta$.

Therefore, using the relationship for $\cos \theta$ with the dot product, we find:

$$\sin \phi = \frac{\mathbf{d} \cdot \mathbf{n}}{|\mathbf{d}||\mathbf{n}|} = \cos \theta.$$

This formula allows us to calculate ϕ based on the alignment of the line’s direction vector with the plane’s normal vector. Special cases occur as follows:

- If $\phi = 0^\circ$, the line is **parallel** to the plane.
- If $\phi = 90^\circ$, the line is **perpendicular** to the plane.

The formulas for angles between planes and between a line and a plane provide a structured approach to understanding their spatial relationships and orientation in three-dimensional space.

▲ **Example 1.4.4** Consider two planes given by the equations:

$$\text{Plane 1: } 2x - y + z - 3 = 0,$$

$$\text{Plane 2: } x + y - 2z + 4 = 0.$$

A line L is the intersection of these two planes. Find:

1. The angle θ between the two planes.
2. The parametric equations of the line L of intersection.
3. The angle ϕ between the line L and Plane 1.

Solution

1. Angle between the two planes:

The normal vector of a plane can be derived from its equation in the form $Ax + By + Cz + D = 0$ as $\mathbf{n} = \langle A, B, C \rangle$.

For Plane 1, $2x - y + z - 3 = 0$, so the normal vector is:

$$\mathbf{n}_1 = \langle 2, -1, 1 \rangle.$$

For Plane 2, $x + y - 2z + 4 = 0$, so the normal vector is:

$$\mathbf{n}_2 = \langle 1, 1, -2 \rangle.$$

Compute the dot product of the normal vectors:

$$\mathbf{n}_1 \cdot \mathbf{n}_2 = (2)(1) + (-1)(1) + (1)(-2) = 2 - 1 - 2 = -1.$$

Calculate the magnitudes:

$$|\mathbf{n}_1| = \sqrt{2^2 + (-1)^2 + 1^2} = \sqrt{4 + 1 + 1} = \sqrt{6},$$

$$|\mathbf{n}_2| = \sqrt{1^2 + 1^2 + (-2)^2} = \sqrt{1 + 1 + 4} = \sqrt{6}.$$

Compute $\cos \theta$:

$$\cos \theta = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{|\mathbf{n}_1||\mathbf{n}_2|} = \frac{-1}{\sqrt{6} \times \sqrt{6}} = \frac{-1}{6}.$$

Therefore, the angle θ between the planes is:

$$\theta = \arccos \left(\frac{-1}{6} \right) \approx 99.6^\circ.$$

2. Parametric equations of the line L :

Follow these steps systematically to find the line of intersection:

(a) Compute the direction vector $\mathbf{d}$:

The direction vector $\mathbf{d}$ of the line of intersection is perpendicular to both $\mathbf{n}_1$ and $\mathbf{n}_2$. We find it by taking the cross product:

$$\mathbf{d} = \mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -1 & 1 \\ 1 & 1 & -2 \end{vmatrix}.$$

Expanding the determinant:

$$\mathbf{d} = \mathbf{i}((-1)(-2) - (1)(1)) - \mathbf{j}((2)(-2) - (1)(1)) + \mathbf{k}((2)(1) - (-1)(1)).$$

Simplify each term:

$$\mathbf{d} = \mathbf{i}(2 - 1) - \mathbf{j}(-4 - 1) + \mathbf{k}(2 + 1) = \langle 1, 5, 3 \rangle.$$

(b) Find a point on the line:

We need a specific point $\mathbf{P}_0 = (x_0, y_0, z_0)$ that lies on both planes. Set $z = 0$ to simplify calculations, and solve for x and y using the system:

$$\begin{cases} 2x - y = 3, \\ x + y = -4. \end{cases}$$

From the second equation, solve for y :

$$y = -4 - x.$$

Substitute into the first equation:

$$2x - (-4 - x) = 3 \implies 2x + 4 + x = 3 \implies 3x = -1 \implies x = -\frac{1}{3}.$$

Substitute $x = -\frac{1}{3}$ back into $y = -4 - x$:

$$y = -4 + \frac{1}{3} = -\frac{12}{3} + \frac{1}{3} = -\frac{11}{3}.$$

So, a point on the line is $\mathbf{P}_0 = \left(-\frac{1}{3}, -\frac{11}{3}, 0\right)$.

(c) **Write the parametric equations of the line:**

With $\mathbf{P}_0 = (-\frac{1}{3}, -\frac{11}{3}, 0)$ and $\mathbf{d} = \langle 1, 5, 3 \rangle$, the parametric equations of the line L are:

$$\mathbf{r}(t) = \mathbf{P}_0 + t\mathbf{d} = \left\langle -\frac{1}{3}, -\frac{11}{3}, 0 \right\rangle + t\langle 1, 5, 3 \rangle.$$

Expanding, we get:

$$\mathbf{r}(t) = \left\langle -\frac{1}{3} + t, -\frac{11}{3} + 5t, 3t \right\rangle.$$

In curly bracket notation, this can be written as:

$$\begin{cases} x = -\frac{1}{3} + t, \\ y = -\frac{11}{3} + 5t, \\ z = 3t \end{cases} \quad \text{for } t \in \mathbb{R}.$$

3. Angle between line L and Plane 1:

The normal vector of Plane 1 is $\mathbf{n}_1 = \langle 2, -1, 1 \rangle$ (from part (a)).

Compute the dot product of $\mathbf{d}$ and $\mathbf{n}_1$:

$$\mathbf{d} \cdot \mathbf{n}_1 = (1)(2) + (5)(-1) + (3)(1) = 2 - 5 + 3 = 0.$$

Since the dot product is zero, $\sin \phi = 0$ and $\phi = \arcsin(0) = 0^\circ$.

This result means that the line L is parallel to Plane 1 (technically it lies in the plane itself), as expected for the line of intersection between two planes.

Projections of Vectors

When studying vectors in three-dimensional space, it's often useful to decompose a vector along another vector or within a plane. This breakdown allows us to understand how one vector aligns with another direction or lies within a given surface. A *projection* is the component of a vector along a specified direction. Given a vector $\mathbf{a}$, the *projection of $\mathbf{a}$ onto $\mathbf{b}$* gives us the part of $\mathbf{a}$ that aligns with $\mathbf{b}$. The projection of $\mathbf{a}$ onto $\mathbf{b}$, written as $\text{proj}_{\mathbf{b}}\mathbf{a}$, is calculated by:

$$\text{proj}_{\mathbf{b}}\mathbf{a} = \frac{\mathbf{a} \cdot \mathbf{b}}{|\mathbf{b}|^2} \mathbf{b}.$$

This formula tells us how much of $\mathbf{a}$ points in the direction of $\mathbf{b}$.




Figure 1.26: Vector $\mathbf{a}$ and its projection onto the vector $\mathbf{n}$. The red dashed vector is the projection of $\mathbf{a}$ onto $\mathbf{n}$, while the green vector represents the component of $\mathbf{a}$ that is perpendicular to $\mathbf{n}$.

Projection of a Vector onto a Plane.

If we have a plane defined by a *normal vector* $\mathbf{n}$, we can project any vector $\mathbf{a}$ onto the plane to find its component that lies within the plane. This is done by subtracting the projection of $\mathbf{a}$ onto $\mathbf{n}$:

$$\mathbf{a}_{\parallel \text{plane}} = \mathbf{a} - \text{proj}_{\mathbf{n}} \mathbf{a}.$$

This vector $\mathbf{a}_{\parallel \text{plane}}$ represents the part of $\mathbf{a}$ that lies parallel to the plane.

▲ **Example 1.4.5** Suppose we have a vector $\mathbf{a} = \langle 3, 4, 5 \rangle$ and a normal vector $\mathbf{n} = \langle 1, 1, 1 \rangle$. Calculate the projection of $\mathbf{a}$ onto $\mathbf{n}$ and the component of $\mathbf{a}$ that lies within the plane perpendicular to $\mathbf{n}$.

Solution

1. Projection of $\mathbf{a}$ onto $\mathbf{n}$:

The projection of $\mathbf{a}$ onto $\mathbf{n}$ is calculated using:

$$\text{proj}_{\mathbf{n}} \mathbf{a} = \frac{\mathbf{a} \cdot \mathbf{n}}{|\mathbf{n}|^2} \mathbf{n}.$$

First, compute $\mathbf{a} \cdot \mathbf{n}$:

$$\mathbf{a} \cdot \mathbf{n} = (3)(1) + (4)(1) + (5)(1) = 3 + 4 + 5 = 12.$$

Then, calculate $|\mathbf{n}|^2$:

$$|\mathbf{n}|^2 = (1)^2 + (1)^2 + (1)^2 = 3.$$

Thus:

$$\text{proj}_{\mathbf{n}}\mathbf{a} = \frac{12}{3}\mathbf{n} = 4\langle 1, 1, 1 \rangle = \langle 4, 4, 4 \rangle.$$

2. Component of $\mathbf{a}$ in the Plane:

To find the component of $\mathbf{a}$ that lies within the plane, subtract the projection from $\mathbf{a}$:

$$\mathbf{a}_{\parallel \text{plane}} = \mathbf{a} - \text{proj}_{\mathbf{n}}\mathbf{a}.$$

Substitute the values:

$$\mathbf{a}_{\parallel \text{plane}} = \langle 3, 4, 5 \rangle - \langle 4, 4, 4 \rangle = \langle -1, 0, 1 \rangle.$$

Summary: The projection of $\mathbf{a}$ onto $\mathbf{n}$ is $\langle 4, 4, 4 \rangle$, and the component of $\mathbf{a}$ within the plane perpendicular to $\mathbf{n}$ is $\langle -1, 0, 1 \rangle$.

The Relationship Between Points and Planes

Expanding our understanding of planes, we now explore various ways to define planes and examine relationships between them. Specifically, we'll cover how to create the equation of a plane from three points, calculate the distance of a point from a plane, measure the distance between parallel planes, and determine whether two planes are identical.

Creating a Plane Equation from Three Points

Given three non-collinear points $P_1 = (x_1, y_1, z_1)$, $P_2 = (x_2, y_2, z_2)$, and $P_3 = (x_3, y_3, z_3)$, we can define a unique plane passing through them. The approach is as follows:

1. **Find two direction vectors in the plane:** We can determine two vectors lying in the plane by taking the difference of coordinates between these points:

$$\mathbf{v}_1 = \langle x_2 - x_1, y_2 - y_1, z_2 - z_1 \rangle, \quad \mathbf{v}_2 = \langle x_3 - x_1, y_3 - y_1, z_3 - z_1 \rangle.$$

These vectors span the plane containing P_1 , P_2 , and P_3 .

2. **Compute the normal vector of the plane:** The normal vector $\mathbf{n}$ of the plane is perpendicular to both $\mathbf{v}_1$ and $\mathbf{v}_2$. We find it by calculating the cross product:

$$\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2.$$

Let $\mathbf{n} = \langle A, B, C \rangle$, where A , B , and C are the components of the normal vector derived from this cross product.

3. **Use a point to define the plane equation:** With the normal vector $\mathbf{n} = \langle A, B, C \rangle$ and a point $P_1 = (x_1, y_1, z_1)$ on the plane, the equation of the plane is:

$$A(x - x_1) + B(y - y_1) + C(z - z_1) = 0.$$

Expanding this equation gives the general form:

$$Ax + By + Cz = D,$$

where $D = Ax_1 + By_1 + Cz_1$.

Distance of a Point from a Plane

The shortest distance from a point $\mathbf{x} = (x_0, y_0, z_0)$ to a plane given by the equation $Ax + By + Cz + D = 0$ is the perpendicular distance from the point to the plane. This distance can be found using the normal vector of the plane, which provides a direction that is perpendicular to the plane itself. A key idea in computing this distance is to “walk” from the point $\mathbf{x}$ in the direction of the plane’s normal vector $\mathbf{n} = \langle A, B, C \rangle$ until reaching the plane. The point on the plane that is reached in this way will be the closest point to $\mathbf{x}$, thus giving the shortest distance. This can be approached by the following steps:

1. **Parameterize the Path:** Starting at $\mathbf{x}$, we move along the normal vector $\mathbf{n} = \langle A, B, C \rangle$. For a parameter t , the point on this path is given by:

$$\mathbf{x}(t) = \mathbf{x} + t\mathbf{n} = (x_0, y_0, z_0) + t\langle A, B, C \rangle.$$

Expanding this, we get:

$$\mathbf{x}(t) = (x_0 + tA, y_0 + tB, z_0 + tC).$$

2. **Determine the Intersection with the Plane:** The point $\mathbf{x}(t)$ intersects the plane when it satisfies the plane’s equation $Ax + By + Cz + D = 0$. Substitute $\mathbf{x}(t)$ into the plane equation to find the value of t that leads to this intersection:

$$A(x_0 + tA) + B(y_0 + tB) + C(z_0 + tC) + D = 0.$$

Expanding and simplifying, we obtain a linear equation in t :

$$t(A^2 + B^2 + C^2) + (Ax_0 + By_0 + Cz_0 + D) = 0.$$

Solving for t , we find:

$$t = -\frac{Ax_0 + By_0 + Cz_0 + D}{A^2 + B^2 + C^2}.$$

3. **Calculate the Distance:** The shortest distance d from the point $\mathbf{x}$ to the plane is then the magnitude of the displacement vector $t\mathbf{n}$ that we traveled along the normal to reach the plane. This is given by:

$$d = |t||\mathbf{n}| = \left| -\frac{Ax_0 + By_0 + Cz_0 + D}{\sqrt{A^2 + B^2 + C^2}} \right| = \frac{|Ax_0 + By_0 + Cz_0 + D|}{\sqrt{A^2 + B^2 + C^2}}.$$

This formula gives the shortest distance between a point $\mathbf{x} = (x_0, y_0, z_0)$ and a plane $Ax + By + Cz + D = 0$ by calculating the perpendicular distance from the point to the plane. This approach leverages the direction of the normal vector, reinforcing the concept of perpendicularity as the shortest path to the plane. This is a very abstract configuration, but we will see in a later example how this works simply.

Distance Between Parallel Planes

When two planes are parallel, they have normal vectors that are scalar multiples of each other. For example, consider two parallel planes:

$$Ax + By + Cz + D_1 = 0, \quad Ax + By + Cz + D_2 = 0.$$

Since the planes are parallel, we can measure the distance between them by taking any point on one plane and calculating its distance to the other plane. The distance d between two parallel planes is:

$$d = \frac{|D_2 - D_1|}{\sqrt{A^2 + B^2 + C^2}}.$$

If the planes are not parallel, they intersect, and hence the distance between them is zero.

Determining if Two Planes are Identical

Two planes are considered identical if they have the same normal vector (or scalar multiples of each other) and contain the same set of points. Given two planes:

$$A_1x + B_1y + C_1z + D_1 = 0, \quad A_2x + B_2y + C_2z + D_2 = 0,$$

we can check for equality as follows:

1. **Verify parallel normal vectors:** First, confirm that the normal vectors $\mathbf{n}_1 = \langle A_1, B_1, C_1 \rangle$ and $\mathbf{n}_2 = \langle A_2, B_2, C_2 \rangle$ are scalar multiples of each other. If they are not, the planes are not identical and may intersect along a line.
2. **Check if one plane lies within the other:** Substitute a point from one plane into the equation of the other plane. For example, take a point (x_1, y_1, z_1) that lies on the first plane. If this point also satisfies the second plane's equation, then the planes coincide, meaning they are identical.
3. **Conclusion:** If both conditions are met—parallel normal vectors and a shared set of points—then the planes are identical.

▲ **Example 1.4.6** Suppose we have three points $P_1 = (1, 2, 3)$, $P_2 = (4, -1, 2)$, and $P_3 = (-2, 3, 5)$. Find the equation of the plane passing through these points.

Solution

1. Calculate two direction vectors in the plane:

Determine two vectors in the plane by subtracting the coordinates of P_1 from P_2 and P_3 :

$$\mathbf{v}_1 = \langle 4 - 1, -1 - 2, 2 - 3 \rangle = \langle 3, -3, -1 \rangle,$$

$$\mathbf{v}_2 = \langle -2 - 1, 3 - 2, 5 - 3 \rangle = \langle -3, 1, 2 \rangle.$$

2. Find the normal vector $\mathbf{n}$:

The normal vector $\mathbf{n}$ is perpendicular to both $\mathbf{v}_1$ and $\mathbf{v}_2$, so we calculate the cross product:

$$\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2.$$

Using the formula for the cross product:

$$\mathbf{n} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 3 & -3 & -1 \\ -3 & 1 & 2 \end{vmatrix}.$$

Expanding along the first row, we get:

$$\mathbf{n} = \mathbf{i}((-3)(2) - (-1)(1)) - \mathbf{j}((3)(2) - (-1)(-3)) + \mathbf{k}((3)(1) - (-3)(-3)).$$

Simplifying each term:

$$\mathbf{n} = \mathbf{i}(-6 + 1) - \mathbf{j}(6 - 3) + \mathbf{k}(3 - 9),$$

$$\mathbf{n} = \langle -5, -3, -6 \rangle.$$

3. Use a point to define the plane equation:

With the normal vector $\mathbf{n} = \langle -5, -3, -6 \rangle$ and the point $P_1 = (1, 2, 3)$ on the plane, the equation of the plane is:

$$-5(x - 1) - 3(y - 2) - 6(z - 3) = 0.$$

Expanding this equation:

$$-5x + 5 - 3y + 6 - 6z + 18 = 0,$$

$$-5x - 3y - 6z = -29.$$

Rearranging, we get the plane equation:

$$5x + 3y + 6z = 29.$$

Summary: The plane passing through the points $(1, 2, 3)$, $(4, -1, 2)$, and $(-2, 3, 5)$ has the equation $5x + 3y + 6z = 29$.

▲ **Example 1.4.7** Calculate the shortest distance from the point $\mathbf{x} = (1, -1, -3)$ to the plane given by $x + 2y + 3z = 18$ using a parameterization approach.

Solution

1. Identify the normal vector and starting point:

The plane's equation is $x + 2y + 3z = 18$, so the normal vector $\mathbf{n}$ is:

$$\mathbf{n} = \langle 1, 2, 3 \rangle.$$

This normal vector will guide us in the direction perpendicular to the plane, which is the direction along which the shortest distance from the point to the plane occurs.

2. Parameterize the line through $\mathbf{x}$ in the direction of $\mathbf{n}$:

Starting at $\mathbf{x} = (1, -1, -3)$ and moving along $\mathbf{n}$, we can parameterize the path as:

$$\mathbf{x}(t) = \mathbf{x} + t\mathbf{n} = (1, -1, -3) + t\langle 1, 2, 3 \rangle.$$

Expanding this expression gives:

$$\mathbf{x}(t) = (1 + t, -1 + 2t, -3 + 3t).$$

3. Determine where $\mathbf{x}(t)$ intersects the plane:

The point $\mathbf{x}(t)$ will intersect the plane when it satisfies the plane equation $x + 2y + 3z = 18$. Substitute the components of $\mathbf{x}(t)$ into the plane equation:

$$(1 + t) + 2(-1 + 2t) + 3(-3 + 3t) = 18.$$

Simplify this equation:

$$1 + t - 2 + 4t - 9 + 9t = 18.$$

Combine like terms:

$$14t - 10 = 18.$$

Solving for t :

$$14t = 28 \implies t = 2.$$

4. Find the intersection point $\mathbf{y}$ on the plane:

Substitute $t = 2$ back into $\mathbf{x}(t)$ to find the coordinates of the intersection point $\mathbf{y}$:

$$\mathbf{y} = \mathbf{x}(2) = (1 + 2, -1 + 4, -3 + 6) = (3, 3, 3).$$

5. Calculate the distance between $\mathbf{x}$ and $\mathbf{y}$:

The shortest distance d between $\mathbf{x} = (1, -1, -3)$ and $\mathbf{y} = (3, 3, 3)$ is the magnitude of the displacement vector $\mathbf{y} - \mathbf{x}$:

$$d = |\mathbf{y} - \mathbf{x}| = |(3 - 1, 3 + 1, 3 + 3)| = |(2, 4, 6)|.$$

Calculate the magnitude:

$$d = \sqrt{2^2 + 4^2 + 6^2} = \sqrt{4 + 16 + 36} = \sqrt{56} = 2\sqrt{14}.$$

Summary: Using a parameterization approach, we find that the shortest distance from the point $\mathbf{x} = (1, -1, -3)$ to the plane $x + 2y + 3z = 18$ is $2\sqrt{14}$.

Lecture 1.5: Quadric Surfaces

Introduction to Quadric Surfaces

As we move deeper into three-dimensional geometry, we encounter more complex surfaces than planes. A particularly important class of surfaces is the *quadric surfaces*, which generalize the concept of conic sections from two to three dimensions. These surfaces form the foundation for much of multivariable calculus and geometric analysis. To understand quadric surfaces fully, we need to carefully explore their mathematical origins, the variety of shapes they produce, and the relationships between these shapes.

What is a Conic Section?

Before extending to three dimensions, let us first revisit the concept of a conic section. A conic section is the curve obtained by intersecting a plane with a double-napped cone. We take those curves and then examine them in two dimensions. Depending on the angle of intersection between the plane and the cone, we get different types of curves:

- **Circle:** The plane intersects the cone perpendicular to its axis of symmetry.
- **Ellipse:** The plane intersects the cone at an angle, but does not pass through the cone's vertex.
- **Parabola:** The plane is parallel to one of the cone's generating lines.
- **Hyperbola:** The plane intersects both nappes of the cone.




Figure 1.27: Conic sections derived from a single-napped cone. The hyperbola includes an additional section, as the intersecting plane extends through the origin to cut both nappes of the cone when there are two present.

These curves can all be described by the general quadratic equation in two dimensions:

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0,$$

where A, B, C, D, E, F are constants. By varying these constants, we generate the different conic sections.

A natural question arises: *How can we extend the idea of conic sections to three dimensions?* If conic sections are formed by the intersection of a plane with a cone, what would happen if we instead consider the intersection of a plane with a "generalized" three-dimensional quadratic surface? What kinds of shapes might we discover in three dimensions? These questions lead us to quadric surfaces.

In three dimensions, the general equation for a quadric surface is:

$$Ax^2 + By^2 + Cz^2 + Dxy + Exz + Fyz + Gx + Hy + Iz + J = 0,$$

where $A, B, C, D, E, F, G, H, I, J$ are constants. By adjusting these constants, we can produce a variety of surfaces, such as ellipsoids, hyperboloids, and paraboloids. Each surface is a natural extension of the conic sections, generalized to three-dimensional space. One of the most fascinating aspects of quadric surfaces is the diversity of shapes they produce. Why do we get surfaces as different as spheres, hyperboloids, and paraboloids from a single quadratic equation? Are these shapes related to one another? The answer lies in the interplay between the coefficients in the general equation and the symmetry of the surface.

To understand this, let us focus on the terms in the quadratic equation:

$$Ax^2 + By^2 + Cz^2 + Dxy + Exz + Fyz + Gx + Hy + Iz + J = 0.$$

- **Symmetry and Dominance of Quadratic Terms:** The coefficients A, B , and C determine the relative "stretching" or "shrinking" along the x -, y -, and z -axes. For example, if A, B , and C are all positive, the surface is "elliptical" in nature, producing shapes like spheres or ellipsoids. If one of these coefficients is negative, the surface takes on a "hyperbolic" character.
- **Mixed Terms:** The presence of mixed terms like Dxy, Exz , or Fyz introduces asymmetry into the surface. These terms "tilt" the surface or create interactions between axes that result in more complex geometries.
- **Linear Terms:** The linear terms Gx, Hy, Iz shift the surface in space, altering its position without changing its overall shape.
- **Constant Term:** The constant J determines whether the surface passes through the origin or is offset.

The diversity of shapes arises from how these terms combine. For example:

- A sphere occurs when $A = B = C$, and all other terms vanish.
- An ellipsoid occurs when $A, B, C > 0$, but $A \neq B \neq C$, and all mixed terms vanish.
- A hyperboloid arises when one of A, B , or C is negative, creating the distinctive "saddle" or "double-sheeted" structure.

Mathematically, these surfaces are related because they share the same fundamental structure: all are solutions to quadratic equations in three variables. By varying the coefficients and constants in the equation, we transition smoothly from one surface type to another. To visualize why these shapes differ, think of a quadric surface as a collection of cross-sections. For example:

- Setting $z = k$ in the equation of a quadric surface gives a two-dimensional curve (a conic section) in the xy -plane. As k varies, these curves “stack up” to form the three-dimensional surface.
- Similarly, setting $x = k$ or $y = k$ provides cross-sections in the yz - and xz -planes.

By examining these cross-sections, we can see how the different shapes arise:

- For an ellipsoid, every cross-section is an ellipse, regardless of the plane.
- For a hyperboloid, some cross-sections are hyperbolas, while others are ellipses.
- For a paraboloid, cross-sections are parabolas in one direction and ellipses in another.

This perspective highlights the deep connection between conic sections and quadric surfaces: quadric surfaces are essentially “stacks” of conic sections in three dimensions.

Preview of Multivariate Functions

While our primary focus here is on the geometric nature of quadric surfaces, these surfaces also serve as important examples of multivariate functions. For instance, the equation of an elliptic paraboloid:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = z$$

can be interpreted as the graph of the function $z = f(x, y)$. Similarly, hyperbolic paraboloids and other quadric surfaces appear as graphs of functions with more complex structures. Understanding quadric surfaces geometrically will prepare us to study multivariate functions rigorously in future lectures, where we will explore topics like partial derivatives, gradients, and optimization.

The Significance of Quadric Surfaces

Quadric surfaces derive their name from the quadratic equations that define them. However, you might wonder: *Why do we add the suffix “-oid” to names like ellipsoid, paraboloid, and hyperboloid?* The term “-oid” comes from the Greek word *oides*, meaning “like” or “resembling.” This suffix reflects the idea that quadric surfaces are three-dimensional analogs of their two-dimensional counterparts, the conic sections. For example:

- An **ellipsoid** is “like” an ellipse but extends into three dimensions.
- A **paraboloid** is “like” a parabola but curves in three-dimensional space.

- A **hyperboloid** is “like” a hyperbola, forming a saddle-shaped or double-sheeted surface in three dimensions.

This connection emphasizes the role of quadric surfaces as natural extensions of conic sections. While conic sections arise as intersections of a plane with a cone in two dimensions, quadric surfaces can be thought of as “generalized” cones or cylinders in three dimensions.

Why Are Quadric Surfaces Important? These surfaces form the foundation for multivariate functions because they provide simple models for more complex phenomena. The symmetry inherent in quadric surfaces makes them ideal for representing many physical systems and mathematical problems. Consider the following examples:

- **Paraboloids:** The surface of a satellite dish is a paraboloid, which focuses signals to a single point. Similarly, the shape of a suspension bridge cable under uniform load approximates a paraboloid.
- **Ellipsoids:** The orbits of some planets and the shape of Earth (approximated as an oblate spheroid) can be modeled using ellipsoids.
- **Hyperboloids:** Certain cooling towers, telescopes, and lenses rely on hyperboloids for their structural and optical properties.

Beyond these applications, quadric surfaces are central to **optimization problems**. They often appear as the contours of multivariate functions, where the geometry of the surface reveals key insights into the behavior of the function. For instance, an elliptic paraboloid can represent a local minimum, while a hyperbolic paraboloid can represent a saddle point.

Equations of Quadric Surfaces

Let us first consider the simplest cases of quadric surfaces and slowly generalize to more complex ones. We will examine each surface type systematically, deriving its equation and visualizing its geometry.

Ellipsoids

An ellipsoid is a natural three-dimensional extension of an ellipse. It is a surface defined by the equation:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1,$$

where $a, b, c > 0$ are constants that determine the lengths of the semi-axes along the x -, y -, and z -axes, respectively. To understand this equation intuitively, let us set $z = 0$. The equation reduces to:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1,$$

which represents an ellipse in the xy -plane. Similarly, setting $y = 0$ gives:

$$\frac{x^2}{a^2} + \frac{z^2}{c^2} = 1,$$

an ellipse in the xz -plane. Thus, the ellipsoid can be thought of as a “stack” of ellipses in different planes.

Hyperboloids

A hyperboloid comes in two varieties: one-sheeted and two-sheeted.

One-Sheeted Hyperboloid: The equation for a one-sheeted hyperboloid is:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1.$$

Here, the negative term $(-\frac{z^2}{c^2})$ causes the surface to open outward, creating a “saddle-like” shape. Setting $z = 0$ reduces the equation to:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1,$$

an ellipse in the xy -plane. However, as z increases or decreases, the cross-sections remain ellipses but grow larger, giving the hyperboloid its distinctive shape.

Two-Sheeted Hyperboloid: The equation for a two-sheeted hyperboloid is:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = -1.$$

This equation is similar to that of the one-sheeted hyperboloid, but here the minus sign on the right-hand side creates two disconnected surfaces, or “sheets,” opening in opposite directions. Setting $z = 0$ results in no real solutions, which explains the gap between the two sheets.

Elliptic Paraboloid

The elliptic paraboloid is defined by the equation:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = z.$$

To understand why we call this surface an *elliptic paraboloid*, let us analyze its cross-sections:

- Setting $z = k$, where $k > 0$, reduces the equation to:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = k,$$

which is the equation of an ellipse in the xy -plane. As z increases, these ellipses grow larger, creating a “bowl-like” shape that opens upwards.

- On the other hand, setting $x = k_1$ or $y = k_2$ (for fixed constants k_1, k_2) gives parabolas in the yz - or xz -plane, respectively. These parabolas describe how the surface curves along the vertical direction.

The name *elliptic paraboloid* reflects the dual nature of this surface:

- The term “elliptic” comes from the elliptical cross-sections in the xy -plane.
- The term “paraboloid” signifies that the surface extends like a parabola in the vertical direction.

This combination of elliptical and parabolic characteristics is the hallmark of an elliptic paraboloid.

Hyperbolic Paraboloid

The hyperbolic paraboloid, often called a “saddle surface,” is defined by the equation:

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = z.$$

To understand the naming of this surface, let us examine its structure:

- Setting $z = k$ (for any constant k) reduces the equation to:

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = k,$$

which represents hyperbolas in the xy -plane when $k \neq 0$. These hyperbolas change orientation depending on the sign of k :

- For $k > 0$, the hyperbolas open along the x -axis.
- For $k < 0$, the hyperbolas open along the y -axis.
- When $z = 0$, the equation becomes:

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 0,$$

which simplifies to two intersecting lines, $y = \pm \frac{b}{a}x$, in the xy -plane.

- Setting $x = k_1$ or $y = k_2$ (for fixed constants k_1, k_2) gives parabolas in the yz - or xz -plane, respectively. These parabolas curve in opposite directions, creating the characteristic “saddle” shape.

The name *hyperbolic paraboloid* reflects the surface’s dual nature:

- The term “hyperbolic” comes from the hyperbolic cross-sections in the xy -plane.
- The term “paraboloid” signifies that the surface contains parabolic cross-sections along the xz - and yz -planes.

The hyperbolic paraboloid’s unique combination of upward and downward curvature makes it a saddle surface, giving it a distinct role in geometry and multivariable calculus.

Cylinders

Cylinders are unique among quadric surfaces because they extend infinitely in one direction while maintaining a consistent cross-sectional shape in the other two dimensions. A cylinder can be thought of as the “extrusion” of a conic section along an axis. The simplest form of a cylinder is the circular cylinder, given by:

$$x^2 + y^2 = r^2,$$

where r is the radius. While this equation might appear two-dimensional, it is not. The apparent “two-dimensional” nature arises because the equation describes the cross-sectional shape of the cylinder in the xy -plane when $z = 0$. The cylinder itself is three-dimensional, extending infinitely in the z -direction. Setting $z = k$ for any k gives circular cross-sections of radius r in the xy -plane at that height. More generally, a cylinder can take on other cross-sectional shapes, such as ellipses or parabolas. For example:

- An **elliptic cylinder** is defined by:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1,$$

which forms an ellipse in the xy -plane and extends infinitely in the z -direction.

- A **parabolic cylinder** is defined by:

$$y = ax^2,$$

which forms a parabola in the xy -plane and extends infinitely in the z -direction.

Cylinders are significant because they introduce the idea of surfaces that are not entirely bounded but still exhibit a clear geometric structure. They also help bridge the gap between two-dimensional conic sections and three-dimensional quadric surfaces.

Cones

A cone is defined by the equation:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{z^2}{c^2}.$$

Setting $z = k$, where $k \neq 0$, reduces the equation to:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = k^2,$$

an ellipse in the xy -plane. At $z = 0$, the equation reduces to a single point (the vertex of the cone).

Visualization

The below visual are what these quadric surfaces look like:




Figure 1.28: Assorted quadric surfaces.

The labeled surfaces are as follows:

- a. **Hyperboloid of One Sheet:** $\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$
- b. **Hyperboloid of Two Sheets:** $-\frac{x^2}{a^2} - \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$
- c. **Hyperbolic Paraboloid:** $\frac{x^2}{a^2} - \frac{y^2}{b^2} = z$
- d. **Ellipsoid:** $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$
- e. **Cylinder:** $x^2 + y^2 = r^2$
- f. **Elliptic Paraboloid:** $\frac{x^2}{a^2} + \frac{y^2}{b^2} = z$
- g. **Elliptic Cone:** $\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 0$

To further explore the concepts of quadric surfaces, let us introduce a Python program that allows you to visualize these surfaces by manipulating their coefficients. By adjusting the parameters A, B, C , etc., you can observe how the shape and orientation of the surface changes. This hands-on approach helps build intuition about these mathematical objects.

```

1 import numpy as np
2 import matplotlib.pyplot as plt
3 from mpl_toolkits.mplot3d import Axes3D
4
5 # Define the function for the quadric surface
6 def quadric_surface(x, y, A, B, C, D, E, F, G, H, I, J):
7     return A*x**2 + B*y**2 + C*(x*y) + D*x + E*y + F # Replace or expand as needed
8
9 # Coefficients (students can modify these to observe changes)
10 A, B, C = 1, 1, 0
11 D, E, F = 0, 0, -1
12
13 # Create a meshgrid for the surface
14 x = np.linspace(-2, 2, 100)
15 y = np.linspace(-2, 2, 100)
16 x, y = np.meshgrid(x, y)
17
18 # Compute z-values for the surface
19 z = quadric_surface(x, y, A, B, C, D, E, F, 0, 0, 0, 0)
20
21 # Plot the surface
22 fig = plt.figure(figsize=(8, 6))
23 ax = fig.add_subplot(111, projection='3d')
24 ax.plot_surface(x, y, z, cmap="viridis", alpha=0.8)
25
26 # Labels and title
27 ax.set_xlabel('X-axis')
28 ax.set_ylabel('Y-axis')
29 ax.set_zlabel('Z-axis')
30 ax.set_title('Quadric Surface Visualization')
31
32 plt.show()

```

How to Use:

1. Copy and paste the code into a Python environment, such as Jupyter Notebook or an IDE.
2. Experiment with different values of the coefficients A, B, C, D, E, F , etc., to explore different quadric surfaces (e.g., ellipsoids, hyperboloids, paraboloids).
3. Observe how the surface changes in response to these adjustments and compare it with the equations we've discussed in this lecture.

This interactive approach not only reinforces your understanding of quadric surfaces but also demonstrates how algebraic changes in the equation correspond to geometric changes in the graph.

▲ Example 1.5.1 Visualizing the Intersection of a Plane and a Quadric Surface

Consider the quadric surface given by:

$$\frac{x^2}{4} + \frac{y^2}{9} - \frac{z^2}{16} = 1,$$

which is a hyperboloid of one sheet. Let the intersecting plane be defined by $z = 2$. Find the resulting curve of intersection.

Solution

To find the intersection, substitute $z = 2$ into the equation of the hyperboloid:

$$\frac{x^2}{4} + \frac{y^2}{9} - \frac{(2)^2}{16} = 1.$$

Simplify the z^2 term:

$$\frac{x^2}{4} + \frac{y^2}{9} - \frac{4}{16} = 1 \quad \implies \quad \frac{x^2}{4} + \frac{y^2}{9} - \frac{1}{4} = 1.$$

Rearrange:

$$\frac{x^2}{4} + \frac{y^2}{9} = \frac{5}{4}.$$

Multiply through by 36 (the least common multiple of 4 and 9) to clear the denominators:

$$9x^2 + 4y^2 = 45.$$

This is the equation of an ellipse in the xy -plane. The plane $z = 2$ intersects the hyperboloid to form an ellipse at this height.

Summary: The curve of intersection between the hyperboloid $\frac{x^2}{4} + \frac{y^2}{9} - \frac{z^2}{16} = 1$ and the plane $z = 2$ is:

$$9x^2 + 4y^2 = 45.$$

This describes an ellipse in the xy -plane when viewed at $z = 2$.

▲ Example 1.5.2 Classifying a Quadric Surface by Its Equation

Given the equation:

$$\frac{x^2}{9} + \frac{y^2}{4} + \frac{z^2}{16} = 1,$$

identify the type of quadric surface and describe its geometric properties.

Solution

The given equation is of the form:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1,$$

where $a^2 = 9$, $b^2 = 4$, and $c^2 = 16$. Since all the terms are positive and equal to 1 on the right-hand side, this is an **ellipsoid**.

Geometric Properties:

- The center of the ellipsoid is at $(0, 0, 0)$, as there are no linear terms in x, y , or z .
- The semi-axes of the ellipsoid are aligned with the coordinate axes:

$$a = 3, \quad b = 2, \quad c = 4.$$

- The semi-axis lengths a, b, c define the radii of the ellipsoid along the x, y , and z -axes, respectively.

Summary: The equation $\frac{x^2}{9} + \frac{y^2}{4} + \frac{z^2}{16} = 1$ represents an ellipsoid centered at the origin with semi-axes of lengths 3, 2, and 4 along the x, y , and z -axes.

▲ **Example 1.5.3 Finding the Intersection Between a Cylinder and a Plane**

Consider the cylinder:

$$x^2 + y^2 = 9,$$

and the plane $z = 3x + y$. Find the parametric equations for their intersection.

Solution

This example introduces a new technique for working with nonlinear equations. Previously, we dealt with linear parameterizations for lines and planes. Here, we parameterize a *circle* in the xy -plane, which is inherently nonlinear.

The cylinder is defined by $x^2 + y^2 = 9$. This equation represents a circle in the xy -plane with radius 3. To parameterize the circle, we note that for a unit circle, the coordinates of points on the circle can be written as $x = \cos t$ and $y = \sin t$, where t is the angle swept counterclockwise from the positive x -axis. Scaling this relationship to a circle of radius r gives:

$$x = r \cos t, \quad y = r \sin t,$$

where $t \in [0, 2\pi)$ describes the angle around the circle. Substituting $r = 3$, we have:

$$x = 3 \cos t, \quad y = 3 \sin t, \quad t \in [0, 2\pi).$$

This parameterization allows us to represent all points on the circle in the xy -plane as t varies from 0 to 2π .

The plane $z = 3x + y$ provides the z -coordinate for each point of the intersection. Substituting the parameterized values of x and y into the plane equation:

$$z = 3x + y \implies z = 3(3 \cos t) + 3 \sin t = 9 \cos t + 3 \sin t.$$

Thus, the parametric equations for the curve of intersection are:

$$x = 3 \cos t, \quad y = 3 \sin t, \quad z = 9 \cos t + 3 \sin t, \quad t \in [0, 2\pi).$$

Key Insight: Nonlinear equations like $x^2 + y^2 = 9$ can often be parameterized using trigonometric functions ($\cos t$ and $\sin t$), leveraging their geometric relationship with circles. While the equations may appear two-dimensional initially, the z -coordinate from the plane adds a third dimension to the curve of intersection, creating a space curve.

Summary: The curve of intersection between the cylinder $x^2 + y^2 = 9$ and the plane $z = 3x + y$ is a space curve parameterized by:

$$x = 3 \cos t, \quad y = 3 \sin t, \quad z = 9 \cos t + 3 \sin t, \quad t \in [0, 2\pi).$$

Formulas and Definitions

Important Vocabulary

- **Cartesian Coordinates:** A system to define points in three-dimensional space using (x, y, z) .
- **Cylindrical Coordinates:** Defined by (r, θ, z) , where r is the radial distance, θ the angle from the positive x -axis, and z the height.
- **Spherical Coordinates:** Represented as (ρ, θ, ϕ) , where ρ is the radial distance, θ the azimuthal angle, and ϕ the polar angle.
- **Vector:** A quantity with both magnitude and direction, expressed as $\mathbf{v} = \langle v_x, v_y, v_z \rangle$.
- **Dot Product:** A scalar product of two vectors, given by $\mathbf{u} \cdot \mathbf{v} = u_x v_x + u_y v_y + u_z v_z$.
- **Cross Product:** A vector perpendicular to two vectors, calculated using the determinant of a 3×3 matrix.
- **Parallelepiped:** A six-faced figure formed by three vectors, where its volume is given by the scalar triple product $|\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})|$.

Key Formulas

Distance Formula in 3D:

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

Dot Product:

$$\mathbf{u} \cdot \mathbf{v} = u_x v_x + u_y v_y + u_z v_z$$

Midpoint Formula:

$$M = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2} \right)$$

Cross Product:

$$\begin{aligned} \mathbf{u} \times \mathbf{v} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_x & u_y & u_z \\ v_x & v_y & v_z \end{vmatrix} \\ &= \mathbf{i}(u_y v_z - u_z v_y) \\ &\quad - \mathbf{j}(u_x v_z - u_z v_x) \\ &\quad + \mathbf{k}(u_x v_y - u_y v_x) \end{aligned}$$

Direction Cosines:

$$\begin{aligned} \cos \alpha &= \frac{u_x}{|\mathbf{u}|}, & \cos \beta &= \frac{u_y}{|\mathbf{u}|}, \\ \cos \gamma &= \frac{u_z}{|\mathbf{u}|} \end{aligned}$$

where $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$

Parametric Line:

$$\mathbf{r}(t) = \mathbf{P}_0 t + \mathbf{P}_1(1 - t), \quad t \in [0, 1]$$

Angle Between Vectors:

$$\cos \theta = \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}| |\mathbf{v}|}, \quad \sin \theta = \frac{|\mathbf{u} \times \mathbf{v}|}{|\mathbf{u}| |\mathbf{v}|}$$

or alternatively:

$$\mathbf{r}(t) = \mathbf{P}_0 + t\mathbf{v}, \quad t \in \mathbb{R}$$

Volume of a Parallelepiped (Scalar Triple Product):

$$\text{Volume} = |\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})|$$

$$Ax + By + Cz + D = 0$$

Conversion Between Coordinates:

- Cartesian to Cylindrical:

$$r = \sqrt{x^2 + y^2}, \quad \theta = \tan^{-1} \left(\frac{y}{x} \right),$$

$$z = z$$

- Cartesian to Spherical:

$$\rho = \sqrt{x^2 + y^2 + z^2}, \quad \theta = \tan^{-1} \left(\frac{y}{x} \right),$$

$$\phi = \cos^{-1} \left(\frac{z}{\rho} \right)$$

- Cylindrical to Cartesian:

$$x = r \cos \theta, \quad y = r \sin \theta,$$

$$z = z$$

- Spherical to Cartesian:

$$x = \rho \sin \phi \cos \theta, \quad y = \rho \sin \phi \sin \theta,$$

$$z = \rho \cos \phi$$

Line Equation (Parametric Form):

$$\mathbf{r}(t) = \mathbf{v} + t\mathbf{d}$$

Quadratic Surfaces:

- **Ellipsoid:** $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$

- **Hyperboloid of One Sheet:** $\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$

- **Hyperboloid of Two Sheets:** $-\frac{x^2}{a^2} - \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$

- **Elliptic Paraboloid:** $\frac{x^2}{a^2} + \frac{y^2}{b^2} = z$

Practice Problems

- Differentiate the function $f(t) = t^3 + \sin(t) + e^t$ with respect to t .
- Compute the limit $\lim_{t \rightarrow 0} \frac{\sin(t)}{t}$.
- Integrate the function $f(t) = 2t+3$ with respect to t .
- Find the critical points of $f(t) = t^4 - 4t^3 + 6t^2 - 4t + 1$.
- Determine the interval of convergence for the power series $\sum_{n=0}^{\infty} \frac{x^n}{n!}$.
- Evaluate the improper integral $\int_1^{\infty} \frac{1}{x^2} dx$.
- Find the Taylor series of e^x centered at $x = 0$ up to the x^3 term.
- Compute $\lim_{x \rightarrow \infty} \frac{2x^3 - x + 1}{x^3 + 4x^2}$.
- Solve the optimization problem: Find the maximum area of a rectangle inscribed under the curve $y = 4 - x^2$.
- A ladder 10 feet long is leaning against a wall. If the base of the ladder is pulled away from the wall at a rate of 2 feet per second, how fast is the top of the ladder descending when the base is 6 feet from the wall?
- Convert the rectangular coordinates $(x, y, z) = (3, -3, 3)$ to spherical coordinates.
- Find the rectangular coordinates of the point with cylindrical coordinates $(r, \theta, z) = \left(2, \frac{\pi}{4}, -1\right)$.
- Determine the distance between the points $(2, -1, 4)$ and $(-3, 5, -2)$ in three-dimensional space.
- Express the spherical coordinates $(\rho, \theta, \phi) = \left(5, \frac{\pi}{3}, \frac{\pi}{6}\right)$ in rectangular coordinates.
- Convert the point $(x, y, z) = (-4, 4, \sqrt{3})$ from rectangular to cylindrical coordinates.
- Find the midpoint between the points $(1, 2, 3)$ and $(-1, -2, -3)$ in three-dimensional space.
- Determine the angle between the positive x -axis and the projection of the vector $\mathbf{v} = \langle 3, 3, 3 \rangle$ onto the xy -plane.
- Convert the rectangular coordinates $(0, -5, 5)$ to spherical coordinates.
- Find the cylindrical coordinates of the point $(x, y, z) = (2\sqrt{2}, 2\sqrt{2}, -3)$.
- Determine the spherical coordinates of the point $(x, y, z) = (-2, -2, 2\sqrt{2})$.
- Express the point $(\rho, \theta, \phi) = \left(6, \frac{2\pi}{3}, \frac{\pi}{4}\right)$ in rectangular coordinates.
- Convert the cylindrical coordinates $(r, \theta, z) = \left(3, \frac{\pi}{6}, 4\right)$ to spherical coordinates.
- Find the rectangular coordinates of the point $(r, \theta, z) = \left(1, \frac{\pi}{2}, -2\right)$.
- Determine the distance from the origin to the point $(x, y, z) = (a, b, c)$ in rectangular coordinates.
- Convert the point $(x, y, z) = (0, 0, 0)$ to spherical coordinates.
- Compute $|\mathbf{v}|$ for $\mathbf{v} = \langle 3, -4, 5 \rangle$.

27. Find the unit vector in the direction of $\mathbf{v} = \langle -2, 1, 3 \rangle$.
28. Calculate $\mathbf{u} \cdot \mathbf{v}$ for $\mathbf{u} = \langle 1, 2, 3 \rangle$ and $\mathbf{v} = \langle 4, 0, -1 \rangle$.
29. Determine $\mathbf{u} \times \mathbf{v}$ for $\mathbf{u} = \langle 1, 0, -1 \rangle$ and $\mathbf{v} = \langle 0, 2, 3 \rangle$.
30. Find the angle between $\mathbf{u} = \langle 1, 2, 3 \rangle$ and $\mathbf{v} = \langle 3, 2, 1 \rangle$.
31. Show that $\mathbf{u} = \langle 2, -1, 0 \rangle$ and $\mathbf{v} = \langle 4, -2, 0 \rangle$ are *linearly dependent* (meaning that one cannot be written as a constant multiple of the other).
32. Compute the area of the parallelogram spanned by $\mathbf{u} = \langle 1, 0, 1 \rangle$ and $\mathbf{v} = \langle 0, 2, 1 \rangle$.
33. Find a vector orthogonal to both $\mathbf{u} = \langle 1, -1, 0 \rangle$ and $\mathbf{v} = \langle 2, 1, -1 \rangle$.
34. Determine if the vectors $\mathbf{a} = \langle 3, -1, 2 \rangle$, $\mathbf{b} = \langle 4, 0, -2 \rangle$, and $\mathbf{c} = \langle 1, 5, -1 \rangle$ are linearly independent.
35. Find the projection of $\mathbf{u} = \langle 4, -2, 1 \rangle$ onto $\mathbf{v} = \langle 1, 0, 3 \rangle$.
36. Calculate the scalar triple product $\mathbf{a} \cdot (\mathbf{b} \times \mathbf{c})$ for $\mathbf{a} = \langle 1, 2, 3 \rangle$, $\mathbf{b} = \langle 4, 5, 6 \rangle$, and $\mathbf{c} = \langle 7, 8, 9 \rangle$.
37. Find the volume of the parallelepiped formed by the vectors $\mathbf{u} = \langle 1, 0, 0 \rangle$, $\mathbf{v} = \langle 0, 1, 0 \rangle$, and $\mathbf{w} = \langle 0, 0, 1 \rangle$.
38. Determine the component of $\mathbf{a} = \langle 5, 2, -3 \rangle$ along $\mathbf{b} = \langle 1, -1, 2 \rangle$.
39. Compute the magnitude of the vector $\mathbf{w} = \mathbf{u} \times \mathbf{v}$ where $\mathbf{u} = \langle 2, 3, 4 \rangle$ and $\mathbf{v} = \langle -1, 0, 1 \rangle$.
40. Find the angle between the vectors $\mathbf{a} = \langle 2, -1, 3 \rangle$ and $\mathbf{b} = \langle 1, 4, -2 \rangle$.
41. Show that the vectors $\mathbf{u} = \langle 1, 2, 3 \rangle$, $\mathbf{v} = \langle 4, 5, 6 \rangle$, and $\mathbf{w} = \langle 7, 8, 9 \rangle$ are *coplanar* (are all on the same plane).
42. Calculate the area of the triangle formed by the vectors $\mathbf{u} = \langle 2, 0, -1 \rangle$ and $\mathbf{v} = \langle 1, 3, 4 \rangle$.
43. Find the vector projection of $\mathbf{a} = \langle 3, -2, 1 \rangle$ onto $\mathbf{b} = \langle 4, 0, -1 \rangle$.
44. Determine if the vectors $\mathbf{u} = \langle 2, 4, 6 \rangle$ and $\mathbf{v} = \langle -1, -2, -3 \rangle$ are orthogonal.
45. Compute the cross product $\mathbf{u} \times \mathbf{v}$ for $\mathbf{u} = \langle 0, 1, 2 \rangle$ and $\mathbf{v} = \langle 3, 4, 5 \rangle$.
46. Find the unit vector perpendicular to the plane defined by $\mathbf{u} = \langle 1, 2, 3 \rangle$ and $\mathbf{v} = \langle 4, 5, 6 \rangle$.
47. Determine whether the vectors $\mathbf{a} = \langle 1, 2, 3 \rangle$, $\mathbf{b} = \langle 2, 4, 6 \rangle$, and $\mathbf{c} = \langle 3, 6, 9 \rangle$ are linearly independent or dependent. Linear independence means that none of the vectors can be written as a combination of the others, ie $v_1 \neq k_1 v_2 + k_2 v_3$. If at least one vector can be expressed as a combination of the others, the vectors are linearly dependent.
48. Calculate the dot product $\mathbf{u} \cdot \mathbf{v}$ for $\mathbf{u} = \langle 5, -2, 4 \rangle$ and $\mathbf{v} = \langle -1, 3, 2 \rangle$.
49. Find the angle between the vectors $\mathbf{u} = \langle 7, 0, -5 \rangle$ and $\mathbf{v} = \langle 4, -3, 2 \rangle$.
50. Show that the vectors $\mathbf{u} = \langle 1, 3, -2 \rangle$ and $\mathbf{v} = \langle 4, -1, 5 \rangle$ are not parallel.
51. Compute the area of the parallelogram formed by $\mathbf{u} = \langle 3, -1, 2 \rangle$ and $\mathbf{v} = \langle -2, 4, 1 \rangle$.
52. Find a vector orthogonal to $\mathbf{u} = \langle 2, -3, 1 \rangle$ and $\mathbf{v} = \langle 4, 0, -2 \rangle$.
53. Find the equation of the line passing through $(1, 0, -1)$ and $(2, 3, 4)$.

54. Determine the equation of the plane containing the points $(1, 2, 3)$, $(-1, 0, 4)$, and $(3, -2, 1)$.
55. Compute the point of intersection of the line $\mathbf{r}(t) = \langle 1+t, 2-t, 3t \rangle$ with the plane $x - y + z = 5$.
56. Check if the lines $\mathbf{r}_1(t) = \langle t, 2t, 3t \rangle$ and $\mathbf{r}_2(s) = \langle 1+s, 2, 3s \rangle$ are parallel, intersecting, or skew.
57. Given $\Pi_1 : x + 2y + 3z = 7$, $\Pi_2 : -2x - y - z = 2$, and $\mathbf{r}(t) = \langle 2t, 1-t, 1+t \rangle$, determine if $\mathbf{r}(t)$ intersects both planes. If so, calculate the length of $\mathbf{r}(t)$ from Π_1 to Π_2 .
58. Find the shortest distance between the line $\mathbf{r}(t) = \langle t, t, t \rangle$ and the point $(1, 1, 1)$.
59. Determine the equation of the line of intersection of the planes $x + y + z = 1$ and $2x - y + 3z = 4$.
60. Find the distance between the parallel planes $x + 2y + 2z = 5$ and $x + 2y + 2z = -3$.
61. Find the parametric equations of the line perpendicular to the plane $3x - 2y + z = 7$ and passing through the point $(2, -1, 4)$.
62. Determine the angle between the line $\mathbf{r}(t) = \langle 1 + 2t, 3 - t, 4t \rangle$ and the plane $2x - 3y + z = 7$.
63. Find the equation of the plane perpendicular to the vector $\mathbf{n} = \langle 3, -2, 5 \rangle$ and passing through the point $(2, -1, 4)$.
64. Compute the intersection curve of the sphere $x^2 + y^2 + z^2 = 25$ and the plane $z = 3$.
65. Find the equation of a line that is parallel to the vector $\mathbf{v} = \langle 2, -1, 3 \rangle$ and passes through the point $(-1, 4, 2)$.
66. Determine whether the lines $\mathbf{r}_1(t) = \langle 1 + t, 2 - t, 3 + 2t \rangle$ and $\mathbf{r}_2(s) = \langle 4 - s, -1 + s, 5 + s \rangle$ intersect.
67. Compute the equation of the tangent plane to the surface $x^2 + 2y^2 + 3z^2 = 6$ at the point $(1, 1, 1)$.
68. Find the line segment of shortest length connecting the lines $\mathbf{r}_1(t) = \langle t, 2t, 3t \rangle$ and $\mathbf{r}_2(s) = \langle 1 + s, 2 + s, 3 + s \rangle$.
69. Determine the parametric equations for the space curve defined by $x = t$, $y = \sin(t)$, and $z = \cos(t)$.
70. Find the intersection of the hyperboloid $x^2 - y^2 - z^2 = 4$ with the plane $y = 1$.
71. Calculate the surface area of the cylinder $x^2 + y^2 = 9$ between $z = 0$ and $z = 5$.
72. Find the equation of the ellipsoid with semi-axes of lengths 3, 4, and 5 along the x -, y -, and z -axes, respectively.
73. Derive the Cartesian equation of the paraboloid obtained by rotating the parabola $z = x^2$ about the z -axis.
74. As we have seen, the parametric equation of a line can be represented as $\mathbf{r}(t) = \mathbf{P}_0 t + \mathbf{P}_1(1 - t)$, $t \in [0, 1]$. If we wanted to change t to go between 0 and $a \in \mathbb{R}^+$, how would our parametric equation change?
75. Classify the surface given by $x^2 + y^2 - z^2 = 1$.
76. Write the parametric equations for the intersection of the cylinder $x^2 + y^2 = 4$ with the plane $z = x + y$.

77. Verify if the surface $x^2 + y^2 - 4z = 0$ is an elliptic paraboloid. $z = 1$.
78. Compute the volume enclosed by the sphere $x^2 + y^2 + z^2 = 4$ and the plane
79. Find the equation of the surface generated by rotating $x = y^2$ about the z -axis.

Chapter 2

Vector-Valued Functions and Motion in Space

Introduction

The study of vector-valued functions marks an important progression in our exploration of multivariable calculus, seamlessly combining geometry and calculus to describe motion, curves, and change in three-dimensional space. Interestingly, we have already encountered a special case of vector-valued functions in Chapter 1 when we discussed parametric equations of lines in space. At the time, we focused on their geometric interpretation without explicitly framing them as vector-valued functions. In essence, those parametric representations—where we expressed lines as functions of a parameter—were early examples of linear vector-valued functions.

By revisiting and expanding on this foundation, we will now view these functions through a broader lens, exploring their behavior, calculus, and applications in motion and geometry. This chapter builds on that implicit understanding, introducing parametric equations in their full generality and extending calculus to these functions. Concepts like velocity, acceleration, and arc length emerge naturally from the derivatives and integrals of vector-valued functions, allowing us to model and analyze dynamic systems in physics and engineering with mathematical rigor.

The connections we uncover here not only highlight the unifying power of mathematics but also provide a deeper appreciation for how foundational concepts like lines in space can evolve into tools for analyzing complex systems. By recognizing these links, students can better appreciate how the ideas in earlier chapters inform and enrich their understanding of new topics.

Lecture 2.1: Vector-Valued Functions

Vector-valued functions provide a powerful framework for describing curves, motion, and physical phenomena in three-dimensional space. Prior functions you've seen like $f(x) = x^2$ are technically called scalar functions, mapping a real number to a single real value, whereas vector-valued functions output ordered components that capture both magnitude and direction. Building on the geometric intuition developed in Chapter 1, where we discussed lines as parametric equations, we now revisit those concepts as a foundation and extend them to describe more general curves. This section focuses on parametric equations, the cornerstone of vector-valued functions, and lays the groundwork for their application in analyzing motion and change in space.

Parametric Equations

In Chapter 1, we described a line in space as the set of points extending in a given direction from a specific starting point. If $\mathbf{P}_0 = \langle x_0, y_0, z_0 \rangle$ is a fixed point on the line and $\mathbf{d} = \langle a, b, c \rangle$ is a direction vector, the position of any point on the line can be expressed as:

$$\mathbf{r}(t) = \mathbf{P}_0 + t\mathbf{d}, \quad t \in \mathbb{R}.$$

Here:

- $\mathbf{P}_0$ specifies the starting point of the line.
- $\mathbf{d}$ indicates the direction and orientation of the line.
- The parameter t determines the location along the line: $t > 0$ moves in the direction of $\mathbf{d}$, $t < 0$ moves in the opposite direction, and $t = 0$ corresponds to the point $\mathbf{P}_0$.

This parametric representation makes clear that a line is defined by a starting point and a direction vector, with the parameter t serving as a simple scalar that scales the direction vector.

Generalizing to Parametric Curves

The simplicity of the line inspires us to generalize this approach to more complex curves. Instead of restricting each coordinate to vary linearly with t , we allow each coordinate to be an independent function of t . A general parametric curve in three-dimensional space can then be expressed as:

$$\mathbf{r}(t) = \langle f(t), g(t), h(t) \rangle, \quad t \in I,$$

where $f(t)$, $g(t)$, and $h(t)$ are continuous functions defined on an interval I . These functions describe the evolution of the curve's x -, y -, and z -coordinates, respectively, as the parameter t varies. This representation of curves as vector-valued functions allows us to extend the linear model of lines to a broader class of paths, capturing motion and shape in space with remarkable flexibility.

Note: In two dimensions, the parametric equation $\mathbf{r}(t) = \langle t, f(t) \rangle$ essentially represents the graph of the function $y = f(x)$. Here, t serves as both the parameter and the x -coordinate, making it straightforward to visualize the curve as the usual plot of y versus x .

Common Parametric Curves

Consider the following examples to illustrate the generalization from lines to more complex curves:

- **Circle in the xy -Plane:** A circle of radius R , centered at the origin, can be parameterized as:

$$\mathbf{r}(t) = \langle R \cos(t), R \sin(t), 0 \rangle, \quad t \in [0, 2\pi].$$

Here, t represents the angle, in radians, that the position vector makes with the positive x -axis.





Figure 2.1: Circle in the xy -plane with radius R .

- **Helix:** A helix is a three-dimensional curve that spirals around an axis. For instance, a helix with radius R and constant upward motion along the z -axis can be expressed as:

$$\mathbf{r}(t) = \langle R \cos(t), R \sin(t), ct \rangle, \quad t \in \mathbb{R},$$

where c determines the rate of ascent along the z -axis.

Figure 2.2: Helix with radius R spiraling around the z -axis.

- **Cycloid:** A cycloid is the path traced by a point on the rim of a wheel as it rolls along a straight line. Assuming the wheel has radius R , and it starts rolling from the origin, the parametric equation for the cycloid is:

$$\mathbf{r}(t) = \langle R(t - \sin(t)), R(1 - \cos(t)), 0 \rangle, \quad t \in \mathbb{R}.$$




Figure 2.3: Cycloid traced by a point on the rim of a rolling wheel.

These examples illustrate how allowing the components of $\mathbf{r}(t)$ to vary independently enables the description of diverse paths in space. The cycloid, for instance, captures a real-world motion, while the helix and circle provide insights into more abstract trajectories.

Properties of Parametric Curves

Understanding the properties of parametric curves is essential for analyzing their behavior. Here, we explore several key properties:

- **Orientation:** One of the primary advantages of parametric equations is their ability to encode the *orientation* of a curve. The parameter t determines the direction of

traversal. For instance, in the parametric equation of a circle:

$$\mathbf{r}(t) = \langle \cos(t), \sin(t), 0 \rangle, \quad t \in [0, 2\pi],$$

the curve is traversed counterclockwise in the xy -plane as t increases. Reversing the parameter (e.g., substituting $t \rightarrow -t$) reverses the orientation, resulting in clockwise traversal.

- **Smoothness:** A parametric curve $\mathbf{r}(t) = \langle f(t), g(t), h(t) \rangle$ is *smooth* if its components $f(t), g(t), h(t)$ are differentiable and $\mathbf{r}'(t) \neq \mathbf{0}$ for all t in the interval of interest. Smoothness ensures that the curve has no abrupt changes in direction or cusps.
- **Reparameterization:** A parametric curve can often be reparameterized to simplify its representation or emphasize certain properties. For example, reparameterizing a curve by its arc length s provides a direct measure of distance traveled along the curve. Arc length reparameterization is a key topic in the calculus of vector-valued functions, as it allows for uniform treatment of curves regardless of their initial parameterization. We will explore this later.

Applications of Parametric Equations

Parametric equations are foundational in physics, engineering, and computer graphics, where they are used to model motion, design structures, and render visual effects. Examples include:

- **Modeling Motion:** The trajectory of a projectile under gravity can be described parametrically as:

$$\mathbf{r}(t) = \langle v_0 \cos(\theta)t, v_0 \sin(\theta)t - \frac{1}{2}gt^2, 0 \rangle,$$

where v_0 is the initial velocity, θ is the launch angle, and g is the acceleration due to gravity.

- **Kinematics of Rigid Bodies:** In robotics, the position and orientation of a robotic arm are described using parametric equations, enabling precise control of movement.
- **Rendering Curves in 3D Graphics:** Parametric equations are used to generate and manipulate curves in computer graphics, such as Bézier curves in animations and CAD design.

As we delve into the calculus of vector-valued functions, the ability to differentiate and integrate parametric equations will allow us to analyze motion and physical systems in unprecedented detail. For instance, consider a particle moving along a winding path. At any instant t , its position $\mathbf{r}(t)$ is determined by the functions $f(t), g(t)$, and $h(t)$. The derivatives of these functions will later help us describe the particle's velocity and acceleration, revealing insights into its behavior at each point along the curve. With this generalization in place, we are now ready to explore the calculus of vector-valued functions, which provides tools to analyze change and motion along these curves.

Calculus of Vector-Valued Functions

Now that we have established the foundation of parametric equations and their representation through vector-valued functions, we move to the calculus of these functions. This section explores the derivatives and integrals of vector-valued functions, tools that enable us to analyze curves and change in space. By extending the familiar principles of single-variable calculus to the vector domain, we unlock a powerful framework for understanding the behavior of dynamic systems.

Derivatives of Vector-Valued Functions

To begin, let us consider the vector-valued function:

$$\mathbf{r}(t) = \langle f(t), g(t), h(t) \rangle,$$

where $f(t)$, $g(t)$, and $h(t)$ are differentiable scalar functions. The derivative of $\mathbf{r}(t)$ with respect to t is defined component-wise as:

$$\mathbf{r}'(t) = \langle f'(t), g'(t), h'(t) \rangle.$$

This derivative represents the rate of change of the vector-valued function with respect to the parameter t . Geometrically, $\mathbf{r}'(t)$ is a vector tangent to the curve defined by $\mathbf{r}(t)$ at the point corresponding to t . Its direction indicates how the curve is changing at that point. For example, if $\mathbf{r}'(t)$ points upward, the curve is ascending; if it points downward, the curve is descending.

Higher derivatives are also defined component-wise. The second derivative of $\mathbf{r}(t)$ is given by:

$$\mathbf{r}''(t) = \langle f''(t), g''(t), h''(t) \rangle,$$

which provides information about the concavity of the curve or how sharply it is bending at each point. These higher-order derivatives are critical for analyzing properties such as increasing or decreasing behavior, inflection points, and curvature.

The concept of component-wise differentiation not only simplifies calculations but also lays the foundation for generalizing change in multidimensional contexts. By treating each coordinate function $f(t)$, $g(t)$, $h(t)$ independently, we maintain a structured approach to understanding how the entire system evolves.

▲ **Example 2.1.1** Suppose a curve is defined by the vector-valued function:

$$\mathbf{r}(t) = \langle t^3, \sin(t), e^t \rangle.$$

Determine whether each component is increasing, decreasing, or concave at $t = \pi$.

Solution To analyze the behavior of each component at $t = \pi$, we compute the first

and second derivatives of $\mathbf{r}(t)$:

$$\mathbf{r}'(t) = \langle 3t^2, \cos(t), e^t \rangle, \quad \mathbf{r}''(t) = \langle 6t, -\sin(t), e^t \rangle.$$

At $t = \pi$, we evaluate the derivatives component-wise:

$$\text{First derivative: } \mathbf{r}'(\pi) = \langle 3\pi^2, \cos(\pi), e^\pi \rangle = \langle 3\pi^2, -1, e^\pi \rangle,$$

$$\text{Second derivative: } \mathbf{r}''(\pi) = \langle 6\pi, -\sin(\pi), e^\pi \rangle = \langle 6\pi, 0, e^\pi \rangle.$$

Component-wise analysis:

- $x(t) = t^3$: The first derivative $3\pi^2 > 0$ indicates $x(t)$ is increasing. The second derivative $6\pi > 0$ confirms $x(t)$ is concave upward.
- $y(t) = \sin(t)$: The first derivative $\cos(\pi) = -1$ indicates $y(t)$ is decreasing. The second derivative $-\sin(\pi) = 0$ suggests no concavity at this point.
- $z(t) = e^t$: The first derivative $e^\pi > 0$ indicates $z(t)$ is increasing. The second derivative $e^\pi > 0$ confirms $z(t)$ is concave upward.

Thus, $x(t)$ and $z(t)$ are increasing and concave upward, while $y(t)$ is decreasing with no concavity at $t = \pi$.

The analysis of derivatives provides a systematic way to study the behavior of vector-valued functions. By extending these principles, we can explore higher-order changes and geometric properties of curves in space.

Integrals of Vector-Valued Functions

Just as derivatives measure rates of change, integrals of vector-valued functions accumulate quantities over an interval. Given a vector-valued function:

$$\mathbf{r}(t) = \langle f(t), g(t), h(t) \rangle,$$

the integral of $\mathbf{r}(t)$ with respect to t is defined component-wise:

$$\int \mathbf{r}(t) dt = \left\langle \int f(t) dt, \int g(t) dt, \int h(t) dt \right\rangle + \mathbf{C},$$

where $\mathbf{C} = \langle C_1, C_2, C_3 \rangle$ is a constant vector of integration.

Integrals of vector-valued functions allow us to reconstruct curves, compute displacements, and analyze the accumulation of quantities across space. This process is inherently component-wise, treating each spatial dimension independently, while combining them to form the overall trajectory or system.

Consider a general situation where a vector-valued function $\mathbf{v}(t)$ represents some change,

such as motion or growth, in three-dimensional space. Integrating $\mathbf{v}(t)$ over time allows us to recover the function $\mathbf{r}(t)$ that describes the system:

$$\mathbf{r}(t) = \int \mathbf{v}(t) dt = \left\langle \int v_x(t) dt, \int v_y(t) dt, \int v_z(t) dt \right\rangle + \mathbf{r}_0,$$

where $\mathbf{r}_0 = \langle x_0, y_0, z_0 \rangle$ is the initial condition or position at $t = 0$.

This approach highlights the power of vector-valued integration in understanding phenomena that depend on independent contributions along multiple dimensions. Similarly, if $\mathbf{a}(t) = \mathbf{r}''(t)$ is a second derivative, integrating $\mathbf{a}(t)$ twice recovers $\mathbf{r}(t)$, with each step of integration requiring the addition of constants reflecting initial conditions.

By framing integrals component-wise, we emphasize the generalizability of these operations to multidimensional systems, while preserving the simplicity and independence of single-variable integration.

▲ **Example 2.1.2** Let $\mathbf{F}(t) = \langle 3t^2, 2\sin t, e^{2t} \rangle$ be a vector-valued function defined on the interval $[0, 1]$.

1. Compute the definite integral $\int_0^1 \mathbf{F}(t) dt$.
2. Verify that $\int_0^1 \mathbf{F}'(t) dt = \mathbf{F}(1) - \mathbf{F}(0)$.
3. Explain why the Fundamental Theorem of Calculus applies component-wise to vector-valued functions.

Solution

1. To compute the definite integral of $\mathbf{F}(t)$, we integrate each component separately:

$$\int_0^1 \mathbf{F}(t) dt = \left\langle \int_0^1 3t^2 dt, \int_0^1 2\sin t dt, \int_0^1 e^{2t} dt \right\rangle.$$

Compute each integral:

First component:

$$\int_0^1 3t^2 dt = [t^3]_0^1 = 1^3 - 0^3 = 1.$$

Second component:

$$\int_0^1 2 \sin t \, dt = -2 \cos t \Big|_0^1 = (-2 \cos 1) - (-2 \cos 0) = (-2 \cos 1) + 2(1) = 2(1 - \cos 1).$$

Third component:

$$\int_0^1 e^{2t} \, dt = \left[\frac{1}{2} e^{2t} \right]_0^1 = \frac{1}{2} (e^2 - e^0) = \frac{1}{2} (e^2 - 1).$$

Therefore,

$$\int_0^1 \mathbf{F}(t) \, dt = \left\langle 1, 2(1 - \cos 1), \frac{1}{2}(e^2 - 1) \right\rangle.$$

2. First, compute $\mathbf{F}'(t)$:

$$\mathbf{F}'(t) = \left\langle \frac{d}{dt} 3t^2, \frac{d}{dt} 2 \sin t, \frac{d}{dt} e^{2t} \right\rangle = \langle 6t, 2 \cos t, 2e^{2t} \rangle.$$

Then compute the integral of $\mathbf{F}'(t)$:

$$\int_0^1 \mathbf{F}'(t) \, dt = \left\langle \int_0^1 6t \, dt, \int_0^1 2 \cos t \, dt, \int_0^1 2e^{2t} \, dt \right\rangle.$$

Compute each integral:

First component:

$$\int_0^1 6t \, dt = [3t^2]_0^1 = 3(1)^2 - 3(0)^2 = 3.$$

Second component:

$$\int_0^1 2 \cos t \, dt = 2 \sin t \Big|_0^1 = 2 \sin 1 - 2 \sin 0 = 2 \sin 1 - 0 = 2 \sin 1.$$

Third component:

$$\int_0^1 2e^{2t} \, dt = [e^{2t}]_0^1 = e^2 - e^0 = e^2 - 1.$$

Now compute $\mathbf{F}(1) - \mathbf{F}(0)$:

$$\mathbf{F}(1) = \langle 3(1)^2, 2 \sin 1, e^{2(1)} \rangle = \langle 3, 2 \sin 1, e^2 \rangle.$$

$$\mathbf{F}(0) = \langle 3(0)^2, 2 \sin 0, e^{2(0)} \rangle = \langle 0, 0, 1 \rangle.$$

Therefore,

$$\mathbf{F}(1) - \mathbf{F}(0) = \langle 3 - 0, 2 \sin 1 - 0, e^2 - 1 \rangle = \langle 3, 2 \sin 1, e^2 - 1 \rangle.$$

Comparing the results:

$$\int_0^1 \mathbf{F}'(t) dt = \langle 3, 2 \sin 1, e^2 - 1 \rangle = \mathbf{F}(1) - \mathbf{F}(0).$$

Thus, we have verified that $\int_0^1 \mathbf{F}'(t) dt = \mathbf{F}(1) - \mathbf{F}(0)$.

3. The Fundamental Theorem of Calculus states that for a real-valued differentiable function f :

$$\int_a^b f'(t) dt = f(b) - f(a).$$

For vector-valued functions, each component is a real-valued function. Therefore, we can apply the Fundamental Theorem of Calculus to each component individually:

$$\begin{aligned} \int_a^b \mathbf{F}'(t) dt &= \left\langle \int_a^b F_1'(t) dt, \int_a^b F_2'(t) dt, \int_a^b F_3'(t) dt \right\rangle \\ &= \langle F_1(b) - F_1(a), F_2(b) - F_2(a), F_3(b) - F_3(a) \rangle = \mathbf{F}(b) - \mathbf{F}(a). \end{aligned}$$

This shows that the Fundamental Theorem of Calculus applies component-wise to vector-valued functions because integration and differentiation are performed separately on each real-valued component function.

Applications of Calculus with Vector-Valued Functions

The calculus of vector-valued functions has a wide range of applications in physics, engineering, and computer science. Some examples include:

- **Kinematics:** Derivatives and integrals of position vectors describe velocity, acceleration, and displacement, essential for analyzing the motion of objects.
- **Electromagnetism:** In physics, vector-valued functions are used to model electric and magnetic fields, with integrals representing quantities like flux.
- **Computer Graphics:** Derivatives of vector-valued functions are used to compute normals and tangents, while integrals are employed in generating smooth curves and surfaces.

By understanding the calculus of vector-valued functions, we gain the ability to analyze and predict the behavior of dynamic systems in space, paving the way for more advanced topics in multivariable calculus.

▲ **Example 2.1.3** Consider the parametric curve defined by

$$\mathbf{r}(t) = \langle e^t, e^{-t}, t \rangle, \quad t \in \mathbb{R}.$$

1. Show that this curve lies on the surface defined by $xy = 1$.
2. Find the equation of the tangent line to the curve at the point where $t = 0$.

Solution

1. To show that the curve lies on the surface $xy = 1$, substitute $x = e^t$ and $y = e^{-t}$ into the equation:

$$xy = e^t \cdot e^{-t} = e^{t-t} = e^0 = 1.$$

Therefore, for all $t \in \mathbb{R}$, the point (x, y, z) lies on the surface $xy = 1$.

2. The tangent vector to the curve at t is given by:

$$\mathbf{r}'(t) = \left\langle \frac{d}{dt}e^t, \frac{d}{dt}e^{-t}, \frac{d}{dt}t \right\rangle = \langle e^t, -e^{-t}, 1 \rangle.$$

At $t = 0$:

$$\mathbf{r}(0) = \langle e^0, e^0, 0 \rangle = \langle 1, 1, 0 \rangle,$$

$$\mathbf{r}'(0) = \langle e^0, -e^0, 1 \rangle = \langle 1, -1, 1 \rangle.$$

Since we are working in three dimensions, we use the parametric equation of a line:

$$\mathbf{r}(s) = \mathbf{P}_0 + s\mathbf{d},$$

where $\mathbf{P}_0$ is a point on the line and $\mathbf{d}$ is the direction vector of the line.

In this context:

$$\mathbf{P}_0 = \mathbf{r}(0) = \langle 1, 1, 0 \rangle, \quad \mathbf{d} = \mathbf{r}'(0) = \langle 1, -1, 1 \rangle.$$

Therefore, the equation of the tangent line at $t = 0$ is:

$$\mathbf{L}(s) = \mathbf{P}_0 + s\mathbf{d} = \langle 1, 1, 0 \rangle + s\langle 1, -1, 1 \rangle.$$

Simplifying, we obtain:

$$\mathbf{L}(s) = \langle 1 + s, 1 - s, 0 + s \rangle = \langle 1 + s, 1 - s, s \rangle.$$

This parametric equation represents the tangent line to the curve at $t = 0$.

Explanation: In three-dimensional space, lines are described using vector equations. The general form is:

$$\mathbf{r}(s) = \mathbf{P}_0 + s\mathbf{d},$$

where $\mathbf{P}_0$ is a fixed point on the line and $\mathbf{d}$ is a direction vector parallel to the line. The parameter s allows us to trace every point along the line. In this problem, $\mathbf{r}(0)$ provides the point of tangency, and $\mathbf{r}'(0)$ gives the direction of the tangent line, which is why we use them in constructing the equation.

Lecture 2.2: Motion Along a Curve

Building on our understanding of vector-valued functions and their derivatives, we now turn to the study of motion along a curve. This section explores how the calculus of vector-valued functions is applied to describe physical motion in space. Specifically, we examine how position, velocity, and acceleration work together to describe the behavior of moving objects, using the calculus of vector-valued functions.

Motion Along a Curve

When an object moves along a curve in three-dimensional space, its position at time t is described by a vector-valued function:

$$\mathbf{r}(t) = \langle x(t), y(t), z(t) \rangle,$$

where $x(t)$, $y(t)$, and $z(t)$ are the coordinates of the object as functions of time. The path traced by $\mathbf{r}(t)$ represents the trajectory of the object. Analyzing this motion relies on the concepts of derivatives and integrals to extract information about how the position vector changes over time and to reconstruct paths from rates of change.

The Velocity Vector $\mathbf{v}(t)$ is defined as the derivative of the position vector:

$$\mathbf{v}(t) = \frac{d\mathbf{r}}{dt} = \langle x'(t), y'(t), z'(t) \rangle.$$

This vector is tangent to the curve at the point $\mathbf{r}(t)$, indicating the direction of motion at time t . Its magnitude represents the speed of the object:

$$|\mathbf{v}(t)| = \sqrt{x'(t)^2 + y'(t)^2 + z'(t)^2}.$$

Geometrically, the velocity vector connects the rate of change of the coordinates x , y , and z , capturing the instantaneous direction and rate at which the object moves along the curve.

The Acceleration Vector $\mathbf{a}(t)$ is defined as the derivative of the velocity vector:

$$\mathbf{a}(t) = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{r}}{dt^2} = \langle x''(t), y''(t), z''(t) \rangle.$$

The acceleration vector quantifies how the velocity vector changes over time, indicating whether the object is speeding up, slowing down, or changing direction. The magnitude of the acceleration vector:

$$|\mathbf{a}(t)| = \sqrt{x''(t)^2 + y''(t)^2 + z''(t)^2},$$

provides the total rate of change of velocity.

Integrals and Motion Reconstruction: In addition to derivatives, integrals play

a fundamental role in understanding motion. The velocity vector $\mathbf{v}(t)$ is often given as the rate of change of position, and integrating $\mathbf{v}(t)$ allows us to reconstruct the position vector:

$$\mathbf{r}(t) = \int \mathbf{v}(t) dt + \mathbf{r}_0,$$

where $\mathbf{r}_0$ is the initial position of the object. Similarly, the acceleration vector $\mathbf{a}(t)$, if known, can be integrated to recover the velocity:

$$\mathbf{v}(t) = \int \mathbf{a}(t) dt + \mathbf{v}_0,$$

where $\mathbf{v}_0$ is the initial velocity of the object. These integrations reveal how the motion accumulates over time, allowing us to describe both the trajectory and the forces acting on the object.

Both velocity and acceleration vectors are essential in understanding the dynamics of motion along a curve. For instance:

- If the velocity vector $\mathbf{v}(t)$ is constant, the object moves along the curve with uniform speed.
- If the acceleration vector $\mathbf{a}(t)$ is zero, the object moves at constant velocity, tracing a straight line in space.
- If $\mathbf{a}(t)$ is not zero, the motion becomes more complex, involving changes in both speed and direction.

The interplay between $\mathbf{v}(t)$, $\mathbf{a}(t)$, and their respective integrals provides insights into real-world motion, such as the trajectories of projectiles or the orbits of celestial bodies. In cases where motion occurs under uniform forces, such as gravity, the integration process enables the reconstruction of entire paths from basic physical laws. These relationships will be expanded upon in subsequent sections, where we explore specific examples and applications of motion in physical systems.

▲ **Example 2.2.1** Suppose a particle moves along a curve with velocity:

$$\mathbf{v}(t) = \langle 3t, 4, t^2 \rangle, \quad t \in [0, 2].$$

Find the displacement of the particle as well as total distance traveled.

Solution To find the displacement of the particle, we integrate the velocity vector:

$$\mathbf{r}(t) = \int \mathbf{v}(t) dt = \left\langle \int 3t dt, \int 4 dt, \int t^2 dt \right\rangle + \mathbf{r}_0.$$

Assuming the particle starts at the origin ($\mathbf{r}_0 = \langle 0, 0, 0 \rangle$), the components of $\mathbf{r}(t)$ are

computed as follows:

$$\begin{aligned}\int 3t \, dt &= \frac{3t^2}{2}, \\ \int 4 \, dt &= 4t, \\ \int t^2 \, dt &= \frac{t^3}{3}.\end{aligned}$$

Thus, the position function is:

$$\mathbf{r}(t) = \left\langle \frac{3t^2}{2}, 4t, \frac{t^3}{3} \right\rangle.$$

At $t = 2$, the displacement is:

$$\mathbf{r}(2) = \left\langle \frac{3(2)^2}{2}, 4(2), \frac{(2)^3}{3} \right\rangle = \left\langle 6, 8, \frac{8}{3} \right\rangle.$$

This vector gives the particle's position relative to its starting point.

To calculate the total distance traveled, we integrate the speed of the particle, which is the magnitude of the velocity:

$$|\mathbf{v}(t)| = \sqrt{(3t)^2 + 4^2 + (t^2)^2}.$$

The total distance is given by:

$$\int_0^2 |\mathbf{v}(t)| \, dt = \int_0^2 \sqrt{(3t)^2 + 4^2 + (t^2)^2} \, dt.$$

Simplifying the expression inside the square root:

$$|\mathbf{v}(t)| = \sqrt{9t^2 + 16 + t^4}.$$

Since this integral does not have a simple closed-form solution, numerical methods, such as Simpson's Rule or a computational tool, are typically used to evaluate it.

Projectile Motion

A classic example of motion along a curve is projectile motion, where an object is launched with an initial speed v_0 at an angle θ above the horizontal plane. The initial velocity vector is given by:

$$\mathbf{v}_0 = \langle v_{x0}, v_{y0}, v_{z0} \rangle = \langle v_0 \cos \theta, 0, v_0 \sin \theta \rangle,$$

assuming motion occurs in the xz -plane (no motion in the y -direction). The acceleration due to gravity acts downward along the z -axis, so the acceleration vector is:

$$\mathbf{a}(t) = \langle 0, 0, -g \rangle,$$

where g is the acceleration due to gravity. Integrating the acceleration vector with respect to time gives the velocity vector:

$$\mathbf{v}(t) = \int \mathbf{a}(t) dt + \mathbf{C}_1 = \langle v_0 \cos \theta, 0, v_0 \sin \theta - gt \rangle,$$

where $\mathbf{C}_1 = \mathbf{v}_0$ is the constant of integration representing the initial velocity. Integrating the velocity vector yields the position vector:

$$\mathbf{r}(t) = \int \mathbf{v}(t) dt + \mathbf{C}_2 = \langle v_0 \cos \theta t + x_0, y_0, v_0 \sin \theta t - \frac{1}{2}gt^2 + z_0 \rangle,$$

where $\mathbf{C}_2 = \langle x_0, y_0, z_0 \rangle$ is the initial position vector.

These equations describe the parabolic trajectory of a projectile. The horizontal motion (x -direction) is uniform, while the vertical motion (z -direction) is uniformly accelerated due to gravity.

The parametric equations for the motion are:

$$\begin{cases} x(t) = v_0 \cos \theta t + x_0, \\ y(t) = y_0, \\ z(t) = v_0 \sin \theta t - \frac{1}{2}gt^2 + z_0. \end{cases}$$

To eliminate t and find the equation of the trajectory, solve for t from $x(t)$:

$$t = \frac{x - x_0}{v_0 \cos \theta}.$$

Substitute t into $z(t)$:

$$z = v_0 \sin \theta \left(\frac{x - x_0}{v_0 \cos \theta} \right) - \frac{1}{2}g \left(\frac{x - x_0}{v_0 \cos \theta} \right)^2 + z_0.$$

Simplify:

$$z = (x - x_0) \tan \theta - \frac{g(x - x_0)^2}{2v_0^2 \cos^2 \theta} + z_0.$$

This equation represents the parabolic path of the projectile in the xz -plane.

Maximum Height

The maximum height $z_{\max}$ of the projectile can be found by analyzing the vertical motion. The vertical component of velocity is:

$$v_z(t) = v_0 \sin \theta - gt.$$

The maximum height occurs when $v_z(t) = 0$:

$$v_0 \sin \theta - gt_{\max} = 0 \implies t_{\max} = \frac{v_0 \sin \theta}{g}.$$

Substitute $t_{\max}$ into $z(t)$ to find $z_{\max}$:

$$z_{\max} = v_0 \sin \theta \left(\frac{v_0 \sin \theta}{g} \right) - \frac{1}{2}g \left(\frac{v_0 \sin \theta}{g} \right)^2 + z_0.$$

Simplify:

$$z_{\max} = \frac{v_0^2 \sin^2 \theta}{g} - \frac{v_0^2 \sin^2 \theta}{2g} + z_0 = \frac{v_0^2 \sin^2 \theta}{2g} + z_0.$$

Angle for Maximum Height

The maximum height $z_{\max}$ depends on $\sin^2 \theta$. Since $\sin \theta$ reaches its maximum value at $\theta = 90^\circ$, the maximum height is achieved when the projectile is launched vertically upward. Therefore, the angle that produces the maximum height is:

$$\theta_{\max_height} = 90^\circ.$$

Maximum Range

The horizontal range R of the projectile (assuming it lands at the same vertical position from which it was launched, i.e., $z = z_0$) is found by determining the time of flight t_{flight} when $z(t_{\text{flight}}) = z_0$:

Set $z(t_{\text{flight}}) = z_0$:

$$v_0 \sin \theta t_{\text{flight}} - \frac{1}{2}gt_{\text{flight}}^2 + z_0 = z_0.$$

Simplify:

$$v_0 \sin \theta t_{\text{flight}} - \frac{1}{2}gt_{\text{flight}}^2 = 0.$$

Factor out $t_{\text{flight}} \neq 0$:

$$t_{\text{flight}} = \frac{2v_0 \sin \theta}{g}.$$

The horizontal range is:

$$R = x(t_{\text{flight}}) - x_0 = v_0 \cos \theta t_{\text{flight}} = v_0 \cos \theta \left(\frac{2v_0 \sin \theta}{g} \right) = \frac{2v_0^2 \sin \theta \cos \theta}{g}.$$

Using the identity $\sin 2\theta = 2 \sin \theta \cos \theta$:

$$R = \frac{v_0^2 \sin 2\theta}{g}.$$




Figure 2.4: A 3D visualization of projectile motion on a golf course. The ball is launched in a parabolic trajectory, reaching the hole marked by the red flag.

Angle for Maximum Range

The maximum range occurs when $\sin 2\theta$ is maximized. Since $\sin 2\theta$ reaches its maximum value of 1 when $2\theta = 90^\circ$, i.e., $\theta = 45^\circ$:

$$\theta_{\text{max.range}} = 45^\circ.$$

Therefore, to achieve the maximum horizontal range, the projectile should be launched at an angle of 45° .

▲ **Example 2.2.2** A golfer is attempting to hit a golf ball over a sandtrap onto the green. The sandtrap is located between 50 meters and 70 meters from the golfer. The green starts at a distance of 70 meters from the golfer. The golfer can hit the ball with an initial speed of $v_0 = 45$ m/s. At what minimum angle θ above the horizontal must the golfer hit the ball to ensure the ball clears the sandtrap and lands on the green? Ignore air resistance and assume the ball lands at the same vertical level from which it was hit.

Solution To ensure the ball clears the sandtrap and lands on the green, the golfer must hit the ball so that it lands at a horizontal distance $x \geq 70$ meters from the starting point.

The horizontal range R of a projectile launched at an angle θ with initial speed v_0 and landing at the same vertical level is given by:

$$R = \frac{v_0^2 \sin 2\theta}{g},$$

where $g = 9.8 \text{ m/s}^2$ is the acceleration due to gravity.

We need to find the minimum angle θ such that $R \geq 70$ meters.

1. **Set up the inequality for the range:**

$$R = \frac{v_0^2 \sin 2\theta}{g} \geq 70.$$

2. **Solve for $\sin 2\theta$:**

$$\sin 2\theta \geq \frac{70g}{v_0^2}.$$

Substitute $v_0 = 45 \text{ m/s}$ and $g = 9.8 \text{ m/s}^2$:

$$\sin 2\theta \geq \frac{70 \times 9.8}{45^2} = \frac{686}{2025} \approx 0.338.$$

3. **Solve for 2θ :**

$$2\theta \geq \sin^{-1}(0.338).$$

Compute $\sin^{-1}(0.338)$:

$$2\theta \geq 19.78^\circ.$$

4. **Solve for θ :**

$$\theta \geq \frac{19.78^\circ}{2} = 9.89^\circ.$$

Conclusion: The golfer must hit the ball at a minimum angle of approximately 9.89° above the horizontal to ensure the ball clears the sandtrap and lands on the green.

Energy Considerations

The initial kinetic energy K of the projectile is:

$$K = \frac{1}{2}mv_0^2,$$

where m is the mass of the projectile. At the maximum height, the kinetic energy is:

$$K_{\text{max.height}} = \frac{1}{2}m(v_0 \cos \theta)^2 = \frac{1}{2}mv_0^2 \cos^2 \theta.$$

The potential energy increase is:

$$\Delta U = mgz_{\text{max}} = mg \left(\frac{v_0^2 \sin^2 \theta}{2g} + z_0 \right) = \frac{1}{2}mv_0^2 \sin^2 \theta + mgz_0.$$

This shows the conservation of mechanical energy:

$$K = K_{\text{max.height}} + \Delta U.$$

▲ Example 2.2.3 An object is launched from the origin ($x_0 = z_0 = 0$) with an initial speed $v_0 = 20$ m/s at an angle $\theta = 45^\circ$.

1. Derive the parametric equations for the motion of the object, assuming acceleration due to gravity is $g = 9.8$ m/s².
2. Find the maximum height of the object using kinematic equations.
3. Determine the total time of flight.
4. Calculate the horizontal range of the projectile.
5. Using energy considerations, calculate the maximum height the object reaches.

Solution**1. Parametric Equations.**

Decompose the initial velocity into horizontal and vertical components:

$$v_{x0} = v_0 \cos \theta = 20 \cos 45^\circ = 20 \left(\frac{\sqrt{2}}{2} \right) = 10\sqrt{2} \text{ m/s},$$

$$v_{z0} = v_0 \sin \theta = 20 \sin 45^\circ = 20 \left(\frac{\sqrt{2}}{2} \right) = 10\sqrt{2} \text{ m/s}.$$

The parametric equations are:

$$x(t) = v_{x0}t = 10\sqrt{2}t,$$

$$z(t) = v_{z0}t - \frac{1}{2}gt^2 = 10\sqrt{2}t - 4.9t^2.$$

2. Maximum Height Using Kinematics.

The maximum height occurs when the vertical velocity becomes zero:

$$v_z(t) = v_{z0} - gt = 0 \implies t_{\max} = \frac{v_{z0}}{g} = \frac{10\sqrt{2}}{9.8} \approx 1.44 \text{ s}.$$

Substitute $t_{\max}$ into $z(t)$:

$$\begin{aligned} z_{\max} &= 10\sqrt{2} \left(\frac{10\sqrt{2}}{9.8} \right) - 4.9 \left(\frac{10\sqrt{2}}{9.8} \right)^2 \\ &= \frac{200}{9.8} - 4.9 \left(\frac{200}{(9.8)^2} \right) \\ &= \frac{200}{9.8} - \frac{200}{9.8} \\ &= \frac{200}{9.8} - \frac{200}{9.8} = 10.204 \text{ m}. \end{aligned}$$

So, the maximum height is approximately 10.2 m.

3. Total Time of Flight.

The total time is when the projectile returns to $z = 0$:

$$z(t) = 0 \implies 10\sqrt{2}t - 4.9t^2 = 0.$$

Factor out $t \neq 0$:

$$t(10\sqrt{2} - 4.9t) = 0 \implies t_{\text{total}} = \frac{10\sqrt{2}}{4.9} \approx 2.04 \text{ s}.$$

4. Horizontal Range.

The horizontal range is:

$$x_{\text{range}} = v_{x0} \times t_{\text{total}} = 10\sqrt{2} \times 2.04 \approx 28.9 \text{ m}.$$

5. Maximum Height Using Energy Considerations.

We will use the conservation of mechanical energy to find $z_{\max}$.

(a) **Initial Kinetic Energy:**

$$K_i = \frac{1}{2}mv_0^2 = \frac{1}{2}m(20)^2 = 200m.$$

(b) **Kinetic Energy at Maximum Height:**

At maximum height, vertical velocity is zero, so:

$$K_{\max} = \frac{1}{2}mv_{x0}^2 = \frac{1}{2}m(10\sqrt{2})^2 = 100m.$$

(c) **Potential Energy at Maximum Height:**

$$U_{\max} = mgz_{\max}.$$

(d) **Conservation of Mechanical Energy:**

$$K_i = K_{\max} + U_{\max}.$$

(e) **Solve for $z_{\max}$:**

$$200m = 100m + mgz_{\max} \implies mgz_{\max} = 100m.$$

Cancel m from both sides:

$$gz_{\max} = 100 \implies z_{\max} = \frac{100}{g} = \frac{100}{9.8} \approx 10.204 \text{ m}.$$

Conclusion: The maximum height calculated using energy considerations is approximately 10.2 m, which matches the result obtained using kinematic equations.

Lecture 2.3: The TNB Frame, Curvature and Torsion

As we extend our study of motion in space, we now introduce the TNB frame—a powerful tool for analyzing the geometry of curves and the motion of objects along those curves. The TNB frame consists of three mutually orthogonal vectors: the tangent vector, the normal vector, and the binormal vector. Together, these vectors form a moving frame that evolves along the curve, providing a dynamic coordinate system to describe its behavior. This section also introduces the Frenet-Serret equations, which govern the interplay between these vectors, and discusses applications of the TNB frame to motion in space.

The TNB Frame

In previous sections, we discussed the derivatives of vector-valued functions, which naturally give rise to the tangent vector ($\mathbf{T}(t)$), and explored the normal vector ($\mathbf{N}(t)$) as a tool for understanding how curves bend. Building upon this foundation, we now introduce the binormal vector ($\mathbf{B}(t)$) to complete the TNB frame. Together, these vectors form an orthonormal basis that evolves along the curve, providing a dynamic coordinate system tied directly to the curve's intrinsic geometry and motion.

Consider a smooth curve parameterized by a vector-valued function:

$$\mathbf{r}(t) = \langle x(t), y(t), z(t) \rangle, \quad t \in I,$$

where $\mathbf{r}'(t) \neq \mathbf{0}$ for all t . The TNB frame is constructed as follows:

- **Tangent Vector ($\mathbf{T}(t)$):** The unit tangent vector is defined as:

$$\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|}.$$

It points in the direction of motion, describing the instantaneous direction of the curve at each point.

- **Normal Vector ($\mathbf{N}(t)$):** The principal normal vector is defined as:

$$\mathbf{N}(t) = \frac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|}.$$

It quantifies how the curve bends by pointing toward the center of curvature at each point.

- **Binormal Vector ($\mathbf{B}(t)$):** The binormal vector is defined by the cross product:

$$\mathbf{B}(t) = \mathbf{T}(t) \times \mathbf{N}(t).$$

It is orthogonal to both $\mathbf{T}(t)$ and $\mathbf{N}(t)$, completing the right-handed orthonormal basis.

These vectors collectively form the TNB frame, providing a coordinate system uniquely tied to the geometry of the curve. The evolution of the TNB frame along the curve encodes information about its curvature and torsion. Below is an example of a TNB frame of a parametric curve that is going around a cylinder:




Figure 2.5: A TNB Frame on a parametric curve around a scaled cylinder

▲ **Example 2.3.1** A particle moves along the space curve:

$$\mathbf{r}(t) = \langle t, t^2, t^3 \rangle, \quad t \in [0, 2].$$

Find the TNB frame vectors $(\mathbf{T}(t), \mathbf{N}(t), \mathbf{B}(t))$ at $t = 1$.

Solution

1. Compute derivatives of $\mathbf{r}(t)$:

$$\mathbf{r}'(t) = \langle 1, 2t, 3t^2 \rangle, \quad \mathbf{r}'(1) = \langle 1, 2, 3 \rangle, \quad |\mathbf{r}'(1)| = \sqrt{1^2 + 2^2 + 3^2} = \sqrt{14}.$$

$$\mathbf{r}''(t) = \langle 0, 2, 6t \rangle, \quad \mathbf{r}''(1) = \langle 0, 2, 6 \rangle.$$

2. Compute $\mathbf{T}(1)$:

$$\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|}, \quad \mathbf{T}(1) = \frac{\langle 1, 2, 3 \rangle}{\sqrt{14}} = \left\langle \frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}} \right\rangle.$$

3. Compute $\mathbf{N}(1)$: Using:

$$\mathbf{N}(t) = \frac{\mathbf{r}''(t) - (\mathbf{r}''(t) \cdot \mathbf{T}(t)) \mathbf{T}(t)}{|\mathbf{r}''(t) - (\mathbf{r}''(t) \cdot \mathbf{T}(t)) \mathbf{T}(t)|},$$

compute the dot product:

$$\mathbf{r}''(1) \cdot \mathbf{T}(1) = \langle 0, 2, 6 \rangle \cdot \left\langle \frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}} \right\rangle = \frac{22}{\sqrt{14}}.$$

Subtract:

$$\mathbf{r}''(1) - (\mathbf{r}''(1) \cdot \mathbf{T}(1)) \mathbf{T}(1) = \langle 0, 2, 6 \rangle - \frac{22}{14} \langle 1, 2, 3 \rangle = \left\langle -\frac{11}{7}, -\frac{8}{7}, \frac{9}{7} \right\rangle.$$

Compute the magnitude:

$$|\mathbf{N}_{\text{numerator}}| = \sqrt{\left(-\frac{11}{7}\right)^2 + \left(-\frac{8}{7}\right)^2 + \left(\frac{9}{7}\right)^2} = \frac{\sqrt{266}}{7}.$$

Thus:

$$\mathbf{N}(1) = \left\langle -\frac{11}{\sqrt{266}}, -\frac{8}{\sqrt{266}}, \frac{9}{\sqrt{266}} \right\rangle.$$

4. Compute $\mathbf{B}(1)$: Using:

$$\mathbf{B}(1) = \mathbf{T}(1) \times \mathbf{N}(1),$$

compute the cross product:

$$\mathbf{B}(1) = \left\langle \frac{3}{\sqrt{266}}, -\frac{3}{\sqrt{266}}, \frac{1}{\sqrt{266}} \right\rangle.$$

At $t = 1$, the TNB frame is:

$$\begin{aligned} \mathbf{T}(1) &= \left\langle \frac{1}{\sqrt{14}}, \frac{2}{\sqrt{14}}, \frac{3}{\sqrt{14}} \right\rangle, \\ \mathbf{N}(1) &= \left\langle -\frac{11}{\sqrt{266}}, -\frac{8}{\sqrt{266}}, \frac{9}{\sqrt{266}} \right\rangle, \\ \mathbf{B}(1) &= \left\langle \frac{3}{\sqrt{266}}, -\frac{3}{\sqrt{266}}, \frac{1}{\sqrt{266}} \right\rangle. \end{aligned}$$

Curvature

Beyond position, velocity, and acceleration, the curvature of a curve provides insights into how sharply the trajectory bends at each point. It quantifies the rate at which the direction

of the curve changes as you move along it. For a smooth curve defined by the vector-valued function $\mathbf{r}(t)$, the curvature $\kappa(t)$ at a point is given by:

$$\kappa(t) = \frac{|\mathbf{r}'(t) \times \mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \frac{|\mathbf{v}(t) \times \mathbf{a}(t)|}{|\mathbf{v}(t)|^3} = \frac{|\mathbf{T}'(s)|}{|\mathbf{r}'(s)|}$$

Here:

- $\mathbf{r}'(t)$ is the first derivative of $\mathbf{r}(t)$, representing the velocity vector.
- $\mathbf{r}''(t)$ is the second derivative of $\mathbf{r}(t)$, representing the acceleration vector.
- The cross product $\mathbf{r}'(t) \times \mathbf{r}''(t)$ captures the tendency of the curve to twist out of a straight path, emphasizing the interaction between velocity and acceleration.

In the case of **plane curves** (curves that lie entirely in a plane), there is a simplified formula for curvature. If the curve is given parametrically by $\mathbf{r}(t) = \langle x(t), y(t) \rangle$, the curvature $\kappa(t)$ can be calculated as:

$$\kappa(t) = \frac{|x'(t)y''(t) - y'(t)x''(t)|}{([x'(t)]^2 + [y'(t)]^2)^{3/2}}$$

This formula is derived from the general vector formula for curvature by evaluating the cross product and magnitudes in two dimensions. When dealing with a single-variable function $y = f(x)$, we can consider it as a plane curve by parameterizing it as:

$$\mathbf{r}(t) = \langle t, f(t) \rangle.$$

In this case, the derivatives are:

$$x(t) = t, \quad y(t) = f(t), \quad x'(t) = 1, \quad y'(t) = f'(t), \quad x''(t) = 0, \quad y''(t) = f''(t).$$

Substituting into the plane curve curvature formula:

$$\kappa(t) = \frac{|x'(t)y''(t) - y'(t)x''(t)|}{([x'(t)]^2 + [y'(t)]^2)^{3/2}} = \frac{|1 \cdot f''(t) - f'(t) \cdot 0|}{([1]^2 + [f'(t)]^2)^{3/2}} = \frac{|f''(t)|}{(1 + [f'(t)]^2)^{3/2}}.$$

Since $x = t$, we can write $\kappa(x)$ as:

$$\kappa(x) = \frac{|f''(x)|}{(1 + [f'(x)]^2)^{3/2}}.$$

This formula gives the curvature of the graph of a function $y = f(x)$ at any point x . This progression shows how the general formula simplifies under specific conditions, making it easier to compute curvature in various contexts.

Curvature has a natural interpretation: it describes how much the direction of the **tangent vector** $\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|}$ changes per unit of **arc length**. This highlights a critical property of curvature: it is independent of the speed at which the curve is traversed.

Whether a particle moves along the curve quickly or slowly, the curvature remains tied solely to the trajectory's geometry.

The plane curve curvature formula emphasizes how curvature depends on the rates of change of the components of the curve. The numerator $x'(t)y''(t) - y'(t)x''(t)$ represents the rate at which the direction is changing, while the denominator normalizes this by the speed to the third power, ensuring that curvature is a geometric property of the curve itself, not dependent on parameterization.

Osculating Circles

The osculating circle at a point on the curve is the unique circle that shares the same tangent and curvature as the curve at that point. Its radius R , known as the *radius of curvature*, is inversely proportional to the curvature:

$$R = \frac{1}{\kappa}.$$

For straight lines, where $\kappa = 0$, the radius of curvature is infinite, reflecting the absence of bending. Conversely, for a tightly curved loop, κ is large, and R is small. The osculating circle provides a geometric representation of curvature, capturing the curve's local behavior with remarkable accuracy. These ideas are not merely theoretical—they allow for practical analysis of paths and trajectories. For example, in designing smooth transitions in roads or roller coasters, one must ensure that the curvature does not exceed safe limits, as excessive bending introduces significant forces.

Motion Context

Curvature also reflects constraints imposed by motion. Consider a car driving along a curved path. The curvature dictates the centripetal force required to maintain the car's trajectory. A sharper turn (higher curvature) demands greater centripetal acceleration, which is proportional to κv^2 , where v is the car's speed. Similarly, in orbital mechanics, the curvature of a planet's trajectory provides insight into the forces acting upon it. The curvature is directly tied to the balance of gravitational pull and the tangential velocity of the orbiting body.

To illustrate the concept of curvature, consider the sine curve $\mathbf{r}(t) = \langle t, t^3 - 3t^2 + t - 2 \rangle$. This curve alternates between regions of high and low curvature. At peaks and troughs, where the curve bends most sharply, the curvature is maximized. In contrast, at inflection points, where the curve transitions between bending upward and downward, the curvature diminishes.




Figure 2.6: A polynomial and its curvature (κ) plotted together. The function is shown in blue, and the curvature is shown in dashed red, with a combined legend for clarity.

In Figure 2.6, the plot illustrates:

- The function $f(x)$, shown as a blue solid line, representing a cubic polynomial. It exhibits inflection points where the curvature changes sign, corresponding to transitions between concave up and concave down behavior.
- The curvature $\kappa(x)$, shown as a red dashed line, which has peaks at $x \approx 0$, $x \approx 1.5$, and $x \approx 2.5$, indicating regions of high curvature.
- Low curvature values for $\kappa(x)$ near $x = -1$, $x = 4$, and other regions where $f(x)$ is nearly linear.

Below is a python snippet that you can use to see a function with its curvature on the same plot.

```

1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 def plot_function_and_curvature(func, x_range, scaling_factor=1):
5     # Generate x values
6     x = np.linspace(x_range[0], x_range[1], 1000)
7     dx = x[1] - x[0] # Assuming uniform spacing
8
9     # Compute y = f(x)
10    y = func(x)
11
12    # First derivative f'(x) using numerical differentiation

```

```

13 f_prime = np.gradient(y, dx)
14
15 # Second derivative f''(x)
16 f_double_prime = np.gradient(f_prime, dx)
17
18 # Compute curvature  $k = |f''(x)| / (1 + [f'(x)]^2)^{3/2}$ 
19 curvature = np.abs(f_double_prime) / (1 + f_prime**2)**1.5
20 curvature *= scaling_factor # Apply scaling if needed
21
22 # Plotting
23 fig, ax1 = plt.subplots(figsize=(10, 6))
24
25 # Plot the function on the left y-axis
26 color1 = 'tab:blue'
27 ax1.set_xlabel('x')
28 ax1.set_ylabel('Function', color=color1)
29 ax1.plot(x, y, label='Function', color=color1, linewidth=2)
30 ax1.tick_params(axis='y', labelcolor=color1)
31 ax1.grid(True)
32
33 # Create a second y-axis for the curvature
34 ax2 = ax1.twinx()
35
36 color2 = 'tab:red'
37 ax2.set_ylabel('Curvature', color=color2)
38 ax2.plot(x, curvature, label='Curvature', color=color2, linestyle='--', linewidth=2)
39 ax2.tick_params(axis='y', labelcolor=color2)
40
41 # Combine legends
42 lines_1, labels_1 = ax1.get_legend_handles_labels()
43 lines_2, labels_2 = ax2.get_legend_handles_labels()
44 ax1.legend(lines_1 + lines_2, labels_1 + labels_2, loc='upper right')
45
46 plt.title('Function and Curvature')
47 plt.show()
48
49 # Example usage with a polynomial function
50 def polynomial_function(x):
51     return x**3 - 3*x**2 + x - 2
52
53 plot_function_and_curvature(polynomial_function, x_range=(-1, 4))

```

▲ **Example 2.3.2** An engineer is designing a curved road described by the parametric equations:

$$\mathbf{r}(t) = \langle t, t^2 \rangle, \quad t \in [0, 5].$$

1. Compute the curvature $\kappa(t)$ of the road at any point t .

2. Determine the point along the road where the curvature is maximized within the interval $t \in [0, 5]$.
3. Discuss how the curvature affects the design and safety of the road.

Solution

1. First, we identify the components of $\mathbf{r}(t)$:

$$x(t) = t, \quad y(t) = t^2.$$

Compute the first derivatives:

$$x'(t) = 1, \quad y'(t) = 2t.$$

Compute the second derivatives:

$$x''(t) = 0, \quad y''(t) = 2.$$

The curvature $\kappa(t)$ for plane curves is given by:

$$\kappa(t) = \frac{|x'(t)y''(t) - y'(t)x''(t)|}{([x'(t)]^2 + [y'(t)]^2)^{3/2}}.$$

Substitute the derivatives:

$$\kappa(t) = \frac{|(1)(2) - (2t)(0)|}{(1^2 + (2t)^2)^{3/2}} = \frac{2}{(1 + 4t^2)^{3/2}}.$$

2. To find where the curvature is maximized, we can analyze $\kappa(t)$:

$$\kappa(t) = \frac{2}{(1 + 4t^2)^{3/2}}.$$

Since the denominator increases with t , the curvature decreases as t increases. Therefore, the maximum curvature occurs at $t = 0$:

$$\kappa_{\max} = \kappa(0) = \frac{2}{(1 + 0)^{3/2}} = 2.$$

3. Implications:

- **Design and Safety:** The highest curvature at $t = 0$ indicates the sharpest bend in the road occurs at the starting point. High curvature requires careful design to ensure vehicles can navigate the curve safely without skidding or overturning.

- **Speed Limits:** Areas with higher curvature may require lower speed limits to maintain safety, as centripetal acceleration increases with curvature and speed.
- **Banking the Road:** Engineers might design the road with banking (tilting the road towards the center of the curve) at high-curvature sections to counteract lateral acceleration and enhance safety.
- **Driver Comfort:** Gradually decreasing curvature along the road (as t increases) provides a smoother driving experience, reducing the risk of accidents due to sudden changes in direction.

Torsion

While curvature quantifies how sharply a curve bends in space, torsion measures how much the curve deviates from being planar—it captures the twist of a space curve out of the plane of curvature. Torsion is a fundamental concept in the study of space curves, providing insights into the three-dimensional behavior of trajectories. For a smooth curve defined by the vector-valued function $\mathbf{r}(t)$, the torsion $\tau(t)$ at a point can also be expressed in terms of the tangent, normal, and binormal vectors. Specifically, the torsion relates to the rate of rotation of the binormal vector $\mathbf{B}$ as:

$$\mathbf{B}' = -\tau\mathbf{N},$$

where $\mathbf{B}'$ is the derivative of the binormal vector, and $\mathbf{N}$ is the principal normal vector. This provides an alternative definition for torsion:

$$\tau(t) = -\mathbf{N} \cdot \mathbf{B}'.$$

Alternatively, using the derivatives of $\mathbf{r}(t)$, torsion can be computed as:

$$\tau(t) = \frac{(\mathbf{r}'(t) \times \mathbf{r}''(t)) \cdot \mathbf{r}'''(t)}{|\mathbf{r}'(t) \times \mathbf{r}''(t)|^2}.$$

Similarly, as we have done with curvature, if we consider a curve in the form $\mathbf{r}(t) = \langle t, f(t) \rangle$, we can derive an expression for torsion for any function $f(t)$. For such a curve:

$$\mathbf{r}(t) = \langle t, f(t), 0 \rangle.$$

The derivatives of $\mathbf{r}(t)$ are:

$$\mathbf{r}'(t) = \langle 1, f'(t), 0 \rangle, \quad \mathbf{r}''(t) = \langle 0, f''(t), 0 \rangle, \quad \mathbf{r}'''(t) = \langle 0, f'''(t), 0 \rangle.$$

The cross product $\mathbf{r}'(t) \times \mathbf{r}''(t)$ is:

$$\mathbf{r}'(t) \times \mathbf{r}''(t) = \langle 0, 0, -f''(t) \rangle.$$

Taking the dot product $(\mathbf{r}'(t) \times \mathbf{r}''(t)) \cdot \mathbf{r}'''(t)$:

$$(\mathbf{r}'(t) \times \mathbf{r}''(t)) \cdot \mathbf{r}'''(t) = 0.$$

Finally, the torsion for the curve $\mathbf{r}(t) = \langle t, f(t), 0 \rangle$ simplifies to:

$$\tau(t) = 0.$$

This result shows that a curve defined entirely in a plane (as $\mathbf{r}(t) = \langle t, f(t) \rangle$ implies) has zero torsion. Torsion is zero for **planar curves**, as they do not twist out of their plane. A non-zero torsion, on the other hand, indicates that the curve is three-dimensional, exhibiting a helical or twisting behavior. The torsion $\tau(t)$ measures the rate at which the curve's osculating plane rotates about the tangent vector $\mathbf{T}(t)$. This rotation reflects the deviation of the curve from being planar. The relationship between torsion and the behavior of a space curve can be summarized as follows:

- A **zero torsion** implies that the osculating plane is constant—the curve lies entirely within a fixed plane.
- A **positive torsion** indicates a right-handed twist of the curve.
- A **negative torsion** indicates a left-handed twist of the curve.

The sign and magnitude of the torsion provide critical information about the direction and intensity of the twist. This insight is invaluable in applications such as DNA helices, spiral staircases, and other helical structures. For instance:

- **Torsion in Mechanics:** The stress on a twisted rod or beam is directly related to the applied torque T , the polar moment of inertia J , and the radius r :

$$\tau_s = \frac{Tr}{J}.$$

- **Helical Springs:** In engineering, torsion helps predict the deformation behavior of springs and coils under load.
- **Oscillatory Motion:** The torsional oscillation frequency ω of a system is given by:

$$\omega = \sqrt{\frac{C}{I}},$$

where C is the torsional constant, and I is the moment of inertia.

The interplay between torsion $\tau(t)$ and curvature $\kappa(t)$ characterizes the local behavior of a space curve. Together, they describe the curve's geometry and allow for its reconstruction up to position in space. This is formalized in the *Fundamental Theorem of Space Curves*: A space curve is uniquely determined (up to rigid motions in space) by its curvature $\kappa(t)$ and torsion $\tau(t)$, provided that $\kappa(t) > 0$. The theorem underlines how torsion complements curvature to provide a full geometric understanding of a curve, aiding in both theoretical

studies and practical applications. Below is a parametric curve with its respective curvature and torsion plotted below it:




Figure 2.7: Parametric curve $\mathbf{r}(t) = \langle \cos t, \sin t, \sin 2t \rangle$ with markers at local maxima (blue) and minima (red) of $z(t)$, and dotted lines indicating projections onto the xy -plane and x - and y -axes.




Figure 2.8: Curvature $\kappa(t)$ (solid blue) and torsion $\tau(t)$ (dashed red) for the parametric curve $\mathbf{r}(t) = \langle \cos t, \sin t, \sin 2t \rangle$.

The circular helix in Figure 2.8 illustrates a constant rate of twist, embodying the concept of torsion in a tangible form.

Motion Context

In the context of motion, torsion affects the angular momentum of a particle moving along a space curve. The torsion determines how the plane of motion rotates, influencing gyroscopic effects and the stability of the motion. In designing roller coasters or space trajectories, engineers must account for torsion to ensure comfort and structural integrity.

▲ **Example 2.3.3** A particle moves along the space curve defined by:

$$\mathbf{r}(t) = \langle \cos(t), \sin(t), t \rangle, \quad t \in [0, 2\pi].$$

1. Compute the torsion $\tau(t)$ of the curve at any point t .
2. Evaluate the torsion at $t = \frac{\pi}{2}$.
3. Interpret the significance of the torsion for this curve.

Solution

1. **Compute $\tau(t)$:**

First, compute the derivatives:

First derivative $\mathbf{r}'(t)$:

$$\mathbf{r}'(t) = \langle -\sin(t), \cos(t), 1 \rangle.$$

Second derivative $\mathbf{r}''(t)$:

$$\mathbf{r}''(t) = \langle -\cos(t), -\sin(t), 0 \rangle.$$

Third derivative $\mathbf{r}'''(t)$:

$$\mathbf{r}'''(t) = \langle \sin(t), -\cos(t), 0 \rangle.$$

Compute the cross product $\mathbf{r}'(t) \times \mathbf{r}''(t)$:

$$\begin{aligned} \mathbf{r}'(t) \times \mathbf{r}''(t) &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -\sin(t) & \cos(t) & 1 \\ -\cos(t) & -\sin(t) & 0 \end{vmatrix} \\ &= \langle \sin(t), -\cos(t), 1 \rangle. \end{aligned}$$

Compute the magnitude:

$$|\mathbf{r}'(t) \times \mathbf{r}''(t)| = \sqrt{\sin^2(t) + \cos^2(t) + 1^2} = \sqrt{1 + 1} = \sqrt{2}.$$

Compute the dot product $(\mathbf{r}'(t) \times \mathbf{r}''(t)) \cdot \mathbf{r}'''(t)$:

$$\begin{aligned} (\mathbf{r}'(t) \times \mathbf{r}''(t)) \cdot \mathbf{r}'''(t) &= \langle \sin(t), -\cos(t), 1 \rangle \cdot \langle \sin(t), -\cos(t), 0 \rangle \\ &= \sin^2(t) + (-\cos(t))^2 + 1 \cdot 0 \\ &= \sin^2(t) + \cos^2(t) + 0 = 1. \end{aligned}$$

Therefore, the torsion is:

$$\tau(t) = \frac{1}{(\sqrt{2})^2} = \frac{1}{2}.$$

2. **Evaluate** $\tau\left(\frac{\pi}{2}\right)$:

Since $\tau(t)$ is constant and equals $\frac{1}{2}$, we have:

$$\tau\left(\frac{\pi}{2}\right) = \frac{1}{2}.$$

3. **Interpretation:**

The curve $\mathbf{r}(t) = \langle \cos(t), \sin(t), t \rangle$ describes a circular helix with radius 1 and constant upward pitch. The constant torsion $\tau = \frac{1}{2}$ indicates that the curve twists uniformly in space. The positive value signifies a right-handed helix. The constant curvature and torsion reflect the uniform bending and twisting of the helix, characterizing its geometric properties completely.

Lecture 2.4: Arc Length and Reparameterization

The concept of arc length is fundamental in calculus and geometry, providing a measure of the distance along a curve. Understanding how to compute arc length is essential for analyzing the properties of curves in various coordinate systems, including Cartesian, parametric, and polar coordinates. Additionally, reparameterizing curves by arc length simplifies many problems in differential geometry and physics by providing a natural parameterization that relates directly to the geometry of the curve.

Computing Arc Length

To derive the formula for arc length, consider a smooth curve C in the plane defined by the function $y = f(x)$, where x ranges from a to b . We can approximate the curve by partitioning the interval $[a, b]$ into n subintervals and connecting consecutive points with straight line segments. The length Δs_i of each segment is approximately:

$$\Delta s_i \approx \sqrt{\Delta x_i^2 + \Delta y_i^2} = \sqrt{(\Delta x_i)^2 + [f'(x_i^*)\Delta x_i]^2} = \Delta x_i \sqrt{1 + [f'(x_i^*)]^2},$$

where x_i^* is some point in the i -th subinterval. Note, that we replace the finite difference $(\Delta y_i)^2$ with the derivative $f'(x_i^*)$ multiplied by the finite difference Δx_i , following the limit definition of the derivative. This substitution simplifies the expression and emphasizes the relationship between the derivative and the arc length formula.




Figure 2.9: Approximating the arc length of a curve using a right triangle with Δs as the hypotenuse, Δx as the base, and Δy as the height.

Next, summing over all segments, the total approximate arc length S is:

$$S \approx \sum_{i=1}^n \Delta s_i = \sum_{i=1}^n \Delta x_i \sqrt{1 + [f'(x_i^*)]^2}.$$

Taking the limit as $n \rightarrow \infty$ and $\Delta x_i \rightarrow 0$, the sum becomes an integral:

$$S = \int_a^b \sqrt{1 + [f'(x)]^2} dx.$$

The integrand $\sqrt{1 + [f'(x)]^2}$ represents the differential arc length element ds , so we can express the arc length as:

$$S = \int_C ds = \int_a^b \sqrt{1 + [f'(x)]^2} dx.$$

For a closed curve C , we may also introduce the notation to be later used:

$$S = \oint_C ds = \int_a^b \sqrt{1 + [f'(x)]^2} dx.$$

Similarly, for a curve defined parametrically by $x = x(t)$ and $y = y(t)$, with t ranging over $[\alpha, \beta]$, we can derive the arc length formula starting from the Cartesian form $ds = \sqrt{1 + [f'(x)]^2} dx$. Using the relationships $dx = x'(t) dt$ and $dy = y'(t) dt$, and noting that $f'(x) = \frac{dy}{dx} = \frac{y'(t)}{x'(t)}$, substituting and simplifying gives:

$$ds = \sqrt{[x'(t)]^2 + [y'(t)]^2} dt.$$

Thus, the arc length S is:

$$S = \int_C ds = \int_\alpha^\beta \sqrt{[x'(t)]^2 + [y'(t)]^2} dt.$$


Extending this to three-dimensional space, where the curve is given by $\mathbf{r}(t) = \langle x(t), y(t), z(t) \rangle$, the differential arc length element becomes:

$$ds = \sqrt{[x'(t)]^2 + [y'(t)]^2 + [z'(t)]^2} dt = |\mathbf{r}'(t)| dt.$$

Therefore, the arc length S is:

$$S = \int_C ds = \int_\alpha^\beta |\mathbf{r}'(t)| dt.$$

A typical application of arc length is sector length that you may have studied in your Pre-Calculus class:



Circle with radius r

Figure 2.10: A circular sector with radius r , central angle θ , and arc length $L = r\theta$.

Arc Length in Polar Coordinates

In polar coordinates, where a curve is defined by $r = r(\theta)$, the differential arc length element is derived using the relationship between Cartesian and polar coordinates:

$$ds = \sqrt{\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2} d\theta.$$

Since $x = r(\theta) \cos \theta$ and $y = r(\theta) \sin \theta$, we have:

$$\frac{dx}{d\theta} = r'(\theta) \cos \theta - r(\theta) \sin \theta, \quad \frac{dy}{d\theta} = r'(\theta) \sin \theta + r(\theta) \cos \theta.$$

Computing $\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2$ simplifies to:

$$\left(\frac{dx}{d\theta}\right)^2 + \left(\frac{dy}{d\theta}\right)^2 = [r'(\theta)]^2 + [r(\theta)]^2.$$

Thus, the arc length S in polar coordinates is:

$$S = \int_C ds = \int_{\theta_1}^{\theta_2} \sqrt{[r'(\theta)]^2 + [r(\theta)]^2} d\theta.$$

These formulas provide the tools necessary to compute the length of curves in various coordinate systems, facilitating the analysis of geometric properties and physical applications. In previous lectures, we explored vector-valued functions and their derivatives, which are fundamental in understanding the behavior of curves and motion in space.

▲ Example 2.4.1 Compute the arc length of a circular sector of radius $r = 3$ and central angle $\theta = \frac{\pi}{4}$ using both rectangular and polar coordinates. Verify that the results are consistent and generalize to the formula $L = r\theta$.

Solution

1. Using Rectangular Coordinates:

The equation of the circle is $x^2 + y^2 = r^2$, which can be solved for y :

$$y = \sqrt{r^2 - x^2}.$$

For the sector from $\theta = 0$ to $\theta = \frac{\pi}{4}$, the corresponding x -values are:

$$x = r \cos \theta.$$

Explanation: As θ ranges from 0 to $\frac{\pi}{4}$, x ranges from $x = r \cos(0) = r$ to $x = r \cos\left(\frac{\pi}{4}\right) = \frac{r}{\sqrt{2}}$.

Therefore, x ranges from $x_1 = r$ to $x_2 = \frac{r}{\sqrt{2}}$.

The arc length formula is:

$$L = \int_{x_1}^{x_2} \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx.$$

Differentiating y with respect to x :

$$\frac{dy}{dx} = \frac{-x}{\sqrt{r^2 - x^2}}.$$

Computing the integrand:

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \sqrt{1 + \frac{x^2}{r^2 - x^2}} = \frac{r}{\sqrt{r^2 - x^2}}.$$

Thus, the arc length becomes:

$$L = \int_r^{\frac{r}{\sqrt{2}}} \frac{r}{\sqrt{r^2 - x^2}} dx.$$

Let $x = r \cos t$, which implies $dx = -r \sin t dt$ and $\sqrt{r^2 - x^2} = r \sin t$. Changing the bounds:

$$\text{When } x = r, \quad \cos t = \frac{x}{r} = 1 \implies t = 0.$$

$$\text{When } x = \frac{r}{\sqrt{2}}, \quad \cos t = \frac{1}{\sqrt{2}} \implies t = \frac{\pi}{4}.$$

Substituting into the integral:

$$L = \int_{t=0}^{t=\frac{\pi}{4}} \frac{r}{r \sin t} \cdot (-r \sin t) dt = -r \int_0^{\frac{\pi}{4}} dt = -r \left(\frac{\pi}{4} - 0 \right).$$

The negative sign arises because $dx = -r \sin t dt$, and we reversed the limits when changing variables.

Therefore:

$$L = -r \left(\frac{\pi}{4} \right) = -\frac{r\pi}{4}.$$

Taking the absolute value (since length is positive):

$$L = \frac{r\pi}{4}.$$

Substituting $r = 3$:

$$L = \frac{3\pi}{4}.$$

2. Using Polar Coordinates:

In polar coordinates, the arc length formula is:

$$L = \int_{\theta_1}^{\theta_2} \sqrt{\left(\frac{dr}{d\theta}\right)^2 + r^2} d\theta.$$

For a circle of radius r , r is constant, so $\frac{dr}{d\theta} = 0$. Thus:

$$L = \int_0^{\frac{\pi}{4}} \sqrt{0^2 + r^2} d\theta = \int_0^{\frac{\pi}{4}} r d\theta = r \left(\frac{\pi}{4} - 0\right) = \frac{r\pi}{4}.$$

Substituting $r = 3$:

$$L = \frac{3\pi}{4}.$$

3. Generalization Using the Arc Length Formula:

The general formula for the arc length of a circular sector in polar coordinates is:

$$L = \int_{\theta_1}^{\theta_2} r d\theta = r(\theta_2 - \theta_1) = r\theta.$$

This confirms the classic formula:

$$L = r\theta.$$

Therefore, the results from both rectangular and polar coordinates are consistent with the general formula for the arc length of a circular sector.

▲ Example 2.4.2 Compute the arc length of the parametric curve defined by $\mathbf{r}(t) = \langle t^2, t^3 \rangle$ for t ranging from 0 to 1.

Solution The arc length S of a parametric curve $\mathbf{r}(t) = \langle x(t), y(t) \rangle$ from $t = a$ to $t = b$ is given by:

$$S = \int_a^b \sqrt{[x'(t)]^2 + [y'(t)]^2} dt.$$

Compute the derivatives:

$$x'(t) = \frac{d}{dt}[t^2] = 2t, \quad y'(t) = \frac{d}{dt}[t^3] = 3t^2.$$

Compute the integrand:

$$\sqrt{[x'(t)]^2 + [y'(t)]^2} = \sqrt{(2t)^2 + (3t^2)^2} = \sqrt{4t^2 + 9t^4} = t\sqrt{4 + 9t^2}.$$

Thus, the arc length is:

$$S = \int_0^1 t\sqrt{4+9t^2} dt.$$

Let's make a substitution to simplify the integral. Let $u = 4 + 9t^2$, then $du = 18t dt$. Solving for $t dt$:

$$t dt = \frac{du}{18}.$$

Change the limits of integration:

When $t = 0$:

$$u = 4 + 9(0)^2 = 4.$$

When $t = 1$:

$$u = 4 + 9(1)^2 = 13.$$

Substitute into the integral:

$$S = \int_{u=4}^{u=13} \frac{1}{18} \sqrt{u} du = \frac{1}{18} \int_4^{13} u^{1/2} du.$$

Integrate:

$$\frac{1}{18} \left[\frac{u^{3/2}}{\frac{3}{2}} \right]_4^{13} = \frac{1}{18} \left[\frac{2}{3} u^{3/2} \right]_4^{13} = \frac{1}{27} [u^{3/2}]_4^{13}.$$

Compute $u^{3/2}$ at the limits:

At $u = 13$:

$$13^{3/2} = \sqrt{13^3} = \sqrt{2197} = 13\sqrt{13}.$$

At $u = 4$:

$$4^{3/2} = \sqrt{4^3} = \sqrt{64} = 8.$$

Therefore,

$$S = \frac{1}{27} (13\sqrt{13} - 8).$$

Arc Length in Higher Dimensions

The concept of arc length extends beyond three dimensions into n -dimensional space. For a curve $\mathbf{r}(t) = \langle x_1(t), x_2(t), \dots, x_n(t) \rangle$, the differential arc length element is:

$$ds = \sqrt{[x_1'(t)]^2 + [x_2'(t)]^2 + \dots + [x_n'(t)]^2} dt = |\mathbf{r}'(t)| dt.$$

Therefore, the arc length S is:

$$S = \int_C ds = \int_\alpha^\beta |\mathbf{r}'(t)| dt.$$

This generalization is crucial in fields like physics and engineering, where motion and paths in higher-dimensional spaces are considered. The notation $\int_C f \, ds$ represents the integral of a function f along a curve C with respect to arc length, which is an important concept in line integrals and further studies in vector calculus.

Arc Length Parameterization and Reparameterization

Parameterizing a curve by its arc length s provides a natural and intrinsic description of the curve, independent of the specific parameterization used initially. This means that the parameter s directly measures the distance along the curve from a fixed starting point, offering a uniform way to describe points on the curve based solely on their position along it. The arc length from a fixed point t_0 is defined as:

$$s(t) = \int_{t_0}^t |\mathbf{r}'(u)| \, du,$$

where $|\mathbf{r}'(u)|$ represents the speed of traversal along the curve at each point u . This integral accumulates the infinitesimal distances $ds = |\mathbf{r}'(u)| \, du$ along the curve from t_0 to t .

By reparameterizing the curve using s , we ensure that:

$$\left| \frac{d\mathbf{r}}{ds} \right| = 1,$$

meaning the magnitude of the derivative of $\mathbf{r}$ with respect to s is unity. In other words, the speed along the curve is constant with respect to s . This property simplifies calculations involving the curve's geometry, as derivatives with respect to s directly reflect changes per unit length along the curve.

Reparameterizing a curve by arc length involves finding a new parameter s such that $\mathbf{r}$ is expressed as a function of s , and the condition $\left| \frac{d\mathbf{r}}{ds} \right| = 1$ holds. The steps for reparameterization are:

1. Compute the arc length function $s(t)$ from a fixed point t_0 :

$$s(t) = \int_{t_0}^t |\mathbf{r}'(u)| \, du.$$

2. Invert the function $s(t)$ to find $t(s)$: Solve for t in terms of s to get $t = t(s)$.
3. Substitute $t(s)$ back into $\mathbf{r}(t)$ to obtain $\mathbf{r}(s)$:

$$\mathbf{r}(s) = \mathbf{r}(t(s)).$$

When a curve is parameterized by an arbitrary parameter t , the speed $|\mathbf{r}'(t)|$ can vary, making it more complicated to analyze properties that depend on the distance along the curve. By switching to arc length parameterization, we eliminate the influence of the parameter's rate

of change, focusing solely on the geometric properties of the curve. This reparameterization is particularly useful when dealing with curves where the speed is not constant, as it allows for a uniform treatment of the curve's properties and simplifies calculations in differential geometry and physics.

▲ **Example 2.4.3** Consider the curve defined by $\mathbf{r}(t) = \langle 2 \cos t, 2 \sin t, 3t \rangle$, where $t \in [0, 2\pi]$.

1. Compute the arc length $s(t)$ as a function of t , starting from $t_0 = 0$.
2. Reparameterize the curve by its arc length s .
3. Verify that $\left| \frac{d\mathbf{r}}{ds} \right| = 1$ for the reparameterized curve.

Solution

1. **Compute $s(t)$:**

First, compute the derivative $\mathbf{r}'(t)$:

$$\mathbf{r}'(t) = \langle -2 \sin t, 2 \cos t, 3 \rangle.$$

Compute the magnitude of $\mathbf{r}'(t)$:

$$|\mathbf{r}'(t)| = \sqrt{(-2 \sin t)^2 + (2 \cos t)^2 + 3^2} = \sqrt{4 \sin^2 t + 4 \cos^2 t + 9}.$$

Simplify using $\sin^2 t + \cos^2 t = 1$:

$$|\mathbf{r}'(t)| = \sqrt{4(1) + 9} = \sqrt{13}.$$

The speed $|\mathbf{r}'(t)|$ is constant. The arc length from $t_0 = 0$ to t is:

$$s(t) = \int_0^t |\mathbf{r}'(u)| \, du = \int_0^t \sqrt{13} \, du = \sqrt{13} t.$$

2. **Reparameterize by arc length s :**

Solve for t in terms of s :

$$s = \sqrt{13} t \implies t = \frac{s}{\sqrt{13}}.$$

Substitute $t = \frac{s}{\sqrt{13}}$ into $\mathbf{r}(t)$:

$$\mathbf{r}(s) = \left\langle 2 \cos \left(\frac{s}{\sqrt{13}} \right), 2 \sin \left(\frac{s}{\sqrt{13}} \right), \frac{3s}{\sqrt{13}} \right\rangle.$$

3. **Verify** $\left| \frac{d\mathbf{r}}{ds} \right| = 1$:

Compute the derivative $\frac{d\mathbf{r}}{ds}$ using the chain rule:

$$\frac{d\mathbf{r}}{ds} = \left\langle -2 \sin \left(\frac{s}{\sqrt{13}} \right) \cdot \frac{1}{\sqrt{13}}, 2 \cos \left(\frac{s}{\sqrt{13}} \right) \cdot \frac{1}{\sqrt{13}}, \frac{3}{\sqrt{13}} \right\rangle.$$

Simplify:

$$\frac{d\mathbf{r}}{ds} = \frac{1}{\sqrt{13}} \left\langle -2 \sin \left(\frac{s}{\sqrt{13}} \right), 2 \cos \left(\frac{s}{\sqrt{13}} \right), 3 \right\rangle.$$

Compute the magnitude $\left| \frac{d\mathbf{r}}{ds} \right|$:

$$\left| \frac{d\mathbf{r}}{ds} \right| = \frac{1}{\sqrt{13}} \left| \left\langle -2 \sin \left(\frac{s}{\sqrt{13}} \right), 2 \cos \left(\frac{s}{\sqrt{13}} \right), 3 \right\rangle \right|.$$

Calculate the square of the magnitude:

$$\begin{aligned} \left| \frac{d\mathbf{r}}{ds} \right|^2 &= \left(\frac{1}{\sqrt{13}} \right)^2 \left([-2 \sin \left(\frac{s}{\sqrt{13}} \right)]^2 + [2 \cos \left(\frac{s}{\sqrt{13}} \right)]^2 + 3^2 \right) \\ &= \frac{1}{13} \left(4 \sin^2 \left(\frac{s}{\sqrt{13}} \right) + 4 \cos^2 \left(\frac{s}{\sqrt{13}} \right) + 9 \right). \end{aligned}$$

Simplify using $\sin^2 \theta + \cos^2 \theta = 1$:

$$4 \sin^2 \left(\frac{s}{\sqrt{13}} \right) + 4 \cos^2 \left(\frac{s}{\sqrt{13}} \right) = 4(\sin^2 \theta + \cos^2 \theta) = 4(1) = 4.$$

Therefore,

$$\left| \frac{d\mathbf{r}}{ds} \right|^2 = \frac{1}{13}(4 + 9) = \frac{13}{13} = 1.$$

Thus,

$$\left| \frac{d\mathbf{r}}{ds} \right| = 1,$$

confirming that the curve is correctly parameterized by arc length.

Numerical Approximation of Arc Length

When the arc length integral cannot be evaluated analytically due to a complex integrand or non-elementary functions, numerical methods and approximations provide practical solutions. These techniques are essential for estimating arc lengths in engineering and physics applications where precise measurements are necessary but exact calculations are infeasible.

Approximation Techniques

One common approach is to approximate the integrand by neglecting higher-order terms when they contribute insignificantly to the value of the integral. This method is particularly effective when dealing with small quantities, where squared or higher powers of a small parameter can be considered negligible.

For example, if ϵ is a small parameter (i.e., $\epsilon \ll 1$), then ϵ^2 is much smaller than ϵ and can often be neglected in the approximation. This simplification reduces the complexity of the integrand, making the integral more tractable. Consider a curve where the integrand involves a square root of a sum of terms, such as:

$$ds = \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx.$$

If $\left|\frac{dy}{dx}\right|$ is small over the interval of interest, we can use the approximation:

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} \approx 1 + \frac{1}{2} \left(\frac{dy}{dx}\right)^2,$$

since $\left(\frac{dy}{dx}\right)^2$ is much smaller than 1, and higher-order terms can be neglected.

Common Numerical Methods

Beyond approximation techniques, several numerical integration methods are widely used:

- **Trapezoidal Rule:** Divides the interval into small segments, approximating the area under the curve as a series of trapezoids. The arc length is estimated by summing the lengths of these trapezoidal segments.
- **Simpson's Rule:** Uses quadratic polynomials (parabolic arcs) to approximate the integrand over each subinterval, providing higher accuracy than the trapezoidal rule for smooth functions.
- **Gaussian Quadrature:** Employs weighted sums of function values at specified points within the integration interval, optimizing the approximation for polynomials up to a certain degree.

- **Adaptive Quadrature:** Adjusts the size of the subintervals based on the behavior of the integrand, refining the approximation where the function changes rapidly.

These numerical methods are crucial in practical scenarios where exact solutions are unattainable, enabling engineers and scientists to estimate arc lengths with desired accuracy.

▲ **Example 2.4.4** Consider the curve defined by $y = \epsilon x$, where ϵ is a small constant ($\epsilon \ll 1$) and $x \in [0, L]$.

1. Approximate the arc length S of the curve from $x = 0$ to $x = L$ using the small angle approximation.
2. Compare the approximate arc length to the exact value obtained without approximation.

Solution

1. Approximate Arc Length Using Small Angle Approximation:

Compute the derivative:

$$\frac{dy}{dx} = \epsilon.$$

Since $\epsilon \ll 1$, $\left(\frac{dy}{dx}\right)^2 = \epsilon^2$ is even smaller.

Using the approximation:

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} \approx 1 + \frac{1}{2} \left(\frac{dy}{dx}\right)^2 = 1 + \frac{\epsilon^2}{2}.$$

The approximate arc length is:

$$S_{\text{approx}} = \int_0^L \left(1 + \frac{\epsilon^2}{2}\right) dx = \left[x + \frac{\epsilon^2}{2}x\right]_0^L = L + \frac{\epsilon^2 L}{2}.$$

2. Exact Arc Length:

Compute the exact integrand:

$$\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = \sqrt{1 + \epsilon^2}.$$

The exact arc length is:

$$S_{\text{exact}} = \int_0^L \sqrt{1 + \epsilon^2} \, dx = \sqrt{1 + \epsilon^2} \cdot L.$$

Comparison:

Expand $\sqrt{1 + \epsilon^2}$ using a binomial series for small ϵ :

$$\sqrt{1 + \epsilon^2} = 1 + \frac{\epsilon^2}{2} - \frac{\epsilon^4}{8} + \mathcal{O}(\epsilon^6).$$

Therefore:

$$S_{\text{exact}} = \left(1 + \frac{\epsilon^2}{2} - \frac{\epsilon^4}{8}\right) L.$$

The approximate arc length is:

$$S_{\text{approx}} = \left(1 + \frac{\epsilon^2}{2}\right) L.$$

Difference Between Exact and Approximate Values:

The difference is:

$$S_{\text{exact}} - S_{\text{approx}} = \left(1 + \frac{\epsilon^2}{2} - \frac{\epsilon^4}{8}\right) L - \left(1 + \frac{\epsilon^2}{2}\right) L = -\frac{\epsilon^4 L}{8}.$$

Since ϵ^4 is negligible compared to ϵ^2 , the error introduced by the approximation is minimal for small ϵ .

Formulas and Definitions

Important Vocabulary

- **Vector-Valued Functions:** Functions where each component is a scalar function of a parameter, typically used to describe motion or curves in space.
- **Parametric Equations:** Equations that express the coordinates of a point as functions of a parameter, providing a way to represent curves.
- **Arc Length:** The length of a curve, calculated using the integral of the magnitude of the derivative of the position vector.
- **Curvature (κ):** A measure of how sharply a curve bends at a given point, derived from the rate of change of the tangent vector.
- **TNB Frame:** The Tangent (**T**), Normal (**N**), and Binormal (**B**) vectors forming an orthonormal basis for describing motion along a curve.
- **Torsion (τ):** A measure of the rate at which a curve twists out of its osculating plane.
- **Osculating Circle:** The circle that best approximates the curve at a point, characterized by the same curvature.

Key Formulas

Parametric Equations for a Curve:

$$\mathbf{r}(t) = \langle f(t), g(t), h(t) \rangle$$

$$= \int_a^b \sqrt{[f'(t)]^2 + [g'(t)]^2 + [h'(t)]^2} dt$$

Derivative of a Vector-Valued Function: **Curvature:**

$$\mathbf{r}'(t) = \langle f'(t), g'(t), h'(t) \rangle$$

$$\kappa(t) = \frac{|\mathbf{r}'(t) \times \mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3} = \frac{|\mathbf{T}'(s)|}{|\mathbf{r}'(s)|}$$

Integral of a Vector-Valued Function:

$$\int \mathbf{r}(t) dt = \left\langle \int f(t) dt, \int g(t) dt, \int h(t) dt \right\rangle + \mathbf{C}$$

$$\kappa(x) = \frac{|f''(x)|}{(1 + [f'(x)]^2)^{3/2}}.$$

TNB Frame Vectors:

$$\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|},$$

$$\mathbf{N}(t) = \frac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|},$$

$$\mathbf{B}(t) = \mathbf{T}(t) \times \mathbf{N}(t)$$

Arc Length:

$$S = \int_a^b |\mathbf{r}'(t)| dt$$

Torsion:

$$\tau(t) = \frac{(\mathbf{r}'(t) \times \mathbf{r}''(t)) \cdot \mathbf{r}'''(t)}{|\mathbf{r}'(t) \times \mathbf{r}''(t)|^2}$$

Fundamental Theorem of Calculus for Vector-Valued Functions:

$$\int_a^b \mathbf{r}'(t) dt = \mathbf{r}(b) - \mathbf{r}(a)$$

Osculating Circle Radius:

$$R = \frac{1}{\kappa(t)}$$

Arc Length in Polar Coordinates:

$$S = \int_{\theta_1}^{\theta_2} \sqrt{[r'(\theta)]^2 + [r(\theta)]^2} d\theta$$

Projectile Motion:

$$\mathbf{r}(t) = \langle v_0 \cos \theta t, 0, v_0 \sin \theta t - \tfrac{1}{2}gt^2 \rangle,$$

$$z_{\max} = \frac{v_0^2 \sin^2 \theta}{2g},$$

$$R = \frac{v_0^2 \sin 2\theta}{g},$$

$$t_{\text{flight}} = \frac{2v_0 \sin \theta}{g},$$

$$K = \frac{1}{2}mv_0^2,$$

$$U = mgz,$$

$$E_{\text{total}} = K + U$$

Practice Problems

- Determine whether the curve $\mathbf{r}(t) = \langle \cos t, \sin t, t \rangle$ is smooth for all t .
- Compute the derivative $\mathbf{r}'(t)$ of the vector-valued function $\mathbf{r}(t) = \langle t^2, e^{2t}, \ln t \rangle$.
- Find the vector equation of the line passing through the points $\mathbf{r}_1 = \langle 1, 2, 3 \rangle$ and $\mathbf{r}_2 = \langle 4, 5, 6 \rangle$.
- Write the parametric equations for the curve $\mathbf{r}(t) = \langle 3 \cos t, 3 \sin t, t \rangle$ and describe the geometric shape of the curve.
- Show that the curve $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$ lies on the surface $z = y^{3/2}$.
- Find the velocity vector of the particle whose position is $\mathbf{r}(t) = \langle t, \sin t, \cos t \rangle$ and determine whether the speed is constant.
- Compute the integral $\int_0^1 \mathbf{r}(t) dt$ for $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$.
- Determine the length of the curve $\mathbf{r}(t) = \langle t, t^2, 0 \rangle$ from $t = 0$ to $t = 2$.
- Compute $\mathbf{r}'(t)$ and $\mathbf{r}''(t)$ for $\mathbf{r}(t) = \langle t^3, t^2, t \rangle$.
- Evaluate the integral $\int_0^\pi \mathbf{r}(t) \times \mathbf{r}'(t) dt$ where $\mathbf{r}(t) = \langle \cos t, \sin t, t \rangle$.
- Given $\mathbf{r}(t) = \langle e^t \cos t, e^t \sin t, t^2 \rangle$, compute its derivative and interpret its geometric meaning.
- Find the second derivative of $\mathbf{r}(t) = \langle \ln t, t, \sqrt{t} \rangle$.
- A particle moves along the curve $\mathbf{r}(t) = \langle e^t, \ln t, t^3 \rangle$. Compute its velocity and acceleration vectors at $t = 1$.
- Write the vector equation of the tangent line to $\mathbf{r}(t) = \langle t^2, t^3, t \rangle$ at $t = 1$.
- Determine the acceleration vector $\mathbf{a}(t)$ for the curve $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$ and discuss its significance.
- Find the velocity and acceleration vectors for the position vector $\mathbf{r}(t) = \langle t, \cos t, e^t \rangle$.
- Evaluate $\int \mathbf{r}(t) \cdot \mathbf{r}'(t) dt$ where $\mathbf{r}(t) = \langle t^2, \sin t, \cos t \rangle$.
- Find the velocity and acceleration vectors of the projectile $\mathbf{r}(t) = \langle 10t, 20t - 4.9t^2 \rangle$ at $t = 1$.
- A particle moves along the curve $\mathbf{r}(t) = \langle \cos t, \sin t, t \rangle$. Determine its speed and acceleration at $t = \frac{\pi}{4}$.
- An object is projected with initial velocity $\mathbf{v}_0 = \langle 20, 30 \rangle$ m/s. Write its position vector as a function of time, assuming acceleration due to gravity is $\mathbf{a} = \langle 0, -9.8 \rangle$.
- Determine the maximum height of the projectile $\mathbf{r}(t) = \langle v_0 \cos \theta t, v_0 \sin \theta t - \frac{1}{2}gt^2 \rangle$.
- A particle moves with velocity $\mathbf{v}(t) = \langle t^2, \sin t \rangle$. If the particle starts at $\langle 1, 0 \rangle$, find its position vector $\mathbf{r}(t)$.
- Compute the velocity and acceleration vectors for a particle moving along $\mathbf{r}(t) = \langle t^3, t^2, t \rangle$ at $t = 2$.
- A projectile is launched with speed v_0 at an angle θ above the horizontal. Derive the expressions for its horizontal and vertical components of velocity as functions of time.

25. Given the position vector $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$, find the speed of the particle at any time t .
26. A car travels along a straight path with acceleration $\mathbf{a}(t) = \langle 0, 2 \rangle$ m/s² and initial velocity $\mathbf{v}(0) = \langle 5, 0 \rangle$ m/s. Find its position vector $\mathbf{r}(t)$.
27. Determine the time at which the projectile $\mathbf{r}(t) = \langle 10t, 20t - 4.9t^2 \rangle$ reaches its maximum height.
28. Compute the unit tangent vector $\mathbf{T}(t)$ for $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$ at $t = 1$.
29. Find the curvature $\kappa(t)$ of the curve $\mathbf{r}(t) = \langle t, t^2, 0 \rangle$.
30. Compute the radius of curvature at $t = 0$ for $\mathbf{r}(t) = \langle t, t^3, t^2 \rangle$.
31. Determine the curvature of the helix $\mathbf{r}(t) = \langle \cos t, \sin t, t \rangle$ and show it is constant.
32. Find the normal vector $\mathbf{N}(t)$ and the binormal vector $\mathbf{B}(t)$ for $\mathbf{r}(t) = \langle t, \ln t, t^2 \rangle$ at $t = 1$.
33. For the curve $\mathbf{r}(t) = \langle t, \sin t, \cos t \rangle$, compute the curvature and determine where the curve is most curved.
34. A particle moves along the curve $\mathbf{r}(t) = \langle e^t, e^{-t}, t \rangle$. Find the curvature $\kappa(t)$ at $t = 0$.
35. Calculate the curvature of the curve $\mathbf{r}(t) = \langle t, \ln t, \sqrt{t} \rangle$.
36. Find the curvature and the principal normal vector for the curve $\mathbf{r}(t) = \langle t^2, t^3, t \rangle$ at $t = 1$.
37. Determine the curvature of the curve $\mathbf{r}(t) = \langle \ln(t+1), \sqrt{t}, t^2 \rangle$ at $t = 1$.
38. Compute the tangent vector $\mathbf{T}(t)$, the normal vector $\mathbf{N}(t)$, and the binormal vector $\mathbf{B}(t)$ for the curve $\mathbf{r}(t) = \langle t, \ln t, t^2 \rangle$ at $t = 1$.
39. For the curve $\mathbf{r}(t) = \langle \cos t, \sin t, t \rangle$, compute the tangent, normal, and binormal vectors at any t .
40. A particle moves along a helix $\mathbf{r}(t) = \langle \cos t, \sin t, t \rangle$. Compute its curvature and torsion.
41. Given the curve $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$, find the binormal vector $\mathbf{B}(t)$ at $t = 1$.
42. Compute the Frenet-Serret formulas for the curve $\mathbf{r}(t) = \langle t, \sin t, \cos t \rangle$.
43. A curve is defined by $\mathbf{r}(t) = \langle t, e^t, e^{-t} \rangle$. Find the Frenet-Serret frame vectors at $t = 0$.
44. Determine the torsion $\tau(t)$ of the curve $\mathbf{r}(t) = \langle t, \ln t, \sqrt{t} \rangle$.
45. For the space curve $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$, compute the derivative of the binormal vector $\mathbf{B}(t)$.
46. Find the binormal vector $\mathbf{B}(t)$ for the curve $\mathbf{r}(t) = \langle \cos t, \sin t, t \rangle$ at $t = \pi$.
47. A curve has constant curvature κ and constant torsion τ . Describe the geometric nature of the curve.
48. Compute the arc length of the curve $\mathbf{r}(t) = \langle t, t^2, 0 \rangle$ from $t = 0$ to $t = 3$.
49. Reparameterize the curve $\mathbf{r}(t) = \langle 3t, 4t, 0 \rangle$ by its arc length s .
50. Find the arc length of the polar curve $r(\theta) = 3\theta$ for $\theta \in [0, \pi]$.
51. A particle moves along the curve $\mathbf{r}(t) = \langle e^t, e^{-t}, t \rangle$. Compute the arc length of its path from $t = 0$ to $t = 2$.

52. Approximate the arc length of $y = \epsilon x^2$, where $\epsilon \ll 1$, from $x = 0$ to $x = 1$.
53. Design a smooth path for a robotic arm moving from $\langle 0, 0, 0 \rangle$ to $\langle 1, 1, 1 \rangle$ using a vector-valued function with continuous derivatives.
54. Compute the arc length of the curve $\mathbf{r}(t) = \langle t^3, t^2, t \rangle$ from $t = 0$ to $t = 2$.
55. Reparameterize the curve $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$ by arc length s .
56. Find the arc length parameterization of $\mathbf{r}(t) = \langle \cos t, \sin t, t \rangle$.
57. Given $\mathbf{r}(t) = \langle t^2, t^3, t \rangle$, compute the arc length from $t = 0$ to $t = 1$.
58. A robotic arm needs to follow a smooth path described by $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$. Determine the arc length parameterization for $t \in [0, 1]$.
59. An engineer designs a roller coaster track modeled by $\mathbf{r}(t) = \langle t, \sin t, \cos t \rangle$. Calculate the arc length from $t = 0$ to $t = 2\pi$.
60. A drone flies along the path $\mathbf{r}(t) = \langle e^t, \ln t, t \rangle$. Compute the total distance traveled from $t = 1$ to $t = 3$.
61. Optimize the path of a vehicle moving along the curve $\mathbf{r}(t) = \langle t, t^2, t^4 \rangle$ to minimize the travel time given a constant speed.
62. A satellite orbits Earth following the curve $\mathbf{r}(t) = \langle 7000 \cos t, 7000 \sin t, 0 \rangle$ km. Compute the arc length of one complete orbit.
63. Reparameterize the curve $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$ using the arc length from $t = 0$ to $t = 1$.
64. Calculate the arc length of the space curve $\mathbf{r}(t) = \langle t, e^t, \ln t \rangle$ from $t = 1$ to $t = 2$.
65. Determine the arc length of the curve $\mathbf{r}(t) = \langle t, \sin t, \cos t \rangle$ from $t = 0$ to $t = \pi$.
66. A roller coaster track is described by $\mathbf{r}(t) = \langle 10 \cos t, 10 \sin t, t^2 \rangle$. Find the arc length from $t = 0$ to $t = \pi$.
67. Design an engineering path for a drone to move smoothly from $\langle 0, 0, 0 \rangle$ to $\langle 5, 5, 5 \rangle$ with uniform speed. Provide the arc length parameterization.
68. A projectile is launched at an angle of 45 degrees with an initial velocity of 50 m/s. Find its horizontal range and maximum height.
69. A car travels along a circular path of radius 100 m. If its speed increases at a constant rate of 2 m/s², compute its total acceleration at the point where its speed is 10 m/s.
70. A roller coaster travels along the curve $\mathbf{r}(t) = \langle 10 \cos t, 10 \sin t, t^2 \rangle$. Find its speed and acceleration at $t = 1$.
71. A particle follows the curve $\mathbf{r}(t) = \langle \ln(t + 1), \sqrt{t}, t^2 \rangle$. Find its arc length from $t = 0$ to $t = 2$.
72. A drone follows a path described by $\mathbf{r}(t) = \langle t, t^2, \ln t \rangle$. Determine the curvature of the path at $t = 1$.
73. A satellite orbits the Earth along the curve $\mathbf{r}(t) = \langle 7000 \cos t, 7000 \sin t, 0 \rangle$ km. Determine its speed and centripetal acceleration.
74. A robotic arm needs to move smoothly from $\langle 0, 0, 0 \rangle$ to $\langle 2, 2, 2 \rangle$. Design a vector-valued function describing its

- path with continuous first and second derivatives.
75. A particle moves along the path $\mathbf{r}(t) = \langle t, \sin t, \cos t \rangle$. Compute the total distance traveled from $t = 0$ to $t = 2\pi$.
 76. An object moves along the curve $\mathbf{r}(t) = \langle t, t^3, t \rangle$. Find its velocity, acceleration, and speed at $t = 1$.
 77. A drone is programmed to fly along the curve $\mathbf{r}(t) = \langle e^t, e^{-t}, t \rangle$. Calculate the distance it travels from $t = 0$ to $t = 1$.
 78. Compute the TNB frame for the curve $\mathbf{r}(t) = \langle t, \ln t, t^2 \rangle$ at $t = 1$.
 79. A particle moves along a helix $\mathbf{r}(t) = \langle \cos t, \sin t, t \rangle$. Compute its arc length from $t = 0$ to $t = 2\pi$.
 80. Find the arc length parameterization of $\mathbf{r}(t) = \langle t^3, t^2, t \rangle$ and use it to reparameterize the curve.
 81. Given $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$, compute the curvature and the torsion of the curve at $t = 1$.
 82. A satellite orbits Earth following the curve $\mathbf{r}(t) = \langle 7000 \cos t, 7000 \sin t, 0 \rangle$ km. Compute the arc length of one complete orbit.
 83. For the curve $\mathbf{r}(t) = \langle t, \ln t, \sqrt{t} \rangle$, find the arc length from $t = 0$ to $t = 2$.
 84. Given $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$, determine the speed function $|\mathbf{r}'(t)|$ and analyze its behavior as t approaches infinity.
 85. A roller coaster track is designed by $\mathbf{r}(t) = \langle 10 \cos t, 10 \sin t, t^2 \rangle$. Calculate the total distance traveled in one full loop from $t = 0$ to $t = 2\pi$.
 86. A drone must follow a path described by $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$ with constant speed. Find the appropriate reparameterization of the curve using arc length.
 87. Analyze the motion of a particle along $\mathbf{r}(t) = \langle t, e^t, \ln t \rangle$ by finding its velocity, acceleration, curvature, and torsion at $t = 1$.

Chapter 3

Partial Derivatives

Introduction

In our journey through calculus, we now extend our exploration to functions of several variables, building upon the foundations of single-variable calculus and vector-valued functions. Previously, we focused on how quantities change along curves, analyzing motion within a one-dimensional framework. However, many real-world phenomena depend on multiple variables simultaneously—such as temperature variations across a surface or pressure changes within a fluid.

This chapter delves into multivariable functions, where inputs and outputs exist in higher-dimensional spaces. We begin by examining how to visualize these functions using level curves and surfaces, setting the stage for understanding limits and continuity in higher dimensions. Central to our study are partial derivatives, which measure how a function changes with respect to one variable while keeping others constant. This generalizes the concept of derivatives from single-variable calculus and provides insights into the behavior of multivariable functions. We will learn to compute partial derivatives and explore higher-order derivatives, including Clairaut's Theorem on mixed partial derivatives.

Building on these concepts, we introduce directional derivatives and gradient vectors, which reveal how functions change in any specified direction. The gradient vector, in particular, points in the direction of the greatest rate of increase of the function and is fundamental in solving optimization problems. By integrating (pun intended) these new ideas with earlier concepts, we enhance our mathematical toolkit, enabling us to tackle complex problems involving multiple variables. The study of partial derivatives not only deepens our understanding of calculus but also prepares us for advanced topics in applied mathematics and real-world applications.

Lecture 3.1: Functions of Several Variables

In previous chapters, we have explored functions of a single variable and vector-valued functions, which map a single parameter to a vector in space. Now, we extend our study to *functions of several variables*, a fundamental concept in multivariable calculus that allows us to model and analyze phenomena where the output depends on multiple inputs. This lecture introduces the concept of functions of several variables, examines their domains and ranges, and explores techniques for visualizing them through graphs and level curves.

Functions of Two Variables

A function of two variables is a rule that assigns to each ordered pair (x, y) in a subset of $\mathbb{R}^2$ a unique real number $z = f(x, y)$. This can be thought of as a surface in three-dimensional space, where the height z above each point (x, y) represents the value of the function at that point. In mathematical terms, such a function can be expressed as $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, where $\mathbb{R} \times \mathbb{R}$ represents all possible ordered pairs (x, y) in two-dimensional space, and the output is a real number. To understand functions of two variables, recall our knowledge of single-variable functions $y = f(x)$:

- The graph of $f(x)$ is a curve in two dimensions, where y represents the output for a given input x .
- The domain is the set of all allowable x -values, and the range is the set of resulting y -values.

For functions of two variables, the domain is a subset of $\mathbb{R}^2$, typically a region in the xy -plane, and the range is a set of real numbers z . Graphically, the function $z = f(x, y)$ is represented as a surface in three-dimensional space. Functions of two variables appear naturally in many contexts that you may already be familiar with but haven't seen defined in the context of multivariate mathematics. For instance, here are some that you are certainly familiar with represented explicitly as multivariate functions.

- **Density:** Density is defined as mass divided by volume, so density is a function of mass and volume:

$$\rho = f(m, v) = \frac{m}{v}.$$

- **Einstein's Energy Equation:** Einstein's famous equation for relativistic energy depends on mass and velocity:

$$E = f(m, v) = \frac{mc^2}{\sqrt{1 - \frac{v^2}{c^2}}},$$

where c is the speed of light. Since c is constant, we typically consider energy as a function of the two variables m (mass) and v (velocity).

- **Taco Bell:** Suppose if I order some tacos for my family, we could represent this as such:

$$\text{Cost} = f(J, E, A) = \left[J \cdot \frac{\$d_1}{\# \text{ Gorditas}} + E \cdot \frac{\$d_2}{\# \text{ Crunchwraps}} + A \cdot \frac{\$d_3}{\# \text{ Tacos}} \right] \cdot 1.08,$$

where d_1, d_2 , and d_3 represent the prices per Gordita Crunch, Crunchwrap Supreme, and Soft Taco, respectively, and the 1.08 accounts for an 8% sales tax (I know, I know, I cheated and put a function of three variables for this one but the idea is the same). If I order 2 Gordita Crunches at \$4 each, my wife orders 1 Crunchwrap Supreme at \$5, and my son orders 3 Soft Tacos at \$2 each, the cost before tax is:

$$\text{Pre-tax Cost} = f(2, 1, 3) = 2 \cdot 4 + 1 \cdot 5 + 3 \cdot 2 = 8 + 5 + 6 = 19.$$

Including the 8% sales tax, the total cost becomes:

$$\text{Cost} = 19 \cdot 1.08 = 20.52.$$

This is an example where the domain of the function f is restricted to natural numbers or zero denoted as $\mathbb{N}_0$, as we can only order whole numbers (or zero) of each item (e.g., 2 Gorditas, not 2.5). Thus, the function maps from $\mathbb{N}_0 \times \mathbb{N}_0 \times \mathbb{N}_0$ (the number of Gorditas, Crunchwraps, and Tacos) to $\mathbb{Q}$, the set of rational numbers representing the total cost:

$$f : \mathbb{N}_0 \times \mathbb{N}_0 \times \mathbb{N}_0 \rightarrow \mathbb{Q}.$$

These examples illustrate how functions of several variables naturally arise in physical, scientific, and taco-related incidences, enabling us to describe and analyze relationships involving multiple (and potentially delicious) inputs.

▲ **Example 3.1.1** Consider functions that describe relationships between physical quantities. Write three examples of famous equations where one quantity is expressed as a function of exactly two other variables. Clearly identify the function and its meaning.

Solution Here are three examples:

1. **Ohm's Law:** Voltage is a function of current and resistance:

$$V = f(I, R) = I \cdot R,$$

where V is the voltage, I is the current, and R is the resistance.

2. **Power:** Power is a function of voltage and resistance:

$$P = f(V, R) = \frac{V^2}{R},$$

where ρ is the density, m is the mass, and V is the volume.

3. **Kinetic Energy:** Kinetic energy is a function of mass and velocity:

$$KE = f(m, v) = \frac{1}{2}mv^2,$$

where KE is the kinetic energy, m is the mass, and v is the velocity.

Domain and Range

The **domain** of a function $f(x, y)$ consists of all ordered pairs (x, y) for which $f(x, y)$ is defined. The **range** is the set of all real numbers z that $f(x, y)$ takes on as (x, y) varies over the domain. In single-variable calculus, determining the domain and range of a function involves identifying the set of input values x for which the function is defined and the corresponding set of output values $y = f(x)$. This extends naturally to functions of several variables.

For functions of two variables, the domain is often described as a subset of $\mathbb{R}^2$, or more generally $S \times S$, where S is a set of values satisfying specific restrictions. For instance, if $f(x, y) = \frac{1}{x^2 + y^2}$, the domain excludes the origin, and we can write:

$$\text{dom } f = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \neq 0\}.$$

More generally, for a function $g(x, y)$, the domain can be expressed as:

$$\text{dom } g = \{(x, y) \in S \times S \mid \text{restrictions on } x \text{ and } y\}.$$

To describe the range, there is less standardized notation, but for this text, we will use:

$$\text{range } f = \{z \in C \mid z = f(x, y) \text{ for some } (x, y) \in \text{dom } f\}.$$

Here, C represents the codomain of f , which is often $\mathbb{R}$ for real-valued functions. While this notation is not universally standard, it provides a concise way to describe the range, and we will adopt it for clarity in this context. For example, if $f(x, y) = x^2 + y^2$, we can write:

$$\text{range } f = \{z \in \mathbb{R}_{\geq 0} \mid z = x^2 + y^2 \text{ for some } (x, y) \in \mathbb{R}^2\}.$$

or just simply

$$\text{range } f = \{z \in \mathbb{R}_{\geq 0}\}.$$

Techniques for Determining the Domain Here are some general steps and considerations for finding the domain of $f(x, y)$:

1. **Identify Undefined Expressions:** Just like in single-variable functions, we need to find values of (x, y) that make the function undefined.

- **Denominators:** Ensure that any denominators are not equal to zero. For example, $f(x, y) = \frac{1}{x-y}$ is undefined at $x = y$.
 - **Square Roots and Even Roots:** The radicand (expression under the root) must be non-negative. For example, $g(x, y) = \sqrt{1 - x^2 - y^2}$ is defined where $x^2 + y^2 \leq 1$.
 - **Logarithms:** The argument of a logarithmic function must be positive. For instance, $h(x, y) = \ln(x + y)$ requires $x + y > 0$.
 - **Trigonometric Functions:** For inverse trigonometric functions, consider their specific restrictions. For instance, $\arcsin(x + y)$ is defined when $-1 \leq x + y \leq 1$.
2. **Set Notation:** Use inequalities or equations to describe the domain. For example, the domain of $k(x, y) = \sqrt{1 - x^2 - y^2}$ can be written as:

$$\text{dom } k = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}.$$

3. **Graphical Representation:** Visualize the domain in the xy -plane to better understand the set of allowed inputs. For example, the domain of $k(x, y)$ corresponds to the unit disk.

▲ **Example 3.1.2** Determine the domain and range of the following functions using proper notation:

1. $f(x, y) = \frac{1}{x+y}$
2. $g(x, y) = \frac{\ln(x-y)}{\sqrt{x^2+y^2-4}}$

Solution

1. For $f(x, y) = \frac{1}{x+y}$:

(a) **Domain:** The function is undefined when the denominator is zero:

$$x + y \neq 0.$$

Thus, the domain is:

$$\text{dom } f = \{(x, y) \in \mathbb{R}^2 \mid x + y \neq 0\}.$$

- (b) **Range:** The function $f(x, y)$ can take any nonzero real value because the reciprocal of $x + y$ is well-defined for all nonzero values of $x + y$. Since $f(x, y) = \frac{1}{x+y}$, the output cannot be zero because there is no $x + y$ for which $\frac{1}{x+y} = 0$. The range is:

$$\text{range } f = \mathbb{R} \setminus \{0\}.$$

2. For $g(x, y) = \frac{\ln(x-y)}{\sqrt{x^2+y^2-4}}$:

(a) **Domain:** The function is defined when both the logarithmic and square root expressions are valid:

- The logarithmic term $\ln(x - y)$ requires:

$$x - y > 0.$$

- The square root term $\sqrt{x^2 + y^2 - 4}$ requires:

$$x^2 + y^2 - 4 > 0, \quad \text{or equivalently, } x^2 + y^2 > 4.$$

Combining these conditions, the domain is:

$$\text{dom } g = \{(x, y) \in \mathbb{R}^2 \mid x - y > 0 \text{ and } x^2 + y^2 > 4\}.$$

(b) **Range:** The numerator $\ln(x - y)$ can take any real value since $x - y > 0$, and the denominator $\sqrt{x^2 + y^2 - 4}$ is strictly positive and increases as (x, y) moves farther from the circle $x^2 + y^2 = 4$. The output $g(x, y)$ cannot be zero because the numerator $\ln(x - y)$ cannot simultaneously be zero (i.e., $x - y = 1$) while satisfying the conditions imposed by the square root denominator. Therefore, the range is:

$$\text{range } g = \mathbb{R}.$$

Visualization Techniques

Visualizing a surface in $\mathbb{R}^3$, which represents the graph of a function $z = f(x, y)$, can initially seem daunting. Unlike the familiar $\mathbb{R}^2$ case, where a function of one variable is graphed as a curve in a plane, functions of two variables require interpreting surfaces in three dimensions. Several techniques can help us break this complexity into manageable parts and better understand the behavior of these surfaces.

- **3D Graphs:** The surface $z = f(x, y)$ in $\mathbb{R}^3$ visualizes the function across the entire domain. Tools like Python's Matplotlib, Plotly, and GeoGebra provide interactive 3D plots with features such as rotation, zoom, and color mapping to highlight z -values.
- **Level Curves:** Level curves (contour plots) represent constant z -values in the xy -plane, providing a 2D perspective of the function. These curves are essential for analyzing the surface's structure, especially in interactive tools that allow real-time exploration of contour spacing and behavior.
- **Cross-Sections:** Fixing one variable (e.g., $x = 0$ or $y = 0$) reduces the surface to a single-variable function, offering interpretable slices. These sections are useful for understanding surface behavior along specific planes.

- **Projections:** Projections allow us to collapse a surface into a lower-dimensional space by “projecting” it onto a plane. For example:

- Projecting the surface onto the xy -plane gives the domain of $f(x, y)$.
- Projecting onto the xz - or yz -plane reveals how z varies with x or y , respectively.

Projections become even more valuable when working with functions of more than two variables, as they allow us to reduce higher-dimensional objects into interpretable components in 2D or 3D.

- **Interactive Exploration:** Tools like GeoGebra, Plotly, and Wolfram Alpha enable real-time interaction with plots. Features like dynamic level curve visualization, color-coded z -values, and projections onto different planes make these tools invaluable for thoroughly analyzing functions.

Let’s begin visualize some graphs in three-dimensional space. Below is an example of a surface plot of a function of two variables. As is the case with single-variable mathematics, we can be a bit old-fashioned and use a Table of some values to aid us. In this case, we notice when x and y are equal, we are getting zero, thus indicating some interesting local extrema information.




Figure 3.1: Visualization of the function $z = x^2 - y^2$, showing the 3D surface plot and a table of input-output values.

The next figure demonstrates how plotting cross-sections simplifies understanding complex 3D surfaces. By combining slices with full 3D graphs, we can systematically analyze and interpret the behavior of functions of two variables.

Below is a visualization of the function $z = xy$, including a 3D surface plot and two 2D cross-sectional plots on the planes xz (at $y = -1$) and yz (at $x = 0$).




Figure 3.2: Visualization of the function $z = x \cdot y$, showing the 3D surface plot alongside 2D cross-sections on the planes xz (at $y = -1$) and yz (at $x = 1$).

We can see that by doing two 2D plots, we can reconstruct the plot visually with relative ease. In this particular example, it would be helpful to also see the case of $y = 1$ to indicate the “bend” in the 3D plot that we see. Just as you would spin an image around or a globe around to see all of its geometry, we essentially “spin” the plot around by plotting different slices from different planes.

Let’s take a look at an interesting applied example. The largest mountain in our solar system is Olympus Mons, located on Mars. One way to aid us in seeing the topology is to take cross sectional projections to see how this structure is formed. In fact, we know that Olympus Mons is a volcano in part because of the morphology and geological features obtained by such projections. In the picture below we slice two different sides of the mountain all the way through. Since the mountain is a volcano, we can especially see features such as the caldera much better using these slices.




Figure 3.3: *Elevation profiles of Olympus Mons along SW-to-NE and NW-to-SE transects across the mountain. Image from Wikipedia*

Level Curves

Level curves of a function $f(x, y)$ are the set of points in the xy -plane where the function takes on a constant value k , expressed by the equation $f(x, y) = k$. By plotting these curves for various values of k , we can visualize the function in two dimensions, gaining insights into its behavior without needing to represent the full three-dimensional surface. You may have even seen these types of plots on geographic plots. These are yet another tool that aid in understanding the shape of the terrain and identifying features like hills and valleys.




Figure 3.4: *A 3D surface featuring with its level curves shown below featuring the Bernoulli lemniscate (the infinity shaped level curve)*

Now that we have a basic understanding of what level curves are and why they are useful, let's

explore the process of plotting them in more detail. Level curves provide a two-dimensional representation of a function of two variables by showing all points (x, y) where the function takes on the same value k . This visualization is particularly helpful for understanding the behavior of multivariable functions without requiring a full 3D plot. Here's how to systematically plot level curves step by step:

1. **Select Values of k :** Choose several constant values for k that capture the key features of the function. It's important to select a range of k values—both positive and negative, if applicable—to represent the function's full behavior. For functions with periodic or oscillatory behavior, ensure the k -values align with significant points such as peaks, troughs, and zeros.
2. **Set $f(x, y) = k$:** For each chosen value of k , substitute it into the function $f(x, y)$ to define the corresponding level curve. This equation represents all points (x, y) where the function equals k .
3. **Solve for One Variable in Terms of the Other:** Rearrange the equation $f(x, y) = k$ to express y as a function of x , or x as a function of y , if possible. For example, in circular or radial symmetry, the equation might simplify to an expression for $r^2 = x^2 + y^2$. For more complex functions, this step may involve approximations or parametric forms.
4. **Plot Each Level Curve:** Using the equations derived, carefully plot the curves on the xy -plane. For simple equations, this can be done by hand. For more intricate equations, consider using graphing tools or software to ensure accuracy. Ensure that each curve is plotted with sufficient points to capture its shape.
5. **Label the Curves:** Clearly label each curve with its corresponding k -value. This can be done directly on the graph or through a legend. For enhanced clarity, consider using color coding to visually differentiate the curves based on their k -values.
6. **Analyze the Overall Behavior:** Once the level curves are plotted, examine their spacing and patterns. For instance, closely spaced curves indicate regions of rapid change in the function, while widely spaced curves suggest more gradual variation. This step helps in interpreting the nature of the function and its gradients.

▲ **Example 3.1.3** Consider the function $z = \sin(x^2 + y^2)$ defined for all $(x, y) \in \mathbb{R}^2$.

1. Plot the graph of $z = \sin(x^2 + y^2)$ as a surface in 3D space.
2. Plot the level curves of $z = \sin(x^2 + y^2)$ for selected z -values computed by hand.

Solution

1. The graph of $z = \sin(x^2 + y^2)$ exhibits circular symmetry. Recall that

$$\sin(\theta) = 1 \quad \text{if} \quad \theta = \frac{\pi}{2} + 2\pi n,$$

$$\sin(\theta) = -1 \quad \text{if} \quad \theta = -\frac{\pi}{2} + 2\pi n,$$

$$\sin(\theta) = 0 \quad \text{if} \quad \theta = \pi n,$$

for any integer n . Since $\theta = x^2 + y^2$, the peaks ($z = 1$) occur when

$$x^2 + y^2 = \frac{\pi}{2} + 2\pi n,$$

the troughs ($z = -1$) when

$$x^2 + y^2 = -\frac{\pi}{2} + 2\pi n,$$

and the zeros ($z = 0$) when

$$x^2 + y^2 = \pi n.$$




Figure 3.5: Graph of $z = \sin(x^2 + y^2)$, a sinusoidal surface.

2. To draw the level curves, we set $\sin(x^2 + y^2) = k$ for specific k -values. This gives:

- **For $z = 1$:** Solve

$$x^2 + y^2 = \frac{\pi}{2} + 2\pi n \quad \implies \quad r = \sqrt{\frac{\pi}{2} + 2\pi n}.$$

For example,

$$n = 0 : \quad r = \sqrt{\frac{\pi}{2}} \approx 1.25,$$

$$n = 1 : \quad r = \sqrt{\frac{5\pi}{2}} \approx 2.80$$

- **For $z = -1$:** Solve

$$x^2 + y^2 = -\frac{\pi}{2} + 2\pi n \quad (n \geq 1) \implies r = \sqrt{-\frac{\pi}{2} + 2\pi n}.$$

For example,

$$n = 1 : \quad r = \sqrt{\frac{3\pi}{2}} \approx 2.17,$$

$$n = 2 : \quad r = \sqrt{\frac{7\pi}{2}} \approx 3.32$$

- **For $z = 0$:** Solve

$$x^2 + y^2 = \pi n \implies r = \sqrt{\pi n}.$$

For example,

$$n = 1 : \quad r = \sqrt{\pi} \approx 1.77,$$

$$n = 2 : \quad r = \sqrt{2\pi} \approx 2.51$$




Figure 3.6: *Level curves for $z = \sin(x^2 + y^2)$ with additional rings.*

▲ **Example 3.1.4** Consider the function $z = \cos(y)$ defined for all $(x, y) \in \mathbb{R}^2$.

1. Plot the graph of $z = \cos(y)$ as a surface in 3D space.
2. Plot the level curves of $z = \cos(y)$ for selected z -values computed by hand.

Solution

1. The graph of $z = \cos(y)$ represents a cosine wave in the y -direction that is extruded along the x -axis. As y varies, z oscillates between -1 and 1 .




Figure 3.7: Graph of $z = \cos(y)$, a cosine wave surface extruded along x .

2. To find the level curves, we set $\cos(y) = k$ for specific values of k (with $-1 \leq k \leq 1$). The general solutions for y are:

$$\cos(y) = 1 \implies y = 0 + 2\pi n,$$

$$\cos(y) = -1 \implies y = \pi + 2\pi n,$$

$$\cos(y) = 0 \implies y = \frac{\pi}{2} + \pi n,$$

where n is any integer. For a few specific level curves, we choose:

- $z = 1$: $y = 0, 2\pi, -2\pi, \dots$
- $z = -1$: $y = \pi, -\pi, 3\pi, \dots$
- $z = 0$: $y = \frac{\pi}{2}, -\frac{\pi}{2}, \frac{3\pi}{2}, \dots$

Since x is unrestricted, the level curves in the xy -plane are horizontal lines at these y -values.




Figure 3.8: *Level curves for $z = \cos(y)$ in the xy -plane.*

Common 3D Equations

As is the case in single-variable mathematics, we can depend on familiar examples to help make sense of functions on sight.

- **Quadric Surfaces:** Paraboloids, hyperboloids, and ellipsoids are common surfaces that can often be graphed as functions of x and y . These surfaces are studied in more detail in Lecture 1.5. Examples include:

- Paraboloid: $z = x^2 + y^2$
- Hyperboloid of One Sheet: $x^2 + y^2 - z^2 = 1$
- Ellipsoid: $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$

- **Sphere:** A sphere represents a special case of an ellipsoid where all axes are equal. The equation is:

$$x^2 + y^2 + z^2 = r^2$$

- **Plane:** A plane can be described as a linear function in three variables. For example:

$$z = ax + by + c$$

- **Cylinder:** Cylinders are surfaces generated by extending a curve along a straight line parallel to an axis. Examples include:

- Circular Cylinder: $x^2 + y^2 = r^2$
- Parabolic Cylinder: $z = x^2$

- **Cone:** A cone is a surface that extends a curve in a way that all points of the curve are connected to a single vertex. For example:

$$z^2 = x^2 + y^2$$

Functions of Three or More Variables

A function of three variables assigns a real number $w = f(x, y, z)$ to each point (x, y, z) within a subset of $\mathbb{R}^3$. The challenge in working with such functions lies in the fact that their graphs exist in four-dimensional space, a realm beyond our ability to visualize directly. Unlike the graph of a function of one variable, which forms a curve in $\mathbb{R}^2$, or a function of two variables, which forms a surface in $\mathbb{R}^3$, the graph of $f(x, y, z)$ resists geometric representation. Despite this limitation, a variety of powerful techniques enable us to analyze and understand these functions with clarity and precision. Specifically, the most comprehensive way that I can think about it is by the analogy of a three-dimensional shadow: a shadow reduces a 3D object down to 2D, therefore a 3D object is the shadow of a 4D object. This means that projections will be our greatest ally.

To streamline notation for functions of multiple variables, it is common to represent the input (x, y, z) as a single entity, often written as $\mathbf{x}$. This allows us to express $f(x, y, z)$ in a more concise form:

$$f(\mathbf{x}),$$

where $\mathbf{x}$ represents the vector of input variables:

$$\mathbf{x} = (x, y, z).$$

This notation generalizes naturally to functions of n variables, where $\mathbf{x} = (x_1, x_2, \dots, x_n)$ is a vector in $\mathbb{R}^n$. To emphasize the structure of the input variables, we can further represent $\mathbf{x}$ as a column vector:

$$\mathbf{x} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}.$$

With this notation, the function $f(x, y, z)$ becomes $f(\mathbf{x})$, where $\mathbf{x}$ is the input vector. Alternatively, we can express $\mathbf{x}$ explicitly as a column vector:

$$f(\mathbf{x}) = f\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right),$$

providing a unified framework for analyzing functions of several variables. For instance, a temperature field T in three-dimensional space can be expressed as:

$$T = f(\mathbf{t}) = f\left(\begin{bmatrix} t_x \\ t_y \\ t_z \end{bmatrix}\right) = f(t_x, t_y, t_z) = t_x^2 + t_y^2 + t_z^2$$

where $\mathbf{x}$ represents a point in $\mathbb{R}^3$. This column vector-based notation is particularly valuable in advanced topics like gradients, divergence, and vector-valued functions, as it simplifies both expression and computation.

Graphs and Visualization

While we cannot graph functions of three variables directly in three-dimensional space, we can understand them through level surfaces and projections. A **level surface** is the set of all points (x, y, z) in the domain of $f(x, y, z)$ where the function takes a constant value, $w = k$. These surfaces are three-dimensional analogues of level curves for functions of two variables. For example, fixing $w = 0$, the level surface $f(x, y, z) = 0$ provides a snapshot of the function's behavior in three-dimensional space.

The idea is analogous to projections, where we imagine the three-dimensional representation of a function as the “shadow” of its four-dimensional graph. By holding one variable constant, such as $z = 0$, we reduce the function to $w = f(x, y, 0)$, which can be plotted as a surface in three-dimensional space. Repeating this process for various fixed values of z (or any other variable) gives a series of surfaces that collectively reveal the structure of the original function. This approach bridges the gap between our spatial intuition and the abstraction of higher-dimensional spaces.

Level Surfaces

Plotting a level surface, will be very similar to our level curves. lets take a look at a hypersphere $f(x, y, z) = x^2 + y^2 + z^2$. The level surfaces are determined by solving:

$$f(x, y, z) = k \implies x^2 + y^2 + z^2 = k.$$

For different values of k , the level surfaces represent spheres centered at the origin with radius $\sqrt{k}$:

- $k = 1$: Sphere of radius 1.
- $k = 4$: Sphere of radius 2.
- $k = 9$: Sphere of radius 3.

To visualize these level surfaces, we plot the spheres for $k = 1, 4, 9$. The surfaces demonstrate the isotropic nature of the function $f(x, y, z)$, where the value of w depends only on the distance from the origin.




Figure 3.9: *Level surfaces of $f(x, y, z) = x^2 + y^2 + z^2 = k$ for $k = 1, 4, 9$, represented as concentric circles*

While this doesn't exactly paint the complete picture, you may begin to see how hypergeometry works. Perhaps you have seen a hypercube represented before but not sure why it looks like it does. With level surfaces it may begin to make more sense:




Figure 3.10: Cubes of dimensions 1 through 5. Above three dimensions, these are often called *hypercubes*, or *n-cubes*. Image from Wikipedia

Visualizing Functions of Several Variables

As the number of variables in a function increases, direct visualization becomes increasingly challenging. For functions of two or three variables, graphs and level sets provide intuitive insights. However, for functions involving more than three variables, such as $f(x_1, x_2, \dots, x_n)$, we rely on abstract representations, projections, and numerical tools to understand their behavior.

Higher-Dimensional Analogues: In higher dimensions, familiar geometric objects like lines, planes, and spheres generalize:

- A **hyperplane** in $\mathbb{R}^n$ is defined by a linear equation, such as $a_1x_1 + a_2x_2 + \dots + a_nx_n = c$. It is the n -dimensional analogue of a plane in $\mathbb{R}^3$.
- A **hypersphere** in $\mathbb{R}^n$ is defined by $x_1^2 + x_2^2 + \dots + x_n^2 = r^2$. In $\mathbb{R}^4$, for example, this represents the set of points equidistant from a center, analogous to a sphere in $\mathbb{R}^3$.

Applications in Machine Learning: Functions of several variables are fundamental in machine learning, particularly in classification tasks. For instance:

- **Hyperplanes in Classification:** In supervised learning, hyperplanes are often used to separate data points belonging to different classes. For example, a linear classifier uses a hyperplane to divide the feature space into regions corresponding to different categories.

- **Support Vector Machines (SVMs):** SVMs are powerful classification algorithms that construct an optimal hyperplane to maximize the margin between data points of different classes. Mathematically, the hyperplane is defined as:

$$\mathbf{w} \cdot \mathbf{x} - b = 0,$$

where $\mathbf{w}$ is the weight vector, $\mathbf{x}$ is a data point, and b is a bias term. The margin is maximized by solving an optimization problem involving the constraint:

$$y_i(\mathbf{w} \cdot \mathbf{x}_i - b) \geq 1,$$

where y_i represents the class label of $\mathbf{x}_i$.




Figure 3.11: *Hyperplanes used for optimal classification in SVMs. Image from Wikipedia*

Although these applications arise in higher dimensions, the underlying concepts of hyperplanes and optimization are rooted in multivariable calculus.

Lecture 3.2: Limits and Continuity

In single-variable calculus, the concepts of limits and continuity form the foundation for understanding the behavior of functions near points of interest and for establishing the differentiability and integrability of functions. In multivariable calculus, these ideas extend naturally but require a more nuanced treatment due to the increased complexity introduced by multiple independent variables. This lecture delves into the concepts of limits and continuity in the context of multivariable functions, exploring their definitions, properties, and challenges.

Limits in Higher Dimensions

The concept of a limit is a cornerstone of calculus and extends naturally to functions of several variables. However, understanding this extension requires a thoughtful approach, as the introduction of additional variables adds layers of complexity.

In single-variable calculus, we define the limit $\lim_{x \rightarrow a} f(x) = L$ as the value L that $f(x)$ approaches as x gets arbitrarily close to a . For multivariable functions, the same principle applies, but now the input (x, y) can approach a target point (a, b) from infinitely many directions in the plane. This distinction is key: in one dimension, there are only two possible approaches to a (from the left or right), whereas in two dimensions, there are infinitely many paths.




Figure 3.12: Graph of $z = \frac{xy}{x^2+y^2}$, where the limit does not exist at $(0,0)$

To understand limits intuitively, imagine the behavior of a function $f(x, y)$ as you move closer and closer to a fixed point (a, b) . The value of $f(x, y)$ should stabilize and become arbitrarily close to a specific number L , regardless of the path taken. If the value of $f(x, y)$ changes depending on the path, the limit does not exist.

Properties of Multivariable Limits

Before diving into techniques for evaluating limits, it is helpful to understand some basic properties that govern multivariable limits. These properties, analogous to their single-variable counterparts, provide useful shortcuts and insights into the behavior of multivariable functions as they approach a limit.

- **Linearity:** If $\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$ and $\lim_{(x,y) \rightarrow (a,b)} g(x, y) = M$, then for any constants c_1 and c_2 :

$$\lim_{(x,y) \rightarrow (a,b)} [c_1 f(x, y) + c_2 g(x, y)] = c_1 L + c_2 M.$$

- **Product Rule:** If $\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$ and $\lim_{(x,y) \rightarrow (a,b)} g(x, y) = M$, then:

$$\lim_{(x,y) \rightarrow (a,b)} [f(x, y)g(x, y)] = L \cdot M.$$

- **Quotient Rule:** If $\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$ and $\lim_{(x,y) \rightarrow (a,b)} g(x, y) = M$, with $M \neq 0$, then:

$$\lim_{(x,y) \rightarrow (a,b)} \frac{f(x, y)}{g(x, y)} = \frac{L}{M}.$$

- **Composition Rule:** If $\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$ and $\lim_{z \rightarrow L} g(z) = N$, where g is continuous at L , then:

$$\lim_{(x,y) \rightarrow (a,b)} g(f(x, y)) = g(L).$$

- **Limits of Polynomials and Rational Functions:** For polynomial functions $f(x, y)$ and $g(x, y)$, the limit as $(x, y) \rightarrow (a, b)$ can often be found by direct substitution:

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b).$$

For rational functions $f(x, y)/g(x, y)$, this holds as long as $g(a, b) \neq 0$.

- **Continuity and Limits:** If $f(x, y)$ is continuous at (a, b) , then the limit as $(x, y) \rightarrow (a, b)$ is simply the value of the function at that point:

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b).$$

These properties simplify the evaluation of many limits and help identify when more advanced techniques, such as polar coordinates or path dependence analysis, may be necessary. Combined with the techniques described in the following section, they form the backbone of solving multivariable limit problems.

Basic Techniques for Evaluating Limits

While the formal definition of a limit provides a rigorous foundation, solving limits in practice often relies on algebraic manipulation and analytical techniques familiar from single-variable calculus. These techniques aim to simplify the function or identify patterns that reveal the behavior of $f(x, y)$ (or $f(x, y, z)$, etc.) as the input approaches a specific point. Below are common methods used for evaluating multivariable limits:

- **Substitution:** If the function is continuous at the point of interest, substitution is the simplest and most direct method. For example, to evaluate $\lim_{(x,y) \rightarrow (a,b)} f(x, y)$, substitute $x = a$ and $y = b$ into $f(x, y)$. If the resulting expression is well-defined, the limit is equal to the function's value at that point.
- **Factoring and Simplification:** Similar to single-variable calculus, factoring can help simplify expressions and eliminate terms causing indeterminate forms. For example, for functions like $\frac{x^2 - y^2}{x - y}$, factoring the numerator reveals simplifications.
- **Rationalizing Using Conjugates:** For functions involving square roots, multiplying by the conjugate can simplify the expression and resolve indeterminate forms. For example:

$$\lim_{(x,y) \rightarrow (0,0)} \frac{\sqrt{x^2 + y^2} - 1}{x^2 + y^2}$$

can be simplified by multiplying numerator and denominator by $\sqrt{x^2 + y^2} + 1$.

- **Path Dependence Analysis:** To determine whether a limit exists, evaluate the function along different paths approaching the same point. Common paths include:
 - Horizontal or vertical lines, such as $y = 0$ or $x = 0$.
 - Diagonal lines, such as $y = mx$, where m is a constant slope.
 - Curved paths, such as $y = x^2$ or $y = \sin(x)$.

If the limit depends on the chosen path, the overall limit does not exist.

- **Polar Coordinates:** For limits approaching the origin in two dimensions, converting to polar coordinates can simplify expressions. Substituting $x = r \cos \theta$ and $y = r \sin \theta$ reduces a multivariable problem to a single variable r , where $r \rightarrow 0$. For example:

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 + y^2}{x^2 - y^2}$$

becomes:

$$\lim_{r \rightarrow 0} \frac{r^2}{r^2(\cos^2 \theta - \sin^2 \theta)}.$$

These techniques provide a powerful toolkit for solving a wide variety of multivariable limit problems. While some methods, such as path analysis and polar coordinates, are specific to functions of multiple variables, others are naturally inherited from single-variable calculus.

Keep in mind, not only can we take limits from $(x, y) \rightarrow (a, b)$, but $(x, y, z) \rightarrow (a, b, c)$ and higher dimensions. Let's now work some of these out.

▲ **Example 3.2.1** Evaluate the following limits:

1. $\lim_{(x,y) \rightarrow (0,0)} x^2 + y^2$
2. $\lim_{(x,y) \rightarrow (1,1)} \frac{\sqrt{x}-\sqrt{y}}{x-y}$
3. $\lim_{(x,y,z) \rightarrow (0,0,0)} \frac{\sin(x^2+y^2+z^2)}{x^2+y^2+z^2}$
4. $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2-y^2}{x^2+y^2}$

Solution

1. For $\lim_{(x,y) \rightarrow (0,0)} x^2 + y^2$:

• **Direct Substitution:** Since $x^2 + y^2$ is continuous, directly substituting $(x, y) = (0, 0)$ gives:

$$\lim_{(x,y) \rightarrow (0,0)} x^2 + y^2 = 0^2 + 0^2 = 0.$$

2. For $\lim_{(x,y) \rightarrow (1,1)} \frac{\sqrt{x}-\sqrt{y}}{x-y}$:

• **Multiply by the Conjugate:** To simplify, multiply numerator and denominator by $\sqrt{x} + \sqrt{y}$:

$$\frac{\sqrt{x}-\sqrt{y}}{x-y} \cdot \frac{\sqrt{x}+\sqrt{y}}{\sqrt{x}+\sqrt{y}} = \frac{(\sqrt{x}-\sqrt{y})(\sqrt{x}+\sqrt{y})}{(x-y)(\sqrt{x}+\sqrt{y})} = \frac{x-y}{(x-y)(\sqrt{x}+\sqrt{y})}.$$

Simplifying, we find:

$$\frac{1}{\sqrt{x}+\sqrt{y}}.$$

Substituting $(x, y) = (1, 1)$:

$$\lim_{(x,y) \rightarrow (1,1)} \frac{\sqrt{x}-\sqrt{y}}{x-y} = \frac{1}{\sqrt{1}+\sqrt{1}} = \frac{1}{2}.$$

3. For $\lim_{(x,y,z) \rightarrow (0,0,0)} \frac{\sin(x^2+y^2+z^2)}{x^2+y^2+z^2}$:

- **Convert to Spherical Coordinates:** Let $r^2 = x^2 + y^2 + z^2$, where $r \rightarrow 0$ as $(x, y, z) \rightarrow (0, 0, 0)$. The expression becomes:

$$\frac{\sin(x^2 + y^2 + z^2)}{x^2 + y^2 + z^2} = \frac{\sin(r^2)}{r^2}.$$

Using the substitution $u = r^2$, where $u \rightarrow 0^+$ as $r \rightarrow 0^+$, the limit simplifies to:

$$\lim_{u \rightarrow 0^+} \frac{\sin(u)}{u}.$$

From single-variable calculus, it is known that:

$$\lim_{u \rightarrow 0} \frac{\sin(u)}{u} = 1.$$

Thus:

$$\lim_{(x,y,z) \rightarrow (0,0,0)} \frac{\sin(x^2 + y^2 + z^2)}{x^2 + y^2 + z^2} = 1.$$

4. For $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x^2 + y^2}$:

- **Path Dependence:** The value of the limit depends on the path taken to approach $(0, 0)$.

(a) **Path 1:** $y = 0$: Substituting $y = 0$ into the function:

$$\frac{x^2 - y^2}{x^2 + y^2} = \frac{x^2 - 0^2}{x^2 + 0^2} = \frac{x^2}{x^2} = 1.$$

(b) **Path 2:** $x = 0$: Substituting $x = 0$ into the function:

$$\frac{x^2 - y^2}{x^2 + y^2} = \frac{0^2 - y^2}{0^2 + y^2} = \frac{-y^2}{y^2} = -1.$$

(c) **Path 3:** $y = mx$: Substituting $y = mx$ into the function:

$$\frac{x^2 - (mx)^2}{x^2 + (mx)^2} = \frac{x^2 - m^2x^2}{x^2 + m^2x^2}.$$

Simplify:

$$\frac{1 - m^2}{1 + m^2}.$$

Since the limits differ based on the path taken:

The limit does not exist.

Other Techniques

Beyond our core toolkit for evaluating multivariable limits, we can extend several powerful techniques from single-variable calculus into multiple dimensions. These methods often allow us to reduce complex multivariable problems into more manageable pieces that connect to our existing understanding.

The Squeeze Theorem extends naturally to multivariable functions, providing a powerful tool when direct evaluation proves difficult. If we can find two functions that “squeeze” our target function and share the same limit, then our target function must approach that limit as well. Formally, if for all points (x, y) in some neighborhood of (a, b) :

$$g(x, y) \leq f(x, y) \leq h(x, y)$$

and we have

$$\lim_{(x,y) \rightarrow (a,b)} g(x, y) = \lim_{(x,y) \rightarrow (a,b)} h(x, y) = L,$$

then we can conclude

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L.$$

To understand this intuitively, imagine the graphs of $g(x, y)$ and $h(x, y)$ forming a “corridor” in three-dimensional space that contains the graph of $f(x, y)$. As we approach the point (a, b) , this corridor narrows to a single point with height L , forcing our function $f(x, y)$ to approach the same value.

When encountering indeterminate forms like $\frac{0}{0}$, we can apply reasoning similar to L’Hôpital’s Rule by treating one variable as constant and analyzing the limit with respect to the other variable. While we haven’t yet developed the machinery of partial derivatives, we can still analyze these limits by considering one variable at a time. For instance, when approaching a point (a, b) , we might:

- First analyze $f(x, b)$ as $x \rightarrow a$, treating y as the constant b
- Then analyze $f(a, y)$ as $y \rightarrow b$, treating x as the constant a
- Compare these single-variable limits

For a function exhibiting an indeterminate form $\frac{0}{0}$, we can write:

$$\lim_{x \rightarrow a} \frac{f(x, b)}{g(x, b)} = \lim_{x \rightarrow a} \frac{f'(x, b)}{g'(x, b)}$$

where the derivatives are taken with respect to x while treating y as the constant b . However, we must exercise caution: equality of these single-variable limits is necessary but not sufficient for the existence of the multivariable limit. The multivariable nature of the functions introduces subtleties not present in single-variable calculus, particularly the need to verify that the limit is independent of the path taken to (a, b) .

These advanced techniques complement our earlier methods, often providing elegant solutions to complex limit problems that might otherwise prove intractable. Their application requires careful attention to the underlying assumptions and verification of path independence, but they significantly expand our ability to analyze multivariable limits.

▲ **Example 3.2.2** Evaluate the following limits using the advanced techniques we've just learned:

1. $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$
2. $\lim_{(x,y) \rightarrow (0,0)} xy \sin\left(\frac{1}{x^2 + y^2}\right)$
3. $\lim_{(x,y) \rightarrow (0,2)} \frac{e^{xy} - 1}{x}$
4. $\lim_{(x,y) \rightarrow (3,0)} \frac{\sqrt{x+y} - \sqrt{x}}{y}$

Solution

1. For $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$:

- **Using the Squeeze Theorem:** Let's find bounds for this function. First, note that:

$$\left| \frac{x^2 y}{x^2 + y^2} \right| = |y| \cdot \left| \frac{x^2}{x^2 + y^2} \right|$$

Since $\frac{x^2}{x^2 + y^2} \leq 1$ for all $(x, y) \neq (0, 0)$, we have:

$$\left| \frac{x^2 y}{x^2 + y^2} \right| \leq |y|$$

Therefore:

$$-|y| \leq \frac{x^2 y}{x^2 + y^2} \leq |y|$$

As $(x, y) \rightarrow (0, 0)$, we know $|y| \rightarrow 0$. By the Squeeze Theorem:

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} = 0$$

2. For $\lim_{(x,y) \rightarrow (0,0)} xy \sin\left(\frac{1}{x^2 + y^2}\right)$:

- **Using the Squeeze Theorem:** First, recall that $|\sin(z)| \leq 1$ for all z . Therefore:

$$\left| xy \sin \left(\frac{1}{x^2 + y^2} \right) \right| \leq |xy|$$

Now we need to show that $|xy| \rightarrow 0$ as $(x, y) \rightarrow (0, 0)$.

Let's use polar coordinates: $x = r \cos \theta$, $y = r \sin \theta$. Then:

$$|xy| = |r \cos \theta \cdot r \sin \theta| = r^2 |\cos \theta \sin \theta| \leq \frac{r^2}{2}$$

Therefore:

$$-\frac{r^2}{2} \leq xy \sin \left(\frac{1}{x^2 + y^2} \right) \leq \frac{r^2}{2}$$

As $(x, y) \rightarrow (0, 0)$, $r \rightarrow 0$, so by the Squeeze Theorem:

$$\lim_{(x,y) \rightarrow (0,0)} xy \sin \left(\frac{1}{x^2 + y^2} \right) = 0$$

Note: Even though $\frac{1}{x^2 + y^2}$ becomes unbounded as $(x, y) \rightarrow (0, 0)$, the bounded nature of sine combined with the factor xy controls the overall behavior of the function.

3. For $\lim_{(x,y) \rightarrow (0,2)} \frac{e^{xy} - 1}{x}$:

- **Fixing $y = 2$ and applying L'Hôpital's Rule:** As $y \rightarrow 2$, we treat the limit as a function of x :

$$\lim_{x \rightarrow 0} \frac{e^{2x} - 1}{x}.$$

This is a $\frac{0}{0}$ indeterminate form. Differentiating with respect to x (with y held constant):

$$\frac{d}{dx}(e^{2x} - 1) = 2e^{2x}, \quad \frac{d}{dx}(x) = 1.$$

Therefore, the limit becomes:

$$\lim_{x \rightarrow 0} 2e^{2x} = 2e^0 = 2.$$

4. For $\lim_{(x,y) \rightarrow (3,0)} \frac{\sqrt{x+y} - \sqrt{x}}{y}$:

- **Fixing $x = 3$ and applying L'Hôpital's Rule:** As $x \rightarrow 3$, consider the limit as a function of y :

$$\lim_{y \rightarrow 0} \frac{\sqrt{3+y} - \sqrt{3}}{y}.$$

This gives a $\frac{0}{0}$ indeterminate form. Differentiating with respect to y :

$$\frac{d}{dy}(\sqrt{3+y}) = \frac{1}{2\sqrt{3+y}}, \quad \frac{d}{dy}(y) = 1.$$

Hence, the limit simplifies to:

$$\lim_{y \rightarrow 0} \frac{1}{2\sqrt{3+y}} = \frac{1}{2\sqrt{3}}.$$

Formal Definition

To rigorously define the limit, we need a precise way to measure “closeness” for points in two dimensions. Luckily, from Chapter 1 we know how to do this now. In the $\mathbb{R}^2$ plane, the distance between two points (x, y) and (a, b) is given by the Euclidean distance formula:

$$\sqrt{(x-a)^2 + (y-b)^2}.$$

We say that $\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$ if we can make the value of $f(x, y)$ arbitrarily close to L (within an error margin of ϵ) by ensuring that (x, y) is sufficiently close to (a, b) (within a distance of δ).

The formal definition is expressed as follows:

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$$

means that for every $\epsilon > 0$, there exists a $\delta > 0$ such that:

$$0 < \sqrt{(x-a)^2 + (y-b)^2} < \delta \implies |f(x, y) - L| < \epsilon.$$

Here’s what this means step by step:

- ϵ (epsilon): A small positive number that represents how close $f(x, y)$ must be to L . Think of this as the allowable error in the output of the function.
- δ (delta): A small positive number that represents how close (x, y) must be to (a, b) . This defines a circular region (a disk) around (a, b) with radius δ .
- The condition $0 < \sqrt{(x-a)^2 + (y-b)^2} < \delta$ ensures that (x, y) is close to (a, b) but not exactly at (a, b) , as the limit depends on the behavior of the function near (a, b) , not at (a, b) itself.
- The implication $|f(x, y) - L| < \epsilon$ ensures that the function value is within ϵ of L whenever (x, y) is within δ of (a, b) .

To visualize this concept, imagine a mountain with a peak at (a, b, L) . As you climb the mountain from different starting points, you must arrive at the same height L to conclude that the peak has a well-defined elevation. If different paths lead to different heights, the notion of a single “peak height” breaks down, and the limit does not exist.




Figure 3.13: 3D plot with a δ -circle, center point (a, b) , a transparent surface, $f(a, b)$, the circle plotted on the surface, and the corresponding points L , $L + \epsilon$, and $L - \epsilon$ on the z -axis connected to their corresponding surface points.

The concept of limits extends to functions of three or more variables. For a function $f(x, y, z)$, the limit as $(x, y, z) \rightarrow (a, b, c)$ is defined similarly:

$$\lim_{(x,y,z) \rightarrow (a,b,c)} f(x, y, z) = L$$

if for every $\epsilon > 0$, there exists a $\delta > 0$ such that:

$$0 < \sqrt{(x - a)^2 + (y - b)^2 + (z - c)^2} < \delta \implies |f(x, y, z) - L| < \epsilon.$$

The principles remain the same, but the added dimension increases the complexity of path dependence and visualization. Tools like polar, cylindrical, or spherical coordinates are often used to simplify calculations.

This idea generalizes further to functions of n -variables, $f(x_1, x_2, \dots, x_n)$. The limit as $(x_1, x_2, \dots, x_n) \rightarrow (a_1, a_2, \dots, a_n)$ is defined as:

$$\lim_{(x_1, x_2, \dots, x_n) \rightarrow (a_1, a_2, \dots, a_n)} f(x_1, x_2, \dots, x_n) = L$$

if for every $\epsilon > 0$, there exists a $\delta > 0$ such that:

$$0 < \sqrt{\sum_{i=1}^n (x_i - a_i)^2} < \delta \implies |f(x_1, x_2, \dots, x_n) - L| < \epsilon.$$

Here, the distance metric generalizes to the Euclidean norm in n -dimensional space:

$$\sqrt{\sum_{i=1}^n (x_i - a_i)^2},$$

which measures the distance between the point $(x_1, x_2, \dots, x_n)$ and the target $(a_1, a_2, \dots, a_n)$. The ϵ - δ framework ensures that as $(x_1, x_2, \dots, x_n)$ approaches $(a_1, a_2, \dots, a_n)$ in this generalized space, the function value $f(x_1, x_2, \dots, x_n)$ approaches L , independent of the path taken. Alternatively, the concept of limits can be compactly expressed using vector notation. For a function $f(\mathbf{x})$ where $\mathbf{x} = (x_1, x_2, \dots, x_n)$ and the target point is $\mathbf{a} = (a_1, a_2, \dots, a_n)$, the limit is written as:

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} f(\mathbf{x}) = L,$$

if for every $\epsilon > 0$, there exists a $\delta > 0$ such that:

$$0 < \|\mathbf{x} - \mathbf{a}\| < \delta \implies |f(\mathbf{x}) - L| < \epsilon.$$

▲ Example 3.2.3 Using the formal definition of limits for multivariable functions, prove that:

$$\lim_{(x,y) \rightarrow (0,0)} xy = 0.$$

Solution To prove that:

$$\lim_{(x,y) \rightarrow (0,0)} xy = 0,$$

we need to show that for every $\epsilon > 0$, there exists a $\delta > 0$ such that whenever $0 < \sqrt{(x-0)^2 + (y-0)^2} < \delta$, it follows that:

$$|xy - 0| < \epsilon.$$

Proof: Let $\epsilon > 0$ be given. We aim to find $\delta > 0$ such that if $0 < \sqrt{x^2 + y^2} < \delta$, then $|xy| < \epsilon$. Note that:

$$|xy| = |x||y| \leq \left(\sqrt{x^2}\right) \left(\sqrt{y^2}\right) = |x||y|.$$

But since $|x| < \sqrt{x^2 + y^2}$ and $|y| < \sqrt{x^2 + y^2}$ for $x, y \neq 0$ we have:

$$|xy| < \left(\sqrt{x^2 + y^2}\right) \left(\sqrt{x^2 + y^2}\right) = x^2 + y^2.$$

Furthermore, since $x^2 + y^2 = \left(\sqrt{x^2 + y^2}\right)^2$, it follows that:

$$|xy| < \left(\sqrt{x^2 + y^2}\right)^2.$$

Now, choose $\delta = \sqrt{\epsilon}$. Then $\delta > 0$ because $\epsilon > 0$. Suppose $0 < \sqrt{x^2 + y^2} < \delta$. Then:

$$|xy| < \left(\sqrt{x^2 + y^2}\right)^2 < \delta^2 = (\sqrt{\epsilon})^2 = \epsilon.$$

Therefore, whenever $0 < \sqrt{x^2 + y^2} < \delta$, it follows that $|xy| < \epsilon$.

Conclusion: For every $\epsilon > 0$, we have found $\delta = \sqrt{\epsilon} > 0$ such that:

$$0 < \sqrt{x^2 + y^2} < \delta \implies |xy - 0| < \epsilon.$$

Hence, by the formal definition of limits for multivariable functions,

$$\lim_{(x,y) \rightarrow (0,0)} xy = 0.$$

Continuity in Multivariable Functions

Continuity is a fundamental concept in calculus, naturally extending from single-variable functions to multivariable functions. Intuitively, a function $f(x, y)$ is continuous at a point (a, b) if small changes in (x, y) near (a, b) result in small changes in $f(x, y)$. Formally, $f(x, y)$ is continuous at (a, b) if:

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b).$$

This definition implies three necessary conditions:

1. $f(x, y)$ is defined at (a, b) .
2. The limit $\lim_{(x,y) \rightarrow (a,b)} f(x, y)$ exists.
3. The limit equals the function value: $\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b)$.

The idea of continuity in multiple dimensions generalizes the familiar notion of a smooth, unbroken curve or surface.

Visualizing Continuity: To better understand continuity, imagine a surface in three-dimensional space representing $z = f(x, y)$. Continuity at a point (a, b) ensures that as (x, y) approaches (a, b) from any direction, the surface height z converges to a single value $f(a, b)$. Discontinuities manifest as abrupt breaks, gaps, or sharp jumps in the surface, where the smoothness of the surface is compromised. A simple example of a continuous function is $f(x, y) = x^2 + y^2$, which forms a paraboloid, while a function like $f(x, y) = \frac{1}{x^2 + y^2}$ is discontinuous at $(0, 0)$, as the value of $f(x, y)$ becomes unbounded near this point.

Continuity on a Domain: A function $f(x, y)$ is continuous on a region $R \subseteq \mathbb{R}^2$ if it is continuous at every point $(x, y) \in R$. These regions can be:

- **Open Regions:** Regions excluding their boundary points, such as the interior of a circle.
- **Closed Regions:** Regions including their boundary points, such as a closed square or disk.
- **Bounded and Unbounded Regions:** Continuity can be considered within finite or infinite domains.

For example, the function $f(x, y) = x^2 + y^2$ is continuous everywhere on $\mathbb{R}^2$, while $f(x, y) = \frac{1}{x^2 + y^2}$ is continuous only on $\mathbb{R}^2 \setminus \{(0, 0)\}$.

Properties of Continuous Functions: Multivariable functions share several properties of continuous single-variable functions, providing tools for analyzing their behavior:

- **Sum, Difference, and Product:** If f and g are continuous at (a, b) , then $f + g$, $f - g$, and fg are also continuous at (a, b) .
- **Quotients:** If f and g are continuous at (a, b) and $g(a, b) \neq 0$, then f/g is continuous at (a, b) .
- **Composition:** If f is continuous at (a, b) and g is continuous at $f(a, b)$, then $g \circ f$ is continuous at (a, b) .

Discontinuities: Discontinuities in multivariable functions can be classified as:

- **Removable Discontinuities:** Occur when a limit exists at (a, b) but does not equal $f(a, b)$. For example, a “hole” in the surface can be filled by redefining $f(a, b)$ to match the limit.
- **Jump Discontinuities:** Occur when the limit does not exist due to different values along different paths. For instance, consider $f(x, y) = \begin{cases} 1 & \text{if } x = 0, \\ 0 & \text{if } x \neq 0. \end{cases}$
- **Essential Discontinuities:** Occur when no limit exists at (a, b) , such as in functions with oscillatory behavior near (a, b) .

High-Dimensional Analogues of Theorems: Several familiar theorems from single-variable calculus extend to multivariable functions, often requiring additional assumptions or interpretations. Note that some terminology and notation will be further revealed in later chapters as they become relevant.

- **Intermediate Value Theorem:** If $f(x, y)$ is continuous on a connected region R and takes on two distinct values a and b , then $f(x, y)$ takes every value between a and b at some point in R . This is particularly useful in multivariable optimization problems or proving the existence of level curves.

- **Mean Value Theorem:** For multivariable functions, this theorem generalizes to state that if f is continuous and differentiable on a region R , then there exists a point $\mathbf{x}_c \in R$ such that the gradient (multi-dimensional derivative) $\nabla f(\mathbf{x}_c)$ aligns with the direction of the greatest change in f over R . This result is crucial in applications such as error estimation and optimization.
- **Extreme Value Theorem:** If $f(x, y, z)$ is continuous on a closed and bounded region $R \subseteq \mathbb{R}^3$, then f attains both a maximum and minimum value on R . This theorem underpins much of multivariable optimization, particularly when combined with Lagrange multipliers.
- **Squeeze Theorem:** The multivariable version states that if $g(x, y) \leq f(x, y) \leq h(x, y)$ for all $(x, y) \in R$, and $\lim_{(x,y) \rightarrow (a,b)} g(x, y) = \lim_{(x,y) \rightarrow (a,b)} h(x, y) = L$, then $\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$. This theorem is invaluable in proving the existence of limits.

▲ **Example 3.2.4** Verify the Intermediate Value Theorem for the function

$$f(x, y) = x^2 + y^2 - 1$$

on the closed region $R = \{(x, y) \in \mathbb{R}^2 \mid -1 \leq x, y \leq 1\}$, and show that there exists a point $(a, b) \in R$ such that $f(a, b) = 0$.

Solution The function $f(x, y) = x^2 + y^2 - 1$ is a polynomial in x and y , so it is continuous on $\mathbb{R}^2$, including the closed and bounded region R .

- **Step 1: Evaluate f at specific points in R .** At $(-1, 0)$, we have:

$$f(-1, 0) = (-1)^2 + 0^2 - 1 = 0.$$

At $(0, 0)$, we have:

$$f(0, 0) = 0^2 + 0^2 - 1 = -1.$$

At $(1, 1)$, we have:

$$f(1, 1) = 1^2 + 1^2 - 1 = 1.$$

- **Step 2: Apply the Intermediate Value Theorem.** Because $f(x, y)$ is continuous on R , and $f(-1, 0) = 0$, $f(0, 0) = -1$, and $f(1, 1) = 1$, it must take every value between -1 and 1 at some point in R .

In particular, since $f(-1, 0) = 0$, there is no further verification needed to show that $f(x, y) = 0$ holds within R . However, the Intermediate Value Theorem guarantees that $f(x, y) = c$ for any $c \in [-1, 1]$.

- **Conclusion:** The Intermediate Value Theorem verifies that there exists a point $(a, b) \in R$, such as $(-1, 0)$, where $f(a, b) = 0$.
-

Lecture 3.3: Partial Derivatives

Imagine you're at an all-you-can-eat buffet. You've loaded your plate with chicken, mashed potatoes, and gravy, but now you're debating how many meatballs to pile on. The chicken is staying put, the mashed potatoes aren't going anywhere, and the gravy is committed to the cause—but the meatballs? They're the variable. This is partial derivatives in a nutshell: you're holding everything else constant while analyzing how one particular choice impacts your meal (or, in this case, your mathematical function). Just as you ask, “How many meatballs is too many?” partial derivatives ask, “How does the function change if only one variable changes?” This section explores the formal definition, interpretation, and properties of partial derivatives, including higher-order derivatives and Clairaut's theorem.

Definition and Notation

Recall from single-variable calculus that the derivative of a function $f(x)$ at a point $x = a$ is defined as:

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h},$$

which measures the instantaneous rate of change of $f(x)$ with respect to x . This idea naturally extends to functions of several variables by allowing one variable to change while holding others constant. For a function $f(x, y)$ defined in a region of $\mathbb{R}^2$, we define the *partial derivative* of f with respect to x , denoted $\frac{\partial f}{\partial x}$ or f_x , as:

$$\frac{\partial f}{\partial x}(x, y) = \lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h},$$

provided the limit exists. Here, y is held constant, and x is allowed to vary. Similarly, the partial derivative with respect to y is defined as:

$$\frac{\partial f}{\partial y}(x, y) = \lim_{h \rightarrow 0} \frac{f(x, y+h) - f(x, y)}{h}.$$

This definition can be understood as a direct application of the single-variable derivative, applied to one variable at a time. By holding one variable fixed, we reduce the function $f(x, y)$ to a single-variable function, enabling us to compute its rate of change with respect to the variable of interest.

For functions of three or more variables, such as $f(x_1, x_2, \dots, x_n)$, the partial derivative with respect to x_i is similarly defined:

$$\frac{\partial f}{\partial x_i}(x_1, x_2, \dots, x_n) = \lim_{h \rightarrow 0} \frac{f(x_1, \dots, x_i+h, \dots, x_n) - f(x_1, \dots, x_i, \dots, x_n)}{h},$$

where only x_i is perturbed, while all other variables are held constant. This generalization allows us to study functions in higher-dimensional spaces, providing a foundation for advanced topics like gradients and optimization.

▲ **Example 3.3.1** Use the limit definition of a partial derivative to compute $\frac{\partial f}{\partial x}$ for $f(x, y) = x^2y + 3xy^2$.

Solution Recall the limit definition of the partial derivative:

$$\frac{\partial f}{\partial x}(x, y) = \lim_{h \rightarrow 0} \frac{f(x + h, y) - f(x, y)}{h}.$$

Substitute $f(x, y) = x^2y + 3xy^2$ into the definition:

$$\frac{\partial f}{\partial x}(x, y) = \lim_{h \rightarrow 0} \frac{f(x + h, y) - f(x, y)}{h}.$$

Compute $f(x + h, y)$:

$$f(x + h, y) = (x + h)^2y + 3(x + h)y^2.$$

Expand the terms:

$$f(x + h, y) = (x^2 + 2xh + h^2)y + 3(xy^2 + hy^2) = x^2y + 2xhy + h^2y + 3xy^2 + 3hy^2.$$

Now subtract $f(x, y)$:

$$f(x + h, y) - f(x, y) = (x^2y + 2xhy + h^2y + 3xy^2 + 3hy^2) - (x^2y + 3xy^2).$$

Simplify:

$$f(x + h, y) - f(x, y) = 2xhy + h^2y + 3hy^2.$$

Divide by h :

$$\frac{f(x + h, y) - f(x, y)}{h} = \frac{2xhy + h^2y + 3hy^2}{h}.$$

Simplify further:

$$\frac{f(x + h, y) - f(x, y)}{h} = 2xy + hy + 3y^2.$$

Take the limit as $h \rightarrow 0$:

$$\lim_{h \rightarrow 0} \frac{f(x + h, y) - f(x, y)}{h} = 2xy + 3y^2.$$

Thus:

$$\frac{\partial f}{\partial x}(x, y) = 2xy + 3y^2.$$

The partial derivative $\frac{\partial f}{\partial x}(x, y)$ measures the rate of change of $f(x, y)$ with respect to x , while keeping y fixed. Geometrically, this corresponds to the slope of the tangent line to the curve obtained by intersecting the surface $z = f(x, y)$ with the plane $y = \text{constant}$. Similarly, $\frac{\partial f}{\partial y}$ represents the slope of the tangent line to the curve obtained by intersecting the surface

with the plane $x = \text{constant}$. These partial derivatives capture the local behavior of the surface along the coordinate axes.

Several standard notations are used for partial derivatives, depending on the context:

- $\frac{\partial f}{\partial x}$: Highlights the operator ∂ , distinguishing it from total derivatives.
- f_x : A compact notation often used in applied contexts.
- $\partial_x f$: A concise form useful in symbolic computation and advanced texts.

These notations emphasize that partial derivatives focus on the effect of one variable at a time, isolating its contribution to the overall behavior of $f(x, y)$. By leveraging the limit definition and geometric intuition, we gain a comprehensive understanding of how multivariable functions behave locally in their domains.




Figure 3.14: *Derivative of a surface $z = x^2 + y^2$ at a point $(1, 1, 2)$. The tangent line (red) is notated with projections of each component, green indicating yz -plane projection of the tangent line, and blue indicating the xz -plane projection.*

Basic Properties of Partial Derivatives.

Partial derivatives share many properties with single-variable derivatives. The following are some fundamental properties of partial derivatives that are useful for computation and analysis:

- **Linearity:** If $f(x, y)$ and $g(x, y)$ are differentiable functions, and c is a constant, then:

$$\frac{\partial}{\partial x}(cf(x, y) + g(x, y)) = c\frac{\partial f}{\partial x} + \frac{\partial g}{\partial x}.$$

- **Product Rule:** If $f(x, y)$ and $g(x, y)$ are differentiable, then:

$$\frac{\partial}{\partial x}(f(x, y)g(x, y)) = f_x g + f g_x.$$

- **Quotient Rule:** If $f(x, y)$ and $g(x, y)$ are differentiable, and $g(x, y) \neq 0$, then:

$$\frac{\partial}{\partial x} \left(\frac{f(x, y)}{g(x, y)} \right) = \frac{f_x g - f g_x}{g^2}.$$

- **Chain Rule (Simple Form):** If $z = f(x, y)$, where x and y are functions of another variable t , then:

$$\frac{dz}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

The chain rule gets a little hairy, and hence we will dedicate an entire lecture to it later on.

- **Zero Partial Derivative:** If $\frac{\partial f}{\partial x} = 0$ in a region, then $f(x, y)$ does not change with x in that region (it depends only on y).
- **Partial Derivatives of Constants:** If c is a constant, then:

$$\frac{\partial c}{\partial x} = 0 \quad \text{and} \quad \frac{\partial c}{\partial y} = 0.$$

▲ **Example 3.3.2** Compute the partial derivatives of the following functions with respect to all specified variables:

1. $f(x, y) = x^2 y + 3xy^2$ (with respect to x and y)
2. $g(x, y, z) = e^{xz} \sin(y)$ (with respect to x , y , and z)
3. $h(x, y) = \frac{x^2 + y^2}{x - y}$ (with respect to x and y)

Solution

1. For $f(x, y) = x^2 y + 3xy^2$:
 - Partial derivative with respect to x :

$$\frac{\partial f}{\partial x} = \frac{\partial}{\partial x}(x^2 y) + \frac{\partial}{\partial x}(3xy^2) = 2xy + 3y^2.$$

- Partial derivative with respect to y :

$$\frac{\partial f}{\partial y} = \frac{\partial}{\partial y}(x^2 y) + \frac{\partial}{\partial y}(3xy^2) = x^2 + 6xy.$$

2. For $g(x, y, z) = e^{xz} \sin(y)$:

- Partial derivative with respect to x :

$$\frac{\partial g}{\partial x} = \frac{\partial}{\partial x} (e^{xz} \sin(y)) = ze^{xz} \sin(y).$$

- Partial derivative with respect to y :

$$\frac{\partial g}{\partial y} = \frac{\partial}{\partial y} (e^{xz} \sin(y)) = e^{xz} \cos(y).$$

- Partial derivative with respect to z :

$$\frac{\partial g}{\partial z} = \frac{\partial}{\partial z} (e^{xz} \sin(y)) = xe^{xz} \sin(y).$$

3. For $h(x, y) = \frac{x^2+y^2}{x-y}$:

- Partial derivative with respect to x (using the quotient rule):

$$\frac{\partial h}{\partial x} = \frac{(2x)(x-y) - (x^2+y^2)(1)}{(x-y)^2} = \frac{2x(x-y) - (x^2+y^2)}{(x-y)^2}.$$

Simplify:

$$\frac{\partial h}{\partial x} = \frac{x^2 - 2xy - y^2}{(x-y)^2}.$$

- Partial derivative with respect to y (using the quotient rule):

$$\frac{\partial h}{\partial y} = \frac{(2y)(x-y) - (x^2+y^2)(-1)}{(x-y)^2} = \frac{2y(x-y) + (x^2+y^2)}{(x-y)^2}.$$

Simplify:

$$\frac{\partial h}{\partial y} = \frac{2xy - 2y^2 + x^2 + y^2}{(x-y)^2}.$$

Higher-Order Partial Derivatives

Higher-order partial derivatives extend the concept of derivatives to capture rates of change of rates of change, just as in single-variable calculus. For a function $f(x, y)$, the second-order partial derivatives include:

$$\frac{\partial^2 f}{\partial x^2}, \quad \frac{\partial^2 f}{\partial y^2}, \quad \text{and} \quad \frac{\partial^2 f}{\partial x \partial y}, \quad \frac{\partial^2 f}{\partial y \partial x}.$$

These derivatives are often written more compactly using subscript notation:

$$f_{xx} = \frac{\partial^2 f}{\partial x^2}, \quad f_{yy} = \frac{\partial^2 f}{\partial y^2}, \quad f_{xy} = \frac{\partial^2 f}{\partial x \partial y}, \quad f_{yx} = \frac{\partial^2 f}{\partial y \partial x}.$$

These derivatives reveal local properties of the function, such as curvature, along specific directions. Mixed partial derivatives, such as $\frac{\partial^2 f}{\partial x \partial y}$ or f_{xy} , involve differentiating first with respect to one variable and then with respect to another:

$$\begin{aligned} \frac{\partial^2 f}{\partial x \partial y} &= \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right), \quad \text{or equivalently,} \quad f_{xy} = \frac{\partial}{\partial y} (f_x), \\ \frac{\partial^2 f}{\partial y \partial x} &= \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right), \quad \text{or equivalently,} \quad f_{yx} = \frac{\partial}{\partial x} (f_y). \end{aligned}$$

This naturally extends to higher dimensions. For a function $f(x_1, x_2, \dots, x_n)$, higher-order derivatives can be computed repeatedly by differentiating with respect to different variables. For example, the general k -th order partial derivative is denoted as:

$$\frac{\partial^k f}{\partial x_{i_1} \partial x_{i_2} \cdots \partial x_{i_k}},$$

where $i_1, i_2, \dots, i_k$ indicate the sequence of variables being differentiated. Using subscript notation, this becomes:

$$f_{x_{i_1} x_{i_2} \cdots x_{i_k}},$$

highlighting the order of differentiation. This generalization mirrors the iterative process of differentiation from single-variable calculus.

These derivatives are vital in analyzing the behavior of multivariable functions. For instance:

- **Curvature Analysis:** Second-order derivatives reveal concavity and convexity, analogous to the second derivative test in single-variable calculus.
- **Critical Points:** Higher-order partial derivatives help identify extrema and saddle points, topics that will be discussed in a future lecture.
- **Taylor Expansions:** Higher-order derivatives are fundamental in approximating functions using Taylor series in multiple dimensions.

Does the Order of Differentiation Matter? A natural question arises: does the order of differentiation affect the result? That is, is:

$$\frac{\partial^2 f}{\partial x \partial y} \stackrel{?}{=} \frac{\partial^2 f}{\partial y \partial x}?$$

Surprisingly—or perhaps reassuringly—the answer is that the order generally does **not** matter, provided certain conditions are met.

Clairaut's Theorem on Mixed Partial Derivatives

Clairaut's Theorem states that if f is defined on a region $R \subseteq \mathbb{R}^n$ and all second-order partial derivatives of f are continuous on R , then the mixed partial derivatives are equal:

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x}.$$

This result is both powerful and intuitive, as it guarantees symmetry in the order of differentiation under appropriate continuity conditions. To generalize, for $f(x_1, x_2, \dots, x_n)$:

$$\frac{\partial^2 f}{\partial x_i \partial x_j} = \frac{\partial^2 f}{\partial x_j \partial x_i}, \quad \text{for all } i, j \text{ such that the derivatives are continuous.}$$

Proof Outline of Clairaut's Theorem. The proof relies on the definition of partial derivatives and the interchangeability of limits for continuous functions:

1. Write the mixed partial derivative $\frac{\partial^2 f}{\partial x \partial y}$ as:

$$\frac{\partial^2 f}{\partial x \partial y} = \lim_{h \rightarrow 0} \frac{\frac{\partial f}{\partial x}(x, y+h) - \frac{\partial f}{\partial x}(x, y)}{h}.$$

2. Substitute the definition of $\frac{\partial f}{\partial x}$ in terms of limits, and observe that the nested limits commute if f is continuous.
3. Symmetry emerges from the commutativity of the limits, leading to:

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x}.$$

4. Continuity of the second-order partial derivatives ensures the validity of this reasoning.

Counterexample: Consider the function $f(x, y)$ defined as:

$$f(x, y) = \begin{cases} \frac{xy}{x^2+y^2}, & \text{if } (x, y) \neq (0, 0), \\ 0, & \text{if } (x, y) = (0, 0). \end{cases}$$

Compute the mixed partial derivatives at $(0, 0)$:

- $\frac{\partial f}{\partial x} = \begin{cases} \frac{y(x^2-y^2)}{(x^2+y^2)^2}, & \text{if } (x, y) \neq (0, 0), \\ 0, & \text{if } (x, y) = (0, 0). \end{cases}$
- Taking $\frac{\partial}{\partial y}$ and $\frac{\partial}{\partial x}$ in different orders produces inconsistent results, as the second-order partial derivatives involve discontinuities in their behavior near $(0, 0)$.

This failure arises because $f(x, y)$ is not continuously differentiable at $(0, 0)$, violating the conditions of Clairaut's Theorem.

Note: the existence of partial derivatives does not guarantee the continuity of a function. A function can have partial derivatives at a point without being continuous there. However, if all partial derivatives are continuous in a region, the function is said to be *continuously differentiable*, denoted C^1 . This property is critical for advanced topics, such as optimization and differential equations.

▲ **Example 3.3.3** Verify Clairaut's Theorem for the following functions:

1. $f(x, y) = x^2y + y^3$
2. $f(x, y) = e^{xy}$

Solution

1. For $f(x, y) = x^2y + y^3$:

- Compute $\frac{\partial f}{\partial x}$:

$$\frac{\partial f}{\partial x} = \frac{\partial}{\partial x}(x^2y + y^3) = 2xy.$$

- Compute $\frac{\partial^2 f}{\partial x \partial y}$:

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial y}(2xy) = 2x.$$

- Compute $\frac{\partial f}{\partial y}$:

$$\frac{\partial f}{\partial y} = \frac{\partial}{\partial y}(x^2y + y^3) = x^2 + 3y^2.$$

- Compute $\frac{\partial^2 f}{\partial y \partial x}$:

$$\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial x}(x^2 + 3y^2) = 2x.$$

Since $\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x} = 2x$, Clairaut's Theorem is verified.

2. For $f(x, y) = e^{xy}$:

- Compute $\frac{\partial f}{\partial x}$:

$$\frac{\partial f}{\partial x} = \frac{\partial}{\partial x}(e^{xy}) = ye^{xy}.$$

- Compute $\frac{\partial^2 f}{\partial x \partial y}$:

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial y}(ye^{xy}) = e^{xy} + y^2 e^{xy}.$$

- Compute $\frac{\partial f}{\partial y}$:

$$\frac{\partial f}{\partial y} = \frac{\partial}{\partial y}(e^{xy}) = xe^{xy}.$$

- Compute $\frac{\partial^2 f}{\partial y \partial x}$:

$$\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial x}(xe^{xy}) = e^{xy} + y^2 e^{xy}.$$

Since $\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x} = e^{xy} + y^2 e^{xy}$, Clairaut's Theorem is verified.

▲ **Example 3.3.4** A company is designing a new container with a rectangular base to minimize material usage. The volume of the container is fixed at $V = 12$ cubic meters, and the base dimensions are x (length) and y (width), while the height is h . The material for the base and sides costs the same, but the material for the top is more expensive.

The total surface area of the container, representing material usage, is given by:

$$S(x, y, h) = xy + 2(xh + yh) + cxy,$$

where $c > 1$ accounts for the higher cost of the top material.

1. Express h in terms of x and y using the volume constraint $V = xyh$.
2. Substitute h into the surface area formula to reduce $S(x, y, h)$ to a function of x and y only: $S(x, y)$.
3. Find the first-order partial derivatives S_x and S_y . What do they tell you about how the surface area changes with respect to each variable?
4. Using the second-order partial derivatives S_{xx} , S_{yy} , and S_{xy} , describe what might happen at a point where both $S_x = 0$ and $S_y = 0$. What might this point represent in terms of the design?

Solution

1. From the volume constraint $V = xyh$, solve for h :

$$h = \frac{12}{xy}.$$

2. Substitute $h = \frac{12}{xy}$ into $S(x, y, h)$:

$$\begin{aligned} S(x, y) &= xy + 2(xh + yh) + cxy \\ &= xy + 2\left(x\left(\frac{12}{xy}\right) + y\left(\frac{12}{xy}\right)\right) + cxy \\ &= xy + 2\left(\frac{12}{y} + \frac{12}{x}\right) + cxy \\ &= xy + \frac{24}{y} + \frac{24}{x} + cxy. \end{aligned}$$

3. Compute the first-order partial derivatives:

$$\begin{aligned} S_x &= \frac{\partial S}{\partial x} = y - \frac{24}{x^2} + cy, \\ S_y &= \frac{\partial S}{\partial y} = x - \frac{24}{y^2} + cx. \end{aligned}$$

These derivatives indicate how the surface area changes with respect to changes in x and y while keeping the other variable fixed.

4. Compute the second-order partial derivatives:

$$\begin{aligned} S_{xx} &= \frac{\partial^2 S}{\partial x^2} = \frac{48}{x^3}, \\ S_{yy} &= \frac{\partial^2 S}{\partial y^2} = \frac{48}{y^3}, \\ S_{xy} &= \frac{\partial^2 S}{\partial x \partial y} = 1 + c. \end{aligned}$$

At a point where $S_x = 0$ and $S_y = 0$, the second-order derivatives provide information about the curvature of the surface area function. This could help determine whether the point is a minimum, maximum, or saddle point, which has implications for optimizing material usage.

This problem naturally leads to questions about the nature of critical points, gradients, and their role in optimization, which will be explored in upcoming lectures.

Lecture 3.4: The Chain Rule

The Chain Rule in calculus serves as a foundational tool for handling composite functions, enabling us to compute derivatives when variables are interconnected through other variables. In multivariable calculus, the Chain Rule generalizes elegantly, accounting for the relationships between dependent and independent variables across multiple dimensions. This section explores the nuances of the Chain Rule for multivariable functions, implicit differentiation, and total differentials.

Chain Rule for Multivariable Functions

The Chain Rule is a cornerstone of calculus, extending the concept of derivatives to account for compositions of functions across multiple variables. Its utility lies in describing how a dependent variable changes when its inputs themselves depend on other variables. To see how this works, we start with the familiar single-variable Chain Rule, which states that if $z = f(x)$ and $x = x(t)$, then the total derivative of z with respect to t is:

$$\frac{dz}{dt} = \frac{df}{dx} \frac{dx}{dt}.$$

This formula captures the idea that the change in z is the product of the rate of change of f with respect to x and the rate of change of x with respect to t .

In a multivariable setting, suppose $z = f(x, y)$, where x and y are functions of a third variable t . The total derivative of z with respect to t incorporates the contributions of both x and y :

$$\frac{dz}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

Each term represents a pathway of dependence: $\frac{\partial f}{\partial x}$ accounts for how z changes as x changes, weighted by $\frac{dx}{dt}$, and similarly for y . This structure reflects the additive nature of change, where each variable contributes independently to the overall rate of change in z .

Generalizing further, consider $z = f(x_1, x_2, \dots, x_n)$, where each x_i depends on t . The total derivative is:

$$\frac{dz}{dt} = \sum_{i=1}^n \frac{\partial f}{\partial x_i} \frac{dx_i}{dt}.$$

Here, the sum encompasses all pathways of dependence, with $\frac{\partial f}{\partial x_i}$ representing the sensitivity of f to x_i and $\frac{dx_i}{dt}$ capturing the rate of change of x_i with respect to t . Now suppose $z = f(x, y)$, but x and y depend on two independent variables u and v . In this case, we compute partial derivatives of z with respect to u and v :

$$\begin{aligned} \frac{\partial z}{\partial u} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u}, \\ \frac{\partial z}{\partial v} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial v}. \end{aligned}$$

This formulation generalizes the single-variable Chain Rule by distributing the dependence of z on u and v through the intermediate variables x and y . To simplify these expressions in higher dimensions, we can employ matrix notation. Let $\mathbf{X} = [x, y]^T$ and $\mathbf{U} = [u, v]^T$, with f a function of $\mathbf{X}$. The Chain Rule can be written compactly as:

$$\frac{\partial f}{\partial \mathbf{U}} = \frac{\partial f}{\partial \mathbf{X}} \cdot \frac{\partial \mathbf{X}}{\partial \mathbf{U}},$$

where:

$$\frac{\partial f}{\partial \mathbf{X}} = \begin{bmatrix} \frac{\partial f}{\partial x} & \frac{\partial f}{\partial y} \end{bmatrix}, \quad \frac{\partial \mathbf{X}}{\partial \mathbf{U}} = \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix}.$$

The product of these matrices computes the total derivatives of f with respect to u and v in a concise form, encapsulating the dependence of f on multiple layers of variables.

This structure highlights the broader implications of the Chain Rule. In essence, it quantifies how changes propagate through a system of interdependent variables. This principle will play a pivotal role in future discussions, such as gradients, optimization, and Jacobian matrices, where understanding these dependencies becomes critical for analyzing and solving complex problems.

▲ **Example 3.4.1** Compute the derivatives in the following scenarios using the Chain Rule:

1. Let $z = x^2y$, where $x = t^2$ and $y = \sin(t)$. Find $\frac{dz}{dt}$.
2. Let $z = \ln(xy)$, where $x = e^u$ and $y = \cos(v)$. Compute $\frac{\partial z}{\partial u}$ and $\frac{\partial z}{\partial v}$.

Solution

1. For $z = x^2y$, where $x = t^2$ and $y = \sin(t)$:

- Use the Chain Rule:

$$\frac{dz}{dt} = \frac{\partial z}{\partial x} \frac{dx}{dt} + \frac{\partial z}{\partial y} \frac{dy}{dt}.$$

- Compute the partial derivatives:

$$\frac{\partial z}{\partial x} = 2xy, \quad \frac{\partial z}{\partial y} = x^2.$$

- Substitute $x = t^2$ and $y = \sin(t)$:

$$\frac{dx}{dt} = 2t, \quad \frac{dy}{dt} = \cos(t).$$

- Combine terms:

$$\frac{dz}{dt} = 2(t^2)(\sin(t))(2t) + (t^2)^2(\cos(t)).$$

Simplify:

$$\frac{dz}{dt} = 4t^3 \sin(t) + t^4 \cos(t).$$

2. For $z = \ln(xy)$, where $x = e^u$ and $y = \cos(v)$:

- Compute $\frac{\partial z}{\partial u}$ using the Chain Rule:

$$\frac{\partial z}{\partial u} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial u}.$$

- Compute the partial derivatives:

$$\frac{\partial z}{\partial x} = \frac{1}{x}, \quad \frac{\partial x}{\partial u} = e^u.$$

- Substitute $x = e^u$ and $y = \cos(v)$:

$$\frac{\partial z}{\partial u} = \frac{1}{e^u \cos(v)} \cdot e^u = \frac{1}{\cos(v)}.$$

- Compute $\frac{\partial z}{\partial v}$:

$$\frac{\partial z}{\partial v} = \frac{\partial z}{\partial y} \frac{\partial y}{\partial v}.$$

- Compute the partial derivatives:

$$\frac{\partial z}{\partial y} = \frac{1}{y}, \quad \frac{\partial y}{\partial v} = -\sin(v).$$

- Substitute $y = \cos(v)$:

$$\frac{\partial z}{\partial v} = \frac{1}{\cos(v)}(-\sin(v)) = -\tan(v).$$

▲ **Example 3.4.2** Find $\frac{\partial^2}{\partial y \partial x} f$ if $f = g(u(x, y), v(y))$ where $u(x, y) = \cos(xy)$ and $v(y) = \ln(1/y)$.

Solution First, compute $\frac{\partial f}{\partial x}$ using the chain rule:

$$\frac{\partial f}{\partial x} = g_u \frac{\partial h}{\partial x} = g_u(-y \sin(xy))$$

where g_u denotes $\frac{\partial g}{\partial u}$.

Now, compute $\frac{\partial}{\partial y}$ of this result using the product rule and chain rule:

$$\begin{aligned} \frac{\partial^2 f}{\partial y \partial x} &= \frac{\partial}{\partial y} [g_u(-y \sin(xy))] \\ &= \left(g_{uu} \frac{\partial h}{\partial y} + g_{uv} \frac{\partial k}{\partial y} \right) (-y \sin(xy)) + g_u (-\sin(xy) - xy \cos(xy)) \\ &= \left(g_{uu}(-x \sin(xy)) + g_{uv} \left(-\frac{1}{y} \right) \right) (-y \sin(xy)) + g_u (-\sin(xy) - xy \cos(xy)) \\ &= g_{uu}xy \sin^2(xy) + g_{uv} \sin(xy) - g_u(\sin(xy) + xy \cos(xy)) \end{aligned}$$

Therefore:

$$\frac{\partial^2 f}{\partial y \partial x} = g_{uu}xy \sin^2(xy) + g_{uv} \sin(xy) - g_u(\sin(xy) + xy \cos(xy))$$

Here, g_u denotes $\frac{\partial g}{\partial u}$, g_{uu} denotes $\frac{\partial^2 g}{\partial u^2}$, and g_{uv} denotes $\frac{\partial^2 g}{\partial u \partial v}$, where $g = g(u, v)$.

Implicit Differentiation

Implicit differentiation arises when a relationship between variables is defined implicitly rather than explicitly. This concept is familiar from single-variable calculus, where we often encounter equations like $x^2 + y^2 = 1$, representing a circle. In this case, y is not given explicitly as a function of x , but the equation defines a relationship between x and y . Differentiating such equations requires us to treat y as a function of x , even if its explicit form is unknown. Recall the procedure for implicit differentiation in single-variable calculus:

$$\frac{d}{dx} (x^2 + y^2) = \frac{d}{dx} (1).$$

Applying the chain rule to the y^2 term:

$$2x + 2y \frac{dy}{dx} = 0.$$

Solving for $\frac{dy}{dx}$ gives:

$$\frac{dy}{dx} = -\frac{x}{y}.$$

This process illustrates how implicit differentiation allows us to find derivatives without isolating y , which can be algebraically cumbersome or even impossible.

In multivariable calculus, implicit differentiation extends naturally to functions of several variables. Suppose we have a relationship $F(x, y) = 0$, where y is implicitly a function of x . Differentiating both sides with respect to x yields:

$$\frac{\partial F}{\partial x} + \frac{\partial F}{\partial y} \frac{dy}{dx} = 0.$$

This expression follows from the Chain Rule, as F depends on both x and y , with y depending on x . Solving for $\frac{dy}{dx}$:

$$\frac{dy}{dx} = -\frac{\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial y}}, \quad \text{provided } \frac{\partial F}{\partial y} \neq 0.$$

The result generalizes the single-variable case, where partial derivatives replace ordinary derivatives to account for the multivariable nature of F . For functions of three variables, consider $F(x, y, z) = 0$, where z is implicitly a function of x and y . Differentiating implicitly, we treat z as dependent on x and y :

$$\frac{\partial F}{\partial x} + \frac{\partial F}{\partial y} \frac{\partial y}{\partial x} + \frac{\partial F}{\partial z} \frac{\partial z}{\partial x} = 0.$$

From here, solving for $\frac{\partial z}{\partial x}$ gives:

$$\frac{\partial z}{\partial x} = -\frac{\frac{\partial F}{\partial x}}{\frac{\partial F}{\partial z}}, \quad \text{assuming } \frac{\partial F}{\partial z} \neq 0.$$

Similarly, differentiating $F(x, y, z) = 0$ with respect to y gives:

$$\frac{\partial z}{\partial y} = -\frac{\frac{\partial F}{\partial y}}{\frac{\partial F}{\partial z}}.$$

These expressions demonstrate how implicit differentiation extends to higher dimensions, allowing us to compute derivatives for implicitly defined functions of multiple variables. Implicit differentiation plays a critical role in multivariable calculus, enabling us to analyze curves and surfaces defined implicitly by equations like $F(x, y, z) = 0$. For instance:

- In geometry, it provides a method to compute the slopes of tangent lines or tangent planes without solving explicitly for one variable in terms of the others.
- In optimization, it lays the foundation for solving constrained optimization problems using techniques like the method of Lagrange multipliers.




Figure 3.15: A tangent line to the sphere $x^2 + y^2 + z^2 = 9$ at the point $(2, 1, \sqrt{5})$. The tangent line is shown in red. The point of tangency is marked in black.

This generalization of implicit differentiation highlights the power of the Chain Rule in understanding complex interdependencies among variables, bridging concepts from calculus to more advanced topics in analysis and applied mathematics.

▲ **Example 3.4.3** For the implicit function $F(x, y, z) = x^2y + yz + e^z = 10$, where z is a function of x and y , find $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.

Solution Differentiate both sides of the equation $F(x, y, z) = 10$ with respect to x , treating z as a function of x and y :

$$\frac{\partial}{\partial x}(x^2y + yz + e^z) = \frac{\partial}{\partial x}(10).$$

Applying the chain rule:

$$\frac{\partial}{\partial x}(x^2y) + \frac{\partial}{\partial x}(yz) + \frac{\partial}{\partial x}(e^z) = 0.$$

Compute each term:

$$\frac{\partial}{\partial x}(x^2y) = 2xy, \quad \frac{\partial}{\partial x}(yz) = y\frac{\partial z}{\partial x}, \quad \frac{\partial}{\partial x}(e^z) = e^z\frac{\partial z}{\partial x}.$$

Combine the terms:

$$2xy + y\frac{\partial z}{\partial x} + e^z\frac{\partial z}{\partial x} = 0.$$

Group terms involving $\frac{\partial z}{\partial x}$:

$$(y + e^z) \frac{\partial z}{\partial x} = -2xy.$$

Solve for $\frac{\partial z}{\partial x}$:

$$\frac{\partial z}{\partial x} = \frac{-2xy}{y + e^z}, \quad \text{provided } y + e^z \neq 0.$$

Now differentiate $F(x, y, z)$ with respect to y :

$$\frac{\partial}{\partial y}(x^2y + yz + e^z) = \frac{\partial}{\partial y}(10).$$

Applying the chain rule:

$$\frac{\partial}{\partial y}(x^2y) + \frac{\partial}{\partial y}(yz) + \frac{\partial}{\partial y}(e^z) = 0.$$

Compute each term:

$$\frac{\partial}{\partial y}(x^2y) = x^2, \quad \frac{\partial}{\partial y}(yz) = z + y \frac{\partial z}{\partial y}, \quad \frac{\partial}{\partial y}(e^z) = e^z \frac{\partial z}{\partial y}.$$

Combine the terms:

$$x^2 + z + (y + e^z) \frac{\partial z}{\partial y} = 0.$$

Group terms involving $\frac{\partial z}{\partial y}$:

$$(y + e^z) \frac{\partial z}{\partial y} = -x^2 - z.$$

Solve for $\frac{\partial z}{\partial y}$:

$$\frac{\partial z}{\partial y} = \frac{-x^2 - z}{y + e^z}, \quad \text{provided } y + e^z \neq 0.$$

Total Differentials

The concept of total differentials arises naturally when we want to understand how a multivariable function changes in response to small changes in its inputs. For a single-variable function $y = f(x)$, the differential $dy = f'(x)dx$ provides a linear approximation to the change in y . Similarly, for a multivariable function $w = f(x_1, x_2, \dots, x_n)$, the total differential dw captures the combined effect of infinitesimal changes in all input variables:

$$dw = \sum_{i=1}^n \frac{\partial f}{\partial x_i} dx_i.$$

Why focus on dw ? The total differential is not just a mathematical curiosity—it is a powerful tool for estimating the change in w based on the rates of change ($\frac{\partial f}{\partial x_i}$) and the input variations (dx_i). It provides a linear approximation that simplifies analysis in scenarios where changes in variables are small but interdependent. For example:

- In **error propagation**, dw quantifies how uncertainties in measurements of $x_1, x_2, \dots, x_n$ affect the overall uncertainty in w .
- In **optimization**, dw aids in understanding the sensitivity of a function to changes in its variables, guiding decisions about which factors matter most.
- In **differential geometry**, dw generalizes to describe changes on curved surfaces and manifolds, forming the backbone of advanced calculus.

By combining the effects of individual variables into one expression, the total differential serves as a concise and unified way to understand the behavior of multivariable functions in the context of infinitesimal changes.

▲ **Example 3.4.4** The temperature T at a point in space is given by $T(x, y, z) = 100 - 2x^2 - y^2 - z^2$, where x, y, z are measured in meters. Suppose a sensor moves from the point $(1, 2, -1)$ to a nearby point $(1.02, 1.98, -1.01)$. Use the total differential to estimate the change in temperature ΔT .

Solution First, compute the partial derivatives of $T(x, y, z)$:

$$\frac{\partial T}{\partial x} = -4x, \quad \frac{\partial T}{\partial y} = -2y, \quad \frac{\partial T}{\partial z} = -2z.$$

At the point $(1, 2, -1)$, evaluate these partial derivatives:

$$\frac{\partial T}{\partial x} = -4(1) = -4, \quad \frac{\partial T}{\partial y} = -2(2) = -4, \quad \frac{\partial T}{\partial z} = -2(-1) = 2.$$

The total differential dT is given by:

$$dT = \frac{\partial T}{\partial x}dx + \frac{\partial T}{\partial y}dy + \frac{\partial T}{\partial z}dz.$$

Substitute $dx = 1.02 - 1 = 0.02$, $dy = 1.98 - 2 = -0.02$, and $dz = -1.01 - (-1) = -0.01$:

$$dT = (-4)(0.02) + (-4)(-0.02) + (2)(-0.01).$$

Simplify:

$$dT = -0.08 + 0.08 - 0.02 = -0.02.$$

Thus, the estimated change in temperature is:

$$\Delta T \approx -0.02^\circ\text{C}.$$

Leibniz Integral Rule

The Leibniz integral rule extends the utility of the Chain Rule to integrals where the limits of integration or the integrand itself depend on a variable. For a function $f(x, t)$, consider the integral:

$$F(x) = \int_{a(x)}^{b(x)} f(x, t) dt.$$

The derivative of $F(x)$ with respect to x is given by:

$$\frac{dF}{dx} = f(x, b(x)) \frac{db}{dx} - f(x, a(x)) \frac{da}{dx} + \int_{a(x)}^{b(x)} \frac{\partial f}{\partial x}(x, t) dt,$$

provided that f and its partial derivative $\frac{\partial f}{\partial x}$ are continuous on the region of integration. This formula encapsulates three contributions to the rate of change of $F(x)$:

- The first term accounts for the upper limit $b(x)$ changing, weighted by the value of the integrand $f(x, t)$ at that limit.
- The second term accounts for the lower limit $a(x)$ changing, weighted by the value of $f(x, t)$ at that limit.
- The third term represents the direct contribution from the dependence of $f(x, t)$ on x , integrated over the interval.

The Leibniz rule is particularly valuable in applications such as:

- Evaluating parametric integrals where the limits or integrand depend on parameters.
- Deriving equations in physics and engineering, such as time-dependent conservation laws.
- Analyzing probability distributions with varying bounds.

This result demonstrates the interplay between differentiation and integration in multivariable contexts, leveraging the Chain Rule to account for variable dependencies within the integral.

Lecture 3.5: Directional Derivatives and Gradient Vectors

In previous lectures, we explored the foundational theories of change, including partial derivatives, total differentials, and the Chain Rule. These tools provided insight into how functions change with respect to individual variables or interconnected variables. However, real-world scenarios rarely involve changes along axes alone. Instead, we often need to understand how a function changes in any arbitrary direction in space. This lecture introduces directional derivatives and gradient vectors, which extend the ideas of partial derivatives and total differentials to arbitrary directions. These tools are critical in understanding rates of change, optimizing functions, and impressing your friends who peer over your shoulder with some of the intense notation we are about to partake.

Directional Derivatives

The directional derivative quantifies the rate of change of a function $f(x, y, z)$ in a specific direction. To motivate its introduction, consider the limitations of partial derivatives: they only describe the rate of change of a function along coordinate axes. But what if we are interested in how f changes in a diagonal direction, or along an arbitrary path? In real-world applications, such as physics or optimization, understanding how a function changes in any given direction is often critical. This leads naturally to the need for directional derivatives. Ponder for a moment: if we know how f changes along the x -, y -, and z -axes, can we combine this information to understand how f changes in an arbitrary direction? This question guides us toward the definition of the directional derivative.

Consider a function f defined in $\mathbb{R}^n$ and a unit vector $\mathbf{u}$ indicating the direction of interest. The directional derivative of f at a point $\mathbf{p} = (x_1, x_2, \dots, x_n)$ in the direction of $\mathbf{u}$ is denoted $D_{\mathbf{u}}f(\mathbf{p})$ and is defined by equipping our increment h in our usual limit definition of derivative with the direction $\mathbf{u}$:

$$D_{\mathbf{u}}f(\mathbf{p}) = \lim_{h \rightarrow 0} \frac{f(\mathbf{p} + h\mathbf{u}) - f(\mathbf{p})}{h}.$$

This definition generalizes the concept of partial derivatives. The numerator, $f(\mathbf{p} + h\mathbf{u}) - f(\mathbf{p})$, represents the change in f as we move from $\mathbf{p}$ along the direction of $\mathbf{u}$ by a small distance h . Dividing by h gives the average rate of change over this small interval, and the limit ensures that we measure the instantaneous rate of change. Let's carefully unpack this definition with an example of notation:

$$D_{\mathbf{u}}f(\mathbf{p}) = \left. \frac{d}{dh} f(\mathbf{p} + h\mathbf{u}) \right|_{h=0}.$$

Here, the notation $\mathbf{p} + h\mathbf{u}$ emphasizes that we are moving from $\mathbf{p}$ in the direction $\mathbf{u}$, scaled by a small step h . Differentiating with respect to h isolates the contribution of the directional movement along $\mathbf{u}$, allowing us to quantify how f changes specifically due to this displacement. While the definition above is mathematically rigorous, it can be challenging to compute in practice due to the dependency of f on multiple variables and the complexity

of evaluating the limit as $h \rightarrow 0$. For example, if $f(x, y) = e^{x^2+y^2}$ and $\mathbf{u} = \begin{bmatrix} \frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \end{bmatrix}$, we need to evaluate:

$$f(\mathbf{p} + h\mathbf{u}) = f\left(\begin{bmatrix} p_x + \frac{h\sqrt{2}}{2} \\ p_y + \frac{h\sqrt{2}}{2} \end{bmatrix}\right) = e^{\left(p_x + \frac{h\sqrt{2}}{2}\right)^2 + \left(p_y + \frac{h\sqrt{2}}{2}\right)^2}.$$

Expanding this expression and differentiating with respect to h involves tedious calculations that are prone to error.




Figure 3.16: A 3D visualization of the function $f(x, y) = x^2 + 3xy + y^2$. The point $\mathbf{p} = (1, 2)$ is marked on the surface, and the blue arrow represents movement in the direction of the unit vector $\mathbf{u} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, illustrating the directional derivative at $\mathbf{p}$.

The Gradient Vector

To fully understand directional derivatives, we must first introduce the gradient vector, a fundamental concept that encapsulates how a function f changes in all coordinate directions simultaneously. The gradient vector provides the foundation upon which the directional derivative is built.

The gradient of a function f , denoted ∇f , is a *vector field* that encodes the partial derivatives of f with respect to each variable. Formally, for $f : \mathbb{R}^n \rightarrow \mathbb{R}$, the gradient vector at a point $\mathbf{p} = (x_1, x_2, \dots, x_n)$ is defined as:

$$\nabla f(\mathbf{p}) = \begin{bmatrix} \frac{\partial f}{\partial x_1}(\mathbf{p}) \\ \frac{\partial f}{\partial x_2}(\mathbf{p}) \\ \vdots \\ \frac{\partial f}{\partial x_n}(\mathbf{p}) \end{bmatrix}.$$

Each component $\frac{\partial f}{\partial x_i}(\mathbf{p})$ measures how f changes as x_i varies, while all other variables are held constant. Thus, the gradient vector consolidates all partial derivatives into a single geometric entity, pointing in the direction of the steepest ascent of f . Let's knock out a quick example before we continue:

▲ **Example 3.5.1** Compute the gradient vector ∇f for the function $f(x, y, z) = x^2y + yz^3 - \sin(xz)$ at the point $\mathbf{p} = (1, 2, 0)$.

Solution The gradient vector is defined as $\nabla f(x, y, z) = \langle f_x, f_y, f_z \rangle$

Compute each partial derivative:

- $\frac{\partial f}{\partial x} = \frac{\partial}{\partial x} (x^2y + yz^3 - \sin(xz)) = 2xy - z \cos(xz).$
- $\frac{\partial f}{\partial y} = \frac{\partial}{\partial y} (x^2y + yz^3 - \sin(xz)) = x^2 + z^3.$
- $\frac{\partial f}{\partial z} = \frac{\partial}{\partial z} (x^2y + yz^3 - \sin(xz)) = 3yz^2 - x \cos(xz).$

Evaluate at $\mathbf{p} = (1, 2, 0)$:

- $\frac{\partial f}{\partial x}(1, 2, 0) = 2(1)(2) - 0 \cos(1 \cdot 0) = 4.$
- $\frac{\partial f}{\partial y}(1, 2, 0) = (1)^2 + (0)^3 = 1.$
- $\frac{\partial f}{\partial z}(1, 2, 0) = 3(2)(0)^2 - 1 \cos(1 \cdot 0) = -1.$

Thus, the gradient vector at $\mathbf{p} = (1, 2, 0)$ is:

$$\nabla f(1, 2, 0) = \begin{bmatrix} 4 \\ 1 \\ -1 \end{bmatrix}.$$

With the gradient in hand, we can rigorously define the directional derivative. The directional derivative $D_{\mathbf{u}}f(\mathbf{p})$ quantifies the rate of change of f at $\mathbf{p}$ in the direction of a unit vector $\mathbf{u}$. It is given by:

$$D_{\mathbf{u}}f(\mathbf{p}) = \nabla f(\mathbf{p}) \cdot \mathbf{u}.$$

This equivalence arises because $\nabla f(\mathbf{p})$ encapsulates all partial derivatives of f at $\mathbf{p}$, and the dot product with $\mathbf{u}$ effectively weights these partial derivatives according to the components of $\mathbf{u}$. Specifically, starting from the definition of the directional derivative:

$$D_{\mathbf{u}}f(\mathbf{p}) = \left. \frac{d}{dh} f(\mathbf{p} + h\mathbf{u}) \right|_{h=0},$$

we recognize that $\mathbf{p} + h\mathbf{u}$ represents a small displacement from $\mathbf{p}$ in the direction of $\mathbf{u}$. Differentiating $f(\mathbf{p} + h\mathbf{u})$ with respect to h involves the chain rule, yielding:

$$\left. \frac{d}{dh} f(\mathbf{p} + h\mathbf{u}) \right|_{h=0} = \nabla f(\mathbf{p}) \cdot \mathbf{u}.$$

Here, the gradient $\nabla f(\mathbf{p})$ acts as a vector of sensitivities, and the dot product projects it onto the direction $\mathbf{u}$, isolating the rate of change along that direction.

To summarize the formula $D_{\mathbf{u}}f(\mathbf{p}) = \nabla f(\mathbf{p}) \cdot \mathbf{u}$:

- The gradient $\nabla f(\mathbf{p})$ encapsulates how f changes in all coordinate directions.
- The dot product $\nabla f(\mathbf{p}) \cdot \mathbf{u}$ projects the gradient onto the direction $\mathbf{u}$, isolating the rate of change along that direction.

Rewriting the directional derivative in expanded form reveals its structure explicitly:

$$D_{\mathbf{u}}f(\mathbf{p}) = \nabla f(\mathbf{p}) \cdot \mathbf{u} = \frac{\partial f}{\partial x_1} u_1 + \frac{\partial f}{\partial x_2} u_2 + \cdots + \frac{\partial f}{\partial x_n} u_n = \sum_i f_{x_i} u_i,$$

where $\mathbf{u} = [u_1, u_2, \dots, u_n]^T$. This expression shows that the directional derivative is a weighted sum of the partial derivatives, with the weights provided by the components of $\mathbf{u}$.

▲ **Example 3.5.2** Let $f(x, y) = x^2 + 3xy + y^2$, and consider the point $\mathbf{p} = (1, 2)$ and the unit vector $\mathbf{u} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}$. Verify that the directional derivative $D_{\mathbf{u}}f(\mathbf{p})$ computed using:

$$D_{\mathbf{u}}f(\mathbf{p}) = \nabla f(\mathbf{p}) \cdot \mathbf{u},$$

is the same as:

$$\left. \frac{d}{dh} f(\mathbf{p} + h\mathbf{u}) \right|_{h=0}.$$

Solution

Step 1: Compute $D_{\mathbf{u}}f(\mathbf{p})$ using the gradient and dot product.

- The gradient of $f(x, y)$ is:

$$\nabla f(x, y) = \begin{bmatrix} 2x + 3y \\ 3x + 2y \end{bmatrix}.$$

At $\mathbf{p} = (1, 2)$:

$$\nabla f(1, 2) = \begin{bmatrix} 2(1) + 3(2) \\ 3(1) + 2(2) \end{bmatrix} = \begin{bmatrix} 8 \\ 7 \end{bmatrix}.$$

- Compute the dot product:

$$D_{\mathbf{u}}f(\mathbf{p}) = \nabla f(\mathbf{p}) \cdot \mathbf{u} = \begin{bmatrix} 8 \\ 7 \end{bmatrix} \cdot \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \frac{8+7}{\sqrt{2}} = \frac{15}{\sqrt{2}}.$$

Step 2: Compute $D_{\mathbf{u}}f(\mathbf{p})$ using the limit definition.

- Substitute $\mathbf{p} + h\mathbf{u}$ into $f(x, y)$:

$$f(\mathbf{p} + h\mathbf{u}) = f\left(1 + \frac{h}{\sqrt{2}}, 2 + \frac{h}{\sqrt{2}}\right).$$

Expand $f(x, y) = x^2 + 3xy + y^2$:

$$f\left(1 + \frac{h}{\sqrt{2}}, 2 + \frac{h}{\sqrt{2}}\right) = \left(1 + \frac{h}{\sqrt{2}}\right)^2 + 3\left(1 + \frac{h}{\sqrt{2}}\right)\left(2 + \frac{h}{\sqrt{2}}\right) + \left(2 + \frac{h}{\sqrt{2}}\right)^2.$$

Combine terms (constant, linear, and quadratic in h):

$$f(\mathbf{p} + h\mathbf{u}) = 11 + \frac{15h}{\sqrt{2}} + \frac{5h^2}{2}.$$

- Compute the derivative with respect to h and evaluate at $h = 0$:

$$\frac{d}{dh}f(\mathbf{p} + h\mathbf{u}) = \frac{15}{\sqrt{2}} + 5h.$$

At $h = 0$:

$$\left.\frac{d}{dh}f(\mathbf{p} + h\mathbf{u})\right|_{h=0} = \frac{15}{\sqrt{2}}.$$

Thus, both methods yield $D_{\mathbf{u}}f(\mathbf{p}) = \frac{15}{\sqrt{2}}$, confirming the equivalence.

A natural question arises: why must $\mathbf{u}$ be a unit vector? The unit length ensures that $D_{\mathbf{u}}f(\mathbf{p})$ measures the rate of change per unit distance traveled along $\mathbf{u}$. If $\mathbf{u}$ is not a unit vector, the directional derivative would also reflect the scaling of $\mathbf{u}$, distorting its geometric interpretation. The gradient vector not only simplifies computations but also provides

profound geometric insights. Key properties of the gradient include:

- **Direction of Maximum Increase:** The gradient vector points in the direction where f increases most rapidly. This follows from the dot product $\nabla f(\mathbf{p}) \cdot \mathbf{u}$, which achieves its maximum value when $\mathbf{u}$ is aligned with $\nabla f(\mathbf{p})$.
- **Magnitude as Rate of Change:** The magnitude $|\nabla f(\mathbf{p})|$ represents the maximum rate of change of f at $\mathbf{p}$.
- **Orthogonality to Level Surfaces:** The gradient is perpendicular to level curves (in two dimensions) or level surfaces (in three dimensions). For example, if $f(x, y, z) = c$ defines a level surface, then ∇f satisfies:

$$\nabla f \cdot \mathbf{v} = 0,$$

where $\mathbf{v}$ is any tangent vector to the level surface.

Now that we have established the gradient, we can revisit the directional derivative from a conceptual standpoint. The gradient encapsulates all partial derivatives and provides the tools to compute how f changes along any direction. The directional derivative extends the idea of partial derivatives by generalizing the rate of change from axis-aligned directions to arbitrary directions. This interplay between the gradient and directional derivatives reflects the power and elegance of multivariable calculus, connecting geometry, algebra, and analysis.

The utility of these concepts will become increasingly evident as we apply them to rate-of-change problems, optimization, and the study of vector fields. The gradient, in particular, acts as a bridge between local behavior (how f changes at a point) and global phenomena (how f behaves across its entire domain).

Applications to Rate of Change Problems

The concepts of directional derivatives and gradients extend naturally to solving real-world problems involving rates of change. Consider the following scenarios:

- **Temperature Distribution:** In a three-dimensional temperature field $T(x, y, z)$, the gradient ∇T indicates the direction in which temperature increases most rapidly. The magnitude of ∇T gives the rate of temperature change in that direction.
- **Flow Problems:** In fluid dynamics, the gradient of a scalar field such as pressure or density can describe the force acting on a fluid particle, guiding its motion.
- **Optimization:** By identifying points where $\nabla f = 0$, we can find critical points that are potential extrema or saddle points of the function f . The directional derivative allows us to analyze changes in any desired direction, facilitating a deeper understanding of the function's behavior.

Gradient Ascent/Descent

Gradient ascent/descent is a fundamental technique used to navigate scalar fields and locate points of maximum or minimum value by following the direction of steepest ascent or descent. At each point on the scalar field $f(x, y)$, the gradient vector ∇f provides the direction in which f increases most rapidly. The ascent/descent process is iterative, with each step determined by moving along the gradient direction to transition from one level curve to the next. This section explains how to compute the step vectors $\mathbf{r}$, which guide this process using the unit vector method.

1. **Understand the Role of the Gradient:** The gradient vector $\nabla f(x, y) = \langle f_x, f_y, f_z \rangle$ points in the direction of maximum increase of the function $f(x, y)$. Its magnitude $|\nabla f|$ reflects the steepness of the ascent at a point.
2. **Define a Step Vector $\mathbf{r}$ Using a Unit Vector:** To ensure consistent step sizes regardless of the gradient's magnitude, define the step vector using a unit vector in the desired direction (ascent or descent):

$$\hat{\mathbf{u}} = \frac{\nabla f(x, y)}{|\nabla f(x, y)|}.$$

This unit vector $\hat{\mathbf{u}}$ maintains the direction of steepest ascent while standardizing the step's magnitude. The step vector $\mathbf{r}$ is then given by:

$$\mathbf{r} = \Delta t \cdot \hat{\mathbf{u}},$$

where Δt is a step size parameter that controls the distance moved in the gradient direction.

3. **Orienting the Gradient Vector:** Depending on whether you aim to perform gradient ascent or gradient descent, orient the gradient vector appropriately by adjusting its sign:
 - **Gradient Ascent:** Use the gradient vector ∇f directly to ascend towards higher function values.
 - **Gradient Descent:** Use the negative gradient vector $-\nabla f$ to descend towards lower function values.

By orienting the gradient vector in the desired direction, the step vector $\mathbf{r}$ guides the ascent or descent process accordingly.

4. **Update the Point of Interest:** Starting from a current point $\mathbf{P}_n = (x_n, y_n)$, compute the next point $\mathbf{P}_{n+1}$ using:

$$\mathbf{P}_{n+1} = \mathbf{P}_n + \mathbf{r}.$$

This equation updates the position by adding the step vector $\mathbf{r}$ to the current coordinates of $\mathbf{P}_n$.

5. **Iterate the Ascent/Descent:** At $\mathbf{P}_{n+1}$, recompute the gradient $\nabla f(\mathbf{P}_{n+1})$ and determine the next step vector $\mathbf{r}$. Repeat the process to continue moving upward (ascent) or downward (descent) along the gradient direction. The iterations trace a path of steepest ascent or descent on the scalar field.
6. **Connecting to Level Curves:** Each point along the ascent/descent path corresponds to a different level curve, defined by $f(x, y) = c$. The ascent/descent process transitions from one level curve to the next, with the gradient at each point providing the direction of movement. The step vector $\mathbf{r}$ guides this transition, and its length is determined by the step size Δt .
7. **Geometric Interpretation:** The ascent/descent path is a curve that moves orthogonal to level curves, following the direction of maximum increase or decrease at each step. The vectors $\mathbf{r}$ represent the incremental displacements along this path, combining direction (from ∇f or $-\nabla f$) and magnitude (scaled by Δt).
8. **Practical Considerations:** While Δt determines the step size, its choice affects both the resolution and speed of the ascent/descent. A small Δt provides finer control and accuracy, while a larger Δt speeds up the process but may overshoot critical points or reduce precision.

Gradient ascent/descent leverages the mathematical properties of the gradient vector to navigate scalar fields systematically. The $\mathbf{r}$ vectors embody this process by encoding the direction and magnitude of each step. By iteratively computing $\mathbf{r}$, we construct a path of steepest ascent or descent that traverses successive level curves and approaches points of maximum or minimum value on $f(x, y)$.




Figure 3.17: An example of gradient ascent on a set of level curves, where $\hat{\mathbf{r}}_i$ indicates the normal vector between level curves.

▲ **Example 3.5.3** Consider the level curves of the function $f(x, y) = 4 - x^2 - y^2$, representing concentric circles centered at the origin. Starting at the point $\mathbf{P}_1 = (0.5, 0.5)$, compute the unit vectors $\hat{\mathbf{r}}_1$ and $\hat{\mathbf{r}}_2$, which point in the direction of steepest descent toward the center of the level curves as we move to the next two inner level curves.

Solution The gradient vector $\nabla f(x, y)$ points radially outward, so the negative gradient $-\nabla f(x, y)$ points inward, toward the center of the level curves. Let's compute $\hat{\mathbf{r}}_1$ and $\hat{\mathbf{r}}_2$ step by step.

1. **Compute the gradient at $\mathbf{P}_1 = (0.5, 0.5)$:**

$$\nabla f(x, y) = \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \end{bmatrix} = \begin{bmatrix} -2x \\ -2y \end{bmatrix}.$$

At $\mathbf{P}_1 = (0.5, 0.5)$:

$$\nabla f(0.5, 0.5) = \begin{bmatrix} -1.0 \\ -1.6 \end{bmatrix}.$$

The negative gradient vector is:

$$-\nabla f(0.5, 0.5) = \begin{bmatrix} 1.0 \\ 1.6 \end{bmatrix}.$$

Compute its magnitude:

$$|-\nabla f(0.5, 0.5)| = \sqrt{(1.0)^2 + (1.6)^2} = \sqrt{1 + 2.56} = \sqrt{3.56} \approx 1.886.$$

The unit vector pointing toward the center is:

$$\hat{\mathbf{r}}_1 = \frac{-\nabla f(0.5, 0.5)}{|-\nabla f(0.5, 0.5)|} = \frac{1}{1.886} \begin{bmatrix} 1.0 \\ 1.6 \end{bmatrix} \approx \begin{bmatrix} 0.53 \\ 0.85 \end{bmatrix}.$$

(Note: For simplicity in plotting, we've approximated $\hat{\mathbf{r}}_1$ to $(-0.6, -0.8)$, maintaining the direction.)

2. **Move to the next points:** Starting from $\mathbf{P}_1 = (0.5, 0.5)$, we move two steps along the direction of $\hat{\mathbf{r}}_1$ and $\hat{\mathbf{r}}_2$.

-First Step ($\mathbf{P}_1$ to $\mathbf{P}_2$):

$$\mathbf{P}_2 = \mathbf{P}_1 + h\hat{\mathbf{r}}_1 = (0.5, 0.5) + 0.3 \cdot (-0.6, -0.8) = (0.5 - 0.18, 0.5 - 0.24) = (0.32, 0.26)$$

$$z_2 = 4 - (0.32)^2 - (0.26)^2 = 4 - 0.1024 - 0.0676 = 3.83$$

Thus, $\mathbf{P}_2 = (0.32, 0.26, 3.83)$.

-Second Step ($\mathbf{P}_2$ to $\mathbf{P}_3$):

$$\mathbf{P}_3 = \mathbf{P}_2 + h\hat{\mathbf{r}}_2 = (0.32, 0.26) + 0.3 \cdot (-0.6, -0.8) = (0.32 - 0.18, 0.26 - 0.24) = (0.14, 0.02)$$

$$z_3 = 4 - (0.14)^2 - (0.02)^2 = 4 - 0.0196 - 0.0004 = 3.98$$

Thus, $\mathbf{P}_3 = (0.14, 0.02, 3.98)$.

3. **Interpretation:** Both $\hat{\mathbf{r}}_1$ and $\hat{\mathbf{r}}_2$ are unit vectors pointing inward, consistent with the symmetry of the concentric level curves. These vectors demonstrate the direction of steepest descent at each step as we move closer to the center of the level curves.

Gradient Descent: Local Surface and Gradient Vectors




Figure 3.18: The gradient ascent of the two vectors found in this problem starting at point $\mathbf{P}_1 = (0.5, 0.5)$.

▲ **Example 3.5.4** A room has a temperature distribution given by $T(x, y, z) = 100 - x^2 - y^2 - z^2$, where T is measured in degrees Celsius, and x, y, z are measured in meters. At the point $\mathbf{p} = (1, 1, 1)$, find:

1. The gradient of T .
2. The direction of the steepest ascent of the temperature.
3. The rate of change of the temperature in the direction of the steepest ascent.

Solution The gradient vector is defined as $\nabla T(x, y, z) = \langle T_x, T_y, T_z \rangle$.

1. Compute each partial derivative:

- $\frac{\partial T}{\partial x} = \frac{\partial}{\partial x} (100 - x^2 - y^2 - z^2) = -2x.$
- $\frac{\partial T}{\partial y} = \frac{\partial}{\partial y} (100 - x^2 - y^2 - z^2) = -2y.$
- $\frac{\partial T}{\partial z} = \frac{\partial}{\partial z} (100 - x^2 - y^2 - z^2) = -2z.$

At $\mathbf{p} = (1, 1, 1)$, the gradient is:

$$\nabla T(1, 1, 1) = \begin{bmatrix} -2(1) \\ -2(1) \\ -2(1) \end{bmatrix} = \begin{bmatrix} -2 \\ -2 \\ -2 \end{bmatrix}.$$

2. The direction of the steepest ascent is given by the unit vector in the direction of $\nabla T(1, 1, 1)$. Compute the magnitude of $\nabla T(1, 1, 1)$:

$$|\nabla T(1, 1, 1)| = \sqrt{(-2)^2 + (-2)^2 + (-2)^2} = \sqrt{12} = 2\sqrt{3}.$$

The unit vector is:

$$\mathbf{u} = \frac{\nabla T(1, 1, 1)}{|\nabla T(1, 1, 1)|} = \frac{1}{2\sqrt{3}} \begin{bmatrix} -2 \\ -2 \\ -2 \end{bmatrix} = \begin{bmatrix} -\frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{3}} \end{bmatrix}.$$

3. The rate of change of the temperature in the direction of the steepest ascent is the magnitude of the gradient:

$$|\nabla T(1, 1, 1)| = 2\sqrt{3}.$$

Thus, the steepest ascent occurs in the direction of $\mathbf{u}$, with a rate of change of $2\sqrt{3}$ degrees Celsius per meter.

Gradient Descent in Machine Learning

Gradient descent is a cornerstone optimization algorithm widely used in machine learning to minimize loss functions, thereby improving model accuracy. At its core, gradient descent iteratively adjusts the model's parameters to find the set of values that result in the lowest possible error between the predicted outputs and the actual data. In machine learning, the objective is often to minimize a loss function (also known as a cost function), which quantifies the difference between the model's predictions and the true values. Common examples include Mean Squared Error (MSE) for regression tasks and Cross-Entropy Loss for classification tasks.

Starting with an initial set of parameters θ_0 (which can be chosen randomly or based on some heuristic), gradient descent proceeds by computing the gradient of the loss function

with respect to each parameter. This gradient indicates the direction of the steepest increase in the loss. To minimize the loss, the algorithm updates each parameter in the opposite direction of the gradient using a unit vector approach:

$$\hat{\mathbf{u}} = \frac{\nabla_{\theta} L(\theta)}{|\nabla_{\theta} L(\theta)|}$$

$$\theta_{\text{new}} = \theta_{\text{old}} - \alpha \cdot \hat{\mathbf{u}}$$

where:

- θ represents the model parameters.
- α is the **learning rate**, a hyperparameter that controls the size of the update steps.
- $\nabla_{\theta} L(\theta)$ is the gradient of the loss function with respect to the parameters.
- $\hat{\mathbf{u}}$ is the **unit vector** in the direction of the gradient, ensuring consistent step sizes.

The process repeats—computing gradients and updating parameters—until the algorithm converges to a minimum of the loss function. Convergence is typically achieved when changes in the loss function become negligible or after a predefined number of iterations. There are several variants of gradient descent tailored to different scenarios, including Batch Gradient Descent, Stochastic Gradient Descent (SGD), and Mini-Batch Gradient Descent.

Gradient descent is favored in machine learning due to its simplicity and efficiency in handling large-scale problems. It scales well with data size and model complexity, making it suitable for training deep neural networks and other complex models. Selecting an appropriate learning rate α is crucial; a learning rate that is too high can cause the algorithm to overshoot minima, while a rate that is too low can result in slow convergence. Techniques like learning rate scheduling and adaptive learning rate algorithms (e.g., Adam, RMSprop) are often employed to address these challenges.

Gradient descent leverages the mathematical properties of the gradient vector to navigate loss landscapes systematically. The step vectors $\mathbf{r}$ embody this process by encoding the direction and magnitude of each update. By iteratively computing $\mathbf{r}$, we construct a path of steepest descent that traverses successive level curves and approaches points of minimum loss on $f(\theta)$.

The example Python code below minimizes a function, in this case $f(x) = x^2$, from a starting value with a learning rate. Obviously, the minimum of this function is 0, and we can see that the algorithm is learning its way down to reach that value.

```

1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 def gradient_descent(func, grad_func, initial_x, learning_rate, num_iterations):
5     x = initial_x
6     x_history = [x]
```

```

7
8     for i in range(num_iterations):
9         grad = grad_func(x)          # Compute the gradient at current x
10        x = x - learning_rate * grad # Update x in the opposite direction of the
    gradient
11        x_history.append(x)           # Record the new x value
12
13        # Optional: Print progress every 10 iterations
14        if (i+1) % 10 == 0:
15            print(f"Iteration {i+1}: x = {x:.4f}, f(x) = {func(x):.4f}")
16
17    return x_history
18
19 # Example usage with the function f(x) = x^2
20 def f(x):
21     return x**2
22
23 def grad_f(x):
24     return 2*x
25
26 # Parameters for gradient descent
27 initial_x = 10          # Starting point
28 learning_rate = 0.1     # Step size
29 num_iterations = 50     # Number of iterations
30
31 # Perform gradient descent
32 x_history = gradient_descent(f, grad_f, initial_x, learning_rate, num_iterations)
33
34 # Plotting the optimization path
35 iterations = range(len(x_history))
36 function_values = [f(x) for x in x_history]
37
38 plt.figure(figsize=(10, 6))
39 plt.plot(iterations, function_values, marker='o', linestyle='-', color='b')
40 plt.title('Gradient Descent Optimization Path')
41 plt.xlabel('Iteration')
42 plt.ylabel('f(x) = x^2')
43 plt.grid(True)
44 plt.show()

```

The graph below shows the journey of the gradient descent minimizing $f(x)$.




Figure 3.19: Gradient descent for minimizing the function $f(x) = x^2$

Lecture 3.6: Tangent Planes and Linear Approximations

In our exploration of multivariable functions, we have examined how these functions change with respect to their independent variables and how to capture this change using partial derivatives. A central concept in single-variable calculus is the linear approximation of a function near a point, often expressed as the tangent line approximation. In multiple dimensions, the analogous idea is the *tangent plane*, which provides a two-dimensional linear approximation to a surface near a given point. In this lecture, we extend our intuition from single-variable calculus to functions of several variables, focusing on tangent planes and linear approximations. We will also connect these ideas to differentials, which enable us to approximate small changes in a function and apply these approximations to estimate errors in measurements and computations.

From Tangent Lines to Tangent Planes

Recall from Single-Variable Calculus: you have a function $y = f(x)$ that is differentiable at a point $x = a$. If we zoom in very close to the graph of f at $(a, f(a))$, the curve appears almost indistinguishable from a straight line. This line is the *tangent line*, and it gives the best linear approximation to f near $x = a$. How do we find this tangent line? We know the slope of the tangent line is given by the derivative $f'(a)$. The point of tangency is $(a, f(a))$. Combining this information, the equation of the tangent line is:

$$y \approx f(a) + f'(a)(x - a).$$

This linear function approximates $f(x)$ well for x near a . In other words, we have a local, linear predictor:

$$f(x) \approx f(a) + f'(a)(x - a).$$

This approximation is not just a convenient guess; it emerges naturally from the definition of the derivative:

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a + h) - f(a)}{h}.$$

If we rearrange this definition for small h :

$$f(a + h) \approx f(a) + f'(a)h,$$

which is exactly our linear approximation. As $h \rightarrow 0$, this approximation becomes increasingly accurate.

Now, consider a function of two variables:

$$z = f(x, y).$$

We want an analogous concept to the tangent line, but now our domain is two-dimensional, and the graph of f is a surface in three-dimensional space. Instead of a line, the natural linear approximation to a surface at a given point is a *tangent plane*.

Suppose f is differentiable at a point (a, b) . Differentiability in two dimensions means that as we move from (a, b) to $(a + \Delta x, b + \Delta y)$, the change in f can be approximated by a linear function of Δx and Δy . Let's express this idea.

In single-variable calculus, differentiability required that:

$$f(a + h) = f(a) + f'(a)h + \epsilon(h), \quad \text{where } \epsilon(h) \rightarrow 0 \text{ as } h \rightarrow 0.$$

In two-variable calculus, if f is differentiable at (a, b) , we have:

$$f(a + \Delta x, b + \Delta y) = f(a, b) + f_x(a, b)\Delta x + f_y(a, b)\Delta y + E(\Delta x, \Delta y),$$

where $f_x(a, b)$ and $f_y(a, b)$ are the partial derivatives of f at (a, b) . Here, Δx and Δy are small increments in x and y (often denoted by h and k in limit definitions). The term $E(\Delta x, \Delta y)$ is an error term that goes to zero faster than $\sqrt{(\Delta x)^2 + (\Delta y)^2}$ as $\Delta x, \Delta y \rightarrow 0$. More precisely:

$$\lim_{(h,k) \rightarrow (0,0)} \frac{E(h,k)}{\sqrt{h^2 + k^2}} = 0.$$

Thus, as $(\Delta x, \Delta y)$ become infinitesimally small, the linear part dominates and the higher-order terms become negligible. This stems from the definition of differentiability, which ensures that the function's behavior is well-approximated by a linear map at the point of tangency.




Figure 3.20: *Left: A global view of $z = \sin(x) \cos(y)$ with a dotted region around $(0, 0)$. Right: A zoomed-in view of the same surface near $(0, 0)$, where it appears nearly flat.*

Focusing only on the linear part gives:

$$f(a + \Delta x, b + \Delta y) = f(a, b) + f_x(a, b)\Delta x + f_y(a, b)\Delta y.$$

Setting $x = a + \Delta x$ and $y = b + \Delta y$, we rewrite this as:

$$f(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b).$$

This defines a plane:

$$z = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b).$$

Let $z_0 = f(a, b)$. Then we can rewrite this as:

$$z - z_0 = f_x(a, b)(x - a) + f_y(a, b)(y - b),$$

which matches the standard form of a plane equation: a constant point (a, b, z_0) plus linear terms in $(x - a)$ and $(y - b)$. Here, the “slopes” $f_x(a, b)$ and $f_y(a, b)$ serve as the coefficients that determine the plane’s tilt in the x - and y -directions. In other words, this **tangent plane** at $(a, b, f(a, b))$ just grazes the surface at that point and provides the best linear approximation, making the surface and the plane virtually indistinguishable when we are very close to (a, b) . Below we have an example of a plane tangent to a surface:




Figure 3.21: *Tangent plane to $z = x^2 + y^2$ at the point $(1, 1, 2)$.*

▲ **Example 3.6.1** Find the equation of the tangent plane to the surface

$$z = f(x, y) = x^2 - 3xy + 4y^2$$

at the point $(2, 1)$.

Solution First, we need the partial derivatives of f with respect to x and y :

$$f_x(x, y) = 2x - 3y, \quad f_y(x, y) = -3x + 8y.$$

Evaluate these at $(2, 1)$:

$$f_x(2, 1) = 2(2) - 3(1) = 4 - 3 = 1, \quad f_y(2, 1) = -3(2) + 8(1) = -6 + 8 = 2.$$

Also, find $f(2, 1)$:

$$f(2, 1) = (2)^2 - 3(2)(1) + 4(1)^2 = 4 - 6 + 4 = 2.$$

The tangent plane is given by:

$$z = f(2, 1) + f_x(2, 1)(x - 2) + f_y(2, 1)(y - 1).$$

Substitute the values:

$$z = 2 + 1(x - 2) + 2(y - 1).$$

Simplify:

$$z = 2 + (x - 2) + 2y - 2 = x + 2y - 2.$$

Thus, the equation of the tangent plane is:

$$z = x + 2y - 2.$$

It doesn't get much harder than that!

Generalization to Higher Dimensions and Hyperplanes

The concept of tangent planes naturally extends to functions of more than two variables. For a function

$$f : \mathbb{R}^n \rightarrow \mathbb{R},$$

differentiability at a point $\mathbf{a} = (a_1, a_2, \dots, a_n)$ means that we can approximate f near $\mathbf{a}$ by a linear function involving the partial derivatives with respect to each of the n variables. Specifically, we have:

$$f(\mathbf{a} + \Delta \mathbf{x}) = f(\mathbf{a}) + \nabla f(\mathbf{a}) \cdot \Delta \mathbf{x} + E(\Delta \mathbf{x}),$$

where $\nabla f(\mathbf{a})$ is the gradient vector of f at $\mathbf{a}$, and $\Delta \mathbf{x}$ is a small displacement vector in $\mathbb{R}^n$. Ignoring the small error term, we obtain a linear approximation:

$$f(\mathbf{a} + \Delta \mathbf{x}) \approx f(\mathbf{a}) + \nabla f(\mathbf{a}) \cdot \Delta \mathbf{x}.$$

The set of points $(\mathbf{x}, z)$ in $\mathbb{R}^{n+1}$ satisfying

$$z = f(\mathbf{a}) + \nabla f(\mathbf{a}) \cdot (\mathbf{x} - \mathbf{a})$$

defines a *tangent hyperplane* to the graph of f at $\mathbf{a}$. For $n = 1$, this hyperplane is just a line (the tangent line). For $n = 2$, it's a plane (the tangent plane). For $n > 2$, the object is an $(n - 1)$ -dimensional hyperplane—a direct generalization of the idea of “flat” linear approximations to higher dimensions.

Hyperplanes in Machine Learning:

In machine learning, hyperplanes appear in various contexts, notably in classification tasks. For instance, linear classifiers such as Support Vector Machines (SVMs) or simple linear regression models rely on hyperplanes to separate data points or fit data. Consider a binary classification problem in $\mathbb{R}^n$. A linear classifier tries to find a hyperplane:

$$\mathbf{w} \cdot \mathbf{x} + b = 0,$$

that best separates data points of different classes. Here, $\mathbf{w}$ is a weight vector (analogous to the gradient vector), and b is a bias term (analogous to a constant offset). The hyperplane defined by this equation divides $\mathbb{R}^n$ into two half-spaces, and the classifier assigns points on one side to one class and points on the other side to the opposite class.

The connection between the tangent plane (or hyperplane) in calculus and the hyperplanes in machine learning is conceptually significant. Both represent the idea of using linear structures to approximate or make decisions about more complex, nonlinear situations. In calculus, the tangent hyperplane locally approximates a nonlinear function. In machine learning, a hyperplane provides a simple linear boundary in a potentially high-dimensional space. Although the goals differ—approximation of functions vs. classification of points—the underlying mathematical object (a linear function defining a hyperplane) is fundamentally the same.

A Simple Linear Classifier in 2D

Consider a simple binary classification task in two dimensions. Suppose each data point is represented by a vector $\mathbf{x} = (x_1, x_2)$ in $\mathbb{R}^2$, and we want to classify it into one of two classes, say “A” or “B”. A very basic linear classifier can be written as:

$$\mathbf{w} \cdot \mathbf{x} + b = 0,$$

where $\mathbf{w} = (w_1, w_2)$ is a weight vector and b is a bias term.

This equation defines a line in the x_1x_2 -plane, which is a 1-dimensional hyperplane (since a line is a hyperplane in $\mathbb{R}^2$). The classifier assigns points on one side of the line to class A and points on the other side to class B. Specifically:

$$\mathbf{w} \cdot \mathbf{x} + b > 0 \implies \text{Class A}, \quad \mathbf{w} \cdot \mathbf{x} + b < 0 \implies \text{Class B}.$$

For example, let's choose $\mathbf{w} = (1, -2)$ and $b = 1$. The decision boundary is:

$$1 \cdot x_1 + (-2) \cdot x_2 + 1 = 0 \implies x_1 - 2x_2 + 1 = 0.$$

Rearranging:

$$x_1 = 2x_2 - 1.$$

This line divides the plane into two regions:

$$x_1 > 2x_2 - 1 \implies \mathbf{x} \text{ is in Class A}, \quad x_1 < 2x_2 - 1 \implies \mathbf{x} \text{ is in Class B}.$$




Figure 3.22: A simple linear classifier in $\mathbb{R}^2$. The line $x_1 = 2x_2 - 1$ separates the plane into two classes. Points labeled A fall on one side, and points labeled B on the other.

Here, the concept of a hyperplane from calculus—a linear construct that approximates behavior locally—appears in a machine learning context as a global decision boundary. Although the goal is different (classifying points rather than approximating a function at a point), the mathematical form is similar: a linear equation representing a flat boundary in space. This exemplifies how linear mathematical objects like lines, planes, and hyperplanes serve as foundational building blocks across various domains, from analyzing rates of change in calculus to making decisions in machine learning.

Linear Approximations: The Bridge Between Theory and Practice

Just as we used the tangent line in one dimension to approximate function values near a point, the tangent plane in two dimensions provides a linear approximation to $f(x, y)$ near (a, b) . This approximation is extremely useful in many contexts:

- **Estimating Values:** If we know $f(a, b)$ and the partial derivatives at (a, b) , we can estimate $f(x, y)$ for (x, y) close to (a, b) without computing f directly.
- **Differentials:** We define differentials dx and dy as small increments in x and y . Then the differential df at (a, b) is:

$$df = f_x(a, b)dx + f_y(a, b)dy.$$

This quantity df represents the best linear prediction for the actual change $\Delta f = f(a + dx, b + dy) - f(a, b)$ for small changes dx and dy . It is the multivariable extension of $dy = f'(a)dx$ from single-variable calculus.

- **Error Estimation:** Suppose you measure x and y with some small possible errors Δx and Δy . Using the tangent plane approximation, you can estimate how these errors propagate into f :

$$\Delta f \approx f_x(a, b)\Delta x + f_y(a, b)\Delta y.$$

This lets you gauge how sensitive the function is to changes in its input variables.

Thus, the progression is:

$$\text{Single variable: Tangent line} \implies f(x) \approx f(a) + f'(a)(x - a).$$

$$\text{Two variables: Tangent plane} \implies f(x, y) \approx f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b).$$

This parallel structure shows that the concept of a linear approximation—so central in understanding functions in one dimension—scales naturally to higher dimensions, giving us a powerful geometric and analytic tool to approximate and understand the behavior of multivariable functions near any given point.

Beyond the Linear Approximation:

In some situations, the linear approximation may not provide sufficient information. For instance, if $\nabla f(a, b) = \mathbf{0}$ (all first-order partial derivatives vanish), then the tangent plane is flat and does not reveal how f changes for small deviations from (a, b) . In such cases, we look to higher-order approximations, much like we use second- or third-order Taylor polynomials in single-variable calculus.

In two variables, a second-order approximation (also known as the quadratic approximation) includes second partial derivatives:

$$\begin{aligned} f(x, y) \approx & f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b) \\ & + \frac{1}{2} [f_{xx}(a, b)(x - a)^2 + 2f_{xy}(a, b)(x - a)(y - b) + f_{yy}(a, b)(y - b)^2]. \end{aligned}$$

These higher-order terms capture curvature and more subtle changes in the function. They can distinguish between extrema and provide more accurate estimates when larger changes in x and y occur.

While the linear approximation (tangent plane) is often the first and most accessible tool, knowing that higher-order expansions exist broadens your ability to model and understand complex phenomena. Whether adjusting manufacturing parameters, forecasting economic indicators, or analyzing stability in physical systems, having the option to include second-order (or higher) terms allows for richer, more nuanced approximations.

▲ **Example 3.6.2** A drone mapping a field captures the elevation of the terrain using the function:

$$f(x, y) = 1000 - (x - 25)^2 - \frac{1}{2}(y - 50)^2,$$

where x and y are coordinates in meters, and $f(x, y)$ gives the elevation in meters. At the reference point $(25, 50)$, the elevation is $f(25, 50) = 1000$ meters. Suppose the drone slightly shifts its measurement to $(26, 48)$.

1. Use a first-order (linear) approximation at $(25, 50)$ to estimate the elevation at $(26, 48)$.

2. Compute the actual elevation at $(26, 48)$ and discuss why the linear approximation fails.
3. Suggest and compute a second-order approximation to improve the estimate.

Solution

1. First-Order Approximation:

The first-order approximation uses the tangent plane to estimate elevation. The general formula for the first-order (linear) approximation of a function $f(x, y)$ at a reference point (a, b) is:

$$f(x, y) \approx f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$$

For the given function:

$$f(x, y) = 1000 - (x - 25)^2 - \frac{1}{2}(y - 50)^2,$$

the partial derivatives are:

$$f_x(x, y) = -2(x - 25), \quad f_y(x, y) = -(y - 50).$$

At the reference point $(25, 50)$:

$$f_x(25, 50) = -2(25 - 25) = 0, \quad f_y(25, 50) = -(50 - 50) = 0.$$

Substituting these values into the first-order approximation formula:

$$f(x, y) \approx f(25, 50) + 0 \cdot (x - 25) + 0 \cdot (y - 50) = 1000.$$

Since both partial derivatives are zero at $(25, 50)$, the linear approximation is a constant:

$$f(x, y) \approx 1000.$$

Using this approximation, the predicted elevation at $(26, 48)$ is:

$$f(26, 48) \approx 1000.$$

2. Actual Elevation:

Compute the exact elevation at $(26, 48)$:

$$f(26, 48) = 1000 - (26 - 25)^2 - \frac{1}{2}(48 - 50)^2.$$

Simplify step by step:

$$f(26, 48) = 1000 - 1 - \frac{1}{2}(4) = 1000 - 1 - 2 = 997.$$

The actual elevation is 997 meters, but the linear approximation predicted 1000. The error occurs because the tangent plane is flat at $(25, 50)$, failing to capture the curvature of the terrain. In fact, our scheme will predict everything to be flat since $f(25, 50)$ is constant. If we are very far up the terrain when this approximation is made, it'll only get much worse the farther out we go.

3. Second-Order Approximation:

To account for curvature, we use a second-order Taylor approximation:

$$f(x, y) \approx f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b) + \frac{1}{2} [f_{xx}(a, b)(x - a)^2 + 2f_{xy}(a, b)(x - a)(y - b) + f_{yy}(a, b)(y - b)^2].$$

Compute the second derivatives:

$$f_{xx}(x, y) = -2, \quad f_{yy}(x, y) = -1, \quad f_{xy}(x, y) = 0.$$

The second-order approximation is:

$$f(x, y) \approx f(25, 50) + \frac{1}{2} [f_{xx}(25, 50)(x - 25)^2 + 2f_{xy}(25, 50)(x - 25)(y - 50) + f_{yy}(25, 50)(y - 50)^2].$$

Substituting the derivatives and evaluating at $(26, 48)$:

$$f(26, 48) \approx 1000 + \frac{1}{2} [-2(1)^2 + 0 + (-1)(-2)^2].$$

Simplify:

$$f(26, 48) \approx 1000 + \frac{1}{2} (-2 - 4) = 1000 - \frac{6}{2} = 1000 - 3 = 997.$$

The second-order approximation recovers the actual value, showing that considering curvature is crucial when first-order terms vanish.

4. Conclusion:

For flat terrain at a critical point (where the gradient is zero), higher-order terms reveal how the elevation changes. The second-order approximation includes curvature information, making it essential for accurate predictions in such cases.

Lecture 3.7: Extrema and Critical Points

One of the central themes in calculus is understanding where a function attains its greatest or least values. In single-variable calculus, this investigation led us to examine critical points—points where the derivative vanishes—and to use second-derivative tests to classify the nature of these points. These critical points allowed us to identify local maxima, local minima, and to note where the function’s concavity changes (potentially at points of inflection). Such insights gave us a clear understanding of the “shape” of a curve along a one-dimensional axis. Because plotting is more challenging in higher dimensions, any analogy or simpler scenario that helps us visualize these concepts is useful. Not only this, but for optimization we will find extrema to be a cornerstone, just as we saw in single variable calculus.

As we move to functions of several variables, the problem becomes richer and more complex. Instead of analyzing a single input-output relationship along a line, we now consider surfaces (or even higher-dimensional objects) defined by multiple input variables. The notion of “increasing” or “decreasing” can no longer be grasped by simply looking to the left or right. We must consider changes in all possible directions around a point (often referred to as examining a *neighborhood*). Consequently, the identification and classification of maxima, minima, and other special points (such as saddle points) demand more sophisticated tools and a broader conceptual framework.

Single-Variable Concepts and Their Limitations

Let’s start by recalling what we know from single-variable calculus. For a differentiable function $f : \mathbb{R} \rightarrow \mathbb{R}$:

- A **critical point** $x = a$ is where $f'(a) = 0$ or $f'(a)$ does not exist.
- At such points, the function might have a local maximum or minimum, or it might simply change its curvature, suggesting a point of inflection (though not necessarily an extremum).
- To distinguish maxima and minima from other behaviors, we use the **second derivative test**:
 - If $f''(a) > 0$, the function is concave up at a , typically indicating a local minimum.
 - If $f''(a) < 0$, the function is concave down at a , typically indicating a local maximum.
 - If $f''(a) = 0$, the test is inconclusive and the behavior around a requires further investigation.
- A **point of inflection** occurs at $x = a$ if the function changes concavity (from concave up to concave down, or vice versa) at that point. To identify such points:
 - Compute $f''(x)$ and find where $f''(x) = 0$ or $f''(x)$ does not exist.

- Check the sign of $f''(x)$ on either side of $x = a$. A change in sign confirms a point of inflection.




Figure 3.23: A single-variable function $f(x) = x^3 - 3x$ illustrating a local maximum, local minimum, and an inflection point. The critical points at $x = -1$ and $x = 1$ correspond to a local maximum and local minimum, respectively, while $x = 0$ is a point of inflection where the curvature changes.

In the figure above, we see an example function $f(x) = x^3 - 3x$ that has:

- A local maximum at $x = -1$.
- A local minimum at $x = 1$.
- An inflection point at $x = 0$, where the graph changes from concave down to concave up.

These principles form the basis of understanding how functions behave near critical points in one dimension. However, when we move to multiple variables, these ideas must be generalized. Instead of a single derivative, we have partial derivatives. Instead of a single second derivative, we have a matrix of second-order partial derivatives (the Hessian). Instead of a single direction to check for concavity changes, we must consider changes in every direction. Because Linear Algebra is not a prerequisite to Multivariate Calculus, we will take this slow, don't worry. The complexity and richness of these scenarios make the transition from one-dimensional to multivariable calculus both challenging and rewarding.

The framework above allowed us to solve a variety of one-dimensional optimization problems. However, this approach heavily relies on the fact that there's only one independent variable. Extending these ideas to multiple variables introduces several challenges:

- **Directionality:** In one dimension, you look left and right. In multiple dimensions, you must consider infinitely many directions from a point.
- **Multidimensional Curvature:** Instead of a single second derivative, we have a whole matrix of second derivatives (the Hessian).

- **Types of Critical Points:** With more directions come new types of stationary behavior. In particular, *saddle points* arise, which are not maxima or minima, but still have zero gradient.

Extrema in Several Variables

Generalizing some of the concepts to higher dimensions comes naturally enough with the notational conventions we have established in previous lectures. Consider a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$. We say that $\mathbf{a} \in \mathbb{R}^n$ is an **extremum** of f if $f(\mathbf{a})$ is a maximum or minimum value of f in some sense. As in one dimension, we distinguish between:

- **Local Maximum in $\mathbb{R}^n$:** $f(\mathbf{a})$ is greater than or equal to $f(\mathbf{x})$ for all points $\mathbf{x}$ in some “small neighborhood” around $\mathbf{a}$.
- **Local Minimum in $\mathbb{R}^n$:** $f(\mathbf{a})$ is less than or equal to $f(\mathbf{x})$ for all points $\mathbf{x}$ in a small neighborhood around $\mathbf{a}$.
- **Global Maximum/Minimum:** The value at $\mathbf{a}$ is the supreme representative—no points in the entire domain beat it (for a max) or undercut it (for a min).

Below we see a 3D plot displaying all these extrema. Because we choose a function on a region $R = [-5, 5] \times [-5, 5]$ that has no division, and thus no points in the plot that go off to $\pm\infty$, we are guaranteed global extrema due the closed and bounded nature of the conditions. We will learn about this as the Extreme Value Theorem for multivariate functions.




Figure 3.24: The surface plot of $f(x, y) = \sin(x)\sin(y)e^{x^2+y^2}$ with annotated extrema. No laughing, please.

However, unlike the single-variable case, “small neighborhood” is not just an interval around a point. In $\mathbb{R}^n$, you can approach a point from infinitely many directions. There can be “flat” regions where the function doesn’t change for a while, disconnected parts of the domain, and

directions in which the function heads off to infinity (no global extrema exist). Thus, while the idea of local and global maxima and minima transfers directly, the complexity of higher dimensions makes the identification and classification of these points more subtle.

A Rigorous Approach

Before we dive into the rigor, we need to define the concepts of **open sets**, **closed sets**, **balls**, and **compactness**. Understanding these definitions is crucial for formalizing our discussions on neighborhoods and extrema in multiple dimensions.

- **Open Set:** A set $U \subset \mathbb{R}^n$ is called *open* if, for every point $\mathbf{a} \in U$, you can move a little bit in any direction from $\mathbf{a}$ and still stay within U . In other words, there are no “edge” points included in U . This is essentially the higher dimensional analog of an open interval (a, b) .
- **Closed Set:** A set $C \subset \mathbb{R}^n$ is called *closed* if it includes all its boundary points. This means that if you have a sequence of points within C that gets closer and closer to some point $\mathbf{x}$, then $\mathbf{x}$ is also included in C . This is essentially the higher dimensional analog of a closed interval $[a, b]$.




Figure 3.25: *Illustration of an open set U with a dotted boundary and a closed set C with a solid boundary.*

- **Open Ball:** An open ball in $\mathbb{R}^n$ centered at $\mathbf{a}$ with radius ϵ is the set of all points $\mathbf{x}$ such that the distance between $\mathbf{x}$ and $\mathbf{a}$ is less than ϵ . It is defined as:

$$B_\epsilon(\mathbf{a}) = \{\mathbf{x} \in \mathbb{R}^n \mid \|\mathbf{x} - \mathbf{a}\| < \epsilon\}.$$

- **Compact Set:** A set $D \subset \mathbb{R}^n$ is called *compact* if it is both closed and bounded. This means that D contains all its boundary points and fits within some finite region of space.

To handle these complexities, we need a more precise definition of what it means to be “close” to a point in multiple dimensions. In one dimension, being close to a meant $|x - a| < \epsilon$ for some $\epsilon > 0$. In $\mathbb{R}^n$, we generalize this by using open balls defined by a norm. A standard

choice is the Euclidean norm $\|\cdot\|$, which represents the usual distance between vectors. Thus, we define:

$$B_\epsilon(\mathbf{a}) = \{\mathbf{x} \in \mathbb{R}^n \mid \|\mathbf{x} - \mathbf{a}\| < \epsilon\}.$$

This $B_\epsilon(\mathbf{a})$ is an *open ball* of radius ϵ centered at $\mathbf{a}$. It serves as the higher-dimensional analogue of a small interval around a point.

With this notation, we can restate the definitions more rigorously:

- **Local Maximum:** A point $\mathbf{a}$ is a local maximum of f if there exists $\epsilon > 0$ such that for all $\mathbf{x} \in B_\epsilon(\mathbf{a})$,

$$f(\mathbf{x}) \leq f(\mathbf{a}).$$

Here, “all $\mathbf{x}$ in $B_\epsilon(\mathbf{a})$ ” means we are checking every direction and every point sufficiently close to $\mathbf{a}$.

- **Local Minimum:** A point $\mathbf{a}$ is a local minimum of f if there exists $\epsilon > 0$ such that for all $\mathbf{x} \in B_\epsilon(\mathbf{a})$,

$$f(\mathbf{x}) \geq f(\mathbf{a}).$$

- **Global Maximum/Minimum:** These are defined similarly to the single-variable case, but we consider the entire domain. A point $\mathbf{a}$ is a global maximum if

$$f(\mathbf{x}) \leq f(\mathbf{a}) \text{ for all } \mathbf{x} \in \text{dom}(f).$$

And similarly for a global minimum.

Using these ϵ -ball/neighborhood definitions makes our concepts more robust. For instance, the notion of local extremum now doesn’t depend on the shape of the domain or the peculiarities of how we approach the point $\mathbf{a}$. Instead, it inherently encodes the idea of “closeness” and “no other nearby points exceeding (or undercutting) the value at $\mathbf{a}$.”

This refined perspective also ties in with deeper results from real analysis. For example, if f is continuous on a compact set $D \subset \mathbb{R}^n$, the Extreme Value Theorem guarantees the existence of global maxima and minima. However, if the domain is not compact, or if f is not nicely behaved, we may fail to find global extrema. The function may “escape” to infinity in some direction, preventing a global maximum or minimum from forming.

Local vs. Global Extrema

The *Extreme Value Theorem* from single-variable calculus ensures that if a continuous function is defined on a closed interval, it attains global maxima and minima. In higher dimensions, there is a corresponding result: if f is continuous on a *compact set* $D \subset \mathbb{R}^n$ (i.e., D is closed and bounded), then f attains both a global maximum and a global minimum on D . However, if the domain is not compact, the situation is more complex:

- If the domain is not bounded, f may “escape” to infinity, never settling on a highest or lowest value.

- If the domain is open or lacks a boundary, local conditions might not guarantee global extremality.

Understanding these nuances is crucial in applications. For example, in optimization problems with constraints, ensuring the constraint set is compact (often via bounding conditions) can help guarantee that a solution (a global extremum) exists.

Consider a two-dimensional function $f(x, y)$ that forms a surface in $\mathbb{R}^3$. A local maximum appears like a peak in some ϵ -ball around (a, b) ; a local minimum looks like a valley. If the surface keeps rising as $x^2 + y^2 \rightarrow \infty$ (like a giant unbounded mountain), no global maxima exist because we can always find higher points. Similarly, if it keeps going downwards without bound, no global minima exist.




Figure 3.26: A surface with a global maximum at the origin. The red dot marks the global maximum at $(0, 0, 3)$. The blue semi-transparent sphere represents a $B_\epsilon([0, 0, 3])$, illustrating a neighborhood within which no other points have a higher value. As we move away from the origin, the surface descends, ensuring that the global maximum is maintained within the $B_\epsilon([0, 0, 3])$.

In the figure above, we see a simple example: $f(x, y) = 3 - 3(x^2 + y^2)$ has a local maximum at $(0, 0)$ where $f(0, 0) = 3$. In fact, this is also a global maximum since the function decreases as we move away from $(0, 0)$, approaching negative infinity without bound. If we had a function that only rose at the origin and flattened out or grew in other directions, we might not have any global maxima or minima, just local ones.

▲ **Example 3.7.1** Consider the Extreme Value Theorem (EVT) in the context of multivariable calculus, which states that if a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuous on a *compact* set D (i.e., D is closed and bounded), then f attains both a global maximum and a global minimum on D .

1. Provide an example of a continuous function defined on an **unbounded** domain in $\mathbb{R}^2$ that does **not** attain a global maximum or a global minimum. Explain why this example illustrates the necessity of the boundedness condition in EVT.
2. Provide an example of a continuous function defined on an **open** (not closed) and **bounded** domain in $\mathbb{R}^2$ that does **not** attain a global maximum or a global minimum. Explain why this example illustrates the necessity of the closedness condition in EVT.

Solution

1. Consider the function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by

$$f(x, y) = x.$$

- **Domain Unbounded:** The domain $\mathbb{R}^2$ is unbounded, meaning it extends infinitely in all directions.
- **Behavior of $f(x, y)$:**
 - As $x \rightarrow \infty$, $f(x, y) \rightarrow \infty$.
 - As $x \rightarrow -\infty$, $f(x, y) \rightarrow -\infty$.
- **Global Extrema:**
 - Global Maximum: f does not attain a global maximum because for any value M , there exists $(x, y) \in \mathbb{R}^2$ with $x > M$.
 - Global Minimum: Similarly, f does not attain a global minimum because for any value m , there exists $(x, y) \in \mathbb{R}^2$ with $x < m$.
- **Conclusion:** This example demonstrates that when the domain is unbounded, a continuous function may fail to attain global extrema. This highlights the necessity of the boundedness condition in the Extreme Value Theorem to ensure the existence of both a global maximum and a global minimum.

2. Consider the function $g : D \rightarrow \mathbb{R}$ defined by

$$g(x, y) = \frac{1}{1 - x^2 - y^2},$$

where the domain $D = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\}$ is the open unit disk.

- **Domain Not Closed:** The domain D is open because it does not include the boundary $x^2 + y^2 = 1$. While D is bounded, it lacks closure.
- **Behavior of $g(x, y)$:**
 - As (x, y) approaches the boundary of D (i.e., as $x^2 + y^2 \rightarrow 1$), the denominator $1 - x^2 - y^2 \rightarrow 0$, causing $g(x, y) \rightarrow \infty$.
 - Within D , $g(x, y)$ is always positive and increases without bound near the boundary.
- **Global Extrema:**
 - Global Maximum: g does not attain a global maximum because for any large value M , there exists $(x, y) \in D$ sufficiently close to the boundary such that $g(x, y) > M$.
 - Global Minimum:
 - * g does attain a global minimum at the center $(0, 0)$, where $g(0, 0) = 1$.
- **Conclusion:** Although g attains a global minimum, it fails to attain a global maximum because the domain D is not closed. This example illustrates that the closedness condition is necessary in the Extreme Value Theorem to ensure that all global extrema are attained.

First Derivative Test for Extrema: Critical Points in Multiple Dimensions

In single-variable calculus, the process of finding candidates for local extrema is straightforward: we compute $f'(x)$, set it equal to zero, and solve for x . Points where $f'(x) = 0$ or $f'(x)$ is undefined are the **critical points**, and each such point is examined to determine whether it corresponds to a local maximum, minimum, or neither. This procedure generalizes naturally to functions of several variables, but with an additional layer of complexity.

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, the analogue to the single-variable derivative is the **gradient** vector:

$$\nabla f(\mathbf{x}) = \left\langle \frac{\partial f}{\partial x_1}(\mathbf{x}), \frac{\partial f}{\partial x_2}(\mathbf{x}), \dots, \frac{\partial f}{\partial x_n}(\mathbf{x}) \right\rangle.$$

The gradient encodes the rate of change of f in each coordinate direction. Setting the gradient to zero:

$$\nabla f(\mathbf{a}) = \mathbf{0}$$

defines the **critical points** or **stationary points** of f . At any such point $\mathbf{a}$, the function f has no instantaneous tendency to increase or decrease in any of the standard coordinate directions.

To find critical points in practice, one must:

1. Compute the partial derivatives $\frac{\partial f}{\partial x_i}$ for each $i = 1, 2, \dots, n$.
2. Solve the system of equations:

$$\frac{\partial f}{\partial x_1}(\mathbf{x}) = 0, \quad \frac{\partial f}{\partial x_2}(\mathbf{x}) = 0, \quad \dots, \quad \frac{\partial f}{\partial x_n}(\mathbf{x}) = 0.$$

Finding solutions may be straightforward for simple functions or can become highly nontrivial for more complicated ones, often requiring numerical methods or careful algebraic manipulation.

3. Identify all points $\mathbf{a}$ that satisfy these equations. Each such $\mathbf{a}$ is a potential extremum, but additional steps are necessary to classify its nature.

Having $\nabla f(\mathbf{a}) = \mathbf{0}$ is a necessary condition for $\mathbf{a}$ to be a local maximum or minimum, but this condition alone does not guarantee the existence of an extremum at $\mathbf{a}$. Consider the single-variable analogy: if $f'(a) = 0$, a might be a max, min, or even a point of inflection. In multiple dimensions, the situation is more nuanced:

- A zero gradient could indicate a local maximum if the function “peaks” at that point.
- It could indicate a local minimum if the function “valleys” at that point.
- It could also indicate a *saddle point*, where the function curves up in some directions and down in others, making the point neither a max nor a min.

The gradient test (setting $\nabla f(\mathbf{a}) = \mathbf{0}$) is like a first filter. It narrows the set of candidate points that could be extrema, but it does not complete the classification. To fully determine the nature of each critical point, we must invoke second-order information, typically via the Hessian matrix, and apply a second derivative test in several variables. Only after analyzing the Hessian (or, more precisely, the definiteness of the Hessian matrix at $\mathbf{a}$) can we decide whether $\mathbf{a}$ is a maximum, a minimum, or a saddle point.

Hence, the first derivative test in multiple dimensions leads us to identify critical points by solving a system of equations $\nabla f(\mathbf{a}) = \mathbf{0}$. While this can be challenging, it is a necessary step towards understanding the structure of the function’s graph, the locations of its extrema, and the detailed shape of its level surfaces.

As n grows, the complexity of setting all partial derivatives to zero increases significantly. In practice, for higher dimensions or more complicated functions, one often relies on numerical methods, optimization algorithms (like gradient descent), or specialized analytic techniques to locate critical points. The first derivative test is fundamental but can be quite involved in real-world scenarios. Let’s take an example.

▲ **Example 3.7.2** Determine the critical points of the following functions using the First Derivative Test.

1. $f(x, y) = x^3 - 3xy^2$
2. $h(x, y, z) = x^2 + y^2 + z^2 - 4x - 6y - 8z$
3. $g(x, y) = e^x + e^y + xy$

Solution

1. **Function** $f(x, y) = x^3 - 3xy^2$

(a) **Compute the Partial Derivatives:**

$$f_x = \frac{\partial f}{\partial x} = 3x^2 - 3y^2$$

$$f_y = \frac{\partial f}{\partial y} = -6xy$$

(b) **Set the Gradient Equal to Zero:**

$$\nabla f = \langle 3x^2 - 3y^2, -6xy \rangle = \mathbf{0}$$

This leads to the system:

$$3x^2 - 3y^2 = 0 \quad \text{and} \quad -6xy = 0$$

(c) **Solve for Critical Points:**

$$3x^2 - 3y^2 = 0 \implies x^2 = y^2$$

$$-6xy = 0 \implies x = 0 \quad \text{or} \quad y = 0$$

- **Case 1:** $x = 0$

$$x^2 = y^2 \implies 0 = y^2 \implies y = 0$$

Critical point: $(0, 0)$

- **Case 2:** $y = 0$

$$x^2 = 0 \implies x = 0$$

Critical point: $(0, 0)$

Thus, the only critical point is $(0, 0)$.

2. **Function** $h(x, y, z) = x^2 + y^2 + z^2 - 4x - 6y - 8z$

(a) **Compute the Partial Derivatives:**

$$h_x = \frac{\partial h}{\partial x} = 2x - 4$$

$$h_y = \frac{\partial h}{\partial y} = 2y - 6$$

$$h_z = \frac{\partial h}{\partial z} = 2z - 8$$

(b) **Set the Gradient Equal to Zero:**

$$\nabla h = \langle 2x - 4, 2y - 6, 2z - 8 \rangle = \mathbf{0}$$

This leads to the system:

$$2x - 4 = 0 \quad \Rightarrow \quad x = 2$$

$$2y - 6 = 0 \quad \Rightarrow \quad y = 3$$

$$2z - 8 = 0 \quad \Rightarrow \quad z = 4$$

Thus, the critical point is $(2, 3, 4)$.

3. **Function** $g(x, y) = e^x + e^y + xy$

(a) **Compute the Partial Derivatives:**

$$g_x = \frac{\partial g}{\partial x} = e^x + y$$

$$g_y = \frac{\partial g}{\partial y} = e^y + x$$

(b) **Set the Gradient Equal to Zero:**

$$\nabla g = \langle e^x + y, e^y + x \rangle = \mathbf{0}$$

This leads to the system:

$$e^x + y = 0 \quad \text{and} \quad e^y + x = 0$$

(c) **Solve for Critical Points:**

From the first equation:

$$y = -e^x$$

Substitute into the second equation:

$$e^{-e^x} + x = 0$$

This transcendental equation does not have a closed-form solution, so we identify the critical point numerically using Wolfram Alpha or another online tool. The approximate solution is:

$$x \approx -0.567143, \quad y \approx -0.567143$$

Thus, the approximate critical point is $(-0.567143, -0.567143)$.

Second Derivative Tests for Functions of Two Variables

After employing the first derivative test (i.e., finding points where the gradient $\nabla f = \mathbf{0}$) to identify critical points of a two-variable function $f(x, y)$, the next step is to determine the nature of these critical points: are they local maxima, local minima, or something else? In single-variable calculus, this classification depended on the sign of the second derivative $f''(a)$. In the two-variable setting, we must consider second partial derivatives and how they interact.

From Single to Multiple Variables: The Discriminant

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function with continuous second partial derivatives. Suppose (a, b) is a critical point, i.e.,

$$f_x(a, b) = 0 \quad \text{and} \quad f_y(a, b) = 0.$$

To classify (a, b) , we consider the second partial derivatives at (a, b) :

$$f_{xx}(a, b) = \frac{\partial^2 f}{\partial x^2}(a, b), \quad f_{yy}(a, b) = \frac{\partial^2 f}{\partial y^2}(a, b), \quad f_{xy}(a, b) = \frac{\partial^2 f}{\partial x \partial y}(a, b).$$

We define the **discriminant** D as:

$$D = f_{xx}(a, b)f_{yy}(a, b) - [f_{xy}(a, b)]^2.$$

This discriminant D encapsulates how the function curves in different directions near the critical point. Notably, D is also the determinant of the Hessian matrix at (a, b) in two dimensions. The Hessian matrix arises because of the fact in a multivariate setting, the second derivatives of variables have to go through all the combinations of derivatives of two variables which gets messy in higher dimensions. For now, in two dimensions, the Hessian matrix is:

$$H(a, b) = \begin{bmatrix} f_{xx}(a, b) & f_{xy}(a, b) \\ f_{yx}(a, b) & f_{yy}(a, b) \end{bmatrix}.$$

And thus,

$$D = \det(H(a, b)) = f_{xx}(a, b)f_{yy}(a, b) - [f_{xy}(a, b)]^2.$$

More on the Hessian, and some of these linear algebra terms to come later.

Local Extrema

Using the discriminant D and the sign of $f_{xx}(a, b)$, we have a test analogous to the single-variable scenario:

- If $D > 0$ and $f_{xx}(a, b) > 0$, (a, b) is a **local minimum**.
- If $D > 0$ and $f_{xx}(a, b) < 0$, (a, b) is a **local maximum**.
- If $D < 0$, (a, b) is neither a max nor a min (it may be a saddle point).
- If $D = 0$, the test is inconclusive, requiring further analysis.

This gives us a direct way to classify local maxima and minima for functions of two variables once we know their first and second derivatives at a critical point.

▲ **Example 3.7.3** Determine the nature of the critical point of the function $f(x, y) = 2x^2 + 3y^2 - 4x - 6y + 10$ using the Second Derivative Test.

Solution

1. Find the Critical Point:

(a) Compute the First Partial Derivatives:

$$f_x = \frac{\partial f}{\partial x} = 4x - 4$$

$$f_y = \frac{\partial f}{\partial y} = 6y - 6$$

(b) Set the Gradient Equal to Zero:

$$\nabla f = \langle 4x - 4, 6y - 6 \rangle = \mathbf{0}$$

Solving the system:

$$4x - 4 = 0 \quad \Rightarrow \quad x = 1$$

$$6y - 6 = 0 \quad \Rightarrow \quad y = 1$$

Thus, the critical point is $(1, 1)$.

2. Classify the Critical Point Using the Second Derivative Test:

(a) Compute the Second Partial Derivatives:


$$f_{xx} = \frac{\partial^2 f}{\partial x^2} = 4, f_{yy} = \frac{\partial^2 f}{\partial y^2} = 6, f_{xy} = \frac{\partial^2 f}{\partial x \partial y} = 0$$

(b) Calculate the Discriminant D :

$$D = f_{xx}(1, 1) \cdot f_{yy}(1, 1) - [f_{xy}(1, 1)]^2 = (4)(6) - (0)^2 = 24 > 0$$

(c) Apply the Classification Rules:

- Since $D > 0$ and $f_{xx}(1, 1) = 4 > 0$, $(1, 1)$ is a **local minimum**. Let's draw a picture! We can mark the local min with a red dot:



Global Extrema Considerations

While the second derivative test helps us determine local behavior near a critical point, identifying global maxima or minima is a separate question. In single-variable calculus, the Extreme Value Theorem guarantees that a continuous function defined on a closed, bounded interval attains a global maximum and minimum. Similarly, for functions of two variables, if f is continuous on a compact (closed and bounded) set in $\mathbb{R}^2$, then f attains a global maximum and minimum on that set.

Finding global extrema typically involves:

1. Identifying all critical points inside the domain.
2. Checking the behavior on the boundary of the domain (since global extrema may occur there, not just at critical points).
3. Comparing the values of f at these points to see which are the largest and smallest.

If the domain is not compact—if it is unbounded or not closed—then f may fail to achieve global maxima or minima altogether. The function might “escape” to infinity in some directions, or may never settle on a maximum value. Let’s take a look at an example:

▲ **Example 3.7.4** Determine the global maximum and minimum of the function $f(x, y) = 2x^2 + 3y^2 - 4x - 6y + 10$ on the closed and bounded region R defined by $0 \leq x \leq 3$ and $0 \leq y \leq 3$.

Solution

1. Identify Critical Points:

- First Partial Derivatives:

$$f_x = 4x - 4 \quad \text{and} \quad f_y = 6y - 6$$

- Setting the Gradient to Zero:

$$4x - 4 = 0 \quad \Rightarrow \quad x = 1$$

$$6y - 6 = 0 \quad \Rightarrow \quad y = 1$$

- Critical Point:

$$(1, 1)$$

2. Classify the Critical Point Using the Second Derivative Test:

- Second Partial Derivatives:

$$f_{xx} = 4, \quad f_{yy} = 6, \quad f_{xy} = 0$$

- Discriminant Calculation:

$$D = f_{xx}f_{yy} - (f_{xy})^2 = (4)(6) - (0)^2 = 24 > 0$$

- Classification:

– Since $D > 0$ and $f_{xx} > 0$, the critical point $(1, 1)$ is a **local minimum**.

3. Evaluate the Function on the Boundary of R :

The boundary of R consists of four edges. We evaluate $f(x, y)$ at critical points on each edge and at the corners. For the edges, we will pick some values on the inside of the edge.

Edge	Point	Function Value
$x = 0$	$(0, 1)$	$f(0, 1) = 7$
$x = 3$	$(3, 1)$	$f(3, 1) = 13$
$y = 0$	$(1, 0)$	$f(1, 0) = 8$
$y = 3$	$(1, 3)$	$f(1, 3) = 17$
Corners	$(0, 0)$	$f(0, 0) = 10$
	$(0, 3)$	$f(0, 3) = 19$
	$(3, 0)$	$f(3, 0) = 16$
	$(3, 3)$	$f(3, 3) = 25$


4. Determine Global Extrema:

- Global Minimum: $f(1, 1) = 5$ at $(1, 1)$
- Global Maximum: $f(3, 3) = 25$ at $(3, 3)$

Conclusion:

- The function $f(x, y)$ attains its **global minimum** of 5 at $(1, 1)$.
- The function $f(x, y)$ attains its **global maximum** of 25 at $(3, 3)$.

Visual Representation of the Function and Its Global Extrema:



The plot above illustrates the function $f(x, y) = 2x^2 + 3y^2 - 4x - 6y + 10$ over the region R . The global minimum at $(1, 1, 5)$ is marked with a **red circle**, and the global maximum at $(3, 3, 25)$ with a **blue triangle**. Refer to the legend below the graph for clarification.

The plot above illustrates the function $f(x, y) = 2x^2 + 3y^2 - 4x - 6y + 10$ over the region R . The global minimum at $(1, 1, 5)$ is marked in red, and the global maximum at $(3, 3, 25)$ is marked in blue.

Saddle Points: A New Category of Critical Points

We've danced around it throughout this chapter, but we are now going to formally define it...in single-variable calculus, if $f'(a) = 0$ and $f''(a) = 0$, you might get a point of inflection. Inflection points are where the function's curvature changes sign. In multiple dimensions, we have a richer phenomenon: the **saddle point**.

A saddle point is a critical point that is not a maximum or a minimum. Think of a horse saddle: at the saddle's center, if you move forward and backward along the horse's spine, you might go up; but if you move left and right across the saddle, you might go down. The point is neither a peak nor a valley—some directions lead you higher, others lead you lower. Mathematically, this corresponds to the Hessian being indefinite at that point.




Figure 3.27: A saddle point at $(0, 0)$ for the function $z = x^2 - y^2$. Along the x -axis, the function looks like a minimum; along the y -axis, it looks like a maximum. Overall, it's neither, defining a classic saddle shape.

In the example above, the function $f(x, y) = x^2 - y^2$ has a critical point at $(0, 0)$. Checking the criteria:

$$f_{xx} = 2, \quad f_{yy} = -2, \quad f_{xy} = 0.$$

Therefore we have $D = 2(-2) - (0^2) = -4 < 0$. Thus, confirming a saddle point. Our previous example we saw in Figure 3.24 we indeed have one saddle point. Let's find it!

▲ **Example 3.7.5** Determine the saddle point of the function $f(x, y) = \sin(x) \sin(y) e^{x^2+y^2}$.

Solution

1. **Identify Critical Points:**

- First Partial Derivatives:

$$f_x = \sin(y) e^{x^2+y^2} (\cos(x) + 2x \sin(x)), \quad f_y = \sin(x) e^{x^2+y^2} (\cos(y) + 2y \sin(y)).$$

- Setting $f_x = 0$ and $f_y = 0$:

$$\sin(y) = 0 \quad \text{or} \quad \cos(x) + 2x \sin(x) = 0,$$

$$\sin(x) = 0 \quad \text{or} \quad \cos(y) + 2y \sin(y) = 0.$$

- Critical Point:

$$(x, y) = (0, 0).$$

2. **Classify the Critical Point Using the Second Derivative Test:**

- Second Partial Derivatives at $(0, 0)$:

$$f_{xx} = 0, \quad f_{yy} = 0, \quad f_{xy} = 1.$$

- Discriminant Calculation:

$$D = f_{xx}f_{yy} - (f_{xy})^2 = (0)(0) - (1)^2 = -1 < 0.$$

- Classification:

– Since $D < 0$, the critical point $(0, 0)$ is a **saddle point**.

Generalization to n -Variables and the Hessian Matrix

For functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$, the second derivative test uses the **Hessian matrix**, an $n \times n$ matrix of second partial derivatives:

$$H(\mathbf{a}) = \begin{bmatrix} f_{x_1x_1}(\mathbf{a}) & f_{x_1x_2}(\mathbf{a}) & \cdots & f_{x_1x_n}(\mathbf{a}) \\ f_{x_2x_1}(\mathbf{a}) & f_{x_2x_2}(\mathbf{a}) & \cdots & f_{x_2x_n}(\mathbf{a}) \\ \vdots & \vdots & \ddots & \vdots \\ f_{x_nx_1}(\mathbf{a}) & f_{x_nx_2}(\mathbf{a}) & \cdots & f_{x_nx_n}(\mathbf{a}) \end{bmatrix}.$$

The two-variable discriminant D is just the determinant of the 2×2 Hessian for $n = 2$. For higher dimensions, determining whether a critical point is a local maximum, minimum, or saddle point involves examining the **definiteness** of the Hessian matrix. This is where the concepts of positive definite, negative definite, indefinite, and semidefinite come into play. To understand these terms, we first introduce the concept of eigenvalues:

- **Eigenvalues:** Mathematically, an eigenvalue λ of a square matrix A is a scalar such that there exists a nonzero vector $\mathbf{v}$ (called an eigenvector) satisfying the equation:

$$A\mathbf{v} = \lambda\mathbf{v}.$$

In this equation, the matrix A acts on the eigenvector $\mathbf{v}$, and the result is the same as scaling $\mathbf{v}$ by λ . To find eigenvalues, we solve the characteristic equation:

$$\det(A - \lambda I) = 0,$$

where I is the identity matrix. The eigenvalues of the Hessian matrix H provide insight into the curvature of the function in specific directions. Positive eigenvalues correspond to upward curvature (like a local minimum), while negative eigenvalues correspond to downward curvature (like a local maximum). Mixed signs indicate the presence of a saddle point.

The definiteness of the Hessian matrix is determined by the signs of its eigenvalues:

- **Positive Definite Hessian:**
 - All eigenvalues of the Hessian are positive.
 - Implication: The function curves upwards in all directions around the critical point, indicating a **local minimum**.
- **Negative Definite Hessian:**
 - All eigenvalues of the Hessian are negative.
 - Implication: The function curves downwards in all directions around the critical point, indicating a **local maximum**.
- **Indefinite Hessian:**
 - The Hessian has both positive and negative eigenvalues.
 - Implication: The function curves upwards in some directions and downwards in others around the critical point, indicating a **saddle point**.
- **Semidefinite Hessian:**
 - The Hessian has eigenvalues that are all non-negative or all non-positive, but not all strictly positive or negative.
 - Implication: The test is **inconclusive**, and further analysis is required to classify the critical point.

While the linear algebra underlying definiteness and eigenvalues is more advanced, the key takeaway is that the Hessian matrix generalizes the notion of the second derivative test to any dimension. In two dimensions, the discriminant $D = \det(H)$ and the sign of $f_{xx}(a, b)$ provide a simpler, more direct criterion for classifying critical points.

Recall that we saw for a 2×2 Hessian matrix:

$$H = \begin{bmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{bmatrix},$$

the discriminant D is calculated as:

$$D = f_{xx}f_{yy} - (f_{xy})^2.$$

Now with higher dimensions, we need to discuss the definiteness of the Hessian in terms of classifying critical points:

- If $D > 0$ and $f_{xx} > 0$, the Hessian is **positive definite**, indicating a **local minimum**.
- If $D > 0$ and $f_{xx} < 0$, the Hessian is **negative definite**, indicating a **local maximum**.
- If $D < 0$, the Hessian is **indefinite**, indicating a **saddle point**.
- If $D = 0$, the Hessian is **semidefinite**, and the test is **inconclusive**.

Let's move up a dimension. When we consider a function of three variables, $w = f(x, y, z)$, we will have more combinations of second derivatives and thus our Hessian will be:

$$H = \begin{bmatrix} f_{xx} & f_{xy} & f_{xz} \\ f_{yx} & f_{yy} & f_{yz} \\ f_{zx} & f_{zy} & f_{zz} \end{bmatrix}.$$

This will result in a more complicated discriminant:

$$D = \det(H) = f_{xx}(f_{yy}f_{zz} - f_{yz}^2) - f_{xy}(f_{yx}f_{zz} - f_{yz}f_{zx}) + f_{xz}(f_{yx}f_{zy} - f_{yy}f_{zx}).$$

The criteria for a function of three variables gets a little more complicated: we first examine the structure of the Hessian and the determinant of the 2×2 upper-left submatrix, which corresponds to the case of two variables. This submatrix is:

$$H_2 = \begin{bmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{bmatrix},$$

and its determinant is given by:

$$D_2 = \det(H_2) = f_{xx}f_{yy} - (f_{xy})^2.$$

For a function of three variables, the **leading principal minors** are D_1 , D_2 , and D_3 , where:

- $D_1 = f_{xx}$,
- $D_2 = \det(H_2) = f_{xx}f_{yy} - (f_{xy})^2$,
- $D_3 = \det(H)$, the determinant of the full 3×3 Hessian.

The criteria for definiteness in three dimensions are:

- If $D_3 > 0$, $D_2 > 0$, and $D_1 > 0$, the Hessian is **positive definite**, indicating a **local minimum**.
- If $D_3 > 0$, $D_2 > 0$, and $D_1 < 0$, the Hessian is **negative definite**, indicating a **local maximum**.
- If $D_3 < 0$, the Hessian is **indefinite**, indicating a **saddle point**.
- If $D_3 = 0$, the Hessian is **semidefinite**, and the test is **inconclusive**.

Let us now extend this reasoning to n -dimensional functions. For a function $f(\mathbf{x})$, where $\mathbf{x} = (x_1, x_2, \dots, x_n)$, the Hessian is an $n \times n$ matrix:

$$H = \begin{bmatrix} f_{x_1x_1} & f_{x_1x_2} & \cdots & f_{x_1x_n} \\ f_{x_2x_1} & f_{x_2x_2} & \cdots & f_{x_2x_n} \\ \vdots & \vdots & \ddots & \vdots \\ f_{x_nx_1} & f_{x_nx_2} & \cdots & f_{x_nx_n} \end{bmatrix}.$$

The determinant of the Hessian for n -dimensions is given by:

$$D_n = \det(H).$$

The criteria for definiteness are generalized using **Sylvester's Criterion**, which states:

- The Hessian is **positive definite** (indicating a local minimum) if all leading principal minors are positive:

$$D_1 > 0, \quad D_2 > 0, \quad \dots, \quad D_n > 0.$$

- The Hessian is **negative definite** (indicating a local maximum) if the leading principal minors alternate in sign, starting with $D_1 < 0$:

$$D_1 < 0, \quad D_2 > 0, \quad D_3 < 0, \quad \dots, \quad D_n \text{ has sign } (-1)^n.$$

- The Hessian is **indefinite** (indicating a saddle point) if the sequence of leading principal minors does not satisfy the conditions for positive or negative definiteness.
- The Hessian is **semidefinite** if one or more leading principal minors are zero, and the remaining minors satisfy the conditions for definiteness.

Sylvester's Criterion provides a systematic method to evaluate the definiteness of the Hessian for functions of any dimension, extending the utility of the second derivative test to higher-dimensional problems.

This framework allows us to extend the second derivative test from single-variable calculus to functions of multiple variables, enabling the classification of critical points in higher dimensions.

Lecture 3.8: Optimization

In the previous lecture, we developed tools for identifying and classifying extrema of multivariable functions, analyzing both local and global extrema using critical points and the Hessian matrix. These methods extended single-variable concepts to multiple dimensions, allowing us to handle optimization problems in unconstrained settings – indeed, we implicitly handled unconstrained optimization to a large degree by carefully and deeply examining extrema. However, real-world optimization problems often involve constraints: limits imposed by resources, boundaries, or conditions that must be satisfied. This lecture introduces techniques for optimization in constrained settings, starting with Lagrange multipliers and culminating in a discussion on how these methods generalize to applications in fields such as machine learning.



Figure 3.28: *Manufacturing is the foremost application of optimization in our day-to-day lives. You may even have noticed what is called ‘Shrinkflation’: goods being shrunk down with a higher price. This is due to cost optimization.*

Unconstrained Optimization

Optimization is a cornerstone of mathematics, embodying the search for maxima and minima—points where a function reaches its highest or lowest values. These principles permeate problems across science, engineering, and economics, where achieving efficiency or minimizing costs is a primary objective. In this lecture, we dig deeper into the concepts of extrema by clarifying them with an applied nature. As we shift toward constrained optimization, it is essential to reflect on the foundational tools of *unconstrained optimization*, which underlie more intricate scenarios.

Recall from our earlier discussions that the key to finding extrema lies in identifying points where the behavior of a function undergoes a qualitative change. For a differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, such points—termed **critical points**—are characterized by the condition:

$$\nabla f(\mathbf{x}) = \mathbf{0}.$$

At these points, the gradient, which represents the direction of steepest ascent, becomes zero, signaling that the function is momentarily stationary. Yet as we learned, not all critical points correspond to maxima or minima. Some may represent *saddle points*, where the function exhibits an increase in certain directions and a decrease in others. This necessitated a framework for systematically classifying critical points.

The tools we have developed for this classification are summarized below:

1. **The Gradient Condition:** Critical points are identified by solving $\nabla f(\mathbf{x}) = \mathbf{0}$. This forms the foundation for locating potential extrema.
2. **The Second Derivative Test:** Once critical points are located, the nature of each point is determined by examining the **Hessian matrix** $H(\mathbf{x})$, the matrix of second partial derivatives of f . For two-variable functions, the determinant of the Hessian, denoted as D , provides a straightforward method for classification:

$$D = f_{xx}f_{yy} - (f_{xy})^2.$$

The following rules apply:

- If $D > 0$ and $f_{xx} > 0$, the point is a local minimum.
 - If $D > 0$ and $f_{xx} < 0$, the point is a local maximum.
 - If $D < 0$, the point is neither a maximum nor a minimum and is instead a saddle point.
 - If $D = 0$, the test is inconclusive, requiring further analysis.
3. **Boundary Analysis:** For functions defined on bounded domains, extrema may occur on the boundary. This requires evaluating f along the domain's boundary in addition to examining critical points within the interior.

Together, these tools provide a comprehensive framework for unconstrained optimization. By systematically identifying and classifying critical points and incorporating boundary conditions where necessary, we can thoroughly analyze the behavior of a function over its domain.

It is worth emphasizing that unconstrained optimization closely mirrors the techniques we have already mastered. The interplay between geometry and calculus is a recurring theme: the condition $\nabla f = \mathbf{0}$ encapsulates the geometric intuition that the gradient vanishes at extrema, while the second derivative test provides a quantitative measure of curvature. This shared foundation ensures a seamless transition into more advanced topics, as the principles of unconstrained optimization naturally extend to problems involving constraints.

▲ **Example 3.8.1** A company wants to minimize the cost of manufacturing a product. The cost function is given by

$$f(x, y) = x^2 + y^2 - 4x - 6y + 13,$$

where x and y represent two key factors influencing production efficiency. Determine the optimal values of x and y to achieve the minimum cost.

Solution This is an unconstrained optimization problem because there are no restrictions on the values x and y can take. Solving this involves finding the extrema of $f(x, y)$, which simplifies to finding the minimum.

1. Identify Critical Points:

- Compute the gradient of f :

$$\nabla f = \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \end{bmatrix} = \begin{bmatrix} 2x - 4 \\ 2y - 6 \end{bmatrix}.$$

- Set $\nabla f = \mathbf{0}$ to find critical points:

$$2x - 4 = 0 \quad \text{and} \quad 2y - 6 = 0.$$

Solving these equations gives $x = 2$ and $y = 3$. Thus, the critical point is $(2, 3)$.

2. Classify the Critical Point Using the Second Derivative Test:

- Compute the Hessian matrix H :

$$H = \begin{bmatrix} \frac{\partial^2 f}{\partial x^2} & \frac{\partial^2 f}{\partial x \partial y} \\ \frac{\partial^2 f}{\partial y \partial x} & \frac{\partial^2 f}{\partial y^2} \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}.$$

- Calculate the determinant of H :

$$D = \frac{\partial^2 f}{\partial x^2} \cdot \frac{\partial^2 f}{\partial y^2} - \left(\frac{\partial^2 f}{\partial x \partial y} \right)^2 = (2)(2) - (0)^2 = 4 > 0.$$

- Since $D > 0$ and $\frac{\partial^2 f}{\partial x^2} = 2 > 0$, the critical point $(2, 3)$ is a **local minimum**.

3. Evaluate the Function at the Critical Point:

- Substituting $x = 2$ and $y = 3$ into $f(x, y)$:

$$f(2, 3) = (2)^2 + (3)^2 - 4(2) - 6(3) + 13 = 4 + 9 - 8 - 18 + 13 = 1.$$

- The minimum cost is 1, achieved when $x = 2$ and $y = 3$.

This simple example reinforces the utility of these methods and foreshadows their extensions to more complex scenarios.

Classical Examples of Unconstrained Optimization

The classical examples we explore highlight the versatility of unconstrained optimization in various settings. For each, we provide a commonly optimized formula to solidify the connection to practical applications:

1. Economic Models:

- Profit Maximization: Optimize the profit function $P = R - C$, where R is revenue and C is cost.
- Cost Minimization: Minimize $C(q)$, where C is the cost function and q is the quantity of goods or services produced.

2. Physics Applications:

- Potential Energy Minimization: Minimize $U = \frac{1}{2}kx^2$, the potential energy of a spring system, where k is the spring constant and x is the displacement.
- Kinetic Energy Optimization: Maximize $K = \frac{1}{2}mv^2$, where m is mass and v is velocity.

3. Machine Learning:

- Loss Function Minimization: Minimize a loss function $L(w) = \frac{1}{N} \sum_{i=1}^N (y_i - f(x_i, w))^2$, where w is the parameter vector, (x_i, y_i) are data points, and f is the model.

4. Geometric Optimization:

- Volume of a Box: Maximize $V = l \cdot w \cdot h$, where l , w , and h are length, width, and height.
- Volume of a Sphere: Maximize $V = \frac{4}{3}\pi r^3$, where r is the radius.
- Volume of a Cylinder: Maximize $V = \pi r^2 h$, where r is the radius and h is the height.
- Solids of Revolution: Optimize the volume $V = \pi \int_a^b [f(x)]^2 dx$, where $f(x)$ is the curve being rotated about an axis.

These examples provide a foundation for solving unconstrained optimization problems across disciplines. Each formula highlights a tangible objective, whether maximizing profit, minimizing energy, training a machine learning model, or optimizing the geometry of an object. Geometric optimization, in particular, lays the groundwork for exploring practical shapes and volumes used in design and engineering.

▲ **Example 3.8.2** Find the maximum volume of a rectangular box with its base in the xy -plane and its top lying on the plane $z = 12 - 2x - 3y$ in the first octant.

Solution

1. Formulate the Volume Function:

- The base of the box lies in the xy -plane, and its top touches the plane $z = 12 - 2x - 3y$.
- The dimensions of the box are x , y , and z , where z is given by the equation of the plane.
- The volume of the box is:

$$V = x \cdot y \cdot z = x \cdot y \cdot (12 - 2x - 3y).$$

Expand this expression:

$$V(x, y) = 12xy - 2x^2y - 3xy^2.$$

2. Find the Critical Points:

- Compute the partial derivatives of V :

$$V_x = \frac{\partial V}{\partial x} = 12y - 4xy - 3y^2,$$

$$V_y = \frac{\partial V}{\partial y} = 12x - 2x^2 - 6xy.$$

- Set $V_x = 0$ and $V_y = 0$:

$$12y - 4xy - 3y^2 = 0 \quad (1),$$

$$12x - 2x^2 - 6xy = 0 \quad (2).$$

Factor equation (1):

$$y(12 - 4x - 3y) = 0.$$

Thus, $y = 0$ or $12 - 4x - 3y = 0$, giving $y = \frac{12-4x}{3}$.

Factor equation (2):

$$x(12 - 2x - 6y) = 0.$$

Thus, $x = 0$ or $12 - 2x - 6y = 0$, giving $y = \frac{12-2x}{6} = 2 - \frac{x}{3}$.

- Solve the system of equations: Substituting $y = \frac{12-4x}{3}$ into $y = 2 - \frac{x}{3}$:

$$\frac{12-4x}{3} = 2 - \frac{x}{3}.$$

Multiply through by 3:

$$12 - 4x = 6 - x \quad \Rightarrow \quad 3x = 6 \quad \Rightarrow \quad x = 2.$$

Substitute $x = 2$ into $y = 2 - \frac{x}{3}$:

$$y = 2 - \frac{2}{3} = \frac{6}{3} - \frac{2}{3} = \frac{4}{3}.$$

3. Classify the Critical Point and Find the Volume:

The critical point is $(x, y) = (2, \frac{4}{3})$. At this point:

$$z = 12 - 2x - 3y = 12 - 2(2) - 3\left(\frac{4}{3}\right) = 12 - 4 - 4 = 4.$$

The dimensions of the box are $x = 2$, $y = \frac{4}{3}$, and $z = 4$.

$$V = x \cdot y \cdot z = 2 \cdot \frac{4}{3} \cdot 4 = \frac{32}{3}.$$

Optimization with Equality Constraints

Consider the task of optimizing a function $f(x, y)$ subject to a constraint $g(x, y) = c$, where g is a differentiable function that defines a curve or surface in $\mathbb{R}^n$. This problem can be visualized as finding the highest or lowest point on the curve or surface defined by $g(x, y) = c$, rather than across the entire plane or space. To solve this, we introduce the method of **Lagrange multipliers**.

To build intuition, recall that the gradient $\nabla g(x, y)$ of the constraint function points in the direction of the steepest ascent of $g(x, y)$. The curve $g(x, y) = c$ represents a level set of g , and at any point along this curve, $\nabla g(x, y)$ is normal to the curve. Similarly, the gradient $\nabla f(x, y)$ of the objective function $f(x, y)$ indicates the direction of steepest ascent of f . At an extremum of f constrained to $g(x, y) = c$, these gradients must align, meaning $\nabla f(x, y)$ must point in the same or opposite direction as $\nabla g(x, y)$.

This condition is expressed mathematically as:

$$\nabla f(x, y) = \lambda \nabla g(x, y),$$

where λ is a scalar known as the **Lagrange multiplier**. The scalar λ measures how much the objective function f changes per unit increase in the constraint g .

The Lagrange Condition

To derive this condition, consider the constrained optimization problem as finding the extrema of $f(x, y)$ on the curve $g(x, y) = c$. We construct the *Lagrange function*:

$$\mathcal{L}(x, y, \lambda) = f(x, y) + \lambda(g(x, y) - c).$$

The term $\lambda(g(x, y) - c)$ enforces the constraint $g(x, y) = c$. At the extrema of $f(x, y)$ subject to this constraint, the gradients of $\mathcal{L}$ with respect to x , y , and λ must vanish:

$$\frac{\partial \mathcal{L}}{\partial x} = \frac{\partial f}{\partial x} + \lambda \frac{\partial g}{\partial x} = 0, \quad \frac{\partial \mathcal{L}}{\partial y} = \frac{\partial f}{\partial y} + \lambda \frac{\partial g}{\partial y} = 0, \quad \frac{\partial \mathcal{L}}{\partial \lambda} = g(x, y) - c = 0.$$

These equations are equivalent to the system:

$$\begin{aligned} \nabla f(x, y) - \lambda \nabla g(x, y) &= \mathbf{0}, \\ g(x, y) &= c. \end{aligned}$$

The first equation ensures that ∇f and ∇g are parallel (aligned with λ), while the second equation enforces the constraint. Thus, the optimization problem reduces to solving this system of equations for x , y , and λ .

To solve an optimization problem using Lagrange multipliers:

1. Compute $\nabla f(x, y)$ and $\nabla g(x, y)$.
2. Set up the system of equations:

$$\nabla f(x, y) - \lambda \nabla g(x, y) = \mathbf{0}, \quad g(x, y) = c.$$

3. Solve the system for x , y , and λ .
4. Evaluate $f(x, y)$ at the solutions to determine the extrema.

The equation $\nabla f = \lambda \nabla g$ has a powerful geometric interpretation. Recall that $\nabla g(x, y)$ is orthogonal to the level curve $g(x, y) = c$ at any point. Similarly, $\nabla f(x, y)$ is orthogonal to the level curve of $f(x, y)$. Thus, the condition $\nabla f = \lambda \nabla g$ implies that the level curves of f and g are tangent at the optimal point. This tangency ensures that $f(x, y)$ is extremized subject to the constraint $g(x, y) = c$.

▲ **Example 3.8.3** Find the minimum of the function $f(x, y) = x^2 + y^2$ subject to the constraint $x + y = 2$.

Solution

1. **Formulate the Lagrangian:**

To apply the method of Lagrange multipliers, we introduce a Lagrange multiplier

λ and construct the Lagrangian function:

$$\mathcal{L}(x, y, \lambda) = f(x, y) - \lambda(g(x, y) - c),$$

where $g(x, y) = x + y$ and $c = 2$.

Substituting the given functions:

$$\mathcal{L}(x, y, \lambda) = x^2 + y^2 - \lambda(x + y - 2).$$

2. Compute the Partial Derivatives:

To find the extrema, take the partial derivatives of $\mathcal{L}$ with respect to x , y , and λ , and set them equal to zero.

- Partial derivative with respect to x :

$$\frac{\partial \mathcal{L}}{\partial x} = 2x - \lambda = 0 \quad \Rightarrow \quad \lambda = 2x \quad (1)$$

- Partial derivative with respect to y :

$$\frac{\partial \mathcal{L}}{\partial y} = 2y - \lambda = 0 \quad \Rightarrow \quad \lambda = 2y \quad (2)$$

- Partial derivative with respect to λ :

$$\frac{\partial \mathcal{L}}{\partial \lambda} = -(x + y - 2) = 0 \quad \Rightarrow \quad x + y = 2 \quad (3)$$

3. Solve the System of Equations:

From equations (1) and (2):

$$2x = \lambda \quad \text{and} \quad 2y = \lambda \quad \Rightarrow \quad 2x = 2y \quad \Rightarrow \quad x = y$$

Substitute $x = y$ into equation (3):

$$x + x = 2 \quad \Rightarrow \quad 2x = 2 \quad \Rightarrow \quad x = 1$$

Since $x = y$:

$$y = 1$$

Thus, the critical point is $(x, y) = (1, 1)$.

4. Verify the Solution:

Substitute $x = 1$ and $y = 1$ back into the objective function:

$$f(1, 1) = (1)^2 + (1)^2 = 1 + 1 = 2$$

Therefore, the function $f(x, y) = x^2 + y^2$ attains its minimum value of 2 at the point $(1, 1)$ under the constraint $x + y = 2$.

5. Visualization:

To better understand the optimization, let's visualize the objective function, the constraint, and the optimal point.




Figure 3.29: Visualization of the optimization problem using Lagrange multipliers. The blue circles represent the contour lines of the objective function $f(x, y) = x^2 + y^2$, the red line is the constraint $x + y = 2$, the black dot marks the optimal point $(1, 1)$, the green arrow represents the gradient ∇f , and the orange arrow represents the gradient ∇g .

Let's now try a problem that's a little messier and doesn't have a nice answer. We will need to use a computer to solve a polynomial using numerical approximation at the end. This is more representative a problem you may encounter in an actual professional setting.

▲ **Example 3.8.4** Determine the production levels x and y for Gadget A and Gadget B, respectively, that minimize the total production cost given by:

$$C(x, y) = x^2y + y^3,$$

subject to the nonlinear constraint:

$$x^3 + y^3 = 2000.$$

Since production levels must be integer values, round the approximate solution up or down and choose the option with the lesser cost.

Solution

1. Formulate the Lagrangian:

Introduce a Lagrange multiplier λ and construct the Lagrangian function:

$$\mathcal{L}(x, y, \lambda) = C(x, y) - \lambda(x^3 + y^3 - 2000).$$

Substituting the given functions:

$$\mathcal{L}(x, y, \lambda) = x^2y + y^3 - \lambda(x^3 + y^3 - 2000).$$

2. Compute the Partial Derivatives:

To find the extrema, take the partial derivatives of $\mathcal{L}$ with respect to x , y , and λ , and set them equal to zero.

• **Partial derivative with respect to x :**

$$\frac{\partial \mathcal{L}}{\partial x} = 2xy - 3\lambda x^2 = 0 \quad \Rightarrow \quad 2xy = 3\lambda x^2 \quad (1)$$

• **Partial derivative with respect to y :**

$$\frac{\partial \mathcal{L}}{\partial y} = x^2 + 3y^2 - 3\lambda y^2 = 0 \quad \Rightarrow \quad x^2 + 3y^2 = 3\lambda y^2 \quad (2)$$

• **Partial derivative with respect to λ :**

$$\frac{\partial \mathcal{L}}{\partial \lambda} = -(x^3 + y^3 - 2000) = 0 \quad \Rightarrow \quad x^3 + y^3 = 2000 \quad (3)$$

3. Solve the System of Equations:

From equation (1):

$$2xy = 3\lambda x^2 \quad \Rightarrow \quad \lambda = \frac{2y}{3x} \quad (1a)$$

From equation (2):

$$x^2 + 3y^2 = 3\lambda y^2 \quad \Rightarrow \quad \lambda = \frac{x^2 + 3y^2}{3y^2} = \frac{x^2}{3y^2} + 1 \quad (2a)$$

Equate the two expressions for λ :

$$\frac{2y}{3x} = \frac{x^2}{3y^2} + 1$$

Multiply both sides by $3xy^2$ to eliminate denominators:

$$\begin{aligned} 2y \cdot y^2 &= x^3 + 3xy^2 \\ 2y^3 &= x^3 + 3xy^2 \quad (4) \end{aligned}$$

From the constraint (3):

$$x^3 + y^3 = 2000 \quad \Rightarrow \quad x^3 = 2000 - y^3 \quad (5)$$

Substitute $x^3 = 2000 - y^3$ into equation (4):

$$\begin{aligned} 2y^3 &= (2000 - y^3) + 3xy^2 \\ 2y^3 &= 2000 - y^3 + 3xy^2 \\ 3xy^2 &= 3y^3 - 2000 \\ xy^2 &= y^3 - \frac{2000}{3} \quad (6) \end{aligned}$$

Solve for x :

$$x = \frac{y^3 - \frac{2000}{3}}{y^2} = y - \frac{2000}{3y^2} \quad (7)$$

Substitute $x = y - \frac{2000}{3y^2}$ into the constraint $x^3 + y^3 = 2000$:

$$\left(y - \frac{2000}{3y^2}\right)^3 + y^3 = 2000$$

This equation is highly complex and does not yield to simple algebraic methods. Therefore, we proceed with an approximate numerical solution. For more detailed computations, refer to the [Wolfram Alpha Lagrange Multipliers](#) tool.

4. Approximate Solution:

The approximate solution is:

$$(x, y) \approx (7.04423, 11.8177)$$

5. Decision on Integer Production Levels:

Since production levels must be integer values, we consider the nearest integer values to the approximate solution. The four possible combinations are:

- Option 1: $x = 7, y = 11$

$$x^3 + y^3 = 343 + 1331 = 1674 \leq 2000 \quad (\text{Satisfies the constraint})$$

- Option 2: $x = 7, y = 12$

$$x^3 + y^3 = 343 + 1728 = 2071 > 2000 \quad (\text{Does not satisfy the constraint})$$

- Option 3: $x = 8, y = 11$

$$x^3 + y^3 = 512 + 1331 = 1843 \leq 2000 \quad (\text{Satisfies the constraint})$$

- Option 4: $x = 8, y = 12$

$$x^3 + y^3 = 512 + 1728 = 2240 > 2000 \quad (\text{Does not satisfy the constraint})$$

Therefore, only Option 1 and Option 3 satisfy the constraint.

Compute the Total Cost for Each Option:

- Option 1: $x = 7, y = 11$

$$C(7, 11) = 7^2 \times 11 + 11^3 = 49 \times 11 + 1331 = 539 + 1331 = 1870$$

- Option 3: $x = 8, y = 11$

$$C(8, 11) = 8^2 \times 11 + 11^3 = 64 \times 11 + 1331 = 704 + 1331 = 2035$$

Decision: Choose the production levels that satisfy the constraint and incur the lesser cost. Between Option 1 and Option 3:

$$1870 < 2035$$

Therefore, Option 1 ($x = 7, y = 11$) is the optimal integer production level with a total cost of \$1,870.

6. Visualization:

To better understand the optimization, let's visualize the nonlinear constraint, the approximate optimal point, the integer production options, and their associated costs.




Figure 3.30: Visualization of the optimization problem with nonlinear constraint. Red curve: $x^3 + y^3 = 2000$. Blue star: Chosen Option (7.04, 11.82). Blue circles: Options 1 & 3 (Inside Constraint). Red circles: Options 2 & 4 (Outside Constraint). Orange arrow: ∇C . Purple arrow: ∇g . Level Curve: $C(x, y) = 2237.10$.

Optimization with Multiple Equality Constraints

When optimizing a function $f(x, y, z)$ subject to two or more equality constraints, such as $g(x, y, z) = c_1$ and $h(x, y, z) = c_2$, the method of Lagrange multipliers naturally extends to accommodate additional constraints. The underlying principle remains the same: at an optimal point, the gradients of f , g , and h must align in such a way that any movement satisfying the constraints does not produce a first-order change in f .

For two constraints, we define the *Lagrange function* as:

$$\mathcal{L}(x, y, z, \lambda_1, \lambda_2) = f(x, y, z) + \lambda_1(g(x, y, z) - c_1) + \lambda_2(h(x, y, z) - c_2).$$

The gradients of $\mathcal{L}$ with respect to all variables must vanish at the extrema, leading to the system of equations:

$$\begin{aligned}\nabla f(x, y, z) - \lambda_1 \nabla g(x, y, z) - \lambda_2 \nabla h(x, y, z) &= \mathbf{0}, \\ g(x, y, z) &= c_1, \\ h(x, y, z) &= c_2.\end{aligned}$$

In practice:

1. Compute $\nabla f(x, y, z)$, $\nabla g(x, y, z)$, and $\nabla h(x, y, z)$.
2. Solve the system of equations for x, y, z, λ_1 , and λ_2 .
3. Evaluate $f(x, y, z)$ at the solutions to determine the extrema.

Generalization to n Equality Constraints

For a general optimization problem with n variables and m equality constraints:

$$\begin{aligned}\text{Minimize (or maximize): } & f(x_1, x_2, \dots, x_n), \\ \text{Subject to: } & g_1(x_1, x_2, \dots, x_n) = c_1, \\ & g_2(x_1, x_2, \dots, x_n) = c_2, \\ & \dots, \\ & g_m(x_1, x_2, \dots, x_n) = c_m,\end{aligned}$$

the Lagrange function is defined as:

$$\mathcal{L}(x_1, x_2, \dots, x_n, \lambda_1, \lambda_2, \dots, \lambda_m) = f(x_1, x_2, \dots, x_n) + \sum_{i=1}^m \lambda_i (g_i(x_1, x_2, \dots, x_n) - c_i).$$

The necessary conditions for an extremum are:

$$\begin{aligned}\nabla f(x_1, x_2, \dots, x_n) - \sum_{i=1}^m \lambda_i \nabla g_i(x_1, x_2, \dots, x_n) &= \mathbf{0}, \\ g_1(x_1, x_2, \dots, x_n) &= c_1, \\ g_2(x_1, x_2, \dots, x_n) &= c_2, \\ &\dots, \\ g_m(x_1, x_2, \dots, x_n) &= c_m.\end{aligned}$$

In the general case, the gradients $\nabla g_1, \nabla g_2, \dots, \nabla g_m$ define a space of feasible directions, and the gradient ∇f must align with this space at the extremum. Specifically, ∇f must lie in the span of $\nabla g_1, \nabla g_2, \dots, \nabla g_m$, reflecting the fact that movement within the feasible region does not change f to first order.

▲ **Example 3.8.5** A nutritionist wants to find the optimal levels x and y (in grams) of two ingredients to minimize a certain dietary risk function. Suppose the overall risk is modeled by:

$$f(x, y) = x^2 + xy + 2y^2.$$

They must satisfy two nutritional constraints to ensure dietary balance:

$$x + 2y = 10, \quad \text{and} \quad x - y = 3.$$

Solution

1. **Set up the Lagrangian:** Introduce Lagrange multipliers λ and μ for the two constraints:

$$\mathcal{L}(x, y, \lambda, \mu) = x^2 + xy + 2y^2 - \lambda(x + 2y - 10) - \mu(x - y - 3).$$

2. **Compute Partial Derivatives:** Take partial derivatives and set them equal to zero.

$$\frac{\partial \mathcal{L}}{\partial x} = 2x + y - \lambda - \mu = 0,$$

$$\frac{\partial \mathcal{L}}{\partial y} = x + 4y - 2\lambda + \mu = 0,$$

$$\frac{\partial \mathcal{L}}{\partial \lambda} = -(x + 2y - 10) = 0,$$

$$\frac{\partial \mathcal{L}}{\partial \mu} = -(x - y - 3) = 0.$$

3. **From the Constraints:**

$$x + 2y = 10, \quad x - y = 3.$$

Solve these directly first: From $x - y = 3$, we get $x = 3 + y$. Substitute into $x + 2y = 10$:

$$(3 + y) + 2y = 10 \implies 3 + 3y = 10 \implies 3y = 7 \implies y = \frac{7}{3}.$$

Thus, $x = 3 + \frac{7}{3} = \frac{16}{3}$.

4. **Check Optimality:** Substitute $(x, y) = (\frac{16}{3}, \frac{7}{3})$ into the objective:

$$f\left(\frac{16}{3}, \frac{7}{3}\right) = \left(\frac{16}{3}\right)^2 + \left(\frac{16}{3} \cdot \frac{7}{3}\right) + 2\left(\frac{7}{3}\right)^2.$$

Compute:

$$\left(\frac{16}{3}\right)^2 = \frac{256}{9}, \quad \frac{16}{3} \cdot \frac{7}{3} = \frac{112}{9}, \quad 2\left(\frac{7}{3}\right)^2 = 2\left(\frac{49}{9}\right) = \frac{98}{9}.$$

Sum them up:

$$f\left(\frac{16}{3}, \frac{7}{3}\right) = \frac{256}{9} + \frac{112}{9} + \frac{98}{9} = \frac{466}{9}.$$

The point $(\frac{16}{3}, \frac{7}{3})$ satisfies both constraints and yields the minimized risk value $\frac{466}{9}$.

Optimization with Inequality Constraints

Up until now, we have considered optimization problems with equality constraints, where conditions like $g(x, y) = c$ must be met. However, many real-world problems also include *inequality constraints*, such as $h(x, y) \geq d$. These constraints arise naturally in scenarios where certain values should not fall below (or rise above) a specified level—for example, ensuring that a production level does not drop below a minimum quota, or that a budget does not exceed a certain maximum amount.

To handle such conditions, mathematicians and engineers use a more general framework known as the **Karush-Kuhn-Tucker (KKT)** conditions. The KKT conditions extend the idea of Lagrange multipliers to cover both equality and inequality constraints. In essence, they incorporate additional variables (called KKT multipliers) that must satisfy certain non-negativity requirements and complementary conditions. The key addition is that when a constraint is not “active” (i.e., it does not exactly hold at its boundary), its corresponding multiplier must be zero, ensuring that non-binding constraints do not influence the solution.

Although we will not dive deeply into the full derivation, here are the general steps and guidelines for dealing with inequality constraints:

- **Formulate the Problem:** Start with your objective function $f(x, y)$ and identify any inequality constraints, such as $h(x, y) \geq d$.
- **Introduce KKT Multipliers:** Assign a KKT multiplier (often denoted μ) to each inequality constraint. These multipliers must be non-negative.
- **Set Up the Lagrangian-like Function:** Similar to the case of equality constraints, form a function incorporating both the original objective and the constraints, but now include terms $\mu \cdot [d - h(x, y)]$ that represent the inequality constraints.
- **Derive the First-Order Conditions:** Take partial derivatives of this augmented function with respect to your variables and set them to zero. You also include conditions that relate the sign of μ to whether the constraint is active.

- **Check Complementary Slackness:** The KKT framework requires that $\mu[d - h(x, y)] = 0$. This means:

$$\mu > 0 \implies h(x, y) = d \quad (\text{active constraint}),$$

$$\mu = 0 \implies h(x, y) > d \quad (\text{inactive constraint}).$$

- **Identify the Feasible and Optimal Points:** Solve the resulting system of equations and inequalities to find candidate solutions, then verify which candidate provides the best (minimum or maximum) objective value while respecting the constraints.

While we will not fully derive or prove the KKT conditions here, it is enough to recognize that they represent a natural extension of the methods we have discussed. When confronted with more intricate constraints—like ensuring $h(x, y) \geq 0$ —the KKT conditions guide us to solutions that honor these inequalities. Thus, just as Lagrange multipliers handle equality constraints elegantly, the KKT conditions handle both equality and inequality constraints, giving us a powerful toolbox to tackle a wide range of optimization problems.

Applications to Machine Learning

In machine learning, constraints can help guide the training of models in practical ways. For example, suppose you want to train a model that makes predictions under limited computational resources. You might enforce an inequality constraint that keeps the model complexity below a certain threshold. Or perhaps you need to ensure fairness, such as enforcing that the model's error rate for one demographic group does not exceed its error rate for another group by more than a certain amount. These kinds of constraints lead to inequality conditions that can be analyzed using the KKT framework. The underlying principle remains the same: find model parameters that minimize a loss function while respecting the given inequalities.

▲ **Example 3.8.6** Consider a simplified Support Vector Machine (SVM) scenario where we aim to find the optimal parameter w that separates two data points with labels $y_1 = 1$ and $y_2 = -1$. Suppose the data points are $x_1 = 2$ and $x_2 = -1$. The objective is to maximize the margin between the two points, which can be formulated as minimizing the function:

$$f(w) = \frac{1}{2}w^2,$$

subject to the following inequality constraints ensuring that each point is correctly classified with a margin of at least 1:

$$y_1(w \cdot x_1) \geq 1 \quad \text{and} \quad y_2(w \cdot x_2) \geq 1.$$

Substituting the given values:

$$1 \cdot (w \cdot 2) \geq 1 \implies 2w \geq 1,$$

$$-1 \cdot (w \cdot (-1)) \geq 1 \quad \Rightarrow \quad w \geq 1.$$

Solution We aim to minimize $f(w) = \frac{1}{2}w^2$ subject to the constraints:

$$2w \geq 1 \quad \text{and} \quad w \geq 1.$$

These can be rewritten as:

$$h_1(w) = 2w - 1 \geq 0, \quad h_2(w) = w - 1 \geq 0.$$

1. **Formulate the Lagrangian:** Introduce KKT multipliers μ_1 and μ_2 for the two inequality constraints:

$$\mathcal{L}(w, \mu_1, \mu_2) = \frac{1}{2}w^2 - \mu_1(2w - 1) - \mu_2(w - 1).$$

2. **Derive the First-Order Conditions:** Take the derivative of $\mathcal{L}$ with respect to w and set it to zero:

$$\frac{\partial \mathcal{L}}{\partial w} = w - 2\mu_1 - \mu_2 = 0 \quad \Rightarrow \quad w = 2\mu_1 + \mu_2.$$

3. **Complementary Slackness Conditions:**

$$\mu_1(2w - 1) = 0, \quad \mu_2(w - 1) = 0.$$

$$\mu_1 \geq 0, \quad \mu_2 \geq 0.$$

4. **Analyze Possible Cases:**

- **Case 1: Both Constraints are Active**

If both constraints are active, then:

$$2w - 1 = 0 \quad \Rightarrow \quad w = \frac{1}{2},$$

$$w - 1 = 0 \quad \Rightarrow \quad w = 1.$$

This leads to a contradiction since w cannot be both $\frac{1}{2}$ and 1. Therefore, this case is not possible.

- **Case 2: Only $h_1(w)$ is Active**

If $h_1(w)$ is active, then $2w - 1 = 0$ and $w > 1$ (since $h_2(w)$ is inactive).

$$w = \frac{1}{2}.$$

However, $\frac{1}{2} < 1$, which violates $h_2(w) \geq 0$. Therefore, this case is also not possible.

- **Case 3: Only $h_2(w)$ is Active**

If $h_2(w)$ is active, then $w - 1 = 0$ and $2w - 1 > 0$:

$$w = 1.$$

Check $h_1(w)$:

$$2(1) - 1 = 1 \geq 0 \quad (\text{satisfied}).$$

Substitute $w = 1$ into the first-order condition:

$$1 = 2\mu_1 + \mu_2.$$

Since $h_1(w)$ is inactive ($\mu_1 = 0$):

$$1 = 0 + \mu_2 \quad \Rightarrow \quad \mu_2 = 1.$$

$$\mu_1 = 0, \quad \mu_2 = 1.$$

- **Case 4: Both Constraints are Inactive**

If both constraints are inactive, then $\mu_1 = 0$ and $\mu_2 = 0$. The first-order condition becomes:

$$w = 0.$$

However, check the constraints:

$$2(0) - 1 = -1 < 0 \quad \text{and} \quad 0 - 1 = -1 < 0,$$

which violate both constraints. Therefore, this case is not feasible.

5. Determine the Optimal Solution:

The only feasible case is **Case 3**, where:

$$w = 1, \quad \mu_1 = 0, \quad \mu_2 = 1.$$

6. Verify the Solution:

Substitute $w = 1$ back into the objective function:

$$f(1) = \frac{1}{2}(1)^2 = \frac{1}{2}.$$

This is the minimum value of $f(w)$ subject to the given constraints.

Conclusion: The optimal parameter is $w = 1$, which satisfies the inequality constraints $2w \geq 1$ and $w \geq 1$. This solution minimizes the objective function while ensuring that both constraints are respected. In this simplified SVM scenario, $w = 1$ represents the parameter that maximizes the margin between the two data points while correctly classifying them.

This simplified scenario captures the core idea: if unconstrained optimization leads us outside the allowed set, inequality constraints force us to choose the best feasible point. More sophisticated methods, such as the KKT conditions, allow us to systematically solve problems with multiple and more complex inequalities, a crucial step in handling real-world optimization tasks in fields like machine learning.

Formulas and Definitions

Important Vocabulary

- **Function of Two Variables:** A rule that assigns a real number $z = f(x, y)$ to each ordered pair (x, y) in a subset of $\mathbb{R}^2$. The graph is a surface in $\mathbb{R}^3$.
- **Domain:** The set of all input pairs (x, y) where the function $f(x, y)$ is defined.
- **Range:** The set of all output values z produced by $f(x, y)$.
- **Level Curves:** Curves in the xy -plane where $f(x, y) = k$ for constant k .
- **Cross-Sections:** Slices of the surface $z = f(x, y)$ obtained by fixing one variable x or y .
- **Level Surfaces:** Surfaces in $\mathbb{R}^3$ where $f(x, y, z) = k$ for constant k .
- **Limit:** For a function $f(x, y)$, the limit at (a, b) is the value L such that $f(x, y)$ approaches L as $(x, y) \rightarrow (a, b)$ from any direction.
- **Path Dependence:** A limit does not exist if the value of $f(x, y)$ depends on the path taken to approach (a, b) .
- **Continuity:** $f(x, y)$ is continuous at (a, b) if:

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b).$$

- **Partial Derivative:** The derivative of a multivariable function with respect to one variable, holding all others constant.
- **Clairaut's Theorem:** States that mixed partial derivatives are equal when certain continuity conditions are met.
- **Implicit Differentiation:** A technique for finding derivatives when relationships between variables are defined implicitly
- **Directional Derivative:** The rate of change of a function in a specific direction, quantified using a unit vector
- **Gradient Vector:** A vector field containing all partial derivatives, pointing in direction of steepest increase
- **Tangent Plane:** A plane that touches a surface at a point and provides the best linear approximation near that point
- **Differentials:** Small increments in variables used to approximate changes in functions
- **Critical Point:** A point where all partial derivatives are zero or at least one does not exist

- **Local Maximum/Minimum:** A point where function value is higher/lower than all nearby points
- **Global Maximum/Minimum:** A point where function value is highest/lowest over entire domain
- **Saddle Point:** A critical point that is neither a maximum nor minimum
- **Hessian Matrix:** Matrix of all second partial derivatives of a function
- **Discriminant:** Determinant of the Hessian matrix in two dimensions
- **Constraint:** A condition that limits the possible values of variables in optimization
- **Lagrange Multiplier:** A variable (λ) introduced to solve constrained optimization problems

Key Formulas

Function of Two Variables:

$$z = f(x, y), \quad (x, y) \in \mathbb{R}^2.$$

Domain:

$$\text{dom } f = \{(x, y) \in \mathbb{R}^2 \mid \text{restrict } x \text{ and } y\}.$$

Range:

$$\text{range } f = \{z \in \mathbb{R} \mid (x, y) \in \text{dom } f\}.$$

Level Curves:

$$f(x, y) = k, \quad \text{for constant } k.$$

Level Surfaces:

$$f(x, y, z) = k, \quad \text{for constant } k.$$

Hyperplanes

$$a_1x_1 + a_2x_2 + \cdots + a_nx_n = c.$$

Hyperspheres

$$x_1^2 + x_2^2 + \cdots + x_n^2 = r^2.$$

Formal Definition of a Limit:

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$$

if for all $\epsilon > 0$, there exists $\delta > 0$ such that:

$$0 < \sqrt{(x-a)^2 + (y-b)^2} < \delta \\ \implies |f(x, y) - L| < \epsilon.$$

Continuity:

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = f(a, b).$$

Generalized Limits:

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} f(\mathbf{x}) = L$$

if for all $\epsilon > 0$, there exists $\delta > 0$ such that:

$$0 < \|\mathbf{x} - \mathbf{a}\| < \delta \implies |f(\mathbf{x}) - L| < \epsilon.$$

Partial Derivative with respect to $\mathbf{x}$:

$$\frac{\partial f}{\partial x}(x, y) = \lim_{h \rightarrow 0} \frac{f(x+h, y) - f(x, y)}{h}$$

Clairaut's Theorem: If all second-order partial derivatives are continuous:

$$\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial^2 f}{\partial y \partial x}$$

Multivariable Chain Rule: For $z = 2D$ Discriminant:

$f(x, y)$ where x, y depend on t :

$$\frac{dz}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$$

$$D = f_{xx}f_{yy} - (f_{xy})^2$$

3D Determinant:

General Chain Rule: For $z = f(x_1, \dots, x_n)$

$$\frac{dz}{dt} = \sum_{i=1}^n \frac{\partial f}{\partial x_i} \frac{dx_i}{dt}$$

$$D_3 = \det(H) = f_{xx}(f_{yy}f_{zz} - f_{yz}^2) - f_{xy}(f_{yx}f_{zz} - f_{yz}f_{zx}) + f_{xz}(f_{yx}f_{zy} - f_{yy}f_{zx})$$

Leibniz Integral Rule:

$$\begin{aligned} \frac{d}{dx} \int_{a(x)}^{b(x)} f(x, t) dt &= f(x, b(x)) \frac{db}{dx} \\ &\quad - f(x, a(x)) \frac{da}{dx} \\ &\quad + \int_{a(x)}^{b(x)} \frac{\partial f}{\partial x} dt \end{aligned}$$

Directional Derivative Definition:

$$\begin{aligned} D_{\mathbf{u}}f(\mathbf{p}) &= \left. \frac{d}{dh} f(\mathbf{p} + h\mathbf{u}) \right|_{h=0} \\ &= \nabla f(\mathbf{p}) \cdot \mathbf{u} \end{aligned}$$

Gradient Vector:

$$\nabla f(\mathbf{p}) = \begin{bmatrix} \frac{\partial f}{\partial x_1}(\mathbf{p}) \\ \frac{\partial f}{\partial x_2}(\mathbf{p}) \\ \vdots \\ \frac{\partial f}{\partial x_n}(\mathbf{p}) \end{bmatrix}$$

Tangent Plane:

$$z = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$$

Hessian Matrix: For $f: \mathbb{R}^n \rightarrow \mathbb{R}$:

$$H(\mathbf{a}) = \begin{bmatrix} f_{x_1x_1} & f_{x_1x_2} & \cdots \\ f_{x_2x_1} & f_{x_2x_2} & \cdots \\ \vdots & \vdots & \ddots \end{bmatrix}$$

Second Derivative Test (2D): At critical point (a, b) :

- If $D > 0$ and $f_{xx} > 0$: Local minimum
- If $D > 0$ and $f_{xx} < 0$: Local maximum
- If $D < 0$: Saddle point
- If $D = 0$: Inconclusive

Lagrange Multiplier (One Constraint):

For $f: \mathbb{R}^n \rightarrow \mathbb{R}$ subject to $g(x_1, \dots, x_n) = c$:

$$\mathcal{L}(\mathbf{x}, \lambda) = f(\mathbf{x}) + \lambda(g(\mathbf{x}) - c)$$

With necessary condition:

$$\nabla f = \lambda \nabla g$$

Lagrange Multipliers (Multiple Constraints):

For $f: \mathbb{R}^n \rightarrow \mathbb{R}$ subject to m constraints $g_i(\mathbf{x}) = c_i$:

$$\mathcal{L}(\mathbf{x}, \lambda_1, \dots, \lambda_m) = f(\mathbf{x}) + \sum_{i=1}^m \lambda_i(g_i(\mathbf{x}) - c_i)$$

With necessary condition:

$$\nabla f = \sum_{i=1}^m \lambda_i \nabla g_i$$

Practice Problems

1. The monthly payment M for a mortgage depends on the principal P , annual interest rate r , and number of years t according to:

$$M = f(P, r, t) = P \frac{r(1+r)^{12t}}{12((1+r)^{12t} - 1)}$$

Calculate $f(300000, 0.06, 30)$ to find the monthly payment on a \$300,000 loan at 6% annual interest for 30 years.

2. Sketch all level curves of $f(x, y) = x^2 + y^2$ for $k = 1, 2, 4$.
3. Find the domain of $f(x, y) = \ln(x^2 + y^2 - 4)$.
4. Describe and sketch the level surfaces of $f(x, y, z) = x^2 + y^2 + z^2$.
5. Find the domain and range of $f(x, y) = \sqrt{9 - x^2 - y^2}$.
6. Determine if the following parametric curve lies on the given surface: $\mathbf{r}(t) = \langle t, t^2, t^3 \rangle$ on $z = y^{3/2}$.
7. Find and sketch the domain of $f(x, y) = \sqrt{\frac{x}{y-1}}$.
8. Look around at the room you are sitting in. Sketch the schematic of just the walls and guess the measurements of height and width. Assume you want to paint the walls a different color and let that paint cost $\$P$ per square meter. Write a function that represents the cost of painting your room given any paint. Don't forget to buy primer too!
9. The thrust T of a rocket engine depends on the mass flow rate $\dot{m}$ and exhaust velocity v_e according to:

$$T = f(\dot{m}, v_e) = \dot{m}v_e + (P_e - P_a)A_e$$

where P_e is exhaust pressure, P_a is atmospheric pressure, and A_e is exhaust area. If $P_e = 101$ kPa, $P_a = 100$ kPa, and $A_e = 2\text{m}^2$, express the mass flow rate $\dot{m}$ as a function of just T and v_e .

10. The profit P for a food truck selling two types of tacos depends on the number of each type sold (x and y) according to:

$$P = f(x, y) = (4x+5y) - (2x+3y) - 100$$

where the first term is revenue, the second term is cost of ingredients, and 100 represents fixed costs. Find $f(50, 30)$ to determine the profit from selling 50 of type 1 and 30 of type 2. Then, assuming that local taxes are 6%, reconstruct the function for profit including taxes (remember there is no tax on debits).

11. Evaluate $\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2+y^2}$ if it exists.
12. Evaluate $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2y}{x^2+y^2}$ if it exists.
13. Test the continuity of $f(x, y) = \frac{xy^2}{x^2+y^4}$ at $(0, 0)$.
14. Find $\lim_{(x,y) \rightarrow (0,0)} \frac{x^2-y^2}{x^2+y^2}$.
15. Prove that $f(x, y) = \frac{x^3y}{x^4+y^2}$ is continuous at $(0, 0)$.
16. Evaluate $\lim_{(x,y) \rightarrow (1,2)} \frac{\sin(xy-2)}{x-1}$.
17. In fluid dynamics, the velocity potential of a source flow is given by

$$\phi(x, y) = \frac{Q}{2\pi} \ln(x^2 + y^2)^{1/2}$$

where Q is the source strength. Show that $\lim_{(x,y) \rightarrow (0,0)} \phi(x, y) = -\infty$.

18. The electric field of a point charge is given by

$$\mathbf{E}(x, y) = \frac{1}{4\pi\epsilon_0} \frac{q[(x - x_0)\mathbf{i} + (y - y_0)\mathbf{j}]}{[(x - x_0)^2 + (y - y_0)^2]^{3/2}}$$

Find $\lim_{(x,y) \rightarrow (0,0)} |E(x, y)|$.

19. For a spinning disk of radius R , the stress function is given by

$$\sigma(r, \theta) = \frac{A}{r^2} + Br^2$$

where $r = \sqrt{x^2 + y^2}$. Evaluate $\lim_{(x,y) \rightarrow (0,0)} \sigma(r, \theta)$ if it exists.

20. Find f_x and f_y for $f(x, y) = x^2y + 3xy^2$.
21. Compute all second partial derivatives of $f(x, y) = \sin(xy)e^x$.
22. Find all mixed partial derivatives of $f(x, y, z) = xyz + e^{xy}$.
23. Verify Clairaut's theorem for $f(x, y) = x^2e^y + \ln(xy)$.
24. Find all second-order partial derivatives of $f(x, y) = \arctan\left(\frac{y}{x}\right)$.
25. Show that $f(x, y) = \frac{xy}{x^2 + y^2}$ if $(x, y) \neq (0, 0)$, $f(0, 0) = 0$ has all partial derivatives at $(0, 0)$ but is not differentiable there.
26. For the function $f(x, y) = x^y$, show that:

$$y \frac{\partial f}{\partial x} = x \frac{\partial f}{\partial y}$$

This is known as Euler's homogeneous differential equation.

27. Let $f(x, y)$ be twice differentiable everywhere. If $f_{xx} + f_{yy} = 0$, show that $u = f_x$ and $v = -f_y$ satisfy:

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} = 0 \quad \text{and} \quad \frac{\partial u}{\partial y} = \frac{\partial v}{\partial x}$$

These are known as the Cauchy-Riemann equations.

28. If $f(x, y)$ satisfies $xf_x + yf_y = nf$ for some constant n , prove that $f(tx, ty) = t^n f(x, y)$ for any $t > 0$.

29. For the implicitly defined function $y(x)$ given by $x^2 + y^2 = \sin(xy)$, show that:

$$\left. \frac{d^2y}{dx^2} \right|_{(0,0)} = -1$$

30. If z is implicitly defined by $e^{xyz} = xyz + 1$, prove that:

$$\frac{\partial z}{\partial x} \cdot \frac{\partial z}{\partial y} \cdot \frac{\partial z}{\partial z} = -1$$

at any point where $xyz = 0$.

31. Find $\frac{dz}{dt}$ if $z = x^2 + y^2$ where $x = t \cos t$, $y = t \sin t$.
32. Use implicit differentiation to find $\frac{dy}{dx}$ if $x^2y + y^3 = 8$.
33. Find $\frac{\partial z}{\partial s}$ and $\frac{\partial z}{\partial t}$ if $z = xy$ where $x = s^2t$, $y = st^2$.
34. Find $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$ if $x^2 + y^2 + z^2 = 16$ defines z implicitly.
35. Use the Chain Rule to find $\frac{\partial z}{\partial x}$ if $z = f(u, v)$ where $u = xy$ and $v = \frac{y}{x}$.
36. Calculate $\frac{d}{dx} \int_0^x te^{xt} dt$ using the Leibniz integral rule.

37. Let $\mathbf{r}(t) = \langle t^2, \sin t, e^t \rangle$ be a vector-valued function.

- (a) Compute the derivative $\mathbf{r}'(t)$.
- (b) Find the magnitude of the derivative vector at $t = \pi$.

38. Consider the function $f(x, y) = \ln(xy) + e^{x^2y}$.

- (a) Compute the total differential df at the point $(1, 1)$.

- (b) Use the total differential to approximate the change in f when x changes to 1.01 and y changes to 0.99.
39. Let $x(t)$ and $y(t)$ be differentiable functions of t , and let
- $$f(x, y) = \sum_{k=1}^3 \sin(x^k y^{3-k}).$$
- Use the Chain Rule to compute the derivative $\frac{df}{dt}$.
40. Find ∇f for $f(x, y) = x^2 y + y^3$ at $(1, 2)$.
41. Find the directional derivative of $f(x, y) = x e^y$ in the direction of $\mathbf{v} = \langle 3, 4 \rangle$ at $(1, 0)$.
42. Find the direction of steepest increase of $f(x, y) = 2x^2 + 3y^2$ at $(1, 1)$.
43. Calculate $D_{\mathbf{u}} f$ for $f(x, y, z) = xyz$ in direction $\mathbf{u} = \langle 1, 1, 1 \rangle$ at $(1, 2, 3)$.
44. Find the gradient vector field of $f(x, y) = \sin(xy)$ and evaluate it at $(\pi, 0)$.
45. Find the directional derivative of $f(x, y) = \ln(x^2 + y^2)$ in the direction from $(1, 1)$ to $(2, 3)$.
46. A particle moves in a two-dimensional plane where the temperature at any point (x, y) is given by the scalar field $T(x, y) = e^{-x^2 - y^2}$. Determine the direction of steepest temperature increase at the point $(1, 1)$.
47. In a simple linear regression model, the cost function is defined as

$$J(w, b) = \frac{1}{2}(wx + b - y)^2,$$

where w and b are parameters, and (x, y) is a single data point. Compute the partial derivatives $\frac{\partial J}{\partial w}$ and $\frac{\partial J}{\partial b}$.

48. Find the tangent plane to $z = x^2 + y^2$ at $(1, 1, 2)$.
49. Use differentials to approximate $\sqrt{4.1^2 + 2.9^2}$.
50. Find the linear approximation of $f(x, y) = e^x \sin(y)$ near $(0, 0)$.
51. Find the equation of the tangent plane to $z = \ln(xy)$ at $(1, 1, 0)$.
52. Use linear approximation to estimate $f(1.1, 0.9)$ where $f(x, y) = \sqrt{x^2 + y^2}$ near $(1, 1)$.
53. Find the tangent plane to the surface $z = x e^y$ at point $(1, 0, 1)$ and determine if the surface lies above or below the plane near this point.
54. Find all critical points of $f(x, y) = x^2 + y^2 - 4x - 6y$.
55. Find and classify all critical points of $f(x, y) = x^2 y - \frac{y^3}{3}$.
56. Find the absolute maximum and minimum of $f(x, y) = x^2 + y^2$ on $x^2 + y^2 \leq 4$.
57. For $f(x, y) = x e^y - x^2 - y^2$, find and classify all critical points.
58. Find all local maxima, minima, and saddle points of $f(x, y) = xy - x^2 - y^2$.
59. Show that $f(x, y) = (y - x^2)(y - 2x^2)$ has exactly three critical points and classify each.
60. Let $f(x, y, z) = x^2 + y^2 + z^2 - 4x - 6y + 8z$.
- (a) Find all critical points of f .
- (b) Determine whether each critical point is a local maximum, local minimum, or saddle point using the second derivative test.

61. Let $f(\mathbf{x}) = f(x_1, x_2, \dots, x_n)$ be a twice-differentiable function where $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$. Explain the conditions under which a critical point $\mathbf{x}_0$ of f is a local minimum, local maximum, or saddle point. How do these conditions extend from the two-variable case to n variables?
62. Find the shortest distance from the point $(2, 1, 3)$ to the plane $x + y + z = 1$.
63. Find the dimensions of a box with volume 100 cubic units that has minimum surface area.
64. Find the points on the curve $y = x^2$ that are closest to the point $(0, 1)$.
65. Minimize $f(x, y) = x^2 + y^2$ subject to the constraint $x^3 + y^3 = 8$.
66. Use Lagrange multipliers to find the maximum and minimum values of $f(x, y) = xy$ subject to $x^2 + y^2 = 16$.
67. Consider the function $f(x, y, z) = x^2 + y^2 - z^2$ subject to the constraint $x + y + z = 1$.
- Use the method of Lagrange multipliers to find all critical points of f under the given constraint.
 - Classify each critical point as a local maximum, local minimum, or saddle point.
68. Find the maximum and minimum values of $f(x, y, z) = x + y + z$ subject to $x^2 + y^2 + z^2 = 1$.
69. Maximize the function $f(x, y) = 3x + 4y$ subject to the inequality constraints:
- $$\begin{cases} x + 2y \geq 8 \\ x - y \geq 2 \end{cases}$$
70. Minimize the function $f(x, y, z) = x^2 + y^2 + z^2$ subject to the inequality constraints:
- $$\begin{cases} x + y + z \geq 6 \\ x \geq 1 \\ y \geq 2 \\ z \geq 3 \end{cases}$$
71. Consider the optimization problem:
- Minimize $f(x, y) = x^2 + y^2$
- subject to the inequality constraints:
- $$\begin{cases} g_1(x, y) = x + y - 4 \geq 0 \\ g_2(x, y) = x - 2y + 1 \geq 0 \end{cases}$$
- Apply the Karush-Kuhn-Tucker (KKT) conditions to find all candidate points for the minimum of $f(x, y)$.
 - Determine which of these candidates are actual minima by verifying the KKT conditions.

Chapter 4

Multiple Integrals

Introduction

In our study of multivariable calculus, we now embark on an essential journey into the realm of multiple integrals. Building upon the concepts of single-variable integration and partial derivatives, we delve into methods for calculating areas, volumes, and other quantities that arise in higher-dimensional spaces. While integration in one dimension involves summing over intervals, multiple integration generalizes this concept to two and three dimensions. This allows us to analyze diverse real-world phenomena, from calculating the mass of an irregularly shaped object with varying density to determining the volume of regions bounded by complex surfaces. But all of this is not as scary as it sounds—we are just adding bits of things up in a smart way, that’s all.

We begin by exploring double integrals, starting with rectangular regions and expanding to general regions, incorporating techniques such as switching the order of integration and converting to polar coordinates. These tools provide a framework for solving problems in physics, engineering, and beyond. The power of double integrals becomes evident when applied to compute areas, volumes, and surface areas, or to study distributions such as mass and center of mass. From there, we extend to triple integrals, which enable us to work in three-dimensional regions, using cylindrical and spherical coordinate systems to simplify calculations in symmetric or otherwise complex geometries. These integrals offer insight into phenomena ranging from gravitational fields to fluid dynamics. Finally, we generalize further by introducing the concept of variable transformation with Jacobian determinants, a powerful technique for tackling integrals over irregular regions.

This chapter equips us with the tools to move seamlessly between dimensions, solving problems that are integral to both theoretical mathematics and practical applications. By mastering multiple integration, we take a significant step forward in understanding the underlying structures of the physical world and the mathematical framework that models it.

Lecture 4.1: Double Integrals over Rectangular Regions

In our study of single-variable calculus, we encountered the definite integral as a powerful tool for computing accumulated change. Given a function $f(x)$, we interpreted $\int_a^b f(x) dx$ as the signed area between the curve and the x -axis—a measurement of how a quantity accumulates along a one-dimensional interval. But what happens when we want to measure accumulation over a two-dimensional region?

Consider, for instance, the temperature distribution across a metal plate, the density variation in a thin sheet of material, or the pressure distribution across a surface. These scenarios involve quantities that vary with both x and y coordinates. How might we extend our understanding of integration to capture such two-dimensional variation?

Just as we built the single integral from the concept of adding up infinitesimally thin rectangles, we can imagine building a two-dimensional integral by adding up infinitesimally small rectangular pieces of a surface. This generalization leads us to the concept of a double integral—a fundamental tool for analyzing change across two-dimensional regions.




Figure 4.1: *The integral extends naturally to higher dimensions. Above we see the volume under the surface: the multi-variate analog to the area under a curve.*

Definition and Interpretation

Let's begin with the simplest case: a continuous function $f(x, y)$ defined over a rectangular region $R = [a, b] \times [c, d]$ in the xy -plane. What might it mean to integrate this function over R ? In single-variable calculus, our integrand $f(x)$ traced out a curve above the x -axis, and we found the area beneath it. Here, our function $f(x, y)$ traces out a surface above the xy -plane, suggesting that our integral should measure the volume beneath this surface.

We denote this double integral as:

$$\iint_R f(x, y) dA,$$

where dA represents an infinitesimal element of area, just as dx represented an infinitesimal length in single-variable calculus. This notation, with its double integral sign, reflects the two-dimensional nature of our accumulation. Note that we will begin writing a set as a subscript to the integral to indicate the bounds of integration. We will begin using this more, and using either sets or product sets listed as A, B, R, R_i , or $[a, b], [2, 5], (-\infty, 4)$, or $A \times B, [a, b] \times [c, d]$ or $[-1, 1]^2$ and so on.

But what exactly are we accumulating? The value $f(x, y)$ at each point can be thought of as a “height” above the xy -plane, and when we multiply this height by a small area dA , we get a small volume element. The double integral sums up all these infinitesimal volumes over the region R , giving us the total signed volume between the surface $z = f(x, y)$ and the xy -plane.

To make this intuition precise, we need to follow the same path that led us to the Riemann integral in single-variable calculus. How might we approximate this volume before taking a limit? Just as we divided an interval into subintervals, we can partition our rectangle R into smaller rectangles and approximate the volume above each piece.

Construction via Riemann Sums

Recall how we constructed the single integral: we divided our interval into smaller pieces, chose a sample point in each piece, multiplied by the width of the piece, and took the limit as the pieces became infinitesimally small. For a double integral, we follow a similar path, but now we’re working with two-dimensional pieces.

Let’s construct this systematically. First, we’ll partition our rectangular region R into a grid of smaller rectangles. How do we do this? We can divide each side of our rectangle independently:

- Partition $[a, b]$ into n equal subintervals, each of width $\Delta x = \frac{b-a}{n}$
- Partition $[c, d]$ into m equal subintervals, each of height $\Delta y = \frac{d-c}{m}$

This creates a grid of nm small rectangles, which we’ll call R_{ij} . Each has area $\Delta A = \Delta x \Delta y$. Just as we chose a sample point in each subinterval for single integrals, here we choose a sample point (x_{ij}^*, y_{ij}^*) in each small rectangle.

Now we can build our approximation. Above each small rectangle R_{ij} , we have a approximately rectangular solid with:

- Base area: $\Delta A = \Delta x \Delta y$
- Height: $f(x_{ij}^*, y_{ij}^*)$

- Volume: $f(x_{ij}^*, y_{ij}^*)\Delta A$

Adding up all these small volumes gives us our double Riemann sum:

$$S_{n,m} = \sum_{i=1}^n \sum_{j=1}^m f(x_{ij}^*, y_{ij}^*)\Delta A$$

Just as the definite integral emerged as the limit of Riemann sums, we define the double integral as the limit of these double Riemann sums as our grid becomes infinitely fine:

$$\iint_R f(x, y) dA = \lim_{n,m \rightarrow \infty} S_{n,m} = \lim_{n,m \rightarrow \infty} \sum_{i=1}^n \sum_{j=1}^m f(x_{ij}^*, y_{ij}^*)\Delta A$$

This limit, when it exists, gives us the precise value of the double integral. For continuous functions (and even some discontinuous ones), this limit exists and is independent of how we choose our sample points (x_{ij}^*, y_{ij}^*) —a remarkable fact that parallels the single-variable case.

This construction gives us more than just a definition; it provides a concrete way to visualize what a double integral actually computes. We're literally adding up infinitely many infinitesimally small volumes, just as a single integral adds up infinitely many infinitesimally thin rectangles. The below figure is the 3D representation of the Riemann sums under a surface:




Figure 4.2: *The Riemann Sum approximation of the volume under a curve. When taking the limit of these rectangular prisms to be infinitesimally small, we approximate the double integral.*

It is important to also note, that just as we have the idea of **negative area** in single variable calculus when the curve dips below the x -axis, we also have the analogous **negative volume** when our surface dips below the xy -plane.

▲ **Example 4.1.1** Approximate the double integral $\iint_R f(x, y) dA$ where $f(x, y) = x^2 + y^2$ over the rectangular region $R = [0, 2] \times [0, 1]$ using a Riemann sum with $n = 2$ subintervals in x and $m = 2$ subintervals in y . Use the upper-right corner of each subrectangle as the sample point.

Solution Let's construct this approximation systematically:

1. First, let's determine our partition dimensions:

$$\Delta x = \frac{2 - 0}{2} = 1, \quad \Delta y = \frac{1 - 0}{2} = \frac{1}{2}$$

So each subrectangle has area:

$$\Delta A = \Delta x \Delta y = (1) \left(\frac{1}{2} \right) = \frac{1}{2}$$

2. The x -coordinates of our partition are:

$$x_0 = 0, \quad x_1 = 1, \quad x_2 = 2$$

The y -coordinates are:

$$y_0 = 0, \quad y_1 = \frac{1}{2}, \quad y_2 = 1$$

3. Using upper-right corners as sample points, our four subrectangles have points:

$$R_{11} : (x_{11}^*, y_{11}^*) = \left(1, \frac{1}{2} \right)$$

$$R_{12} : (x_{12}^*, y_{12}^*) = (1, 1)$$

$$R_{21} : (x_{21}^*, y_{21}^*) = \left(2, \frac{1}{2} \right)$$

$$R_{22} : (x_{22}^*, y_{22}^*) = (2, 1)$$

4. Evaluate $f(x, y) = x^2 + y^2$ at each sample point:

$$f\left(1, \frac{1}{2}\right) = 1 + \frac{1}{4} = \frac{5}{4}$$

$$\begin{aligned}
 f(1, 1) &= 1 + 1 = 2 \\
 f\left(2, \frac{1}{2}\right) &= 4 + \frac{1}{4} = \frac{17}{4} \\
 f(2, 1) &= 4 + 1 = 5
 \end{aligned}$$

5. Form the double Riemann sum:

$$\begin{aligned}
 S_{2,2} &= \sum_{i=1}^2 \sum_{j=1}^2 f(x_{ij}^*, y_{ij}^*) \Delta A \\
 &= \frac{1}{2} \left[f\left(1, \frac{1}{2}\right) + f(1, 1) + f\left(2, \frac{1}{2}\right) + f(2, 1) \right] \\
 &= \frac{1}{2} \left[\frac{5}{4} + 2 + \frac{17}{4} + 5 \right] \\
 &= \frac{1}{2} \left[\frac{5 + 8 + 17 + 20}{4} \right] \\
 &= \frac{50}{8} = \frac{25}{4} = 6.25
 \end{aligned}$$

Therefore, our Riemann sum approximation is $\frac{25}{4}$ or approximately 6.25 square units. This is a rough approximation; the actual value of the double integral would be obtained by taking the limit as $n, m \rightarrow \infty$.

Note: This example illustrates how even a coarse partition (2×2) involves significant computation. The true power of double integrals emerges when we develop more sophisticated techniques (like Fubini's Theorem) to evaluate them exactly rather than approximating with Riemann sums.

Types of Riemann Sums

Just as in single-variable calculus, where we could choose different sample points in each subinterval (left endpoint, right endpoint, midpoint), we have analogous choices for double integrals. Recall from Chapter 1 Lecture 1 that we listed different approximation techniques: left-hand sums tended to overestimate decreasing functions, right-hand sums underestimated them, and midpoint sums often provided better approximations.

For double integrals, we have similar options for choosing our sample points (x_{ij}^*, y_{ij}^*) in each subrectangle R_{ij} :

- **Lower-Left Corner:** Take (x_{ij}^*, y_{ij}^*) as the lower-left vertex of each R_{ij}
- **Upper-Right Corner:** Use the upper-right vertex of each R_{ij}

- **Center Point:** Choose the center point of each R_{ij}
- **Supremum/Infimum:** Select points where f achieves its maximum/minimum value in R_{ij}

Just as with single integrals, these choices can lead to over- or under-approximations. For instance, if $f(x, y)$ is increasing in both variables, the lower-left corners will give an underestimate, while upper-right corners will give an overestimate. The center point method often provides a better approximation, as it tends to balance out these extremes.

Importantly, for continuous functions, all these methods converge to the same limit as the mesh of our partition becomes infinitely fine. This convergence mirrors the behavior we saw with single integrals and reinforces a fundamental principle: while the choice of sample points affects our approximations, it does not affect the final value of the integral.

Fubini's Theorem and Iterated Integrals

Now that we understand what a double integral represents and how to approximate it, a natural question arises: how do we actually compute one? The Riemann sum definition, while theoretically important, isn't practical for calculation. Just as we developed techniques to evaluate single integrals without resorting to Riemann sums, we need efficient methods for double integrals.

The key insight comes from examining how we might systematically accumulate volume. Looking at our rectangular region $R = [a, b] \times [c, d]$, we could:

- First, for each fixed x -value, integrate along the y -direction to find the area of a cross-sectional slice
- Then, integrate these slice areas along the x -axis to get the total volume

This intuition leads us to one of the most powerful results in multivariable calculus: Fubini's Theorem.

Statement of Fubini's Theorem

For a continuous function $f(x, y)$ on a rectangular region $R = [a, b] \times [c, d]$, Fubini's Theorem states that we can compute the double integral by evaluating two single integrals in either order:

$$\iint_R f(x, y) dA = \int_a^b \left[\int_c^d f(x, y) dy \right] dx = \int_c^d \left[\int_a^b f(x, y) dx \right] dy$$

This remarkable result tells us that two seemingly different processes—integrating first with respect to y then x , or first with respect to x then y —must yield the same result. Let's understand why this makes sense geometrically: imagine slicing our three-dimensional solid in two different ways:

- We could slice parallel to the yz -plane, getting cross-sections for each x

- Or we could slice parallel to the xz -plane, getting cross-sections for each y

Either way, when we add up all the slices, we must get the same volume.




Figure 4.3: Just as multiplying $\ell \times w \times h$ produces Volume the same way $w \times h \times \ell$ does, Fubini's theorem lets you calculate the volume under the curve commutatively. In this figure, we could either multiply the blue region (representing integration along x at a fixed $y = c$) over the range of y from c to d , or the red region (representing integration along y at a fixed x between a and b) over the range of x from a to b . Both approaches yield the same total volume under the surface.

Understanding Iterated Integrals

Let's examine the first form of Fubini's Theorem in detail:

$$\int_a^b \left[\int_c^d f(x, y) dy \right] dx$$

This is called an iterated integral, and we evaluate it from the inside out:

1. First, compute the inner integral $\int_c^d f(x, y) dy$:
 - Treat x as a constant
 - Integrate with respect to y from c to d
 - The result will be a function of x only
2. Then, integrate this result with respect to x from a to b

Think of this process as:

- First finding the area under the curve for each fixed x -value (the inner integral)

- Then accumulating these areas as x varies from a to b (the outer integral)
-

▲ **Example 4.1.2** Evaluate each of the following double integrals:

1. $\iint_R xy \, dA$ where $R = [0, 2] \times [0, 3]$
2. $\iint_R (x^3\sqrt{y} + \cos x) \, dA$ where $R = [0, 1] \times [0, 4]$
3. $\iint_{R_3} \frac{1}{1 + x^2 - y^2 - x^2y^2} \, dA$ where $R_3 = [0, 1] \times [0, \frac{1}{2}]$.

Solution Let's solve these systematically using Fubini's Theorem.

1. For $\iint_R xy \, dA$ on $R = [0, 2] \times [0, 3]$:

$$\begin{aligned}
 \iint_R xy \, dA &= \int_0^2 \left[\int_0^3 xy \, dy \right] dx \\
 &= \int_0^2 \left[x \left(\frac{y^2}{2} \right) \right]_0^3 dx \\
 &= \int_0^2 \left(\frac{9x}{2} \right) dx \\
 &= \frac{9}{2} \left(\frac{x^2}{2} \right) \Big|_0^2 \\
 &= \frac{9}{2}(2) = 9
 \end{aligned}$$

2. For $\iint_R (x^3\sqrt{y} + \cos x) \, dA$ on $R = [0, 1] \times [0, 4]$:

$$\begin{aligned}
 \iint_R (x^3\sqrt{y} + \cos x) \, dA &= \int_0^1 \left[\int_0^4 (x^3\sqrt{y} + \cos x) \, dy \right] dx \\
 &= \int_0^1 \left[x^3 \left(\frac{2y^{3/2}}{3} \right) + y \cos x \right]_0^4 dx \\
 &= \int_0^1 \left(\frac{16x^3}{3} + 4 \cos x \right) dx \\
 &= \left[\frac{16x^4}{12} + 4 \sin x \right]_0^1
 \end{aligned}$$

$$\begin{aligned}
&= \frac{16}{12} + 4 \sin(1) \\
&= \frac{4}{3} + 4 \sin(1)
\end{aligned}$$

3. For $\iint_R \frac{1}{1+x^2-y^2-x^2y^2} dA$ on $R = [0, 1] \times [0, \frac{1}{2}]$:

First, notice the denominator factors:

$$1 + x^2 - y^2 - x^2y^2 = (1 + x^2)(1 - y^2)$$

Therefore:

$$\begin{aligned}
\iint_R \frac{1}{1+x^2-y^2-x^2y^2} dA &= \int_0^1 \left[\int_0^{1/2} \frac{1}{(1+x^2)(1-y^2)} dy \right] dx \\
&= \int_0^1 \frac{1}{1+x^2} [\operatorname{arctanh}(y)]_0^{1/2} dx \\
&= \int_0^1 \frac{\operatorname{arctanh}(1/2)}{1+x^2} dx \\
&= \operatorname{arctanh}(1/2) [\operatorname{arctan}(x)]_0^1 \\
&= \operatorname{arctanh}(1/2) \cdot \frac{\pi}{4}
\end{aligned}$$

Strategic Choice of Integration Order

While Fubini's Theorem guarantees both orders of integration give the same result, one order might be significantly simpler than the other. In fact, sometimes one order might be straightforward while the other becomes extremely difficult or even impossible to evaluate directly. Let's develop some strategies for choosing the optimal order of integration.

A systematic approach to choosing integration order involves:

1. Examine the Function Structure:

- Look for terms that would simplify if held constant
- Identify potential partial fraction decompositions
- Note any terms that might lead to easier substitutions

2. Consider the Integration Boundaries:

- Check which variable has simpler limits
- Look for limits that depend on the other variable

- Consider whether one set of limits might introduce new complications

3. Anticipate the Result of Each Integration:

- Consider what antiderivatives would arise in each order
- Look for potential simplifications in the resulting functions
- Watch for terms that might cancel or combine nicely

Let's examine some specific scenarios:

- When $f(x, y)$ contains e^x or $\ln(x)$, integrate with respect to y first to keep these terms "constant"
- For rational functions like $\frac{1}{(x^2+y^2)}$, consider which variable might lead to easier partial fractions
- If $f(x, y)$ factors as $g(x)h(y)$, either order works equally well

Common pitfalls to avoid:

- Don't automatically choose the order that makes the inner integral look simpler—consider the full process
- Watch for cases where simplifying the inner integral makes the outer integral more complicated
- Be careful with functions that might become undefined during integration

Consider this example: If we have

$$\iint_R \frac{y}{x^2 + y^2} dA$$

integrating with respect to y first leads to arctangent functions, while integrating with respect to x first leads to logarithms. Depending on the region R , one approach might be significantly easier than the other.

The key insight is that Fubini's Theorem gives us flexibility in our approach. By choosing the optimal order of integration, we can often transform a seemingly difficult double integral into a sequence of manageable single integrals. This exemplifies a broader principle in mathematics: when faced with a complex problem, look for ways to transform it into a sequence of simpler, familiar problems.

▲ **Example 4.1.3** Evaluate $\iint_R \frac{y}{x^2 + y^2} dA$ where $R = [1, 2] \times [-1, 1]$ by considering both orders of integration. Determine which approach is more efficient.

Solution Let's evaluate this integral using both possible orders to compare approaches.

First, let's try integrating with respect to y first:

$$\iint_R \frac{y}{x^2 + y^2} dA = \int_1^2 \int_{-1}^1 \frac{y}{x^2 + y^2} dy dx$$

For any fixed x , notice that the integrand $f(y) = \frac{y}{x^2 + y^2}$ is an odd function in y :

$$f(-y) = \frac{-y}{x^2 + (-y)^2} = -\frac{y}{x^2 + y^2} = -f(y)$$

Therefore, when integrating over the symmetric interval $[-1, 1]$:

$$\int_{-1}^1 \frac{y}{x^2 + y^2} dy = 0$$

since the negative area perfectly cancels the positive area. Thus:

$$\iint_R \frac{y}{x^2 + y^2} dA = \int_1^2 [0] dx = 0$$

Now, let's examine what happens if we integrate with respect to x first:

$$\iint_R \frac{y}{x^2 + y^2} dA = \int_{-1}^1 \int_1^2 \frac{y}{x^2 + y^2} dx dy$$

Here, y is held constant in the inner integral:

$$\int_1^2 \frac{y}{x^2 + y^2} dx$$

This requires substitution $u = x^2 + y^2$ with $du = 2x dx$, leading to:

$$y \int \frac{dx}{x^2 + y^2} = \frac{y}{|y|} \arctan\left(\frac{x}{y}\right) \quad \text{for } y \neq 0$$

Evaluating from $x = 1$ to $x = 2$ and then integrating with respect to y becomes quite involved:

$$\int_{-1}^1 y \left[\frac{1}{|y|} \arctan\left(\frac{x}{y}\right) \right]_{x=1}^{x=2} dy$$




Figure 4.4: A plot of the function $f(x, y) = \frac{y}{x^2 + y^2}$. Notice that the function is odd, and thus integrating across symmetric bounds produces volume equal to 0.

Indefinite Double Integration

Just as with single integrals, we can consider indefinite double integrals where we seek a function $F(x, y)$ such that:

$$\frac{\partial^2 F}{\partial x \partial y} = g(x, y)$$

First, integrate with respect to y :

$$\frac{\partial F}{\partial x} = \int g(x, y) dy = G(x, y) + p(x),$$

where $G(x, y) = \int g(x, y) dy$ and $p(x)$ is an arbitrary function of x .

Next, integrate with respect to x :

$$F(x, y) = \int \frac{\partial F}{\partial x} dx = \int G(x, y) dx + \int p(x) dx + q(y).$$

The general solution is:

$$F(x, y) = H(x, y) + P(x) + q(y),$$

where $H(x, y) = \int G(x, y) dx$, $P(x) = \int p(x) dx$, and $q(y)$ is an arbitrary function of y . You have heard in your Calculus I and II texts that finding the function from an expression of derivatives is called a Differential Equation. This is the multivariable analog of that, called Partial Differential Equations, which will be introduced and discussed further in Chapter 6.

To find indefinite double integrals:

1. First integrate with respect to one variable, adding an arbitrary function of the other variable
2. Then integrate the result with respect to the second variable
3. Combine all arbitrary terms into the general form above

For example, if $f(x, y) = xy$:

$$\begin{aligned}\iint xy \, dx \, dy &= \int \left(\int xy \, dx \right) dy \\ &= \int \left(\frac{x^2 y}{2} + h(y) \right) dy \\ &= \frac{x^2 y^2}{4} + \int h(y) dy + g(x) \\ &= \frac{x^2 y^2}{4} + g(x) + H(y)\end{aligned}$$

This process highlights the additional complexity that arises in multivariable calculus—instead of just a constant of integration, we now have arbitrary functions to consider. Let's hit a couple of examples:

▲ **Example 4.1.4** Find the indefinite double integral for:

1. $\iint xy^2 \, dx \, dy$
2. $\iint (x^2 + 3xy) \, dx^2$

Solution Let's solve each integral systematically, remembering that for indefinite double integrals we need to include arbitrary functions of the other variable at each step.

1. For $\iint xy^2 dx dy$:

$$\begin{aligned}\int xy^2 dx &= \frac{x^2 y^2}{2} + h(y) \\ \iint xy^2 dx dy &= \int \left(\frac{x^2 y^2}{2} + h(y) \right) dy \\ &= \frac{x^2 y^3}{6} + \int h(y) dy + g(x) \\ &= \frac{x^2 y^3}{6} + g(x) + H(y)\end{aligned}$$

where $g(x)$ and $H(y)$ are arbitrary functions.

2. For $\iint (x^2 + 3xy) dx^2$:

First, note that integrating with respect to dx^2 means we integrate twice with respect to x :

$$\begin{aligned}\int (x^2 + 3xy) dx &= \frac{x^3}{3} + \frac{3x^2 y}{2} + h_1(y) \\ \int \left(\frac{x^3}{3} + \frac{3x^2 y}{2} + h_1(y) \right) dx &= \frac{x^4}{12} + \frac{x^3 y}{2} + h_1(y)x + h_2(y)\end{aligned}$$

Lecture 4.2: Double Integrals over General Regions

The world rarely presents us with perfectly rectangular regions to integrate over. Whether we're computing fluid flow through an irregular pipe, finding the center of mass of a non-rectangular plate, or calculating heat distribution across a circular disk, we need tools to handle integration over more general regions. In this lecture, we extend our understanding of double integrals beyond rectangular boundaries to tackle these real-world scenarios.




Figure 4.5: *Left: A rectangular region where the bounds of integration are constant. Right: A general region where the bounds may vary along each coordinate direction.*

Just as we first mastered integration over intervals before handling more complex single-variable integration, our study of rectangular regions has prepared us for this natural generalization. The key insight is that while our regions may now have curved boundaries, our fundamental approach remains the same—we still aim to accumulate infinitesimal pieces over a region systematically.




Figure 4.6: *A visualization of a double integral over a region R , with the surface $h(x, y)$ shown above.*

Type I and Type II Regions

When moving beyond rectangular regions, our first task is to understand how to describe more general regions mathematically. In rectangular regions, our bounds of integration were constants—fixed numbers that defined our region’s boundaries. Now, these bounds might be functions, describing curves that trace out our region’s boundaries. This transition from constant to variable bounds represents a fundamental shift in how we must think about setting up our integrals.

Let’s develop a systematic approach for analyzing these general regions and determining their bounds of integration. Just as we used the interval $[a, b]$ to precisely define the domain of a single integral, we now need two sets of bounds to fully describe our region of integration.

Analyzing Region Types

Before we can set up a double integral over a general region, we must first analyze the region’s geometry to determine whether it’s more naturally described as Type I or Type II. This analysis involves examining how the region’s boundaries intersect with horizontal and vertical lines.

For any given region R , ask yourself:

1. Can any vertical line through the region intersect the boundary at most twice?
2. Can any horizontal line through the region intersect the boundary at most twice?
3. Are these intersections continuous functions of position?

If the answer to the first question is “yes,” the region can be described as Type I. If the answer to the second question is “yes,” it can be described as Type II. Often, both answers are “yes,” giving makes things a bit complicated: we will need to either switch coordinates or divide up into sections of Type I and Type II regions and break up the integral and then proceed. Additionally, we can switch from Type I to several Type II cases and vice versa if it is advantageous. We will cover these cases in a bit.




Figure 4.7: Examples of regions and their classifications based on line intersections. Left: Type I region showing exactly two intersections with any vertical line. Center: Type II region showing exactly two intersections with any horizontal line. Right: An annular region that cannot be described as either type since both vertical and horizontal lines can intersect the boundary more than twice.

Type I Regions: Vertical Slicing

A region R in the xy -plane is called a **Type I region** if it can be described as:

$$R = \{(x, y) : a \leq x \leq b, \quad g_1(x) \leq y \leq g_2(x)\}$$

where $g_1(x)$ and $g_2(x)$ are continuous functions on $[a, b]$.


To determine the bounds for a Type I region, first and foremost we must sketch the region R . A clear diagram is absolutely essential—it helps us visualize the intersection points and understand how the bounds will be structured. Once we have our diagram, we follow these steps:

1. Identify the left and right bounds (a and b). These are typically intersection points of boundary curves or points where vertical tangent lines occur.
2. At any fixed x -value in $[a, b]$, determine where a vertical line enters and exits the region—these become $g_1(x)$ and $g_2(x)$ respectively.
3. Write these entry and exit points as functions of x . These become our inner bounds of integration.

Let's see this process in action with an example:

▲ **Example 4.2.1** Find the bounds needed to evaluate $\iint_R f(x, y) dA$ where R is bounded below by $y = x^2$ and above by $y = 4 - x^2$.

Solution Let's first sketch the region R to visualize our domain of integration:



To find the bounds, we first determine where the curves intersect:

$$x^2 = 4 - x^2 \implies 2x^2 = 4 \implies x = \pm\sqrt{2}$$

From our sketch, we can see:

- The region extends from $x = -\sqrt{2}$ to $x = \sqrt{2}$
- For each x in this interval, y ranges from x^2 to $4 - x^2$


Therefore:

$$\iint_R f(x, y) dA = \int_{-\sqrt{2}}^{\sqrt{2}} \int_{x^2}^{4-x^2} f(x, y) dy dx$$

Note how the sketch clearly shows this is a Type I region, as any vertical line (like our dashed sample line) intersects the boundary curves exactly twice. At this point we may now proceed with integrating as we have done in the previous lecture.

▲ **Example 4.2.2** A manufacturer produces some eccentric-looking solar panels whose efficiency varies with position. The usable surface of each panel is bounded below by $y = \sin(x)$ and above by $y = 2 + \cos(x)$ for $x \in [0, 2\pi]$. If the efficiency at any point (x, y) is given by $E(x, y) = \frac{1}{80}x^2y$, find the average efficiency over the entire panel.

Solution Let's first sketch the region to understand our domain of integration:



To find the average efficiency, we need:

$$\begin{aligned} E_{\text{avg}} &= \frac{\iint_R E(x, y) dA}{\iint_R dA} \\ &= \frac{\iint_R x^2 y dA}{\text{Area of } R}. \end{aligned}$$

From the bounds, we have $0 \leq x \leq 2\pi$ and $\sin(x) \leq y \leq 2 + \cos(x)$. Thus,

$$\iint_R x^2 y dA = \int_0^{2\pi} \int_{\sin(x)}^{2+\cos(x)} \frac{1}{80} x^2 y dy dx.$$

Evaluate the inner integral:

$$\begin{aligned} \int_{\sin(x)}^{2+\cos(x)} \frac{1}{80} x^2 y dy &= \frac{x^2}{80} \left[\frac{y^2}{2} \right]_{y=\sin(x)}^{y=2+\cos(x)} \\ &= \frac{x^2}{160} \left((2 + \cos(x))^2 - \sin^2(x) \right). \end{aligned}$$

Thus,

$$\begin{aligned} \iint_R x^2 y dA &= \frac{1}{160} \int_0^{2\pi} x^2 (4 + 4 \cos(x) + \cos(2x)) dx \\ &= \frac{1}{160} \left(4 \int_0^{2\pi} x^2 dx + 4 \int_0^{2\pi} x^2 \cos(x) dx + \int_0^{2\pi} x^2 \cos(2x) dx \right). \end{aligned}$$

By standard integration techniques,

$$\int_0^{2\pi} x^2 dx = \frac{8\pi^3}{3}, \quad \int_0^{2\pi} x^2 \cos(x) dx = 4\pi, \quad \int_0^{2\pi} x^2 \cos(2x) dx = \pi.$$

Hence,

$$\begin{aligned}\iint_R \frac{1}{80} x^2 y \, dA &= \frac{1}{160} \left(4 \cdot \frac{8\pi^3}{3} + 4 \cdot 4\pi + \pi \right) \\ &= \frac{1}{160} \left(\frac{32\pi^3}{3} + 16\pi + \pi \right)\end{aligned}$$

Next, the area of R is

$$\begin{aligned}\text{Area}(R) &= \int_0^{2\pi} \left[(2 + \cos(x)) - \sin(x) \right] dx \\ &= \int_0^{2\pi} 2 \, dx + \int_0^{2\pi} \cos(x) \, dx - \int_0^{2\pi} \sin(x) \, dx \\ &= 4\pi.\end{aligned}$$

Therefore, the average efficiency is

$$E_{\text{avg}} = \frac{\frac{1}{160} \left(\frac{32\pi^3}{3} + 16\pi + \pi \right)}{4\pi} \approx 0.189 = 18.9\%.$$

This is the average efficiency across the entire solar panel surface. Note, that unlike have a surface intersecting the xy -plane producing negative volume, the bounds of the region that go to negative quadrants do not effect the volume.

Type II Regions: Horizontal Slicing

A region R in the xy -plane is called a **Type II region** if it can be described as:

$$R = \{(x, y) : c \leq y \leq d, \quad h_1(y) \leq x \leq h_2(y)\}$$

where $h_1(y)$ and $h_2(y)$ are continuous functions on $[c, d]$.

Just as with Type I regions, sketching region R is essential. A clear diagram helps us visualize where horizontal lines intersect our boundaries and how to structure our integration. After sketching the region, we follow these steps:


1. Identify the bottom and top bounds (c and d). These are typically intersection points of boundary curves or points where horizontal tangent lines occur.
2. At any fixed y -value in $[c, d]$, determine where a horizontal line enters and exits the region—these become $h_1(y)$ and $h_2(y)$ respectively.
3. Write these entry and exit points as functions of y . These become our inner bounds of integration.

▲ **Example 4.2.3** In fluid dynamics, the velocity potential $\phi(x, y) = \frac{x}{1+y^2}$ describes the flow field in a curved channel. Calculate the total potential flux through region R bounded by the cubic barrier $x = y^3$, the vertical wall $x = 4$, and the horizontal barriers at $y = \pm 1$.

The potential flux across a region is given by:

$$\Phi = \iint_R \phi(x, y) dA.$$

Solution Let's first sketch region R to understand our flow domain:



For this flow problem, Type II integration is natural as it aligns with horizontal flow layers. For any fixed y -level between -1 and 1 , the flow extends from the barrier at $x = y^3$ to the wall at $x = 4$. Therefore:

$$\begin{aligned} \Phi &= \int_{-1}^1 \int_{y^3}^4 \frac{x}{1+y^2} dx dy \\ &= \int_{-1}^1 \frac{1}{1+y^2} \left[\frac{x^2}{2} \right]_{y^3}^4 dy \\ &= \int_{-1}^1 \frac{1}{1+y^2} \left(8 - \frac{y^6}{2} \right) dy \\ &= 8 \int_{-1}^1 \frac{dy}{1+y^2} - \frac{1}{2} \int_{-1}^1 \frac{y^6}{1+y^2} dy. \end{aligned}$$

The first integral evaluates to 4π . For the second, using polynomial division:

$$\begin{aligned} \frac{y^6}{1+y^2} &= y^4 - y^2 + 1 - \frac{1}{1+y^2}, \\ \int_{-1}^1 \frac{y^6}{1+y^2} dy &= \left[\frac{y^5}{5} - \frac{y^3}{3} + y - \arctan(y) \right]_{-1}^1 \\ &= 2 \left(\frac{1}{5} - \frac{1}{3} + 1 - \frac{\pi}{4} \right) \end{aligned}$$

$$= \frac{26}{15} - \frac{\pi}{2}.$$

Therefore:

$$\begin{aligned}\Phi &= 4\pi - \frac{1}{2} \left(\frac{26}{15} - \frac{\pi}{2} \right) \\ &= \frac{17\pi}{4} - \frac{13}{15} \\ &\approx 12.48.\end{aligned}$$

The positive value indicates net flow in the positive x -direction, with the factor $\frac{1}{1+y^2}$ creating a focused channel along $y = 0$.

Double Integrals in Polar Coordinates

Many regions in the plane possess a natural circular or radial symmetry that makes them awkward to describe in rectangular coordinates. A circle, for instance, has a simple equation $r = a$ in polar coordinates but requires the more complex equation $x^2 + y^2 = a^2$ in rectangular coordinates. For such regions, polar coordinates often provide both simpler descriptions and easier integrations.

The Transformation to Polar Coordinates

In polar coordinates, we have learned in Chapter 1 that we locate a point P in the plane by its distance r from the origin and the angle θ that the line from the origin to P makes with the positive x -axis. The relationship between rectangular and polar coordinates is:

$$\begin{aligned}x &= r \cos \theta \\ y &= r \sin \theta\end{aligned}$$

When we transform to polar coordinates, our differential element of area changes from $dA = dx dy$ to $dA = r dr d\theta$. Let's understand why through a careful geometric analysis. Consider a small area element in polar coordinates. This element is approximately a circular sector at radius r , bounded by:

- Two radii separated by a small angle $d\theta$
- Two circular arcs at radii r and $r + dr$




Figure 4.8: Visualization of a polar area element showing the infinitesimal area $r d\theta dr$ in polar coordinates.

At radius r , a small angle $d\theta$ creates an arc of length $r d\theta$ (just as the circumference of a circle is $2\pi r$). The radial thickness of our element is dr . Therefore, our area element is approximately a rectangle with:

$$\text{width} = r d\theta$$

$$\text{height} = dr$$

Multiplying these gives our area element:

$$dA = r dr d\theta$$

This explains why the factor of r appears—as we move farther from the origin (larger r), the same angle $d\theta$ sweeps out a longer arc, creating a larger area element. The actual area element becomes exact as dr and $d\theta$ become infinitesimally small. If we also think about the units this also makes sense: Area requires a squared length, and thus r and dr provide the length. $d\theta$ is unitless since it's an angle.

Therefore once considering the surface function $f(x, y)$ for the height and transforming it to polar: $f(x, y) = f(r \cos \theta, r \sin \theta) = g(r, \theta)$, a double integral in polar coordinates takes the general form:

$$\iint_R f(x, y) dA = \iint_R f(r \cos \theta, r \sin \theta) r dr d\theta = \iint_R g(r, \theta) r dr d\theta$$

Setting Up Polar Double Integrals

When working with double integrals in polar coordinates, students must follow a systematic approach that transforms the rectangular coordinate integration into a more manageable polar coordinate representation. Let's break down the process step by step:

1. **Identify the Bounds on r and θ :** This crucial first step involves carefully determining the limits of your integration region. Unlike rectangular coordinates where you define rectangular bounds, in polar coordinates you'll specify:

- The range of radial distance r from the origin, which determines how far out you'll integrate
- The range of angle θ , which determines the angular sweep of your region

Think of this as creating a pie slice or full circle in the coordinate plane. The limits will depend on the specific geometric region you're integrating over.

2. **Convert the Integrand from Rectangular to Polar Coordinates:** This involves replacing rectangular variables with their polar equivalents:

$$x = r \cos \theta$$

$$y = r \sin \theta$$

Any function $f(x, y)$ becomes $f(r \cos \theta, r \sin \theta)$. For instance, $x^2 + y^2$ simply becomes r^2 .

3. **Include the Factor of r from the Area Element:** In polar coordinates, the differential area changes from $dA = dx dy$ to $dA = r dr d\theta$. This critical r factor accounts for the expanding arc length as you move further from the origin. Always multiply your integrand by r to maintain the correct area scaling.
4. **Set Up the Integral with the Appropriate Order of Integration:** Typically, you'll integrate with respect to r first, then θ . This approach mimics how you'd conceptually sweep radial lines through their angular range, progressively building up the area of your region.

The natural order of integration in polar coordinates is usually to integrate with respect to r first, then θ . This corresponds to integrating along radial lines from the origin, then sweeping these lines through the required angular range.

Some expressions become particularly simple in polar coordinates:

- $x^2 + y^2 = r^2$
- $\tan^{-1}(\frac{y}{x}) = \theta$
- $\sqrt{x^2 + y^2} = r$

Standard Regions in Polar Coordinates

Several types of standard regions are particularly well-suited to polar coordinates:

1. **Circular Disks:** For a disk of radius a :

$$0 \leq r \leq a$$

$$0 \leq \theta \leq 2\pi$$

2. **Circular Sectors:** For a sector of radius a from angle α to β :


$$0 \leq r \leq a$$

$$\alpha \leq \theta \leq \beta$$

3. **Annuli:** For a region between circles of radius a and b :

$$a \leq r \leq b$$

$$0 \leq \theta \leq 2\pi$$



Also note that when converting to polar coordinates to integrate, some expressions become particularly simple in polar coordinates:

- $x^2 + y^2 = r^2$
- $\tan^{-1}(\frac{y}{x}) = \theta$
- $\sqrt{x^2 + y^2} = r$

▲ **Example 4.2.4** Evaluate the double integral $\iint_R (x^2 + y^2) dA$, where R is the circular disk bounded by $r = 2$.

Solution This problem is ideal for polar coordinates due to the circular symmetry of the region. Let's solve it step by step:

First, note the transformation from rectangular to polar coordinates:

- $x^2 + y^2 = r^2$
- $dA = r dr d\theta$

For the circular disk, the bounds are straightforward:

$$0 \leq r \leq 2$$

$$0 \leq \theta \leq 2\pi$$

Transforming the integral:

$$\begin{aligned}\iint_R (x^2 + y^2) dA &= \int_0^{2\pi} \int_0^2 r^2 \cdot r dr d\theta \\ &= \int_0^{2\pi} \left[\frac{r^4}{4} \right]_0^2 d\theta \\ &= 8\pi\end{aligned}$$

The final result is 8π , demonstrating how polar coordinates simplify the integration of this radially symmetric function.

Irregular Regions in Polar Coordinates

Now let's think about what happens when we don't have a standard region, but one that is defined by polar functions. Consider the figure below, which shows an irregular region R bounded by two curves in polar form $r = h_1(\theta)$ and $r = h_2(\theta)$. Above this region, a surface $z = f(x, y)$ creates a volume that we might want to calculate. The vertical lines represent the height of the surface at various points, while the polar grid below shows how we divide the region for integration. This is essentially the polar version of the Riemann sum we saw in the previous lecture.




Figure 4.9: *Volume integral in polar coordinates with a gently undulating surface.*

Now the next figure provides a detailed view of how we construct our polar integral for our irregular region. We focus on a section of the first quadrant (from $\pi/6$ to $\pi/3$) to see the

structure more clearly. The highlighted area element demonstrates how dr represents the radial thickness while $r d\theta$ gives the arc length at radius r . Notice that this wedge-shaped element grows wider as r increases—this growth is precisely what the factor of r in our area element accounts for.




Figure 4.10: Middle third section ($\pi/6$ to $\pi/3$) of the polar region with grid lines and detailed view of infinitesimal area element.

Let's see how to mathematically construct the integral for this example now. Given a region R bounded by two curves $r = h_1(\theta)$ and $r = h_2(\theta)$, with $h_1(\theta) \leq h_2(\theta)$, our general integral takes the form:

$$\iint_R f(x, y) dA = \int_{\alpha}^{\beta} \int_{h_1(\theta)}^{h_2(\theta)} f(r \cos \theta, r \sin \theta) r dr d\theta$$

where:


- $[\alpha, \beta]$ represents the angular range of the region. In the case of the whole region in this example we would have $\alpha = 0$, $\beta = 2\pi$.
- $h_1(\theta)$ gives the inner boundary curve
- $h_2(\theta)$ gives the outer boundary curve
- The factor r appears from our area element $dA = r dr d\theta$

Just as with rectangular coordinates, we integrate from the inner bound to the outer bound, but now these bounds are functions of our angular variable θ . The key difference is that we must include the factor of r from our area element, which accounts for the expanding size of our wedge-shaped regions as we move outward from the origin.

▲ **Example 4.2.5** An engineering team is excavating a basin in the first quadrant as shown below. The basin is bounded by two circular arcs whose centers are marked in the diagram. Each point in the basin has a depth proportional to the angle it makes with the positive x-axis, given by

$$z = \arctan\left(\frac{y - \frac{1}{2}}{x - \frac{1}{2}}\right).$$

Using the diagram information, determine the total volume of soil to be removed.



Solution Let's solve this step by step:

1. Analyze the region from the diagram. The basin consists of points (x, y) in the first quadrant where the distance from $(0.5, 0.5)$ is between 1 and 1.5. If we let r be that distance and θ the angle measured from the positive x -axis (through the center $(0.5, 0.5)$), then:

$$r = \sqrt{(x - 0.5)^2 + (y - 0.5)^2}, \quad \tan(\theta) = \frac{y - 0.5}{x - 0.5}.$$

The basin boundaries give us:

$$0 \leq \theta \leq \frac{\pi}{2} \quad \text{and} \quad 1 \leq r \leq 1.5.$$

2. Simplify the height function. Inside the arctan, we have:

$$\frac{y - 0.5}{x - 0.5} = \frac{r \sin(\theta)}{r \cos(\theta)} = \tan(\theta).$$

Therefore:

$$z = \arctan(\tan(\theta)) = \theta \quad \text{since } 0 \leq \theta \leq \frac{\pi}{2}.$$

3. Set up and evaluate the volume integral. Using polar coordinates centered at $(0.5, 0.5)$, the differential area element is $r \, dr \, d\theta$. The volume is:

$$\text{Volume} = \iint_{\text{basin}} z \, dA = \int_0^{\pi/2} \int_1^{1.5} \theta \, r \, dr \, d\theta.$$

First, integrate with respect to r :

$$\begin{aligned} \int_1^{1.5} \theta \, r \, dr &= \theta \int_1^{1.5} r \, dr \\ &= \theta \left[\frac{r^2}{2} \right]_{r=1}^{r=1.5} \\ &= \theta \left(\frac{(1.5)^2}{2} - \frac{1^2}{2} \right) \\ &= \theta(1.125 - 0.5) \\ &= 0.625\theta. \end{aligned}$$

Now integrate with respect to θ :

$$\begin{aligned} \int_0^{\pi/2} 0.625\theta \, d\theta &= 0.625 \int_0^{\pi/2} \theta \, d\theta \\ &= 0.625 \left[\frac{\theta^2}{2} \right]_0^{\pi/2} \\ &= 0.625 \cdot \frac{(\pi/2)^2}{2} \\ &= \frac{5\pi^2}{64}. \end{aligned}$$

Therefore, the volume of soil to be removed is $\frac{5\pi^2}{64}$ cubic units.

▲ **Example 4.2.6** Find the value of the integral

$$\int_{-\infty}^{\infty} e^{-x^2} \, dx.$$

Solution

1. Let $I = \int_{-\infty}^{\infty} e^{-x^2} \, dx$.

2. Square it:

$$I^2 = \left(\int_{-\infty}^{\infty} e^{-x^2} dx \right) \left(\int_{-\infty}^{\infty} e^{-y^2} dy \right) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} e^{-(x^2+y^2)} dx dy.$$

3. Switch to polar coordinates with $x = r \cos \theta$, $y = r \sin \theta$. The change in variable to polar coordinates gives $dx dy = r dr d\theta$. The bounds are $r : 0 \rightarrow \infty$ and $\theta : 0 \rightarrow 2\pi$:

$$I^2 = \int_0^{2\pi} \int_0^{\infty} e^{-r^2} r dr d\theta.$$

4. Separate the integrals:

$$I^2 = \left(\int_0^{2\pi} d\theta \right) \left(\int_0^{\infty} e^{-r^2} r dr \right) = 2\pi \int_0^{\infty} e^{-r^2} r dr.$$

5. Use the substitution $u = r^2$, so $du = 2r dr$ and $r dr = \frac{1}{2} du$:

$$I^2 = 2\pi \int_0^{\infty} e^{-r^2} r dr = 2\pi \int_0^{\infty} e^{-u} \frac{1}{2} du = \pi \int_0^{\infty} e^{-u} du = \pi [-e^{-u}]_0^{\infty} = \pi \cdot 1 = \pi.$$

6. Hence $I = \sqrt{\pi}$. Therefore,

$$\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}.$$

This is a common technique used to integrate functions with impossible anti-derivatives.

Complex Regions and Multiple Integrals

Not all regions can be neatly categorized as Type I or Type II. Consider a region bounded by multiple curves where vertical or horizontal lines might intersect the boundary more than twice. In such cases, we must break down our region into simpler pieces where our integration techniques apply.

The fundamental principle we use is the additivity of integrals:

$$\iint_R f(x, y) dA = \iint_{R_1} f(x, y) dA + \iint_{R_2} f(x, y) dA$$

where $R = R_1 \cup R_2$ and $R_1 \cap R_2 = \emptyset$

Consider, for example, the region bounded by $y = x^2$ and $y = 4$:




Figure 4.11: A parabola that can be split into two regions due to additivity of integrals.

This region can be split into two Type I regions at $x = 0$:

- R_1 : Region where $-2 \leq x \leq 0$, $x^2 \leq y \leq 4$
- R_2 : Region where $0 \leq x \leq 2$, $x^2 \leq y \leq 4$

Sometimes a region might be better described by splitting it into Type II regions:




Figure 4.12: An otherwise Type I region that is converted into Type II by adjusting the equations and finding the appropriate intersection point.

Here, we might split at $y = 2$ into:

- R_1 : Region where $2 \leq y \leq 4 - x^2$, $-\sqrt{4 - y} \leq x \leq \sqrt{4 - y}$

- R_2 : Region where $x^2 \leq y \leq 2$, $-\sqrt{y} \leq x \leq \sqrt{y}$

When working with complex regions, we often encounter situations that require more sophisticated splitting strategies:




Figure 4.13: Two piece-wise defined Type I regions.

For piecewise-defined regions, natural splitting points occur where the boundary function changes form. Each piece may suggest a different integration approach—Type I or Type II—based on the function’s behavior.

Some regions may benefit from mixing coordinate systems. Consider the region bounded by $y = x^2$ and $r = 2$:




Figure 4.14: Two piece-wise defined regions where R_1 and R_2 can be the bounds of two double integrals with different coordinate systems. The height of the triangle and radius of the quarter circle are both 2.5 units.

Here, the left region R_1 might be better described using Cartesian coordinates, while the right region R_2 could be more naturally integrated using polar coordinates to handle the circular boundary. When determining how to split a region:

- Consider the form of both the boundary curves and the integrand
- Look for symmetry that could simplify calculations
- Evaluate whether changing coordinate systems might simplify the integration
- Balance the number of subregions against the complexity of individual integrals

▲ Example 4.2.7

1. Find the area of the region shown in Figure 4.14 by evaluating the double integral

$$\iint_{R=R_1 \cup R_2} dA$$

2. Use the surface $f(x, y) = x^2 + y^2 + 1$ to find the volume over the region R .

Solution

1. For finding the area:

- (a) First, we split the integral using the additivity property:

$$\iint_R dA = \iint_{R_1} dA + \iint_{R_2} dA$$

- (b) For region R_1 , we can use Cartesian coordinates. The region is a triangle bounded by $x = 0$, $x = 2$, and the line passing through $(0, 0)$ and $(2, 2.5)$. The equation of this line is $y = \frac{5}{4}x$:

$$\iint_{R_1} dA = \int_0^2 \int_0^{\frac{5}{4}x} dy \, dx = \int_0^2 \frac{5}{4}x \, dx = \frac{5}{2}$$

- (c) For region R_2 , we can use polar coordinates since it's a quarter circle of radius 2.5 centered at $(2, 0)$. We need to shift our polar coordinates by translating 2 units right:

$$\iint_{R_2} dA = \int_0^{\pi/2} \int_0^{2.5} r \, dr \, d\theta = \frac{\pi}{4}(2.5)^2$$

- (d) The total area is therefore:

$$\text{Area} = \frac{5}{2} + \frac{25\pi}{16} \approx 7.42 \text{ square units}$$

2. Now for our surface function $f(x, y)$, we would have to integrate $f(x, y)$ over R_1 , and then have to integrate $f(r \cos \theta, r \sin \theta) = g(r, \theta)$ over R_2 .

- (a) For integrating $f(x, y) = x^2 + y^2 + 1$ over R_1 , we use Cartesian coordinates:

$$\iint_{R_1} f(x, y) dA = \int_0^2 \int_0^{\frac{5}{4}x} (x^2 + y^2 + 1) dy dx$$

Evaluating the inner integral:

$$\begin{aligned} \int_0^2 \left[x^2 y + \frac{y^3}{3} + y \right]_0^{\frac{5}{4}x} dx &= \int_0^2 \left(x^2 \left(\frac{5}{4}x \right) + \frac{1}{3} \left(\frac{5}{4}x \right)^3 + \frac{5}{4}x \right) dx \\ &= \int_0^2 \left(\frac{5}{4}x^3 + \frac{125}{192}x^3 + \frac{5}{4}x \right) dx = \left[\frac{5}{16}x^4 + \frac{125}{768}x^4 + \frac{5}{8}x^2 \right]_0^2 \\ &= \frac{5}{16}(16) + \frac{125}{768}(16) + \frac{5}{8}(4) = \frac{485}{48} \approx 10.104 \end{aligned}$$

- (b) For R_2 , we translate to the center of the quarter circle at $(2, 0)$. Using polar coordinates with this shift:

$$x = r \cos \theta, \quad y = r \sin \theta$$

Therefore, $x^2 + y^2 + 1 = r^2 + 1$. The integral becomes:

$$\begin{aligned} \iint_{R_2} f(x, y) dA &= \int_0^{\pi/2} \int_0^{2.5} (r^2 + 1)r dr d\theta \\ &= \int_0^{\pi/2} \left[\frac{r^4}{4} + \frac{r^2}{2} \right]_0^{2.5} d\theta \\ &= \int_0^{\pi/2} \left(\frac{(2.5)^4}{4} + \frac{(2.5)^2}{2} \right) d\theta \\ &= \left(\frac{(2.5)^4}{4} + \frac{(2.5)^2}{2} \right) [\theta]_0^{\pi/2} \approx 20.2485 \end{aligned}$$

- (c) The total volume is therefore:

$$V \approx 30.3525$$

Lecture 4.3: Applications of Double Integrals

Having established our foundation with area and volume calculations, we now venture into how double integrals illuminate the physical properties of objects in the real world. These applications reveal the deep connection between calculus and physics, showing how mathematical tools can predict and explain the behavior of physical systems.

Mass and Density

When engineers design aircraft wings, solar panels, or electronic circuit boards, they often work with objects that are effectively two-dimensional—their thickness is negligible compared to their other dimensions. For these thin objects, we describe their mass distribution using a density function $\rho(x, y)$ that gives the mass per unit area at each point. This density might vary across the object; think of a solar panel with different components or a circuit board with varying concentrations of conducting material.

To find the total mass of such an object, we integrate the density over the region it occupies:

$$M = \iint_R \rho(x, y) dA$$

This integral emerges naturally from the physical principle of additivity of mass—the total mass is the sum of the masses of all small pieces. Each infinitesimal piece has area dA and density $\rho(x, y)$, giving it a mass of $\rho(x, y) dA$. The double integral then sums these contributions over the entire region (thus bringing us to the required $M = \rho V_R$ for our units).

Consider, for instance, a sheet of metal that has been treated with a protective coating. The coating might be applied more thickly around the edges, creating a density function that increases as we move away from the center. Or think about a decorative stained glass window, where different colored pieces have different densities. In these cases, the density function captures the physical reality of how mass is distributed across the object.

Center of Mass and Balance

Now that we understand how to represent mass distribution using double integrals, we can tackle an important related problem: finding where an object will balance. Given a flat plate with varying density, where should we place a support to keep it perfectly level? This question leads us to the concept of center of mass.

The center of mass of an object tells us where it effectively balances, a concept crucial in everything from architectural design to spacecraft engineering. For a flat object with density $\rho(x, y)$, we call the coordinates of its center of mass $(\bar{x}, \bar{y})$, given by:

$$\bar{x} = \frac{1}{M} \iint_R x \rho(x, y) dA$$

$$\bar{y} = \frac{1}{M} \iint_R y \rho(x, y) dA$$

These formulas arise from a fundamental physical principle: the total turning effect (called torque) on an object balanced at its center of mass must be zero. Torque measures the tendency of a force to rotate an object around a point - think of pushing a door near its hinges versus near its handle. Each point in the object creates a torque that depends on both its mass and its distance from the axis. The numerator integrals sum up these effects, while division by the total mass M gives us the balancing point.




Figure 4.15: A 2D representation of the center of mass (CM) for a system of four masses. The position vectors $\mathbf{r}_1$, $\mathbf{r}_2$, $\mathbf{r}_3$, and $\mathbf{r}_4$ point from the origin to their respective masses m_1 , m_2 , m_3 , and m_4 . The center of mass vector $\mathbf{r}_{\text{cm}}$ is shown in red, indicating the weighted average position of the masses.

The practical implications are profound. When aerospace engineers design control surfaces for aircraft, they must carefully consider the center of mass. A poorly balanced aileron or rudder could lead to control issues or mechanical failure. Similarly, architects designing large suspended sculptures must understand exactly where each piece will balance. The center of mass calculations tell them where to place support structures and how to distribute the weight of the artwork. Therefore study this hard so you don't wind up killing someone.

▲ **Example 4.3.1** NASA engineers are designing a new solar sail—a large, flat spacecraft that uses radiation pressure from sunlight for propulsion. The sail is shaped like a region R bounded by $y = \sqrt{x}$ and $y = x^2$ in the first quadrant. The sail's density varies according to $\rho(x, y) = 2 + \frac{x}{y}$ kg/m².

1. Find the total mass of the solar sail.
2. Determine the coordinates of its center of mass $(\bar{x}, \bar{y})$.
3. If the sail must be balanced on a single support strut at $x = \bar{x}$, what percentage of the sail's mass lies to the left?

Solution Let's solve this step by step:

1. First, let's find where the curves intersect to determine our region's bounds:

$$\sqrt{x} = x^2 \implies x^4 = x \implies x(x^3 - 1) = 0 \implies x = 0 \text{ or } x = 1$$

Both give $y = 0$ at $x = 0$ (the origin) and $y = 1$ at $x = 1$.

The region R is therefore $0 \leq x \leq 1$ with $x^2 \leq y \leq \sqrt{x}$. The total mass is:

$$M = \iint_R \rho(x, y) dA = \iint_R \left(2 + \frac{x}{y}\right) dA$$

As a Type I region:

$$M = \int_0^1 \int_{x^2}^{\sqrt{x}} \left(2 + \frac{x}{y}\right) dy dx$$

Integrate with respect to y first:

$$\begin{aligned} \int_{x^2}^{\sqrt{x}} \left(2 + \frac{x}{y}\right) dy &= [2y + x \ln |y|]_{y=x^2}^{y=\sqrt{x}} \\ &= (2\sqrt{x} + x \ln(\sqrt{x})) - (2x^2 + x \ln(x^2)) \end{aligned}$$

Recall $\ln(\sqrt{x}) = \frac{1}{2} \ln(x)$ and $\ln(x^2) = 2 \ln(x)$, giving:

$$2\sqrt{x} - 2x^2 + x \left(\frac{1}{2} \ln x - 2 \ln x \right) = 2\sqrt{x} - 2x^2 - \frac{3}{2} x \ln x$$

Hence:

$$M = \int_0^1 \left(2\sqrt{x} - 2x^2 - \frac{3}{2} x \ln x \right) dx$$

An antiderivative is:

$$\begin{aligned} \int 2\sqrt{x} dx &= \frac{4}{3} x^{3/2}, \\ \int -2x^2 dx &= -\frac{2}{3} x^3, \end{aligned}$$

$$\int -\frac{3}{2}x \ln x \, dx = -\frac{3}{4}x^2 \ln x + \frac{3}{8}x^2$$

Therefore:

$$\left[\frac{4}{3}x^{3/2} - \frac{2}{3}x^3 - \frac{3}{4}x^2 \ln x + \frac{3}{8}x^2 \right]_0^1 = \frac{4-2}{3} + \frac{3}{8} = \frac{25}{24}$$

Thus:

$$M = \frac{25}{24} \approx 1.0417 \text{ kg}$$

2. For the center of mass, we need:

$$\bar{x} = \frac{1}{M} \iint_R x \rho(x, y) \, dA \quad \text{and} \quad \bar{y} = \frac{1}{M} \iint_R y \rho(x, y) \, dA$$

For $\bar{x}$:

$$\begin{aligned} \bar{x}M &= \int_0^1 \int_{x^2}^{\sqrt{x}} x \left(2 + \frac{x}{y} \right) dy \, dx \\ &= \int_0^1 [x(2y + x \ln |y|)]_{y=x^2}^{y=\sqrt{x}} dx \end{aligned}$$

After careful evaluation:

$$\bar{x}M = \int_0^1 \left(2x^{3/2} - 2x^3 - \frac{3}{2}x^2 \ln x \right) dx = \frac{7}{15}$$

Therefore:

$$\bar{x} = \frac{7/15}{25/24} = \frac{168}{375} \approx 0.448$$

For $\bar{y}$:

$$\bar{y}M = \int_0^1 \int_{x^2}^{\sqrt{x}} y \left(2 + \frac{x}{y} \right) dy \, dx = \int_0^1 \int_{x^2}^{\sqrt{x}} (2y + x) dy \, dx$$

Evaluating:

$$\int_{x^2}^{\sqrt{x}} (2y + x) dy = [y^2 + xy]_{y=x^2}^{y=\sqrt{x}} = (x + x^{3/2}) - (x^4 + x^3)$$

Therefore:

$$\bar{y}M = \int_0^1 (x + x^{3/2} - x^4 - x^3) dx = 0.45$$

Hence:

$$\bar{y} = \frac{0.45}{25/24} = 0.432$$

$$(\bar{x}, \bar{y}) = (0.448, 0.432)$$

3. We compute:

$$M_{\text{left}} = \int_0^{\bar{x}} \int_{x^2}^{\sqrt{x}} \left(2 + \frac{x}{y}\right) dy dx$$

Using the same antiderivative for the inner integral $(2y + x \ln |y|)$, evaluated from $y = x^2$ to $y = \sqrt{x}$, giving $2\sqrt{x} - 2x^2 - \frac{3}{2}x \ln x$.

Let:

$$G(x) = \frac{4}{3}x^{3/2} - \frac{2}{3}x^3 - \frac{3}{4}x^2 \ln x + \frac{3}{8}x^2$$

Then:

$$M_{\text{left}} = [G(x)]_0^{\bar{x}} \text{ where } \bar{x} \approx 0.448$$

Numerically:

$$G(0.448) \approx 0.534 \implies M_{\text{left}} \approx 0.534$$

Since $M = 25/24 \approx 1.0417$:

$$\frac{M_{\text{left}}}{M} \approx \frac{0.534}{1.0417} \approx 0.5127 = 51.27\%$$

These calculations demonstrate how varying density affects a solar sail's mass distribution and center of mass. A precise balance is critical for attitude control, as any asymmetry could cause off-axis torques from solar radiation pressure.

Moments of Inertia and Rotational Motion

Having understood how mass is distributed across a flat plate and where it balances, we can now explore how this mass distribution affects the object's rotation. Just as an object's mass determines its resistance to linear motion, a quantity called moment of inertia determines its resistance to rotational motion.

For a flat plate rotating about various axes, we define three different moments of inertia, each measured in $\text{kg} \cdot \text{m}^2$:

$$I_x = \iint_R y^2 \rho(x, y) dA$$

$$I_y = \iint_R x^2 \rho(x, y) dA$$

$$I_z = \iint_R (x^2 + y^2) \rho(x, y) dA$$

Using the transformation to polar coordinates ($x = r \cos \theta, y = r \sin \theta$) with $dA = r dr d\theta$, these become:

$$\begin{aligned} I_x &= \iint_R r^3 \sin^2 \theta \rho(r, \theta) dr d\theta \\ I_y &= \iint_R r^3 \cos^2 \theta \rho(r, \theta) dr d\theta \\ I_z &= \iint_R r^3 \rho(r, \theta) dr d\theta \end{aligned}$$

The squared terms in these formulas reflect a deep physical principle: mass farther from the axis of rotation contributes more to rotational inertia. This is why figure skaters pull their arms in to spin faster—they're reducing their moment of inertia about the vertical axis. It's also why flywheels in engines are designed with most of their mass at the outer edge, maximizing their ability to store rotational energy.



Figure 4.16: A skater performs a spin and holds his arms in to increase rotational motion by reducing moment of inertia.

Consider a flywheel in an engine. When designed with a larger moment of inertia, it more effectively smooths out the engine's power delivery by storing and releasing rotational energy. This application shows why the r^2 term appears in polar form: mass at twice the radius contributes four times as much to the moment of inertia. Engineers use these calculations to design flywheels that achieve the desired balance between energy storage capacity and overall weight.

Moments of inertia appear in several key equations in physics:

- Rotational Kinetic Energy: $E = \frac{1}{2}I\omega^2$, where ω is angular velocity
 - Torque: $\tau = I\alpha$, where α is angular acceleration
 - Angular Momentum: $L = I\omega$
 - Period of Rotation: $T = 2\pi\sqrt{\frac{I}{k}}$ for a torsional oscillator with spring constant k
-

▲ **Example 4.3.2** Elijah the Prophet, a legendary DJ, is designing a custom turntable platter—the rotating disc that holds vinyl records. The platter is shaped like an annular region (a washer shape) with inner radius $r = 1$ inch and outer radius $r = 6$ inches. To ensure smooth playback, the platter’s density increases quadratically with radius according to $\rho(r) = kr^2$ ounces/inch², where k is a constant. If the total mass of the platter is 20 ounces:

1. Find the value of k .
2. Calculate the moment of inertia I_z about the turntable’s axis of rotation.
3. If the turntable spins at $33\frac{1}{3}$ RPM (rotations per minute), find its rotational kinetic energy in inch-ounces.

Solution

1. First, let’s find k using the total mass equation in polar coordinates:

$$M = \int_0^{2\pi} \int_1^6 \rho(r)r \, dr \, d\theta = \int_0^{2\pi} \int_1^6 kr^3 \, dr \, d\theta = 20$$

Evaluating the integrals:

$$\begin{aligned} 2\pi k \left[\frac{r^4}{4} \right]_1^6 &= 20 \\ 2\pi k \left(\frac{1296}{4} - \frac{1}{4} \right) &= 20 \\ 2\pi k \left(324 - \frac{1}{4} \right) &= 20 \end{aligned}$$

Therefore:

$$k = \frac{20}{2\pi(323.75)} \approx 0.0098 \text{ oz/inch}^4$$

2. The moment of inertia about the z -axis is:

$$I_z = \int_0^{2\pi} \int_1^6 r^2 \rho(r) r \, dr \, d\theta = \int_0^{2\pi} \int_1^6 k r^5 \, dr \, d\theta$$

Substituting our value of k :

$$I_z = 2\pi k \left[\frac{r^6}{6} \right]_1^6 = 2\pi \cdot \frac{20}{2\pi(323.75)} \cdot \left(\frac{46656}{6} - \frac{1}{6} \right)$$

$$I_z = \frac{20}{323.75} \cdot 7776 \approx 480 \text{ oz} \cdot \text{inch}^2$$

3. The angular velocity at $33\frac{1}{3}$ RPM is:

$$\omega = \frac{100}{3} \cdot \frac{2\pi}{60} = \frac{10\pi}{9} \text{ rad/s}$$

The rotational kinetic energy is:

$$E = \frac{1}{2} I_z \omega^2 = \frac{1}{2} (480) \left(\frac{10\pi}{9} \right)^2 \approx 2926 \text{ oz} \cdot \text{inch}^2/\text{s}^2$$

This high moment of inertia helps maintain consistent playback speed, as the platter's substantial rotational inertia resists changes in angular velocity. The quadratic density increase toward the outer edge—similar to putting weights around a turntable's perimeter—maximizes this stabilizing effect.

▲ Example 4.3.3 A robotic arm at a manufacturing plant rotates components on a flat mounting plate. The rectangular plate measures $40 \text{ cm} \times 30 \text{ cm}$ and has a base density of $\rho_0 = 2 \text{ g/cm}^2$. To optimize the rotation dynamics, the plate's density varies according to $\rho(x, y) = \rho_0(1 + \frac{x^2}{100}) \text{ g/cm}^2$, where x is measured in centimeters from the plate's vertical centerline. The system rotates with an initial angular velocity of $\omega_0 = 2\pi \text{ rad/s}$ but must decelerate to a stop within time t using a constant torque τ .

1. Calculate the moment of inertia I_z about the central vertical axis.
2. Using the rotational form of Newton's Second Law, $\tau = I\alpha$, find the required torque to bring the plate to rest in 3 seconds.
3. What is the total angular displacement during this deceleration? Use the equation $\theta = \omega_0 t + \frac{1}{2} \alpha t^2$.

Solution

1. Setting up coordinates with the origin at the center of the plate, our region is $R = [-20, 20] \times [-15, 15]$ in centimeters. For I_z :

$$\begin{aligned} I_z &= \iint_R (x^2 + y^2) \rho_0 \left(1 + \frac{x^2}{100}\right) dA \\ &= \rho_0 \int_{-20}^{20} \int_{-15}^{15} (x^2 + y^2) \left(1 + \frac{x^2}{100}\right) dy dx \end{aligned}$$

Breaking this into parts and evaluating:

$$I_z = 2(754,000) = 1,508,000 \text{ g} \cdot \text{cm}^2$$

Converting to SI units:

$$I_z = 1,508,000 \times 10^{-7} = 0.1508 \text{ kg} \cdot \text{m}^2$$

2. For constant deceleration from ω_0 to 0 in time t :

$$\alpha = \frac{\omega_{\text{final}} - \omega_0}{t} = \frac{0 - 2\pi}{3} = -\frac{2\pi}{3} \text{ rad/s}^2$$

Using $\tau = I\alpha$:

$$\tau = 0.1508 \cdot \left(-\frac{2\pi}{3}\right) = -0.316 \text{ N} \cdot \text{m}$$

3. For the total angular displacement:

$$\begin{aligned} \theta &= \omega_0 t + \frac{1}{2} \alpha t^2 = 2\pi(3) + \frac{1}{2} \left(-\frac{2\pi}{3}\right) (3)^2 \\ &= 6\pi - \frac{9\pi}{2} = \frac{3\pi}{2} \text{ rad} = 270^\circ \end{aligned}$$

Average Values over Two-Dimensional Regions

We took a sneak peak of this concept when we looked at average efficiency in the previous lecture – When analyzing quantities that vary over a region, we often need to find their average values. Just as the average of discrete values is found by summing and dividing by the count, the average value of a continuous function $f(x, y)$ over a region R is found by

integrating and dividing by the region's area:

$$f_{avg} = \frac{\iint_R f(x, y) dA}{\iint_R dA}$$

This formula has a natural physical interpretation: the denominator $\iint_R dA$ represents the total area of region R , while the numerator accumulates the value of f over that same region. For example, if $f(x, y)$ represents temperature at each point on a metal plate, this formula gives the plate's average temperature.

For non-uniform materials where density varies across the region, we often use a weighted average:

$$f_{avg} = \frac{\iint_R f(x, y)\rho(x, y) dA}{\iint_R \rho(x, y) dA}$$

Here, $\rho(x, y)$ acts as a weighting function, giving more importance to regions of higher density. This weighted form finds frequent use in engineering applications, from calculating the center of mass of a plate with varying density to determining average stress in composite materials.

Both formulas extend naturally to polar coordinates through the appropriate change of variables, with $dA = r dr d\theta$. This proves particularly useful when working with circular or radially symmetric regions where the function or density varies with distance from a central point.

Surface Area and Physical Interfaces

The motivation for surface area begins with the arc length formula that we learned in Chapter 2. Recall that for a curve $y = f(x)$, we found its arc length by considering how differential changes in x and y contribute to the length of an infinitesimal segment:

$$ds = \sqrt{dx^2 + dy^2} = \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

This gave us the arc length formula $L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$.

For a surface $z = f(x, y)$, we follow similar reasoning. Consider a small patch of surface above a region dA in the xy -plane. As we move in both x and y directions, the surface rises according to its partial derivatives. The actual area of this patch must account for these slopes in both directions, leading to:

$$dA_{\text{surface}} = \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} dA$$

Integrating over the region R gives us the total surface area:

$$A = \iint_R \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} dA$$




Figure 4.17: The surface area element dS arises from the cross product of tangent vectors $\vec{u} = (dx, 0, \frac{\partial f}{\partial x} dx)$ and $\vec{v} = (0, dy, \frac{\partial f}{\partial y} dy)$. Their cross product gives $dS = \|\vec{u} \times \vec{v}\| = \sqrt{1 + \left(\frac{\partial f}{\partial x}\right)^2 + \left(\frac{\partial f}{\partial y}\right)^2} dx dy$.

If our surface is better described using polar coordinates, we can transform this formula. For $z = f(r, \theta)$, the surface area becomes:

$$A = \iint_R \sqrt{1 + \left(\frac{\partial f}{\partial r}\right)^2 + \frac{1}{r^2} \left(\frac{\partial f}{\partial \theta}\right)^2} r dr d\theta$$

Thus, this shows us that surface area is just arc-length in three dimensions. This pattern also generalizes. For a *hypersurface* in n dimensions given by a function of $n - 1$ variables, the hypersurface area becomes:

$$A = \int_R \sqrt{1 + \sum_{i=1}^{n-1} \left(\frac{\partial f}{\partial x_i}\right)^2} dA$$

This generalization finds applications in physics and engineering. The bad thing about these formulae is that the integral is usually too hard to solve, so we will frequently have to rely on numerical methods and/or the all seeing eye: Wolfram Alpha.

Let us now explore these formulae with a concrete example:

▲ **Example 4.3.4** In architecture, hyperbolic paraboloids (saddle surfaces) are used to create striking modern designs that combine aesthetics with structural efficiency. Consider a surface given by $z = \frac{x^2}{4} - \frac{y^2}{9}$ over the region $R = [-1, 1] \times [-2, 2]$.

1. Set up (but do not evaluate) the surface area integral.
2. Use the fact that this integral evaluates to approximately 8.56 square units to

discuss the efficiency of this architectural form.

Solution Let's solve this step by step:

1. First, we compute the partial derivatives:

$$\frac{\partial z}{\partial x} = \frac{x}{2}, \quad \frac{\partial z}{\partial y} = -\frac{2y}{9}$$

The surface area is given by:

$$A = \iint_R \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dA$$

Substituting the partial derivatives:

$$A = \iint_R \sqrt{1 + \frac{x^2}{4} + \frac{4y^2}{81}} dA$$

Since the integrand $\sqrt{1 + \frac{x^2}{4} + \frac{4y^2}{81}}$ is even in both x and y , we can integrate over the first quadrant and multiply by 4:

$$A = 4 \int_0^1 \int_0^2 \sqrt{1 + \frac{x^2}{4} + \frac{4y^2}{81}} dy dx$$

2. The numerical evaluation of this integral yields $A \approx 8.56$ square units. This relatively small surface area compared to the region covered (8 square units in the xy -plane) demonstrates why hyperbolic paraboloids are efficient architectural forms. The surface area is only about 7% more than the flat projection, making them materially economical while their curvature provides structural advantages through geometric stiffness.

Lecture 4.4: Triple Integrals

Having explored integration over regions in the plane, we now extend our understanding to higher dimensions. Recall that double integrals allowed us to find areas under surfaces by integrating a function $f(x, y)$ over a region R , producing outputs in units of “function $\times$ area”. For instance, when integrating a constant height function $f(x, y) = h$ over a region R , we obtained volume: $V = h \iint_R dA$.

Triple integrals follow this pattern but work in three dimensions. When we integrate a function $f(x, y, z)$ over a solid region E in space, our output will have units of “function $\times$ volume”. At first glance, this might suggest we are limited to producing quantities measured in spatial units to the fourth power—a type of hypervolume. However, just as we saw with density functions producing mass ($\text{g/cm}^3 \times \text{cm}^3 = \text{g}$), the integrand can carry rates that combine with volume in various ways.

Consider, for example, the second moment of inertia tensor in engineering, which has units of $\text{kg} \cdot \text{m}^2$. When analyzing rotating bodies, we might integrate $\rho(x, y, z)(y^2 + z^2)$ to find the moment of inertia about the x -axis. Here, the density (kg/m^3) combines with squared distance terms (m^2) and volume (m^3) to produce the correct units. Or in dynamics, when studying jerk (the rate of change of acceleration), we might integrate a function with units m/s^3 over a volume to find total change in momentum rate. The power of triple integrals lies in this flexibility to handle various physical quantities through appropriate choice of integrand, practical implications of which will be explored in more detailed in the coming lectures in this chapter. The good news for us in this introduction is that we will use almost identical machinery to what we used with double integrals – there is not many more layers of complexity to add when generalizing double integrals to triple and beyond.

Definition and Computation

Consider a solid region E in three-dimensional space and a function $f(x, y, z)$ defined over this region. Just as we built double integrals from the concept of summing over rectangles, we construct triple integrals by summing over small rectangular boxes called volume elements. Each volume element has dimensions Δx , Δy , and Δz , giving it a volume of $\Delta V = \Delta x \Delta y \Delta z$.

When constructing our Riemann sum, we must choose a sample point (x_i^*, y_i^*, z_i^*) within each box. Where single integrals required choosing a point on a line segment and double integrals required choosing a point in a rectangle, triple integrals require selecting a point within a rectangular box. This point could be at any corner of the box (there are eight corners), at the center of the box, or at any other location within the box. For each such box with its chosen sample point, we compute $f(x_i^*, y_i^*, z_i^*) \Delta V_i$.

As all dimensions of these boxes approach zero (that is, as Δx , Δy , and Δz all approach

zero), their sum approaches the triple integral:

$$\iiint_E f(x, y, z) dV = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*, y_i^*, z_i^*) \Delta V_i$$

Just as with double integrals, different choices of sample points might give different approximations, but all reasonable choices converge to the same limit as the box dimensions approach zero. The differential volume element $dV = dx dy dz$ represents the limiting case of ΔV as all dimensions approach zero. The units of dV will always be cubic units (cubic meters, cubic centimeters, etc.), while the units of $f(x, y, z)$ could vary depending on the application.

For a rectangular box $E = [a, b] \times [c, d] \times [p, q]$, we can write an iterated integral:

$$\iiint_E f(x, y, z) dV = \int_a^b \int_c^d \int_p^q f(x, y, z) dz dy dx$$

This can be evaluated in any order—Fubini's Theorem extends naturally to three dimensions:

$$\int_a^b \int_c^d \int_p^q f(x, y, z) dz dy dx = \int_p^q \int_c^d \int_a^b f(x, y, z) dx dy dz$$

and similarly for any other permutation of the variables. The choice of order, while not affecting the result, often significantly impacts the complexity of computation.

For example, if we're computing the mass of a solid where density varies with height, integrating with respect to z first might simplify our work. If instead the density varies with distance from an axis, we might prefer cylindrical coordinates—a system we'll explore shortly.

Types of Integration Regions

When integrating over general three-dimensional regions, we must carefully analyze the geometry to determine appropriate bounds of integration. Just as with double integrals, we can often view a region from different perspectives, leading to distinct but equivalent ways to set up our integral. Let's examine the three fundamental types of regions we encounter.

Type I Regions: These regions are defined by viewing the solid from the side, looking perpendicular to the xy -plane. The region is bounded above and below by surfaces $z = g_2(x, y)$ and $z = g_1(x, y)$ respectively, where these surfaces sit above a region R in the xy -plane. For such regions:

$$\iiint_E f(x, y, z) dV = \iint_R \int_{g_1(x, y)}^{g_2(x, y)} f(x, y, z) dz dA$$




Figure 4.18: A *Type I* region bounded by two surfaces above a region R in the xy -plane.

Type II Regions: These regions are viewed by looking perpendicular to the yz -plane, bounded by surfaces $x = h_1(y, z)$ and $x = h_2(y, z)$ over a region R in the yz -plane. Here:

$$\iiint_E f(x, y, z) dV = \iint_R \int_{h_1(y, z)}^{h_2(y, z)} f(x, y, z) dx dA$$





Figure 4.19: A *Type II* region bounded by surfaces viewed from the yz -plane.

Type III Regions: These regions are viewed by looking perpendicular to the xz -plane, bounded by surfaces $y = k_1(x, z)$ and $y = k_2(x, z)$ over a region R in the xz -plane:

$$\iiint_E f(x, y, z) dV = \iint_R \int_{k_1(x, z)}^{k_2(x, z)} f(x, y, z) dy dA$$

Figure 4.20: A Type III region bounded by surfaces viewed from the xz -plane.

▲ Example 4.4.1

1. Find $\iiint_E (x^2 + y^2) dV$ where E is the rectangular box $[0, 1] \times [1, 2] \times [-1, 3]$.
2. Find $\iiint_E xyz dV$ where E is the region bounded below by the xy -plane ($z = 0$) and above by the plane $z = 2 - x - y$ in the first octant.

Solution

1. For the rectangular box:




Figure 4.21: Problem 1: Rectangular box region

$$\begin{aligned}
\iiint_E (x^2 + y^2) dV &= \int_0^1 \int_1^2 \int_{-1}^3 (x^2 + y^2) dz dy dx \\
&= \int_0^1 \int_1^2 (x^2 + y^2)[z]_{-1}^3 dy dx \\
&= 4 \int_0^1 \int_1^2 (x^2 + y^2) dy dx \\
&= 4 \int_0^1 \left[x^2 y + \frac{y^3}{3} \right]_1^2 dx \\
&= 4 \int_0^1 \left(x^2(2-1) + \frac{8}{3} - \frac{1}{3} \right) dx \\
&= 4 \left[\frac{x^3}{3} + \frac{7x}{3} \right]_0^1 = \frac{32}{3}
\end{aligned}$$

2. For the region in the first octant:




Figure 4.22: Problem 2: Region bounded by plane $z = 2 - x - y$ and xy -plane

The bounds are in the first octant bounded below by the xy -plane ($z = 0$) and above by the plane $z = 2 - x - y$. From the xy -plane projection, we see x ranges from 0 to 2, and for each fixed x , y ranges from 0 to $2 - x$. For each fixed x and y , z then ranges from 0 to $2 - x - y$, giving the triple integral's bounds.

$$\iiint_E xyz dV = \int_0^2 \int_0^{2-x} \int_0^{2-x-y} xyz dz dy dx$$

$$\begin{aligned}
&= \int_0^2 \int_0^{2-x} xy \left[\frac{z^2}{2} \right]_0^{2-x-y} dy dx \\
&= \frac{1}{2} \int_0^2 \int_0^{2-x} xy(2-x-y)^2 dy dx
\end{aligned}$$

Let $u = 2 - x - y$, so $y = 2 - x - u$ and $dy = -du$. When $y = 0$, $u = 2 - x$; when $y = 2 - x$, $u = 0$. This gives:

$$\begin{aligned}
&= \frac{1}{2} \int_0^2 x \int_{2-x}^0 (2-x-u)u^2(-du) dx \\
&= \frac{1}{2} \int_0^2 x \left[(2-x)\frac{u^3}{3} - \frac{u^4}{4} \right]_0^{2-x} dx \\
&= \frac{1}{24} \int_0^2 x(2-x)^4 dx = \frac{4}{45}
\end{aligned}$$

▲ **Example 4.4.2** Find $\iiint_E 1 dV$ where E is the region bounded by the intersection of two perpendicular unit cylinders $x^2 + y^2 = 1$ and $x^2 + z^2 = 1$.

Solution




Figure 4.23: The Steinmetz solid, formed by the intersection of two perpendicular unit cylinders $x^2 + y^2 = 1$ and $x^2 + z^2 = 1$.

Due to the symmetry of the region across all coordinate planes, we can evaluate the integral in the first octant and multiply by 8. In the first octant, x ranges from 0 to 1, and for each fixed x , both y and z range from 0 to $\sqrt{1-x^2}$ due to the circular cross-sections of the cylinders (found by solving our two equations for y and z):

$$\begin{aligned}\iiint_E 1 \, dV &= 8 \int_0^1 \int_0^{\sqrt{1-x^2}} \int_0^{\sqrt{1-x^2}} 1 \, dz \, dy \, dx \\ &= 8 \int_0^1 \int_0^{\sqrt{1-x^2}} \sqrt{1-x^2} \, dy \, dx \\ &= 8 \int_0^1 (1-x^2) \, dx \\ &= 8 \left[x - \frac{x^3}{3} \right]_0^1 = 8 \left(1 - \frac{1}{3} \right) = \frac{16}{3}\end{aligned}$$

The choice of which type to use for a given region often depends on the geometry of the bounding surfaces and the nature of the integrand. Many complex regions might require breaking the integral into parts and using different types for each part. Just as with double integrals, the goal is to choose the type that leads to the most straightforward computation.

However, for certain geometries—particularly those with circular or spherical symmetry—we'll find that alternative coordinate systems provide more natural representations. These systems, cylindrical and spherical coordinates, will be our next topic of study.

Coordinate Systems for Triple Integration

The choice of coordinate system can dramatically simplify a triple integral computation. Just as we found polar coordinates useful for double integrals over circular regions, cylindrical and spherical coordinates provide natural frameworks for three-dimensional regions with certain types of symmetry. Let's examine these alternative coordinate systems and understand how they transform our integration process.

Cylindrical Coordinates

As we learned in Chapter 1, cylindrical coordinates extend polar coordinates into three dimensions by adding a height coordinate. A point (x, y, z) in rectangular coordinates is represented by coordinates (r, θ, z) in the cylindrical system:

$$\begin{aligned}x &= r \cos \theta \\ y &= r \sin \theta \\ z &= z\end{aligned}$$

The coordinate r represents the distance from the point to the z -axis, θ measures the angle from the positive x -axis in the xy -plane, and z gives the height above the xy -plane. This system naturally describes surfaces like cylinders, where r is constant, and cones, where z is proportional to r .

When converting a triple integral to cylindrical coordinates, we must account for how volume elements transform. A small rectangular box $dx dy dz$ becomes a curved wedge $r dr d\theta dz$. The factor of r appears for the same reason it did in double integrals—as we move farther from the z -axis, the same angular increment $d\theta$ sweeps out a larger arc. Thus:

$$\iiint_E f(x, y, z) dV = \iiint_E f(r \cos \theta, r \sin \theta, z) r dr d\theta dz$$




Figure 4.24: Representation of a volume element in cylindrical coordinates (r, θ, z) , where r is the radial distance from the z -axis, θ is the azimuthal angle in the xy -plane, and z is the height along the z -axis.

The bounds of integration depend on the region E . Common patterns include:

$$\begin{array}{ll} 0 \leq r \leq R & \text{(solid cylinder)} \\ 0 \leq \theta \leq 2\pi & \text{(full revolution)} \\ a \leq z \leq b & \text{(fixed height)} \end{array}$$

Spherical Coordinates

Spherical coordinates offer an even more radical departure from rectangular coordinates, particularly useful for regions with spherical symmetry. A point (x, y, z) is represented by

coordinates (ρ, θ, ϕ) , where:

$$\begin{aligned}x &= \rho \sin \phi \cos \theta \\y &= \rho \sin \phi \sin \theta \\z &= \rho \cos \phi\end{aligned}$$

Here, ρ represents the distance from the origin to the point, θ measures the angle from the positive x -axis in the xy -plane (as in cylindrical coordinates), and ϕ measures the angle from the positive z -axis. Note that ϕ is often called the polar angle, while θ is the azimuthal angle.

The volume element transformation in spherical coordinates is more complex than in cylindrical coordinates. A small volume element becomes $\rho^2 \sin \phi d\rho d\theta d\phi$. The factor $\rho^2 \sin \phi$ can be understood geometrically:

- The ρ^2 term accounts for the radial growth in three dimensions
- The $\sin \phi$ term accounts for the “thinning” of the volume element near the poles

Therefore, our triple integral becomes:

$$\iiint_E f(x, y, z) dV = \iiint_E f(\rho \sin \phi \cos \theta, \rho \sin \phi \sin \theta, \rho \cos \phi) \rho^2 \sin \phi d\rho d\theta d\phi$$




Figure 4.25: Representation of a volume element in spherical coordinates (ρ, ϕ, θ) , where ρ is the radial distance, ϕ is the azimuthal angle in the xy -plane, and θ is the polar angle from the z -axis. The volume element is defined by $\rho^2 \sin \theta d\rho d\phi d\theta$, illustrating the differential components $d\rho$, $\rho d\theta$, and $\rho \sin \theta d\phi$.

Typical bounds in spherical coordinates follow these patterns:

$$\begin{array}{ll} 0 \leq \rho \leq R & \text{(solid sphere)} \\ 0 \leq \theta \leq 2\pi & \text{(full revolution)} \\ 0 \leq \phi \leq \pi & \text{(full sphere)} \end{array}$$

The choice between cylindrical and spherical coordinates often depends on the region's geometry and the integrand's form. Cylindrical coordinates excel for regions with circular cross-sections parallel to the xy -plane, while spherical coordinates are natural for regions bounded by spheres or cones centered at the origin. Both systems can dramatically simplify integrals that would be challenging in rectangular coordinates, particularly when the integrand involves terms like $\sqrt{x^2 + y^2}$ or $\sqrt{x^2 + y^2 + z^2}$.

▲ Example 4.4.3

1. Find the volume of a right circular cone with height h and radius r using cylindrical coordinates.
2. Evaluate $\iiint_E (x^2 + y^2 + z^2) dV$ where E is the solid sphere of radius 2 centered at the origin.

Solution

1. Let's first visualize what's going on:




Figure 4.26: A right circular cone of height h and radius r . The red dashed circle represents a horizontal cross-section at height z with radius $R(z)$, demonstrating how the radius varies linearly with height.

For a right circular cone, we first need to understand how the radius varies with height. Consider any horizontal cross-section at height z from the base. This creates two similar triangles: the full cone (radius r , height h), and the smaller cone above our cross-section (radius $R(z)$, height $h - z$). By the properties of similar triangles:

$$\frac{R(z)}{r} = \frac{h - z}{h}$$

Solving for $R(z)$, we get $R(z) = \frac{r}{h}(h - z)$. This equation shows that the radius decreases linearly as we move up the cone, becoming zero at the top ($z = h$) and equal to r at the base ($z = 0$). In cylindrical coordinates, we integrate over the angle θ from 0 to 2π as it sweeps around the xy -plane, the height z from 0 to h , and for each fixed z , the radius ρ from 0 to $R(z) = \frac{r}{h}(h - z)$:

$$\begin{aligned} V &= \iiint_E 1 \, dV = \int_0^{2\pi} \int_0^h \int_0^{r(h-z)/h} \rho \, d\rho \, dz \, d\theta \\ &= \int_0^{2\pi} \int_0^h \left[\frac{\rho^2}{2} \right]_0^{r(h-z)/h} dz \, d\theta \\ &= \int_0^{2\pi} \int_0^h \frac{r^2(h-z)^2}{2h^2} dz \, d\theta \\ &= 2\pi \int_0^h \frac{r^2(h-z)^2}{2h^2} dz \\ &= \frac{\pi r^2}{h^2} \int_0^h (h-z)^2 dz \\ &= \frac{\pi r^2}{h^2} \left[-\frac{(h-z)^3}{3} \right]_0^h \\ &= \frac{\pi r^2}{h^2} \left[0 - \left(-\frac{h^3}{3} \right) \right] = \frac{1}{3} \pi r^2 h \end{aligned}$$

2. The integrand $x^2 + y^2 + z^2$ suggests the use of spherical coordinates. Recall the transformations:

$$x = \rho \sin \phi \cos \theta, \quad y = \rho \sin \phi \sin \theta, \quad z = \rho \cos \phi$$

The volume element in spherical coordinates is:

$$dV = \rho^2 \sin \phi \, d\rho \, d\phi \, d\theta$$

For a sphere centered at the origin with radius 2, we integrate over the angle θ from 0 to 2π , the polar angle ϕ from 0 to π to cover the full sphere north to south, and the radius ρ from 0 to 2. Thus our integral becomes:

$$\begin{aligned}
 \iiint_E (x^2 + y^2 + z^2) dV &= \iiint_E \rho^2 dV \\
 &= \int_0^{2\pi} \int_0^\pi \int_0^2 \rho^2 \cdot \rho^2 \sin \phi d\rho d\phi d\theta \\
 &= \int_0^{2\pi} \int_0^\pi \int_0^2 \rho^4 \sin \phi d\rho d\phi d\theta \\
 &= \int_0^{2\pi} \int_0^\pi \left[\frac{\rho^5}{5} \right]_0^2 \sin \phi d\phi d\theta \\
 &= \int_0^{2\pi} \int_0^\pi \frac{32}{5} \sin \phi d\phi d\theta \\
 &= \frac{32}{5} \int_0^{2\pi} [-\cos \phi]_0^\pi d\theta \\
 &= \frac{32}{5} \int_0^{2\pi} [(-\cos \pi) - (-\cos 0)] d\theta = \frac{128\pi}{5}
 \end{aligned}$$

▲ **Example 4.4.4** An engineer is designing a cylindrical support beam that needs to be contained entirely within a spherical dome. If the dome has radius R and the beam has radius a (where $a < R$), find the volume of the portion of the beam that lies within the dome.

Mathematically, find the volume of the region bounded by:

- Sphere: $x^2 + y^2 + z^2 = R^2$
- Cylinder: $x^2 + y^2 = a^2$

Solution The symmetry of this region suggests using cylindrical coordinates. First, let's understand our bounds:

In cylindrical coordinates:

- The cylinder equation becomes $r = a$

- The sphere equation becomes $r^2 + z^2 = R^2$, which gives us $z = \pm\sqrt{R^2 - r^2}$

Therefore, our volume integral will have:

- $0 \leq r \leq a$ (from center to cylinder wall)
- $0 \leq \theta \leq 2\pi$ (full circular cross-section)
- $-\sqrt{R^2 - r^2} \leq z \leq \sqrt{R^2 - r^2}$ (sphere bounds)




Figure 4.27: *Intersection of a sphere (radius R) and cylinder (radius a). The gray circles represent the sphere's cross-sections, the blue lines show the cylinder, and the red circles indicate where the cylinder intersects the sphere at heights $\pm\sqrt{R^2 - a^2}$. This intersection forms the boundary of our integration region.*

Now we can set up our integral:

$$\begin{aligned}
 V &= \iiint_E 1 \, dV = \int_0^{2\pi} \int_0^a \int_{-\sqrt{R^2 - r^2}}^{\sqrt{R^2 - r^2}} r \, dz \, dr \, d\theta \\
 &= \int_0^{2\pi} \int_0^a r [z]_{-\sqrt{R^2 - r^2}}^{\sqrt{R^2 - r^2}} \, dr \, d\theta
 \end{aligned}$$

$$= \int_0^{2\pi} \int_0^a 2r\sqrt{R^2 - r^2} \, dr \, d\theta$$

For the r integral, let $u = R^2 - r^2$, so $du = -2r \, dr$:

$$\begin{aligned} V &= 2\pi \int_0^a 2r\sqrt{R^2 - r^2} \, dr \\ &= -2\pi \int_{R^2}^{R^2 - a^2} \sqrt{u} \, du \\ &= -2\pi \left[\frac{2u^{3/2}}{3} \right]_{R^2}^{R^2 - a^2} \\ &= \frac{4\pi}{3} [R^3 - (R^2 - a^2)^{3/2}] \end{aligned}$$

This formula tells us that when $a = R$, the volume becomes $\frac{4\pi R^3}{3}$ (full sphere), and when a approaches 0, the volume approaches 0 (infinitely thin beam), as expected.

Lecture 4.5: Change of Variables in Multiple Integrals

Throughout our study of multiple integrals, we've seen how choosing the right coordinate system can dramatically simplify our calculations. When working with circular regions, polar coordinates turned our complicated bounds into simple inequalities. For cylindrical shells, cylindrical coordinates made our integration straightforward. And for spherical objects, spherical coordinates proved invaluable.

Let's examine what actually happens in these coordinate transformations. When we switch from rectangular coordinates (x, y) to polar coordinates (r, θ) , we use the equations:

$$\begin{aligned}x &= r \cos \theta \\y &= r \sin \theta\end{aligned}$$

And when we make this change, we learned that our area element transforms from $dA = dx dy$ to $dA = r dr d\theta$. Similarly, in spherical coordinates, our volume element becomes $dV = \rho^2 \sin \phi d\rho d\theta d\phi$. These patterns suggest a deeper principle at work—one that explains why these particular factors appear and generalizes to any coordinate transformation.

Just as in single-variable calculus, where substitution requires adjusting our differential element:

$$\int f(x) dx = \int f(g(t))g'(t) dt$$

we need a way to adjust our area and volume elements in multiple dimensions. This adjustment, which we'll find involves determinants of matrices of partial derivatives, provides a powerful tool for handling integrals over more general regions.

The Jacobian Determinant

Recall from Chapter 3, Lecture 7 and 8 we briefly mentioned matrices and their determinants. Think of moving vectors around in space - we might stretch them, rotate them, or both. When we do these operations, each component (x and y) of our vector changes in a way that depends on both original components. A matrix organizes these relationships. For instance, a 2×2 matrix A has the form:

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

Associated with every square matrix is a number called its determinant. For a 2×2 matrix, this determinant is:

$$\det(A) = a_{11}a_{22} - a_{12}a_{21}$$

For a 3×3 matrix, we can compute the determinant by expanding along any row or column. For example, expanding along the first row:

$$\det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = a_{11} \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$$

A linear transformation is a function that preserves addition and scaling—for example, the transformation that rotates all points by a fixed angle or scales all distances by a fixed factor. The determinant of a linear transformation’s matrix tells us exactly how that transformation scales areas or volumes. If the determinant is 2, areas are doubled; if it’s $1/3$, volumes are reduced to one-third their original size.




Figure 4.28: A visualization of how the Jacobian determinant measures area scaling under a transformation. When mapping from region Z to region X , the determinant of the inverse transformation (J_f^{-1}) tells us the factor by which areas are scaled. Different parts of the region may be scaled by different factors, as shown by the red ($\det J_f^{-1} = \frac{1}{4}$) and green ($\det J_f^{-1} = -\frac{1}{4}$) regions. The negative determinant indicates a reflection.

When we transform from one coordinate system to another, we’re generally making a nonlinear transformation—like our polar coordinate transformation where $x = r \cos \theta$ and $y = r \sin \theta$. However, at each point, we can approximate this nonlinear transformation by a linear one that tells us how small regions near that point are transformed. This linear approximation is given by the Jacobian matrix, and its determinant tells us the local scaling factor for areas or volumes.

Consider a transformation that maps points (u, v) in a region R' to points (x, y) in a region R :

$$\begin{aligned}x &= x(u, v) \\ y &= y(u, v)\end{aligned}$$

The Jacobian matrix contains all the partial derivatives describing how small changes in (u, v) affect (x, y) :

$$J = \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix}$$

To understand why this matrix appears, consider what happens to a small rectangular region in the uv -plane with sides of length du and dv . When we transform to xy -coordinates:

- A step of du in the u -direction changes x by $\frac{\partial x}{\partial u} du$ and y by $\frac{\partial y}{\partial u} du$
- A step of dv in the v -direction changes x by $\frac{\partial x}{\partial v} dv$ and y by $\frac{\partial y}{\partial v} dv$

These transformed vectors define a parallelogram in the xy -plane whose area is given by the absolute value of the determinant of J times $du dv$. This is exactly what we need to convert our area element from $du dv$ to $dx dy$.

The Jacobian determinant, denoted $|J|$ or $\left| \frac{\partial(x,y)}{\partial(u,v)} \right|$, is therefore:

$$|J| = \det \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix} = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$

This same principle extends to three dimensions. For transformations mapping (u, v, w) to (x, y, z) , the Jacobian determinant becomes:

$$|J| = \det \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{bmatrix}$$

This determinant, computed by the expansion rule along any row or column, gives us the local scaling factor for volumes under our transformation. It's this factor that explains the mysterious terms (r and $\rho^2 \sin \phi$) that appeared in our coordinate transformations earlier in the chapter. Let's collect all the coordinate transformations we've studied, showing both the transformations and their associated Jacobian matrices and determinants:

Polar Coordinates:

$$\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \end{aligned}$$

$$J = \begin{bmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{bmatrix}, \quad |J| = r, \quad dA = r dr d\theta$$

Cylindrical Coordinates:

$$\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \\ z &= z \end{aligned}$$

$$J = \begin{bmatrix} \cos \theta & -r \sin \theta & 0 \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad |J| = r, \quad dV = r dr d\theta dz$$

Spherical Coordinates:

$$\begin{aligned}x &= \rho \sin \phi \cos \theta \\y &= \rho \sin \phi \sin \theta \\z &= \rho \cos \phi\end{aligned}$$

$$J = \begin{bmatrix} \sin \phi \cos \theta & \rho \cos \phi \cos \theta & -\rho \sin \phi \sin \theta \\ \sin \phi \sin \theta & \rho \cos \phi \sin \theta & \rho \sin \phi \cos \theta \\ \cos \phi & -\rho \sin \phi & 0 \end{bmatrix}, \quad |J| = \rho^2 \sin \phi, \quad dV = \rho^2 \sin \phi \, d\rho \, d\theta \, d\phi$$

▲ **Example 4.5.1** Compute the Jacobian determinant for toroidal coordinates, where the transformation is given by:

$$\begin{aligned}x &= \frac{R \sinh \tau}{\cosh \tau - \cos \sigma} \cos \phi \\y &= \frac{R \sinh \tau}{\cosh \tau - \cos \sigma} \sin \phi \\z &= \frac{R \sin \sigma}{\cosh \tau - \cos \sigma}\end{aligned}$$

Here, R represents the distance from the origin to the center of the tube, τ and σ are the toroidal coordinates, and ϕ is the azimuthal angle in the xy -plane.

Solution To find the Jacobian determinant, we first compute all partial derivatives and organize them in our matrix:

$$J = \begin{bmatrix} \frac{\partial x}{\partial \tau} & \frac{\partial x}{\partial \sigma} & \frac{\partial x}{\partial \phi} \\ \frac{\partial y}{\partial \tau} & \frac{\partial y}{\partial \sigma} & \frac{\partial y}{\partial \phi} \\ \frac{\partial z}{\partial \tau} & \frac{\partial z}{\partial \sigma} & \frac{\partial z}{\partial \phi} \end{bmatrix}$$

Computing each partial derivative:

$$\begin{aligned}\frac{\partial x}{\partial \tau} &= \frac{R \cosh \tau (\cosh \tau - \cos \sigma) - R \sinh \tau \sinh \tau}{(\cosh \tau - \cos \sigma)^2} \cos \phi = \frac{R}{\cosh \tau - \cos \sigma} \cos \phi \\ \frac{\partial x}{\partial \sigma} &= \frac{R \sinh \tau \sin \sigma}{(\cosh \tau - \cos \sigma)^2} \cos \phi \\ \frac{\partial x}{\partial \phi} &= -\frac{R \sinh \tau}{\cosh \tau - \cos \sigma} \sin \phi \\ \frac{\partial y}{\partial \tau} &= \frac{R}{\cosh \tau - \cos \sigma} \sin \phi \\ \frac{\partial y}{\partial \sigma} &= \frac{R \sinh \tau \sin \sigma}{(\cosh \tau - \cos \sigma)^2} \sin \phi\end{aligned}$$

$$\begin{aligned}
\frac{\partial y}{\partial \phi} &= \frac{R \sinh \tau}{\cosh \tau - \cos \sigma} \cos \phi \\
\frac{\partial z}{\partial \tau} &= \frac{R \sin \sigma \sinh \tau}{(\cosh \tau - \cos \sigma)^2} \\
\frac{\partial z}{\partial \sigma} &= \frac{R \cos \sigma (\cosh \tau - \cos \sigma) + R \sin \sigma \sin \sigma}{(\cosh \tau - \cos \sigma)^2} = \frac{R(\cosh \tau)}{(\cosh \tau - \cos \sigma)^2} \\
\frac{\partial z}{\partial \phi} &= 0
\end{aligned}$$

Therefore, the Jacobian matrix is:

$$J = \begin{bmatrix} \frac{R}{\cosh \tau - \cos \sigma} \cos \phi & \frac{R \sinh \tau \sin \sigma}{(\cosh \tau - \cos \sigma)^2} \cos \phi & -\frac{R \sinh \tau}{\cosh \tau - \cos \sigma} \sin \phi \\ \frac{R}{\cosh \tau - \cos \sigma} \sin \phi & \frac{R \sinh \tau \sin \sigma}{(\cosh \tau - \cos \sigma)^2} \sin \phi & \frac{R \sinh \tau}{\cosh \tau - \cos \sigma} \cos \phi \\ \frac{R \sin \sigma \sinh \tau}{(\cosh \tau - \cos \sigma)^2} & \frac{R \cosh \tau}{(\cosh \tau - \cos \sigma)^2} & 0 \end{bmatrix}$$

To compute the determinant $|J|$, we can factor out $\frac{R}{(\cosh \tau - \cos \sigma)^2}$ from the first two rows and the first two columns, simplifying the calculation:

$$|J| = \frac{R^2 \sinh \tau}{(\cosh \tau - \cos \sigma)^3}$$

Therefore, the volume element in toroidal coordinates is:

$$dV = \frac{R^2 \sinh \tau}{(\cosh \tau - \cos \sigma)^3} d\tau d\sigma d\phi$$

The Change of Variables Formula

With the Jacobian determinant in hand, we can state the change of variables formula. For double integrals:

$$\iint_R f(x, y) dx dy = \iint_{R'} f(x(u, v), y(u, v)) |J| du dv$$

For triple integrals:

$$\iiint_E f(x, y, z) dx dy dz = \iiint_{E'} f(x(u, v, w), y(u, v, w), z(u, v, w)) |J| du dv dw$$

The absolute value in these formulas ensures our integral remains positive even when our transformation involves a reflection or orientation change. The formula tells us that to change variables, we:

1. Express the original variables in terms of the new ones

2. Substitute these expressions into the integrand
3. Multiply by the absolute value of the Jacobian determinant
4. Express the region bounds in terms of the new variables

Techniques for Finding Transformations

When faced with a complicated region of integration, several systematic approaches can help identify useful transformations:

1. **Identify the Region Type:** Look at the equations defining your region's boundaries:
 - Quadratic terms like $x^2 + y^2$ suggest circular regions
 - Terms like $\frac{x^2}{a^2} + \frac{y^2}{b^2}$ suggest elliptical regions
 - Linear terms like $ax + by = c$ indicate straight lines
2. **Look for Patterns:** Common boundary equations suggest specific transformations:
 - For ellipses $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, try $x = au$, $y = bv$
 - For rotated regions, try $x = u \cos \theta - v \sin \theta$, $y = u \sin \theta + v \cos \theta$
 - For skewed regions, try $x = u + kv$, $y = v$




Figure 4.29: Transforming an elliptical region $\frac{x^2}{9} + y^2 = 1$ into a unit circle $u^2 + v^2 = 1$ using the transformation $x = 3u$, $y = v$. The Jacobian determinant $|J| = 3$ accounts for the area scaling.

When applying these transformations, verify that:

- The boundary equations simplify in the new coordinates
- The Jacobian determinant is nonzero in your region
- The transformed region has simpler bounds for integration

Conditions for Valid Transformations

Not every transformation that simplifies our region or integrand is valid for change of variables. For the formula to work, our transformation must be:

- One-to-one (each point in R corresponds to exactly one point in R')
- Continuously differentiable (all partial derivatives exist and are continuous)
- Have a nonzero Jacobian determinant throughout the region

These conditions ensure our transformation preserves the essential structure of the region and allows us to meaningfully compute the transformed integral. They also guarantee that our transformation has an inverse, allowing us to map back to our original coordinates if needed.

The nonzero Jacobian condition is particularly important—it ensures our transformation doesn’t “collapse” any part of the region, which would make the transformation fail to be one-to-one. This explains why polar coordinates have a singularity at the origin (where $r = 0$ and the Jacobian vanishes) and why we must handle such points carefully.

▲ **Example 4.5.2** Consider the integral $\iint_R (x+y)^2 dx dy$ where R is the parallelogram with vertices at $(0,0)$, $(2,0)$, $(1,2)$, and $(3,2)$. Use a linear transformation to convert this to an integral over a rectangle.

Solution First, let’s understand how to find an appropriate transformation. Looking at our parallelogram:




Figure 4.30: We seek a transformation $x = f(u,v)$, $y = g(u,v)$ that “un-shears” the parallelogram into a rectangle by shifting each horizontal slice back by an amount proportional to its height.

To find our transformation, we observe that each horizontal line in our parallelogram is shifted right as we move up. Specifically:

- At $y = 0$, the line segment goes from $x = 0$ to $x = 2$

- At $y = 2$, the line segment goes from $x = 1$ to $x = 3$

This suggests each horizontal line is shifted right by half its height. To undo this shift, we need to subtract $\frac{y}{2}$ from each x -coordinate. Therefore, if we let:

$$\begin{aligned}u &= x - \frac{y}{2} \\ y &= v\end{aligned}$$

Solving for x and y gives our transformation:

$$\begin{aligned}x &= u + \frac{v}{2} \\ y &= v\end{aligned}$$

This transformation maps the parallelogram to a rectangle with vertices at $(0, 0)$, $(2, 0)$, $(0, 2)$, and $(2, 2)$. To verify this:

- When $(u, v) = (0, 0)$: $(x, y) = (0, 0)$
- When $(u, v) = (2, 0)$: $(x, y) = (2, 0)$
- When $(u, v) = (0, 2)$: $(x, y) = (1, 2)$
- When $(u, v) = (2, 2)$: $(x, y) = (3, 2)$

The Jacobian matrix for this transformation is:

$$J = \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix} = \begin{bmatrix} 1 & \frac{1}{2} \\ 0 & 1 \end{bmatrix}$$

Therefore:

$$|J| = \det \begin{bmatrix} 1 & \frac{1}{2} \\ 0 & 1 \end{bmatrix} = 1$$

The integrand becomes:

$$(x + y)^2 = \left(u + \frac{v}{2} + v\right)^2 = \left(u + \frac{3v}{2}\right)^2$$

Thus our transformed integral is:

$$\begin{aligned}\iint_R (x + y)^2 dx dy &= \iint_{R'} \left(u + \frac{3v}{2}\right)^2 \cdot 1 du dv \\ &= \int_0^2 \int_0^2 \left(u + \frac{3v}{2}\right)^2 du dv \\ &= \int_0^2 \int_0^2 \left(u^2 + 3uv + \frac{9v^2}{4}\right) du dv\end{aligned}$$

$$\begin{aligned}
&= \int_0^2 \left[\frac{u^3}{3} + \frac{3u^2v}{2} + \frac{9v^2u}{4} \right]_0^2 dv \\
&= \int_0^2 \left(\frac{8}{3} + 6v + \frac{9v^2}{2} \right) dv \\
&= \left[\frac{8v}{3} + 3v^2 + \frac{3v^3}{2} \right]_0^2 \\
&= \frac{16}{3} + 12 + 12 = \frac{88}{3}
\end{aligned}$$

Note that $|J| = 1$ made this transformation particularly nice, as we didn't need to multiply by any extra factors. The transformation simplified our region (making it rectangular) while keeping the integrand relatively manageable.

▲ **Example 4.5.3** An astronomer is studying the density of a spiral galaxy. The density at any point is given by

$$\rho(x, y) = e^{-\left(\frac{x^2}{4} + y^2\right)} e^{-\tan^{-1}\left(\frac{2y}{x}\right)}$$

where (x, y) represents the position in kiloparsecs from the galactic center. Find the total mass contained within the elliptical region bounded by $\frac{x^2}{4} + y^2 = 1$ using appropriate changes of variables.

Solution Let's first understand our region of integration. We have an ellipse defined by $\frac{x^2}{4} + y^2 = 1$. To simplify the integration, we'll transform this ellipse into a unit circle through a change of variables. Ellipse to circle transformations are not too hard. we just need to make the coefficient on x^2 to be 1 and transform y to v .

Consider the linear transformation:

$$\begin{aligned}
x &= 2u \\
y &= v
\end{aligned}$$

We will have the same sort of region transformation as seen in Figure 4.29. When we substitute these into the equation of our ellipse:

$$\frac{(2u)^2}{4} + v^2 = \frac{4u^2}{4} + v^2 = u^2 + v^2 = 1$$

we verify that our elliptical region transforms into the unit circle $u^2 + v^2 = 1$.

The Jacobian matrix for this transformation is:

$$J = \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$$

Therefore, $|J| = 2$. Under this transformation, the density function becomes:

$$\rho(u, v) = e^{-(u^2+v^2)} e^{-\tan^{-1}(\frac{v}{u})}$$

To further simplify integration over the unit circle, we can use polar coordinates:

$$\begin{aligned} u &= r \cos \theta \\ v &= r \sin \theta \end{aligned}$$

The Jacobian for this polar transformation is r . Substituting these into our density function:

$$\rho(r, \theta) = e^{-r^2} e^{-\theta}$$

Combining both changes of variables with their respective Jacobians, our mass integral becomes:

$$M = 2 \int_0^{2\pi} \int_0^1 e^{-r^2} e^{-\theta} r \, dr \, d\theta$$

This integral is set up in a way that separates the variables, potentially simplifying evaluation:

$$M = 2 \left(\int_0^{2\pi} e^{-\theta} \, d\theta \right) \left(\int_0^1 e^{-r^2} r \, dr \right)$$

While the integral over r can be evaluated using substitution, the integral over θ is straightforward:

$$\int_0^{2\pi} e^{-\theta} \, d\theta = [-e^{-\theta}]_0^{2\pi} = -(e^{-2\pi} - 1) = 1 - e^{-2\pi}$$

The integral over r can be evaluated using substitution $w = r^2$, $dw = 2r \, dr$:

$$\int_0^1 e^{-r^2} r \, dr = \frac{1}{2} \int_0^1 e^{-w} \, dw = \frac{1}{2} [-e^{-w}]_0^1 = \frac{1}{2}(1 - e^{-1})$$

Therefore, the total mass is:

$$M = 2 (1 - e^{-2\pi}) \left(\frac{1}{2} (1 - e^{-1}) \right) = (1 - e^{-2\pi}) (1 - e^{-1})$$

Lecture 4.6: Applications of Triple Integrals

Having developed the machinery for evaluating triple integrals in various coordinate systems, we now explore how these tools illuminate real-world physical phenomena. Just as double integrals helped us analyze flat objects with varying properties, triple integrals allow us to study three-dimensional bodies and their characteristics. The applications we'll encounter demonstrate how calculus provides precise mathematical descriptions of physical reality.

Mass and Density in Three Dimensions

Consider a solid object occupying a region E in space. Like we described in Lecture 3 of this chapter, real materials often have density that varies from point to point. A wooden beam might have varying density due to its grain structure; a block of metal might have impurities or intentional composition gradients; even the atmosphere has density that changes with altitude. We can represent this variation using a density function $\rho(x, y, z)$ that gives the mass per unit volume at each point.

To find the total mass of such an object, we follow the same reasoning that led us to double integrals: we partition our region into small volume elements, multiply each element's volume by its density, and sum over all elements. In the limit, this process yields:

$$M = \iiint_E \rho(x, y, z) dV$$

This integral emerges naturally from the physical principle that mass is additive—the total mass is the sum of the masses of all small pieces. Each infinitesimal piece has volume dV and density $\rho(x, y, z)$, giving it mass $\rho(x, y, z) dV$. The triple integral then sums these contributions over the entire region.

Center of Mass and Balance

Just as with flat plates, understanding where a three-dimensional object balances is crucial for engineering applications. The center of mass of a solid object tells us where it effectively balances and how it will behave when rotating. For an object with density $\rho(x, y, z)$, we find its center of mass coordinates $(\bar{x}, \bar{y}, \bar{z})$ using:

$$\begin{aligned}\bar{x} &= \frac{1}{M} \iiint_E x \rho(x, y, z) dV \\ \bar{y} &= \frac{1}{M} \iiint_E y \rho(x, y, z) dV \\ \bar{z} &= \frac{1}{M} \iiint_E z \rho(x, y, z) dV\end{aligned}$$

These formulas arise from the physical principle of torque balance. Each point in the object creates a torque proportional to both its mass and its distance from the axis. The numerator integrals sum up these torque contributions, while division by the total mass M gives us

the balancing point. This generalizes our earlier work with double integrals to fully three-dimensional scenarios.



Figure 4.31: A balancing bird toy demonstrates the principle of center of mass. While often called the center of gravity, these terms are equivalent on Earth where gravitational field strength is approximately uniform. The bird's center of mass is carefully positioned below its support point (the beak) via a non-uniform density throughout the toy, creating a stable equilibrium. When disturbed, the bird will oscillate around this equilibrium point until returning to its stable position—a principle fundamental to both engineering design and physical analysis.

▲ **Example 4.6.1** A metal foundry is producing a cylindrical rod with a height of $L = 1$ meter and a radius of $R = 0.5$ meters. Due to the cooling process, the density of the rod varies with height according to $\rho(x, y, z) = kz^2$, where z measures the height from the base of the cylinder. The total mass of the rod is measured to be $M = 10$ kg.

1. Determine the value of the constant k that describes the density distribution.
2. Calculate the z -coordinate of the center of mass of the rod from the base.

Solution

1. **Determine k :**

The total mass of the rod is given by:

$$M = \iiint_E \rho(x, y, z) \, dV.$$

In cylindrical coordinates, $x^2 + y^2 = r^2$, and $dV = r dr d\theta dz$. The density becomes $\rho(r, \theta, z) = kz^2$, and the region E is defined by $r \in [0, R]$, $\theta \in [0, 2\pi]$, and $z \in [0, L]$. Substituting:

$$M = \int_0^L \int_0^{2\pi} \int_0^R kz^2 r dr d\theta dz.$$

Separate the integrals:

$$M = k \int_0^L z^2 dz \int_0^{2\pi} d\theta \int_0^R r dr.$$

Compute each integral:

$$\int_0^{2\pi} d\theta = 2\pi, \quad \int_0^R r dr = \left[\frac{r^2}{2} \right]_0^R = \frac{R^2}{2}, \quad \int_0^L z^2 dz = \left[\frac{z^3}{3} \right]_0^L = \frac{L^3}{3}.$$

Substitute into the equation for M :

$$M = k \cdot \frac{L^3}{3} \cdot 2\pi \cdot \frac{R^2}{2}.$$

Simplify:

$$M = k \cdot \frac{\pi R^2 L^3}{3}.$$

Solve for k :

$$k = \frac{3M}{\pi R^2 L^3}.$$

Substitute $M = 10$, $R = 0.5$, and $L = 1$:

$$k = \frac{3 \cdot 10}{\pi \cdot (0.5)^2 \cdot 1^3} = \frac{30}{\pi \cdot 0.25} = \frac{120}{\pi}.$$

2. Find the center of mass:

The z -coordinate of the center of mass is:

$$\bar{z} = \frac{1}{M} \iiint_E z \rho(x, y, z) dV.$$

In cylindrical coordinates:

$$\bar{z} = \frac{1}{M} \int_0^L \int_0^{2\pi} \int_0^R z(kz^2)r dr d\theta dz.$$

Separate the integrals:

$$\bar{z} = \frac{k}{M} \int_0^L z^3 dz \int_0^{2\pi} d\theta \int_0^R r dr.$$

Compute each integral:

$$\int_0^{2\pi} d\theta = 2\pi, \quad \int_0^R r \, dr = \frac{R^2}{2}, \quad \int_0^L z^3 \, dz = \left[\frac{z^4}{4} \right]_0^L = \frac{L^4}{4}.$$

Substitute:

$$\bar{z} = \frac{k}{M} \cdot \frac{L^4}{4} \cdot 2\pi \cdot \frac{R^2}{2}.$$

Simplify:

$$\bar{z} = \frac{k\pi R^2 L^4}{4M}.$$

Substituting $k = \frac{120}{\pi}$, $M = 10$, $R = 0.5$, and $L = 1$:

$$\bar{z} = \frac{\frac{120}{\pi} \cdot \pi \cdot (0.5)^2 \cdot 1^4}{4 \cdot 10}.$$

Simplify:

$$\bar{z} = \frac{120 \cdot 0.25}{40} = \frac{30}{40} = 0.75 \text{ meters}.$$

The center of mass of the rod is located 0.75 meters from the base.

Moments and Products of Inertia in Three Dimensions

When an object rotates, its resistance to changes in rotational motion depends not just on its total mass but on how that mass is distributed. Think of a figure skater pulling in their arms to spin faster, or a diver tucking into a ball to increase their rotation speed. These effects are described by quantities called moments and products of inertia.

For rotation about the coordinate axes, we define the moments of inertia:

$$\begin{aligned} I_x &= \iiint_E (y^2 + z^2) \rho(x, y, z) \, dV \\ I_y &= \iiint_E (x^2 + z^2) \rho(x, y, z) \, dV \\ I_z &= \iiint_E (x^2 + y^2) \rho(x, y, z) \, dV \end{aligned}$$

Each moment measures how hard it is to rotate the object about that axis. Notice the squared terms - mass farther from the axis contributes more to the moment of inertia, just as it's harder to spin a dumbbell held at arm's length than one held close to your body.

The products of inertia capture how rotation about one axis affects rotation about others:

$$I_{xy} = \iiint_E xy \rho(x, y, z) \, dV$$

$$I_{yz} = \iiint_E yz\rho(x, y, z) dV$$

$$I_{xz} = \iiint_E xz\rho(x, y, z) dV$$

When an object rotates, its resistance to changes in rotational motion depends not just on its total mass but on how that mass is distributed. This distribution is characterized by moments and products of inertia that we can arrange in a special 3×3 grid called the inertia tensor:

$$\begin{bmatrix} I_{xx} & -I_{xy} & -I_{xz} \\ -I_{yx} & I_{yy} & -I_{yz} \\ -I_{zx} & -I_{zy} & I_{zz} \end{bmatrix}$$

Though this appears as a matrix (a 2-dimensional array of numbers), it represents something more fundamental in physics. Just as vectors (1-dimensional arrays) are special cases of matrices, matrices themselves are special cases of tensors - mathematical objects that preserve relationships between physical quantities regardless of how we measure them. Consider a spinning top - its rotational behavior remains the same whether we measure it using xyz coordinates or any other system. The inertia tensor $\mathbf{I}$ captures this fundamental behavior by relating angular velocity to angular momentum in a way that remains true under any coordinate transformation. While the mathematics of tensors extends beyond this text, we can work with $\mathbf{I}$ as a matrix while appreciating its deeper physical significance.

This structure explains why there exist three special directions called principal axes where rotation is simplest - imagine spinning a book along its length, width, or thickness. When rotating about these axes, the object spins cleanly without wobbling. While finding these directions involves concepts from linear algebra that we'll encounter in later courses, understanding their existence helps explain why gymnasts and divers choose particular axes for their rotations.

These physical insights manifest in several fundamental equations. For any rotating object in 3D space, with angular velocity $\boldsymbol{\omega} = (\omega_x, \omega_y, \omega_z)$ and angular acceleration $\boldsymbol{\alpha} = (\alpha_x, \alpha_y, \alpha_z)$, we can describe its motion using the inertia tensor:

- Angular momentum:

$$\begin{bmatrix} L_x \\ L_y \\ L_z \end{bmatrix} = \begin{bmatrix} I_{xx} & -I_{xy} & -I_{xz} \\ -I_{yx} & I_{yy} & -I_{yz} \\ -I_{zx} & -I_{zy} & I_{zz} \end{bmatrix} \begin{bmatrix} \omega_x \\ \omega_y \\ \omega_z \end{bmatrix}$$

- Rotational kinetic energy:

$$E = \frac{1}{2} \begin{bmatrix} \omega_x & \omega_y & \omega_z \end{bmatrix} \begin{bmatrix} I_{xx} & -I_{xy} & -I_{xz} \\ -I_{yx} & I_{yy} & -I_{yz} \\ -I_{zx} & -I_{zy} & I_{zz} \end{bmatrix} \begin{bmatrix} \omega_x \\ \omega_y \\ \omega_z \end{bmatrix}$$

- Torque:

$$\begin{bmatrix} \tau_x \\ \tau_y \\ \tau_z \end{bmatrix} = \begin{bmatrix} I_{xx} & -I_{xy} & -I_{xz} \\ -I_{yx} & I_{yy} & -I_{yz} \\ -I_{zx} & -I_{zy} & I_{zz} \end{bmatrix} \begin{bmatrix} \alpha_x \\ \alpha_y \\ \alpha_z \end{bmatrix}$$

When an object is symmetric or rotates about a principal axis, the off-diagonal terms (I_{xy}, I_{yz}, I_{xz}) become zero and these equations simplify to the scalar forms seen earlier. Common examples include a cylinder spinning about its axis or a sphere rotating about any diameter.

These matrix operations work as follows:

Matrix-Vector Multiplication: When multiplying a 3×3 matrix by a 3×1 vector:

$$\begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} ax + by + cz \\ dx + ey + fz \\ gx + hy + iz \end{bmatrix}$$

Vector-Matrix-Vector Multiplication: For a row vector times matrix times column vector:

$$\begin{bmatrix} x & y & z \end{bmatrix} \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = ax^2 + (b + d)xy + (c + g)xz + (e)y^2 + (f + h)yz + (i)z^2$$

This explains why the kinetic energy expression $E = \frac{1}{2} \boldsymbol{\omega}^T \mathbf{I} \boldsymbol{\omega}$ produces a scalar value - it's a vector-matrix-vector product that combines all the moments and products of inertia with the angular velocity components. The T over $\boldsymbol{\omega}$ represents the *transpose* which switches the component indices, ie, x_{ij} becomes x_{ji} . Note that this can also change the structure, especially seen in the kinetic energy expression where the dimensions of $\boldsymbol{\omega}$ are 3×1 where $\boldsymbol{\omega}^T$ has dimensions 1×3 which are not the same and have different structural meanings.




Figure 4.32: A visualization of matrix operations. Notice the change in dimensionality. For any matrix multiplication we will have $A_{m \times n} \cdot B_{n \times p} = C_{m \times p}$. Note that the column dimension of A must equal the row dimension of B or else you cannot perform the matrix operation

Matrices naturally emerge in multivariate calculus as they elegantly encode how quantities in multiple dimensions relate to and transform each other, just as we've seen with moments

of inertia describing rotational motion. While we've only scratched the surface here, a deeper study of linear algebra reveals powerful tools for understanding everything from coordinate transformations to optimization problems in higher dimensions. As you continue your mathematical journey, we strongly encourage exploring linear algebra to gain deeper insight into the beautiful structures underlying multivariate calculus.

Force and Work in Space

When a force varies continuously throughout a region of space, triple integrals help us calculate its effects. Consider a force field $\mathbf{F}(x, y, z)$ acting on a solid object. Just as with mass distribution, we can integrate the components of the force field over the region to find the total force exerted on the object in each direction:

$$\begin{aligned} F_x &= \iiint_E F_x(x, y, z) dV, \\ F_y &= \iiint_E F_y(x, y, z) dV, \\ F_z &= \iiint_E F_z(x, y, z) dV. \end{aligned}$$

When calculating work, the approach differs as work involves the displacement of the object under the force field. For a displacement along a path C , the work done by the force field is given by the line integral:

$$W = \int_C \mathbf{F} \cdot d\mathbf{r},$$

where $d\mathbf{r} = (dx, dy, dz)$ represents an infinitesimal displacement vector along the path. Expanding the dot product, we have:

$$\mathbf{F} \cdot d\mathbf{r} = F_x dx + F_y dy + F_z dz.$$

This expression calculates the infinitesimal work done by each component of the force along its respective displacement direction. In scenarios where the force is distributed over a volume and each point in the volume moves with a velocity field $\mathbf{v}(x, y, z)$, the *power density* (rate of doing work per unit volume) can be expressed as:

$$P(x, y, z) = \mathbf{F}(x, y, z) \cdot \mathbf{v}(x, y, z).$$

The total work done over a period of time can then be obtained by integrating the power density over the entire volume:

$$W = \iiint_E \mathbf{F}(x, y, z) \cdot \mathbf{v}(x, y, z) dV.$$

This framework is essential in physics and engineering applications. For example, when calculating the work done by an electric field $\mathbf{E}$ on a charge distribution $\rho(x, y, z)$ moving with velocity $\mathbf{v}(x, y, z)$, the integral takes the form:

$$W = \iiint_E \rho(x, y, z) (\mathbf{E} \cdot \mathbf{v}) dV,$$

where $\mathbf{v}$ is the local velocity vector at each point in the charge distribution. Here, the triple integral sums contributions to the total work throughout the entire region of charge. This process models how electric forces interact with volumetric charge distributions in systems such as capacitors or plasma clouds.

In Chapter 5, we will refine these concepts further by introducing line integrals and exploring how work is more directly computed along specific paths within a vector field. For now, understand that triple integrals are invaluable for modeling distributed forces and power densities across volumes.

▲ **Example 4.6.2** A magnetic field $\mathbf{B}(x, y, z) = (y^2, xz, xy)$ tesla acts on a conductive fluid in the unit cube $[0, 1]^3$. The fluid has charge density $\rho(x, y, z) = e^{-(x^2+y^2+z^2)}$ (coulombs/m³) and velocity $\mathbf{v}(x, y, z) = (z, x, y)$ (m/s).

Find:

1. The total magnetic force on the fluid given the differential Lorentz force $d\mathbf{F} = \rho(\mathbf{r})(\mathbf{v} \times \mathbf{B}) dV$ and force density $\mathbf{f}(\mathbf{r}) = \rho(\mathbf{r})[\mathbf{v}(\mathbf{r}) \times \mathbf{B}(\mathbf{r})]$
2. The work done by the magnetic field in one second, given that instantaneous power is $dP = \mathbf{F} \cdot \mathbf{v}$

Solution

1. Total Magnetic Force on the Fluid

Computing the cross product:

$$\mathbf{v} \times \mathbf{B} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ z & x & y \\ y^2 & xz & xy \end{vmatrix} = (x^2y - xyz, y^3 - xyz, xz^2 - xy^2)$$

Hence the total force components are:

$$F_x = \iiint_{E=[0,1]^3} e^{-(x^2+y^2+z^2)} (x^2y - xyz) dV$$

$$F_y = \iiint_{E=[0,1]^3} e^{-(x^2+y^2+z^2)} (y^3 - xyz) dV$$

$$F_z = \iiint_{E=[0,1]^3} e^{-(x^2+y^2+z^2)} (xz^2 - xy^2) dV$$

2. Work Done by the Magnetic Field in One Second

For a fluid element, the differential power is:

$$dP = [\rho(\mathbf{v} \times \mathbf{B})] \cdot \mathbf{v} dV = \rho(\mathbf{v} \times \mathbf{B}) \cdot \mathbf{v} dV$$

Because $\mathbf{v} \times \mathbf{B}$ is always perpendicular to $\mathbf{v}$, their dot product is zero:

$$(\mathbf{v} \times \mathbf{B}) \cdot \mathbf{v} = 0$$

Thus the magnetic field does no work on the fluid:

$$W = \iiint_E \rho[(\mathbf{v} \times \mathbf{B}) \cdot \mathbf{v}] dV = 0$$

▲ **Example 4.6.3** A force field $\mathbf{F}(x, y, z) = (yz, xz, xy)$ newtons acts on a volume of gas expanding in the unit cube $[0, 1]^3$. The gas particles move with velocity

$$\mathbf{v}(x, y, z) = (x, y, z) \quad (\text{m/s}).$$

Find:

1. The total force on the gas volume
2. The work done by the force field on the gas, given that instantaneous power is $dP = \mathbf{F} \cdot \mathbf{v} dV$

Solution

1. Total Force Components

The total force is found by integrating each component over the volume:

$$\begin{aligned} F_x &= \iiint_{[0,1]^3} yz dV = \int_0^1 \int_0^1 \int_0^1 yz dx dy dz \\ &= \int_0^1 \int_0^1 yz dy dz = \int_0^1 \frac{z}{2} dz = \frac{1}{4} \\ F_y &= \iiint_{[0,1]^3} xz dV = \int_0^1 \int_0^1 \int_0^1 xz dx dy dz \end{aligned}$$

$$\begin{aligned}
&= \int_0^1 \int_0^1 \frac{z}{2} dy dz = \frac{1}{4} \\
F_z &= \iiint_{[0,1]^3} xy dV = \int_0^1 \int_0^1 \int_0^1 xy dx dy dz \\
&= \int_0^1 \int_0^1 \frac{y}{2} dy dz = \frac{1}{4}
\end{aligned}$$

Therefore $\mathbf{F}_{total} = (\frac{1}{4}, \frac{1}{4}, \frac{1}{4})$ newtons.

2. Work Done by the Force Field

The instantaneous power at each point is:

$$\begin{aligned}
\mathbf{F} \cdot \mathbf{v} &= (yz, xz, xy) \cdot (x, y, z) \\
&= xyz + xyz + xyz \\
&= 3xyz
\end{aligned}$$

Therefore the total work done is:

$$\begin{aligned}
W &= \iiint_{[0,1]^3} 3xyz dV \\
&= 3 \int_0^1 \int_0^1 \int_0^1 xyz dx dy dz \\
&= 3 \int_0^1 \int_0^1 \frac{y}{2} z dy dz \\
&= 3 \int_0^1 \frac{z}{4} dz = \frac{3}{8} \text{ joules}
\end{aligned}$$

Average Values and Distributions

Just as we found average values using double integrals, we can find the average value of a function $f(x, y, z)$ over a three-dimensional region E :

$$f_{avg} = \frac{\iiint_E f(x, y, z) dV}{\iiint_E dV}$$

This concept proves invaluable in various physical contexts:

- Temperature distribution in a solid body, where $f(x, y, z)$ represents temperature

- Concentration of a solute in a solution, where $f(x, y, z)$ gives concentration
- Stress distribution in a structural element, where $f(x, y, z)$ describes stress intensity

The denominator $\iiint_E dV$ represents the volume of region E , making this formula a natural extension of the average value theorem to three dimensions. When working with non-uniform materials, we often need weighted averages that account for varying density:

$$f_{avg} = \frac{\iiint_E f(x, y, z) \rho(x, y, z) dV}{\iiint_E \rho(x, y, z) dV}$$

This weighted form finds particular use in materials science and engineering, where we might need to account for how material properties vary throughout a component. For instance, in a composite material, both density and thermal conductivity might vary spatially, affecting the overall thermal behavior of the object.

▲ **Example 4.6.4** A spherical blast furnace of radius 3 meters operates at extremely high temperatures. The temperature (in Celsius) at any point (x, y, z) in the furnace varies according to:

$$T(x, y, z) = 1500e^{-(x^2+y^2+z^2)/9} + 500$$

The density of the gas inside the furnace varies with height according to:

$$\rho(x, y, z) = e^{-z/3} \quad (\text{kg/m}^3)$$

Find:

1. The average temperature in the furnace
2. The density-weighted average temperature, accounting for the fact that hotter regions tend to have different densities

Solution

1. First, note the volume of our region E , a sphere of radius 3:

$$\iiint_E dV = \frac{4}{3}\pi(3^3) = 36\pi$$

The average temperature is:

$$T_{avg} = \frac{\iiint_E T(x, y, z) dV}{\iiint_E dV} = \frac{\iiint_E T(x, y, z) dV}{36\pi}$$

Converting to spherical coordinates (r, ϕ, θ) with

$$x = r \sin \phi \cos \theta, \quad y = r \sin \phi \sin \theta, \quad z = r \cos \phi, \quad dV = r^2 \sin \phi \, dr \, d\phi \, d\theta,$$

we have $x^2 + y^2 + z^2 = r^2$, so the temperature becomes

$$T(r) = 1500 e^{-\frac{r^2}{9}} + 500.$$

Therefore,

$$\iiint_E T(x, y, z) \, dV = \int_0^{2\pi} \int_0^\pi \int_0^3 \left(1500 e^{-\frac{r^2}{9}} + 500 \right) r^2 \sin \phi \, dr \, d\phi \, d\theta.$$

The angular parts give a factor of 4π , so

$$\iiint_E T(x, y, z) \, dV = 4\pi \left[1500 \int_0^3 r^2 e^{-\frac{r^2}{9}} \, dr + 500 \int_0^3 r^2 \, dr \right].$$

Evaluating,

$$\int_0^3 r^2 \, dr = 9 \quad \text{and} \quad \int_0^3 r^2 e^{-\frac{r^2}{9}} \, dr \approx 10.69.$$

Hence,

$$1500 \cdot 10.69 + 500 \cdot 9 = 1500 \cdot 10.69 + 4500 = 16035 + 4500 = 20535.$$

Multiplying by 4π gives:

$$20535 \cdot 4\pi = 82140\pi.$$

So

$$T_{\text{avg}} = \frac{82140\pi}{36\pi} \approx 2282 \text{ }^\circ\text{C}.$$

2. The density-weighted average temperature is

$$T_{\text{avg}} = \frac{\iiint_E T(x, y, z) \rho(x, y, z) \, dV}{\iiint_E \rho(x, y, z) \, dV}.$$

In spherical coordinates, $z = r \cos \phi$, so

$$\rho(r, \phi) = e^{-\frac{r \cos \phi}{3}}.$$

The resulting integrals involve products of exponentials in r and $\cos \phi$, so we evaluate them numerically using programming or Wolfram Alpha. One finds

$$T_{\text{avg}} \approx 2220 \text{ }^\circ\text{C}.$$

The density-weighted average is lower because the hotter central regions (larger r^2 term in $e^{-r^2/9}$) coincide with regions where ρ is relatively smaller for higher z ; effectively, less “weight” is assigned to the hottest points.

Formulas and Definitions

Important Vocabulary

- **Double Integral:** A mathematical tool that measures accumulation over a two-dimensional region, used to find quantities like area, volume, mass, and center of mass.
- **Double Integral Type I Region:** A region R in the (x, y) -plane of the form $R = \{(x, y) : a \leq x \leq b, g_1(x) \leq y \leq g_2(x)\}$, where vertical lines intersect the region's boundary exactly twice.
- **Double Integral Type II Region:** A region R in the (x, y) -plane of the form $R = \{(x, y) : c \leq y \leq d, h_1(y) \leq x \leq h_2(y)\}$, where horizontal lines intersect the region's boundary exactly twice.
- **Fubini's Theorem:** States that a double or triple integral can be evaluated as repeated integrals in any order over a rectangular region.
- **Triple Integral:** An extension of double integrals to three dimensions, used to find volumes, mass, and other physical quantities in $\mathbb{R}^3$.
- **Triple Integral Type I Region:** A solid region E bounded above and below by surfaces $z = g_2(x, y)$ and $z = g_1(x, y)$ over a region R in the (x, y) -plane, where vertical lines intersect the boundary exactly twice.
- **Triple Integral Type II Region:** A solid region E bounded by surfaces $x = h_1(y, z)$ and $x = h_2(y, z)$ over a region R in the (y, z) -plane, where lines parallel to the x -axis intersect the boundary exactly twice.
- **Triple Integral Type III Region:** A solid region E bounded by surfaces $y = k_1(x, z)$ and $y = k_2(x, z)$ over a region R in the (x, z) -plane, where lines parallel to the y -axis intersect the boundary exactly twice.
- **Jacobian Determinant:** A measure of how a coordinate transformation affects area or volume elements, denoted $|J|$.
- **Moment of Inertia:** A measure of an object's resistance to rotational motion about an axis, denoted I_x , I_y , or I_z .
- **Product of Inertia:** A measure of coupling between rotations about different axes, denoted I_{xy} , I_{yz} , or I_{xz} .
- **Center of Mass:** The point $(\bar{x}, \bar{y}, \bar{z})$ where an object's mass can be considered concentrated for certain calculations.
- **Inertia Tensor:** A 3×3 matrix describing how mass is distributed in a rotating object.

Key Formulas

Double Integrals

$$\text{Volume} = \iint_R f(x, y) dA$$

$$\text{Mass} = \iint_R \rho(x, y) dA$$

$$\text{Average} = \frac{\iint_R f(x, y) dA}{\iint_R dA}$$

Center of Mass (2D)

$$\bar{x} = \frac{1}{M} \iint_R x \rho(x, y) dA$$

$$\bar{y} = \frac{1}{M} \iint_R y \rho(x, y) dA$$

Polar Coordinates

$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$dA = r dr d\theta$$

Triple Integrals

$$\text{Mass} = \iiint_E \rho(x, y, z) dV$$

$$\text{Average} = \frac{\iiint_E f(x, y, z) dV}{\iiint_E dV}$$

Center of Mass (3D)

$$\bar{x} = \frac{1}{M} \iiint_E x \rho(x, y, z) dV$$

$$\bar{y} = \frac{1}{M} \iiint_E y \rho(x, y, z) dV$$

$$\bar{z} = \frac{1}{M} \iiint_E z \rho(x, y, z) dV$$

Cylindrical Coordinates

$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$z = z$$

$$dV = r dr d\theta dz$$

Spherical Coordinates

$$x = \rho \sin \phi \cos \theta$$

$$y = \rho \sin \phi \sin \theta$$

$$z = \rho \cos \phi$$

$$dV = \rho^2 \sin \phi d\rho d\theta d\phi$$

Moments of Inertia

$$I_x = \iiint_E (y^2 + z^2) \rho dV$$

$$I_y = \iiint_E (x^2 + z^2) \rho dV$$

$$I_z = \iiint_E (x^2 + y^2) \rho dV$$

Products of Inertia

$$I_{xy} = \iiint_E xy \rho dV$$

$$I_{yz} = \iiint_E yz \rho dV$$

$$I_{xz} = \iiint_E xz \rho dV$$

Jacobian Determinant (2D)

$$|J| = \det \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{bmatrix}$$

Jacobian Determinant (3D)

$$|J| = \det \begin{bmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} & \frac{\partial x}{\partial w} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} & \frac{\partial y}{\partial w} \\ \frac{\partial z}{\partial u} & \frac{\partial z}{\partial v} & \frac{\partial z}{\partial w} \end{bmatrix}$$

Change of Variables Formula

$$\iint_R f(x, y) \, dx \, dy = \iint_{R'} f(u, v) |J| \, du \, dv$$

$$\iiint_E f \, dx \, dy \, dz = \iiint_{E'} f |J| \, du \, dv \, dw$$

Force and Work

$$F_x = \iiint_E F_x(x, y, z) \, dV$$

$$F_y = \iiint_E F_y(x, y, z) \, dV$$

$$F_z = \iiint_E F_z(x, y, z) \, dV$$

$$W = \iiint_E \mathbf{F} \cdot \mathbf{v} \, dV$$

Practice Problems

1. Evaluate $\iint_R (2x + 3y) dA$ where $R = [0, 2] \times [1, 3]$.

2. Find $\iint_R xy^2 dA$ where $R = [-1, 1] \times [0, 2]$.

3. Calculate $\iint_R \sin(x + y) dA$ where $R = [0, \pi] \times [0, \pi]$.

4. Consider a rectangular sheet $[0, a] \times [0, b]$ with temperature distribution $T(x, y) = \sin(\pi x/a) \sin(\pi y/b)$. Find its average temperature.

5. The force on an electrical conductor depends on position according to $F(x, y) = Be^{-x^2-y^2}$ newtons per square meter, where B is magnetic field strength. Find the total force on a rectangular conductor $[-1, 1] \times [-1, 1]$ when $B = 2$.

6. For $\iint_R f(x, y) dA = 4$ where $R = [0, 1] \times [0, 1]$, and $\iint_R xf(x, y) dA = 3$, find $\iint_R x^2 f(x, y) dA$.

7. Evaluate $\iint_R \frac{1}{(1+x^2)(1+y^2)} dA$ where $R = [0, 1] \times [0, 1]$.

8. Use a Riemann sum to approximate

$$\iint_{[0,2] \times [1,3]} 2xy dA$$

by dividing both dimensions into 2 equal parts (a 2×2 grid). Take the *mid-point* of each subrectangle as the sample point. Show the subdivision, identify

each sample point, and compute the resulting sum. Do *not* convert this into an iterated integral directly; follow the Riemann sum framework:

$$S_{2,2} = \sum_{i=1}^2 \sum_{j=1}^2 f(x_{ij}^*, y_{ij}^*) \Delta A.$$

9. Consider the function $f(x, y) = x - y$ over the rectangular region $[0, 1] \times [0, 2]$.

(a) Set up $\iint_R (x - y) dA$ explicitly as an iterated integral.

(b) Evaluate the integral and interpret the result in terms of *negative* and *positive* volume, given that $x - y$ dips below the xy -plane in some parts of R .

10. Evaluate using Fubini's Theorem:

$$\iint_R (3 + 2x) dA,$$

where $R = [1, 2] \times [0, 1]$. Show both orders of integration explicitly and confirm they produce the same numerical result.

11. Let $f(x, y) = e^x \sin(y)$. Compute

$$\iint_{[0,\pi] \times [-1,1]} e^x \sin(y) dA$$

by factoring the integral into a product of two single-variable integrals. Show each step clearly.

12. Determine which order of integration is simpler, and then evaluate:

$$\iint_{[0,1] \times [2,3]} \frac{x}{1+x^2+y^2} dA.$$

Discuss briefly why integrating with respect to y first (or vice versa) might be advantageous. Verify the final result.

13. Consider

$$\iint_{[1,2] \times [0,2]} \frac{1}{(x+1)(y+2)} dA.$$

- (a) Rewrite the integrand using partial fractions if helpful.
- (b) Evaluate the resulting iterated integral using Fubini's Theorem.
14. Briefly explain why the choice of sample points (lower-left, upper-right, midpoint, or max/min) does not affect the *limit* of the Riemann sum for a *continuous* function over a rectangular region. In your explanation, address how increasing the partition's fineness guarantees convergence to a single value. (No detailed calculations required; this is conceptual.)
15. Show that

$$\iint_{[-1,1] \times [-2,2]} \frac{xy}{x^2 + y^2} dA = 0$$

by leveraging symmetry arguments rather than brute-force integration. Hint: For fixed x , consider the integrand's behavior as y varies symmetrically around 0.

16. Recall that the average value of
- $f(x, y)$
- over
- R
- is

$$f_{\text{avg}} = \frac{1}{\text{Area}(R)} \iint_R f(x, y) dA.$$

Compute the average value of $f(x, y) = \sin(xy)$ over $R = [0, 2] \times [0, 2]$. Then simplify your final expression if possible.

17. Find the general form of the indefinite double integral:

$$\iint x^2 y \, dx \, dy.$$

Explicitly show how you introduce an arbitrary function of one variable at each step.

18. Compute the indefinite double integral:

$$\iint (3x + 2xy^2) \, dx \, dy,$$

carefully indicating the arbitrary functions that arise. Provide your final answer as

$$F(x, y) + g(x) + h(y).$$

19. Show that for any continuous function
- $f(x, y)$
- on
- $[a, b] \times [c, d]$
- :

$$\iint_R f(x, y) dA = \frac{(b-a)(d-c)}{4} [f(a, c) + f(a, d) + f(b, c) + f(b, d)]$$

if and only if $f(x, y)$ is bilinear (linear in each variable and of the form $f(x, y) = ax + by + cxy + d$).

20. Prove that if
- $f(x, y)$
- and
- $g(x, y)$
- are continuous on
- $[a, b] \times [c, d]$
- , then:

$$\left(\iint_R f(x, y) dA \right) \left(\iint_R g(x, y) dA \right) \leq (b-a)(d-c) \iint_R f(x, y)g(x, y) dA$$

21. For a quantum particle in a box
- $[0, a] \times [0, b]$
- , the probability density is:

$$|\psi(x, y)|^2 = \frac{4}{ab} \sin^2\left(\frac{\pi x}{a}\right) \sin^2\left(\frac{\pi y}{b}\right)$$

Show that the total probability equals 1.

22. Sketch the region bounded below by
- $y = 0$
- and above by
- $y = 3x$
- for
- $0 \leq x \leq 2$
- . Write down (but do not evaluate) the double integral

$$\iint_R (x + y) dA$$

as a Type I region, stating all limits clearly.

23. Consider the region R bounded on the left by $x = 1$ and on the right by $x = y + 3$, with $-2 \leq y \leq 2$. Express

$$\iint_R 5 \, dA$$

as a Type II double integral, but do not integrate. Show the bounds for y and x explicitly.

24. A region R is described by $0 \leq x \leq 4$ and $x^2 \leq y \leq 8 + x$.
- Explain why R is Type I in x .
 - Attempt to describe the same R as Type II: find the interval for y and the functions $h_1(y)$, $h_2(y)$.

No need to integrate; just show the correct bounds for each description.

25. Find the area of the region bounded by $y = x^2$ and $y = 6 - x$ for $-1 \leq x \leq 3$. Write and evaluate

$$\iint_R 1 \, dA.$$

If necessary, split into subregions to handle where the curves intersect.

26. Let R be the region with $0 \leq x \leq 2$ and $x \leq y \leq 4$. Evaluate

$$\iint_R (1 + xy) \, dA$$

using:

- Type I bounds in the order $dy \, dx$,
 - Type II bounds in the order $dx \, dy$.
- Confirm both give the same result.

27. A region R in the plane is bounded by $y = -2$, $y = 2$, and $x = (y + 1)^2$. Compute

$$\iint_R (4 - x - y) \, dA$$

using a Type II setup. First, identify $h_1(y)$ and $h_2(y)$, then integrate.

28. Use polar coordinates to evaluate the double integral

$$\iint_R (r^2) \, dA$$

where R is the disk of radius 3 centered at the origin. Clearly indicate the r and θ bounds.

29. Set up (but do not evaluate) the double integral for

$$\iint_R (x^2 + y^2) \, dA$$

over the sector where $0 \leq r \leq 2$, $\frac{\pi}{4} \leq \theta \leq \frac{3\pi}{4}$. Show each step of your conversion to polar form.

30. Compute

$$\iint_R (3 + x + y) \, dA$$

where R is the region between the circles $r = 2$ and $r = 3$ (centered at the origin). Express x and y in terms of r, θ , then evaluate fully.

31. A region R consists of the portion of the parabola $y = x^2$ from $x = 0$ to $x = 1$, together with the quarter-circle of radius 1 to complete a closed region.

- Explain why it may be simpler to split R into two parts: R_1 for the parabola region (Cartesian) and R_2 for the quarter circle (Polar).
- Write the integral for

$$\iint_R (1 + x^2 + y^2) \, dA$$

as $\iint_{R_1} \cdots + \iint_{R_2} \cdots$, specifying each region's setup.

32. A region R is bounded by $r = 1 + \cos \theta$ and $r = 2 + \sin \theta$ for $0 \leq \theta \leq \frac{\pi}{2}$. Write (but do not evaluate) the integral

$$\iint_R \sqrt{x^2 + y^2} dA$$

in polar coordinates. Clearly label the inner and outer radius functions and the range of θ .

33. The region in the first quadrant bounded by $y = x^3$, $y = 4$, and $x = 3$ is not entirely Type I due to the x^3 shape for small x . Show how to split this region into two simpler Type I parts, and set up

$$\iint_R (2x + y) dA$$

so that each subregion can be integrated in the form $\int_a^b \int_{g_1(x)}^{g_2(x)} \dots dy dx$.

34. A thin plate occupies the region $0 \leq y \leq x^2$ for $0 \leq x \leq 3$. Its density is $\rho(x, y) = 5 + x$.

- Write down the double integral for the total mass.
- Write down the integrals for the moments M_x and M_y , and hence the coordinates of the center of mass $(\bar{x}, \bar{y})$.
- You may leave the integrals in un-simplified form, or carry them through to a final numeric result (your choice).

35. A region R is bounded by the cardioid $r = 2 + 2\cos(\theta)$ in polar coordinates.

- Write an integral in polar form to find the area of R .
- Convert that same region R to a Type I or II region in Cartesian coordinates (this will look complicated). Briefly argue why polar

is more convenient for actual integration.

36. The region R is bounded by $x = 0$, $y = 0$, $x^2 + y^2 = 4$, and $y = 4 - x$. This region is part circular arc, part straight line, part coordinate axes.

- Sketch R carefully. Identify obvious subregions if needed (e.g. one part might be simpler in polar, another part simpler in Cartesian).
- Propose a setup for

$$\iint_R (x + y) dA$$

employing your chosen splits and coordinate systems. No need to fully evaluate, but write the integrals clearly.

37. A thin plate occupies the rectangular region $R = [0, 3] \times [0, 2]$ in the xy -plane. Its (uniform) density is $\rho_0 = 5 \text{ kg/m}^2$.

- Write a double integral to find the plate's total mass.
- Evaluate the integral explicitly to obtain M .

38. Consider a right triangle in the first quadrant with vertices at $(0, 0)$, $(4, 0)$, and $(4, 3)$. The plate has constant density $\rho_0 = 2 \text{ kg/m}^2$.

- Find the total mass of the plate.
- Compute its center of mass $(\bar{x}, \bar{y})$.
- Briefly interpret why $\bar{x}$ is more than halfway along the x -base if ρ_0 is uniform.

39. A rectangular plate extends over $0 \leq x \leq 2$, $0 \leq y \leq 3$, and has density $\rho(x, y) = 5 + x$ (in g/cm^2).

- Set up the double integral for the total mass.

- (b) Evaluate it and give a numeric answer in grams.
40. A metal plate occupies the region R bounded by $x = 0$, $y = 0$, and $y = 4 - x^2$ in the first quadrant. The temperature at each point is $T(x, y) = 10 + xy$ (in $^{\circ}\text{C}$).
- (a) Write a double integral for the average temperature T_{avg} over R .
- (b) Evaluate the integral to get a numerical value for T_{avg} .

41. A semi-circular sheet of radius 3 (centered at the origin) lies above the x -axis, so $y = \sqrt{9 - x^2}$, $-3 \leq x \leq 3$, $0 \leq y \leq \sqrt{9 - x^2}$. The density is constant, $\rho_0 = 1$. Show that

$$I_x = \iint_R y^2 \rho_0 dA$$

can be computed more easily in polar coordinates. Write and evaluate the polar integral, where $r \in [0, 3]$, $\theta \in [0, \pi]$.

42. A uniform circular plate of radius R has density ρ_0 . Using polar coordinates, show that

$$I_z = \iint_R (x^2 + y^2) \rho_0 dA = \rho_0 \pi \frac{R^4}{2}.$$

No advanced derivation needed; just set up and solve the integral quickly.

43. A disc of radius 2 centered at the origin has a density that increases linearly with r , given by $\rho(r) = 4 + r$ (in some consistent units).
- (a) Write the polar-coordinate integral for the total mass M .
- (b) Compute M .
- (c) Find the average density across the disc, $\rho_{\text{avg}} = \frac{1}{\text{Area}} \iint_R \rho(r) dA$.

44. Let $z = x^2 + 3y^2$ be a paraboloid defined over the rectangle $0 \leq x \leq 2$, $0 \leq y \leq 1$.

- (a) Write the formula for its surface area:

$$A = \iint_R \left(1 + \left(\frac{\partial z}{\partial x} \right)^2 + \left(\frac{\partial z}{\partial y} \right)^2 \right)^{1/2} dA.$$

- (b) Substitute the appropriate partial derivatives and set up (but do not necessarily evaluate) the integral explicitly.

45. A solid region in the first octant is bounded below by the plane $z = 0$, on one side by $x = 0$, on another side by $y = 0$, and on top by the plane $x + y + z = 6$. Write a triple integral for

$$\iiint_E (2xy - z) dV$$

using an order of integration that you find convenient.

46. Find

$$\iiint_E 1 dV$$

for the region E bounded by the parabolic cylinder $z = x^2$, the planes $x = 0$ and $y = 2$, and the plane $z = 4$. Split or rearrange as needed to ensure correct bounds, then evaluate to obtain a numerical volume.

47. A tetrahedron in the first octant has one vertex at the origin and its three edges on the coordinate axes meet the plane $2x + 3y + z = 6$. Compute

$$\iiint_E (x^2 + yz) dV$$

.

48. The solid E is in cylindrical coordinates:

$$0 \leq r \leq 3, \quad 0 \leq \theta \leq \pi, \quad -1 \leq z \leq 2.$$

Set up (but do not evaluate) the integral

$$\iiint_E (3r^2 + z) r \, dr \, d\theta \, dz.$$

Clearly specify the bounds for r , θ , z .

49. Find

$$\iiint_E (x^2 + y^2 + z^2) \, dV$$

where E is the solid ball of radius 4 centered at the origin, using spherical coordinates. Include the factor $\rho^2 \sin \phi$ and show each step.

50. A region is defined by the intersection of the spheres $x^2 + y^2 + z^2 = 4$ and $x^2 + y^2 + (z-2)^2 = 4$. Write a spherical-coordinate integral to compute its volume. Give precise bounds for ρ, θ, ϕ , but do not do the final integration.

51. Evaluate

$$\iiint_E \rho(x, y, z) \, dV$$

where $\rho(x, y, z) = 3x$ is a density function, and E is the region in the first octant bounded by $x^2 + y^2 + z^2 = 9$. Specify whether you choose cylindrical or spherical coordinates, and solve the integral step by step.

52. A region R in the xy -plane is given by the circle $x^2 + y^2 \leq 4$. Set up the integral

$$\iint_R (4 - x^2 - y^2) \, dx \, dy$$

using the transformation to polar coordinates, identifying the Jacobian factor. Do not evaluate.

53. Consider the integral

$$\iint_R \exp((x+y)^2) \, dx \, dy$$

where R is the parallelogram with vertices $(0, 0)$, $(2, 0)$, $(3, 2)$, and $(1, 2)$. Propose a linear change of variables to convert R into a rectangle. Write down the new integral in the transformed variables, along with its Jacobian determinant.

54. A region in the uv -plane is the rectangle $0 \leq u \leq 3$, $0 \leq v \leq 2$. Under the transformation

$$x = 2u + 3v, \quad y = u - 2v,$$

the image region in (x, y) -space is a parallelogram.

- (a) Find the Jacobian $|J|$ of this mapping.
(b) Write

$$\iint_R (x + 2y) \, dx \, dy$$

in terms of u and v , indicating proper bounds but not solving.

55. A transformation from (r, θ) to (x, y) is given by

$$x = r^2 \cos \theta, \quad y = r^2 \sin \theta.$$

Find the Jacobian determinant $\left| \frac{\partial(x, y)}{\partial(r, \theta)} \right|$ and rewrite

$$\iint_R \sqrt{x^2 + y^2} \, dx \, dy$$

over the region $0 \leq r \leq 1$, $0 \leq \theta \leq 2\pi$.

56. Use the transformation

$$u = \frac{x-y}{\sqrt{2}}, \quad v = \frac{x+y}{\sqrt{2}}$$

to rewrite the double integral

$$\iint_R 1 \, dx \, dy$$

where R is the square with vertices $(0,0)$, $(2,0)$, $(2,2)$, $(0,2)$. Make sure to express the new region and the Jacobian in your final integral.

57. In three dimensions, consider

$$\iiint_E (x + 2y - z) \, dx \, dy \, dz,$$

where E is the solid cylinder $x^2 + y^2 \leq 4$ with $0 \leq z \leq 5$. Use cylindrical coordinates $(x = r \cos \theta, y = r \sin \theta, z = z)$ to change variables. Include the Jacobian factor and the correct bounds for r, θ, z , but do not evaluate.

58. Consider the integral $\iint_R (x + y) \, dx \, dy$ where R is the parallelogram with vertices at $(0,0)$, $(2,0)$, $(1+a, 2+a)$, and $(3+a, 2+a)$ for $a > 0$. Use a linear transformation to convert this to an integral over a rectangle.

59. Transform the integral

$$\iiint_E (x^2 + y^2 + z^2) \, dx \, dy \, dz$$

where E is the region $\{(x, y, z) \mid x^2 + y^2 + z^2 \leq 9\}$, using spherical coordinates. Clearly write each step, showing how dV becomes $\rho^2 \sin \phi \, d\rho \, d\phi \, d\theta$, and specify limits.

60. For the integral

$$\iiint_E 4xyz \, dx \, dy \, dz$$

where E is bounded by the surfaces $u = x - y$, $v = y - 2z$, $w = x + z = 3$ (plus planes to close the region), propose a linear transformation

$$(u, v, w) \mapsto (x, y, z)$$

with a nonzero Jacobian. Write the integral in (u, v, w) -space (no need for the precise region if it's complicated; just demonstrate how to set up the change of variables and compute its Jacobian).

61. Consider the following 3D coordinate system (sometimes called *elliptic cylindrical coordinates*) defined by parameters (μ, ν, z) mapping to rectangular coordinates (x, y, z) via

$$\begin{cases} x = a \cosh(\mu) \cos(\nu), \\ y = a \sinh(\mu) \sin(\nu), \\ z = z, \end{cases}$$

where a is a positive constant, $\mu \in [0, \infty)$, $\nu \in [0, 2\pi)$, and $z \in (-\infty, \infty)$.

- (a) Show the Jacobian matrix $J = \frac{\partial(x, y, z)}{\partial(\mu, \nu, z)}$.
- (b) Compute the determinant $|\frac{\partial(x, y, z)}{\partial(\mu, \nu, z)}|$ and interpret how it affects a volume element dV .

62. Find the total mass of a right circular cone of height 3 and base radius 1, where the density at each point is proportional to its height: $\rho(x, y, z) = kz$. Express your answer in terms of k .
63. A solid hemisphere of radius 2 is centered at the origin with $z \geq 0$. Its density varies according to $\rho(x, y, z) = 1 + z^2$. Write a triple integral for the total mass (no need to evaluate) using spherical coordinates.

64. A rectangular box $0 \leq x \leq 1$, $0 \leq y \leq 2$, $0 \leq z \leq 3$ contains a material with density $\rho(x, y, z) = x + y + z$.
- Set up the integral for the total mass.
 - Determine the z -coordinate of its center of mass (no need for $\bar{x}$ or $\bar{y}$).
65. A solid occupies the ellipsoid $\frac{x^2}{9} + \frac{y^2}{4} + \frac{z^2}{1} \leq 1$. Its density is $\rho(x, y, z) = \sqrt{x^2 + y^2 + z^2}$. Propose a suitable change of variables to simplify the region to a sphere in (u, v, w) -coordinates. Write the transformed integral for the total mass, including the Jacobian factor.
66. A force field $\mathbf{F}(x, y, z) = (yz, xz, xy)$ acts on the solid cylinder $x^2 + y^2 \leq 4$ for $0 \leq z \leq 5$.
- Find the total force on the cylinder by integrating each component over the volume.
 - If each point in the cylinder moves with velocity $\mathbf{v}(x, y, z) = (x, y, z)$, find the work done in one second.
67. Compute the moment of inertia I_z about the z -axis for a uniform sphere of radius R and density ρ_0 . Use spherical coordinates. Show intermediate steps but leave your final answer in simplified symbolic form.
68. A solid tetrahedron is formed by the coordinate planes ($x = 0$, $y = 0$, $z = 0$) and the plane $x + 2y + 3z = 6$. Its density is $\rho(x, y, z) = 2x + 5y + z$.
- Find the total mass of the tetrahedron.
 - Determine its center of mass $(\bar{x}, \bar{y}, \bar{z})$.
69. A region in $\mathbb{R}^3$ is bounded by $r = 2$ in cylindrical coordinates and z ranging from -1 to 3 . Let $\rho(r, \theta, z) = 2 + z$ be the density.
- Write the integrals for I_x and I_y (the moments of inertia about x - and y -axes).
 - Briefly comment on why I_{xy} might be zero by symmetry (no explicit computation required).
70. A solid with density $\rho(x, y, z) = e^{-z}$ occupies the region $x^2 + y^2 + z^2 \leq 9$.
- Convert to spherical coordinates.
 - Write the triple integral for the total mass (no need to evaluate).
 - Describe how you might adapt the integral to find the density-weighted average value of z .

Chapter 5

Vector Calculus

Introduction

In our mathematical journey thus far, we have built a foundation with vectors, developed sophistication with partial derivatives, and mastered the art of multiple integration. Now, we stand at the threshold of vector calculus, where these powerful tools converge to unveil the deeper structures of multidimensional mathematics. Vector calculus represents the natural evolution of our studies, allowing us to analyze not just quantities that vary over space, but also the flow, circulation, and flux of vector fields that pervade the physical world. This but another step into the idea of generalization we have been encouraging in our studies so thus far.

The concepts we explored in earlier chapters—vector operations, partial derivatives, and multiple integrals—now serve as building blocks for understanding more complex phenomena. Our facility with multiple integrals transforms into the ability to evaluate line and surface integrals, enabling us to calculate work done along paths and flux through surfaces. The geometric intuition we developed for vectors becomes essential as we visualize and analyze vector fields, from simple gradient fields to the intricate patterns of electromagnetic forces.

This chapter introduces fundamental theorems that bridge different dimensions of integration—Green’s Theorem, Stokes’ Theorem, and the Divergence Theorem. These powerful results establish profound connections between different types of integrals, helping us understand how local properties of vector fields relate to their global behavior. As we progress, we will see how the seemingly abstract concepts from previous chapters find concrete expression in the language of vector calculus, providing a gateway to more advanced mathematics and its applications in science and engineering. The journey ahead promises to be both challenging and illuminating, as we develop the ability to analyze and describe the fundamental patterns that shape our physical world.

Lecture 5.1: Vector Fields

Our journey through multivariable calculus has led us through scalar functions, where we mapped points in space to single numerical values. For instance, we could assign a temperature to each point in a room, or a pressure value to each point in the atmosphere. But many physical quantities we encounter can't be fully described by just a single number—they need both a magnitude and a direction.

Think about the wind blowing across a vast plain. At each point in space, the wind has both a speed (magnitude) and a direction. Or imagine iron filings scattered near a magnet—at each point, there's a magnetic force acting with a specific strength and orientation. These are examples where we need to assign not just a number, but a vector to each point in space.




Figure 5.1: A magnet pulling iron filings is a physical representation of a vector field. each point around the magnet has a magnitude of force and direction emanating from the magnet

Let's build this idea carefully. Start by imagining a single point (x_0, y_0) in the plane. At this point, we want to attach a vector—perhaps representing wind velocity. This vector will have both an x -component and a y -component, which we might call $P(x_0, y_0)$ and $Q(x_0, y_0)$ respectively. The vector at this point would then be:

$$\mathbf{v}_{(x_0, y_0)} = P(x_0, y_0)\mathbf{i} + Q(x_0, y_0)\mathbf{j}$$

Now imagine doing this for every point in the plane. As we move to different points (x, y) , both components of our vector might change—the wind might blow in different directions and with different speeds at different locations. We're no longer just looking at a single vector, but at a function that assigns a vector to every point. This is the essence of a vector field.

Definition and Basic Properties

Formally, a **vector field** $\mathbf{F}$ in $\mathbb{R}^n$ is a function that assigns to each point $\mathbf{p}$ in its domain a vector $\mathbf{F}(\mathbf{p})$. Just as scalar fields mapped points to numbers, vector fields map points to

vectors. In two dimensions, this means associating each point (x, y) with a two-dimensional vector:

$$\mathbf{F}(x, y) = P(x, y)\mathbf{i} + Q(x, y)\mathbf{j}$$

Here, $P(x, y)$ gives the x -component of the vector at any point (x, y) , while $Q(x, y)$ gives the y -component. To understand how this works in practice, let's break down the process of evaluating a vector field at specific points.

Consider a simple vector field $\mathbf{F}(x, y) = x\mathbf{i} + y\mathbf{j}$. Here, $P(x, y) = x$ and $Q(x, y) = y$. To plot vectors in this field:

x	y	Vector Tail at	$P(x, y)$	$Q(x, y)$
1	1	(1,1)	1	1
2	1	(2,1)	2	1
1	2	(1,2)	1	2
-1	1	(-1,1)	-1	1

At each point (x, y) , we:

1. Place the tail of our vector at the point (x, y)
2. Move $P(x, y)$ units in the $\mathbf{i}$ direction
3. Move $Q(x, y)$ units in the $\mathbf{j}$ direction

For example, at the point $(2, 1)$, we place our vector tail at $(2, 1)$, then move 2 units in the $\mathbf{i}$ direction and 1 unit in the $\mathbf{j}$ direction, resulting in a vector pointing to $(4, 2)$.

The idea extends naturally to three dimensions. At each point (x, y, z) in space, we now need three components to specify a vector:

$$\mathbf{F}(x, y, z) = P(x, y, z)\mathbf{i} + Q(x, y, z)\mathbf{j} + R(x, y, z)\mathbf{k}$$

For instance, in the three-dimensional vector field $\mathbf{F}(x, y, z) = z\mathbf{i} + x\mathbf{j} + y\mathbf{k}$:

x	y	z	$P(x, y, z)$	$Q(x, y, z)$	$R(x, y, z)$
1	2	3	3	1	2
2	1	0	0	2	1
-1	0	2	2	-1	0

At each point (x, y, z) , we:

1. Place the tail of our vector at the point (x, y, z)
2. Move $P(x, y, z)$ units in the $\mathbf{i}$ direction
3. Move $Q(x, y, z)$ units in the $\mathbf{j}$ direction
4. Move $R(x, y, z)$ units in the $\mathbf{k}$ direction

For example, at the point $(1, 2, 3)$, the vector starts at $(1, 2, 3)$ and has components $\langle 3, 1, 2 \rangle$, meaning it extends 3 units in the x -direction, 1 unit in the y -direction, and 2 units in the z -direction.

The functions P , Q , and R are called the **component functions** of the vector field. Each represents the magnitude of the field's projection onto one of the coordinate axes. Just as coordinates tell us where we are in space, these component functions tell us what vector we'll find when we get there.

Just as we studied continuity and differentiability for scalar functions—concepts that told us how “smoothly” the function varied as we moved around—these ideas extend naturally to vector fields. A vector field is continuous if all its component functions are continuous, meaning the vectors change smoothly as we move from point to point. Similarly, a vector field is differentiable if all its component functions are differentiable, allowing us to analyze rates of change in the field's behavior.

Visualizing Vector Fields

After understanding how to evaluate vector fields at individual points, our next step is visualizing them across regions of space. When we plot a vector field, we draw arrows at carefully chosen points to represent the field's behavior. Each arrow has two key properties:

- The direction of the arrow shows where the vector field points at that location
- The length (or magnitude) of the arrow indicates the vector's strength at that point

Let's develop this systematically. Consider the vector field $\mathbf{F}(x, y) = -y\mathbf{i} + x\mathbf{j}$. To begin visualizing this field, we'll evaluate it at several points using a grid:


x	y	Vector Tail at	$P(x, y) = -y$	$Q(x, y) = x$
1	1	(1,1)	-1	1
1	-1	(1,-1)	1	1
-1	-1	(-1,-1)	1	-1
-1	1	(-1,1)	-1	-1

For each point in our table, we plot a straight arrow by:

1. Starting at the tail point (x, y)
2. Drawing a straight line in the direction $\langle P(x, y), Q(x, y) \rangle$
3. Optionally normalize it by dividing by $\sqrt{P(x, y)^2 + Q(x, y)^2}$

It's crucial to note that regardless of the vector field, each arrow we draw is always straight—even if the overall pattern suggests rotation or curvature. Each arrow represents the instantaneous direction of the field at a single point.

For example, here's how we would visualize this field:



Notice how even with just these four points, we can begin to see the rotational pattern emerging in the field. Together the arrows suggest a counterclockwise circulation around the origin. Here's an example of a complete field generated systematically for $\mathbf{F}(x, y) = -y\mathbf{i} + x\mathbf{j}$:






Figure 5.2: *The circular pattern emerges from the collective behavior of many straight arrows, each showing the instantaneous direction at its starting point. This is analogous to how the tangent line approximates a curve at a single point—each vector arrow shows the field's direction at just one point in space.*

When we plot many more points, patterns emerge in the vector field. These patterns help us identify important features:


- **Sources:** Points where vectors emanate outward, like water springing from a fountain. For example, in the field $\mathbf{F}(x, y) = x\mathbf{i} + y\mathbf{j}$, the origin is a source:




- **Sinks:** Points where vectors converge inward, like water draining from a bathtub. The field $\mathbf{F}(x, y) = -x\mathbf{i} - y\mathbf{j}$ has a sink at the origin:



- **Circulation:** Regions where vectors suggest rotational motion, like our example field $\mathbf{F}(x, y) = -y\mathbf{i} + x\mathbf{j}$. Remember though—each individual arrow is still straight!



- **Saddle Points:** Points exhibiting both attractive and repulsive behavior, like the origin in $\mathbf{F}(x, y) = x\mathbf{i} - y\mathbf{j}$:



Notice how the colors in these plots indicate the magnitude of the vectors at each point. Brighter colors represent stronger vectors, while darker colors show regions where the field is weaker. The origin $(0,0)$ plays a special role in each of these fields, serving as the central point around which the field's behavior is organized.

Below is a Python script that can be used to visualize vector fields since it requires a lot of computation. Use it to aid you in your sketches for the examples to come.

```

1 import numpy as np
2 import matplotlib.pyplot as plt
3
4 # Define the functions P(x, y) and Q(x, y)
5 def P(x, y):
6     return -y # Example: -y
7
8 def Q(x, y):
9     return x # Example: x
10
11 # Create the grid
12 x, y = np.meshgrid(np.linspace(-10, 10, 25), np.linspace(-10, 10, 25))
13
14 # Compute the vector components
15 u = P(x, y)
16 v = Q(x, y)
17
18 # Normalize vectors to minimize arrow crossings
19 magnitude = np.sqrt(u**2 + v**2)
20 u = u / magnitude
21 v = v / magnitude
22
23 # Plot the vector field
24 plt.quiver(x, y, u, v, angles='xy', scale_units='xy', scale=1, color='blue', width
25           =0.0025)
26 plt.xlim(-10, 10)
27 plt.ylim(-10, 10)
28 plt.xlabel('x')
29 plt.ylabel('y')
30 plt.title('2D Vector Field')
31 plt.grid()
32 plt.show()

```

Understanding Vector Field Normalization

When visualizing vector fields, we often face a choice between showing the true magnitudes of our vectors and normalizing them to unit length. This decision fundamentally impacts what information we can glean from our visualization. Just as we might choose different coordinate systems based on the symmetries of a problem, we must thoughtfully consider when to normalize our vector fields.

To normalize a vector field $\mathbf{F}(x, y) = P(x, y)\mathbf{i} + Q(x, y)\mathbf{j}$, we divide by its magnitude at each point:

$$\mathbf{F}_{\text{norm}}(x, y) = \frac{P(x, y)}{\sqrt{P(x, y)^2 + Q(x, y)^2}}\mathbf{i} + \frac{Q(x, y)}{\sqrt{P(x, y)^2 + Q(x, y)^2}}\mathbf{j}$$

This transformation preserves the direction of each vector while scaling them all to unit length. Consider the field $\mathbf{F}(x, y) = (x^2 - y^2)\mathbf{i} + (2xy)\mathbf{j}$. In its normalized form, the quadratic growth in magnitude disappears, revealing purely directional information about the field. Alternatively, without normalization the magnitudes would cause eventual overlapping which would make the visualization unwieldy.

The decision to normalize a vector field should be guided by our analytical goals. Normalization is particularly valuable when:

1. **Directional Analysis is Primary:** When we need to understand the flow patterns or trajectories in our field, normalization helps by eliminating the potentially distracting variations in magnitude. This is especially useful in studying fluid flow patterns, magnetic field lines, or gravitational field directions.
2. **Visual Clarity is Crucial:** In fields with large magnitude variations, like our quadratic example, smaller vectors can become nearly invisible compared to larger ones. Normalization ensures each vector contributes equally to our visual understanding.
3. **Pattern Recognition is Key:** Normalized fields often reveal symmetries, circulation patterns, and topological features that might be obscured by magnitude variations. For instance, the rotational nature of our quadratic field becomes much clearer after normalization.

However, normalization isn't always appropriate. We should maintain the original magnitudes when:

1. **Magnitude Information is Essential:** In many physical applications, vector magnitude carries crucial information. A velocity field in fluid dynamics, for instance, needs its magnitude preserved to show where flow is fastest or slowest.
2. **Physical Accuracy Matters:** For quantitative analysis or physical simulations, preserving true vector magnitudes is often crucial. For example, when studying force fields, the strength of the force is just as important as its direction.

Just as a topographic map might show both elevation contours and gradient directions, we can often benefit from examining both normalized and unnormalized versions of a vector field. The normalized field reveals pure directional patterns, while the unnormalized field preserves magnitude information. Together, they provide a complete picture of the field's behavior.

▲ **Example 5.1.1** Classify and visualize the following vector fields:

1. $\mathbf{F}(x, y) = \frac{x}{\sqrt{x^2 + y^2}}\mathbf{i} + \frac{y}{\sqrt{x^2 + y^2}}\mathbf{j}$

$$2. \mathbf{F}(x, y) = \frac{x^2 - y^2}{\sqrt{(x^2 - y^2)^2 + (2xy)^2}} \mathbf{i} + \frac{2xy}{\sqrt{(x^2 - y^2)^2 + (2xy)^2}} \mathbf{j}$$

Solution Let's analyze each field systematically:

$$1. \text{ For } \mathbf{F}(x, y) = \frac{x}{\sqrt{x^2 + y^2}} \mathbf{i} + \frac{y}{\sqrt{x^2 + y^2}} \mathbf{j}:$$


Let's first understand what this field does:

- The denominator $\sqrt{x^2 + y^2}$ represents the distance from the origin to point (x, y)
- At each point (x, y) , dividing by this distance normalizes our vector to unit length
- The field gives us a way to point outward from the origin in every direction

Let's evaluate at eight points around the origin:

(x, y)	$\sqrt{x^2 + y^2}$	$P(x, y)$	$Q(x, y)$	Direction
(1, 0)	1	1	0	Right
(1, 1)	$\sqrt{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	NE diagonal
(0, 1)	1	0	1	Up
(-1, 1)	$\sqrt{2}$	$-\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	NW diagonal
(-1, 0)	1	-1	0	Left
(-1, -1)	$\sqrt{2}$	$-\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	SW diagonal
(0, -1)	1	0	-1	Down
(1, -1)	$\sqrt{2}$	$\frac{1}{\sqrt{2}}$	$-\frac{1}{\sqrt{2}}$	SE diagonal


Here's a rough sketch showing these eight unit vectors:



Several important properties emerge from our analysis:

- Every vector has length 1 (they're normalized)
- Each vector points directly away from the origin
- The field is undefined at the origin (division by zero)
- The pattern has radial symmetry about the origin

The computer-generated vector field looks like this:



This is a **source field** with unit vectors. All vectors point outward from the origin with equal magnitude. It's similar to the electric field of a positive point charge or the field lines emerging from a magnetic monopole (if one existed).

2. For $\mathbf{F}(x, y) = \frac{x^2 - y^2}{\sqrt{(x^2 - y^2)^2 + (2xy)^2}} \mathbf{i} + \frac{2xy}{\sqrt{(x^2 - y^2)^2 + (2xy)^2}} \mathbf{j}$


This normalized version of our previous field will help us understand its pure directional behavior. At each point, we're taking the same direction but scaling it to have unit length. Let's understand how we got here:

- The original field was $\mathbf{F}(x, y) = (x^2 - y^2)\mathbf{i} + (2xy)\mathbf{j}$
- Its magnitude at any point is $\sqrt{(x^2 - y^2)^2 + (2xy)^2}$
- Dividing by this magnitude normalizes each vector to length 1

Let's evaluate at carefully chosen points. Since we're normalizing, what matters most is the ratio between components rather than their absolute values:

(x, y)	$\sqrt{(x^2 - y^2)^2 + (2xy)^2}$	$P(x, y)$	$Q(x, y)$	Direction
$(1, 0)$	1	1	0	Right
$(1, 1)$	2	0	1	Up
$(0, 1)$	1	-1	0	Left
$(-1, 1)$	2	0	1	Up
$(-1, 0)$	1	-1	0	Left
$(-1, -1)$	2	0	-1	Down
$(0, -1)$	1	1	0	Right
$(1, -1)$	2	0	-1	Down

Let's sketch these unit vectors, being careful to show points at different distances from the origin:




These additional points help reveal even more patterns in the field:

- The same directional pattern appears at all distances from the origin - notice how points like $(1, 1)$ and $(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}})$ show identical behavior despite being at different distances
- Along rays from the origin (like the points $(1, 0)$ and $(2, 0)$), vectors point in exactly the same direction

- The transition between vertical and horizontal flow is smooth and consistent at all distances
- Each point's vector direction depends only on the ratio of y to x , not on their absolute values

The computer-generated vector field looks like this:



This normalized version of the field reveals its pure directional behavior. Now we can see clearly how the field creates a twisted circulation pattern, with vectors smoothly transitioning between vertical flow along diagonals and horizontal flow near the axes. This type of normalization is often useful in physics when we care more about the direction of a force or flow than its magnitude, such as in studying the directional patterns of complex fluid flows.

Conservative Vector Fields

When studying vector fields, one of the most fundamental questions we can ask is whether the field represents a quantity that depends only on position, or one that depends on how we got to that position. This distinction leads us to the crucial concept of conservative vector fields.

A vector field $\mathbf{F}$ is called **conservative** if there exists a scalar function f , called a **potential**

function, such that:

$$\mathbf{F} = \nabla f$$

In two dimensions, this means:

$$P(x, y) = \frac{\partial f}{\partial x} \quad \text{and} \quad Q(x, y) = \frac{\partial f}{\partial y}$$

And in three dimensions:

$$P(x, y, z) = \frac{\partial f}{\partial x}, \quad Q(x, y, z) = \frac{\partial f}{\partial y}, \quad \text{and} \quad R(x, y, z) = \frac{\partial f}{\partial z}$$

The term “conservative” comes from the fact that these fields conserve energy—there’s no loss or gain of energy when moving along a closed path. This leads to several critical properties:

- The work done by a conservative field along any path between two points depends only on the endpoints, not on the path taken
- The curl of a conservative field is zero (a fact we’ll explore in detail later)
- Line integrals of conservative fields around closed paths are always zero

When a field is not conservative, we call it **non-conservative** or **rotational**. Understanding whether a field is conservative or non-conservative has profound implications in physics and engineering:

- **Conservative Fields:**

- Gravitational fields in a uniform gravitational field
- Electrostatic fields
- Pressure gradients in static fluids

- **Non-Conservative Fields:**

- Magnetic fields
- Friction forces
- Velocity fields in rotating fluids

Testing for Conservative Fields

Not all vector fields are conservative. The key test involves examining the field’s partial derivatives. In two dimensions, if $\mathbf{F}(x, y) = P\mathbf{i} + Q\mathbf{j}$ is defined on a simply connected domain, then $\mathbf{F}$ is conservative if and only if:

$$\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$$

This condition arises from the equality of mixed partial derivatives of the potential function:

$$\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial^2 f}{\partial x \partial y}$$

In three dimensions, a vector field $\mathbf{F}(x, y, z) = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$ is conservative if and only if:

$$\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}, \quad \frac{\partial P}{\partial z} = \frac{\partial R}{\partial x}, \quad \frac{\partial Q}{\partial z} = \frac{\partial R}{\partial y}$$

The significance of conservative fields extends far beyond these mathematical conditions. Consider these practical implications:

1. **Energy Conservation:** In conservative systems, energy is perfectly conserved. A ball rolling in a gravitational field (ignoring friction) will reach the same height regardless of the path it takes.
2. **Path Independence:** The work done by conservative forces depends only on start and end positions. This means we can choose the most convenient path for our calculations.
3. **Potential Functions:** Conservative fields allow us to define potential energy functions, greatly simplifying many physical calculations. Instead of computing complicated path integrals, we can just evaluate the potential function at endpoints.
4. **Equilibrium Analysis:** The existence of a potential function helps identify stable and unstable equilibrium points in physical systems.

In contrast, non-conservative fields exhibit fundamentally different behavior:

1. They can do different amounts of work along different paths
2. They can add or remove energy from a system
3. They often represent dissipative forces or fields that depend on velocity
4. They cannot be derived from a potential function

This distinction between conservative and non-conservative fields is crucial in physics and engineering. For instance, when designing energy systems, knowing whether forces are conservative helps predict whether energy will be conserved or dissipated. In fluid dynamics, identifying non-conservative velocity fields helps understand where energy losses occur. Even in economics, conservative fields can model ideal market behaviors, while non-conservative fields might better represent real markets with friction and inefficiencies.

▲ **Example 5.1.2** Let's see what kind of fields Example 5.1.1 had (with the second example being changed to non-normalized for ease of differentiation). Please classify them as either Conservative or Non-Conservative:

1. The normalized radial field $\mathbf{F}(x, y) = \frac{x}{\sqrt{x^2+y^2}}\mathbf{i} + \frac{y}{\sqrt{x^2+y^2}}\mathbf{j}$

2. The quadratic field $\mathbf{F}(x, y) = (x^2 - y^2)\mathbf{i} + (2xy)\mathbf{j}$

Determine whether each field is conservative by testing the condition $\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$.

Solution Let's analyze each field systematically:

1. For the normalized radial field, let:

$$P(x, y) = \frac{x}{\sqrt{x^2 + y^2}} \quad \text{and} \quad Q(x, y) = \frac{y}{\sqrt{x^2 + y^2}}$$

First, compute $\frac{\partial P}{\partial y}$ using the quotient rule:

$$\begin{aligned} \frac{\partial P}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{x}{\sqrt{x^2 + y^2}} \right) \\ &= x \cdot \frac{\partial}{\partial y} \left(\frac{1}{\sqrt{x^2 + y^2}} \right) \\ &= x \cdot \left(-\frac{1}{2}(x^2 + y^2)^{-3/2} \cdot 2y \right) \\ &= -\frac{xy}{(x^2 + y^2)^{3/2}} \end{aligned}$$

Next, compute $\frac{\partial Q}{\partial x}$:

$$\begin{aligned} \frac{\partial Q}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{y}{\sqrt{x^2 + y^2}} \right) \\ &= y \cdot \frac{\partial}{\partial x} \left(\frac{1}{\sqrt{x^2 + y^2}} \right) \\ &= y \cdot \left(-\frac{1}{2}(x^2 + y^2)^{-3/2} \cdot 2x \right) \\ &= -\frac{xy}{(x^2 + y^2)^{3/2}} \end{aligned}$$

Since $\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$, this field is conservative. Indeed, we can find its potential function: $f(x, y) = \sqrt{x^2 + y^2}$. This explains why this normalized field points radially outward—it represents the direction of steepest increase of the distance function from the origin.

2. For the quadratic field, let:

$$P(x, y) = x^2 - y^2 \quad \text{and} \quad Q(x, y) = 2xy$$

Computing the partial derivatives:

$$\begin{aligned}\frac{\partial P}{\partial y} &= \frac{\partial}{\partial y}(x^2 - y^2) = -2y \\ \frac{\partial Q}{\partial x} &= \frac{\partial}{\partial x}(2xy) = 2y\end{aligned}$$

Since $\frac{\partial P}{\partial y} = -2y \neq 2y = \frac{\partial Q}{\partial x}$, this field is non-conservative. This aligns with our earlier observation of its rotational behavior—the field exhibits a circulation pattern that cannot be derived from any potential function. If we were to compute the work done by this field along different paths between the same points, we would get different results.

Lecture 5.2: Line Integrals

After our exploration of vector fields, we now turn to one of the most fundamental operations we can perform on them: integration along curves. Just as we previously extended single-variable integration to multiple dimensions through multiple integrals, we'll now develop the tools to integrate both scalar functions and vector fields along paths in space. This concept will prove crucial in physics, engineering, and advanced mathematics.

Line Integrals of Scalar Functions

Before diving into vector fields, let's start with a simpler scenario: integrating a scalar function along a curve. Imagine walking along a curved path on a hilly terrain. At each point, there's an elevation—a scalar value associated with your position. If we wanted to calculate the “total elevation” experienced along our path, we'd need a line integral.

Recall from Chapter 2 Lecture 4 that we developed the concept of arc length for a parametric curve. Given a curve C parameterized by $\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j}$ for $a \leq t \leq b$, we found its length using:

$$L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \int_a^b \left\| \frac{d\mathbf{r}}{dt} \right\| dt$$

Now imagine that at each point along this curve, we're not just measuring distance but also evaluating some scalar function $f(x, y)$. We can think of this as “weighting” each infinitesimal piece of arc length by the value of f at that point. This leads naturally to the line integral:

$$\int_C f(x, y) ds = \int_a^b f(x(t), y(t)) \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

We note that this will yield us an area, because we are multiplying the arc length by a height function. Let's break down the components of this integral:

- $f(x(t), y(t))$ evaluates our scalar function at each point along C
- $\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$ gives us the infinitesimal arc length ds
- Their product $f(x(t), y(t)) ds$ represents the contribution of each tiny piece of the curve

This geometric interpretation helps us understand several key properties:

1. **Reparametrization Invariance:** The value of the line integral doesn't depend on how we parameterize C . Different parameters will adjust both $f(x(t), y(t))$ and $\frac{d\mathbf{r}}{dt}$, but their product remains constant.
2. **Direction Independence:** For scalar line integrals, the direction we traverse C doesn't matter. This is because ds is always positive, measuring pure distance rather than directed movement.




Figure 5.3: Geometric interpretation of a scalar line integral $\int_C f(x, y) ds$. The curve C (shown in red) lies on the surface $z = f(x, y)$, while its projection onto the xy -plane is shown as a dashed magenta curve. The shaded “curtain” between these curves represents the accumulation of the function values along the path, with each vertical height $f(x, y)$ being weighted by the infinitesimal arc length ds .

3. **Physical Meaning:** The integral accumulates the product of the scalar field and distance traveled, giving us meaningful physical quantities:

- Mass = $\int_C \rho(x, y) ds$ where ρ is linear density
- Cost = $\int_C c(x, y) ds$ where c is cost per unit length
- Work = $\int_C F(x, y) ds$ where F is scalar force magnitude

The computation process follows these steps:

1. Express the curve C parametrically: $\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j}$
2. Calculate $\frac{dx}{dt}$ and $\frac{dy}{dt}$
3. Compose f with the parametrization: $f(x(t), y(t))$
4. Form the integrand: $f(x(t), y(t))\sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}$
5. Integrate with respect to t from a to b

You may encounter an alternative notation for line integrals, particularly in complex analysis and physics: the contour integral. Written as $\oint_C f(x, y) ds$, this notation emphasizes that we’re integrating along a specific path or contour in the plane. The circle in the integral symbol traditionally suggests a closed path—one that returns to its starting point—though

the notation is now commonly used for open paths as well. While we'll primarily use $\int_C$ notation in our development of vector calculus, recognizing the contour notation is crucial as it appears frequently in applications, especially when dealing with conservative fields and path independence.

▲ **Example 5.2.1** Evaluate the line integral $\int_C f(x, y) ds$ where $f(x, y) = y - 2x$ along:

1. The curve C_1 given by $\mathbf{r}(t) = (2t)\mathbf{i} + (3t)\mathbf{j}$, $0 \leq t \leq 1$
2. The curve C_2 given by $\mathbf{r}(t) = (\cos t)\mathbf{i} + (2 \sin t)\mathbf{j}$, $0 \leq t \leq 2\pi$

Solution Let's evaluate each integral systematically:

1. For curve C_1 :

First, we need:

- $\frac{dx}{dt} = 2$, $\frac{dy}{dt} = 3$
- $f(x(t), y(t)) = 3t - 4t = -t$
- $ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \sqrt{4 + 9} dt = \sqrt{13} dt$

Therefore:

$$\begin{aligned} \int_C f(x, y) ds &= \int_0^1 (-t)(\sqrt{13}) dt \\ &= -\sqrt{13} \int_0^1 t dt \\ &= -\sqrt{13} \left[\frac{t^2}{2} \right]_0^1 \\ &= -\frac{\sqrt{13}}{2} \end{aligned}$$

2. For curve C_2 :

We need:

- $\frac{dx}{dt} = -\sin t$, $\frac{dy}{dt} = 2 \cos t$
- $f(x(t), y(t)) = 2 \sin t - 2 \cos t$
- $ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \sqrt{\sin^2 t + 4 \cos^2 t} dt$

Therefore:

$$\begin{aligned}\oint_C f(x, y) ds &= \int_0^{2\pi} (2 \sin t - 2 \cos t) \sqrt{\sin^2 t + 4 \cos^2 t} dt \\ &= 0\end{aligned}$$

We see in this example where the height function goes negative for part of the region, and thus the total area is 0 from equal positive and negative contributions.

Contours

To evaluate line integrals effectively, we need a systematic understanding of the curves—or contours—along which we integrate; we will not always be given the parameterization of the curve like in the last two examples. A contour in the plane is simply a continuous curve that can be parameterized, but the nature of these curves (whether they're open or closed, smooth or piecewise) significantly impacts our computational approach.

Definition: A *contour* C is a curve in the plane that can be described by a continuous vector-valued function $\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j}$ over some interval $[a, b]$. We classify contours into several types:

- **Open Contours:** Curves where the endpoints are distinct: $\mathbf{r}(a) \neq \mathbf{r}(b)$
- **Closed Contours:** Curves that return to their starting point: $\mathbf{r}(a) = \mathbf{r}(b)$
- **Simple Contours:** Curves that don't intersect themselves (except possibly at end-points)
- **Piecewise Contours:** Curves composed of multiple smooth segments




Figure 5.4: *Different types of contours. The piecewise contour shows junction points where segments meet.*

To work with these contours, we need appropriate parameterizations. Here are the fundamental parameterizations that form our computational toolkit:

Table 5.1: Common Parameterized Curves in Line Integrals and Their Arc Length Elements

Curve Type	Parameterization	Parameter Range	Arc Length Element
Line Segment	$\mathbf{r}(t) = (1 - t)(x_1, y_1) + t(x_2, y_2)$ or $(x_1 + t(x_2 - x_1), y_1 + t(y_2 - y_1))$	$0 \leq t \leq 1$	$ds = \sqrt{(\Delta x)^2 + (\Delta y)^2} dt$
Circle	$\mathbf{r}(t) = (a + r \cos t, b + r \sin t)$	$0 \leq t \leq 2\pi$	$ds = r dt$
Circular Arc	$\mathbf{r}(t) = (a + r \cos t, b + r \sin t)$	$\alpha \leq t \leq \beta$	$ds = r dt$
Ellipse	$\mathbf{r}(t) = (a \cos t, b \sin t)$	$0 \leq t \leq 2\pi$	$ds = \sqrt{a^2 \sin^2 t + b^2 \cos^2 t} dt$
Parabola	$\mathbf{r}(t) = (t, at^2)$	$t_1 \leq t \leq t_2$	$ds = \sqrt{1 + 4a^2 t^2} dt$
Any Function	$\mathbf{r}(t) = (t, f(t))$	$t_1 \leq t \leq t_2$	$ds = \sqrt{1 + (f'(t))^2} dt$

For piecewise contours, we use the principle of additivity: when a contour C is composed of several pieces $C_1, C_2, \dots, C_n$, we can break down the integral:

$$\int_C f(x, y) ds = \sum_{i=1}^n \int_{C_i} f(x, y) ds$$

This fundamental property transforms complex integration problems into a series of simpler calculations over basic curves. When working with piecewise contours:

1. Identify the junction points where different segments meet
2. Choose appropriate parameterizations for each segment
3. Ensure consistent orientation throughout the contour
4. Apply the additivity principle to sum the contributions




Figure 5.5: *Breaking down a complex contour into basic elements: line segments (C_1 , C_2 , C_4) and a circular arc (C_3). The arrows indicate the orientation of integration.*

The choice of parameterization affects our calculations but not the final result. When selecting parameterizations:

- Choose parameterizations that simplify the integrand when possible
- Ensure smooth transitions at junction points
- Consider symmetry to simplify calculations
- Remember that different parameterizations of the same curve yield the same integral value

This systematic approach to understanding and working with contours provides the foundation for evaluating line integrals in practice. Whether we're computing work done by a force field or finding the mass of a curved wire, these tools allow us to break complex problems into manageable pieces.

▲ **Example 5.2.2** Consider the scalar function $f(x, y) = x^2 + y$.

1. Compute $\int_C f(x, y) ds$ where C is the path from $(0, 0)$ to $(1, 1)$ consisting of:
 - A quarter circle of radius 1 from $(0, 0)$ to $(1, 1)$
 - A vertical line segment from $(1, 0)$ to $(1, 1)$
2. Compute $\int_C f(x, y) ds$ where C is the curve $y = x^2$ from $(0, 0)$ to $(1, 1)$.

Solution

1. Let's break this into two parts using $\int_C f(x, y) ds = \int_{C_1} f(x, y) ds + \int_{C_2} f(x, y) ds$




Figure 5.6: The piecewise path consisting of a quarter circle (C_1) centered at $(1,1)$ followed by a vertical line segment (C_2).

For C_1 (quarter circle):

- Parameterization: $\mathbf{r}(t) = (1 + \cos t)\mathbf{i} + \sin t\mathbf{j}$, $\frac{\pi}{2} \leq t \leq \pi$
- $\frac{dx}{dt} = -\sin t$, $\frac{dy}{dt} = \cos t$
- $ds = \sqrt{(-\sin t)^2 + (\cos t)^2} dt = dt$
- $f(x(t), y(t)) = (1 + \cos t)^2 + (\sin t) = 1 + 2\cos t + \cos^2 t + \sin t$

The integral over C_1 becomes:

$$\int_{C_1} f(x, y) ds = \int_{\pi/2}^{\pi} (1 + 2\cos t + \cos^2 t + \sin t) dt = \frac{3\pi}{4} - 1 \approx 1.3562$$

For C_2 (vertical line):


- Parameterization: $\mathbf{r}(t) = \mathbf{i} + t\mathbf{j}$, $0 \leq t \leq 1$
- $f(x(t), y(t)) = 1 + t$
- $ds = dt$

Therefore:

$$\begin{aligned} \int_{C_2} f(x, y) ds &= \int_0^1 (1 + t) dt \\ &= \left[t + \frac{t^2}{2}\right]_0^1 = \frac{3}{2} \end{aligned}$$

The total integral is the sum $\frac{3\pi}{4} - 1 + \frac{3}{2} \approx 2.8562$.

2. For the parabola $y = x^2$:

Figure 5.7: The parabolic path $y = x^2$.

- Parameterization: $\mathbf{r}(t) = t\mathbf{i} + t^2\mathbf{j}$, $0 \leq t \leq 1$
- $f(x(t), y(t)) = t^2 + t^2 = 2t^2$
- $\frac{dx}{dt} = 1$, $\frac{dy}{dt} = 2t$
- $ds = \sqrt{1 + 4t^2} dt$

The integral kind of sucks:

$$\int_C f(x, y) ds = \int_0^1 2t^2 \sqrt{1 + 4t^2} dt$$

We get

$$\frac{1}{32} \left(18\sqrt{5} - \sinh^{-1}(2) \right) \approx 1.2127$$

Line Integrals of Vector Fields

A natural avenue to go down would be combining line integrals with vector fields and study their interactions. While scalar line integrals measure accumulation of a scalar quantity along a path (like computing mass from density), vector field line integrals measure how much a vector field “works with” or “works against” our motion along a path. This notion arises naturally in physics when computing work done by a force field.

Consider a particle moving through a force field $\mathbf{F}(x, y) = P(x, y)\mathbf{i} + Q(x, y)\mathbf{j}$. At each point, the force vector has both magnitude and direction. The work done by this force depends on:

- The strength of the force at each point
- How much the force points in our direction of travel
- The distance traveled




Figure 5.8: At each point along the path, the contribution to the line integral depends on how much the vector field aligns with the direction of travel.

This physical intuition leads us to define the line integral of a vector field:

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_C P(x, y) dx + Q(x, y) dy$$

When parameterized by t , this becomes:

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt = \int_a^b \left[P(x(t), y(t)) \frac{dx}{dt} + Q(x(t), y(t)) \frac{dy}{dt} \right] dt.$$

Or more generally:

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_a^b \left[\sum_{i=1}^n F_i(x_1(t), \dots, x_n(t)) \frac{dx_i}{dt} \right] dt$$

The dot product $\mathbf{F} \cdot d\mathbf{r}$ captures precisely what we want:

- The magnitude of $\mathbf{F}$ affects the contribution
- The cosine of the angle between $\mathbf{F}$ and $d\mathbf{r}$ determines how much $\mathbf{F}$ “helps” or “hinders” motion
- Integration sums these contributions along the entire path

This differs fundamentally from scalar line integrals in several ways:

1. **Direction Matters:** Unlike scalar line integrals where ds is always positive, $d\mathbf{r}$ has direction. Reversing path direction changes the sign of the integral:

$$\int_{-C} \mathbf{F} \cdot d\mathbf{r} = - \int_C \mathbf{F} \cdot d\mathbf{r}$$

2. **Path Dependence:** The integral may depend on the path taken, not just endpoints. This leads to the rich theory of conservative versus non-conservative fields.
3. **Physical Interpretation:** Vector field line integrals naturally represent:
 - Work done by force fields
 - Circulation in fluid flows
 - Magnetic flux through surfaces

The computation process reflects these key differences:

1. Parameterize the curve $\mathbf{r}(t)$. Be sure to carefully label the values for which t runs since direction matters.
2. Calculate $\frac{dx}{dt}$ and $\frac{dy}{dt}$ or for n -many dimensions, $\frac{dx_i}{dt}$.
3. Evaluate $P(x(t), y(t))$ and $Q(x(t), y(t))$ or for n -many dimensions, $F_i(x_1(t), \dots, x_n(t))$. Note: for higher dimensions we denote the unit vector $\mathbf{e}_i$ or $\hat{\mathbf{e}}_i$ instead of $\mathbf{i}, \mathbf{j}, \dots$ etc.
4. Form the integrand $P\frac{dx}{dt} + Q\frac{dy}{dt}$, or for n -many dimensions, $\sum_i F_i(x_1(t), \dots, x_n(t))\frac{dx_i}{dt}$
5. Integrate with respect to t , paying careful attention to orientation

▲ **Example 5.2.3** Consider the vector field $\mathbf{F}(x, y) = \frac{-y}{\sqrt{x^2+y^2}}\mathbf{i} + \frac{x}{\sqrt{x^2+y^2}}\mathbf{j}$. Evaluate $\int_C \mathbf{F} \cdot d\mathbf{r}$ where C is the circle of radius 2 centered at the origin:

1. Traversed counterclockwise
2. Traversed clockwise

Solution Let's have a visualization to aid us:




Figure 5.9: The circle C traversed in both directions. The vector field $\mathbf{F}$ (shown in red) is tangent to circles centered at the origin.

For both cases, we'll parameterize the circle but change our parameter range to reflect direction.

1. For counterclockwise traversal:

- Parameterization: $\mathbf{r}(t) = (2 \cos t)\mathbf{i} + (2 \sin t)\mathbf{j}$, $0 \leq t \leq 2\pi$
- $\frac{dx}{dt} = -2 \sin t$, $\frac{dy}{dt} = 2 \cos t$
- $P(x(t), y(t)) = \frac{-2 \sin t}{\sqrt{4 \cos^2 t + 4 \sin^2 t}} = -\sin t$
- $Q(x(t), y(t)) = \frac{2 \cos t}{\sqrt{4 \cos^2 t + 4 \sin^2 t}} = \cos t$

Therefore:

$$\begin{aligned}
 \int_C \mathbf{F} \cdot d\mathbf{r} &= \int_0^{2\pi} [(-\sin t)(-2 \sin t) + (\cos t)(2 \cos t)] dt \\
 &= \int_0^{2\pi} (2 \sin^2 t + 2 \cos^2 t) dt \\
 &= \int_0^{2\pi} 2 dt \\
 &= 4\pi
 \end{aligned}$$

2. For clockwise traversal:

- Parameterization: $\mathbf{r}(t) = (2 \cos t)\mathbf{i} + (2 \sin t)\mathbf{j}$, $2\pi \leq t \leq 0$
- This is equivalent to the negative of the counterclockwise case:

$$\int_{-C} \mathbf{F} \cdot d\mathbf{r} = - \int_C \mathbf{F} \cdot d\mathbf{r} = -4\pi$$

This negative result also makes sense mathematically because of the integral property $\int_a^b f = -\int_b^a f$.

▲ **Example 5.2.4** Consider the vector field $\mathbf{F}(x, y, z) = (y^2)\mathbf{i} + (2xy)\mathbf{j} + (x + z)\mathbf{k}$. Evaluate $\int_C \mathbf{F} \cdot d\mathbf{r}$ along the piecewise path C from $A(0, 0, 0)$ to $D(1, 1, 1)$ consisting of:

- C_1 : From $A(0, 0, 0)$ to $B(1, 0, 0)$ along the x -axis
- C_2 : From $B(1, 0, 0)$ to $C(1, 1, 0)$ parallel to the y -axis
- C_3 : From $C(1, 1, 0)$ to $D(1, 1, 1)$ parallel to the z -axis




Figure 5.10: The path of the curves (a) with the 3D vector field plotted separately using Python (b)

Solution We'll evaluate the line integral along each segment and sum the results:

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_{C_1} \mathbf{F} \cdot d\mathbf{r} + \int_{C_2} \mathbf{F} \cdot d\mathbf{r} + \int_{C_3} \mathbf{F} \cdot d\mathbf{r}$$

We will also make use of

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_a^b \left[P \frac{dx}{dt} + Q \frac{dy}{dt} + R \frac{dz}{dt} \right] dt.$$

1. Along C_1 (from A to B):

- Parameterization: $\mathbf{r}(t) = t\mathbf{i} + 0\mathbf{j} + 0\mathbf{k}$, $0 \leq t \leq 1$
- $\frac{d\mathbf{r}}{dt} = \mathbf{i}$
- Along this path: $y = 0$, $z = 0$, and $x = t$
- Plug into $\mathbf{F} = (y^2)\mathbf{i} + (2xy)\mathbf{j} + (x+z)\mathbf{k}$ to get $\mathbf{F}(\mathbf{r}(t)) = (0)\mathbf{i} + (0)\mathbf{j} + (t)\mathbf{k}$

Therefore:

$$\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_0^1 [(0)(1) + (0)(0) + (t)(0)] dt = 0$$

2. Along C_2 (from B to C):

- Parameterization: $\mathbf{r}(t) = 1\mathbf{i} + t\mathbf{j} + 0\mathbf{k}$, $0 \leq t \leq 1$
- $\frac{d\mathbf{r}}{dt} = \mathbf{j}$
- Along this path: $x = 1$, $z = 0$, and $y = t$
- Plug into $\mathbf{F} = (y^2)\mathbf{i} + (2xy)\mathbf{j} + (x+z)\mathbf{k}$ to get $\mathbf{F}(\mathbf{r}(t)) = (t^2)\mathbf{i} + (2t)\mathbf{j} + (1)\mathbf{k}$

Therefore:

$$\begin{aligned} \int_{C_2} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 [(t^2)(0) + (2t)(1) + (1)(0)] dt \\ &= \int_0^1 2t dt = [t^2]_0^1 = 1 \end{aligned}$$

3. Along C_3 (from C to D):

- Parameterization: $\mathbf{r}(t) = 1\mathbf{i} + 1\mathbf{j} + t\mathbf{k}$, $0 \leq t \leq 1$
- $\frac{d\mathbf{r}}{dt} = \mathbf{k}$
- Along this path: $x = 1$, $y = 1$, and $z = t$
- Plug into $\mathbf{F} = (y^2)\mathbf{i} + (2xy)\mathbf{j} + (x+z)\mathbf{k}$ to get $\mathbf{F}(\mathbf{r}(t)) = (1)\mathbf{i} + (2)(1)\mathbf{j} + (1+t)\mathbf{k}$

Therefore:

$$\begin{aligned} \int_{C_3} \mathbf{F} \cdot d\mathbf{r} &= \int_0^1 [(1)(0) + (2)(0) + (1+t)(1)] dt \\ &= \int_0^1 (1+t) dt = [t + \frac{t^2}{2}]_0^1 = \frac{3}{2} \end{aligned}$$

The total line integral is:

$$\int_C \mathbf{F} \cdot d\mathbf{r} = 0 + 1 + \frac{3}{2} = \frac{5}{2}$$

Potential Functions and Conservative Fields

Let's now explore a special class of vector fields that yield remarkable properties. Consider a vector field $\mathbf{F}(x, y)$ defined on a simply connected domain D . A vector field is conservative if it can be written as the gradient of some scalar function $f(x, y)$:

$$\mathbf{F}(x, y) = \nabla f = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j}$$

This means that $P(x, y) = \frac{\partial f}{\partial x}$ and $Q(x, y) = \frac{\partial f}{\partial y}$. The requirement of a simply connected domain is crucial here—it ensures that any closed path can be continuously deformed to a point, which will prove essential for the path independence of our line integrals. Let's see what happens when we compute a line integral in such a field. Consider a curve C lying entirely in D and parameterized by $\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j}$ from $t = a$ to $t = b$.

Starting with the vector field line integral:

$$\begin{aligned} \int_C \mathbf{F} \cdot d\mathbf{r} &= \int_C P(x, y) dx + Q(x, y) dy \\ &= \int_a^b \left[P(x(t), y(t)) \frac{dx}{dt} + Q(x(t), y(t)) \frac{dy}{dt} \right] dt \end{aligned}$$

Now, because $\mathbf{F}$ is conservative on our simply connected domain D , we can substitute $P = \frac{\partial f}{\partial x}$ and $Q = \frac{\partial f}{\partial y}$:

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_a^b \left[\frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} \right] dt$$

The key insight is recognizing that this is the chain rule for the total derivative of f with respect to t :

$$\begin{aligned} \int_C \mathbf{F} \cdot d\mathbf{r} &= \int_a^b \frac{d}{dt} [f(x(t), y(t))] dt \\ &= f(x(b), y(b)) - f(x(a), y(a)) \\ &= f(B) - f(A) \end{aligned}$$

This derivation reveals the Fundamental Theorem of Line Integrals:

$$\int_C \mathbf{F} \cdot d\mathbf{r} = f(B) - f(A)$$

Note that this result, valid on our simply connected domain D , demonstrates that the line integral depends only on the endpoints A and B , not on the particular path chosen—a powerful consequence of both the conservative nature of $\mathbf{F}$ and the simple connectivity of D . This looks remarkably similar to our good old friend, the Fundamental Theorem of Calculus II.

To build some physical intuition, think of climbing a mountain. The work (force $\times$ displacement) done against gravity (a conservative field) depends only on your starting and ending elevations, not the path you take—whether you climb straight up or take a winding trail. The simple connectivity of the domain ensures there are no “holes” in the mountain that would force you to take fundamentally different paths (imagine if there were a bottomless pit you had to walk around). Meanwhile, the conservative nature of the field ensures that no energy is gained or lost in cycles—if you end where you started, the total work is zero, just as walking around the mountain at the same elevation requires no net work against gravity.

The importance of simply connected domains becomes even more profound when we delve into complex integration in Chapter 7. There, this concept forms the foundation of Cauchy’s Integral Theorem and leads to the remarkable theory of complex analytic functions, where the implications of path independence reach far deeper than in the real case.




Figure 5.11: In a conservative field, the line integral equals the change in potential function value, regardless of path taken. Level curves of f (gray circles) are perpendicular to the field vectors.

This powerful result has several profound implications:

1. **Path Independence:** A vector field $\mathbf{F}$ is path-independent if and only if it’s conservative, meaning the line integral depends only on the endpoints A and B , not on the path taken between them. This is because we’re simply measuring the total change in the potential function f . Remember to check for the conservation by comparing

all combinations of the partial derivatives. For a vector field in n dimensions with components $F_1, F_2, \dots, F_n$:

$$\frac{\partial F_i}{\partial x_j} = \frac{\partial F_j}{\partial x_i} \quad \text{for all } i, j \in \{1, \dots, n\}, i \neq j$$

For example, in three dimensions ($n = 3$ dimensions), this gives us:

$$\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}, \quad \frac{\partial P}{\partial z} = \frac{\partial R}{\partial x}, \quad \frac{\partial Q}{\partial z} = \frac{\partial R}{\partial y}$$

2. **Closed Paths:** If C is a closed path (where $A = B$), then:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = f(A) - f(A) = 0$$

The circle in the integral symbol reminds us we're integrating around a closed loop.

3. **Equipotential Curves:** Along any curve where f is constant (level curves of f), the line integral is zero because there's no change in potential.

In physics, this theorem explains why conservative forces (like gravity) do work that depends only on initial and final positions, not the path taken. The potential function f might represent Gravitational potential energy, Electric potential in an electrostatic field, Pressure potential in a static fluid, and more.

This connection between conservative vector fields and potential functions provides both computational simplification and deep physical insight. When we can identify a potential function, complicated line integral calculations reduce to simple evaluation at endpoints.


Let's take some examples of these bad boys now:

▲ **Example 5.2.5** For each of the following vector fields:

- Check if the field is conservative
- If conservative, find its potential function
- Evaluate the given line integral


Vector fields and regions:

1. $\mathbf{F}_1(x, y) = (3x^2 + y)\mathbf{i} + (x + 4y^3)\mathbf{j}$ on $\mathbb{R}^2$
Evaluate $\oint_C \mathbf{F}_1 \cdot d\mathbf{r}$ along the closed curve C shown:



2. $\mathbf{F}_2(x, y) = e^y \mathbf{i} + xe^y \mathbf{j}$ on $\mathbb{R}^2$

Evaluate $\int_C \mathbf{F}_2 \cdot d\mathbf{r}$ from $A(0, 0)$ to $B(1, 2)$ along the curve shown:



3. $\mathbf{F}_3(x, y) = \frac{-y}{x^2+y^2} \mathbf{i} + \frac{x}{x^2+y^2} \mathbf{j}$ on $\mathbb{R}^2 \setminus \{(0, 0)\}$

Evaluate $\oint_C \mathbf{F}_3 \cdot d\mathbf{r}$ along any circle centered at the origin.

Solution

1. For $\mathbf{F}_1$:

- Check if conservative by comparing partial derivatives:

$$\frac{\partial P}{\partial y} = 1 = \frac{\partial Q}{\partial x}$$

- Since these are equal and the domain is simply connected, $\mathbf{F}_1$ is conservative
- Find potential function by integrating P with respect to x :

$$f(x, y) = \int (3x^2 + y) dx = x^3 + xy + g(y)$$

$$\begin{aligned}\frac{\partial f}{\partial y} &= x + g'(y) = x + 4y^3 \quad (\text{comparing with } Q) \\ g(y) &= y^4 + C\end{aligned}$$

- Therefore, $f(x, y) = x^3 + xy + y^4 + C$
- For any closed curve over a conservative vector: $\oint_C \mathbf{F}_1 \cdot d\mathbf{r} = 0$

2. For $\mathbf{F}_2$:

- Check partial derivatives:

$$\frac{\partial P}{\partial y} = e^y = \frac{\partial Q}{\partial x}$$

- Field is conservative on $\mathbb{R}^2$ (simply connected domain)
- Find potential function:

$$\begin{aligned}f(x, y) &= \int e^y dx = xe^y + g(y) \\ \frac{\partial f}{\partial y} &= xe^y + g'(y) = xe^y \quad (\text{comparing with } Q) \\ g(y) &= C\end{aligned}$$

- Therefore, $f(x, y) = xe^y + C$
- Line integral from $A(0, 0)$ to $B(1, 2)$:

$$\begin{aligned}\int_C \mathbf{F}_2 \cdot d\mathbf{r} &= f(B) - f(A) \\ &= [f(1, 2)] - [f(0, 0)] \\ &= [(1)e^2 + C] - [(0)e^0 + C] \\ &= e^2 + C - C \\ &= e^2\end{aligned}$$

Note how the constant C cancels in the difference.

3. For $\mathbf{F}_3$:

- Check partial derivatives:

$$\begin{aligned}\frac{\partial P}{\partial y} &= \frac{-x^2 + y^2}{(x^2 + y^2)^2} \\ \frac{\partial Q}{\partial x} &= \frac{-x^2 + y^2}{(x^2 + y^2)^2}\end{aligned}$$

- Although partial derivatives are equal, domain is not simply connected (origin removed)
- Therefore $\mathbf{F}_3$ is not conservative
- For a circle of radius r centered at origin, parameterize on $t \in [0, 2\pi]$ using:

$$x = r \cos \theta, \quad y = r \sin \theta$$

- Calculating $\mathbf{F}_3 \cdot d\mathbf{r}$:

$$\begin{aligned}\mathbf{F}_3 &= \frac{-y}{r^2}\mathbf{i} + \frac{x}{r^2}\mathbf{j} = \frac{-\sin \theta}{r}\mathbf{i} + \frac{\cos \theta}{r}\mathbf{j} \\ d\mathbf{r} &= \frac{dx}{d\theta}d\theta\mathbf{i} + \frac{dy}{d\theta}d\theta\mathbf{j} = -r \sin \theta d\theta\mathbf{i} + r \cos \theta d\theta\mathbf{j}\end{aligned}$$

- Taking the dot product:

$$\begin{aligned}\mathbf{F}_3 \cdot d\mathbf{r} &= \left(\frac{-\sin \theta}{r}\right)(-r \sin \theta d\theta) + \left(\frac{\cos \theta}{r}\right)(r \cos \theta d\theta) \\ &= \sin^2 \theta d\theta + \cos^2 \theta d\theta \\ &= (\sin^2 \theta + \cos^2 \theta)d\theta = d\theta\end{aligned}$$

- Therefore, for any circle centered at origin:

$$\oint_C \mathbf{F}_3 \cdot d\mathbf{r} = \int_0^{2\pi} 1 d\theta = 2\pi$$

regardless of the radius r

Lecture 5.3: Green's Theorem

The relationship between derivatives and integrals stands as one of the most profound insights in mathematics. We've seen this connection in various forms: the Fundamental Theorem of Calculus linking derivatives to single integrals, and the relationship between partial derivatives and multiple integrals in higher dimensions. Now we explore another manifestation of this deep connection—Green's Theorem—which relates line integrals around a closed curve to double integrals over the region it encloses. This relationship not only provides computational power but also reveals fundamental properties of vector fields and their behavior in the plane.

The path to understanding Green's Theorem will leverage several powerful tools from our mathematical arsenal. The Mean Value Theorem will help us analyze vector field behavior over small intervals, while Taylor's Theorem will allow us to approximate vector components with increasing accuracy. These classical results, combined with careful limiting arguments, will reveal how local vector field properties accumulate to create global patterns of circulation and flux, which refers to the flow of the vector field through a surface.

Our journey begins with an intuitive understanding through fluid flow, moves through rigorous mathematical development, and culminates in a theorem that bridges boundary behavior with interior properties. Just as the Fundamental Theorem of Calculus shows how differentiation and integration are inverse processes, Green's Theorem will demonstrate how boundary circulation captures the cumulative effect of interior rotation. This lecture represents an important leap in mathematical formalism for advanced theorem's that you may not have encountered before. Don't get caught up too much in the detail, but stop to appreciate it how we lay out the architecture.

Motivation and Mathematical Foundation

Before diving into the mathematics, let's build some intuition. Imagine a lake with a current described by a vector field $\mathbf{F}(x, y) = P(x, y)\mathbf{i} + Q(x, y)\mathbf{j}$. When studying this flow, we might be interested in two fundamental quantities:

- The circulation of the water around the lake's boundary
- How much the water rotates or “swirls” at each point inside the lake

The first quantity measures boundary behavior through a line integral $\oint_C \mathbf{F} \cdot d\mathbf{r}$, while the second examines interior rotation through derivatives $\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}$. Green's Theorem reveals that these seemingly different measurements are intimately connected—the total circulation around the boundary exactly equals the sum of all the interior rotation.

To understand Green's Theorem from a rigorous mathematical perspective, let's begin to build the framework piece by piece using tools we know. Consider a simple closed curve C with four constituent parts $C_1, \dots, C_4$ in the plane, oriented counterclockwise, that bounds a region D . We can think about this curve in two complementary ways:

1. As the boundary where we might evaluate a line integral
2. As the edge of a region where we might compute a double integral




Figure 5.12: A region D bounded by a curve $C = C_1 + C_2 + C_3 + C_4$. We can begin to heuristically see the relationship between the line integral and the volume double integral.

Let's explore why these perspectives might be related. Consider breaking our region into tiny rectangles with sides Δx and Δy . For each rectangle:

- The line integral around its boundary involves terms like $P\Delta x$ and $Q\Delta y$
- The double integral over its area involves terms like $\frac{\partial Q}{\partial x}\Delta x\Delta y$ and $\frac{\partial P}{\partial y}\Delta x\Delta y$

This suggests a possible connection between:

$$\text{Boundary Terms: } P\Delta x + Q\Delta y$$

and

$$\text{Area Terms: } \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \Delta x \Delta y$$




Figure 5.13: Differential element showing contributions of a vector field (P, Q) to the line integral around the boundary of the rectangular area element $dx dy$.

To formalize this intuition, let's examine what happens when we sum the line integral contributions around a single small rectangle with vertices at (x, y) , $(x + \Delta x, y)$, $(x + \Delta x, y + \Delta y)$, and $(x, y + \Delta y)$:

$$\begin{aligned} \oint_{\text{rect}} P dx + Q dy &= \int_x^{x+\Delta x} P(s, y) ds + \int_y^{y+\Delta y} Q(x + \Delta x, t) dt \\ &\quad - \int_{x+\Delta x}^x P(s, y + \Delta y) ds - \int_{y+\Delta y}^y Q(x, t) dt. \end{aligned}$$

Using the Mean Value Theorem and Taylor's Theorem, we expand $Q(x + \Delta x, y)$ and $P(x, y + \Delta y)$ around (x, y) :

$$\begin{aligned} Q(x + \Delta x, y) &= Q(x, y) + \frac{\partial Q}{\partial x} \Delta x + O(\Delta x^2), \\ P(x, y + \Delta y) &= P(x, y) + \frac{\partial P}{\partial y} \Delta y + O(\Delta y^2). \end{aligned}$$

For higher-order terms ($O(\Delta x^2)$, for example), note that these terms vanish in the limit as $\Delta x, \Delta y \rightarrow 0$ due to their smaller magnitude compared to Δx or Δy . After careful algebraic manipulation, the line integral around the rectangle simplifies to:

$$\oint_{\text{rect}} P dx + Q dy = \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \Delta x \Delta y + O(\Delta x^2 \Delta y + \Delta x \Delta y^2).$$

Here, $\Delta x \Delta y$ represents the leading-order contribution proportional to the area of the rectangle, while the higher-order terms vanish faster than $\Delta x \Delta y$ as $\Delta x, \Delta y \rightarrow 0$. This dominant term transitions to the differential $dx dy$ in the integral when taking the limit over all rectangles.

The key insight is this: if we partition the entire region D into such small rectangles and sum their contributions, the following occurs:

- Interior edges cancel in pairs, as they are traversed in opposite directions.
- Only the boundary edges survive, forming the boundary ∂D of the region D .
- The sum of area terms provides an approximation to the double integral over D .

Taking the limit as the partition becomes infinitely fine, let the partition consist of n rectangles, each with area $\Delta A_i = \Delta x_i \Delta y_i$. For each rectangle R_i , the local line integral can be expressed as:

$$\oint_{\partial R_i} P dx + Q dy = \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \Delta A_i + O(\Delta x_i^2 \Delta y_i + \Delta x_i \Delta y_i^2).$$

Summing over all rectangles, we obtain:

$$\begin{aligned}
\sum_{i=1}^n \oint_{\partial R_i} P dx + Q dy &= \sum_{i=1}^n \left[\left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \Delta A_i + O(\Delta x_i^2 \Delta y_i + \Delta x_i \Delta y_i^2) \right] \\
&= \sum_{i=1}^n \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \Delta A_i + \sum_{i=1}^n O(\Delta x_i^2 \Delta y_i + \Delta x_i \Delta y_i^2).
\end{aligned}$$

Let $\delta = \max_i \{\Delta x_i, \Delta y_i\}$ denote the mesh size of the partition. The error term is then bounded by:

$$\left| \sum_{i=1}^n O(\Delta x_i^2 \Delta y_i + \Delta x_i \Delta y_i^2) \right| \leq M \delta \text{Area}(D),$$

for some constant M . As $\delta \rightarrow 0$, the error term vanishes, leaving:

$$\lim_{\delta \rightarrow 0} \sum_{i=1}^n \oint_{\partial R_i} P dx + Q dy = \lim_{\delta \rightarrow 0} \sum_{i=1}^n \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \Delta A_i.$$

On the left-hand side, the sum of line integrals reduces to the line integral around the boundary of the region:

$$\oint_{\partial D} P dx + Q dy.$$

On the right-hand side, the Riemann sum transitions to the double integral over D :

$$\lim_{\delta \rightarrow 0} \sum_{i=1}^n \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \Delta A_i = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA.$$

Through this limiting process, the local relationships derived for small rectangles extend to the entire region D . As the partition becomes infinitely fine ($\delta \rightarrow 0$), the sum of approximations converges to exact values: the line integral around the boundary of D is precisely balanced by the double integral of the curl over the region. This establishes **Green's Theorem**:

$$\oint_{\partial D} \mathbf{F} \cdot d\mathbf{r} = \oint_{\partial D} P dx + Q dy = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA.$$

This theorem stands as a fundamental bridge between different types of integration, much like how the Fundamental Theorem of Calculus I connects derivatives and integrals in single-variable calculus. What's even crazier about this theorem is that it solves so many physics problems without addressing and assuming a lot of physical assumptions. As Richard Dawkins once put it, the potency of a law is the strength of what it proves divided by what it assumes.

We can now substitute line integrals for double integrals wherever it proves advantageous—a frequent and powerful simplification! Do you see anything interesting about the integrand? Anything that connects to Chapter 1 vector concepts? We will soon make an important

generalization of Green's Theorem, but as an exercise, try to see if you can adjust this formula for any dimension n .

Let's consider one more intuitive example. Imagine a region enclosed by a smooth, oriented curve C , as shown in the diagram. Inside this curve, we have several small, uniformly distributed loops, all oriented in the same direction. Each of these loops can be thought of as representing a local rotational field or a "microscopic circulation" at each point in the region. The arrows on the boundary C indicate the macroscopic circulation along the curve's orientation. By Green's Theorem, the total circulation along C is equal to the sum of these individual microscopic rotations, effectively connecting the behavior of the field inside the region to its boundary.

▲ Example 5.3.1 Consider the vector field $\mathbf{F}(x, y) = (x^3y^2)\mathbf{i} + (x^2y^3)\mathbf{j}$. Evaluate $\oint_C \mathbf{F} \cdot d\mathbf{r}$, where C is the triangle with vertices at $(0, 0)$, $(1, 0)$, and $(0, 1)$, oriented counterclockwise.

Solution We'll solve this two ways: first directly using the line integral, and then using Green's Theorem.

1. Method 1: Direct Line Integral

Split the boundary C into three segments:

- C_1 : From $(0, 0)$ to $(1, 0)$,
- C_2 : From $(1, 0)$ to $(0, 1)$,
- C_3 : From $(0, 1)$ to $(0, 0)$.

Let's crunch these now (even though if you look at the vector field you may already know the answer...)

(a) C_1 : From $(0, 0)$ to $(1, 0)$ Parameterize C_1 :

$$\mathbf{r}_1(t) = (t, 0), \quad 0 \leq t \leq 1, \quad \text{so } \frac{d\mathbf{r}_1}{dt} = (1, 0).$$

Evaluate $\mathbf{F} \cdot d\mathbf{r}$ along C_1 :

$$\mathbf{F}(x, y) = (x^3y^2, x^2y^3), \quad \mathbf{F}(\mathbf{r}_1(t)) = (t^3 \cdot 0^2, t^2 \cdot 0^3) = (0, 0).$$

Thus:

$$\int_{C_1} \mathbf{F} \cdot d\mathbf{r} = \int_0^1 (0)(1) + (0)(0) dt = 0.$$

(b) C_2 : From $(1, 0)$ to $(0, 1)$ Parameterize C_2 :

$$\mathbf{r}_2(t) = (1 - t, t), \quad 0 \leq t \leq 1, \quad \frac{d\mathbf{r}_2}{dt} = (-1, 1).$$

Evaluate $\mathbf{F} \cdot d\mathbf{r}$ along C_2 :

$$\mathbf{F}(x, y) = (x^3y^2, x^2y^3), \quad \mathbf{F}(\mathbf{r}_2(t)) = ((1 - t)^3t^2, (1 - t)^2t^3).$$

$$\mathbf{F} \cdot \frac{d\mathbf{r}_2}{dt} = ((1 - t)^3t^2)(-1) + ((1 - t)^2t^3)(1).$$

Simplify:

$$\mathbf{F} \cdot \frac{d\mathbf{r}_2}{dt} = -t^2(1 - t)^3 + t^3(1 - t)^2.$$

Integrate:

$$\int_{C_2} \mathbf{F} \cdot d\mathbf{r} = \int_0^1 (-t^2(1 - 3t + 3t^2 - t^3) + t^3(1 - 2t + t^2)) dt.$$

Expand:

$$\begin{aligned} -t^2(1 - 3t + 3t^2 - t^3) &= -t^2 + 3t^3 - 3t^4 + t^5, \\ t^3(1 - 2t + t^2) &= t^3 - 2t^4 + t^5. \end{aligned}$$

Combine terms:

$$\int_{C_2} \mathbf{F} \cdot d\mathbf{r} = \int_0^1 (-t^2 + 4t^3 - 5t^4 + 2t^5) dt.$$

Integrate term by term:

$$\int_0^1 \mathbf{F} \cdot d\mathbf{r} = \left[-\frac{t^3}{3} + t^4 - \frac{5t^5}{5} + \frac{2t^6}{6} \right]_0^1 = -\frac{1}{3} + 1 - 1 + \frac{1}{3} = 0.$$

(c) C_3 : From $(0, 1)$ to $(0, 0)$ Parameterize C_3 :

$$\mathbf{r}_3(t) = (0, 1 - t), \quad 0 \leq t \leq 1, \quad \frac{d\mathbf{r}_3}{dt} = (0, -1).$$

Evaluate $\mathbf{F} \cdot d\mathbf{r}$ along C_3 :

$$\mathbf{F}(x, y) = (x^3y^2, x^2y^3), \quad \mathbf{F}(\mathbf{r}_3(t)) = (0^3(1 - t)^2, 0^2(1 - t)^3) = (0, 0).$$

Thus:

$$\int_{C_3} \mathbf{F} \cdot d\mathbf{r} = \int_0^1 (0)(0) + (0)(-1) dt = 0.$$

The total line integral is:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \int_{C_1} \mathbf{F} \cdot d\mathbf{r} + \int_{C_2} \mathbf{F} \cdot d\mathbf{r} + \int_{C_3} \mathbf{F} \cdot d\mathbf{r} = 0 + 0 + 0 = 0.$$

2. Method 2: Green's Theorem

Using Green's Theorem:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA.$$

Compute:

$$\frac{\partial Q}{\partial x} = 2xy^3, \quad \frac{\partial P}{\partial y} = 2x^3y, \quad \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} = 2xy^3 - 2x^3y.$$

Set up the integral:

$$\iint_D (2xy^3 - 2x^3y) dA = \int_0^1 \int_0^{1-x} (2xy^3 - 2x^3y) dy dx.$$

Evaluate the inner integral:

$$\int_0^{1-x} (2xy^3 - 2x^3y) dy = \left[2x \frac{y^4}{4} - 2x^3 \frac{y^2}{2} \right]_0^{1-x}.$$

Substituting $y = 1 - x$:

$$\int_0^{1-x} (2xy^3 - 2x^3y) dy = \frac{x(1-x)^4}{2} - x^3(1-x)^2.$$

Finally, integrate with respect to x :

$$\int_0^1 \left(\frac{x(1-x)^4}{2} - x^3(1-x)^2 \right) dx = 0.$$

Both methods confirm:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 0.$$

Now we don't always have to transform line integrals into double integrals. We can use Green's Theorem in both directions. This bidirectional relationship provides a powerful com-

putational tool, allowing us to choose the most elegant solution path. Sometimes, seemingly complex double integrals can be dramatically simplified by recognizing that they represent the integrand of some vector field's differentiated component's differences—a pattern that transforms our volume calculation into a boundary computation.

▲ **Example 5.3.2** In computational fluid dynamics, understanding vorticity patterns often requires calculating complex volume integrals. Consider a basin modeled by the region D bounded by $y = x^2$ and $y = 2 - x^2$ for $x \in [-1, 1]$. A vorticity field in this basin is given by $f(x, y) = xe^y - y \cos(x)$. Our task is to calculate the total vorticity by evaluating $\iint_D f(x, y) dA$.




Figure 5.14: The basin region D bounded by $y = x^2$ and $y = 2 - x^2$. Blue arrows indicate local vorticity patterns, demonstrating the symmetric flow structure about the vertical axis. This visualization helps us understand why the total vorticity integrates to $8[\sin(1) - \cos(1)]$.

Solution Rather than tackle this double integral directly, we can exploit the structure of Green's Theorem to transform our volumetric computation into a boundary analysis:

1. Recognition of Pattern

Our vorticity field $f(x, y) = xe^y - y \cos(x)$ mirrors the curl formula $\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}$ in Green's Theorem:

$$\iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA = \oint_{\partial D} P dx + Q dy$$

2. Construction of Potential Functions

We identify functions P and Q satisfying our pattern:

$$\begin{aligned}P(x, y) &= -xe^y \\ Q(x, y) &= -y \sin(x)\end{aligned}$$

Verification through partial derivatives:

$$\begin{aligned}\frac{\partial Q}{\partial x} &= -y \cos(x) \\ -\frac{\partial P}{\partial y} &= xe^y\end{aligned}$$

$$\text{Therefore: } \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} = xe^y - y \cos(x) = f(x, y)$$

3. Boundary Analysis

Our region's boundary ∂D comprises:

- Upper curve: $y = 2 - x^2$, traversed from $x = -1$ to $x = 1$
- Lower curve: $y = x^2$, traversed from $x = 1$ to $x = -1$

4. Line Integral Formulation

Along the upper curve ($y = 2 - x^2$):

$$\int_{-1}^1 [-xe^{2-x^2} + 2x(2 - x^2) \sin(x)] dx$$

Along the lower curve ($y = x^2$):

$$\int_{-1}^1 [xe^{x^2} + 2x^3 \sin(x)] dx$$

Combining yields:

$$\int_{-1}^1 [x(e^{x^2} - e^{2-x^2}) + 4x \sin(x)] dx$$

5. Symmetry Exploitation

Two key observations simplify our computation:

- $x(e^{x^2} - e^{2-x^2})$ is odd on $[-1, 1]$, contributing zero to the integral
- $4x \sin(x)$ exhibits even symmetry, allowing reduction to $[0, 1]$

Therefore:

$$\begin{aligned}\iint_D f(x, y) dA &= 8 \int_0^1 x \sin(x) dx \\ &= 8[\sin(1) - \cos(1)] \\ &\approx 2.4096\end{aligned}$$

Physical Interpretation: The final value of approximately 2.4096 reveals several key insights about our vorticity field:

- The positive result indicates a net counterclockwise rotation in the basin
- The moderate magnitude suggests balanced vorticity distribution
- The symmetric basin geometry is reflected in our integration technique
- Local rotational effects combine to produce measurable global circulation

Big O Notation

When we work with limits and approximations in calculus, we often encounter situations where we want to describe how quickly certain terms become negligible as a variable approaches zero. Big O notation provides a precise language for describing these behaviors.

In its simplest form, when we write $f(x) = O(g(x))$ as $x \rightarrow a$, we mean that the function $f(x)$ grows no faster than $g(x)$ near a . More precisely, there exist constants $M > 0$ and $\delta > 0$ such that:

$$|f(x)| \leq M|g(x)| \quad \text{whenever } |x - a| < \delta$$

For example, in our analysis of Green's Theorem, we encountered terms like:

$$O(\Delta x^2 \Delta y + \Delta x \Delta y^2)$$

This notation tells us that as Δx and Δy approach zero, these "error terms" become negligible much faster than the main terms ($\Delta x \Delta y$) in our approximation.

A powerful illustration of this notation appears in the limit definition of the derivative. Consider finding $\frac{d}{dx}(x^n)$ using first principles:

$$\begin{aligned}\frac{d}{dx}(x^n) &= \lim_{h \rightarrow 0} \frac{(x+h)^n - x^n}{h} \\ &= \lim_{h \rightarrow 0} \frac{x^n + nx^{n-1}h + \frac{n(n-1)}{2}x^{n-2}h^2 + O(h^3) - x^n}{h} \\ &= \lim_{h \rightarrow 0} \left(nx^{n-1} + \frac{n(n-1)}{2}x^{n-2}h + O(h^2) \right)\end{aligned}$$

$$= nx^{n-1}$$

Here, $O(h^3)$ captures all terms with powers of h greater than 2 in the binomial expansion, and after dividing by h , these terms become $O(h^2)$. As $h \rightarrow 0$, these higher-order terms vanish, leaving us with our familiar result.

Think of Big O notation as a way of classifying the "speed" at which functions approach zero:

- $O(\Delta x)$ approaches zero linearly
- $O(\Delta x^2)$ approaches zero quadratically (much faster)
- $O(\Delta x^3)$ approaches zero even faster

In our proof of Green's Theorem, when we write:

$$Q(x + \Delta x, y) = Q(x, y) + \frac{\partial Q}{\partial x} \Delta x + O(\Delta x^2)$$

we're saying that the error in our linear approximation shrinks quadratically as $\Delta x \rightarrow 0$. This precision in describing approximation errors is crucial for establishing the validity of our limiting arguments.




Figure 5.15: Comparison of convergence rates as $\Delta x \rightarrow 0$. Each curve represents a different order of convergence, demonstrating how higher-order terms approach zero more rapidly. This visualization helps us understand why we can neglect higher-order terms in limit computations.

This notation proves invaluable in calculus as it allows us to focus on dominant terms in approximations and streamline complex limit computations by isolating key terms.

Lecture 5.4: Curl and Divergence

Vector fields appear throughout mathematics and physics—they might represent fluid flow, gravitational forces, or electric fields. While we can visualize these fields through arrows showing direction and magnitude at each point, we often need more sophisticated tools to analyze their behavior. Two fundamental concepts help us understand how vector fields behave locally: curl, which measures rotation, and divergence, which measures expansion or contraction. These operators provide powerful tools for analyzing fluid flow, electromagnetic fields, and many other physical phenomena.

Way back in Chapter 1, we encountered the cross product of vectors—a operation that produced a vector perpendicular to two given vectors, with magnitude related to the parallelogram they formed. This geometric interpretation of the cross product will now take on new life as we use it to measure rotation in vector fields. Similarly, the dot product, which measured how much vectors pointed in the same direction, will help us quantify how much vector fields expand or contract.

The beauty of these concepts lies in how they bridge local and global behaviors. Just as derivatives in single-variable calculus measure instantaneous rates of change, curl and divergence measure instantaneous rotation and expansion in vector fields. And just as integration gives us global information about functions, theorems involving curl and divergence connect local and global properties of vector fields. We will also see an important generalization of Green's Theorem which we touched upon briefly after our proof.

Curl: Measuring Rotation

Imagine stirring a cup of coffee. As your spoon moves through the liquid, it creates a swirling motion—the coffee rotates around points in the fluid. This everyday observation leads us to a profound question in vector calculus: How can we mathematically measure the “amount of rotation” at any point in a vector field?

In the previous lecture, we approached rotation from a global perspective through Green's Theorem, relating the total rotation within a region to the behavior of the field along its boundary. Now we'll zoom in and examine rotation at individual points, stripping away the accumulation of rotational effects to understand the fundamental local behavior. Just as derivatives in single-variable calculus helped us understand rates of change at a point, we'll develop tools to understand rotation at a point. This local perspective will prove invaluable especially in fluid dynamics to electromagnetic theory, for which we will take a few examples.

To approach this systematically, let's start with perhaps the simplest rotating system: a rigid disk spinning with angular velocity ω . From basic physics, we know that a point at radius r from the center moves with speed $v = \omega r$. This linear relationship between speed and radius creates a vector field of velocities:




Figure 5.16: To measure rotation at point P_0 , we probe the vector field $\mathbf{F}$ using a counterclockwise-oriented test circle (red, dashed). The background shows the normalized vector field. At each point on our test circle, we'll compare how well the vector field aligns with the circle's orientation.

Given any vector field $\mathbf{F}$ and any point P_0 in its domain, we can investigate rotation by examining how $\mathbf{F}$ interacts with a geometric probe - specifically, a counterclockwise-oriented 'test' circle we draw centered at P_0 . When we examine the vector field values $\mathbf{F}$ at points along this circle, we notice that some vectors align more closely with our circle's orientation than others. This alignment pattern (or lack thereof) reveals the rotational tendency of $\mathbf{F}$ at P_0 . We can capture this behavior precisely through a line integral, measuring how much $\mathbf{F}$ "follows along" our test circle C_r of radius r :

$$\oint_{C_r} \mathbf{F} \cdot d\mathbf{r}$$

Here, the dot product naturally compares the field's alignment with our circular path at each point, while the integral sums these contributions around the entire circle. But here's the clever part: if we want to measure the "rotation per unit area," we should divide by the circle's area (similar to how we calculated our average efficiency problem from Chapter 4):

$$\frac{1}{\pi r^2} \oint_{C_r} \mathbf{F} \cdot d\mathbf{r}$$

Now, what happens as we make our circle smaller and smaller? We get closer to measuring the pure rotational tendency at point P :

$$\lim_{r \rightarrow 0} \frac{1}{\pi r^2} \oint_{C_r} \mathbf{F} \cdot d\mathbf{r}$$




Figure 5.17: As we shrink the circular path around point P_0 , we measure the local rotational tendency of the field.

Through careful mathematical analysis, this limit equals:

$$\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}.$$

To see this *deep breath*, parameterize the circle C_r of radius r centered at $P_0 = (x_0, y_0)$ using $x = x_0 + r \cos \theta$ and $y = y_0 + r \sin \theta$, where $\theta \in [0, 2\pi]$. The line integral for $\mathbf{F} = P\mathbf{i} + Q\mathbf{j}$ becomes:

$$\oint_{C_r} \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} (P(x_0 + r \cos \theta, y_0 + r \sin \theta)(-r \sin \theta) + Q(x_0 + r \cos \theta, y_0 + r \sin \theta)(r \cos \theta)) d\theta.$$

Expanding P and Q about (x_0, y_0) using a first-order Taylor approximation:

$$P(x_0 + r \cos \theta, y_0 + r \sin \theta) \approx P(x_0, y_0) + \frac{\partial P}{\partial x} r \cos \theta + \frac{\partial P}{\partial y} r \sin \theta,$$

$$Q(x_0 + r \cos \theta, y_0 + r \sin \theta) \approx Q(x_0, y_0) + \frac{\partial Q}{\partial x} r \cos \theta + \frac{\partial Q}{\partial y} r \sin \theta.$$

Substitute these approximations into the integral. Expanding and collecting terms, we observe that the contributions from $P(x_0, y_0)$ and $Q(x_0, y_0)$ involve factors of $\sin \theta$ and $\cos \theta$, respectively:

$$\int_0^{2\pi} P(x_0, y_0)(-r \sin \theta) d\theta = 0, \quad \int_0^{2\pi} Q(x_0, y_0)(r \cos \theta) d\theta = 0,$$

because the integrals of $\sin \theta$ and $\cos \theta$ over $[0, 2\pi]$ are zero.

Next, consider the first-order terms involving $\frac{\partial P}{\partial x}$, $\frac{\partial P}{\partial y}$, $\frac{\partial Q}{\partial x}$, and $\frac{\partial Q}{\partial y}$:

$$\int_0^{2\pi} \left(\frac{\partial P}{\partial x} r^2 \cos \theta (-\sin \theta) + \frac{\partial P}{\partial y} r^2 \sin^2 \theta - \frac{\partial Q}{\partial x} r^2 \cos^2 \theta - \frac{\partial Q}{\partial y} r^2 \cos \theta \sin \theta \right) d\theta.$$

The terms proportional to $\cos \theta \sin \theta$ also vanish because:

$$\int_0^{2\pi} \cos \theta \sin \theta d\theta = 0.$$

The remaining terms simplify using:

$$\int_0^{2\pi} \cos^2 \theta d\theta = \int_0^{2\pi} \sin^2 \theta d\theta = \pi.$$

Thus, the integral reduces to:

$$\oint_{C_r} \mathbf{F} \cdot d\mathbf{r} = r^2 \pi \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right).$$

Dividing by πr^2 and taking the limit as $r \rightarrow 0$, we obtain:

$$\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}.$$

This result can be generalized to three dimensions by considering a vector field $\mathbf{F} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$ and a small, oriented closed loop in space. Using a similar limiting process, we find that the rotational tendency is captured by the *curl* of $\mathbf{F}$:

$$\text{curl } \mathbf{F} = \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) \mathbf{i} + \left(\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x} \right) \mathbf{j} + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \mathbf{k} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix} = \nabla \times \mathbf{F}$$

Conveniently, the cross product provides an elegant closed form for the curl. In three dimensions, for a vector field

$$\mathbf{F}(x, y, z) = P(x, y, z) \mathbf{i} + Q(x, y, z) \mathbf{j} + R(x, y, z) \mathbf{k},$$

the curl is defined as

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix} = \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) \mathbf{i} + \left(\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x} \right) \mathbf{j} + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \mathbf{k}.$$

For two-dimensional vector fields, where the field is given by $\mathbf{F}(x, y) = P(x, y) \mathbf{i} + Q(x, y) \mathbf{j}$ (i.e. with $R(x, y) = 0$), the curl reduces to a vector having only a z -component:

$$\nabla \times \mathbf{F} = \left(0, 0, \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right).$$

It is common in the plane to refer to the scalar

$$\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}$$

as the curl, with a positive value indicating counterclockwise rotation and a negative value indicating clockwise rotation.




Figure 5.18: Patterns of curl in vector fields. Left: Positive curl ($\nabla \times \mathbf{F} > 0$) corresponds to counterclockwise rotation. Right: Negative curl ($\nabla \times \mathbf{F} < 0$) corresponds to clockwise rotation. The direction and magnitude of the curl quantify the rotational tendency of the field at a given point. When we say the curl is positive or negative in this case, we are referring to the $\mathbf{k}$ component.

We have several situations that arise where there is no rotation (also called irrotation). Below we depict them:




Figure 5.19: Vector field patterns that have 0 curl. The arrows clearly show no local rotation about the point.

Green's Theorem Reformulated

Green's Theorem is intimately related to the curl of a vector field. In two dimensions, for a vector field $\mathbf{F}(x, y) = P(x, y)\mathbf{i} + Q(x, y)\mathbf{j}$, we recognize that the full three-dimensional curl reduces to:

$$\nabla \times \mathbf{F} = \left(0, 0, \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right).$$

Thus, we often denote the scalar $\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}$ as the curl, which measures the rotational tendency (about the z -axis) of the field at each point. Substituting this scalar curl into Green's Theorem, we obtain:

$$\oint_{\partial D} \mathbf{F} \cdot d\mathbf{r} = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right) dA.$$

This formulation connects the circulation around the boundary ∂D to the net rotation within the region D .

▲ Example 5.4.1 The electric field in a region of space exhibits complex oscillatory behavior. Consider a simplified model of such a field given by:

$$\mathbf{F}(x, y, z) = e^x \cos(y)\mathbf{i} + e^y \sin(z)\mathbf{j} + e^z \cos(x)\mathbf{k}$$

Calculate curl $\mathbf{F}$ and interpret its physical meaning.

Solution Let's compute the curl of the vector field $\mathbf{F}(x, y, z)$ systematically.

1. Identify the Components:

$$P = e^x \cos(y), \quad Q = e^y \sin(z), \quad R = e^z \cos(x)$$

2. Recall the Formula for Curl: In three dimensions, the curl of $\mathbf{F}$ is given by:

$$\text{curl } \mathbf{F} = \nabla \times \mathbf{F} = \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z}\right)\mathbf{i} + \left(\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x}\right)\mathbf{j} + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y}\right)\mathbf{k}$$

3. Compute Each Partial Derivative:

• **For the $\mathbf{i}$ Component:**

$$\frac{\partial R}{\partial y} = \frac{\partial}{\partial y}(e^z \cos(x)) = 0$$

$$\frac{\partial Q}{\partial z} = \frac{\partial}{\partial z}(e^y \sin(z)) = e^y \cos(z)$$

Thus, the $\mathbf{i}$ component is: $0 - e^y \cos(z) = -e^y \cos(z)$

• For the **j** Component:

$$\frac{\partial P}{\partial z} = \frac{\partial}{\partial z} (e^x \cos(y)) = 0$$

$$\frac{\partial R}{\partial x} = \frac{\partial}{\partial x} (e^z \cos(x)) = -e^z \sin(x)$$

Thus, the **j** component is: $0 - (-e^z \sin(x)) = e^z \sin(x)$

• For the **k** Component:

$$\frac{\partial Q}{\partial x} = \frac{\partial}{\partial x} (e^y \sin(z)) = 0$$

$$\frac{\partial P}{\partial y} = \frac{\partial}{\partial y} (e^x \cos(y)) = -e^x \sin(y)$$

Thus, the **k** component is: $0 - (-e^x \sin(y)) = e^x \sin(y)$

4. **Assemble the Curl:** Combining the components, we obtain:

$$\text{curl } \mathbf{F} = -e^y \cos(z)\mathbf{i} + e^z \sin(x)\mathbf{j} + e^x \sin(y)\mathbf{k}$$

▲ **Example 5.4.2** Recall that a vector field is called conservative if it can be written as the gradient of some scalar function. That is, $\mathbf{F} = \nabla f$ for some scalar function f . Prove that every conservative vector field is irrotational (has curl equal to zero) in three dimensions. Assume f is twice differentiable.

Solution Let's build this from our understanding of gradients and curl:

1. First, let's write out what it means for $\mathbf{F}$ to be conservative. In three dimensions:

$$\mathbf{F} = \nabla f = \frac{\partial f}{\partial x}\mathbf{i} + \frac{\partial f}{\partial y}\mathbf{j} + \frac{\partial f}{\partial z}\mathbf{k}$$

2. This means our components are:

$$P = \frac{\partial f}{\partial x}$$

$$Q = \frac{\partial f}{\partial y}$$

$$R = \frac{\partial f}{\partial z}$$

3. Now, let's examine $\text{curl } \mathbf{F} = \nabla \times \mathbf{F}$. We'll check each component:

(a) **i** component:

$$\begin{aligned}\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} &= \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial z} \right) - \frac{\partial}{\partial z} \left(\frac{\partial f}{\partial y} \right) \\ &= \frac{\partial^2 f}{\partial y \partial z} - \frac{\partial^2 f}{\partial z \partial y} = 0\end{aligned}$$

(b) **j** component:

$$\begin{aligned}\frac{\partial P}{\partial z} - \frac{\partial R}{\partial x} &= \frac{\partial}{\partial z} \left(\frac{\partial f}{\partial x} \right) - \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial z} \right) \\ &= \frac{\partial^2 f}{\partial z \partial x} - \frac{\partial^2 f}{\partial x \partial z} = 0\end{aligned}$$

(c) **k** component:

$$\begin{aligned}\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} &= \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right) - \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right) \\ &= \frac{\partial^2 f}{\partial x \partial y} - \frac{\partial^2 f}{\partial y \partial x} = 0\end{aligned}$$

Each component vanishes due to the equality of mixed partial derivatives (Clairaut's Theorem), assuming f is twice continuously differentiable.

Divergence: Measuring Flow

Just as curl measured rotation by examining how vectors align with circular paths, divergence measures the tendency of a vector field to “spread out” or “compress” at a point. Think of a crowd of people dispersing from a central location—the flow of people creates a pattern of expansion. Or imagine water being sucked down a drain—here the flow pattern shows compression. How can we mathematically capture this intuitive notion of expansion and contraction?

Consider any vector field $\mathbf{F}$ and a point P_0 in its domain. To measure the field's expansive tendency at P_0 , we can use another geometric probe—this time drawing a small closed curve (like a circle) around P_0 and examining how much the field flows across this boundary. But rather than measuring alignment with the boundary (as we did with curl), we'll measure flow through the boundary:




Figure 5.20: To measure divergence at point P_0 , we examine how much the vector field $\mathbf{F}$ flows across a small test curve (red). The outward unit normal vectors $\mathbf{n}$ help us measure the net outward flow.

At each point on our test curve, we can use the dot product $\mathbf{F} \cdot \mathbf{n}$ to measure how much the field flows outward (where $\mathbf{n}$ is the outward unit normal vector). Positive values indicate outward flow, negative values indicate inward flow. The line integral of this outward flux along our curve (ds denoting infinitesimal arc length) gives us the total net outward flow:

$$\oint_{C_r} \mathbf{F} \cdot \mathbf{n} ds$$

Similar to our development of curl, if we want to measure the “flow per unit area”, we should divide by the area enclosed by our curve:

$$\frac{1}{\pi r^2} \oint_{C_r} \mathbf{F} \cdot \mathbf{n} ds$$

This ratio gives us profound insight into the field’s behavior. Consider what happens if we shrink our test circle, making r approach zero:

$$\lim_{r \rightarrow 0} \frac{1}{\pi r^2} \oint_{C_r} \mathbf{F} \cdot \mathbf{n} ds$$

This limit measures the pure expansive tendency at point P_0 - in fluid dynamics, it would tell us whether fluid is being created (positive divergence) or destroyed (negative divergence) at that point. For an incompressible fluid like water, this divergence must be zero everywhere - you can’t create or destroy water in the flow!




Figure 5.21: As we shrink our test curve around P_0 , we measure the local expansive tendency of the field. The normal vectors (red) measure outward flow at each point.

Let's pull out another can of math whoop-ass to get our definition of divergence and start with the aforementioned flux integral around a small circle C_r centered at P_0 :

$$\oint_{C_r} \mathbf{F} \cdot \mathbf{n} ds$$

In polar coordinates our unit normal vector from the origin is $\mathbf{n} = \langle \cos \theta, \sin \theta \rangle$. Thus we have:

$$\mathbf{F} \cdot \mathbf{n} = P \cos \theta + Q \sin \theta$$

Express $\mathbf{F}$ in terms of its components and expand to first order using Taylor Series expansion:

$$\mathbf{F} \cdot \mathbf{n} \approx \left(P + \frac{\partial P}{\partial x} r \cos \theta + \frac{\partial P}{\partial y} r \sin \theta \right) \cos \theta + \left(Q + \frac{\partial Q}{\partial x} r \cos \theta + \frac{\partial Q}{\partial y} r \sin \theta \right) \sin \theta$$

Integrate over the circle and retain the leading terms:

$$\oint_{C_r} \mathbf{F} \cdot \mathbf{n} ds \approx \pi r^2 \left(\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} \right)$$

Divide by the area πr^2 and take the limit as $r \rightarrow 0$:

$$\lim_{r \rightarrow 0} \frac{1}{\pi r^2} \oint_{C_r} \mathbf{F} \cdot \mathbf{n} ds = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} = \nabla \cdot \mathbf{F}$$

$$\operatorname{div} \mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y}$$

The physical interpretation is striking: $\frac{\partial P}{\partial x}$ measures how much the horizontal component changes as we move horizontally, while $\frac{\partial Q}{\partial y}$ measures how much the vertical component changes as we move vertically. Their sum tells us the total spreading out of the field at that

point.




Figure 5.22: Fundamental divergence patterns in vector fields. The divergence measures the net flux per unit area. Left: A source field exhibits positive divergence as vectors emanate outward, indicating fluid expansion. Center: A sink field shows negative divergence as vectors converge inward, representing compression. Right: An incompressible flow field maintains zero divergence, preserving volume as fluid moves. Note how the vector patterns visually represent the mathematical concept of divergence through their geometric arrangement.

In three dimensions, divergence maintains its scalar nature:

$$\operatorname{div} \mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$$

▲ **Example 5.4.3** In fluid dynamics, the velocity field of flow through a pipe often exhibits interesting divergence patterns. Consider the following velocity field modeling fluid flow near a constriction:

$$\mathbf{F}(x, y, z) = (2x)\mathbf{i} + (-y)\mathbf{j} + (-z)\mathbf{k}$$

Calculate the divergence of this field and explain what it reveals about the fluid flow.

Solution This ain't too bad.

1. First, recall that in three dimensions:

$$\operatorname{div} \mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$$

2. For our velocity field:

$$\frac{\partial P}{\partial x} = \frac{\partial}{\partial x}(2x) = 2$$

$$\frac{\partial Q}{\partial y} = \frac{\partial}{\partial y}(-y) = -1$$

$$\frac{\partial R}{\partial z} = \frac{\partial}{\partial z}(-z) = -1$$

3. Therefore:

$$\operatorname{div} \mathbf{F} = 2 - 1 - 1 = 0$$

4. This zero divergence tells us something profound about our flow:

- The fluid is incompressible - no fluid is being created or destroyed
- This matches what we expect in a pipe constriction - as the fluid is squeezed inward in two directions, it must speed up in the third direction to maintain the same volume flow rate




Figure 5.23: *The fluid speeds up where the pipe narrows (indicated by longer arrows) and slows down where it widens, maintaining constant volume throughout.*

This example illustrates a key principle in fluid dynamics: for an incompressible fluid, the divergence of the velocity field must be zero everywhere. In our case, the field automatically satisfies this condition through careful balance of its components.

Vector Operators and Their Compositions

We have now encountered three fundamental vector operators: the curl ($\nabla \times \mathbf{F} = \operatorname{curl} \mathbf{F}$), divergence ($\nabla \cdot \mathbf{F} = \operatorname{div} \mathbf{F}$), and from Chapter 3, the gradient of scalar functions (∇f , we will also notate this as $\operatorname{grad} f$). The gradient, which we've also used implicitly in conservative fields converts a scalar field into a vector field by measuring the direction and magnitude of steepest increase:

$$\operatorname{grad} f = \nabla f = \frac{\partial f}{\partial x} \mathbf{i} + \frac{\partial f}{\partial y} \mathbf{j} + \frac{\partial f}{\partial z} \mathbf{k}$$

These operators form a rich algebraic system with several remarkable properties:

- **Curl of a Gradient:** For any scalar field f (assuming sufficient smoothness),

$$\text{curl}(\text{grad } f) = \nabla \times (\nabla f) = \mathbf{0}$$

This explains why conservative fields (fields that are gradients) are always irrotational!

- **Divergence of a Curl:** For any vector field $\mathbf{F}$,

$$\text{div}(\text{curl } \mathbf{F}) = \nabla \cdot (\nabla \times \mathbf{F}) = 0$$

This means no matter how a field rotates, it never creates or destroys mass in doing so.

- **Second-order Compositions:** The Laplacian emerges when we take divergence of gradient:

$$\text{div}(\text{grad } f) = \nabla \cdot (\nabla f) = \nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}$$

We have some intimate connections between Scalar Fields and Vector Fields that can be intertwined with these three operators:




Figure 5.24: *The relationships between vector operators. Gradient maps scalar to vector fields, divergence does the reverse, and curl maps vector fields to vector fields.*

Some key identities that often prove useful:

1. For scalar fields f and g :

$$\nabla(fg) = f\nabla g + g\nabla f$$

2. For a vector field $\mathbf{F}$:

$$\nabla \cdot (f\mathbf{F}) = f(\nabla \cdot \mathbf{F}) + \mathbf{F} \cdot (\nabla f)$$

3. The vector triple product:

$$\nabla \times (\nabla \times \mathbf{F}) = \nabla(\nabla \cdot \mathbf{F}) - \nabla^2 \mathbf{F}$$

These relationships help explain many physical phenomena. For instance, Maxwell's equations in electromagnetics can be written elegantly using these operators:

- $\nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0}$ (Gauss's law)

- $\nabla \cdot \mathbf{B} = 0$ (No magnetic monopoles)
- $\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$ (Faraday's law)
- $\nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t}$ (Ampère's law)

The fact that $\nabla \cdot (\nabla \times \mathbf{F}) = 0$ ensures magnetic field lines always form closed loops (no magnetic monopoles), while $\nabla \times (\nabla f) = \mathbf{0}$ explains why electric fields in electrostatics can be written as the gradient of a potential.

FTOC Reformulated

The Fundamental Theorem of Calculus can be extended naturally to parametric curves in higher dimensions. Given a scalar field f and a curve $\mathbf{r}(t)$ for $a \leq t \leq b$, we can integrate the gradient along the curve to recover the total change in f :

$$\int_C \nabla f \cdot d\mathbf{r} = f(\mathbf{r}(b)) - f(\mathbf{r}(a))$$

This elegant formulation shows that for any curve, integrating the rate of change of a function along that curve yields the total change in the function. The geometric meaning becomes clear when we recall that ∇f points in the direction of steepest increase of f , and the dot product with $d\mathbf{r}$ measures how much we move in that direction.

Just as the original FTOC connected differentiation and integration in one dimension, this version connects them along any path through space—a profound generalization that bridges single-variable and multivariable calculus.

▲ **Example 5.4.4** Vector operator identities often prove useful in physics and engineering. Derive expressions for:

1. $\nabla \cdot (\mathbf{F} \times \mathbf{G})$ where $\mathbf{F}$ and $\mathbf{G}$ are vector fields
2. $\nabla \times (f\mathbf{F})$ where f is a scalar field and $\mathbf{F}$ is a vector field

Solution

1. For $\nabla \cdot (\mathbf{F} \times \mathbf{G})$:

- Write $\mathbf{F} = F_1\mathbf{i} + F_2\mathbf{j} + F_3\mathbf{k}$ and $\mathbf{G} = G_1\mathbf{i} + G_2\mathbf{j} + G_3\mathbf{k}$
- The cross product is:

$$\begin{aligned} \mathbf{F} \times \mathbf{G} &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ F_1 & F_2 & F_3 \\ G_1 & G_2 & G_3 \end{vmatrix} \\ &= (F_2G_3 - F_3G_2)\mathbf{i} + (F_3G_1 - F_1G_3)\mathbf{j} + (F_1G_2 - F_2G_1)\mathbf{k} \end{aligned}$$

- Take divergence:

$$\begin{aligned}\nabla \cdot (\mathbf{F} \times \mathbf{G}) &= \frac{\partial}{\partial x}(F_2G_3 - F_3G_2) + \frac{\partial}{\partial y}(F_3G_1 - F_1G_3) \\ &\quad + \frac{\partial}{\partial z}(F_1G_2 - F_2G_1)\end{aligned}$$

- Expand using product rule:

$$\begin{aligned}&= \left(\frac{\partial F_2}{\partial x}G_3 + F_2\frac{\partial G_3}{\partial x} - \frac{\partial F_3}{\partial x}G_2 - F_3\frac{\partial G_2}{\partial x} \right) \\ &+ \left(\frac{\partial F_3}{\partial y}G_1 + F_3\frac{\partial G_1}{\partial y} - \frac{\partial F_1}{\partial y}G_3 - F_1\frac{\partial G_3}{\partial y} \right) \\ &+ \left(\frac{\partial F_1}{\partial z}G_2 + F_1\frac{\partial G_2}{\partial z} - \frac{\partial F_2}{\partial z}G_1 - F_2\frac{\partial G_1}{\partial z} \right)\end{aligned}$$

- Collect terms and recognize the dot product pattern:

$$\nabla \cdot (\mathbf{F} \times \mathbf{G}) = \mathbf{G} \cdot (\nabla \times \mathbf{F}) - \mathbf{F} \cdot (\nabla \times \mathbf{G})$$

2. For $\nabla \times (f\mathbf{F})$:

- Write out component-wise, using product rule:

$$\begin{aligned}\nabla \times (f\mathbf{F}) &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ fF_1 & fF_2 & fF_3 \end{vmatrix} \\ &= \left(\frac{\partial}{\partial y}(fF_3) - \frac{\partial}{\partial z}(fF_2) \right) \mathbf{i} \\ &\quad + \left(\frac{\partial}{\partial z}(fF_1) - \frac{\partial}{\partial x}(fF_3) \right) \mathbf{j} \\ &\quad + \left(\frac{\partial}{\partial x}(fF_2) - \frac{\partial}{\partial y}(fF_1) \right) \mathbf{k}\end{aligned}$$

- Apply product rule to each term:

$$\nabla \times (f\mathbf{F}) = f(\nabla \times \mathbf{F}) + \nabla f \times \mathbf{F}$$

These identities are particularly useful in fluid dynamics and electromagnetics. For example, in ideal fluid flow, the term $\nabla \times (f\mathbf{F})$ appears when studying vorticity transport, while $\nabla \cdot (\mathbf{F} \times \mathbf{G})$ arises in the study of helicity in fluid flows.

Vector Operators in Different Coordinate Systems

So far, we've worked exclusively in Cartesian coordinates, where our formulas for gradient, divergence, and curl took relatively simple forms. However, many physical problems are more naturally expressed in other coordinate systems. For instance, studying planetary motion is easier in spherical coordinates, while analyzing flow through a pipe often uses cylindrical coordinates.

Here's how our vector operators transform in common coordinate systems:

Operator	Cylindrical (r, θ, z)	Spherical (r, θ, ϕ)
∇f	$\frac{\partial f}{\partial r} \mathbf{e}_r + \frac{1}{r} \frac{\partial f}{\partial \theta} \mathbf{e}_\theta + \frac{\partial f}{\partial z} \mathbf{e}_z$	$\frac{\partial f}{\partial r} \mathbf{e}_r + \frac{1}{r} \frac{\partial f}{\partial \theta} \mathbf{e}_\theta + \frac{1}{r \sin \theta} \frac{\partial f}{\partial \phi} \mathbf{e}_\phi$
$\nabla \cdot \mathbf{F}$	$\frac{1}{r} \frac{\partial}{\partial r}(r F_r) + \frac{1}{r} \frac{\partial F_\theta}{\partial \theta} + \frac{\partial F_z}{\partial z}$	$\frac{1}{r^2} \frac{\partial}{\partial r}(r^2 F_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta}(\sin \theta F_\theta) + \frac{1}{r \sin \theta} \frac{\partial F_\phi}{\partial \phi}$
$\nabla \times \mathbf{F}$	$\begin{pmatrix} \frac{1}{r} \frac{\partial F_z}{\partial \theta} - \frac{\partial F_\theta}{\partial z} \\ \frac{\partial F_r}{\partial z} - \frac{\partial F_z}{\partial r} \\ \frac{1}{r} \frac{\partial}{\partial r}(r F_\theta) - \frac{1}{r} \frac{\partial F_r}{\partial \theta} \end{pmatrix}$	$\begin{pmatrix} \frac{1}{r \sin \theta} \left(\frac{\partial}{\partial \theta}(\sin \theta F_\phi) - \frac{\partial F_\theta}{\partial \phi} \right) \\ \frac{1}{r} \left(\frac{1}{\sin \theta} \frac{\partial F_r}{\partial \phi} - \frac{\partial}{\partial r}(r F_\phi) \right) \\ \frac{1}{r} \left(\frac{\partial}{\partial r}(r F_\theta) - \frac{\partial F_r}{\partial \theta} \right) \end{pmatrix}$

Table 5.2: Vector operators in cylindrical and spherical coordinates. Note how factors of r and $\sin \theta$ appear to account for the geometry.

These expressions might look intimidating, but they arise naturally from the geometry of these coordinate systems. For instance, the factor of $\frac{1}{r}$ in cylindrical coordinates accounts for the fact that the length of a circular arc increases with radius. Similarly, the $\sin \theta$ terms in spherical coordinates reflect the convergence of longitude lines at the poles.

One particularly useful fact: the expressions for divergence ensure that Gauss's theorem works in any coordinate system—the divergence always represents the flux per unit volume, regardless of how we measure that volume.

▲ **Example 5.4.5** The electric field due to an infinite straight wire carrying current I is given in cylindrical coordinates by:

$$\mathbf{E}(r, \theta, z) = \frac{k}{r} \mathbf{e}_r$$

where k is a constant. Verify that this field is irrotational (curl-free) and compute its divergence.

Solution Let's approach this systematically using our cylindrical coordinate formulas:

1. First, let's identify our components:

$$E_r = \frac{k}{r}, \quad E_\theta = 0, \quad E_z = 0$$

2. To compute curl $\mathbf{E}$, we need all three components:

$$(\nabla \times \mathbf{E})_r = \frac{1}{r} \frac{\partial E_z}{\partial \theta} - \frac{\partial E_\theta}{\partial z} = 0$$

$$(\nabla \times \mathbf{E})_\theta = \frac{\partial E_r}{\partial z} - \frac{\partial E_z}{\partial r} = 0$$

$$(\nabla \times \mathbf{E})_z = \frac{1}{r} \frac{\partial}{\partial r}(rE_\theta) - \frac{1}{r} \frac{\partial E_r}{\partial \theta} = 0$$

Therefore, $\nabla \times \mathbf{E} = \mathbf{0}$, confirming the field is irrotational.

3. For divergence:

$$\nabla \cdot \mathbf{E} = \frac{1}{r} \frac{\partial}{\partial r}(rE_r) + \frac{1}{r} \frac{\partial E_\theta}{\partial \theta} + \frac{\partial E_z}{\partial z}$$

$$= \frac{1}{r} \frac{\partial}{\partial r}(k) + 0 + 0$$

$$= 0$$

The zero divergence everywhere except at $r = 0$ (where the field is undefined) tells us this is a valid solution to Laplace's equation in free space. This matches what we expect for an electric field in a region with no charges. The irrotational nature confirms that this field can be written as the gradient of a scalar potential.

Lecture 5.5: Surface Integrals

Surface integrals are the natural evolution of mathematical curiosity—because apparently, integrating along lines wasn’t ambitious enough. While line integrals let us accumulate quantities along a path, surface integrals take things up a notch by summing contributions across two-dimensional surfaces in three-dimensional space. Useful, if you’ve ever found yourself pondering electromagnetic fields, fluid flow through membranes, or heat transfer across boundaries during your morning tea.

Parameterized Surfaces

Consider the humble soap bubble, floating serenely until my toddler son Andy, an agent of chaos, pokes it with unbridled enthusiasm, leaving only a fleeting memory and some soapy residue. The bubble’s surface—its tension, the way it reflects light, and the air flowing around it—offers a wealth of complexity. Mathematically, analyzing such phenomena requires surface integrals. These integrals come in two flavors: those measuring accumulation over a surface and those measuring flow through a surface (like calculating how much water trickles through a filter). And there you have it: bubbles, toddlers, and integrals, all proving that life’s simplest joys can also be mind-bendingly complicated.

Recall that a parametric curve in space is given by a vector-valued function of one parameter:

$$\mathbf{r}(t) = x(t)\mathbf{i} + y(t)\mathbf{j} + z(t)\mathbf{k}$$

As t varies, this function traces out a one-dimensional path through space. The single parameter t gives us freedom to move in one dimension—along the curve. To describe a surface, we need more freedom of movement. We need to be able to move in two independent directions at any point. This naturally leads us to functions of two parameters:

$$\mathbf{r}(u, v) = x(u, v)\mathbf{i} + y(u, v)\mathbf{j} + z(u, v)\mathbf{k}$$

Think of u and v as coordinates on a “parameter plane”. Each point (u, v) in this plane maps to a point on our surface in three-dimensional space. As we vary both parameters, we sweep out a two-dimensional surface rather than a one-dimensional curve.




Figure 5.25: The parametric mapping $\mathbf{r}(u, v)$ transforms regions in the parameter plane to pieces of surface in three-dimensional space.

When working with surfaces, we often need to convert between parametric and scalar representations. Moving from parametric to scalar form involves writing out how x , y , and z depend on parameters, then systematically eliminating those parameters through substitution and algebra. We must carefully track any parameter restrictions as these become domain constraints in our final scalar equation. Let's see some examples:

▲ **Example 5.5.1** Let's practice converting between parametric and scalar representations of surfaces. These problems progress in difficulty, with the last one presenting a more intricate challenge.

1. Convert the parametric surface

$$\mathbf{r}(u, v) = (u \cos v)\mathbf{i} + (u \sin v)\mathbf{j} + v\mathbf{k}$$

where $u \geq 0$ and $0 \leq v \leq 2\pi$, to scalar form.

2. Convert the parametric surface

$$\mathbf{r}(u, v) = (\sinh u \cos v)\mathbf{i} + (\sinh u \sin v)\mathbf{j} + (\cosh u)\mathbf{k}$$

where $u \geq 0$ and $0 \leq v \leq 2\pi$, to scalar form.

3. Convert the parametric surface

$$\mathbf{r}(u, v) = \left(u\sqrt{1-v^2}\right)\mathbf{i} + (uv)\mathbf{j} + \left(\frac{1}{1-v^2}\right)\mathbf{k}$$

where $-1 < v < 1$ and $u > 0$, to scalar form.

Solution

1. For the first parametric surface:

- The component equations are:

$$x = u \cos v$$

$$y = u \sin v$$

$$z = v$$

- Square and add the x and y equations to eliminate v :

$$x^2 + y^2 = u^2(\cos^2 v + \sin^2 v) = u^2$$

- Since $z = v$, the scalar equation is:

$$x^2 + y^2 = z^2$$

The domain constraint is $x^2 + y^2 \geq 0$ since $u \geq 0$.

2. For the second parametric surface:

- The component equations are:

$$x = \sinh u \cos v$$

$$y = \sinh u \sin v$$

$$z = \cosh u$$

- Square and add the x and y equations to eliminate v :

$$x^2 + y^2 = \sinh^2 u (\cos^2 v + \sin^2 v) = \sinh^2 u$$

- Use the hyperbolic identity $\cosh^2 u - \sinh^2 u = 1$ to express $\sinh^2 u$ in terms of z :

$$x^2 + y^2 = \cosh^2 u - 1 = z^2 - 1$$

- The scalar equation is:

$$x^2 + y^2 + 1 = z^2$$

The domain constraint is $z \geq 1$ since $u \geq 0$.

3. For the third parametric surface:

- The component equations are:

$$x = u\sqrt{1-v^2}$$

$$y = uv$$

$$z = \frac{1}{1-v^2}$$

- Solve for u in terms of x and v :

$$u = \frac{x}{\sqrt{1-v^2}}$$

- Substitute u into $y = uv$:

$$y = \frac{xv}{\sqrt{1-v^2}}$$

- Solve for v^2 in terms of z using the third equation:

$$z = \frac{1}{1-v^2} \Rightarrow 1-v^2 = \frac{1}{z} \Rightarrow v^2 = 1 - \frac{1}{z}$$

Note: Since $-1 < v < 1$ and $u > 0$, we have $z > 1$.

- Substitute $v^2 = 1 - \frac{1}{z}$ into the equation for y :

$$y = \frac{xv}{\sqrt{1-v^2}} = \frac{xv}{\sqrt{\frac{1}{z}}} = xv\sqrt{z}$$

Since $v^2 = 1 - \frac{1}{z}$, we have:

$$v = \pm \sqrt{1 - \frac{1}{z}}$$

Therefore:

$$y = \pm x\sqrt{z}\sqrt{1 - \frac{1}{z}} = \pm x\sqrt{z\left(1 - \frac{1}{z}\right)} = \pm x\sqrt{z-1}$$

- Square both sides to eliminate the square root and the sign:

$$y^2 = x^2(z-1)$$

- Rearrange to obtain the scalar equation:

$$x^2z - x^2 - y^2 = 0$$

Or, equivalently:

$$x^2z = x^2 + y^2$$

Domain Constraint:

$$z > 1$$

Moving from scalar to parametric form requires choosing appropriate parameters based on the surface's geometry. For explicit surfaces, we often use x, y as parameters directly. For cylindrical surfaces, parameters such as height and angle are more appropriate, while spherical coordinates are typically used for spherical surfaces. After selecting the parameters, the coordinates are expressed in terms of these parameters, and their ranges are determined to ensure proper coverage of the surface.

Common strategies for parameterization include the following:

- **Level Curve Method:** This method involves holding one variable constant (e.g., z in the surface $z = f(x, y)$) to obtain level curves. These curves represent intersections

of the surface with horizontal planes and can be parameterized independently. The parameters are then combined to describe the entire surface.

- **Trace Method:** This method examines the intersections of the surface with planes parallel to coordinate planes, such as $x = c$ or $y = c$. Each intersection results in a curve (called a trace) that can be parameterized. By varying the fixed variable and combining the parameterizations, we obtain the parametric form of the surface.
- **Direct Substitution:** For explicitly defined surfaces (e.g., $z = f(x, y)$), x and y can be used directly as parameters, and z is computed from the surface equation.

For the latter, here are some basic forms of parameterization:

1. For $x = f(y, z)$: Use y, z as parameters

$$\mathbf{r}(y, z) = f(y, z)\mathbf{i} + y\mathbf{j} + z\mathbf{k}$$

2. For $y = f(x, z)$: Use x, z as parameters

$$\mathbf{r}(x, z) = x\mathbf{i} + f(x, z)\mathbf{j} + z\mathbf{k}$$

3. For $z = f(x, y)$: Use x, y as parameters

$$\mathbf{r}(x, y) = x\mathbf{i} + y\mathbf{j} + f(x, y)\mathbf{k}$$

These forms are particularly useful when our surface is given by an equation that can be solved for one variable.

▲ **Example 5.5.2** Let's explore three fundamental techniques for working with surface representations. These problems demonstrate the key strategies for converting between parametric and scalar forms.

1. Convert the scalar surface $x^2 + y^2 = 4z^2$ to parametric form using the **level curve method**.
2. Convert the elliptic paraboloid $x = 5y^2 + 2z^2 - 10$ to parametric form using the **trace method**, considering only the portion in front of the yz -plane (where $x \geq 0$).

Solution

1. For the cone $x^2 + y^2 = 4z^2$ (using the **level curve method**):
 - Consider fixing $z = c$, which gives a circle in the x, y -plane:

$$x^2 + y^2 = 4c^2$$

- Parameterize the level curve as:

$$x = 2z \cos \theta, \quad y = 2z \sin \theta$$

- Combine the parameterization with the variable z :

$$\mathbf{r}(z, \theta) = (2z \cos \theta)\mathbf{i} + (2z \sin \theta)\mathbf{j} + z\mathbf{k}$$

where $z \in \mathbb{R}$ and $0 \leq \theta \leq 2\pi$.

2. For the elliptic paraboloid $x = 5y^2 + 2z^2 - 10$ (using the **trace method**):

- Consider fixing $y = c$, which gives a parabola in the xz -plane:

$$x = 5c^2 + 2z^2 - 10$$

- Parameterize x in terms of z and y and z :

$$\mathbf{r}(y, z) = (5y^2 + 2z^2 - 10)\mathbf{i} + y\mathbf{j} + z\mathbf{k}$$

- The domain is constrained by $x \geq 0$, which implies:

$$5y^2 + 2z^2 \geq 10 \quad \text{or equivalently} \quad \frac{y^2}{2} + \frac{z^2}{5} \geq 1$$

Note how each solution leverages different geometric insights: exponential spiral structure in problem 1, circular level curves in problem 2, and parabolic traces in problem 3. The choice of parameterization emerges naturally from each surface's inherent geometric properties.

Tangent Planes and Surface Normal

At each point on our surface, we can move in two independent directions, determined by our parameters u and v . When we vary u slightly (keeping v fixed), we move in the direction of:

$$\mathbf{r}_u = \frac{\partial \mathbf{r}}{\partial u} = \frac{\partial x}{\partial u}\mathbf{i} + \frac{\partial y}{\partial u}\mathbf{j} + \frac{\partial z}{\partial u}\mathbf{k}$$

Similarly, varying v (keeping u fixed) moves us in the direction:

$$\mathbf{r}_v = \frac{\partial \mathbf{r}}{\partial v} = \frac{\partial x}{\partial v}\mathbf{i} + \frac{\partial y}{\partial v}\mathbf{j} + \frac{\partial z}{\partial v}\mathbf{k}$$

These vectors $\mathbf{r}_u$ and $\mathbf{r}_v$ are tangent to the surface and span the tangent plane at each point. Their cross product:

$$\mathbf{n} = \mathbf{r}_u \times \mathbf{r}_v$$

gives us a vector normal to the surface which we can use to construct the plane equation.




Figure 5.26: At each point P on the surface, the vectors $\mathbf{r}_u$ and $\mathbf{r}_v$ span the tangent plane, while their cross product $\mathbf{n}$ gives the normal direction.

The magnitude of this normal vector has special significance:

$$\|\mathbf{r}_u \times \mathbf{r}_v\| = \text{area of parallelogram spanned by } \mathbf{r}_u \text{ and } \mathbf{r}_v$$

This quantity will appear later in our surface integrals because it relates area in the parameter plane to actual surface area in three-dimensional space. When we integrate over a surface, we'll need to account for how the surface might be stretched or compressed relative to the parameter domain.

Now, let's apply this geometric understanding to calculate surface area. Consider how a small rectangular region in the parameter plane, with sides Δu and Δv , maps to our surface. As these lengths become infinitesimal (du and dv), this region approaches a parallelogram formed by the vectors $\mathbf{r}_u du$ and $\mathbf{r}_v dv$. The area of this parallelogram gives us our fundamental area element:

$$dA = \|\mathbf{r}_u \times \mathbf{r}_v\| du dv$$

To find the total surface area, we integrate this element over our parameter domain R :

$$\text{Area} = \iint_R \|\mathbf{r}_u \times \mathbf{r}_v\| du dv$$

This integral captures how our surface stretches and deforms relative to the flat parameter domain. The cross product $\mathbf{r}_u \times \mathbf{r}_v$ not only gives us the surface normal but also, through its magnitude, tells us exactly how much area expansion or compression occurs at each point. This area element will prove fundamental when we begin integrating scalar functions over surfaces.

▲ **Example 5.5.3** Find the surface area of the portion of the paraboloid $z = x^2 + y^2$ that lies inside the cylinder $x^2 + y^2 = 4$.

Solution Let's solve this systematically:

1. Using polar coordinates, parameterize the surface:

$$\mathbf{r}(r, \theta) = r \cos \theta \mathbf{i} + r \sin \theta \mathbf{j} + r^2 \mathbf{k}$$

where $0 \leq r \leq 2$ and $0 \leq \theta \leq 2\pi$

2. Calculate partial derivatives:

$$\mathbf{r}_r = \cos \theta \mathbf{i} + \sin \theta \mathbf{j} + 2r \mathbf{k}$$

$$\mathbf{r}_\theta = -r \sin \theta \mathbf{i} + r \cos \theta \mathbf{j}$$

3. Their cross product $\mathbf{r}_r \times \mathbf{r}_\theta$ is:

$$\begin{aligned} \mathbf{r}_r \times \mathbf{r}_\theta &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \cos \theta & \sin \theta & 2r \\ -r \sin \theta & r \cos \theta & 0 \end{vmatrix} \\ &= (-2r^2 \cos \theta) \mathbf{i} + (-2r^2 \sin \theta) \mathbf{j} + r \mathbf{k} \end{aligned}$$

4. Calculate the magnitude:

$$\begin{aligned} \|\mathbf{r}_r \times \mathbf{r}_\theta\| &= \sqrt{4r^4 \cos^2 \theta + 4r^4 \sin^2 \theta + r^2} \\ &= \sqrt{4r^4(\cos^2 \theta + \sin^2 \theta) + r^2} \\ &= \sqrt{4r^4 + r^2} \\ &= r\sqrt{4r^2 + 1} \end{aligned}$$

5. Set up and evaluate the surface area integral:

$$\begin{aligned} \text{Area} &= \int_0^{2\pi} \int_0^2 r\sqrt{4r^2 + 1} \, dr \, d\theta \\ &= 2\pi \int_0^2 r\sqrt{4r^2 + 1} \, dr \end{aligned}$$

6. For the inner integral, let $u = 4r^2 + 1$:

$$\frac{du}{dr} = 8r \implies r \, dr = \frac{du}{8}$$

$$\begin{aligned}
\text{Area} &= 2\pi \int_1^{17} \frac{\sqrt{u}}{8} du \\
&= \frac{2\pi}{12} [u^{3/2}]_1^{17} \\
&= \frac{2\pi}{12} (17^{3/2} - 1) \\
&\approx 36.175 \text{ square units}
\end{aligned}$$

Surface Integrals of Scalar Functions

Just as we integrated functions over curves in Chapter 2 using arc length, we now face the challenge of integrating over curved surfaces. Then, we needed to account for how curves stretched as we moved along them; now, we must account for how surfaces stretch in two dimensions.

Consider a curved metal sheet with density $\rho(x, y, z)$ varying over its surface. To find its total mass, we need to sum up the mass of infinitesimal surface pieces:

$$\text{Mass} = \iint_S \rho(x, y, z) dS$$

where dS represents an infinitesimal surface area element. This exemplifies a general surface integral for any scalar function $f(x, y, z)$:

$$\iint_S f(x, y, z) dS$$

To compute such integrals, we need to understand how surface area relates to our parameters. Recall that for arc length, we had:

$$ds = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2 + \left(\frac{dz}{dt}\right)^2} dt$$

which measured how much a curve stretched as we moved along it. For surfaces, we need an analogous formula for area. When we parameterize a surface with parameters u and v :

$$\mathbf{r}(u, v) = x(u, v)\mathbf{i} + y(u, v)\mathbf{j} + z(u, v)\mathbf{k}$$

a small rectangular region $[u, u + du] \times [v, v + dv]$ maps to a tiny parallelogram on our surface. The sides of this parallelogram are given by the tangent vectors $\mathbf{r}_u du$ and $\mathbf{r}_v dv$. The area of this parallelogram is:

$$dS = \|\mathbf{r}_u \times \mathbf{r}_v\| du dv$$

Since we've parameterized our surface, each point (x, y, z) is given by $\mathbf{r}(u, v)$. Therefore, our scalar function $f(x, y, z)$ must be composed with $\mathbf{r}$ to give a function of our parameters. Therefore, our surface integral becomes:

$$\iint_S f(x, y, z) dS = \iint_D f(\mathbf{r}(u, v)) \left\| \frac{\partial \mathbf{r}}{\partial u} \times \frac{\partial \mathbf{r}}{\partial v} \right\| du dv$$

where D is the parameter domain.

A common special case occurs when our surface is the graph of a function $z = g(x, y)$. Here, we can use:

$$\mathbf{r}(x, y) = x\mathbf{i} + y\mathbf{j} + g(x, y)\mathbf{k}$$

Then:

$$\begin{aligned}\mathbf{r}_x &= \mathbf{i} + \frac{\partial g}{\partial x} \mathbf{k} \\ \mathbf{r}_y &= \mathbf{j} + \frac{\partial g}{\partial y} \mathbf{k}\end{aligned}$$

Their cross product is:

$$\mathbf{r}_x \times \mathbf{r}_y = -\frac{\partial g}{\partial x} \mathbf{i} - \frac{\partial g}{\partial y} \mathbf{j} + \mathbf{k}$$

Therefore:

$$\|\mathbf{r}_x \times \mathbf{r}_y\| = \sqrt{1 + \left(\frac{\partial g}{\partial x}\right)^2 + \left(\frac{\partial g}{\partial y}\right)^2}$$

Our surface integral for a graph becomes:

$$\iint_S f(x, y, z) dS = \iint_D f(x, y, g(x, y)) \sqrt{1 + \left(\frac{\partial g}{\partial x}\right)^2 + \left(\frac{\partial g}{\partial y}\right)^2} dx dy$$

This formula makes intuitive sense: when our surface is nearly horizontal, the partial derivatives are small and $dS \approx dx dy$. When the surface becomes steeper, the partial derivatives grow larger, increasing dS to account for the additional surface area of the slope.

▲ Example 5.5.4 A curved solar panel is shaped like the paraboloid $z = 4 - x^2 - y^2$ for $-1 \leq x \leq 1$, $-1 \leq y \leq 1$. The efficiency of the panel at each point (x, y, z) is given by $f(x, y, z) = \frac{z}{4}$ (as a percentage of maximum efficiency). Find the average efficiency over the entire panel surface.

Solution To find the average efficiency, we'll integrate $f(x, y, z)$ over the surface and divide by the total surface area.

1. First parameterize the surface using x, y as parameters:

$$\mathbf{r}(x, y) = x\mathbf{i} + y\mathbf{j} + (4 - x^2 - y^2)\mathbf{k}$$

2. Calculate partial derivatives:

$$\mathbf{r}_x = \mathbf{i} + (-2x)\mathbf{k}$$

$$\mathbf{r}_y = \mathbf{j} + (-2y)\mathbf{k}$$

3. Calculate their cross product:

$$\begin{aligned}\mathbf{r}_x \times \mathbf{r}_y &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & -2x \\ 0 & 1 & -2y \end{vmatrix} \\ &= (2x)\mathbf{i} + (2y)\mathbf{j} + \mathbf{k}\end{aligned}$$

4. Find the magnitude:

$$\|\mathbf{r}_x \times \mathbf{r}_y\| = \sqrt{4x^2 + 4y^2 + 1}$$

5. Compose f with our parameterization:

$$f(\mathbf{r}(x, y)) = \frac{4 - x^2 - y^2}{4}$$

6. Set up the integral for total efficiency:

$$\iint_S f \, dS = \int_{-1}^1 \int_{-1}^1 \frac{4 - x^2 - y^2}{4} \sqrt{4x^2 + 4y^2 + 1} \, dx \, dy$$

7. Set up the integral for surface area:

$$\iint_S dS = \int_{-1}^1 \int_{-1}^1 \sqrt{4x^2 + 4y^2 + 1} \, dx \, dy$$

8. The average efficiency is:

$$\text{Average Efficiency} = \frac{\iint_S f \, dS}{\iint_S dS} \approx 0.823$$

▲ **Example 5.5.5** A hemispherical dome of radius 3 meters has a coating whose density varies with height according to $\rho(x, y, z) = z$ kg/m². Find the total mass of the coating.

Solution Let's solve this systematically, paying special attention to determining our bounds.

1. First, we recognize this is part of a sphere. Since it's a hemisphere:

$$x^2 + y^2 + z^2 = 9, \quad z \geq 0$$

2. For a sphere, spherical coordinates provide a natural parameterization:

$$\mathbf{r}(\phi, \theta) = 3 \sin \phi \cos \theta \mathbf{i} + 3 \sin \phi \sin \theta \mathbf{j} + 3 \cos \phi \mathbf{k}$$

3. Determine bounds carefully:

- For θ : We need full rotation, so $0 \leq \theta \leq 2\pi$
- For ϕ : Since $z = 3 \cos \phi \geq 0$, we need $0 \leq \phi \leq \frac{\pi}{2}$

4. Calculate partial derivatives:

$$\begin{aligned}\mathbf{r}_\phi &= 3 \cos \phi \cos \theta \mathbf{i} + 3 \cos \phi \sin \theta \mathbf{j} - 3 \sin \phi \mathbf{k} \\ \mathbf{r}_\theta &= -3 \sin \phi \sin \theta \mathbf{i} + 3 \sin \phi \cos \theta \mathbf{j}\end{aligned}$$

5. Calculate cross product:

$$\|\mathbf{r}_\phi \times \mathbf{r}_\theta\| = 9 \sin \phi$$

6. Express density in terms of parameters:

$$\rho(\mathbf{r}(\phi, \theta)) = 3 \cos \phi$$

7. Set up the mass integral and evaluate:

$$\begin{aligned}M &= \iint_S \rho(x, y, z) dS \\ &= \int_0^{2\pi} \int_0^{\pi/2} (3 \cos \phi)(9 \sin \phi) d\phi d\theta \\ &= 27 \int_0^{2\pi} \int_0^{\pi/2} \cos \phi \sin \phi d\phi d\theta \\ &= 27 \int_0^{2\pi} \left[\frac{\sin^2 \phi}{2} \right]_0^{\pi/2} d\theta \\ &= 27 \int_0^{2\pi} \frac{1}{2} d\theta \\ &= 27\pi \text{ kg}\end{aligned}$$




Figure 5.27: The hemispherical dome with spherical coordinates (ϕ, θ) shown.

The total mass of the coating is 27π kilograms. Note how understanding the geometry helped us determine correct bounds: the hemisphere required $\phi \leq \frac{\pi}{2}$, while needing the full surface required the complete range of θ .

Surface Integrals of Vector Fields

When integrating a vector field over a surface, we need a way to account for both the magnitude of the field and its direction relative to the surface at each point. The dot product with the surface normal provides exactly what we need—it measures the component of the field that passes through the surface, rather than just moving along it.




Figure 5.28: To integrate a vector field $\mathbf{F}$ over a surface S , we must consider both the field's magnitude and its alignment with the surface normal $\mathbf{n}$ at each point. The red vectors represent the vector field, while blue vectors show surface normals.

When working with a surface given by $z = h(x, y)$, we can derive a simpler form for the flux integral. For such a surface, we have a natural parameterization $\mathbf{r}(x, y) = x\mathbf{i} + y\mathbf{j} + h(x, y)\mathbf{k}$. The tangent vectors are:

$$\begin{aligned}\mathbf{r}_x &= \mathbf{i} + \frac{\partial h}{\partial x} \mathbf{k} \\ \mathbf{r}_y &= \mathbf{j} + \frac{\partial h}{\partial y} \mathbf{k}\end{aligned}$$

Taking their cross product:

$$\mathbf{r}_x \times \mathbf{r}_y = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & \frac{\partial h}{\partial x} \\ 0 & 1 & \frac{\partial h}{\partial y} \end{vmatrix} = -\frac{\partial h}{\partial x} \mathbf{i} - \frac{\partial h}{\partial y} \mathbf{j} + \mathbf{k}$$

For a vector field $\mathbf{F} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$, the flux integral becomes:

$$\begin{aligned} \iint_S \mathbf{F} \cdot \mathbf{n} \, dS &= \iint_D \mathbf{F} \cdot \left(-\frac{\partial h}{\partial x} \mathbf{i} - \frac{\partial h}{\partial y} \mathbf{j} + \mathbf{k} \right) dA \\ &= \iint_D \left(-P \frac{\partial h}{\partial x} - Q \frac{\partial h}{\partial y} + R \right) dA \end{aligned}$$

For a general parametric surface $\mathbf{r}(u, v)$, the surface normal comes from the cross product of tangent vectors:

$$\mathbf{r}_u \times \mathbf{r}_v = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial x}{\partial u} & \frac{\partial y}{\partial u} & \frac{\partial z}{\partial u} \\ \frac{\partial x}{\partial v} & \frac{\partial y}{\partial v} & \frac{\partial z}{\partial v} \end{vmatrix}$$

Therefore:

$$\iint_S \mathbf{F} \cdot d\mathbf{S} = \iint_D \mathbf{F} \cdot (\mathbf{r}_u \times \mathbf{r}_v) \, du \, dv$$

This formulation naturally extends our understanding from scalar surface integrals. Just as we needed surface area to integrate scalar functions, we need oriented surface elements to properly account for vector quantities. The cross product $\mathbf{r}_u \times \mathbf{r}_v$ provides both the area element and the direction needed to measure the field's contribution through the surface.

Orientability and Choice of Normal

A crucial concept in surface integrals is orientation. A surface is **orientable** if we can consistently define a normal vector at each point. Most familiar surfaces (spheres, graphs of functions) are orientable, but some (like the Möbius strip) are not. For orientable surfaces, we must choose one of two possible orientations:

- For closed surfaces (like spheres), we typically choose the outward normal
- For surfaces that are graphs $z = g(x, y)$, we typically choose the upward normal
- For general surfaces, we must specify our choice of orientation explicitly

The choice of orientation affects the sign of our surface integrals. If we reverse the orientation (switch from $\mathbf{n}$ to $-\mathbf{n}$), the integral changes sign:

$$\iint_{-S} \mathbf{F} \cdot d\mathbf{S} = - \iint_S \mathbf{F} \cdot d\mathbf{S}$$

This mirrors our experience with line integrals, where reversing direction negated the integral. Understanding orientation becomes crucial when we apply surface integrals to physical problems and when we develop Stokes' and Gauss's theorems in subsequent lectures.

▲ **Example 5.5.6** A cylindrical air filter in a ventilation system has radius 2 meters and height 3 meters. The air flow through the filter is modeled by the vector field $\mathbf{F}(x, y, z) = \frac{x\mathbf{i} + y\mathbf{j}}{x^2 + y^2}$ (measured in m/s). Find the total rate of air flow through the side surface of the filter.

Solution Let's solve this systematically:

1. The side of the cylinder can be parameterized as:

$$\mathbf{r}(\theta, z) = (2 \cos \theta)\mathbf{i} + (2 \sin \theta)\mathbf{j} + z\mathbf{k}$$

where $0 \leq \theta \leq 2\pi$ and $0 \leq z \leq 3$

2. Calculate $\mathbf{r}_\theta$ and $\mathbf{r}_z$:

$$\mathbf{r}_\theta = (-2 \sin \theta)\mathbf{i} + (2 \cos \theta)\mathbf{j}$$

$$\mathbf{r}_z = \mathbf{k}$$

3. Their cross product (pointing outward) is:

$$\mathbf{r}_\theta \times \mathbf{r}_z = (2 \cos \theta)\mathbf{i} + (2 \sin \theta)\mathbf{j}$$

4. Evaluate $\mathbf{F}$ at surface points:

$$\begin{aligned} \mathbf{F}(\mathbf{r}(\theta, z)) &= \frac{2 \cos \theta \mathbf{i} + 2 \sin \theta \mathbf{j}}{4} \\ &= \frac{1}{2}(\cos \theta \mathbf{i} + \sin \theta \mathbf{j}) \end{aligned}$$

5. Set up flux integral:

$$\begin{aligned} \text{Flow Rate} &= \iint_S \mathbf{F} \cdot d\mathbf{S} \\ &= \int_0^3 \int_0^{2\pi} \frac{1}{2}(\cos \theta \mathbf{i} + \sin \theta \mathbf{j}) \cdot (2 \cos \theta \mathbf{i} + 2 \sin \theta \mathbf{j}) d\theta dz \\ &= \int_0^3 \int_0^{2\pi} (\cos^2 \theta + \sin^2 \theta) d\theta dz \end{aligned}$$

6. Evaluate:

$$\text{Flow Rate} = \int_0^3 \int_0^{2\pi} 1 d\theta dz$$

$$\begin{aligned} &= \int_0^3 2\pi \, dz \\ &= 6\pi \text{m}^3/\text{s} \end{aligned}$$

The total flow rate through the filter is $6\pi \approx 18.85$ cubic meters per second. Note that the flow field is designed to model radially outward flow, as would occur in a cylindrical air filter.

Lecture 5.6: Stokes' Theorem

After exploring line integrals, surface integrals, and Green's Theorem, we now arrive at one of the most profound results in vector calculus: **Stokes' Theorem**. Just as Green's Theorem connects a line integral around a closed curve to a double integral over the region it bounds, Stokes' Theorem generalizes this idea to surfaces in three dimensions. It unifies geometric intuition with the behavior of vector fields, revealing how circulation around a boundary curve is related to the curl of a field across the surface.




Figure 5.29: *Visualization of Stokes' Theorem: a portion of the unit sphere Σ emerging from a fluid with outward-pointing normals $\mathbf{n}$. The boundary curve $\partial\Sigma$ is oriented counterclockwise, showing how circulation along the boundary relates to the curl flux through the surface.*

An essential part of applying Stokes' Theorem is choosing a consistent **orientation**. In Green's Theorem, the boundary curve is naturally oriented counterclockwise. For a surface in three-dimensional space, we must pair the boundary's orientation with the normal vector via the right-hand rule. This consistency in orientation is crucial for the theorem's validity.

In our discussion of Green's Theorem, we used an analogy of water flowing around a lake, where circulation around the boundary equaled the total “rotation” within the region. This concept extends to three dimensions. Imagine a curved surface, such as a soap bubble (as in Figure 5.29), with fluid flowing around its edge. Just as rotation in the interior determined boundary circulation in Green's Theorem, so too does the rotational behavior in three dimensions determine the circulation along the boundary curve of a surface.

To make this precise, recall from our previous lecture that when working with surfaces, we measure flow by taking the dot product of the vector field with the surface normal:

$$\text{Flux through surface} = \iint_S \mathbf{F} \cdot \mathbf{n} \, dS.$$

We'll apply this same idea to measure rotation: we take the dot product of the curl (which

measures rotation) with the surface normal.

Consider a small patch of surface bounded by a tiny closed curve. At each point on this curve, the vector field might align perfectly with the curve's direction (maximum circulation), be perpendicular (no circulation), or fall somewhere in between. The total effect—how much the field “follows along” with the boundary—is exactly what we found the line integral measures:

$$\oint_C \mathbf{F} \cdot d\mathbf{r}.$$

Meanwhile, the curl vector at each point indicates the direction and strength of rotation. Only its component parallel to the surface normal contributes to boundary circulation, so we take the dot product:

$$(\text{curl } \mathbf{F}) \cdot \mathbf{n}.$$

Integrating this over the surface gives the total rotation contributing to boundary circulation:

$$\iint_S (\text{curl } \mathbf{F}) \cdot \mathbf{n} dS.$$

Stokes' Theorem tells us these two quantities are equal: the circulation around any boundary matches the total rotation through the surface it bounds. This insight generalizes both Green's Theorem and the Fundamental Theorem of Calculus, showing how local rotation (at each point) accumulates to produce global circulation (around the boundary). Let's get a little more technical now.

The Theorem and Its Meaning

Let Σ be a piecewise smooth, oriented surface in $\mathbb{R}^3$ bounded by a simple closed curve C (where $C = \partial\Sigma$). Let $\mathbf{F}$ be a vector field with continuous first partial derivatives defined on an open region containing Σ . Then Stokes' Theorem states:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_{\Sigma} (\text{curl } \mathbf{F}) \cdot \mathbf{n} d\Sigma.$$

Here, C inherits its orientation from Σ via the right-hand rule: placing your right thumb along $\mathbf{n}$ makes your fingers curl in the positive direction of C . The curl operator, which we studied extensively in Lecture 4, is given by:

$$\text{curl } \mathbf{F} = \nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix}.$$

This result generalizes several theorems we've encountered:

- **Fundamental Theorem of Calculus:** When applied to a curve in one dimension
- **Green's Theorem:** When applied to a region in the plane (as we saw in our reformulation using curl in Lecture 4 of this chapter)

- **Conservative Field Tests:** Demonstrating that curl-free fields have path-independent line integrals

When applying Stokes' Theorem in practice, we need concrete coordinate expressions. Recall that curl in coordinates is:

$$\text{curl } \mathbf{F} = \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) \mathbf{i} - \left(\frac{\partial R}{\partial x} - \frac{\partial P}{\partial z} \right) \mathbf{j} + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \mathbf{k}$$

For a surface with unit normal $\mathbf{n} = \langle n_x, n_y, n_z \rangle$, the crucial dot product becomes:

$$(\text{curl } \mathbf{F}) \cdot \mathbf{n} = \left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) n_x - \left(\frac{\partial R}{\partial x} - \frac{\partial P}{\partial z} \right) n_y + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) n_z$$

We have several fundamental situations to consider:

1. **Level Surface Case:** If Σ is given by $f(x, y, z) = c$:

- The gradient ∇f provides a natural normal vector:

$$\nabla f = \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right\rangle$$

- Normalize to get the unit normal:

$$\mathbf{n} = \frac{\nabla f}{\|\nabla f\|}$$

- The surface integral becomes:

$$\iint_{\Sigma} (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, d\Sigma = \iint_{\Sigma} (\nabla \times \mathbf{F}) \cdot \frac{\nabla f}{\|\nabla f\|} \, d\Sigma$$

- Now if the surface Σ can be written as $z = g(x, y)$, we have:

$$d\Sigma = \sqrt{1 + \left(\frac{\partial z}{\partial x} \right)^2 + \left(\frac{\partial z}{\partial y} \right)^2} \, dx \, dy$$

More generally, if Σ is given by $f(x, y, z) = c$ with $\frac{\partial f}{\partial z} \neq 0$, then:

$$d\Sigma = \frac{\|\nabla f\|}{|\partial f / \partial z|} \, dx \, dy$$

These formulas match when we express z in terms of f . Thus, the complete surface integral $\iint_{\Sigma} (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, d\Sigma$ for a level surface becomes:

$$\iint_D (\nabla \times \mathbf{F}) \cdot \frac{\nabla f}{|\partial f / \partial z|} \, dx \, dy$$

Or expanded out fully:

$$\iint_D \left[\left(\frac{\partial R}{\partial y} - \frac{\partial Q}{\partial z} \right) \frac{\partial f}{\partial x} - \left(\frac{\partial R}{\partial x} - \frac{\partial P}{\partial z} \right) \frac{\partial f}{\partial y} + \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) \frac{\partial f}{\partial z} \right] \frac{dx \, dy}{|\partial f / \partial z|}$$

We can handle that jawn.

2. Parameterized Surface Case: For Σ given by $\mathbf{r}(u, v) = \langle x(u, v), y(u, v), z(u, v) \rangle$:

- The surface element is:

$$d\Sigma = \|\mathbf{r}_u \times \mathbf{r}_v\| du dv$$

- The unit normal is:

$$\mathbf{n} = \frac{\mathbf{r}_u \times \mathbf{r}_v}{\|\mathbf{r}_u \times \mathbf{r}_v\|}$$

- The integral becomes:

$$\iint_{\Sigma} (\nabla \times \mathbf{F}) \cdot \mathbf{n} d\Sigma = \iint_D [(\nabla \times \mathbf{F})(\mathbf{r}(u, v))] \cdot (\mathbf{r}_u \times \mathbf{r}_v) du dv$$

In practice, we choose whichever representation makes our computation simpler. The power of Stokes' Theorem lies in its guarantee that:

$$\oint_{\partial\Sigma} \mathbf{F} \cdot d\mathbf{r} = \iint_{\Sigma} (\nabla \times \mathbf{F}) \cdot \mathbf{n} d\Sigma$$

This equality means we can compute whichever integral is easier and deduce the other. The choice often depends on:

- The geometry of the surface Σ
- The complexity of the vector field $\mathbf{F}$
- Whether boundary or surface integration seems more tractable
- The particular physical or geometric insights we seek

Orientation and Boundary Relationships

The relationship between surface orientation and boundary orientation is crucial. Given an oriented surface Σ , its boundary curve $\partial\Sigma$ inherits an orientation through the right-hand rule:

- Orient your right thumb along the surface normal $\mathbf{n}$
- Your curled fingers indicate the positive direction around $\partial\Sigma$

This convention ensures consistency between the surface integral of curl (measuring rotation through the surface) and the line integral around the boundary (measuring total circulation). If we reverse the surface's orientation (switching $\mathbf{n}$ to $-\mathbf{n}$), we must also reverse the boundary orientation to maintain the theorem's validity:

$$\oint_{-C} \mathbf{F} \cdot d\mathbf{r} = \iint_{-\Sigma} (\text{curl } \mathbf{F}) \cdot \mathbf{n} d\Sigma = - \iint_{\Sigma} (\text{curl } \mathbf{F}) \cdot \mathbf{n} d\Sigma$$

▲ **Example 5.6.1** Let Σ be the portion of the paraboloid $z = x^2 + y^2$ that lies below the plane $z = 1$. Let $\mathbf{F}(x, y, z) = y^2 \mathbf{i} + z^2 \mathbf{j} + x^2 \mathbf{k}$. Evaluate $\oint_C \mathbf{F} \cdot d\mathbf{r}$ where C is the boundary curve oriented counterclockwise when viewed from above.

Solution We will use Stokes' Theorem to evaluate the line integral:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_{\Sigma} (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, d\Sigma$$

where $\mathbf{n}$ is the unit normal vector to the surface Σ .

1. Understanding the Surface Σ :

- The surface Σ is defined by the paraboloid $z = x^2 + y^2$ bounded above by the plane $z = 1$.
- The boundary curve C is the intersection of the paraboloid and the plane, which occurs when $x^2 + y^2 = 1$ at $z = 1$. Thus, C is the circle $x^2 + y^2 = 1$ in the plane $z = 1$.
- The surface Σ can be expressed as the level surface $f(x, y, z) = z - x^2 - y^2 = 0$.

2. Calculating the Curl of $\mathbf{F}$:

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y^2 & z^2 & x^2 \end{vmatrix}$$

Expanding the determinant:

$$\begin{aligned} (\nabla \times \mathbf{F})_x &= \frac{\partial x^2}{\partial y} - \frac{\partial z^2}{\partial z} = 0 - 2z = -2z \\ (\nabla \times \mathbf{F})_y &= \frac{\partial y^2}{\partial z} - \frac{\partial x^2}{\partial x} = 0 - 2x = -2x \\ (\nabla \times \mathbf{F})_z &= \frac{\partial z^2}{\partial x} - \frac{\partial y^2}{\partial y} = 0 - 2y = -2y \end{aligned}$$

Therefore:

$$\nabla \times \mathbf{F} = -2z \mathbf{i} - 2x \mathbf{j} - 2y \mathbf{k}$$

3. Determining the Unit Normal Vector $\mathbf{n}$:

The gradient of $f(x, y, z) = z - x^2 - y^2$ is:

$$\nabla f = \langle -2x, -2y, 1 \rangle$$

The unit normal vector is:

$$\mathbf{n} = \frac{\nabla f}{\|\nabla f\|} = \frac{\langle -2x, -2y, 1 \rangle}{\sqrt{1 + 4x^2 + 4y^2}}$$

4. Setting Up the Surface Integral:

Applying Stokes' Theorem:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_{\Sigma} (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, d\Sigma$$

Substituting the expressions for $\nabla \times \mathbf{F}$ and $\mathbf{n}$:

$$(\nabla \times \mathbf{F}) \cdot \mathbf{n} = (-2z)(-2x) + (-2x)(-2y) + (-2y)(1) = 4zx + 4xy - 2y$$

Since $z = x^2 + y^2$ on Σ :

$$4zx + 4xy - 2y = 4x(x^2 + y^2) + 4xy - 2y = 4x^3 + 4xy^2 + 4xy - 2y$$

Thus, the integral becomes:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_D (4x^3 + 4xy^2 + 4xy - 2y) \, dx \, dy$$

where D is the unit disk $x^2 + y^2 \leq 1$.

5. Determining the Integral Bounds:

To evaluate the integral over the unit disk D , switch to polar. Thus we have

- r ranges from 0 to 1.
- θ ranges from 0 to 2π .

Substituting $x = r \cos \theta$ and $y = r \sin \theta$ into the integrand:

$$\begin{aligned} 4x^3 &= 4(r \cos \theta)^3 = 4r^3 \cos^3 \theta \\ 4xy^2 &= 4(r \cos \theta)(r \sin \theta)^2 = 4r^3 \cos \theta \sin^2 \theta \\ 4xy &= 4(r \cos \theta)(r \sin \theta) = 4r^2 \cos \theta \sin \theta \\ -2y &= -2(r \sin \theta) = -2r \sin \theta \end{aligned}$$

Therefore

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \int_0^{2\pi} \int_0^1 (4r^3 \cos^3 \theta + 4r^3 \cos \theta \sin^2 \theta + 4r^2 \cos \theta \sin \theta - 2r \sin \theta) r \, dr \, d\theta$$

6. Evaluating the Integral Using Symmetry:

Observe the integrand terms:

- $4r^4 \cos^3 \theta$ is an odd function in $\cos \theta$.
- $4r^4 \cos \theta \sin^2 \theta$ is an odd function in $\cos \theta$.
- $4r^3 \cos \theta \sin \theta$ is an odd function in $\cos \theta$ and $\sin \theta$.
- $-2r^2 \sin \theta$ is an odd function in $\sin \theta$.

Thus:

- $\int_0^{2\pi} \cos^3 \theta \, d\theta = 0$
- $\int_0^{2\pi} \cos \theta \sin^2 \theta \, d\theta = 0$
- $\int_0^{2\pi} \cos \theta \sin \theta \, d\theta = 0$
- $\int_0^{2\pi} \sin \theta \, d\theta = 0$

Therefore, each term in the integrand integrates to zero over the interval 0 to 2π :

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 0$$

7. **Conclusion:** The value of the line integral $\oint_C \mathbf{F} \cdot d\mathbf{r}$ is 0. This result is consistent with the symmetry of the vector field and the surface, leading to cancellation of the circulation around the boundary curve.

▲ **Example 5.6.2** Let $\mathbf{F}(x, y, z) = (y^2 + z^2)\mathbf{i} + (z^2 + x^2)\mathbf{j} + (x^2 + y^2)\mathbf{k}$. Compute the line integral $\oint_C \mathbf{F} \cdot d\mathbf{r}$ where C is the intersection of the plane $x + y + z = 1$ and the cylinder $x^2 + y^2 = 1$, oriented counterclockwise when viewed from above. Verify your answer using Stokes' Theorem.

Solution We'll solve this problem by computing both sides of Stokes' Theorem:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_{\Sigma} (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, d\Sigma$$

where Σ is the portion of the plane $x + y + z = 1$ enclosed by the curve C .

1. Direct Computation of the Line Integral:

First, we need to parameterize the curve C , which is the intersection of the plane $x + y + z = 1$ and the cylinder $x^2 + y^2 = 1$.

Let's parameterize the cylinder as $x = \cos t$ and $y = \sin t$ for $0 \leq t \leq 2\pi$. From the plane equation, we get $z = 1 - x - y = 1 - \cos t - \sin t$.

So, our parameterization of C is:

$$\mathbf{r}(t) = \cos t \mathbf{i} + \sin t \mathbf{j} + (1 - \cos t - \sin t) \mathbf{k}, \quad 0 \leq t \leq 2\pi$$

To compute the line integral, we need $d\mathbf{r}/dt$:

$$\frac{d\mathbf{r}}{dt} = -\sin t \mathbf{i} + \cos t \mathbf{j} + (\sin t - \cos t) \mathbf{k}$$

Now, let's evaluate $\mathbf{F}$ along the curve. On C , we have $x^2 + y^2 = 1$ and $z = 1 - \cos t - \sin t$, so:

$$\begin{aligned} \mathbf{F}(\mathbf{r}(t)) &= (\sin^2 t + (1 - \cos t - \sin t)^2) \mathbf{i} \\ &\quad + ((1 - \cos t - \sin t)^2 + \cos^2 t) \mathbf{j} \\ &\quad + (x^2 + y^2) \mathbf{k} \\ &= (\sin^2 t + (1 - \cos t - \sin t)^2) \mathbf{i} \\ &\quad + ((1 - \cos t - \sin t)^2 + \cos^2 t) \mathbf{j} \\ &\quad + (1) \mathbf{k} \end{aligned}$$

The dot product is:

$$\begin{aligned} \mathbf{F} \cdot \frac{d\mathbf{r}}{dt} &= -(\sin^2 t + (1 - \cos t - \sin t)^2) \sin t \\ &\quad + ((1 - \cos t - \sin t)^2 + \cos^2 t) \cos t \\ &\quad + (1)(\sin t - \cos t) \end{aligned}$$

When integrated over $[0, 2\pi]$, terms with $\sin t$, $\cos t$, $\sin^3 t$, $\cos^3 t$, and other odd powers or mixed products all evaluate to zero due to their periodic nature. Therefore:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 0$$

Let's verify this using Stokes' Theorem.

2. Computation Using Stokes' Theorem:

First, we calculate the curl of $\mathbf{F}$:

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y^2 + z^2 & z^2 + x^2 & x^2 + y^2 \end{vmatrix}$$

Computing each component:

$$\begin{aligned}
 (\nabla \times \mathbf{F})_x &= \frac{\partial}{\partial y}(x^2 + y^2) - \frac{\partial}{\partial z}(z^2 + x^2) \\
 &= 2y - 2z \\
 (\nabla \times \mathbf{F})_y &= \frac{\partial}{\partial z}(y^2 + z^2) - \frac{\partial}{\partial x}(x^2 + y^2) \\
 &= 2z - 2x \\
 (\nabla \times \mathbf{F})_z &= \frac{\partial}{\partial x}(z^2 + x^2) - \frac{\partial}{\partial y}(y^2 + z^2) \\
 &= 2x - 2y
 \end{aligned}$$

Therefore:

$$\nabla \times \mathbf{F} = (2y - 2z)\mathbf{i} + (2z - 2x)\mathbf{j} + (2x - 2y)\mathbf{k}$$

Now, we need the unit normal vector to the plane $x + y + z = 1$. The gradient of the plane is:

$$\nabla(x + y + z - 1) = \langle 1, 1, 1 \rangle$$

The unit normal is:

$$\mathbf{n} = \frac{\langle 1, 1, 1 \rangle}{\|\langle 1, 1, 1 \rangle\|} = \frac{\langle 1, 1, 1 \rangle}{\sqrt{3}} = \frac{1}{\sqrt{3}}\langle 1, 1, 1 \rangle$$

Computing the dot product:

$$\begin{aligned}
 (\nabla \times \mathbf{F}) \cdot \mathbf{n} &= \frac{1}{\sqrt{3}}((2y - 2z) + (2z - 2x) + (2x - 2y)) \\
 &= \frac{1}{\sqrt{3}}(0) = 0
 \end{aligned}$$

Since the dot product is zero, the surface integral is:

$$\iint_{\Sigma} (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, d\Sigma = 0$$

Therefore, by Stokes' Theorem:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 0$$

3. Verification Through Direct Integration:

To verify this result more explicitly, let's break down the line integral component by component:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \oint_C (y^2 + z^2) dx + (z^2 + x^2) dy + (x^2 + y^2) dz$$

Using our parameterization:

$$dx = -\sin t \, dt$$

$$dy = \cos t \, dt$$

$$dz = (\sin t - \cos t) \, dt$$

And we know that on C , $x^2 + y^2 = 1$ and $z = 1 - \cos t - \sin t$. So:

$$\begin{aligned} \oint_C (y^2 + z^2) dx &= \int_0^{2\pi} (\sin^2 t + (1 - \cos t - \sin t)^2) \cdot (-\sin t) \, dt \\ \oint_C (z^2 + x^2) dy &= \int_0^{2\pi} ((1 - \cos t - \sin t)^2 + \cos^2 t) \cdot \cos t \, dt \\ \oint_C (x^2 + y^2) dz &= \int_0^{2\pi} (1) \cdot (\sin t - \cos t) \, dt \end{aligned}$$

For the last integral:

$$\begin{aligned} \int_0^{2\pi} (\sin t - \cos t) \, dt &= [-\cos t - \sin t]_0^{2\pi} \\ &= (-\cos(2\pi) - \sin(2\pi)) - (-\cos(0) - \sin(0)) \\ &= (-1 - 0) - (-1 - 0) = 0 \end{aligned}$$

For the first two integrals, when expanded, they contain terms with $\sin t$, $\cos t$, $\sin^2 t \sin t$, $\cos^2 t \cos t$, etc. The integrals of these terms over $[0, 2\pi]$ all vanish due to their periodic nature.

Therefore, the direct calculation confirms:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 0$$

4. Conclusion:

We have verified Stokes' Theorem by calculating both sides of the equation and finding that:

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_{\Sigma} (\nabla \times \mathbf{F}) \cdot \mathbf{n} \, d\Sigma = 0$$

This result makes sense geometrically. The curl vector $\nabla \times \mathbf{F} = (2y - 2z)\mathbf{i} + (2z - 2x)\mathbf{j} + (2x - 2y)\mathbf{k}$ is orthogonal to our normal vector $\mathbf{n} = \frac{1}{\sqrt{3}}\langle 1, 1, 1 \rangle$. Consequently, the flux of the curl through the surface is zero, making the circulation around the boundary also zero.

▲ **Example 5.6.3** Let $\mathbf{F}(x, y, z) = (x^3 + \sin y)\mathbf{i} + (y^3 + xz)\mathbf{j} + (z^3 - xy)\mathbf{k}$ and let V be the region bounded by the hemisphere $x^2 + y^2 + z^2 = a^2$, $z \geq 0$, and the disk $x^2 + y^2 \leq a^2$ in the xy -plane. Verify the Divergence Theorem by computing both:

1. The surface integral $\oint_S \mathbf{F} \cdot \mathbf{n} dS$
2. The volume integral $\iiint_V \operatorname{div} \mathbf{F} dV$

Solution We'll verify the Divergence Theorem by computing both integrals separately.

1. Volume Integral

First, calculate $\operatorname{div} \mathbf{F}$:

$$\begin{aligned} \operatorname{div} \mathbf{F} &= \frac{\partial}{\partial x}(x^3 + \sin y) + \frac{\partial}{\partial y}(y^3 + xz) + \frac{\partial}{\partial z}(z^3 - xy) \\ &= 3x^2 + 0 + 3y^2 + 0 + 3z^2 + 0 \\ &= 3x^2 + 3y^2 + 3z^2 \end{aligned}$$

Using spherical coordinates:

- $x = r \sin \phi \cos \theta$, $y = r \sin \phi \sin \theta$, $z = r \cos \phi$
- $dV = r^2 \sin \phi dr d\phi d\theta$
- Bounds: $0 \leq r \leq a$, $0 \leq \phi \leq \frac{\pi}{2}$, $0 \leq \theta \leq 2\pi$

Substituting the spherical coordinates:

$$\begin{aligned} \operatorname{div} \mathbf{F} &= 3(r \sin \phi \cos \theta)^2 + 3(r \sin \phi \sin \theta)^2 + 3(r \cos \phi)^2 \\ &= 3r^2 \sin^2 \phi \cos^2 \theta + 3r^2 \sin^2 \phi \sin^2 \theta + 3r^2 \cos^2 \phi \\ &= 3r^2 \sin^2 \phi + 3r^2 \cos^2 \phi \\ &= 3r^2(\sin^2 \phi + \cos^2 \phi) \\ &= 3r^2 \end{aligned}$$

The volume integral becomes:

$$\begin{aligned}
 \iiint_V \operatorname{div} \mathbf{F} dV &= \int_0^{2\pi} \int_0^{\pi/2} \int_0^a (3r^2) r^2 \sin \phi dr d\phi d\theta \\
 &= 3 \int_0^{2\pi} \int_0^{\pi/2} \int_0^a r^4 \sin \phi dr d\phi d\theta \\
 &= 3 \int_0^{2\pi} d\theta \int_0^{\pi/2} \sin \phi d\phi \int_0^a r^4 dr \\
 &= 3 \cdot 2\pi \cdot 1 \cdot \frac{a^5}{5} \\
 &= \frac{6\pi a^5}{5}
 \end{aligned}$$

2. Surface Integral

The boundary surface S consists of two parts:

- S_1 : The hemispherical surface $x^2 + y^2 + z^2 = a^2$, $z \geq 0$
- S_2 : The circular disk $x^2 + y^2 \leq a^2$ in the xy -plane

For S_1 (hemisphere):

- The outward normal is $\mathbf{n}_1 = \frac{\langle x, y, z \rangle}{a}$
- Using spherical coordinates:

$$\iint_{S_1} \mathbf{F} \cdot \mathbf{n}_1 dS = \frac{6\pi a^5}{5}$$

For S_2 (disk in the xy -plane):

- The outward normal is $\mathbf{n}_2 = -\mathbf{k}$
- The flux integral evaluates to zero due to odd functions in θ :

$$\iint_{S_2} \mathbf{F} \cdot \mathbf{n}_2 dS = 0$$

The total surface integral is:

$$\oiint_S \mathbf{F} \cdot \mathbf{n} dS = \frac{6\pi a^5}{5} + 0 = \frac{6\pi a^5}{5}$$

Since both the surface integral and volume integral equal $\frac{6\pi a^5}{5}$, the Divergence Theorem is verified for this case.

Physical Interpretation: This example demonstrates how the Divergence Theorem connects the flow characteristics of a vector field throughout a volume to its behavior across the boundary. The field has a uniform expansion rate of $3r^2$ at each point, creating a predictable outward flux proportional to the fifth power of the hemisphere's radius.

Physical Interpretation

Stokes' Theorem provides the mathematical foundation for fundamental laws in electromagnetism. Two of these laws, discovered experimentally in the 19th century, take the form of circulation integrals that become elegant applications of our theorem.

The first is Faraday's Law of Induction, which describes how a changing magnetic field creates an electric field. If we have a loop of wire in space, changes in the magnetic field passing through the loop will induce an electromotive force (EMF) - essentially a voltage that drives current around the wire. Mathematically:

$$\oint_C \mathbf{E} \cdot d\mathbf{r} = -\frac{d}{dt} \iint_S \mathbf{B} \cdot \mathbf{n} dS$$

The left side measures the EMF around the loop (the path integral of the electric field), while the right side measures how quickly the magnetic field passing through the surface is changing. The negative sign tells us the induced EMF opposes the change causing it.

The second fundamental law is Ampère's Law, which tells us that electric currents create magnetic fields:

$$\oint_C \mathbf{B} \cdot d\mathbf{r} = \mu_0 \iint_S \mathbf{J} \cdot \mathbf{n} dS$$

Here $\mathbf{J}$ is the current density (current per unit area) and μ_0 is a physical constant. The left side measures the circulation of the magnetic field around any closed path, while the right side measures the total current passing through any surface bounded by that path. We saw this law in action in our previous example with the straight wire.

When James Clerk Maxwell studied these equations in the 1860s, he realized something profound. Ampère's Law seemed incomplete - it suggested that whenever current flows, magnetic fields circle around it. But what about cases where current is interrupted, like a capacitor charging? Maxwell proposed adding a new term:

$$\oint_C \mathbf{B} \cdot d\mathbf{r} = \mu_0 \iint_S \left(\mathbf{J} + \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} \right) \cdot \mathbf{n} dS$$

The new term $\epsilon_0 \frac{\partial \mathbf{E}}{\partial t}$ represents changing electric fields. Maxwell realized that not only do electric currents create magnetic fields, but changing electric fields do too.




Figure 5.30: This image shows the magnetic force F_2 between two parallel current-carrying wires. The currents (I_1 and I_2) flow through conductors separated by a distance d , with B_1 representing the magnetic field generated by the lower wire. The conductors have length ℓ , illustrating the Ampère-force law and how parallel currents create magnetic fields that interact with each other.

This modification completed a beautiful symmetry in nature: changing magnetic fields create electric fields (Faraday), and changing electric fields create magnetic fields (Maxwell). These insights, expressed mathematically through Stokes' Theorem, led to our modern understanding of electromagnetic waves - disturbances in electric and magnetic fields that propagate through space at the speed of light.

The mathematical structure revealed by Stokes' Theorem - relating circulation around a boundary to flux through a surface - appears repeatedly in physics. Whether studying electromagnetic fields, fluid flows, or gravitational fields, this fundamental connection between local rotation and global circulation proves indispensable for understanding natural phenomena.

▲ **Example 5.6.4** A long straight wire carries a current I in the positive z -direction, creating a magnetic field $\mathbf{B}$. Using Ampère's law and Stokes' Theorem:

1. Find the magnetic field $\mathbf{B}$ at any point (x, y) .
2. Verify your answer by directly calculating $\oint_C \mathbf{B} \cdot d\mathbf{r}$ around a circular loop of radius a centered on the wire.

Solution

1. Let's use our basic definitions from above and what we know about Stokes' Theorem:

- (a) First, recall Ampère's law in differential form:

$$\text{curl } \mathbf{B} = \mu_0 \mathbf{J}$$

where $\mathbf{J}$ is the current density. We know from physics that:

- The field lines should form concentric circles around the wire
- The field strength should decrease with distance from the wire
- The field direction follows the right-hand rule relative to current

- (b) This suggests a field of the form:

$$\mathbf{B}(x, y) = \frac{k}{x^2 + y^2} (-y\mathbf{i} + x\mathbf{j})$$

where k is a constant to be determined.

- (c) Let's choose a circular disk Σ of radius a centered on the wire at height $z = 0$:

- Its boundary C is oriented counterclockwise when viewed from above
- By Stokes' Theorem: $\oint_C \mathbf{B} \cdot d\mathbf{r} = \iint_{\Sigma} (\text{curl } \mathbf{B}) \cdot \mathbf{n} d\Sigma$
- The right side must equal $\mu_0 I$ by Ampère's law

2. Now let's verify it.

- (a) Let's calculate the line integral directly:

- Parameterize C as $\mathbf{r}(t) = (a \cos t)\mathbf{i} + (a \sin t)\mathbf{j}$, $0 \leq t \leq 2\pi$
- Along C , $\mathbf{B} = \frac{k}{a^2}(-a \sin t \mathbf{i} + a \cos t \mathbf{j})$
- $d\mathbf{r} = (-a \sin t \mathbf{i} + a \cos t \mathbf{j}) dt$

- (b) Therefore:

$$\begin{aligned} \oint_C \mathbf{B} \cdot d\mathbf{r} &= \int_0^{2\pi} \frac{k}{a^2} (a^2 \sin^2 t + a^2 \cos^2 t) dt \\ &= \int_0^{2\pi} k dt = 2\pi k = \mu_0 I \end{aligned}$$

- (c) Solving for k :

$$k = \frac{\mu_0 I}{2\pi}$$

(d) Thus, the magnetic field is:

$$\mathbf{B}(x, y) = \frac{\mu_0 I}{2\pi(x^2 + y^2)} (-y\mathbf{i} + x\mathbf{j})$$

3. Important notes on orientation:

- The orientation of C and the upward normal $\mathbf{n}$ to Σ are related by the right-hand rule
- If we reverse the surface orientation (flip $\mathbf{n}$), we must also reverse C 's orientation
- The physical field is independent of our choice of orientation
- Any surface Σ' with boundary C gives the same result, showing path independence

Generalization and Abstraction

Time to get a little crazy. Throughout our studies of calculus, we've encountered several seemingly different theorems that share a remarkable pattern:

1. **Fundamental Theorem of Calculus:**

$$\int_a^b \frac{df}{dx} dx = f(b) - f(a)$$

2. **Green's Theorem:**

$$\oint_C (P dx + Q dy) = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA$$

3. **Stokes' Theorem:**

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_{\Sigma} (\text{curl } \mathbf{F}) \cdot \mathbf{n} d\Sigma$$

4. **Divergence Theorem:**

$$\oiint_S \mathbf{F} \cdot \mathbf{n} dS = \iiint_V (\text{div } \mathbf{F}) dV$$

In each case, we integrate an object over a boundary and relate it to an integral over the region that the boundary encloses. To unify these results, we introduce several new concepts. First, we introduce the notion of a **differential form**. Differential forms generalize the functions and expressions we integrate, incorporating geometric information such as orientation. In one dimension, we integrate expressions like

$$f(x) dx,$$

where dx is a 1-form—it takes a tangent vector and returns the component of that vector in the x -direction. In two dimensions, when integrating along a curve (a 1-dimensional manifold in $\mathbb{R}^2$), we use a 1-form such as

$$\omega = P dx + Q dy.$$

On the other hand, when integrating over a region (a 2-dimensional manifold), we use a 2-form such as

$$\eta = dx \wedge dy,$$

where the wedge product $\wedge$ is antisymmetric ($dx \wedge dy = -dy \wedge dx$) and encodes orientation. More generally, a k -form on an n -dimensional space is an object that can be integrated over a k -dimensional surface. The **exterior derivative** d is an operator that maps k -forms to $(k + 1)$ -forms:

- It takes a function f (a 0-form) to its differential df (a 1-form).
- It takes a 1-form like $\omega = P dx + Q dy$ to the 2-form

$$d\omega = \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx \wedge dy.$$

- It similarly maps 2-forms to 3-forms, and so on.

With these tools, we can state one unified theorem:

Generalized Stokes' Theorem: For any differential form ω defined on a smooth, oriented manifold Ω with boundary $\partial\Omega$,

$$\int_{\Omega} d\omega = \int_{\partial\Omega} \omega.$$

This elegant formula encapsulates our earlier theorems:

- For the Fundamental Theorem of Calculus, Ω is an interval $[a, b]$ and $\omega = f dx$.
- For Green's Theorem, we consider a 1-form $\omega = P dx + Q dy$ defined on a region $D \subset \mathbb{R}^2$; its exterior derivative is the 2-form

$$d\omega = \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx \wedge dy,$$

so that $\int_{\partial D} \omega$ equals $\iint_D d\omega$.

- For classical Stokes' Theorem, Ω is a surface and ω is a 1-form corresponding to a vector field.
- For the Divergence Theorem, Ω is a volume and ω is a 2-form dual to the vector field.

The power of this formulation lies in its abstraction: it applies to any manifold—any space that locally resembles $\mathbb{R}^n$ —and reveals the deep relationship between integration and differentiation via the boundary operator. This unification appears throughout mathematics and physics, from electromagnetism to quantum field theory.

Two key properties underpin this framework:

1. $d^2 = 0$: The exterior derivative applied twice always yields zero.
2. $\partial^2 = 0$: The boundary of a boundary is always empty.

These properties ensure the theorems interlock perfectly, regardless of the space's dimension or shape, and they lay the groundwork for the rich theory of de Rham cohomology, which studies the topology of spaces via differential forms. Even if we don't explore these advanced concepts further, recognizing that all our integration theorems arise from a single underlying principle helps explain their common structure and wide applicability. I leave you with this meme I made:




Figure 5.31: *The master of all: Stokes' Theorem*

Lecture 5.7: Divergence Theorem

After our exploration of Stokes' Theorem and surface integrals, we now arrive at another fundamental result that connects the behavior of vector fields inside a region to their behavior on the boundary. The Divergence Theorem (also known as Gauss's Theorem) relates the flux of a vector field across a closed surface to the total divergence within the volume it encloses. This relationship proves crucial in physics—particularly in electromagnetism and fluid dynamics—where it helps us understand how fields “flow” through space.

Just as Stokes' Theorem generalized Green's Theorem to surfaces in three dimensions, the Divergence Theorem extends this pattern further, connecting surface integrals to volume integrals. This progression reveals a deep pattern in vector calculus: integrals over boundaries relate to integrals over the regions they bound, regardless of dimension. We'll explore this pattern more fully after developing the theorem itself.

Statement and Meaning

Throughout our studies of vector calculus, we've encountered divergence as a way to measure how much a vector field “spreads out” at each point. A positive divergence indicates a source—a point where field lines emerge—while negative divergence signifies a sink where field lines converge. Now we'll discover a profound connection between this local behavior and the global flow across boundaries.

Consider a single small volume element in three-dimensional space, with a vector field $\mathbf{F}$ flowing through it. At each face of this volume, we can measure the flux—how much of the field passes through that face. The flux through each face is given by the dot product of $\mathbf{F}$ with the outward normal vector, multiplied by the area of the face. If we sum these contributions over all faces:

$$\sum_{\text{faces}} \mathbf{F} \cdot \mathbf{n} d\Phi$$

where $\mathbf{n}$ points outward from each face and $d\Phi$ represents the area of a face element, we get the net outward flux. For a small enough volume element dV , this flux should approximately equal the divergence within the volume multiplied by the volume:

$$\sum_{\text{faces}} \mathbf{F} \cdot \mathbf{n} d\Phi \approx (\operatorname{div} \mathbf{F}) dV$$

This local relationship suggests something deeper. Just as Green's Theorem connected boundary circulation to interior curl, here we see flux through boundaries connecting to interior divergence. Consider dividing an entire region into smaller subvolumes, as shown in Figure 5.32. When we add up all these small volume contributions, interior faces cancel in pairs (since adjacent volumes share faces with opposite orientations), leaving only the boundary faces.




Figure 5.32: (a) Adjacent subvolumes V_i and V_j sharing an interface Φ through which the field flows. (b) Complete subdivision of a volume showing all interfaces (green). Interior interfaces cancel in pairs, leaving only boundary contributions.

When two subvolumes V_i and V_j share an interface Φ , as shown in Figure 5.32(a), the flux through this interface appears twice in our sum—once for each subvolume but with opposite signs, since what flows out of V_i must flow into V_j . As we sum over all subvolumes, these interior contributions cancel in pairs, leaving only the boundary faces, as visualized in Figure 5.32(b). Taking the limit as our subdivision becomes infinitesimal, we arrive at:

$$\oiint_S \mathbf{F} \cdot \mathbf{n} dS = \iiint_V \operatorname{div} \mathbf{F} dV$$

This is the Divergence Theorem. The left side measures total flux across the boundary surface S , while the right side accumulates the divergence throughout the volume V . Their equality tells us something profound: the net outward flow across any closed surface exactly equals the total “source strength” within.

To make this concrete, imagine a fluid flowing through space with velocity field $\mathbf{F}$. The surface integral $\oiint_S \mathbf{F} \cdot \mathbf{n} dS$ measures how much fluid escapes through the boundary per unit time. Meanwhile, $\operatorname{div} \mathbf{F}$ at each point tells us the rate at which fluid is being created (if positive) or destroyed (if negative) at that point, so $\iiint_V \operatorname{div} \mathbf{F} dV$ gives the total rate of fluid generation within the volume. Their equality expresses a fundamental conservation principle: what flows out must be created within.

Gauss’s Law

This relationship extends far beyond fluid flow. In electromagnetism, when $\mathbf{F}$ is the electric field $\mathbf{E}$, the theorem becomes Gauss’s law:

$$\oiint_S \mathbf{E} \cdot \mathbf{n} dS = \frac{1}{\epsilon_0} \iiint_V \rho dV$$

where ρ is charge density and ϵ_0 is a physical constant. Here $\operatorname{div} \mathbf{E} = \rho/\epsilon_0$, relating the spreading of electric field lines to the presence of charge. Electric field lines begin on positive charges and end on negative charges—another manifestation of this deep connection between boundary behavior and interior sources. This behavior is illustrated in Figure 5.33.




Figure 5.33: Electric field lines (shown in green) emerging from a positive charge and terminating on a negative charge. The divergence $\nabla \cdot \mathbf{F}$ is positive near the source and negative near the sink. For any closed surface Ω (shown in blue) with surface element dS and outward normal $\hat{\mathbf{n}}$, the total flux through the surface equals the net charge enclosed, scaled by ϵ_0 .

The figure shows how field lines emanate from positive charges (sources where $\nabla \cdot \mathbf{F} > 0$) and converge at negative charges (sinks where $\nabla \cdot \mathbf{F} < 0$). For any closed surface Ω enclosing these charges, the net flux through surface elements dS with outward normal $\mathbf{n}$ exactly equals the enclosed charge divided by ϵ_0 . This provides a powerful method for calculating electric fields: if we can find a surface where symmetry makes the flux calculation simple, we can determine the field without directly solving differential equations.

Just as Green's Theorem connected line integrals to double integrals in the plane, the Divergence Theorem bridges surface integrals and volume integrals in space. This progression reveals a pattern: integrals over boundaries relate to integrals over the regions they bound, with the relationship governed by derivatives of the fields involved. We'll explore this pattern more fully after examining the theorem's applications and computational techniques.

Orientation and Boundary Conditions

For the Divergence Theorem, orientation proves crucial. The surface normal $\mathbf{n}$ must point outward from the enclosed volume. This convention ensures:

- Positive flux represents outward flow
- Negative flux represents inward flow
- The total signed flux captures net flow across the boundary

For a surface enclosing a volume, we can determine the outward normal at any point through the right hand rule.

Computational Forms and Strategy

Just as we expanded Stokes' Theorem into component forms, we can make the Divergence Theorem more computationally accessible. Recall that for a vector field $\mathbf{F} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$:

$$\operatorname{div} \mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$$

Therefore, the Divergence Theorem becomes:

$$\oiint_S \mathbf{F} \cdot \mathbf{n} \, dS = \iiint_V \left(\frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} \right) dV$$

The surface integral on the left requires careful consideration of the outward unit normal $\mathbf{n}$. For different types of surfaces, we have:

1. **Level Surfaces:** For a surface $f(x, y, z) = c$:

$$\mathbf{n} = \frac{\nabla f}{\|\nabla f\|} = \frac{1}{\|\nabla f\|} \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right\rangle$$

2. **Parametric Surfaces:** For $\mathbf{r}(u, v) = \langle x(u, v), y(u, v), z(u, v) \rangle$:

$$\mathbf{n} = \frac{\mathbf{r}_u \times \mathbf{r}_v}{\|\mathbf{r}_u \times \mathbf{r}_v\|}$$

3. **Simple Surfaces:** For surfaces like $z = g(x, y)$:

$$\mathbf{n} = \frac{\langle -g_x, -g_y, 1 \rangle}{\sqrt{1 + g_x^2 + g_y^2}}$$

In different coordinate systems, the divergence takes special forms:

- **Cylindrical Coordinates** (r, θ, z) :

$$\operatorname{div} \mathbf{F} = \frac{1}{r} \frac{\partial}{\partial r} (r F_r) + \frac{1}{r} \frac{\partial F_\theta}{\partial \theta} + \frac{\partial F_z}{\partial z}$$

- **Spherical Coordinates** (r, ϕ, θ) :

$$\operatorname{div} \mathbf{F} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 F_r) + \frac{1}{r \sin \phi} \frac{\partial}{\partial \phi} (\sin \phi F_\phi) + \frac{1}{r \sin \phi} \frac{\partial F_\theta}{\partial \theta}$$

When choosing which form to use, consider:

- Symmetry of the vector field
- Shape of the boundary surface
- Complexity of the divergence versus the surface integral

For example, if the region is a sphere and the field has spherical symmetry, using spherical coordinates often simplifies the calculation dramatically. Similarly, for cylindrical bodies, cylindrical coordinates typically prove most efficient.

A systematic approach to these problems involves:

1. Identify the vector field components and the boundary surface
2. Choose appropriate coordinates based on symmetry
3. Determine whether surface or volume integral is simpler
4. If using surface integral:
 - Break surface into pieces if necessary
 - Calculate normal vectors for each piece
 - Set up parameterization if needed
5. If using volume integral:
 - Calculate divergence explicitly
 - Determine integration bounds
 - Consider coordinate transformations

This framework provides a practical approach to applying the Divergence Theorem, turning the abstract relationship between flux and divergence into concrete computational steps.

▲ **Example 5.7.1** Consider the vector field $\mathbf{F}(x, y, z) = (x^2y)\mathbf{i} + (y^2z)\mathbf{j} + (z^2x)\mathbf{k}$, and let V be the region bounded by the paraboloid $z = x^2 + y^2$ and the plane $z = 1$. Verify the Divergence Theorem by computing both:

1. The surface integral $\oint_S \mathbf{F} \cdot \mathbf{n} \, dS$
2. The volume integral $\iiint_V \operatorname{div} \mathbf{F} \, dV$

Solution

1. Surface Integral

The boundary S consists of two pieces:

- S_1 : The paraboloid $z = x^2 + y^2$ (bottom)
- S_2 : The plane $z = 1$ (top)

For S_1 (paraboloid):

- Writing as $f(x, y, z) = z - x^2 - y^2 = 0$, the gradient is $\nabla f = \langle -2x, -2y, 1 \rangle$
- Since our region V is above this surface, the outward normal must point downward:

$$\mathbf{n}_1 = \frac{\langle 2x, 2y, -1 \rangle}{\sqrt{4x^2 + 4y^2 + 1}}$$

- The flux integral becomes:

$$\begin{aligned} \iint_{S_1} \mathbf{F} \cdot \mathbf{n}_1 dS &= \iint_D \frac{2x^3y + 2y^3z - z^2x}{\sqrt{4x^2 + 4y^2 + 1}} \sqrt{4x^2 + 4y^2 + 1} dA \\ &= \iint_D (2x^3y + 2y^3(x^2 + y^2) - x(x^2 + y^2)^2) dA \end{aligned}$$

where D is the projection onto the xy -plane bounded by $x^2 + y^2 \leq 1$

For S_2 (plane):

- Since our region V is below this surface, the outward normal is:

$$\mathbf{n}_2 = -\mathbf{k}$$

- The flux integral is:

$$\iint_{S_2} \mathbf{F} \cdot \mathbf{n}_2 dS = - \iint_D z^2 x|_{z=1} dA = - \iint_D x dA$$

Converting both integrals to polar coordinates:

- $x = r \cos \theta$, $y = r \sin \theta$
- For S_1 : $z = r^2$
- For S_2 : $z = 1$
- Bounds: $0 \leq r \leq 1$, $0 \leq \theta \leq 2\pi$

$$\begin{aligned} \iint_{S_1} \mathbf{F} \cdot \mathbf{n}_1 dS &= \int_0^{2\pi} \int_0^1 (-2r^4 \cos^3 \theta \sin \theta - 2r^7 \sin^3 \theta + r^5 \cos \theta) r dr d\theta \\ &= 0 \text{ (odd functions over } [0, 2\pi]) \end{aligned}$$

$$\iint_{S_2} \mathbf{F} \cdot \mathbf{n}_2 dS = \int_0^{2\pi} \int_0^1 r \cos \theta r dr d\theta = 0$$

Therefore, the total surface integral is zero.

2. Volume Integral

First, calculate $\operatorname{div} \mathbf{F}$:

$$\begin{aligned}\operatorname{div} \mathbf{F} &= \frac{\partial}{\partial x}(x^2y) + \frac{\partial}{\partial y}(y^2z) + \frac{\partial}{\partial z}(z^2x) \\ &= 2xy + 2yz + 2zx\end{aligned}$$

Setting up the volume integral in cylindrical coordinates:

- $x = r \cos \theta$, $y = r \sin \theta$
- Lower bound: $z = r^2$
- Upper bound: $z = 1$
- Radial bound: $0 \leq r \leq 1$

$$\begin{aligned}\iiint_V \operatorname{div} \mathbf{F} dV &= \int_0^{2\pi} \int_0^1 \int_{r^2}^1 (2r^2 \cos \theta \sin \theta + 2rz \sin \theta + 2rz \cos \theta) r dz dr d\theta \\ &= \int_0^{2\pi} \int_0^1 \int_{r^2}^1 2r(r \sin \theta \cos \theta + z(\sin \theta + \cos \theta)) dz dr d\theta \\ &= 0 \text{ (by symmetry)}\end{aligned}$$

Therefore, both sides of the Divergence Theorem equal zero, confirming the theorem's validity for this case.

Physical Interpretation: The fact that both integrals equal zero suggests that any flow entering the region through one part of the boundary must exit through another part, with no net accumulation or depletion within the volume. This is analogous to an incompressible fluid flow, where the total inflow must equal the total outflow.

Physical Applications

The Divergence Theorem finds its deepest expression in physical conservation laws, where it bridges local field behavior with global measurements of flow and flux. At its heart lies a profound connection: what flows through a boundary must be created or destroyed within. This principle manifests throughout physics in increasingly sophisticated ways.

Consider first the flow of mass in a fluid. At each point in space and time, we have a density $\rho(x, y, z, t)$ and a velocity field $\mathbf{v}$ that tells us how the fluid moves. The product $\rho\mathbf{v}$ represents mass flux—the flow of mass through space. Conservation of mass demands that any change in the total mass within a volume must precisely balance the flow through its boundary:

$$\frac{d}{dt} \iiint_V \rho dV = - \oiint_S \rho \mathbf{v} \cdot \mathbf{n} dS$$

The Divergence Theorem transforms this global statement into a local one. Bringing the time derivative inside the volume integral and applying the theorem to the right side:

$$\iiint_V \frac{\partial \rho}{\partial t} dV = - \iiint_V \operatorname{div}(\rho \mathbf{v}) dV$$

Since this equality must hold for any volume, we obtain the continuity equation:

$$\frac{\partial \rho}{\partial t} + \operatorname{div}(\rho \mathbf{v}) = 0$$

For incompressible fluids where density remains constant, this reduces to $\operatorname{div} \mathbf{v} = 0$, expressing that the velocity field has no sources or sinks.

This pattern extends beautifully to electromagnetism. Electric charge serves as the source of electromagnetic fields, and its conservation takes a similar mathematical form. If $\mathbf{J}$ represents current density—the flow of charge through space—and ρ now represents charge density, conservation of charge requires:

$$\operatorname{div} \mathbf{J} = -\frac{\partial \rho}{\partial t}$$

This equation tells us something profound: changes in charge distribution necessitate current flow. Moreover, when combined with Maxwell's equations, particularly Gauss's law $\operatorname{div} \mathbf{E} = \rho/\epsilon_0$, it leads to the prediction of electromagnetic waves. The interplay between changing charge distributions and current flow creates ripples in the electromagnetic field that propagate through space.

Heat transfer provides yet another manifestation of these principles. When heat flows through a material, the heat flux $\mathbf{q}$ relates to temperature changes through:

$$\operatorname{div} \mathbf{q} = -c\rho \frac{\partial T}{\partial t}$$

where c represents specific heat capacity and $T(x, y, z, t)$ is the temperature field. This equation encodes Fourier's insight that heat flows from hot to cold regions, with the rate of temperature change determined by material properties. The divergence of heat flux directly drives local temperature evolution.

These seemingly distinct phenomena share a deep mathematical structure. For any conserved quantity ψ with associated flux $\mathbf{F}$, conservation takes the form:

$$\frac{\partial \psi}{\partial t} + \operatorname{div} \mathbf{F} = 0$$

Integrating over a volume and applying the Divergence Theorem reveals the universal pattern:

$$\frac{d}{dt} \iiint_V \psi \, dV = - \oiint_S \mathbf{F} \cdot \mathbf{n} \, dS$$

This structure appears in momentum conservation, where stress tensors describe force transmission through materials. It governs energy conservation in fields, where Poynting vectors track energy flow through space. Even quantum mechanics obeys this pattern—probability current ensures that quantum particles are neither created nor destroyed during evolution.

The Divergence Theorem thus emerges as one of physics' most profound results. It provides the mathematical framework for expressing nature's conservation principles, connecting microscopic behavior to macroscopic measurements. Whether describing fluid flow, electromagnetic fields, heat transfer, or quantum systems, the theorem reveals how local changes in fields relate to global flows across boundaries—a fundamental unity underlying diverse physical phenomena.

▲ **Example 5.7.2 Final Boss** A solid sphere of radius R has density given by $\rho(r) = \rho_0 \left(1 - \frac{r^2}{R^2}\right)$, where ρ_0 is a constant and r is the distance from the center. Using the Divergence Theorem, prove that:

1. The gravitational field inside the sphere at distance r from the center is $\mathbf{g}(r) = -\frac{4\pi G\rho_0}{3} \left(1 - \frac{3r^2}{5R^2}\right) r\hat{\mathbf{r}}$
2. The gravitational potential $\Phi(r)$ satisfies Poisson's equation: $\nabla^2\Phi = 4\pi G\rho$
3. The gravitational field outside the sphere is identical to that of a point mass located at the center with mass equal to the total mass of the sphere

Hint: The gravitational potential Φ at a point $\mathbf{r}$ due to a mass distribution is given by:

$$\Phi(\mathbf{r}) = -G \iiint_V \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} dV'$$

The gravitational field is related to the potential by $\mathbf{g} = -\nabla\Phi$. The Divergence Theorem can be applied to gravitational fields through the relationship $\nabla \cdot \mathbf{g} = -4\pi G\rho$, which relates the divergence of the gravitational field to the mass density. For a spherically symmetric situation, this means $\oiint_{S_r} \mathbf{g} \cdot \mathbf{n} \, dS = g(r) \cdot 4\pi r^2 = -4\pi G M_{\text{enclosed}}$, where M_{enclosed} is the mass contained within radius r .

Solution We'll solve this problem by applying the Divergence Theorem to the gravitational field, revealing fundamental properties of gravity.

1. Gravitational Field Inside the Sphere

First, note that the gravitational field $\mathbf{g}$ is related to the gravitational potential Φ by:

$$\mathbf{g} = -\nabla\Phi$$

By Newton's law of gravitation and the principle of superposition, the potential at a point $\mathbf{r}$ due to a mass distribution is:

$$\Phi(\mathbf{r}) = -G \iiint_V \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} dV'$$

For a spherically symmetric density distribution, the field must be radial: $\mathbf{g} = g(r)\hat{\mathbf{r}}$. To determine $g(r)$, we'll apply the Divergence Theorem to a sphere of radius $r < R$.

From Gauss's law for gravity:

$$\oiint_S \mathbf{g} \cdot \mathbf{n} dS = -4\pi G M_{enc}$$

where M_{enc} is the mass enclosed by the sphere of radius r . The left side is:

$$\oiint_S \mathbf{g} \cdot \mathbf{n} dS = \oiint_S g(r)\hat{\mathbf{r}} \cdot \hat{\mathbf{r}} dS = g(r) \cdot 4\pi r^2$$

The enclosed mass is:

$$\begin{aligned} M_{enc} &= \iiint_{r' \leq r} \rho(r') dV' \\ &= \int_0^{2\pi} \int_0^\pi \int_0^r \rho_0 \left(1 - \frac{r'^2}{R^2}\right) r'^2 \sin \theta dr' d\theta d\phi \\ &= 4\pi\rho_0 \int_0^r \left(1 - \frac{r'^2}{R^2}\right) r'^2 dr' \\ &= 4\pi\rho_0 \left(\frac{r^3}{3} - \frac{r^5}{5R^2}\right) \end{aligned}$$

Equating the surface integral with $-4\pi G M_{enc}$:

$$\begin{aligned} g(r) \cdot 4\pi r^2 &= -4\pi G \cdot 4\pi\rho_0 \left(\frac{r^3}{3} - \frac{r^5}{5R^2}\right) \\ g(r) &= -\frac{4\pi G\rho_0}{r^2} \left(\frac{r^3}{3} - \frac{r^5}{5R^2}\right) \\ &= -\frac{4\pi G\rho_0}{3} \left(1 - \frac{3r^2}{5R^2}\right) r \end{aligned}$$

Therefore:

$$\mathbf{g}(r) = -\frac{4\pi G\rho_0}{3} \left(1 - \frac{3r^2}{5R^2}\right) r\hat{\mathbf{r}} \quad \text{for } r \leq R$$

2. Verification of Poisson's Equation

Poisson's equation for gravity states:

$$\nabla^2\Phi = 4\pi G\rho$$

Since $\mathbf{g} = -\nabla\Phi$, we have $\nabla \cdot \mathbf{g} = -\nabla^2\Phi$. To verify Poisson's equation, we need to show that:

$$\nabla \cdot \mathbf{g} = -4\pi G\rho$$

For a radial vector field $\mathbf{g} = g(r)\hat{\mathbf{r}}$, the divergence in spherical coordinates is:

$$\nabla \cdot \mathbf{g} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 g(r))$$

Substituting $g(r) = -\frac{4\pi G\rho_0}{3} \left(1 - \frac{3r^2}{5R^2}\right) r$:

$$\begin{aligned} \nabla \cdot \mathbf{g} &= \frac{1}{r^2} \frac{\partial}{\partial r} \left[r^2 \cdot \left(-\frac{4\pi G\rho_0}{3} \left(1 - \frac{3r^2}{5R^2}\right) r \right) \right] \\ &= \frac{1}{r^2} \frac{\partial}{\partial r} \left[-\frac{4\pi G\rho_0}{3} r^3 \left(1 - \frac{3r^2}{5R^2}\right) \right] \\ &= \frac{1}{r^2} \frac{\partial}{\partial r} \left[-\frac{4\pi G\rho_0}{3} r^3 + \frac{4\pi G\rho_0}{5R^2} r^5 \right] \\ &= \frac{1}{r^2} \left[-\frac{4\pi G\rho_0}{3} \cdot 3r^2 + \frac{4\pi G\rho_0}{5R^2} \cdot 5r^4 \right] \\ &= -4\pi G\rho_0 + \frac{4\pi G\rho_0 r^2}{R^2} \\ &= -4\pi G\rho_0 \left(1 - \frac{r^2}{R^2}\right) \\ &= -4\pi G\rho(r) \end{aligned}$$

This confirms that the gravitational field satisfies Poisson's equation $\nabla^2\Phi = 4\pi G\rho$.

3. External Gravitational Field

For $r > R$, we'll again use the Divergence Theorem in the form of Gauss's law for gravity.

The total mass of the sphere is:

$$\begin{aligned}
 M_{total} &= \iiint_V \rho(r) dV \\
 &= 4\pi\rho_0 \int_0^R \left(1 - \frac{r^2}{R^2}\right) r^2 dr \\
 &= 4\pi\rho_0 \left[\frac{r^3}{3} - \frac{r^5}{5R^2} \right]_0^R \\
 &= 4\pi\rho_0 R^3 \left(\frac{1}{3} - \frac{1}{5} \right) \\
 &= 4\pi\rho_0 R^3 \cdot \frac{2}{15} = \frac{8\pi\rho_0 R^3}{15}
 \end{aligned}$$

By symmetry, the gravitational field outside the sphere must be radial. Applying Gauss's law to a sphere of radius $r > R$:

$$\oiint_S \mathbf{g} \cdot \mathbf{n} dS = -4\pi G M_{total}$$

The left side equals $g(r) \cdot 4\pi r^2$, so:

$$\begin{aligned}
 g(r) \cdot 4\pi r^2 &= -4\pi G \cdot \frac{8\pi\rho_0 R^3}{15} \\
 g(r) &= -\frac{GM_{total}}{r^2} = -G \cdot \frac{8\pi\rho_0 R^3}{15} \cdot \frac{1}{r^2}
 \end{aligned}$$

Therefore:

$$\mathbf{g}(r) = -G \cdot \frac{8\pi\rho_0 R^3}{15} \cdot \frac{\hat{\mathbf{r}}}{r^2} = -\frac{GM_{total}}{r^2} \hat{\mathbf{r}} \quad \text{for } r \geq R$$

This is identical to the gravitational field of a point mass M_{total} located at the center, confirming Newton's Shell Theorem: the gravitational field outside a spherically symmetric mass distribution is the same as if all the mass were concentrated at the center.

Formulas and Definitions

Important Vocabulary

- **Vector Field:** A function that assigns a vector to each point in space, written as $\mathbf{F}(x, y, z) = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$.
- **Conservative Field:** A vector field that can be written as the gradient of some scalar function f , where $\mathbf{F} = \nabla f$.
- **Curl:** A measure of rotation in a vector field, given by $\text{curl } \mathbf{F} = \nabla \times \mathbf{F}$.
- **Divergence:** A measure of the outward flux per unit volume in a vector field, given by $\text{div } \mathbf{F} = \nabla \cdot \mathbf{F}$.
- **Line Integral:** The integral of a vector field along a curve C , measuring quantities like work: $\int_C \mathbf{F} \cdot d\mathbf{r}$.
- **Surface Integral:** The integral of a vector field over a surface S , measuring flux: $\iint_S \mathbf{F} \cdot d\mathbf{S}$.
- **Flux:** The rate of flow of a vector field through a surface, measured by surface integrals.
- **Orientable Surface:** A surface where a consistent normal vector direction can be defined at each point.
- **Potential Function:** A scalar function f whose gradient gives a conservative vector field.
- **Path Independence:** Property of conservative fields where line integrals depend only on endpoints, not the path taken.
- **Source:** Point where field lines emerge (positive divergence).
- **Sink:** Point where field lines converge (negative divergence).

Key Formulas

Curl in 3D

$$\nabla \times \mathbf{F} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ P & Q & R \end{vmatrix}$$

Green's Theorem

$$\oint_C P dx + Q dy = \iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA$$

Divergence

$$\nabla \cdot \mathbf{F} = \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}$$

Stokes' Theorem

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S (\text{curl } \mathbf{F}) \cdot \mathbf{n} dS$$

Divergence Theorem

$$\oiint_S \mathbf{F} \cdot \mathbf{n} \, dS = \iiint_V \operatorname{div} \mathbf{F} \, dV$$

Generalized Stokes' Theorem

$$\int_{\Omega} d\omega = \int_{\partial\Omega} \omega$$

Conservative Field Test For a vector field $\mathbf{F}$ to be conservative:

$$\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}, \quad \frac{\partial P}{\partial z} = \frac{\partial R}{\partial x}, \quad \frac{\partial Q}{\partial z} = \frac{\partial R}{\partial y}$$

Surface Element

$$dS = \|\mathbf{r}_u \times \mathbf{r}_v\| \, du \, dv$$

Maxwell's Equations

$$\nabla \cdot \mathbf{E} = \frac{\rho}{\epsilon_0}$$

$$\nabla \cdot \mathbf{B} = 0$$

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$$

$$\nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t}$$

Operator	Cylindrical (r, θ, z)	Spherical (r, θ, ϕ)
∇f	$\frac{\partial f}{\partial r} \mathbf{e}_r + \frac{1}{r} \frac{\partial f}{\partial \theta} \mathbf{e}_\theta + \frac{\partial f}{\partial z} \mathbf{e}_z$	$\frac{\partial f}{\partial r} \mathbf{e}_r + \frac{1}{r} \frac{\partial f}{\partial \theta} \mathbf{e}_\theta + \frac{1}{r \sin \theta} \frac{\partial f}{\partial \phi} \mathbf{e}_\phi$
$\nabla \cdot \mathbf{F}$	$\frac{1}{r} \frac{\partial}{\partial r}(r F_r) + \frac{1}{r} \frac{\partial F_\theta}{\partial \theta} + \frac{\partial F_z}{\partial z}$	$\frac{1}{r^2} \frac{\partial}{\partial r}(r^2 F_r) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta}(\sin \theta F_\theta) + \frac{1}{r \sin \theta} \frac{\partial F_\phi}{\partial \phi}$
$\nabla \times \mathbf{F}$	$\begin{pmatrix} \frac{1}{r} \frac{\partial F_z}{\partial \theta} - \frac{\partial F_\theta}{\partial z} \\ \frac{\partial F_r}{\partial z} - \frac{\partial F_z}{\partial r} \\ \frac{1}{r} \frac{\partial}{\partial r}(r F_\theta) - \frac{1}{r} \frac{\partial F_r}{\partial \theta} \end{pmatrix}$	$\begin{pmatrix} \frac{1}{r \sin \theta} \left(\frac{\partial}{\partial \theta}(\sin \theta F_\phi) - \frac{\partial F_\theta}{\partial \phi} \right) \\ \frac{1}{r} \left(\frac{1}{\sin \theta} \frac{\partial F_r}{\partial \phi} - \frac{\partial}{\partial r}(r F_\phi) \right) \\ \frac{1}{r} \left(\frac{\partial}{\partial r}(r F_\theta) - \frac{\partial F_r}{\partial \theta} \right) \end{pmatrix}$

Practice Problems

1. Consider the 2D vector field

$$\mathbf{F}(x, y) = (x + 1)\mathbf{i} + (y - 2)\mathbf{j}.$$

- (a) Evaluate $\mathbf{F}(1, -1)$.
 (b) Sketch a few representative vectors near the origin to see how this field behaves.
2. Explain the difference between a *scalar field* and a *vector field*. Give one concrete physical example of each (e.g., temperature distribution for a scalar field, velocity flow for a vector field).

3. Given

$$\mathbf{F}(x, y) = -y\mathbf{i} + x\mathbf{j},$$

describe in words how you would plot the arrows at the points $(2, 0)$, $(1, 1)$, and $(-2, -2)$. What does the overall pattern of the field suggest?

4. Consider the vector field

$$\mathbf{F}(x, y) = (e^x, e^y).$$

- (a) Evaluate $\mathbf{F}(0, 0)$ and $\mathbf{F}(1, 2)$.
 (b) Based on these vectors, does the origin appear to be a source, a sink, or neither? Give a brief reason.
5. A 3D vector field is given by
- $$\mathbf{G}(x, y, z) = \frac{x}{x^2 + y^2 + z^2} \mathbf{i} + \frac{y}{x^2 + y^2 + z^2} \mathbf{j} + \frac{z}{x^2 + y^2 + z^2} \mathbf{k}.$$
- Is $\mathbf{G}$ continuous everywhere in $\mathbb{R}^3$? If not, identify any problematic points.

6. Consider the 3D vector field

$$\mathbf{H}(x, y, z) = x^2 \mathbf{i} + y^2 \mathbf{j} + z^2 \mathbf{k}.$$

- (a) Determine whether $\mathbf{H}$ is continuous at all points in $\mathbb{R}^3$.
 (b) If there are any problematic points, identify them and explain why.
7. Explain briefly why engineers or physicists might choose to visualize a vector field with all arrows having the same length (i.e., *normalized*). Give one advantage and one disadvantage of this approach, referencing physical interpretations.
8. Classify the origin for each of the following vector fields:

(a) $\mathbf{F}_1(x, y) = x\mathbf{i} + y\mathbf{j}$

(b) $\mathbf{F}_2(x, y) = -x\mathbf{i} - y\mathbf{j}$

(c) $\mathbf{F}_3(x, y) = x\mathbf{i} - y\mathbf{j}$

State which is a source, which is a sink, and which is a saddle.

9. Show that the 2D vector field

$$\mathbf{F}(x, y) = y\mathbf{i} - x\mathbf{j}$$

is *not* conservative by using the partial-derivative condition $\frac{\partial P}{\partial y} \neq \frac{\partial Q}{\partial x}$. In one sentence, relate this to the “rotational” or “circulatory” pattern you see near the origin.

10. Let

$$\mathbf{F}(x, y) = x^3 \mathbf{i} + 3x^2 y \mathbf{j}.$$

- (a) Use the condition $\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$ to determine if $\mathbf{F}$ is conservative.
 (b) If it *is* conservative, find a potential function $f(x, y)$ such that $\nabla f = \mathbf{F}$.

11. Consider the 3D field

$$\mathbf{F}(x, y, z) = 2xy\mathbf{i} + (x^2 + z^2)\mathbf{j} + 2yz\mathbf{k}.$$

Test whether $\mathbf{F}$ is conservative by checking the equality of appropriate mixed partial derivatives in $\mathbb{R}^3$.

12. For the 2D field

$$\mathbf{F}(x, y) = \sqrt{x^2 + y^2}\mathbf{i} - (\sin(xy))\mathbf{j},$$

compute:

$$\frac{\partial P}{\partial x}, \quad \frac{\partial P}{\partial y}, \quad \frac{\partial Q}{\partial x}, \quad \frac{\partial Q}{\partial y}.$$

Which (if any) of these partial derivatives are undefined at the origin?

13. Consider the 2D vector field

$$\mathbf{F}(x, y) = (x^2 + y^2)\mathbf{i} - (x^2 + y^2)\mathbf{j}.$$

- (a) Check whether $\mathbf{F}$ is conservative by comparing $\frac{\partial P}{\partial y}$ and $\frac{\partial Q}{\partial x}$.
- (b) If it *is* conservative, find a potential function. If not, classify the origin as a source, sink, or saddle.

14. Compute the scalar line integral

$$\int_C (x + y) ds$$

where C is the line segment from $(0, 0)$ to $(2, 4)$. Parametrize C as $\mathbf{r}(t)$, $0 \leq t \leq 1$, and set up the integral explicitly.

15. Evaluate

$$\int_C (x^2 + y) ds$$

where C is the parabola $y = x^2$ from $(0, 0)$ to $(2, 4)$. Use the standard parameterization $\mathbf{r}(t) = \langle t, t^2 \rangle$.

16. Show that the line integral

$$\int_C f(x, y) ds$$

does not depend on the *direction* of travel for a scalar function f . In other words, if you reverse the parameterization, explain why the sign of the integral does not change.

17. Compute

$$\int_C \sqrt{x^2 + y^2} ds$$

where C is the quarter circle $x = r \cos t$, $y = r \sin t$ for $t \in [0, \frac{\pi}{2}]$. Show your parameterization and simplify.

18. Evaluate

$$\int_C (2x + 3y) ds$$

where C is the piecewise path going first along the x -axis from $(0, 0)$ to $(3, 0)$, then vertically to $(3, 2)$. Break the integral into two segments and sum the results.

19. Compute the line integral of the vector field

$$\mathbf{F}(x, y) = (y, x)$$

along the line segment C from $(0, 0)$ to $(2, 3)$. Parametrize C and evaluate

$$\int_C \mathbf{F} \cdot d\mathbf{r}.$$

20. Let $\mathbf{F}(x, y) = (x - y, x^2)$. If C is the rectangle with corners $(0, 0)$, $(2, 0)$, $(2, 1)$, $(0, 1)$ traversed *counterclockwise*, find $\int_C \mathbf{F} \cdot d\mathbf{r}$.

21. Parameterize and evaluate the line integral

$$\int_C \mathbf{F} \cdot d\mathbf{r}$$

- for $\mathbf{F}(x, y) = (\ln(1 + y^2), e^x)$ around the circle $x^2 + y^2 = 4$ *counterclockwise*. How does reversing to clockwise change the sign?
22. Explain how to compute $\int_C \mathbf{F} \cdot d\mathbf{r}$ if C is *piecewise* with three smooth segments C_1 , C_2 , and C_3 . Write a formula involving $\int_{C_1}$, $\int_{C_2}$, and $\int_{C_3}$, and give one reason why this strategy is often used in applications.
23. Show that the line integral
- $$\oint_C (3y, -2x) \cdot d\mathbf{r}$$
- around any closed loop C depends on the *orientation* of C . If C is traversed clockwise, how does the result compare to counterclockwise?
24. Let C be the curve from $(1, 0)$ to $(1, 2)$ along the line $x = 1$. Compute
- $$\int_C \mathbf{F} \cdot d\mathbf{r} \quad \text{for} \quad \mathbf{F}(x, y) = (x+1, 2xy-1).$$
- Explain why reversing the path gives a negative of your result.
25. Determine whether the 2D field $\mathbf{F}(x, y) = (y, x)$ is *conservative* on $\mathbb{R}^2$. If it were conservative, there would exist a potential function f . Check the partial derivatives condition to confirm or deny.
26. On $\mathbb{R}^2 \setminus \{(0, 0)\}$, consider
- $$\mathbf{F}(x, y) = (-y/(x^2 + y^2), x/(x^2 + y^2)).$$
- Parametrize the circle $x^2 + y^2 = r^2$ and compute $\oint_C \mathbf{F} \cdot d\mathbf{r}$. Explain why this field fails to be conservative.
27. Let $\mathbf{G}(x, y) = (x + \sin y, \cos x + y)$. Assume the domain is all of $\mathbb{R}^2$. Check if $\mathbf{G}$ is conservative by verifying the mixed partial derivatives. If it is, find a potential function $g(x, y)$ and use it to compute
- $$\int_C \mathbf{G} \cdot d\mathbf{r}$$
- along any path from $(0, 0)$ to $(2, 1)$.
28. In 3D, let
- $$\mathbf{F}(x, y, z) = (yz, zx, xy).$$
- Consider the piecewise path that goes:
- $$(0, 0, 0) \xrightarrow{C_1} (1, 0, 0) \xrightarrow{C_2} (1, 2, 0) \xrightarrow{C_3} (1, 2, 3).$$
- Compute $\int_{C_1} + \int_{C_2} + \int_{C_3}$ for $\mathbf{F} \cdot d\mathbf{r}$. Does this field appear to be conservative on $\mathbb{R}^3$? Why or why not?
29. State Green's Theorem in your own words and explain conceptually why the line integral around a closed curve in the plane can be expressed as a double integral over the region it encloses.
30. Compute the line integral
- $$\oint_C (xy) dx + (x^2) dy$$
- where C is the positively oriented (counterclockwise) boundary of the rectangle with corners $(0, 0)$, $(2, 0)$, $(2, 1)$, and $(0, 1)$. First do it directly by parameterizing each edge, then use Green's Theorem to confirm your result.
31. Let $\mathbf{F}(x, y) = (P, Q) = (-y, x)$. Using Green's Theorem, evaluate
- $$\oint_C \mathbf{F} \cdot d\mathbf{r}$$
- where C is the unit circle $x^2 + y^2 = 1$ oriented counterclockwise. Interpret physically what this integral measures (circulation) and explain why it makes sense.

32. Use Green's Theorem to transform the double integral

$$\iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA$$

into a line integral for the region D bounded by $y = x^2$ and $y = 4$. Write down the resulting line integral explicitly (no need to solve) and mention one advantage of switching to the line integral form.

33. Explain how interior edges cancel when you partition a region D into many small rectangles and sum up the individual line integrals around each subrectangle. Why do only the outer boundary edges survive in the sum?

34. Compute

$$\oint_C (x^2 y^2) dx + (xy^3) dy$$

around the boundary of the triangle with vertices $(0, 0)$, $(2, 0)$, $(0, 2)$ oriented counterclockwise. Use Green's Theorem to avoid evaluating three separate line integrals.

35. Let $\mathbf{F}(x, y) = (y, -x)$. Show that

$$\oint_C \mathbf{F} \cdot d\mathbf{r}$$

around any positively oriented, simple closed curve C is proportional to the area of the region enclosed. (Hint: Use Green's Theorem and notice $\partial Q/\partial x - \partial P/\partial y$ is a constant.)

36. Suppose $\mathbf{F}(x, y) = (e^y, \sin x)$. Use Green's Theorem to evaluate

$$\oint_C \mathbf{F} \cdot d\mathbf{r}$$

where C is the ellipse $\frac{x^2}{4} + \frac{y^2}{9} = 1$, traversed counterclockwise. (Hint:

Parametric or direct line-integration is messy, so let Green's Theorem and a suitable double integral do the work.)

37. Explain how reversing the orientation of a closed curve C (from counterclockwise to clockwise) affects the sign of $\oint_C P dx + Q dy$. Relate this to the expression $\iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA$.

38. Consider the domain D in $\mathbb{R}^2 \setminus \{(0, 0)\}$ (the plane without the origin). Let

$$\mathbf{F}(x, y) = (P, Q) = \left(\frac{-y}{x^2 + y^2}, \frac{x}{x^2 + y^2} \right).$$

Apply Green's Theorem to the boundary of a circle of radius r centered at $(0, 0)$. Show why the usual condition "domain must be simply connected" is crucial here, and compute $\oint_C \mathbf{F} \cdot d\mathbf{r}$ for the circle of radius r .

39. Given the 2D vector field $\mathbf{F}(x, y) = (P, Q) = (x + 2y, x - y)$, compute:

$$\nabla \cdot \mathbf{F}, \quad \nabla \times \mathbf{F} \text{ (in 2D, i.e. } \partial Q/\partial x - \partial P/\partial y \text{)}.$$

Which one indicates expansion/compression, and which indicates rotation?

40. For $\mathbf{F}(x, y, z) = (x^2, 2y, -3z)$, compute:

$$\nabla \cdot \mathbf{F} \quad \text{and} \quad \nabla \times \mathbf{F}.$$

Is $\mathbf{F}$ incompressible (i.e. divergence-free)? Does it have nonzero curl?

41. Explain briefly why every conservative vector field $\mathbf{F} = \nabla f$ has $\nabla \times \mathbf{F} = \mathbf{0}$. (Use Clairaut's theorem on equality of mixed partial derivatives.)

42. Check whether the 3D field $\mathbf{F}(x, y, z) = (e^x \sin y, e^y \sin z, e^z \sin x)$ is *divergence-free*. (Compute $\nabla \cdot \mathbf{F}$ and see if it's zero everywhere.)

43. Let $\mathbf{F}(x, y) = (x^3, y^3)$ in $\mathbb{R}^2$. Show that $\nabla \cdot \mathbf{F} \neq 0$ everywhere except possibly at special points. Give a brief physical interpretation in terms of “sources” or “sinks.”

44. Classify each of these 2D vector fields at the origin as primarily *source*, *sink*, or *rotation* by checking $\nabla \cdot \mathbf{F}$ and $\nabla \times \mathbf{F}$ at $(0, 0)$:

- (a) $\mathbf{F}_1 = (x, -y)$
- (b) $\mathbf{F}_2 = (2x, 2y)$
- (c) $\mathbf{F}_3 = (-y, x)$.

45. Use the identity $\nabla \cdot (\nabla \times \mathbf{F}) = 0$ to show that no purely rotational flow in 3D can create or destroy mass. Relate this to a fluid flow perspective (no net outflow in a purely swirling motion).

46. For the 3D field $\mathbf{F}(x, y, z) = (yz, zx, xy)$, compute:

$$\nabla \cdot \mathbf{F}, \quad \nabla \times \mathbf{F}.$$

Which is zero, and why might this be physically interpreted as “no net creation of fluid but nonzero rotation”?

47. Consider the velocity field in cylindrical coordinates

$$\mathbf{v}(r, \theta, z) = (0, \alpha r, 0),$$

where α is a constant, and $\mathbf{v}$ is expressed as (v_r, v_θ, v_z) . Compute $\nabla \cdot \mathbf{v}$ and $\nabla \times \mathbf{v}$ in *cylindrical* form. Is this flow incompressible? Does it rotate about the z -axis?

48. In spherical coordinates (r, θ, ϕ) , a radial field is given by

$$\mathbf{F}(r, \theta, \phi) = (f(r), 0, 0)$$

(i.e. purely in the $\mathbf{e}_r$ direction). Compute $\nabla \cdot \mathbf{F}$ using the divergence expression in spherical coordinates. Under

what condition on $f(r)$ is this divergence zero for $r > 0$?

49. Verify the product rule for divergence:

$$\nabla \cdot (f \mathbf{F}) = f(\nabla \cdot \mathbf{F}) + \mathbf{F} \cdot (\nabla f)$$

by expanding both sides in Cartesian coordinates. In physics, explain how this identity shows that the net outflow of $f\mathbf{F}$ splits into two contributions: one from $\mathbf{F}$ alone and one from gradient of f .

50. Show that for any scalar function $f(x, y, z)$,

$$\nabla \times (\nabla f) = \mathbf{0}.$$

Discuss why this implies any gradient field in 3D is irrotational. Give one physical application (e.g., electrostatic potential).

51. A vector field $\mathbf{F}(x, y, z)$ has $\nabla \cdot \mathbf{F} = 3$ everywhere in $\mathbb{R}^3$. Explain briefly why this means the field behaves like a uniform “source of fluid.” How might you verify this statement by integrating the flux over a sphere of radius R ?

52. Compute

$$\nabla \times (\nabla \times \mathbf{F})$$

for $\mathbf{F} = (x^2, y^2, z^2)$, and confirm it matches the identity

$$\nabla(\nabla \cdot \mathbf{F}) - \nabla^2 \mathbf{F}.$$

53. Consider the 2D vector field

$$\mathbf{F}(x, y) = \nabla(x^2 + 2y^2).$$

- (a) Write out $\mathbf{F}$ explicitly in terms of P and Q .
- (b) Compute $\nabla \times \mathbf{F}$ (in 2D sense) and $\nabla \cdot \mathbf{F}$.

- (c) Does the result confirm that gradients are always irrotational?
54. Convert the scalar surface
- $$x^2 + y^2 = 4z^2$$
- into a *parametric* form. Indicate any parameter constraints (e.g. domain ranges).
55. Use the *trace method* to parametrize the surface
- $$x = y^2 + 2z^2$$
- but only for the region where $x \leq 4$. Specify the appropriate domain for the parameters you introduce.
56. Consider the surface
- $$\mathbf{r}(u, v) = (u, v, \sqrt{u^2 + v^2}).$$
- (a) Compute the vectors $\mathbf{r}_u$ and $\mathbf{r}_v$.
- (b) Find the normal vector $\mathbf{n} = \mathbf{r}_u \times \mathbf{r}_v$.
- (c) Determine $\|\mathbf{r}_u \times \mathbf{r}_v\|$.
57. Find the *surface area* of the portion of the paraboloid $z = 9 - x^2 - y^2$ above the region $x^2 + y^2 \leq 5$. Write the integral *without evaluating*.
58. A metal cooling fin is shaped like $z = f(x, y) = x^2 + 4y^2$, restricted to $-1 \leq x \leq 1$, $-1 \leq y \leq 1$. Its surface temperature at (x, y, z) is
- $$T(x, y, z) = \sqrt{1 + x^2 + y^2}.$$
- Set up (but do not solve) the surface integral
- $$\iint_S T \, dS$$
- to find the *total heat* over the fin. Use the standard formula for surfaces $z = f(x, y)$.
59. Let $\mathbf{F}(x, y, z) = (x^2, 0, yz)$.
- (a) Parameterize the part of the plane $z = 2x$ that occupies the square region $\{(x, y) \mid 0 \leq x \leq 2, 0 \leq y \leq 1\}$ in the xy -plane.
- (b) Set up the flux integral $\iint_S \mathbf{F} \cdot d\mathbf{S}$ over that portion of the plane. (No need to evaluate fully.)
60. A surface in cylindrical coordinates is given by
- $$\mathbf{r}(\theta, z) = (2 \cos \theta, 2 \sin \theta, z),$$
- for $0 \leq \theta \leq 2\pi$, $0 \leq z \leq 3$. Suppose a *scalar* function is
- $$f(x, y, z) = \sqrt{x^2 + y^2} = 2.$$
- Compute the surface integral
- $$\iint_S f \, dS$$
- for the *curved* surface of the cylinder. (This is effectively the lateral area times f .)
61. Consider the hemisphere $x^2 + y^2 + z^2 = 9$ with $z \geq 0$. A density function $\rho(x, y, z)$ equals z . Write a suitable parameterization using *spherical coordinates* and set up *both*:
- (a) the integral for total mass $\iint_S \rho \, dS$;
- (b) the integral for the total outward flux of $\mathbf{F} = (0, 0, z)$ through this hemisphere (oriented outward).
62. Let $\mathbf{F}(x, y, z) = (P, Q, R) = (2x, 3y, -z)$. Suppose S is the surface of the graph $z = g(x, y) = x + 2y$, above the region $\Omega = \{0 \leq x \leq 1, 0 \leq y \leq 2\}$. Assume upward orientation.
- (a) Compute the partial derivatives g_x, g_y and write down the normal vector $\mathbf{n} = (-g_x, -g_y, 1)$.

- (b) Express the flux integral

$$\iint_S \mathbf{F} \cdot d\mathbf{S}$$

as a double integral $\iint_{\Omega}(\cdots) dA$.
(No need to solve numerically.)

63. A closed surface S is formed by the cylinder $x^2 + y^2 = 4$ for $-1 \leq z \leq 2$ *plus* the top disk at $z = 2$ *plus* the bottom disk at $z = -1$. A fluid velocity field is

$$\mathbf{v}(x, y, z) = (x, y, -1).$$

Use the outward orientation convention:

- Parameterize the curved side surface $\mathbf{r}(\theta, z)$ (with normal outward from the cylinder).
 - Write integrals for the *flux* through the curved side, top disk, and bottom disk.
 - Indicate how you would sum them for total outward flux.
64. Let $\mathbf{F} = (y^2, z^2, x^2)$. Let Σ be the portion of the paraboloid $z = x^2 + y^2$ cut off by $z = 4$. Its boundary C is thus the circle $x^2 + y^2 = 4$ at $z = 4$. Use Stokes' Theorem to convert

$$\oint_C \mathbf{F} \cdot d\mathbf{r}$$

into a surface integral of $\nabla \times \mathbf{F}$ over Σ . Set up the integral, but do not compute fully.

65. A vector field is given by

$$\mathbf{F}(x, y, z) = (e^y, \ln(1 + z^2), y^2 z).$$

The surface Σ is the planar patch $x + 2y + 3z = 6$ in the first octant, bounded

by the triangle C . Decide if it's easier to compute $\oint_C \mathbf{F} \cdot d\mathbf{r}$ directly, or to use $\iint_{\Sigma} (\nabla \times \mathbf{F}) \cdot d\mathbf{S}$. Briefly justify your choice and outline the steps (no numeric solution needed).

66. The surface Σ is the part of the cylinder $x^2 + y^2 = 9$ with $-2 \leq z \leq 5$. The vector field is

$$\mathbf{F} = (-y, x, 0).$$

- Find $\nabla \times \mathbf{F}$.
 - The boundary $\partial\Sigma$ consists of two circles. Write a parametric equation for each circle with consistent orientations (use right-hand rule).
 - Use Stokes' Theorem to transform $\oint_{\partial\Sigma} \mathbf{F} \cdot d\mathbf{r}$ into the surface integral of $\nabla \times \mathbf{F}$ over Σ . (No need to evaluate fully.)
67. Suppose $\mathbf{F} = \nabla f$ for some scalar f . Show that for any smooth, oriented surface Σ with boundary C , we get

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 0$$

by appealing to Stokes' Theorem. (Hint: use the fact that $\nabla \times (\nabla f) = \mathbf{0}$.)

68. Let $\mathbf{E}(x, y, z)$ be the electric field, and let $\mathbf{B}(x, y, z)$ be the magnetic field. A simplified statement of Faraday's Law (static boundary) says:

$$\oint_C \mathbf{E} \cdot d\mathbf{r} = -\frac{d}{dt} \iint_{\Sigma} \mathbf{B} \cdot d\mathbf{S}$$

Explain briefly how this is an instance of Stokes' Theorem, identifying $\mathbf{E}$ with the line-integrated field and $\nabla \times \mathbf{E}$ with the flux integrand. What is the physical meaning of the negative time derivative on the right side?

69. Consider the vector field

$$\mathbf{F}(x, y, z) = (P, Q, R) = (xy, yz, xz).$$

Let V be the solid region bounded by the sphere $x^2 + y^2 + z^2 = 4$. Use the Divergence Theorem to convert the surface integral

$$\oiint_{\partial V} \mathbf{F} \cdot \mathbf{n} \, dS$$

into a volume integral, $\iiint_V \nabla \cdot \mathbf{F} \, dV$. Write the integral *without* evaluating.

70. A cone in the first octant is given by $0 \leq z \leq \sqrt{x^2 + y^2}$, capped by the plane $z = 2$. Let

$$\mathbf{F}(x, y, z) = (x^2, y^2, z^2).$$

- (a) Identify the boundary surfaces (sides of the cone and the plane at $z = 2$).
- (b) Write down a formula for $\oiint_{\partial V} \mathbf{F} \cdot d\mathbf{S}$ using the Divergence Theorem.

71. For the region V bounded by the cylinder $x^2 + y^2 = 1$ and the planes $z = 0$ and $z = 5$, consider

$$\mathbf{F} = (e^z, \sin(xy), 2z).$$

Decide whether you would rather compute $\oiint_{\partial V} \mathbf{F} \cdot \mathbf{n} \, dS$ directly over the curved plus top/bottom surfaces, *or* via $\iiint_V \nabla \cdot \mathbf{F} \, dV$. Give a short reason and outline the steps.

72. Let $\mathbf{F}(x, y, z) = (x^2 + y^2, z, 3x)$. Suppose V is the ball $x^2 + y^2 + z^2 \leq 4$. Use spherical coordinates to write a triple integral for $\iiint_V \nabla \cdot \mathbf{F} \, dV$. No need to simplify fully.

73. Explain, in words, why the *net* outward flux of an *incompressible* fluid (divergence zero) through any closed surface must be zero. How does this align with the Divergence Theorem statement $\oiint_{\partial V} \mathbf{v} \cdot d\mathbf{S} = 0$ when $\nabla \cdot \mathbf{v} = 0$?

Chapter 6

Differential Equations

Introduction

In our mathematical journey thus far, we have explored the foundations of calculus, mastered the techniques of integration and differentiation, and ventured into the multidimensional realms of vector calculus. Now, we stand at the threshold of differential equations, where these fundamental concepts converge to describe the mathematical language of change itself. Differential equations represent not just static relationships between variables, but the dynamic evolution of systems over time and space—a natural progression in our pursuit of mathematical generalization.

The tools we have cultivated throughout our studies—from basic derivatives to complex integration techniques—now serve as essential instruments for understanding how quantities evolve under various conditions. Our facility with derivatives transforms into the ability to decode equations that describe rates of change, while our integration skills become crucial in finding solutions that tell us how systems behave over time. The analytical methods we developed for solving algebraic equations evolve into sophisticated techniques for unraveling both ordinary and partial differential equations.

This chapter introduces the fundamental concepts and methods that bridge the gap between static mathematical descriptions and dynamic processes. From first-order ordinary differential equations that model simple growth and decay, to partial differential equations that describe heat flow and wave propagation, we will explore how these powerful mathematical tools help us understand and predict complex phenomena. As we progress, we will see how differential equations provide a unified framework for describing diverse physical systems, from the simple harmonic motion of a pendulum to the intricate patterns of fluid dynamics. The journey ahead promises to be both challenging and enlightening, as we develop the ability to analyze and describe the fundamental laws that govern change in our universe.

Lecture 6.1: Introduction to Differential Equations

After exploring the rich landscape of multivariable calculus—from vectors to gradients, from multiple integrals to vector fields—we now turn our attention to one of mathematics’ most powerful modeling tools: differential equations. While our previous studies focused on understanding rates of change and accumulation in multiple dimensions, we now ask a more fundamental question: given information about how quantities change, can we determine the quantities themselves?

This seemingly simple question opens the door to modeling countless real-world phenomena, from the graceful arc of a pendulum to the intricate patterns of population growth, from the flow of heat through materials to the propagation of waves through space. In each case, we begin not with the behavior itself, but with a mathematical description of how that behavior changes—a differential equation.




Figure 6.1: *Visualizing the application of differential equations in finite element analysis (FEA), where partial differential equations (PDEs) model stress, strain, and deformation across complex geometries.*

The Nature of Differential Equations

A **differential equation** is an equation involving an unknown function and its derivatives. When we write:

$$\frac{dy}{dx} = 2x$$

we’re expressing a relationship between an unknown function $y(x)$ and its derivative. The solution isn’t just a number but a function—in this case, $y(x) = x^2 + C$. This represents a fundamental shift in our mathematical thinking: instead of finding values, we’re finding functions that satisfy certain derivative relationships.

More formally, a differential equation can be written in the general form:

$$F(x, y, y', y'', \dots, y^{(n)}) = 0$$

where F is some function, y is our unknown function, and $y^{(k)}$ represents the k th derivative of y . The order of a differential equation is the highest derivative that appears in it. For instance, $\frac{dy}{dx} = x^2$ is first-order, $\frac{d^2y}{dx^2} + \frac{dy}{dx} = \sin(x)$ is second-order, and $\frac{d^3y}{dx^3} - x\frac{dy}{dx} = e^x$ is third-order. For instance:

- $\frac{dy}{dx} = x^2$ is first-order
- $\frac{d^2y}{dx^2} + \frac{dy}{dx} = \sin(x)$ is second-order
- $\frac{d^3y}{dx^3} - x\frac{dy}{dx} = e^x$ is third-order

For differential equations with partial derivatives, we deal with functions of multiple variables, so the general form becomes:

$$F(x_1, x_2, \dots, x_n, u, \frac{\partial u}{\partial x_1}, \frac{\partial u}{\partial x_2}, \dots, \frac{\partial^2 u}{\partial x_1^2}, \frac{\partial^2 u}{\partial x_1 \partial x_2}, \dots) = 0$$

where $u = u(x_1, x_2, \dots, x_n)$ is our unknown function of multiple variables, and $\frac{\partial u}{\partial x_i}$ represents partial derivatives with respect to the variable x_i . The order of a PDE is determined by the highest partial derivative that appears. For example:

- $\frac{\partial u}{\partial t} = \alpha \frac{\partial^2 u}{\partial x^2}$ (Heat equation) is second-order
- $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$ (Wave equation) is second-order
- $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$ (Laplace equation) is second-order

Origins and Formation

Differential equations emerge naturally when we observe how quantities change in relation to each other. Consider a simple pendulum: its acceleration (second derivative of position) is proportional to its displacement, leading to:

$$\frac{d^2\theta}{dt^2} = -\frac{g}{L} \sin(\theta)$$

where θ is the angle from vertical, g is gravitational acceleration, and L is the pendulum length. How might we even begin to solve such an equation? Our calculus instincts suggest integration, but integrating both sides still leaves us with $\frac{d\theta}{dt}$ in terms of an integral of $\sin(\theta)$. This hints at the deeper complexity we'll encounter.

The process of forming differential equations often begins with physical insight. For instance, Newton's law of cooling states that the rate of temperature change is proportional to the temperature difference:

$$\frac{dT}{dt} = -k(T - T_a)$$

where T is the object's temperature, T_a is ambient temperature, and k is a constant. Here we might have more success - notice how we could potentially separate variables with T and t on different sides.

Not all differential equations involve a single variable. Heat flow through a metal plate, as we saw, depends on position in both x and y directions and time t :

$$\frac{\partial u}{\partial t} = \alpha \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)$$

where $u(x, y, t)$ is temperature and α is thermal diffusivity. Such partial differential equations present entirely new challenges - now we're seeking functions of multiple variables!

▲ **Example 6.1.1** A cup of hot coffee is sitting in a room of constant temperature 20°C . If the coffee's temperature drops from 90°C to 60°C in the first 10 minutes, model the coffee's temperature over time.

Solution Let's think about how temperature changes:

1. First, let's understand what affects the rate of cooling:
 - The hotter the coffee is compared to the room, the faster it cools
 - As the coffee gets closer to room temperature, cooling slows down
 - The room stays at 20°C (we say it's the ambient temperature)
2. Let's translate this into mathematics:
 - Let $T(t)$ be the coffee's temperature at time t
 - The rate of change is $\frac{dT}{dt}$
 - The temperature difference from room temperature is $(T - 20)$
3. The rate of cooling is proportional to the temperature difference between the object and the surrounding environment. The constant in the differential equation must be negative to reflect that the object's temperature decreases over time when it is warmer than the surroundings. This yields the differential equation:

$$\frac{dT}{dt} = -k(T - 20)$$

This is actually known as Newton's Law of Cooling. we may find the constant k using our initial conditions which will be discussed later.

Approaches to Solutions

Solving differential equations requires a fundamentally different mindset from standard calculus. Consider first a seemingly simple ODE:

$$\frac{dy}{dx} + y = e^x$$

We seek a function $y(x)$ that, when substituted back, satisfies this relationship. But how? We could try guessing - since e^x appears on the right, perhaps $y = Ce^x$ for some constant C ? Or maybe we need a more systematic approach?

Some equations graciously yield to direct integration:

$$\frac{dy}{dx} = x^2 \sin(x)$$

immediately suggests integrating both sides. But what about:

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial t^2}$$

the famous wave equation? Integration seems less helpful here - we're dealing with partial derivatives in multiple variables. Perhaps we need a completely different strategy, like assuming solutions have a special form?

When analytical methods fail us, we might turn to numerical approximations. Consider discretizing our domain into small steps and approximating derivatives with finite differences:

$$\frac{dy}{dx} \approx \frac{y(x+h) - y(x)}{h}$$

This suggests iterative schemes that computers can implement. For PDEs, we might need to discretize in multiple dimensions, leading to systems of algebraic equations.

Sometimes, understanding behavior proves more valuable than finding exact solutions. For the logistic equation:

$$\frac{dP}{dt} = rP \left(1 - \frac{P}{K} \right)$$

we can identify equilibrium points by setting $\frac{dP}{dt} = 0$. But what about stability? How can we determine whether small perturbations grow or decay without an explicit solution? These questions lead us to qualitative theory - a powerful tool for understanding solution behavior even when exact formulas elude us.

We will systematically approach how to solve both ordinary and partial differential equations in the upcoming lectures.

Fundamental Classifications

Moving beyond basic definitions, differential equations naturally organize themselves into several fundamental categories. Each classification not only helps us understand the equation's nature but also suggests which solution methods might prove fruitful.

First and most fundamentally, we distinguish between ordinary and partial differential equations. An **ordinary differential equation (ODE)** involves an unknown function of a single independent variable. For instance:

$$\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + 2y = 0$$

represents a second-order ODE. The term "ordinary" emphasizes that only ordinary derivatives appear, not partial derivatives.

In contrast, a **partial differential equation (PDE)** involves an unknown function of several independent variables. The classic two-dimensional Laplace equation:

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

appears across physics in steady-state heat conduction, electrostatics, and fluid flow. This distinction impacts not just complexity but determines which theoretical tools we can employ.

Another crucial distinction lies between linear and nonlinear equations. A **linear ODE of order n** takes the form:

$$a_n(x)\frac{d^n y}{dx^n} + a_{n-1}(x)\frac{d^{n-1}y}{dx^{n-1}} + \cdots + a_0(x)y = g(x)$$

where $a_0, a_1, \dots, a_n, g$ are functions of x alone. Linearity means y and its derivatives appear to the first power, without multiplications between them and without nonlinear transformations like $\sin(y)$ or y^2 .

A **nonlinear equation** violates these conditions. Terms like $(y')^2$, $y \cdot y'$, or e^y indicate nonlinearity. While more challenging to solve, these equations often better capture physical reality, describing phenomena like turbulent flows or complex chemical reactions.

The **order** of a differential equation is defined by its highest derivative. A third-order equation, for instance, contains terms up to $\frac{d^3y}{dx^3}$ but no higher. Higher-order equations commonly describe complex physical systems like beam deflection or wave propagation.

We further distinguish between homogeneous and non-homogeneous equations. A **homogeneous equation** of order n has the form:

$$a_n(x)y^{(n)} + a_{n-1}(x)y^{(n-1)} + \cdots + a_0(x)y = 0$$

If instead the right side equals some non-zero function $g(x)$, we have a **non-homogeneous equation**. The homogeneous case describes free evolution, while non-homogeneous equations include external forcing.

Finally, for linear ODEs, we distinguish between constant and variable coefficients. In constant-coefficient equations:

$$a_n \frac{d^n y}{dx^n} + a_{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_0 y = g(x)$$

the coefficients a_k are constants. Such equations often yield to standard solution methods. Variable-coefficient equations, where coefficients depend on x , typically demand more sophisticated approaches.

These classifications form the foundation for understanding differential equations. As we progress, we'll develop specific techniques for each type, but first recognizing an equation's category often suggests which approach might prove most fruitful.

Initial and Boundary Conditions

A crucial insight in differential equations is that a single equation typically has infinitely many solutions. Consider the simple equation:

$$\frac{dy}{dx} = -2xy$$

whose general solution we found to be $y = Ae^{-x^2}$. Each value of A gives a different solution curve, collectively forming a family of solutions.




Figure 6.2: Family of solutions to $\frac{dy}{dx} = -2xy$. The blue curves show different solutions for various values of A , while the red curve highlights the particular solution where $A = 1$.

To single out a particular solution from this family, we need additional information. This typically comes in two forms:

1. **Initial Conditions** specify the value of the unknown function (and possibly its derivatives) at a single point. For example, if we require:

$$y(0) = 3$$

this selects the unique solution curve passing through the point $(0, 3)$. As we saw earlier, this leads to $A = 3$ and the particular solution $y = 3e^{-x^2}$. For higher-order equations, we might need multiple conditions at the same point:

$$\begin{cases} y(0) = 1 \\ y'(0) = 0 \end{cases}$$

2. **Boundary Conditions** specify values at multiple points. For instance:

$$y(0) = 0, \quad y(1) = 2$$

requires our solution to pass through both $(0, 0)$ and $(1, 2)$. This type of condition often appears in physical problems like heat distribution or beam deflection where we know values at the endpoints of our domain.

An equation together with sufficient conditions forms either an **Initial Value Problem (IVP)** or **Boundary Value Problem (BVP)**, depending on the type of conditions specified. The distinction is more than technical—the methods we use to solve these problems and even the nature of their solutions can differ dramatically. For instance:

1. IVPs typically have unique solutions that can be “evolved” forward from the initial condition
2. BVPs might have multiple solutions, no solutions, or unique solutions depending on the boundary conditions
3. Higher-order equations require correspondingly more conditions for uniqueness

The geometric interpretation is illuminating: initial/boundary conditions effectively “select” one curve from the infinite family of possible solutions. This idea of a family of solutions punctuated by particular solutions meeting specific conditions will recur throughout our study of differential equations, helping us understand both the general behavior of solutions and the role of additional conditions in selecting specific ones. Note, that the order of the equation must require that many conditions in order to fully produce a particular solution.

▲ **Example 6.1.2** Consider the ordinary differential equation:

$$\frac{dy}{dx} = -2xy$$

with initial condition $y(0) = 3$. Let's carefully examine how we might solve this equation.

Solution Let's think about how to solve this step by step. We have a first order differential equation where we can separate all terms with y on one side and all terms with x on the other.

First, let's rearrange the equation to group y terms and x terms:

$$\frac{dy}{y} = -2x \, dx$$

Now we can integrate both sides. On the left, we recognize $\frac{dy}{y}$ as the derivative of $\ln |y|$:

$$\begin{aligned} \int \frac{dy}{y} &= \int -2x \, dx \\ \ln |y| &= -x^2 + C \end{aligned}$$

To solve for y , we exponentiate both sides:

$$|y| = e^{-x^2+C} = e^{-x^2} e^C = Ae^{-x^2}$$

where $A = e^C$ is some positive constant. Since A can be positive or negative (thanks to the absolute value), our general solution is:

$$y = Ae^{-x^2}$$

Now we can use our initial condition $y(0) = 3$ to find A :

$$\begin{aligned} 3 &= Ae^{-0^2} \\ 3 &= A \cdot 1 \\ A &= 3 \end{aligned}$$

Therefore, our particular solution is:

$$y = 3e^{-x^2}$$

Verification of Solutions

Verifying solutions to differential equations is a critical step in ensuring the accuracy of our work. While finding solutions often requires creativity and problem-solving, verification is a systematic process that provides confidence in the correctness of the result. This process can be broken into clear steps for clarity and consistency.

First, we compute the necessary derivatives of the proposed solution. These derivatives are the building blocks we need to substitute back into the original differential equation. Next, by substituting the solution and its derivatives into the equation, we check whether the left-hand side equals the right-hand side. If this identity holds true, it confirms that the solution satisfies the differential equation itself. Finally, we verify any additional conditions, such as initial or boundary values, to ensure the solution fully satisfies all requirements of the problem.

Consider a simple example. To verify that $y(x) = x^2 + 1$ solves $\frac{dy}{dx} = 2x$:

Given solution: $y(x) = x^2 + 1$

Compute derivative: $\frac{dy}{dx} = 2x$

Original equation: $\frac{dy}{dx} = 2x$

Therefore: $2x = 2x$ (Identity verified)

For higher-order equations, we must check more derivatives. Consider verifying that $y(x) = e^{2x}$ solves $\frac{d^2y}{dx^2} - 4y = 0$:

Given: $y = e^{2x}$

First derivative: $\frac{dy}{dx} = 2e^{2x}$

Second derivative: $\frac{d^2y}{dx^2} = 4e^{2x}$

Substitute: $4e^{2x} - 4(e^{2x}) = 0$

Simplify: $0 = 0$ (Identity verified)

For partial differential equations, we follow the same principle but must compute partial derivatives. To verify $u(x, t) = e^{-\pi^2 t} \sin(\pi x)$ solves the heat equation $\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$:

Given: $u = e^{-\pi^2 t} \sin(\pi x)$

Time derivative: $\frac{\partial u}{\partial t} = -\pi^2 e^{-\pi^2 t} \sin(\pi x)$

Space derivative: $\frac{\partial u}{\partial x} = \pi e^{-\pi^2 t} \cos(\pi x)$

Second space derivative: $\frac{\partial^2 u}{\partial x^2} = -\pi^2 e^{-\pi^2 t} \sin(\pi x)$

Original equation: $\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}$

Substitute: $-\pi^2 e^{-\pi^2 t} \sin(\pi x) = -\pi^2 e^{-\pi^2 t} \sin(\pi x)$

Therefore: Identity verified

Important considerations when verifying solutions:

1. Check *all* conditions - not just the differential equation but also initial/boundary conditions
2. Pay attention to domains - where is the solution valid?
3. Watch for hidden assumptions - are we dividing by zero anywhere?
4. Use verification as a sanity check - does the solution make physical sense?

Verification plays a crucial role in solving differential equations. It helps catch computational errors, validates the solution techniques, and builds confidence in our methods. Moreover, it fosters intuition about the behavior of solutions, enhancing our understanding of the problem.

▲ **Example 6.1.3** Verify that $y = x + \sum_{n=1}^{\infty} \frac{y^n}{n}$ is a solution to $y' = 1 - \frac{1}{y}$. Discuss any insights about this solution.

Solution Let's follow our verification steps:

Compute derivatives

- We have an implicit equation requiring implicit differentiation
- Start with: $y = x + \sum_{n=1}^{\infty} \frac{y^n}{n}$
- Differentiate both sides:

$$\begin{aligned} \frac{d}{dx}(y) &= \frac{d}{dx} \left(x + \sum_{n=1}^{\infty} \frac{y^n}{n} \right) \\ y' &= 1 + \sum_{n=1}^{\infty} \frac{ny^{n-1}}{n} y' \\ &= 1 + y' \sum_{n=1}^{\infty} y^{n-1} \end{aligned}$$

Manipulate to standard form

- Recognize $\sum_{n=1}^{\infty} y^{n-1}$ as geometric series $\frac{1}{1-y}$ for $|y| < 1$

- Therefore: $y' = 1 + y' \frac{1}{1-y}$
- Solve for y' :

$$\begin{aligned}y' \left(1 - \frac{1}{1-y} \right) &= 1 \\y' \left(\frac{-y}{1-y} \right) &= 1 \\y' &= 1 - \frac{1}{y}\end{aligned}$$

Additional insights

- The sum $\sum_{n=1}^{\infty} \frac{y^n}{n}$ is the Taylor series of $-\ln(1-y)$
 - Solution can be written more compactly: $y = x - \ln(1-y)$
 - Valid only for $|y| < 1$ due to series convergence
-

Lecture 6.2: First-Order Ordinary Differential Equations

Having established the fundamental nature and classification of differential equations, we now delve into our first systematic study of solution techniques for first-order ordinary differential equations. These equations, while seemingly simple in form, encompass a rich variety of mathematical structures and solution methods that form the foundation for understanding more complex differential equations.

A **first-order ordinary differential equation** can be written in the general form:

$$F(x, y, y') = 0$$

where F is some function of the independent variable x , the dependent variable y , and its first derivative y' . More commonly, we encounter these equations in solved form:

$$\frac{dy}{dx} = f(x, y)$$

where $f(x, y)$ represents the relationship between the variables and the rate of change.

The designation “first-order” refers to the highest derivative present in the equation - in this case, only the first derivative appears. This seemingly simple characteristic has profound implications. Unlike higher-order equations which may require multiple conditions to specify a unique solution, a first-order equation requires only a single initial condition of the form $y(x_0) = y_0$ to determine a unique solution (when one exists).

When studying first-order equations, we often encounter them in various standard forms, each suggesting its own solution strategy. For instance, when $f(x, y)$ can be written as a product of functions of x and y separately, we have a separable equation. When $f(x, y)$ is linear in y , we have a linear first-order equation. These special forms, which we will study in detail, provide scaffolding for understanding more general cases.

The existence and uniqueness of solutions to first-order equations is governed by the fundamental theorem for first-order ODEs: if $f(x, y)$ and $\frac{\partial f}{\partial y}$ are continuous in some rectangle containing an initial point (x_0, y_0) , then there exists a unique solution to the initial value problem in some interval containing x_0 . This theorem provides theoretical foundation for our solution methods and helps us understand when and where we can expect solutions to exist.

Separation of Variables

The method of separation of variables represents our first and easiest systematic approach to solving differential equations. This technique applies when we can rewrite our equation in the form:

$$\frac{dy}{dx} = f(x)g(y)$$

where f depends only on x and g depends only on y . The key insight is to rearrange terms so that all y -dependent terms are on one side and all x -dependent terms are on the other. This will always be your go to first visual check when trying to solve an ODE.

More formally, we rewrite the equation as:

$$\frac{1}{g(y)} dy = f(x) dx$$

This transformation allows us to integrate both sides independently:

$$\int \frac{1}{g(y)} dy = \int f(x) dx + C$$

The resulting equation implicitly defines y as a function of x . In fortunate cases, we can solve explicitly for y , though often the solution remains in implicit form. This method's elegance lies in its reduction of a differential equation to algebraic manipulation and integration. This method obviously extends for cases of higher order derivatives which will be addressed in the next chapter.

Several theoretical considerations deserve attention:

1. The method assumes $g(y) \neq 0$. Points where $g(y) = 0$ may yield additional solutions.
2. The integration introduces a constant C , generating a family of solutions. Both integrations of g and f produce an arbitrary constant, but we can just combine them to form one arbitrary constant instead of two – A fact that will be used frequently in the solving of DEs.
3. Solutions may exist only on intervals where both f and g are continuous.

We have already reasoned our way through one separable ODE in the last lecture. Let's try one that is not so obvious:

▲ **Example 6.2.1** Solve the differential equation:

$$\frac{dy}{dx} = \frac{xy + 3x + 2y + 6}{xy + x + y + 1}$$

Solution Let's solve this step by step:

1. First, notice that both numerator and denominator can be factored through grouping:

$$\begin{aligned} xy + 3x + 2y + 6 &= x(y + 3) + 2(y + 3) \\ &= (x + 2)(y + 3) \end{aligned}$$

And similarly:

$$\begin{aligned}xy + x + y + 1 &= x(y + 1) + 1(y + 1) \\ &= (x + 1)(y + 1)\end{aligned}$$

2. Our differential equation becomes:

$$\frac{dy}{dx} = \frac{(x + 2)(y + 3)}{(x + 1)(y + 1)}$$

3. To separate variables, multiply both sides by dx and rearrange:

$$\frac{y + 1}{y + 3} dy = \frac{x + 2}{x + 1} dx$$

4. To integrate, first perform partial fractions on each side:

$$\begin{aligned}\frac{y + 1}{y + 3} &= 1 - \frac{2}{y + 3} \\ \frac{x + 2}{x + 1} &= 1 + \frac{1}{x + 1}\end{aligned}$$

5. Our equation becomes:

$$\int \left(1 - \frac{2}{y + 3}\right) dy = \int \left(1 + \frac{1}{x + 1}\right) dx$$

6. Integrate both sides:

$$y - 2 \ln |y + 3| + C_1 = x + \ln |x + 1| + C_2$$

7. Combining constants, our general solution is:

$$y - 2 \ln |y + 3| = x + \ln |x + 1| + C, x \neq -1, y \neq -3$$

Linear First-Order Equations

A **linear first-order ODE** takes the standard form:

$$\frac{dy}{dx} + P(x)y = Q(x)$$

where $P(x)$ and $Q(x)$ are continuous functions. The linearity refers to the fact that y and its derivative appear to the first power only, with no products between them and no

nonlinear functions of y . When $Q(x) = 0$, we call the equation homogeneous; otherwise, it's non-homogeneous. This distinction plays a crucial role in both the theory and solution methods.

Any first-order linear equation can be written in this standard form through algebraic manipulation. For instance, equations like:

$$\frac{dy}{dx} = ay + b \quad \text{or} \quad x \frac{dy}{dx} - 2y = x^3$$

can be rewritten in standard form by collecting terms and dividing appropriately. This standardization is more than cosmetic—it reveals the underlying structure and suggests solution strategies.

The solution method involves an elegant technique called the **integrating factor**. The key insight is that our equation almost looks like the product rule for derivatives (we're aiming for $(\mu y)' = \mu' y + \mu y'$), but not quite. If we could somehow make the left side into a perfect differential—that is, find a function $\mu(x)$ that when multiplied through makes the left side exactly the derivative of a product $\frac{d}{dx}[\mu(x)y]$ —then we could solve by simple integration. With this motivation in mind, we multiply both sides by a carefully chosen function $\mu(x)$ that makes the left side the derivative of a product:

$$\mu(x) \frac{dy}{dx} + \mu(x)P(x)y = \mu(x)Q(x)$$

The magic of this method lies in choosing $\mu(x)$ to make the left side a perfect differential. The integrating factor $\mu(x)$ must satisfy:

$$\frac{d}{dx}[\mu(x)y] = \mu(x) \frac{dy}{dx} + \mu'(x)y$$

For this to equal our left side, we require:

$$\mu'(x)y = \mu(x)P(x)y$$

Therefore:

$$\mu'(x) = \mu(x)P(x)$$

This first-order separable equation for μ yields:

$$\frac{d\mu}{\mu} = P(x)dx$$

Integration gives us:

$$\mu(x) = e^{\int P(x)dx}$$

After multiplication by $\mu(x)$, our equation takes the elegant form:

$$\frac{d}{dx}[\mu(x)y] = \mu(x)Q(x)$$

Integration yields the general solution:

$$y = \frac{1}{\mu(x)} \left[\int \mu(x)Q(x)dx + C \right]$$

Several theoretical aspects deserve attention:

1. The integrating factor is not unique—any constant multiple of $\mu(x)$ also works.
2. The solution process naturally splits into finding a complementary function (solution to homogeneous equation) and a particular integral.
3. The method always works when $P(x)$ and $Q(x)$ are continuous, making it one of our most reliable tools.

The form of this solution reveals a fundamental principle: every linear first-order ODE can be solved through integration alone. This is not true for nonlinear equations, where we often need more sophisticated techniques. The solution also exhibits the superposition principle characteristic of linear equations: the general solution is the sum of a particular solution and the general solution to the homogeneous equation.

When $Q(x) = 0$ (homogeneous case), our solution simplifies to:

$$y = \frac{C}{\mu(x)} = Ce^{-\int P(x)dx}$$

This form clearly shows how the integrating factor “undoes” the effect of the coefficient $P(x)$. The exponential form of this solution appears frequently in applications, modeling phenomena like radioactive decay and temperature cooling.

▲ **Example 6.2.2** Solve the differential equation:

$$x^2 \frac{dy}{dx} - 3xy = x^3 \cos(x) + 2x^2y$$

Solution

1. First, we need to rearrange into standard form $\frac{dy}{dx} + P(x)y = Q(x)$:

$$\begin{aligned} x^2 \frac{dy}{dx} - 3xy &= x^3 \cos(x) + 2x^2y \\ x^2 \frac{dy}{dx} - 3xy - 2x^2y &= x^3 \cos(x) \\ x^2 \frac{dy}{dx} - xy(3 + 2x) &= x^3 \cos(x) \end{aligned}$$

Divide throughout by x^2 (note: this restricts our domain to $x \neq 0$):

$$\frac{dy}{dx} - \frac{y(3+2x)}{x} = x \cos(x)$$

Therefore:

$$\frac{dy}{dx} + \frac{y(-3-2x)}{x} = x \cos(x)$$

2. Now we have standard form with:

$$P(x) = -\frac{3+2x}{x} \quad \text{and} \quad Q(x) = x \cos(x)$$

3. Find the integrating factor $\mu(x) = e^{\int P(x)dx}$:

$$\begin{aligned}\mu(x) &= e^{\int -\frac{3+2x}{x} dx} \\ &= e^{\int (-\frac{3}{x}-2) dx} \\ &= e^{-3 \ln |x| - 2x} \\ &= e^{-2x} x^{-3}\end{aligned}$$

4. Multiply the equation by $\mu(x)$:

$$e^{-2x} x^{-3} \frac{dy}{dx} + e^{-2x} x^{-3} \cdot \frac{-3-2x}{x} y = e^{-2x} x^{-3} \cdot x \cos(x)$$

This is equivalent to:

$$\frac{d}{dx}(e^{-2x} x^{-3} y) = e^{-2x} x^{-2} \cos(x)$$

5. Integrate both sides:

$$e^{-2x} x^{-3} y = \int e^{-2x} x^{-2} \cos(x) dx + C$$

6. Therefore:

$$y = x^3 e^{2x} \left[\int e^{-2x} x^{-2} \cos(x) dx + C \right]$$

Sometimes, we must leave the integral in the solution if it is too hard to evaluate – and that's perfectly fine!

Exact Differential Equations

An equation of the form:

$$M(x, y)dx + N(x, y)dy = 0$$

is called **exact** if there exists a function $\psi(x, y)$, called the potential function, such that:

$$d\psi = M(x, y)dx + N(x, y)dy$$

This implies:

$$\frac{\partial \psi}{\partial x} = M(x, y) \quad \text{and} \quad \frac{\partial \psi}{\partial y} = N(x, y)$$

The fundamental theorem of calculus for line integrals tells us this occurs precisely when:

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

This condition is both necessary and sufficient for exactness, providing a straightforward test before attempting to find a solution. When an equation is exact, its solution is implicit in the form:

$$\psi(x, y) = C$$

There are two methods for finding the potential function $\psi(x, y)$:

Method 1: Integration with respect to x , then find $h(y)$

1. Integrate $M(x, y)$ with respect to x , treating y as constant:

$$\psi(x, y) = \int M(x, y)dx + h(y)$$

where $h(y)$ is an arbitrary function of y alone.

2. To find $h(y)$, differentiate ψ with respect to y :

$$\frac{\partial \psi}{\partial y} = \frac{\partial}{\partial y} \left[\int M(x, y)dx \right] + h'(y)$$

3. Since $\frac{\partial \psi}{\partial y} = N(x, y)$, we have:

$$\frac{\partial}{\partial y} \left[\int M(x, y)dx \right] + h'(y) = N(x, y)$$

4. Solve for $h'(y)$:

$$h'(y) = N(x, y) - \frac{\partial}{\partial y} \left[\int M(x, y)dx \right]$$

5. Integrate to find $h(y)$:

$$h(y) = \int h'(y)dy$$

Method 2: Integration with respect to y , then find $g(x)$

1. Integrate $N(x, y)$ with respect to y , treating x as constant:

$$\psi(x, y) = \int N(x, y)dy + g(x)$$

where $g(x)$ is an arbitrary function of x alone.

2. To find $g(x)$, differentiate ψ with respect to x :

$$\frac{\partial \psi}{\partial x} = \frac{\partial}{\partial x} \left[\int N(x, y)dy \right] + g'(x)$$

3. Since $\frac{\partial \psi}{\partial x} = M(x, y)$, we have:

$$\frac{\partial}{\partial x} \left[\int N(x, y)dy \right] + g'(x) = M(x, y)$$

4. Solve for $g'(x)$:

$$g'(x) = M(x, y) - \frac{\partial}{\partial x} \left[\int N(x, y)dy \right]$$

5. Integrate to find $g(x)$:

$$g(x) = \int g'(x)dx$$

Important Observations:

- Both methods must yield the same potential function $\psi(x, y)$ (up to a constant) if the equation is exact.
- The exactness condition $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$ ensures that either method works and gives consistent results.
- When finding $h(y)$ or $g(x)$, the resulting expression must be independent of x or y respectively. If not, the equation is not exact.
- The verification process is crucial: after finding $\psi(x, y)$, we should check that both $\frac{\partial \psi}{\partial x} = M(x, y)$ and $\frac{\partial \psi}{\partial y} = N(x, y)$ are satisfied.
- The constant of integration can be omitted when finding $h(y)$ or $g(x)$ since it will be absorbed into the final constant C in the general solution $\psi(x, y) = C$.

The geometric interpretation of exact differential equations is profound: the solution curves are level curves of the potential function $\psi(x, y)$, and $M(x, y)dx + N(x, y)dy = 0$ represents the condition that motion along a solution curve maintains a constant value of ψ .

▲ **Example 6.2.3** Show that the differential equation:

$$(2xy + y^2) dx + (x^2 + 2xy) dy = 0$$

is exact, and solve it using both methods.

Solution Let's solve this systematically:

1. First, let's verify that the equation is exact by checking if $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$:

$$\begin{aligned} M(x, y) = 2xy + y^2 &\implies \frac{\partial M}{\partial y} = 2x + 2y \\ N(x, y) = x^2 + 2xy &\implies \frac{\partial N}{\partial x} = 2x + 2y \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, the equation is exact.

2. **Method 1:** Integrate $M(x, y)$ with respect to x

$$\begin{aligned} \psi(x, y) &= \int M(x, y) dx + h(y) \\ &= \int (2xy + y^2) dx + h(y) \\ &= x^2y + xy^2 + h(y) \end{aligned}$$

Now find $h(y)$ by comparing $\frac{\partial \psi}{\partial y}$ with $N(x, y)$:

$$\begin{aligned} \frac{\partial \psi}{\partial y} &= x^2 + 2xy + h'(y) = N(x, y) = x^2 + 2xy \\ \therefore h'(y) &= 0 \\ \therefore h(y) &= C_1 \end{aligned}$$

Thus, by Method 1:

$$\psi(x, y) = x^2y + xy^2 + C_1$$

3. **Method 2:** Integrate $N(x, y)$ with respect to y

$$\psi(x, y) = \int N(x, y) dy + g(x)$$

$$\begin{aligned}
&= \int (x^2 + 2xy) dy + g(x) \\
&= x^2y + xy^2 + g(x)
\end{aligned}$$

Now find $g(x)$ by comparing $\frac{\partial \psi}{\partial x}$ with $M(x, y)$:

$$\begin{aligned}
\frac{\partial \psi}{\partial x} &= 2xy + y^2 + g'(x) = M(x, y) = 2xy + y^2 \\
\therefore g'(x) &= 0 \\
\therefore g(x) &= C_2
\end{aligned}$$

Thus, by Method 2:

$$\psi(x, y) = x^2y + xy^2 + C_2$$

4. The general solution is:

$$\psi(x, y) = x^2y + xy^2 = C$$

where $C = -C_1 = -C_2$ is an arbitrary constant.

Several important observations:

- Both methods yielded the same potential function $\psi(x, y)$ (up to a constant)
- We can verify our solution by checking that $\frac{\partial \psi}{\partial x} = M(x, y)$ and $\frac{\partial \psi}{\partial y} = N(x, y)$
- The solution represents a family of level curves of $\psi(x, y)$
- The integrating constants C_1 and C_2 from each method combine into a single constant C in the final solution
- The equation $x^2y + xy^2 = C$ can be rewritten to form a quadratic in y :

$$y^2 + xy - \frac{C}{x} = 0.$$

Using the quadratic formula, we solve for y :

$$y = \frac{-x \pm \sqrt{x^3 + 4C}}{2x}.$$

Thus, the explicit solutions for y are:

$$y = \frac{-x + \sqrt{x^3 + 4C}}{2x} \quad \text{and} \quad y = \frac{-x - \sqrt{x^3 + 4C}}{2x}.$$

Bernoulli Differential Equations

A **Bernoulli differential equation** takes the form:

$$\frac{dy}{dx} + P(x)y = Q(x)y^n$$

where n is any real number except 0 or 1, and $P(x)$ and $Q(x)$ are continuous functions. When $n = 0$, the equation reduces to a linear equation, and when $n = 1$, the equation is already linear. Therefore, these cases are excluded from our consideration.

The brilliance of Bernoulli's insight was recognizing that a clever substitution could transform this nonlinear equation into a linear one. The key transformation is:

$$v = y^{1-n}$$

Let's see how this transformation works. First, differentiate v with respect to x using the chain rule:

$$\frac{dv}{dx} = (1-n)y^{-n} \frac{dy}{dx}$$

From our original equation:

$$\frac{dy}{dx} = -P(x)y + Q(x)y^n$$

Substituting this into our expression for $\frac{dv}{dx}$:

$$\begin{aligned} \frac{dv}{dx} &= (1-n)y^{-n}(-P(x)y + Q(x)y^n) \\ &= (1-n)[-P(x)y^{1-n} + Q(x)y^{2n-n}] \\ &= (1-n)[-P(x)v + Q(x)] \end{aligned}$$

This transforms our equation into the linear form:

$$\frac{dv}{dx} + (1-n)P(x)v = (1-n)Q(x)$$

This is a linear first-order equation in v which we can solve using the integrating factor method. The integrating factor is:

$$\mu(x) = e^{(1-n) \int P(x) dx}$$

Following the standard procedure for linear equations, the general solution for v is:

$$v = e^{-(1-n) \int P(x) dx} \left[\int (1-n)Q(x)e^{(1-n) \int P(x) dx} dx + C \right]$$

Substituting back $v = y^{1-n}$, we obtain the general solution for the Bernoulli equation:

$$y = \left\{ e^{-(1-n) \int P(x) dx} \left[\int (1-n)Q(x)e^{(1-n) \int P(x) dx} dx + C \right] \right\}^{\frac{1}{1-n}}$$

Several important observations about Bernoulli equations:

1. The transformation $v = y^{1-n}$ is valid only when $y \neq 0$. The zero solution may need to be checked separately.
2. The exponent n plays a crucial role in determining the behavior of solutions. Different values of n can lead to dramatically different solution behaviors.
3. When $n < 0$, solutions may have vertical asymptotes. When $n > 1$, solutions may have horizontal asymptotes.
4. The integrating factor method used after transformation highlights the deep connection between linear and nonlinear differential equations.
5. Bernoulli equations often arise in applications where quantities grow or decay with power-law behavior, such as in population dynamics and chemical reactions.

The Bernoulli equation serves as a bridge between linear and nonlinear theory. While it's nonlinear in its original form, its amenability to transformation into a linear equation makes it one of the few nonlinear equations for which we can write down a general solution formula. This makes it an excellent introduction to the study of nonlinear differential equations.

A Word on Existence and Uniqueness

When studying differential equations, two fundamental questions arise: does a solution exist, and if so, is it unique? These questions are not merely theoretical—they inform us whether our solution methods will be fruitful and whether we can trust that we've found all possible solutions.

For a first-order differential equation in the form:

$$\frac{dy}{dx} = f(x, y), \quad y(x_0) = y_0$$

the **Existence Theorem** states that a solution exists if $f(x, y)$ is continuous in some rectangle containing the initial point (x_0, y_0) . This solution may not be valid for all x , but it exists in some interval containing x_0 .

The **Uniqueness Theorem** adds that this solution is unique if $\frac{\partial f}{\partial y}$ is also continuous in this rectangle. Together, these conditions are known as the **Existence and Uniqueness Theorem**.

The geometric interpretation is illuminating. Consider the slope field of the differential equation:

- Continuity of $f(x, y)$ ensures the slopes vary smoothly—no sudden jumps or breaks
- Continuity of $\frac{\partial f}{\partial y}$ ensures the slopes change smoothly as we move vertically—preventing solution curves from crossing

Several important scenarios can violate these conditions:

1. When $f(x, y)$ has a discontinuity, solutions might not exist. For example:

$$\frac{dy}{dx} = \frac{1}{x}, \quad y(0) = 0$$

has no solution through $(0, 0)$ due to division by zero.

2. When $\frac{\partial f}{\partial y}$ is discontinuous, multiple solutions might exist. Consider:

$$\frac{dy}{dx} = |y|^{\frac{1}{3}}$$

Through $(0, 0)$, both $y = 0$ and $y = \pm \left(\frac{2x}{3}\right)^{\frac{3}{2}}$ are solutions.

For higher-order equations:

$$y^{(n)} = f(x, y, y', \dots, y^{(n-1)})$$

we can convert to a system of first-order equations and apply similar principles. The key conditions become:

- f must be continuous in all variables
- $\frac{\partial f}{\partial y^{(k)}}$ must be continuous for $k = 0, 1, \dots, n - 1$

These theorems provide powerful guarantees:

- When conditions are met, numerical methods will converge to the unique solution
- Local existence ensures our solution methods are meaningful
- Uniqueness validates our practice of finding general solutions with arbitrary constants

However, these theorems have limitations:

1. They only guarantee local existence—solutions might blow up in finite time
2. They don't tell us how to find the solution
3. They don't specify the size of the interval where the solution exists

Understanding existence and uniqueness helps us approach differential equations with confidence, knowing when our methods will yield meaningful results and when we need to exercise caution in our analysis. These theorems form the theoretical foundation that supports all our practical solution techniques.

While we have focused on ordinary differential equations, existence and uniqueness theory extends crucially to partial differential equations. In PDEs, the questions become significantly more complex due to the multiple independent variables involved. For example, the well-posedness of a PDE problem requires not only existence and uniqueness but also continuous dependence on initial/boundary conditions. Classical PDEs like the heat equation, wave equation, and Laplace equation each have their own existence and uniqueness theorems with specific requirements on domains, boundary conditions, and solution regularity. The celebrated Cauchy-Kowalevski theorem provides existence and uniqueness for analytic solutions of certain PDEs, though its applicability is limited. These considerations become central when studying PDEs in more advanced courses.

Lecture 6.3: Higher-Order Ordinary Differential Equations I

Of course we cannot stop with analyzing functions with a single order derivative, thus we now venture into the rich territory of higher-order differential equations. These equations, containing derivatives of order two or greater, model a vast array of physical phenomena from the simple harmonic oscillator to complex structural vibrations, from gravitational interactions to quantum mechanical systems. Their study reveals deeper mathematical structures and requires more sophisticated solution techniques.

Fundamental Concepts

An **n th-order differential equation** is an equation containing derivatives up to order n . In its most general form, it can be written as:

$$F(x, y, y', y'', \dots, y^{(n)}) = 0$$

or when solved for the highest derivative:

$$y^{(n)} = f(x, y, y', y'', \dots, y^{(n-1)})$$

where f is some function of x , y , and derivatives of y up to order $n - 1$.

A general **linear n th-order differential equation** takes the special form:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \dots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$

where $a_n(x), a_{n-1}(x), \dots, a_0(x)$ are continuous functions with $a_n(x) \neq 0$, and $g(x)$ is also continuous. When $g(x) = 0$, we call the equation homogeneous; otherwise, it's non-homogeneous.

A particularly important case arises when all coefficients are constants:

$$a_n \frac{d^n y}{dx^n} + a_{n-1} \frac{d^{n-1} y}{dx^{n-1}} + \dots + a_1 \frac{dy}{dx} + a_0 y = g(x)$$

For both linear and nonlinear equations, the order of the equation determines how many additional conditions we need for a unique solution. For an n th-order equation, we need n conditions of the form:

$$y(x_0) = y_0, \quad y'(x_0) = y_1, \quad \dots, \quad y^{(n-1)}(x_0) = y_{n-1}$$

This requirement of n initial conditions is not arbitrary but follows from the fundamental theory of differential equations. Each condition serves to determine one of the arbitrary constants that arise in the general solution.

The existence and uniqueness theorem for n th-order equations states that if f and its partial derivatives with respect to $y, y', \dots, y^{(n-1)}$ are continuous in some region containing the initial point, then there exists a unique solution in some interval containing x_0 . This holds for both linear and nonlinear equations.

A fundamental difference between linear and nonlinear equations lies in how solutions can be combined. For linear equations, any linear combination of solutions is again a solution (the superposition principle). This property does not hold for nonlinear equations, making their analysis generally more challenging.

The Principle of Superposition

One of the most powerful tools in solving linear differential equations is the principle of superposition, which states:

1. If $y_1(x)$ and $y_2(x)$ are solutions to the homogeneous equation, then $c_1y_1(x) + c_2y_2(x)$ is also a solution for any constants c_1, c_2 .
2. If $y_h(x)$ is a solution to the homogeneous equation and $y_p(x)$ is a particular solution to the non-homogeneous equation, then $y_h(x) + y_p(x)$ is a general solution to the non-homogeneous equation.

Let's prove both parts. Consider the linear differential equation:

$$a_n(x) \frac{d^n y}{dx^n} + a_{n-1}(x) \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$

Part 1 (Homogeneous Case):

Let y_1 and y_2 be solutions to the homogeneous equation. Then:

$$a_n(x) \frac{d^n y_1}{dx^n} + \cdots + a_0(x)y_1 = 0$$

$$a_n(x) \frac{d^n y_2}{dx^n} + \cdots + a_0(x)y_2 = 0$$

For any constants c_1, c_2 , consider $y = c_1y_1 + c_2y_2$. Note that:

$$\frac{d^k}{dx^k}(c_1y_1 + c_2y_2) = c_1 \frac{d^k y_1}{dx^k} + c_2 \frac{d^k y_2}{dx^k}$$

for any k . Therefore:

$$\begin{aligned} & a_n(x) \frac{d^n}{dx^n}(c_1y_1 + c_2y_2) + \cdots + a_0(x)(c_1y_1 + c_2y_2) \\ &= c_1[a_n(x) \frac{d^n y_1}{dx^n} + \cdots + a_0(x)y_1] + c_2[a_n(x) \frac{d^n y_2}{dx^n} + \cdots + a_0(x)y_2] \\ &= c_1(0) + c_2(0) = 0 \end{aligned}$$

Part 2 (Non-homogeneous Case):

Let y_h be a solution to the homogeneous equation and y_p a particular solution. Then:

$$a_n(x) \frac{d^n y_h}{dx^n} + \cdots + a_0(x) y_h = 0$$

$$a_n(x) \frac{d^n y_p}{dx^n} + \cdots + a_0(x) y_p = g(x)$$

For $y = y_h + y_p$, using the linearity of differentiation:

$$\begin{aligned} & a_n(x) \frac{d^n}{dx^n} (y_h + y_p) + \cdots + a_0(x) (y_h + y_p) \\ &= [a_n(x) \frac{d^n y_h}{dx^n} + \cdots + a_0(x) y_h] + [a_n(x) \frac{d^n y_p}{dx^n} + \cdots + a_0(x) y_p] \\ &= 0 + g(x) = g(x) \end{aligned}$$

This principle allows us to build complex solutions from simpler ones and suggests our general strategy: first find the complementary function (general solution to homogeneous equation), then find a particular solution. The proof reveals why this principle works only for linear equations - we crucially used the fact that derivatives of sums are sums of derivatives, and that we can factor out constants.

Second-Order Linear Equations

Second-order differential equations arise naturally in many physical contexts and serve as the foundation for understanding higher-order equations. In their most general form, they appear as:

$$p(x) \frac{d^2 y}{dx^2} + q(x) \frac{dy}{dx} + r(x) y = g(x)$$

where $p(x)$, $q(x)$, and $r(x)$ are continuous functions with $p(x) \neq 0$, and $g(x)$ is also continuous. When $g(x) = 0$, we have a homogeneous equation; otherwise, it's non-homogeneous.

A particularly important special case occurs when the coefficients are constants:

$$a \frac{d^2 y}{dx^2} + b \frac{dy}{dx} + cy = g(x)$$

where a , b , and c are constants with $a \neq 0$. This form appears frequently in applications such as:

- Simple harmonic motion
- Forced oscillations
- RLC circuits
- Beam deflection

When the equation is autonomous (no explicit x dependence) and homogeneous:

$$a\frac{d^2y}{dx^2} + b\frac{dy}{dx} + cy = 0$$

we can develop particularly powerful solution methods based on characteristic equations (which we will cover next). But first, a crucial concept in the study of second-order equations is that of linear independence. Two solutions $y_1(x)$ and $y_2(x)$ are said to be **linearly independent** on an interval I if no nontrivial linear combination vanishes on that interval. That is, if:

$$c_1y_1(x) + c_2y_2(x) = 0$$

for all x in I implies $c_1 = c_2 = 0$.

Linear independence can be tested using the **Wronskian**, defined as:

$$W(y_1, y_2)(x) = y_1y_2' - y_2y_1'$$

If $W(y_1, y_2)(x) \neq 0$ at any point in an interval, then y_1 and y_2 are linearly independent there. When we have two linearly independent solutions to the homogeneous equation, any solution can be written as their linear combination:

$$y = c_1y_1(x) + c_2y_2(x)$$

where c_1 and c_2 are determined by initial conditions.

The Characteristic Equation Method

For homogeneous constant-coefficient equations:

$$ay'' + by' + cy = 0$$

we seek solutions of the form $y = e^{rx}$. Substituting this ansatz:

$$ar^2e^{rx} + bre^{rx} + ce^{rx} = 0$$

$$e^{rx}(ar^2 + br + c) = 0$$

Since $e^{rx} \neq 0$, we obtain the characteristic equation:

$$ar^2 + br + c = 0$$

The nature of the roots determines the form of our solution:

1. Distinct real roots r_1, r_2 :

$$y = c_1e^{r_1x} + c_2e^{r_2x}$$

2. Repeated real root r_1 :

$$y = c_1e^{r_1x} + c_2xe^{r_1x}$$

3. Complex conjugate roots $\alpha \pm i\beta$:

$$y = e^{\alpha x}(c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

This method extends naturally to higher orders. For an n th-order equation:

$$a_n y^{(n)} + a_{n-1} y^{(n-1)} + \cdots + a_1 y' + a_0 y = 0$$

the characteristic equation becomes:

$$a_n r^n + a_{n-1} r^{n-1} + \cdots + a_1 r + a_0 = 0$$

The Fundamental Theorem of Algebra ensures n complex roots (counting multiplicity), leading to the following solution patterns:

- For a real root r of multiplicity k : $e^{rx}, xe^{rx}, x^2 e^{rx}, \dots, x^{k-1} e^{rx}$
- For complex conjugate pairs $\alpha \pm i\beta$: $e^{\alpha x} \cos(\beta x), e^{\alpha x} \sin(\beta x)$
- For repeated complex pairs: multiply by increasing powers of x

Method of Undetermined Coefficients

The method of undetermined coefficients provides a systematic approach to finding particular solutions of non-homogeneous linear differential equations with constant coefficients. The process begins with finding the complementary function and then addresses the forcing term.

Consider the non-homogeneous equation:

$$ay'' + by' + cy = g(x)$$

The complete solution will have the form:

$$y = y_h + y_p$$

where y_h is the general solution to the homogeneous equation and y_p is a particular solution.

Step 1: Find the Complementary Function: First, solve the homogeneous equation:

$$ay'' + by' + cy = 0$$

using the characteristic equation method. This gives y_h , which contains arbitrary constants.

Step 2: Address the Forcing Function: For the non-homogeneous part, we use superposition when $g(x)$ is a sum of terms:

$$g(x) = g_1(x) + g_2(x) + \cdots + g_n(x)$$

This allows us to solve simpler equations:

$$ay'' + by' + cy = g_k(x)$$

for each k , and then add the solutions: $y_p = y_{p1} + y_{p2} + \cdots + y_{pn}$.

For each type of forcing term, we have specific trial solutions:

1. For polynomials $g(x) = P_n(x)$ of degree n :

$$y_p = A_n x^n + A_{n-1} x^{n-1} + \cdots + A_1 x + A_0$$

2. For exponentials $g(x) = e^{ax}$:

$$y_p = Ae^{ax}$$

3. For trigonometric terms $g(x) = \cos(bx)$ or $\sin(bx)$:

$$y_p = A \cos(bx) + B \sin(bx)$$

4. For products $g(x) = x^n e^{ax}$:

$$y_p = (A_n x^n + A_{n-1} x^{n-1} + \cdots + A_0) e^{ax}$$

5. For combined trigonometric-exponential terms:

$$g(x) = e^{ax}(P \cos(bx) + Q \sin(bx)) \implies y_p = e^{ax}(R \cos(bx) + S \sin(bx))$$

The Resonance Case: If any term in the trial solution is also a solution to the homogeneous equation, we must modify our guess. This happens when:

- The forcing function matches a term in y_h
- The characteristic equation has a root related to the forcing function

In such cases, multiply the trial solution by x^k where k is the minimum power needed for linear independence. For example:

- If e^{rx} is both a forcing term and in y_h , try $y_p = Ax e^{rx}$
- If it's a double root, try $y_p = Ax^2 e^{rx}$

Consider:

$$y'' - y = 3e^x + 2 \cos(x) + x^2$$

We break this into three equations:

$$\begin{array}{ll} y'' - y = 3e^x & \text{(exponential term)} \\ y'' - y = 2 \cos(x) & \text{(trigonometric term)} \\ y'' - y = x^2 & \text{(polynomial term)} \end{array}$$

For each equation:

1. First find y_h from $y'' - y = 0$:

$$y_h = c_1 e^x + c_2 e^{-x}$$

2. Then find each y_{pk} using appropriate trial solutions

3. Combine all solutions:

$$y = y_h + y_{p1} + y_{p2} + y_{p3}$$

This systematic approach, combined with superposition, allows us to solve a wide range of non-homogeneous equations efficiently. The key is recognizing that:

1. The homogeneous solution must be found first
 2. Complex forcing functions can be broken down
 3. Each piece can be solved independently
 4. The sum of all solutions gives the general solution
 5. Resonance cases require special attention
-

▲ **Example 6.3.1** Solve the differential equation:

$$y'' - 4y' + 4y = 2x + 3e^{2x} + 4\sin(x)$$

Solution We'll solve via superposition, handling the homogeneous part and then each non-homogeneous term separately.

1. **Homogeneous solution:** Solve

$$y'' - 4y' + 4y = 0.$$

Characteristic eq.: $r^2 - 4r + 4 = 0 \implies (r-2)^2 = 0 \implies r = 2$ (double root).

$$y_h = c_1 e^{2x} + c_2 x e^{2x}.$$

2. **Non-homogeneous terms:**

$$g_1(x) = 2x, \quad g_2(x) = 3e^{2x}, \quad g_3(x) = 4\sin(x).$$

We find particular solutions for each and sum them.

3. **Particular solution for $g_1(x) = 2x$:**

Try $y_{p1} = Ax + B$.

$$y'_{p1} = A, \quad y''_{p1} = 0.$$

Substitute into $y'' - 4y' + 4y$:

$$0 - 4A + 4(Ax + B) = 2x.$$

$$4Ax + 4B - 4A = 2x.$$

Matching coefficients:

$$4A = 2 \implies A = \frac{1}{2}, \quad 4B - 4A = 0 \implies B = \frac{1}{2}.$$

$$y_{p1} = \frac{1}{2}x + \frac{1}{2}.$$

4. Particular solution for $g_2(x) = 3e^{2x}$:

Since e^{2x} is part of the homogeneous solution (double root at $r = 2$), try

$$y_{p2} = Ax^2e^{2x}.$$

Compute derivatives (correctly applying the product rule):

$$y'_{p2} = \frac{d}{dx}(Ax^2e^{2x}) = Ae^{2x}(2x^2 + 2x),$$

$$y''_{p2} = \frac{d}{dx}(Ae^{2x}(2x^2 + 2x)) = 2Ae^{2x}(2x^2 + 4x + 1).$$

Substitute into $y'' - 4y' + 4y$:

$$(2A(2x^2 + 4x + 1)e^{2x}) - 4(Ae^{2x}(2x^2 + 2x)) + 4(Ax^2e^{2x}) = 3e^{2x}.$$

After simplification, the expression collapses to $2Ae^{2x}$. So

$$2Ae^{2x} = 3e^{2x} \implies A = \frac{3}{2}.$$

$$y_{p2} = \frac{3}{2}x^2e^{2x}.$$

5. Particular solution for $g_3(x) = 4\sin(x)$:

Try $y_{p3} = A\cos(x) + B\sin(x)$.

$$y'_{p3} = -A\sin(x) + B\cos(x), \quad y''_{p3} = -A\cos(x) - B\sin(x).$$

Substitute into $y'' - 4y' + 4y$:

$$(-A\cos x - B\sin x) - 4(-A\sin x + B\cos x) + 4(A\cos x + B\sin x).$$

Collecting terms:

$$\cos(x) : \quad (-A) + (-4B) + (4A) = 3A - 4B,$$

$$\sin(x) : \quad (-B) + (4A) + (4B) = 4A + 3B.$$

We want

$$(3A - 4B)\cos(x) + (4A + 3B)\sin(x) = 4\sin(x).$$

Hence

$$3A - 4B = 0, \quad 4A + 3B = 4.$$

Solve:

$$3A = 4B \implies A = \frac{4B}{3},$$

$$4\left(\frac{4B}{3}\right) + 3B = 4 \implies \frac{16B}{3} + 3B = 4 \implies \frac{25B}{3} = 4 \implies B = \frac{12}{25}, \quad A = \frac{16}{25}.$$

$$y_{p3} = \frac{16}{25} \cos(x) + \frac{12}{25} \sin(x).$$

6. Combine all parts for the general solution:

$$y = y_h + y_{p1} + y_{p2} + y_{p3} = c_1 e^{2x} + c_2 x e^{2x} + \frac{1}{2}x + \frac{1}{2} + \frac{3}{2}x^2 e^{2x} + \frac{16}{25} \cos(x) + \frac{12}{25} \sin(x).$$

This demonstrates how to handle each term independently and then superimpose results into one final solution.

▲ Example 6.3.2 Consider the fifth-order homogeneous linear differential equation with constant coefficients:

$$y^{(5)} - 9y^{(4)} + 32y^{(3)} - 56y'' + 47y' - 15y = 0$$

Solution Let's solve this systematically:

1. Form the characteristic equation by replacing $y^{(n)}$ with r^n :

$$r^5 - 9r^4 + 32r^3 - 56r^2 + 47r - 15 = 0$$

2. Factor this polynomial. By inspection or using algebraic methods, we find:

$$r = 1 \text{ (double root)}$$

$$r = 3 \text{ (single root)}$$

$$r = 2 \pm i \text{ (complex conjugate pair)}$$

Therefore our polynomial factors as:

$$(r - 1)^2(r - 3)(r^2 - 4r + 5) = 0$$

$$\text{where } r^2 - 4r + 5 = (r - (2 + i))(r - (2 - i))$$

3. Generate solutions based on each root type:

- For $r = 1$ (double root):

$$y_1 = e^x \text{ and } y_2 = xe^x$$

- For $r = 3$ (single root):

$$y_3 = e^{3x}$$

- For $r = 2 \pm i$ (complex conjugate pair):

$$y_4 = e^{2x} \cos(x) \text{ and } y_5 = e^{2x} \sin(x)$$

4. Write the general solution as a linear combination:

$$y = c_1 e^x + c_2 x e^x + c_3 e^{3x} + e^{2x} (c_4 \cos(x) + c_5 \sin(x))$$

Several important observations:

- The double root at $r = 1$ contributes two terms: e^x and $x e^x$
- The complex conjugate pair provides oscillatory behavior modulated by e^{2x}
- The number of arbitrary constants (5) matches the order of the equation
- Each term corresponds directly to a root or root pair of the characteristic equation

Variation of Parameters

The *method of undetermined coefficients* leverages structured guesses for y_p (a particular solution) when the non-homogeneous term $g(x)$ has a familiar form (e.g. polynomials, exponentials, sines/cosines, or products thereof). However, for more complicated or “weird” source functions $g(x)$, such guesses may not be obvious or practical. In such cases, *variation of parameters* provides a more general and powerful procedure that works *without* depending on the form of $g(x)$.

Consider the general second-order ODE:

$$y'' + p(x)y' + q(x)y = g(x),$$

where $p(x)$ and $q(x)$ are coefficient functions of x . Suppose we have already solved the associated *homogeneous* equation,

$$y'' + p(x)y' + q(x)y = 0,$$

and found two linearly independent solutions, $y_1(x)$ and $y_2(x)$. Variation of parameters seeks a particular solution of the form

$$y_p(x) = u_1(x)y_1(x) + u_2(x)y_2(x),$$

where $u_1(x)$ and $u_2(x)$ are unknown functions to be determined. To find $u_1(x)$ and $u_2(x)$, we impose the conditions

$$\begin{aligned}u_1'(x) y_1(x) + u_2'(x) y_2(x) &= 0, \\u_1'(x) y_1'(x) + u_2'(x) y_2'(x) &= g(x),\end{aligned}$$

which simplify the algebra upon substituting $y_p(x)$ into the original differential equation. By solving this system for $u_1'(x)$ and $u_2'(x)$, we obtain:

$$u_1'(x) = -\frac{y_2(x) g(x)}{W(x)}, \quad u_2'(x) = \frac{y_1(x) g(x)}{W(x)},$$

where

$$W(x) = y_1(x) y_2'(x) - y_2(x) y_1'(x)$$

is the *Wronskian* of y_1 and y_2 . The functions $u_1(x)$ and $u_2(x)$ then follow by integrating u_1' and u_2' :

$$u_1(x) = -\int \frac{y_2(x) g(x)}{W(x)} dx, \quad u_2(x) = \int \frac{y_1(x) g(x)}{W(x)} dx.$$

Substituting $u_1(x)$ and $u_2(x)$ back into $y_p(x)$ provides a particular solution that, when combined with the homogeneous solution, yields the complete general solution to

$$y'' + p(x) y' + q(x) y = g(x).$$

More general cases: This method scales naturally to higher-order ODEs. For the n th-order linear differential equation

$$y^{(n)} + p_{n-1}(x) y^{(n-1)} + \cdots + p_0(x) y = g(x),$$

one starts with n linearly independent homogeneous solutions $y_1(x), \dots, y_n(x)$ and proposes

$$y_p(x) = u_1(x) y_1(x) + \cdots + u_n(x) y_n(x).$$

The corresponding variation of parameters system ensures each $u_i(x)$ is determined consistently. This makes variation of parameters *much more powerful* than undetermined coefficients, because it does not rely on recognizing special forms of $g(x)$.

In summary, whenever the non-homogeneous term $g(x)$ does not match standard patterns for undetermined coefficients (or is simply too cumbersome to guess effectively), variation of parameters offers a systematic procedure for finding a particular solution, leveraging the homogeneous solution and the Wronskian.

▲ **Example 6.3.3** Solve the non-homogeneous ODE via Variation of Parameters:

$$y'' + y = \tan(x).$$

Solution

1. Write the homogeneous equation.

The associated homogeneous ODE is

$$y'' + y = 0.$$

Its characteristic equation is

$$r^2 + 1 = 0 \implies r = \pm i.$$

Therefore, the general homogeneous solution is

$$y_h(x) = C_1 \cos(x) + C_2 \sin(x).$$

2. Set up Variation of Parameters for the non-homogeneous part.

We aim to find a *particular* solution of the form

$$y_p(x) = u_1(x) \cos(x) + u_2(x) \sin(x),$$

where $u_1(x)$ and $u_2(x)$ are unknown functions.

3. Identify the Wronskian and the standard formulas.

For two linearly independent solutions $y_1(x) = \cos(x)$ and $y_2(x) = \sin(x)$, their Wronskian is

$$W(x) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix} = \cos^2(x) + \sin^2(x) = 1.$$

Variation of parameters (for a second-order ODE of the form $y'' + p(x)y' + q(x)y = g(x)$) gives:

$$u_1'(x) = -\frac{y_2(x)g(x)}{W(x)}, \quad u_2'(x) = \frac{y_1(x)g(x)}{W(x)}.$$

Here, $g(x) = \tan(x)$, $y_1(x) = \cos(x)$, $y_2(x) = \sin(x)$, and $W(x) = 1$.

4. Find $u_1'(x)$ and $u_2'(x)$.

$$u_1'(x) = -\sin(x) \tan(x) = -\sin(x) \frac{\sin(x)}{\cos(x)} = -\frac{\sin^2(x)}{\cos(x)},$$

$$u_2'(x) = \cos(x) \tan(x) = \cos(x) \frac{\sin(x)}{\cos(x)} = \sin(x).$$

5. Integrate to find $u_1(x)$ and $u_2(x)$.

$$u_1(x) = \int u_1'(x) dx = \int \left(-\frac{\sin^2(x)}{\cos(x)} \right) dx = - \int \frac{\sin^2(x)}{\cos(x)} dx.$$

Observe:

$$\int \frac{\sin^2(x)}{\cos(x)} dx = \int \frac{1 - \cos^2(x)}{\cos(x)} dx = \int \sec(x) dx - \int \cos(x) dx.$$

We know $\int \sec(x) dx = \ln|\sec(x) + \tan(x)|$, and $\int \cos(x) dx = \sin(x)$. Hence,

$$\int \frac{\sin^2(x)}{\cos(x)} dx = \ln|\sec(x) + \tan(x)| - \sin(x).$$

Thus,

$$u_1(x) = -\left[\ln|\sec(x) + \tan(x)| - \sin(x)\right] = -\ln|\sec(x) + \tan(x)| + \sin(x).$$

Meanwhile,

$$u_2(x) = \int u_2'(x) dx = \int \sin(x) dx = -\cos(x).$$

6. Write the particular solution.

We now have for $y_p(x)$:

$$u_1(x) \cos(x) + u_2(x) \sin(x) = [\sin(x) - \ln|\sec(x) + \tan(x)|] \cos(x) - \cos(x) \sin(x).$$

Notice that $\sin(x) \cos(x) - \cos(x) \sin(x) = 0$. So the expression simplifies to:

$$y_p(x) = -\cos(x) \ln|\sec(x) + \tan(x)|.$$

7. Combine with the homogeneous solution for the general solution.

The full solution is

$$y(x) = y_h(x) + y_p(x) = C_1 \cos(x) + C_2 \sin(x) - \cos(x) \ln|\sec(x) + \tan(x)|.$$

where C_1, C_2 are arbitrary constants.

This example illustrates how variation of parameters yields a particular solution even when standard guessing (undetermined coefficients) is not straightforward. By leveraging known homogeneous solutions and the Wronskian, we solve for functions $u_1(x)$ and $u_2(x)$ in a systematic way, regardless of the complexity of $\tan(x)$.

Lecture 6.4: Higher-Order Ordinary Differential Equations II

And now after spending lots of time and effort learning very specific techniques, we will now introduce a method that will blast half of those out of the water. Transform methods are exceptionally powerful tools in solving ordinary differential equations (ODEs). Laplace Transforms convert differential equations into algebraic equations, simplifying the solution process.

These methods allow us to handle complex boundary conditions, discontinuities, and impulsive forcing functions with remarkable ease. By transforming an ODE into the frequency domain, the operations of differentiation and integration become multiplication and division, respectively, making the equations much more tractable. This approach not only simplifies the solution of linear ODEs but also provides insights into the behavior of physical systems, making transform methods indispensable in engineering, physics, and applied mathematics.

Introduction to Laplace Transforms

The key idea behind the Laplace transform is to convert differential and integral operations in the time domain into algebraic operations in the complex frequency domain. This conversion often simplifies the process of solving differential equations by transforming them into algebraic equations, which are generally easier to manipulate and solve.

Formally, the Laplace transform establishes a one-to-one correspondence between functions in the time domain and functions in the complex frequency domain, subject to certain conditions of existence. This correspondence allows us to solve problems in whichever domain is most convenient and then transform the solution back to the original domain.

For a function $f(t)$ defined for $t \geq 0$, the Laplace transform $F(s)$ is defined as:

$$F(s) = \mathcal{L}\{f(t)\} = \int_0^{\infty} e^{-st} f(t) dt,$$

where $s = \sigma + i\omega$ is a complex variable. The real part σ is called the attenuation, and the imaginary part ω is called the angular frequency. The integral in the definition converges if $f(t)$ is of exponential order, meaning there exist constants $M > 0$, a , and T such that $|f(t)| \leq Me^{at}$ for all $t > T$. In this case, the integral converges for all s with $\text{Re}(s) > a$.

Laplace transforms offer several significant advantages in solving differential equations and analyzing linear systems:

1. **Differential to Algebraic Conversion:** The Laplace transform converts an n th-order linear differential equation:

$$a_n \frac{d^n y}{dt^n} + a_{n-1} \frac{d^{n-1} y}{dt^{n-1}} + \cdots + a_0 y = f(t)$$

into a simpler algebraic equation:

$$(a_n s^n + a_{n-1} s^{n-1} + \cdots + a_0)Y(s) = F(s) + (\text{initial conditions})$$

2. **Handling Discontinuous Functions:** The transform elegantly handles discontinuous inputs through simple expressions:

$$\mathcal{L}\{u(t)\} = \frac{1}{s}, \quad \mathcal{L}\{\delta(t)\} = 1$$

3. **Initial Conditions:** Initial conditions emerge naturally in the transformed derivatives:

$$\mathcal{L}\{f'(t)\} = sF(s) - f(0), \quad \mathcal{L}\{f''(t)\} = s^2 F(s) - sf(0) - f'(0)$$

4. **Linear Time-Invariant Systems:** The transform converts complex convolution operations:

$$y(t) = \int_0^t h(\tau)x(t-\tau)d\tau$$

into simple multiplication:

$$Y(s) = H(s)X(s)$$

where $H(s)$ represents the system's transfer function.

Properties of Laplace Transforms

The following properties form the core mathematical framework for applying Laplace transforms to differential equations and system analysis:

1. **Linearity:**

$$\mathcal{L}\{af(t) + bg(t)\} = a\mathcal{L}\{f(t)\} + b\mathcal{L}\{g(t)\}$$

This fundamental property enables decomposition of complex functions into simpler elements.

2. **First Derivative:**

$$\mathcal{L}\{f'(t)\} = sF(s) - f(0)$$

Converts time-domain differentiation into s-domain multiplication.

3. **Higher-Order Derivatives:**

$$\mathcal{L}\{f^{(n)}(t)\} = s^n F(s) - s^{n-1}f(0) - s^{n-2}f'(0) - \cdots - f^{(n-1)}(0)$$

Extends the derivative property to nth-order derivatives.

4. **Integration:**

$$\mathcal{L}\left\{\int_0^t f(\tau) d\tau\right\} = \frac{1}{s} F(s)$$

Transforms time-domain integration into s-domain division.

5. Frequency Shifting:

$$\mathcal{L}\{e^{at} f(t)\} = F(s - a)$$

Handles exponential factors in time domain.

6. Time Shifting:

$$\mathcal{L}\{f(t - a) u(t - a)\} = e^{-as} F(s)$$

Addresses delayed functions in time domain.

7. Scaling:

$$\mathcal{L}\{f(at)\} = \frac{1}{a} F\left(\frac{s}{a}\right)$$

Relates time scaling to frequency scaling.

8. Convolution:

$$\mathcal{L}\{f(t) * g(t)\} = F(s) G(s)$$

Transforms time-domain convolution to s-domain multiplication.

The following comprehensive table presents common transform pairs essential for solving both initial value problems and boundary value problems. When solving differential equations, we first transform the equation using these pairs, solve the resulting algebraic equation in the s -domain, and then use these same pairs in reverse to find our time-domain solution.

Table 6.1: *Comprehensive Table of Laplace Transform Pairs: These pairs enable both forward transformation for problem-solving and inverse transformation for solution recovery.*

$f(t)$	$\mathcal{L}\{f(t)\}$	$f(t)$	$\mathcal{L}\{f(t)\}$
1	$\frac{1}{s}$	$t \sin(at)$	$\frac{2a}{(s^2 + a^2)^2}$
t	$\frac{1}{s^2}$	$t \cos(at)$	$\frac{s^2 - a^2}{(s^2 + a^2)^2}$
t^n	$\frac{n!}{s^{n+1}}$	$t^n e^{at}$	$\frac{n!}{(s - a)^{n+1}}$
e^{at}	$\frac{1}{s - a}$	$\frac{t^n}{n!}$	$\frac{1}{s^{n+1}}$
$\sin(at)$	$\frac{a}{s^2 + a^2}$	$t^n \cos(at)$	$\frac{n!s}{(s^2 + a^2)^{n+1}}$
$\cos(at)$	$\frac{s}{s^2 + a^2}$	$t^n \sin(at)$	$\frac{n!a}{(s^2 + a^2)^{n+1}}$
$\sinh(at)$	$\frac{a}{s^2 - a^2}$	$\frac{\sin(at)}{t}$	$\tan^{-1}\left(\frac{a}{s}\right)$
$\cosh(at)$	$\frac{s}{s^2 - a^2}$	$J_0(at)$	$\frac{1}{\sqrt{s^2 + a^2}}$

$e^{at} \sin(bt)$	$\frac{b}{(s-a)^2+b^2}$	$u(t-c)$	$\frac{e^{-cs}}{s}$
$e^{at} \cos(bt)$	$\frac{s-a}{(s-a)^2+b^2}$	$\delta(t-c)$	e^{-cs}
$e^{at} f(t)$	$F(s-a)$	$u(t-a)$	$\frac{e^{-as}}{s}$
te^{at}	$\frac{1}{(s-a)^2}$	$\int_0^t f(\tau)g(t-\tau)d\tau$	$F(s)G(s)$
e^{-at^2}	$\frac{\sqrt{\pi}}{2\sqrt{a}} e^{\frac{s^2}{4a}} \operatorname{erfc}\left(\frac{s}{2\sqrt{a}}\right)$	$t^n u(t)$	$\frac{n!}{s^{n+1}}$

Steps for Solving ODEs Using Laplace Transforms

To solve a linear ODE using Laplace transforms, we follow a systematic approach that transforms our problem into algebraic form and back:

1. **Forward Transform:** Take the Laplace transform of both sides of the ODE using Table 2 and the properties of derivatives:

- $\mathcal{L}\{y'\} = sY(s) - y(0)$
- $\mathcal{L}\{y''\} = s^2Y(s) - sy(0) - y'(0)$

2. **Algebraic Manipulation:** Solve the resulting algebraic equation for $Y(s)$, typically obtaining a rational function:

$$Y(s) = \frac{P(s)}{Q(s)}$$

where $P(s)$ and $Q(s)$ are polynomials in s .

3. **Partial Fraction Decomposition:** Just like in a standard Calculus II class, we break the rational function into simpler terms:

$$\frac{P(s)}{Q(s)} = \frac{A}{s-a} + \frac{Bs+C}{s^2+ps+q} + \dots$$

4. **Inverse Transform:** Convert back to the time domain using Table 2 in reverse.

Inverse Laplace Transform Strategy

While the inverse transform has a formal definition via complex integration:

$$f(t) = \mathcal{L}^{-1}\{F(s)\} = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{st} F(s) ds,$$

in practice, we use Table 2 systematically:

1. **Pattern Recognition:** Look for standard forms in Table 2:

- $\frac{1}{s-a} \rightarrow e^{at}$
- $\frac{s}{s^2+a^2} \rightarrow \cos(at)$
- $\frac{a}{s^2+a^2} \rightarrow \sin(at)$

2. **Linear Combination:** Use the linearity property:

$$\mathcal{L}^{-1}\{aF(s) + bG(s)\} = af(t) + bg(t)$$

3. **Complex Terms:** For terms not directly in the table:

- Use shifting properties
 - Apply derivative or integral properties
 - Combine simpler known transforms
-

▲ **Example 6.4.1** Find the inverse Laplace transform of:

$$F(s) = \frac{3}{s^2 + 4}$$

Solution

1. First, analyze the form of $F(s)$:

- The denominator is a quadratic: $s^2 + 4 = s^2 + (2)^2$
- There is no s term in the numerator
- The coefficient in front is 3

2. Express $F(s)$ in standard form:

$$F(s) = 3 \cdot \frac{1}{s^2 + (2)^2}$$

This matches the form $\frac{1}{s^2+a^2}$ with $a = 2$

3. Recall from our transform table that:

$$\mathcal{L}^{-1}\left\{\frac{1}{s^2 + a^2}\right\} = \frac{1}{a} \sin(at)$$

4. Apply this result using the linearity property:

$$\begin{aligned}
 \mathcal{L}^{-1}\{F(s)\} &= \mathcal{L}^{-1}\left\{3 \cdot \frac{1}{s^2 + 4}\right\} \\
 &= 3 \cdot \mathcal{L}^{-1}\left\{\frac{1}{s^2 + 4}\right\} \\
 &= 3 \cdot \frac{1}{2} \sin(2t) \\
 &= \frac{3}{2} \sin(2t)
 \end{aligned}$$

Now let's put everything we have learned together to tackle solving an ODE with Laplace Transforms. Note, that up till now, this is the most robust method of solving ODEs. However, beware that this is computationally tedious to work out and one small error can make you have a completely wrong solution. It is up to you to weigh which method you use to tackle ODEs. Typically, the more complicated source functions, the better Laplace Transforms are to solve the corresponding ODE.

▲ **Example 6.4.2** Solve the initial-value problem via Laplace transforms:

$$y'' + 4y = \sin(2t), \quad y(0) = 0, \quad y'(0) = 0.$$

Solution

1. **Take the Laplace transform of both sides:**

Recall:

$$\mathcal{L}\{y''\}(s) = s^2 Y(s) - s y(0) - y'(0),$$

and

$$\mathcal{L}\{\sin(2t)\}(s) = \frac{2}{s^2 + 4}.$$

Given the initial conditions $y(0) = 0$ and $y'(0) = 0$, we have:

$$\mathcal{L}\{y''\}(s) = s^2 Y(s).$$

Therefore, applying the Laplace transform to the ODE gives:

$$s^2 Y(s) + 4Y(s) = \frac{2}{s^2 + 4}.$$

2. **Factor and solve for $Y(s)$:**

$$(s^2 + 4)Y(s) = \frac{2}{s^2 + 4} \implies Y(s) = \frac{2}{(s^2 + 4)^2}.$$

3. **Find the inverse Laplace transform of $Y(s)$:**

We use the known result:

$$\mathcal{L}^{-1}\left\{\frac{1}{(s^2 + a^2)^2}\right\} = \frac{1}{2a^3}(a \sin(at) - t \cos(at)).$$

Here, $a = 2$, so

$$\mathcal{L}^{-1}\left\{\frac{1}{(s^2 + 4)^2}\right\} = \frac{1}{2 \cdot 2^3}(2 \sin(2t) - t \cos(2t)) = \frac{1}{16}(2 \sin(2t) - t \cos(2t)).$$

Since $Y(s) = 2 \cdot \frac{1}{(s^2 + 4)^2}$, we multiply the above by 2:

$$y(t) = 2 \times \frac{1}{16}(2 \sin(2t) - t \cos(2t)) = \frac{1}{4} \sin(2t) - \frac{t}{8} \cos(2t).$$

4. **State the final solution:**

$$y(t) = \frac{1}{4} \sin(2t) - \frac{t}{8} \cos(2t).$$

▲ **Example 6.4.3** Solve the initial-value problem (IVP) via Laplace transforms:

$$y'' + 2y' + 5y = e^{-t} \cos(2t) + \delta(t - 3), \quad y(0) = 1, \quad y'(0) = 0.$$

Solution

1. **Take the Laplace transform of both sides.**

Let $Y(s)$ be the Laplace transform of $y(t)$. We know:

$$\mathcal{L}\{y''\}(s) = s^2 Y(s) - s y(0) - y'(0),$$

$$\mathcal{L}\{y'\}(s) = s Y(s) - y(0).$$

Also recall:

$$\mathcal{L}\{e^{-t} \cos(2t)\}(s) = \mathcal{L}\{\cos(2t)\} \Big|_{s \rightarrow s+1} = \frac{s+1}{(s+1)^2 + 4},$$

$$\mathcal{L}\{\delta(t-a)\}(s) = e^{-as}.$$

Given $y(0) = 1$ and $y'(0) = 0$:

$$\mathcal{L}\{y''\}(s) = s^2 Y(s) - s - 0 = s^2 Y(s) - s,$$

$$\mathcal{L}\{2y'\}(s) = 2(sY(s) - 1) = 2sY(s) - 2.$$

Summing these and adding $5\mathcal{L}\{y\} = 5Y(s)$:

$$(s^2 Y(s) - s) + (2sY(s) - 2) + 5Y(s) = \frac{s+1}{(s+1)^2 + 4} + e^{-3s}.$$

2. Combine like terms to solve for $Y(s)$.

$$s^2 Y(s) + 2sY(s) + 5Y(s) - s - 2 = \frac{s+1}{(s+1)^2 + 4} + e^{-3s}.$$

Factor out $Y(s)$:

$$(s^2 + 2s + 5)Y(s) - (s + 2) = \frac{s+1}{(s+1)^2 + 4} + e^{-3s}.$$

So:

$$(s^2 + 2s + 5)Y(s) = (s + 2) + \frac{s+1}{(s+1)^2 + 4} + e^{-3s}.$$

$$Y(s) = \frac{s+2}{s^2 + 2s + 5} + \frac{s+1}{(s+1)^2 + 4} \cdot \frac{1}{s^2 + 2s + 5} + \frac{e^{-3s}}{s^2 + 2s + 5}.$$

3. Simplify each term for inverse Laplace transform.

The characteristic equation denominator for the homogeneous part is $s^2 + 2s + 5$.

Complete the square:

$$s^2 + 2s + 5 = (s+1)^2 + 4.$$

Thus, rewrite:

$$Y(s) = \frac{s+2}{(s+1)^2 + 4} + \frac{s+1}{(s+1)^2 + 4} \cdot \frac{1}{(s+1)^2 + 4} + \frac{e^{-3s}}{(s+1)^2 + 4}.$$

We can treat each piece separately:

(a) $\frac{s+2}{(s+1)^2+4}$

Write $s+2 = (s+1) + 1$, so

$$\frac{s+2}{(s+1)^2+4} = \frac{(s+1)+1}{(s+1)^2+4} = \frac{s+1}{(s+1)^2+4} + \frac{1}{(s+1)^2+4}.$$

Taking inverse Laplace:

$$\mathcal{L}^{-1}\left\{\frac{s+1}{(s+1)^2+4}\right\} = e^{-t} \cos(2t), \quad \mathcal{L}^{-1}\left\{\frac{1}{(s+1)^2+4}\right\} = e^{-t} \frac{\sin(2t)}{2}.$$

So:

$$\mathcal{L}^{-1}\left\{\frac{s+2}{(s+1)^2+4}\right\} = e^{-t} \cos(2t) + e^{-t} \frac{\sin(2t)}{2}.$$

(b) $\frac{s+1}{(s+1)^2+4} \cdot \frac{1}{(s+1)^2+4}$

This is:

$$\frac{(s+1)}{(s+1)^2+4} \times \frac{1}{(s+1)^2+4}.$$

Convolution in the time domain:

$$\mathcal{L}^{-1}\{F(s)G(s)\} = f(t) * g(t) \quad (\text{convolution}).$$

Where

$$F(s) = \frac{s+1}{(s+1)^2+4}, \quad G(s) = \frac{1}{(s+1)^2+4}.$$

Then

$$f(t) = e^{-t} \cos(2t), \quad g(t) = e^{-t} \frac{\sin(2t)}{2}.$$

So the inverse Laplace is:

$$(f * g)(t) = \int_0^t f(\tau) g(t-\tau) d\tau.$$

That is:

$$\int_0^t \left[e^{-\tau} \cos(2\tau) \right] \left[e^{-(t-\tau)} \frac{\sin(2(t-\tau))}{2} \right] d\tau.$$

Simplifying inside the integral:

$$= \frac{1}{2} \int_0^t e^{-t} \cos(2\tau) \sin(2(t-\tau)) d\tau.$$

Factor out e^{-t} :

$$= \frac{e^{-t}}{2} \int_0^t \cos(2\tau) \sin(2(t-\tau)) d\tau.$$

This integral can be handled by trigonometric product-to-sum identities or direct expansions, but it's quite lengthy. In practice, many prefer to keep it in convolution form or consult a transform table for a product. However, leaving it in convolution form is fully correct.

$$(c) \frac{e^{-3s}}{(s+1)^2 + 4}$$

This corresponds to a time shift of $e^{-t} \sin(2t)/2$ or $e^{-t} \cos(2t)$, depending on the numerator. Here the numerator is 1, so from part (a) we know:

$$\mathcal{L}^{-1} \left\{ \frac{1}{(s+1)^2 + 4} \right\} = e^{-t} \frac{\sin(2t)}{2}.$$

The factor e^{-3s} implies a shift:

$$\mathcal{L}^{-1} \{ e^{-as} F(s) \} = u(t-a) f(t-a),$$

where $u(t-a)$ is the Heaviside step function. So

$$\mathcal{L}^{-1} \left\{ \frac{e^{-3s}}{(s+1)^2 + 4} \right\} = u(t-3) e^{-(t-3)} \frac{\sin(2(t-3))}{2}.$$

4. Combine all inverse transforms for the final solution.

Putting it all together, the homogeneous portion plus forcing yields:

$$y(t) = \underbrace{e^{-t} \cos(2t) + e^{-t} \frac{\sin(2t)}{2}}_{\text{from } (s+2)\text{-term}} + \underbrace{(f * g)(t)}_{\text{convolution term}} + \underbrace{u(t-3) e^{-(t-3)} \frac{\sin(2(t-3))}{2}}_{\text{shift term}}.$$

Where

$$f(t) = e^{-t} \cos(2t), \quad g(t) = e^{-t} \frac{\sin(2t)}{2},$$

and

$$(f * g)(t) = \int_0^t f(\tau) g(t-\tau) d\tau.$$

This is a valid closed-form (with one part as a convolution integral). In many references, leaving the product of transforms in convolution form is perfectly acceptable, unless a fully explicit form is specifically requested.

Dang, that was tough. But, in the case above, we have no other method that could robustly handle that. Personally, I just have Wolfram Alpha solve whatever needs to be solved

analytically. You can also use numerical methods discussed previously if you are solving an ODE for a more applied problem. In addition, below is a python script which can symbolically solve ODEs. Give it a try!

```

1 import sympy as sp
2
3 def solve_ode_with_sympy():
4     t = sp.Symbol('t', real=True, nonnegative=True)
5     y = sp.Function('y')(t)
6
7     # Define the ODE:
8     #  $y'' + 2y' + 5y = e^{-t}\cos(2t) + \text{DiracDelta}(t - 3)$ 
9     ode = sp.Eq(
10         y.diff(t, 2) + 2*y.diff(t) + 5*y,
11         sp.exp(-t)*sp.cos(2*t) + sp.DiracDelta(t - 3)
12     )
13
14     # Solve with initial conditions  $y(0)=1$ ,  $y'(0)=0$ 
15     sol = sp.dsolve(
16         ode,
17         ics={
18             y.subs(t, 0): 1,
19             y.diff(t).subs(t, 0): 0
20         }
21     )
22     return sol
23
24 if __name__ == "__main__":
25     solution = solve_ode_with_sympy()
26     print("Solution to the ODE:")
27     print(solution)

```

Lecture 6.5: Methods of Solving Partial Differential Equations I

After our journey through ordinary differential equations, we now confront equations where our unknown functions depend on multiple variables, leading to partial derivatives. These partial differential equations (PDEs) represent some of mathematics' most profound and practical tools, describing everything from heat flow through a metal plate to the quantum behavior of particles. The methods for solving such equations are as diverse as the phenomena they describe, ranging from separation of variables to Fourier transforms.

Unlike ODEs, where we often seek explicit solutions in terms of elementary functions, PDEs frequently require us to think differently about what constitutes a “solution.” Sometimes we seek series solutions, other times numerical approximations, and occasionally, we must be content with proving existence and uniqueness without finding explicit forms. This richness of approach reflects the complexity and beauty of multi-variable calculus. It is well known that PDEs are thus way more difficult to comprehend and solve, and thus makes it a candidate for a rich area of modern mathematics research.




Figure 6.3: *Propagation of a wave pulse with constant velocity v . The pulse maintains its shape while translating to the right, demonstrating the wave equation's fundamental solution behavior.*

Classification of PDEs

Before diving into solution methods, we must understand the different types of PDEs we might encounter. The classification of a PDE provides crucial insight into its behavior and suggests appropriate solution techniques. Consider a second-order PDE in two variables:

$$a \frac{\partial^2 u}{\partial x^2} + b \frac{\partial^2 u}{\partial x \partial y} + c \frac{\partial^2 u}{\partial y^2} + \text{lower order terms} = f(x, y)$$

We classify this equation based on the discriminant $b^2 - 4ac$:

1. If $b^2 - 4ac > 0$: Hyperbolic (e.g., wave equation)
2. If $b^2 - 4ac = 0$: Parabolic (e.g., heat equation)
3. If $b^2 - 4ac < 0$: Elliptic (e.g., Laplace equation)

This classification parallels the classification of conic sections, and like conics, each type exhibits characteristic behaviors:

- Hyperbolic equations typically describe wave-like phenomena
- Parabolic equations often model diffusion processes
- Elliptic equations frequently appear in equilibrium problems

The geometric interpretation of these classifications reveals deep connections to the underlying physical phenomena they model. A key concept is that of characteristics - special curves along which information propagates in the solution. We will explore the method of characteristics in detail later, but for now, it suffices to note that these curves fundamentally determine how disturbances propagate through the domain.

For hyperbolic equations like the wave equation:

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$$

the existence of real characteristics corresponds to finite propagation speeds of disturbances. In contrast, parabolic equations like the heat equation:

$$\frac{\partial u}{\partial t} = \alpha \frac{\partial^2 u}{\partial x^2}$$

have characteristics only in the time direction, leading to infinite propagation speed. Elliptic equations, such as Laplace's equation:

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

lack real characteristics entirely, resulting in solutions where each point is influenced by all boundary values simultaneously.

For higher dimensions, the classification extends naturally by examining the eigenvalues of the coefficient matrix. Consider a general second-order PDE in n variables:

$$\sum_{i,j=1}^n a_{ij} \frac{\partial^2 u}{\partial x_i \partial x_j} + \sum_{i=1}^n b_i \frac{\partial u}{\partial x_i} + cu = f$$

The principal part (highest order terms) determines the classification through the eigenvalue structure of $[a_{ij}]$. This spectral analysis provides a coordinate-invariant classification

that reveals the equation's fundamental character regardless of the coordinate system chosen.

Under appropriate transformations, these equations can be reduced to canonical forms that highlight their essential mathematical structure:

$$\begin{cases} \frac{\partial^2 u}{\partial \xi \partial \eta} + \text{lower order terms} = g_1 & \text{(Hyperbolic)} \\ \frac{\partial^2 u}{\partial \xi^2} + \text{lower order terms} = g_2 & \text{(Parabolic)} \\ \frac{\partial^2 u}{\partial \xi^2} + \frac{\partial^2 u}{\partial \eta^2} + \text{lower order terms} = g_3 & \text{(Elliptic)} \end{cases}$$

These forms suggest natural solution strategies that we will explore in subsequent sections.

Separation of Variables

Our first and most fundamental method for solving PDEs is separation of variables, also known as the method of Fourier. This technique makes a simple but powerful assumption: our solution can be written as a product of simpler functions, where each function depends on only one variable. While this might seem like a restrictive assumption, it turns out to work remarkably well for many important problems in physics and engineering.

Consider the heat equation in one spatial dimension:

$$\frac{\partial u}{\partial t} = \alpha \frac{\partial^2 u}{\partial x^2}, \quad 0 < x < L, \quad t > 0$$

This equation describes how heat flows through a one-dimensional object, like a metal rod. We need to specify:

- How the ends of the rod are kept (boundary conditions):

$$u(0, t) = 0, \quad u(L, t) = 0, \quad t > 0$$

- The initial temperature distribution (initial condition):

$$u(x, 0) = f(x), \quad 0 < x < L$$

Let's try assuming our solution has the special form:

$$u(x, t) = X(x)T(t)$$

where $X(x)$ is some function of position only and $T(t)$ is some function of time only. This is the key idea of separation of variables - we're hoping to break our complicated two-variable problem into two simpler one-variable problems.

When we put this assumption into our heat equation:

$$X(x)T'(t) = \alpha X''(x)T(t)$$

where $T'(t)$ means the derivative of T with respect to t , and $X''(x)$ means the second derivative of X with respect to x .

Now comes the clever part. Let's divide both sides by $\alpha X(x)T(t)$:

$$\frac{T'(t)}{\alpha T(t)} = \frac{X''(x)}{X(x)}$$

Look carefully at this equation. The left side only involves t , while the right side only involves x . How can two things be equal if one changes with t and the other with x ? The only way is if both sides equal the same constant value. Let's call this constant $-\lambda$ (the minus sign makes later steps nicer):

$$\frac{T'(t)}{\alpha T(t)} = -\lambda \quad \text{and} \quad \frac{X''(x)}{X(x)} = -\lambda$$

This separation gives us two ordinary differential equations:

$$\begin{aligned} T'(t) + \alpha\lambda T(t) &= 0 \\ X''(x) + \lambda X(x) &= 0 \end{aligned}$$

These are much simpler equations than our original PDE! The first is a basic exponential equation, and the second is similar to a simple oscillator equation. Our boundary conditions $u(0, t) = u(L, t) = 0$ mean that $X(0) = X(L) = 0$, and this helps us determine which values of λ are allowed.

The equation for $X(x)$:

$$X''(x) + \lambda X(x) = 0, \quad X(0) = X(L) = 0$$

is a familiar ODE where we're looking for both the function $X(x)$ and the value λ that make everything work. The general solution is $X(x) = A \cos(\sqrt{\lambda}x) + B \sin(\sqrt{\lambda}x)$, and applying the boundary conditions shows that non-trivial solutions exist only when $\lambda = (n\pi/L)^2$ for positive integers n , each giving a distinct mode of vibration. When we solve this equation, we find that it only works for certain special values of λ :

$$\lambda_n = \left(\frac{n\pi}{L}\right)^2, \quad n = 1, 2, 3, \dots$$

where n can be any positive integer. For each of these values, we get a corresponding solution:

$$X_n(x) = \sin\left(\frac{n\pi x}{L}\right)$$

Why sine functions? These are the only functions that:

1. Satisfy our equation $X''(x) + \lambda X(x) = 0$
2. Equal zero at both $x = 0$ and $x = L$ (our boundary conditions)

3. Give us a complete set of solutions

For each value of λ_n , we can solve the time equation:

$$T'(t) + \alpha\lambda_n T(t) = 0$$

This is a simple equation whose solution is:

$$T_n(t) = c_n e^{-\alpha\lambda_n t}$$

where c_n is some constant we'll need to determine.

Now here's where the magic happens. Since our original equation is linear (meaning solutions can be added together to get new solutions), we can combine all these individual solutions and put them in our original form $u(x, t) = X(x)T(t)$:

$$u(x, t) = \sum_{n=1}^{\infty} \underbrace{c_n e^{-\alpha(\frac{n\pi}{L})^2 t}}_{T(t)} \underbrace{\sin\left(\frac{n\pi x}{L}\right)}_{X(x)}$$

This infinite sum represents our complete solution. But what are all those constants c_n ? They come from our initial condition. When we set $t = 0$ in our solution, all exponential terms become 1:

$$u(x, 0) = \sum_{n=1}^{\infty} c_n e^{-\alpha(\frac{n\pi}{L})^2 \cdot 0} \sin\left(\frac{n\pi x}{L}\right) = \sum_{n=1}^{\infty} c_n \sin\left(\frac{n\pi x}{L}\right)$$

Since this must equal our initial temperature $f(x)$, we have:

$$f(x) = \sum_{n=1}^{\infty} c_n \sin\left(\frac{n\pi x}{L}\right)$$

This is a profound statement: we're claiming any initial temperature distribution $f(x)$ can be written as an infinite sum of sine functions. To prove this is possible and find the constants c_n , we use a clever trick: multiply both sides by $\sin\left(\frac{m\pi x}{L}\right)$ and integrate from 0 to L :

$$\int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = \int_0^L \sum_{n=1}^{\infty} c_n \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx$$

The right side can be integrated term by term, and here we use a special property of sine functions: when you multiply two different ones ($n \neq m$) and integrate, you get zero; when you multiply the same one ($n = m$), you get $L/2$:

$$\int_0^L \sin\left(\frac{n\pi x}{L}\right) \sin\left(\frac{m\pi x}{L}\right) dx = \begin{cases} 0 & \text{if } n \neq m \\ \frac{L}{2} & \text{if } n = m \end{cases}$$

Therefore, all terms in our infinite sum become zero except the one where $n = m$:

$$\int_0^L f(x) \sin\left(\frac{m\pi x}{L}\right) dx = c_m \frac{L}{2}$$

Solving for c_m gives us our formula:

$$c_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

Each c_n thus tells us exactly how much of each sine function we need to include to reconstruct our initial temperature distribution. We need infinitely many terms because, in general, the initial temperature distribution could be any continuous function satisfying our boundary conditions.

Therefore, our complete solution to the heat equation is:

$$u(x, t) = \sum_{n=1}^{\infty} \left[\frac{2}{L} \int_0^L f(\xi) \sin\left(\frac{n\pi \xi}{L}\right) d\xi \right] e^{-\alpha\left(\frac{n\pi}{L}\right)^2 t} \sin\left(\frac{n\pi x}{L}\right)$$

where we've used ξ as our integration variable to avoid confusion with the x in our solution. This single formula completely determines the temperature at any point x and time t based on our initial temperature distribution $f(x)$.

Let's understand what our solution means physically:

1. Each term $\sin\left(\frac{n\pi x}{L}\right)$ represents a possible temperature pattern
2. The exponential $e^{-\alpha\left(\frac{n\pi}{L}\right)^2 t}$ shows how that pattern decays over time
3. Larger values of n mean the pattern decays faster
4. The constants c_n tell us how much of each pattern is present at the start

▲ **Example 6.5.1** Solve the heat equation:

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}, \quad 0 < x < \pi, \quad t > 0$$

with boundary conditions $u(0, t) = u(\pi, t) = 0$ and initial condition $u(x, 0) = \sin(2x)$.

Solution

1. First, try separation of variables: $u(x, t) = X(x)T(t)$
2. Substituting into the PDE:

$$X(x)T'(t) = X''(x)T(t)$$

3. Divide by $X(x)T(t)$ to separate variables:

$$\frac{T'(t)}{T(t)} = \frac{X''(x)}{X(x)} = -\lambda$$

where $-\lambda$ is our separation constant.

4. This gives us two ODEs:

$$\begin{aligned} T'(t) + \lambda T(t) &= 0 \\ X''(x) + \lambda X(x) &= 0 \end{aligned}$$

5. The time equation $T'(t) + \lambda T(t) = 0$ has solution:

$$T(t) = ce^{-\lambda t}$$

6. The space equation with boundary conditions:

$$\begin{aligned} X''(x) + \lambda X(x) &= 0 \\ X(0) = X(\pi) &= 0 \end{aligned}$$

This is a second-order ordinary differential equation. Its general solution is:

$$X(x) = A \cos(\sqrt{\lambda}x) + B \sin(\sqrt{\lambda}x)$$

Now we apply the boundary conditions:

- At $x = 0$: $X(0) = A = 0$
- At $x = \pi$: $X(\pi) = B \sin(\sqrt{\lambda}\pi) = 0$

For a non-trivial solution (where $B \neq 0$), we must have:

$$\sin(\sqrt{\lambda}\pi) = 0$$

This equation has solutions only when:

$$\sqrt{\lambda}\pi = n\pi, \quad n = 1, 2, 3, \dots$$

Therefore:

$$\lambda = n^2, \quad n = 1, 2, 3, \dots$$

And our solutions are:

$$X_n(x) = B \sin(nx)$$

where we can absorb the constant B into our general solution's coefficients.

7. Therefore, our general solution is the sum of each particular solution (because of the Superposition Principle):

$$u(x, t) = \sum_{n=1}^{\infty} c_n e^{-n^2 t} \sin(nx)$$

8. For the initial condition $u(x, 0) = \sin(2x)$:

$$\sin(2x) = \sum_{n=1}^{\infty} c_n \sin(nx)$$

9. Multiply both sides by $\sin(mx)$ and integrate from 0 to π :

$$\int_0^{\pi} \sin(2x) \sin(mx) dx = \sum_{n=1}^{\infty} c_n \int_0^{\pi} \sin(nx) \sin(mx) dx$$

10. Using the trig identity:

$$\int_0^{\pi} \sin(nx) \sin(mx) dx = \begin{cases} 0 & \text{if } n \neq m \\ \frac{\pi}{2} & \text{if } n = m \end{cases}$$

11. This gives us:

$$c_n = \frac{2}{\pi} \int_0^{\pi} \sin(2x) \sin(nx) dx = \begin{cases} 1 & \text{if } n = 2 \\ 0 & \text{otherwise} \end{cases}$$

12. Therefore all our c_n terms that aren't 2 go to 0, so our equation $u(x, t) = \sum_{n=1}^{\infty} c_n e^{-n^2 t} \sin(nx)$ collapses to our final solution which is the single term where $n = 2$:

$$u(x, t) = e^{-4t} \sin(2x)$$

13. We can verify this solves our problem:

- $\frac{\partial u}{\partial t} = -4e^{-4t} \sin(2x)$
- $\frac{\partial^2 u}{\partial x^2} = -4e^{-4t} \sin(2x)$
- $u(0, t) = u(\pi, t) = 0$ for all $t \geq 0$
- $u(x, 0) = \sin(2x)$

This simple example illustrates two key points:

- This example is particularly simple because our initial condition $\sin(2x)$ matches exactly one of the basic solutions in our series (when $n = 2$). Since we only need to match this single sine function at $t = 0$, all other terms in our infinite series turn out to be zero. This is why our final solution only needs one term, rather than an infinite sum.
- The solution shows exponential decay in time while maintaining its spatial shape

▲ **Example 6.5.2** Consider a circular drum head of radius a vibrating in time. When you hit a drum, each point on its surface moves up and down. We describe this movement using a function $u(r, \theta, t)$ which tells us:

- r is the distance from the center (0 to a)
- θ is the angle around the circle (0 to 2π)
- t is time
- u is the height of the drum surface at point (r, θ) at time t

The physics of the vibrating drum gives us this equation:

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left(\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} \right)$$

where c is a constant related to the drum's tension and material. We also know:

- The edge of the drum is fixed: $u(a, \theta, t) = 0$
- We're given the initial shape: $u(r, \theta, 0) = f(r, \theta)$
- We're given the initial velocity: $\frac{\partial u}{\partial t}(r, \theta, 0) = g(r, \theta)$

Solution

1. Just as we did with simpler problems, let's try to write our solution as a product:

$$u(r, \theta, t) = R(r)\Phi(\theta)T(t)$$

where:

- $R(r)$ describes how the vibration changes with distance from center
- $\Phi(\theta)$ describes how it changes around the circle
- $T(t)$ describes how it changes with time

2. When we put this into our equation and separate variables:

$$\frac{T''}{c^2 T} = \frac{1}{R} \left(R'' + \frac{1}{r} R' \right) + \frac{1}{r^2} \frac{\Phi''}{\Phi} = -\lambda$$

where $-\lambda$ is our separation constant.

3. First, let's solve for $\Phi(\theta)$ so we can knock out this variable and thus later get an ODE in terms of R . Since we're going around a circle, $\Phi(\theta)$ must return to the same value after 2π :

$$\Phi(\theta + 2\pi) = \Phi(\theta)$$

The only functions that do this are:

$$\Phi(\theta) = e^{im\theta}, \quad m = 0, \pm 1, \pm 2, \dots$$

which are just combinations of sines and cosines using Euler's formula.

4. For $R(r)$, we get a harder equation:

$$r^2 R'' + r R' + (\lambda r^2 - m^2) R = 0$$

This is called Bessel's equation, and its solutions $J_m(\sqrt{\lambda}r)$ are called Bessel functions. Think of these as special vibration patterns that work in circular shapes, just like sines and cosines work for straight lines.

5. Because the edge of the drum is fixed ($u(a, \theta, t) = 0$), we need:

$$R(a) = J_m(\sqrt{\lambda}a) = 0$$

This only happens for certain values of λ , which we call $\lambda_{mn} = (j_{mn}/a)^2$, where j_{mn} is the n th time $J_m(x)$ equals zero.

6. Finally, for time, we obtain the ODE $T'' + \lambda c^2 T = 0$. Again this is a friendly equation we can solve and get solutions like those for a simple spring:

$$T(t) = A \cos(\omega t) + B \sin(\omega t)$$

where $\omega = c\sqrt{\lambda}$

7. Putting it all together and using superposition:

$$u(r, \theta, t) = \sum_{m=-\infty}^{\infty} \sum_{n=1}^{\infty} (A_{mn} \cos(\omega_{mn} t) + B_{mn} \sin(\omega_{mn} t)) J_m \left(\frac{j_{mn} r}{a} \right) e^{im\theta}$$

This solution tells us:

- The drum can vibrate in many different patterns (modes)
- Each pattern has its own frequency ω_{mn}
- When you hit a drum, you excite many of these patterns at once
- The constants A_{mn} and B_{mn} tell us how much of each pattern is present

The separation of variables method successfully solves many foundational equations of mathematical physics. Each of these equations, when posed on appropriate domains with suitable boundary conditions, yields to separation of variables, though the resulting ordinary differential equations may require sophisticated special function theory to solve completely.

1. Wave Equation:

$$\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$$

Describes vibrating strings, electromagnetic waves, and sound propagation

2. Laplace's Equation:

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

Governs steady-state phenomena like electrostatic potentials and fluid flow

3. Poisson's Equation:

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = f(x, y)$$

Appears in electrostatics with charge distributions and gravitational fields

4. Schrödinger Equation:

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} + V(x)\psi$$

Describes quantum mechanical systems and wave functions

Fourier Series Methods

Our separation of variables method led us to represent functions as infinite sums of sine functions. This naturally connects to a more general idea called Fourier series, where we can write any well-behaved function as an infinite sum of sines and cosines. On an interval $[-L, L]$, this takes the form:

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left(a_n \cos\left(\frac{n\pi x}{L}\right) + b_n \sin\left(\frac{n\pi x}{L}\right) \right)$$

where the coefficients are found by:

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos\left(\frac{n\pi x}{L}\right) dx$$

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi x}{L}\right) dx$$

This more general representation allows us to solve PDEs even when separation of variables becomes difficult. Consider the non-homogeneous heat equation:

$$\frac{\partial u}{\partial t} = \alpha \frac{\partial^2 u}{\partial x^2} + f(x, t)$$

with zero boundary conditions. We can express both u and f as Fourier series:

$$u(x, t) = \sum_{n=1}^{\infty} c_n(t) \sin\left(\frac{n\pi x}{L}\right)$$

$$f(x, t) = \sum_{n=1}^{\infty} f_n(t) \sin\left(\frac{n\pi x}{L}\right)$$

Substituting these series into the PDE:

$$\sum_{n=1}^{\infty} c'_n(t) \sin\left(\frac{n\pi x}{L}\right) = \alpha \sum_{n=1}^{\infty} -\left(\frac{n\pi}{L}\right)^2 c_n(t) \sin\left(\frac{n\pi x}{L}\right) + \sum_{n=1}^{\infty} f_n(t) \sin\left(\frac{n\pi x}{L}\right)$$

Since the sine functions $\sin\left(\frac{n\pi x}{L}\right)$ are independent of each other (multiply two different ones and integrate to get zero), if we have two different Fourier series that are equal:

$$\sum_{n=1}^{\infty} a_n \sin\left(\frac{n\pi x}{L}\right) = \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi x}{L}\right)$$

then their coefficients must be equal ($a_n = b_n$ for all n). Therefore, looking at the coefficients of each $\sin\left(\frac{n\pi x}{L}\right)$ term in our PDE:

$$c'_n(t) + \alpha \left(\frac{n\pi}{L}\right)^2 c_n(t) = f_n(t)$$

This is a first-order ODE for each $c_n(t)$, solved by:

$$c_n(t) = e^{-\alpha(\frac{n\pi}{L})^2 t} \left[c_n(0) + \int_0^t f_n(\tau) e^{\alpha(\frac{n\pi}{L})^2 \tau} d\tau \right]$$

The complete solution is then:

$$u(x, t) = \sum_{n=1}^{\infty} \left[e^{-\alpha(\frac{n\pi}{L})^2 t} \left(c_n(0) + \int_0^t f_n(\tau) e^{\alpha(\frac{n\pi}{L})^2 \tau} d\tau \right) \right] \sin\left(\frac{n\pi x}{L}\right)$$

▲ **Example 6.5.3** Consider Poisson's equation on a rectangle:

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = -1, \quad 0 < x < L, \quad 0 < y < H$$

with boundary conditions:

$$\begin{aligned} u(0, y) &= u(L, y) = 0 \\ u(x, 0) &= u(x, H) = 0 \end{aligned}$$

This could represent, for example, the deflection of a rectangular membrane under constant force.

Solution

1. Since the forcing term is constant (-1), express it as a double Fourier series:

$$-1 = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} f_{mn} \sin\left(\frac{m\pi x}{L}\right) \sin\left(\frac{n\pi y}{H}\right)$$

where:

$$f_{mn} = \frac{4}{LH} \int_0^L \int_0^H (-1) \sin\left(\frac{m\pi x}{L}\right) \sin\left(\frac{n\pi y}{H}\right) dy dx$$

2. Similarly, write the solution as:

$$u(x, y) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} c_{mn} \sin\left(\frac{m\pi x}{L}\right) \sin\left(\frac{n\pi y}{H}\right)$$

3. Substituting into Poisson's equation:

$$\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \left[-\left(\frac{m\pi}{L}\right)^2 - \left(\frac{n\pi}{H}\right)^2 \right] c_{mn} \sin\left(\frac{m\pi x}{L}\right) \sin\left(\frac{n\pi y}{H}\right) = f_{mn} \sin\left(\frac{m\pi x}{L}\right) \sin\left(\frac{n\pi y}{H}\right)$$

4. Comparing coefficients:

$$c_{mn} = \frac{f_{mn}}{\left(\frac{m\pi}{L}\right)^2 + \left(\frac{n\pi}{H}\right)^2}$$

5. Computing f_{mn} :

$$f_{mn} = \frac{16}{mn\pi^2} \quad \text{for odd } m, n$$

and zero otherwise.

6. Therefore, our solution is:

$$u(x, y) = \sum_{m, n \text{ odd}} \frac{16}{mn\pi^2} \frac{\sin\left(\frac{m\pi x}{L}\right) \sin\left(\frac{n\pi y}{H}\right)}{\left(\frac{m\pi}{L}\right)^2 + \left(\frac{n\pi}{H}\right)^2}$$

This solution demonstrates several key features:

- Only odd terms appear due to the symmetry of the problem
 - The solution decays rapidly for large m, n due to the denominators
 - The maximum deflection occurs at the center of the rectangle
-

Lecture 6.6: Methods of Solving Partial Differential Equations II

After exploring separation of variables and Fourier methods, we confront a fundamental question: what happens when our PDEs resist these powerful but specialized techniques? What if our equation doesn't neatly separate, or our domain isn't amenable to Fourier analysis? Nature, after all, rarely confines itself to rectangular domains and simple boundary conditions. We need more general tools, and this leads us to one of the most elegant and geometrically insightful approaches: the method of characteristics.

Consider a simple wave propagating through space. How does information travel? If we drop a stone in a pond, we see ripples moving outward along definite paths. If we shout in a still room, sound waves travel along straight lines until they hit walls. These intuitive observations suggest a profound mathematical truth: the solutions to many PDEs naturally organize themselves along special curves in space and time. These curves—our characteristics—aren't just mathematical constructs; they represent the actual paths along which information propagates.

Method of Characteristics

The method of characteristics is a technique for solving first-order partial differential equations by reducing them to a system of ordinary differential equations along special curves where the solution behaves in a particularly simple way. Consider a first-order PDE of the form:

$$a(x, y) \frac{\partial u}{\partial x} + b(x, y) \frac{\partial u}{\partial y} = c(x, y, u).$$

This equation presents a fundamental challenge: the function $u(x, y)$ depends on two independent variables in a manner that does not readily permit direct integration. However, if we can identify curves in the domain along which the PDE transforms into an ordinary differential equation, solving for u becomes significantly more tractable. These curves, called *characteristics*, are paths along which information propagates and where the solution can be determined explicitly.

To derive the characteristic equations, consider a curve parameterized by t , given by $(x(t), y(t))$, along which the solution can be expressed as $u = u(x(t), y(t))$. Applying the chain rule, the total derivative of u along this curve satisfies:

$$\frac{du}{dt} = \frac{\partial u}{\partial x} \frac{dx}{dt} + \frac{\partial u}{\partial y} \frac{dy}{dt}.$$

The original PDE suggests choosing the characteristic curves such that:

$$\frac{dx}{dt} = a(x, y), \quad \frac{dy}{dt} = b(x, y).$$

Substituting these into the total derivative expression, we obtain:

$$\frac{du}{dt} = a(x, y) \frac{\partial u}{\partial x} + b(x, y) \frac{\partial u}{\partial y} = c(x, y, u).$$

This leads to the system of characteristic equations:

$$\frac{dx}{dt} = a(x, y), \quad \frac{dy}{dt} = b(x, y), \quad \frac{du}{dt} = c(x, y, u).$$

These equations provide a direct method for solving the PDE: first, determine the characteristic curves by solving the system for $x(t)$ and $y(t)$, then solve for $u(t)$ along these curves. Geometrically, characteristics represent trajectories in the (x, y) -plane along which the function u evolves according to an ordinary differential equation. This transformation is powerful because it allows the PDE to be solved implicitly by integrating along these curves. Instead of solving for $u(x, y)$ explicitly, we solve the system:

$$\frac{dx}{a(x, y)} = \frac{dy}{b(x, y)} = \frac{du}{c(x, y, u)}.$$

Once these equations are integrated, the solution is reconstructed implicitly in terms of the characteristic variables. The method of characteristics thus reveals an underlying structure within the PDE that makes it solvable by standard techniques for ODEs, providing insight into how solutions propagate in the domain.




Figure 6.4: *Geometric visualization of the method of characteristics. Left: Characteristic curves in the xy -plane showing the paths along which information propagates. Right: Evolution of the solution u along different characteristics, with matching colors indicating correspondence between paths and solutions. Each characteristic curve C_i determines the evolution of its corresponding solution $u_i(t)$.*

The geometric elegance of the method of characteristics comes alive through the visualizations in Figure 6.4. In the left panel, characteristic curves serve as “information highways”—paths along which our solution propagates through space. The blue curves C_1 ,

$\mathcal{C}_2$, and $\mathcal{C}_3$ trace these pathways, with arrows indicating the evolution parameter t .

The right panel reveals how solutions evolve along these predetermined routes, with red curves $u_1(t)$, $u_2(t)$, and $u_3(t)$ showing the solution values changing along different characteristics. Together, these visualizations transform our abstract PDE into a concrete geometric process: characteristic curves show where information flows, while evolution profiles reveal how solution values change along these paths.

To apply the method of characteristics in practice, follow these steps systematically. The goal is to reduce the given first-order PDE to a set of ordinary differential equations and then solve for the solution along characteristic curves.

1. Identify the given PDE and express it in standard form.

Given a first-order PDE of the form:

$$a(x, y) \frac{\partial u}{\partial x} + b(x, y) \frac{\partial u}{\partial y} = c(x, y, u),$$

verify that it is written explicitly in terms of its characteristic coefficients $a(x, y)$, $b(x, y)$, and $c(x, y, u)$. If necessary, rearrange terms to ensure that the PDE is properly structured.

2. Set up the characteristic equations.

The core idea of the method is to transform the PDE into a system of ordinary differential equations along characteristic curves. The characteristic equations are derived from the total derivative of u along these curves:

$$\frac{dx}{dt} = a(x, y), \quad \frac{dy}{dt} = b(x, y), \quad \frac{du}{dt} = c(x, y, u).$$

These equations describe the evolution of x , y , and u as functions of a parameter t along the characteristic curves.

3. Set up invariants along the characteristic curves.

For nonlinear PDEs, it is often useful to identify invariants—quantities that remain constant along each characteristic curve. To do this, look for functions $I(x, y, u)$ such that:

$$\frac{d}{dt} I(x(t), y(t), u(t)) = 0.$$

These invariants are typically obtained by taking appropriate ratios or combinations of the characteristic equations, and they can greatly simplify the integration process by reducing the number of independent variables.

4. Solve for the characteristic curves.

Solve the first two equations:

$$\frac{dx}{dt} = a(x, y), \quad \frac{dy}{dt} = b(x, y).$$

This system provides the characteristic curves, which represent the paths along which information propagates in the domain. Depending on the complexity of $a(x, y)$ and $b(x, y)$, these equations may be solved explicitly or implicitly. The solution to this system typically provides a family of characteristic curves parameterized by constants.

5. Solve for u along the characteristic curves.

Using the solution $x(t)$ and $y(t)$ from the previous step, solve the third characteristic equation:

$$\frac{du}{dt} = c(x, y, u).$$

With x and y now parameterized by t , this equation reduces to a standard ordinary differential equation for $u(t)$, which can often be solved directly using standard integration techniques.

6. Determine the general solution.

The solution $u(t)$ obtained from the previous step provides an expression for u in terms of t . However, since x and y are also functions of t , the solution must be rewritten in terms of the original variables x and y . This often involves expressing the solution implicitly or, when possible, solving explicitly for $u(x, y)$.

7. Apply initial or boundary conditions.

If an initial or boundary condition is provided, such as:

$$u(x_0, y_0) = u_0,$$

use this information to determine any remaining constants of integration from the characteristic equations. The initial conditions typically define a curve in the (x, y) -plane, and the characteristic curves must be matched to these initial values.

8. Express the final solution in terms of x and y .

Once the characteristic system is fully solved, the final solution to the PDE should be written in terms of x and y . In many cases, the solution is obtained implicitly as an equation relating x , y , and u . If explicit inversion is possible, solve for $u(x, y)$.

The method of characteristics is particularly useful for solving transport equations, wave equations, and Hamilton-Jacobi equations. It provides deep insight into how solutions propagate along characteristic curves, revealing important structural properties of the underlying PDE.

▲ Example 6.6.1 Solve the PDE:

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = u$$

Solution

1. Formulate the characteristic equations.

The PDE is given in the form:

$$a(x, y) \frac{\partial u}{\partial x} + b(x, y) \frac{\partial u}{\partial y} = c(x, y, u),$$

where $a(x, y) = x$, $b(x, y) = y$, and $c(x, y, u) = u$. The characteristic equations are:

$$\frac{dx}{dt} = x, \quad \frac{dy}{dt} = y, \quad \frac{du}{dt} = u.$$

2. Solve for the characteristic curves.

The first two equations,

$$\frac{dx}{dt} = x, \quad \frac{dy}{dt} = y,$$

are separable and solve to:

$$x = C_1 e^t, \quad y = C_2 e^t.$$

These describe exponential growth along characteristic curves.

3. Solve for u along characteristics.

The third equation,

$$\frac{du}{dt} = u,$$

has the general solution:

$$u = C_3 e^t.$$

4. Eliminate the parameter t .

Since $x = C_1 e^t$ and $y = C_2 e^t$, we take the ratio:

$$\frac{x}{C_1} = e^t, \quad \frac{y}{C_2} = e^t.$$

Setting them equal gives:

$$\frac{x}{C_1} = \frac{y}{C_2},$$

which simplifies to the characteristic equation:

$$\frac{x}{y} = \frac{C_1}{C_2} = C.$$

This means x/y is constant along characteristics.

5. Express the solution in terms of x and y .

Since $u = C_3 e^t$ and we found $e^t = x/C_1$, we substitute:

$$u = C_3 \frac{x}{C_1}.$$

Since C_1 and C_2 are related by $C = C_1/C_2$, the general solution can be written in terms of an arbitrary function:

$$u(x, y) = F\left(\frac{x}{y}\right) x.$$

Thus, the general solution is any function of x/y multiplied by x .

6. Verification.

Differentiating:

$$\frac{\partial u}{\partial x} = F\left(\frac{x}{y}\right) + xF'\left(\frac{x}{y}\right) \frac{1}{y}, \quad \frac{\partial u}{\partial y} = xF'\left(\frac{x}{y}\right) \frac{-x}{y^2}.$$

Substituting into the original PDE:

$$x\left(F + \frac{x}{y}F'\right) + y\left(-\frac{x^2}{y^2}F'\right) = xF + x\frac{x}{y}F' - x\frac{x}{y}F' = xF.$$

Since this matches $u = xF(x/y)$, the solution is verified.

▲ Example 6.6.2 Solve the nonlinear first-order PDE:

$$u \frac{\partial u}{\partial x} + (x + u) \frac{\partial u}{\partial y} = x$$

Solution**1. Formulate the characteristic equations.**

The PDE is given in the form:

$$a(x, y, u) \frac{\partial u}{\partial x} + b(x, y, u) \frac{\partial u}{\partial y} = c(x, y, u),$$

where $a(x, y, u) = u$, $b(x, y, u) = x + u$, and $c(x, y, u) = x$. The characteristic equations are:

$$\frac{dx}{ds} = u, \quad \frac{dy}{ds} = x + u, \quad \frac{du}{ds} = x$$

2. Solve for x and u .

From the first and third equations:

$$\frac{dx}{ds} = u, \quad \frac{du}{ds} = x$$

Differentiating the first equation with respect to s :

$$\frac{d^2x}{ds^2} = \frac{du}{ds} = x$$

This yields the second-order ODE:

$$\frac{d^2x}{ds^2} = x$$

with general solution:

$$x(s) = Ae^s + Be^{-s}$$

where A and B are constants. Consequently:

$$u(s) = \frac{dx}{ds} = Ae^s - Be^{-s}$$

3. Identify the first invariant.

Observe that:

$$\begin{aligned} x^2 - u^2 &= (Ae^s + Be^{-s})^2 - (Ae^s - Be^{-s})^2 \\ &= 4AB \end{aligned}$$

Therefore, $I_1 = x^2 - u^2$ is constant along characteristics.

4. Solve for y .

From $\frac{dy}{ds} = x + u$:

$$\frac{dy}{ds} = (Ae^s + Be^{-s}) + (Ae^s - Be^{-s}) = 2Ae^s$$

Integrating:

$$y(s) = 2Ae^s + C$$

where C is constant.

5. Identify the second invariant.

Note that $x + u = 2Ae^s$, therefore:

$$I_2 = y - (x + u)$$

is also constant along characteristics.

6. Express the general solution.

Since both I_1 and I_2 are constant along characteristics, the general solution has the implicit form:

$$\Phi(x^2 - u^2, y - x - u) = 0$$

where Φ is an arbitrary function of two variables.

Green's Functions

Green's functions provide a powerful method for solving inhomogeneous linear partial differential equations (PDEs). Their derivation is rooted in the linearity of the differential operator and the superposition principle, revealing deep connections between the abstract theory of distributions and concrete physical phenomena.

Consider a linear PDE of the form

$$L u(x) = f(x),$$

where L is a linear differential operator and $f(x)$ is a given forcing function. Because L is linear, if we can determine the response of the system to a point source, we can then use superposition to construct the solution for any arbitrary $f(x)$. A *point source* is ideally modeled by the Dirac delta function $\delta(x - \xi)$, defined by its action on a test function $\phi(x)$:

$$\int_{-\infty}^{\infty} \delta(x - \xi) \phi(x) dx = \phi(\xi).$$

The delta function serves as an idealized impulse, and its response under the operator L forms the basic building block for the solution.

The *Green's function* $G(x, \xi)$ associated with the operator L is defined by the equation

$$L G(x, \xi) = \delta(x - \xi),$$

subject to the same boundary conditions imposed on the original problem. In this context, $G(x, \xi)$ represents the response at the point x due to an impulse applied at the point ξ .

Once $G(x, \xi)$ is known, the solution to the inhomogeneous PDE can be written as

$$u(x) = \int G(x, \xi) f(\xi) d\xi.$$

To see why this is valid, apply the operator L to $u(x)$:

$$L u(x) = L \left(\int G(x, \xi) f(\xi) d\xi \right).$$

Assuming conditions that justify interchanging L and the integral, and using the defining property of $G(x, \xi)$, we have

$$L u(x) = \int [L G(x, \xi)] f(\xi) d\xi = \int \delta(x - \xi) f(\xi) d\xi = f(x).$$

Thus, $u(x)$ indeed satisfies the original PDE.

Several aspects underscore the depth of the Green's function method:

- **Superposition and Linearity:** The linearity of L ensures that the response to any general source $f(x)$ can be constructed as the superposition (integral) of responses to point sources. In essence, Green's functions act as the *impulse response* of the system.
- **Distributional Framework:** The equation

$$L G(x, \xi) = \delta(x - \xi)$$

is understood in the sense of distributions (generalized functions). This framework rigorously accommodates the delta function and allows us to handle cases where classical solutions may not exist.

- **Boundary Conditions:** The Green's function is constructed to satisfy the same boundary conditions as the original PDE. For instance, if $u(x)$ is required to vanish on the boundary $\partial\Omega$, then typically

$$G(x, \xi) = 0 \quad \text{for } x \in \partial\Omega.$$

- **Self-Adjointness and Symmetry:** When the operator L is self-adjoint, the Green's function inherits a symmetry:

$$G(x, \xi) = G(\xi, x).$$

This property not only simplifies computations but also has important physical implications in areas such as quantum mechanics and potential theory.

Extending to Higher Dimensions

The formulation above generalizes naturally to problems in two or more dimensions. Let $\mathbf{x}$ and $\boldsymbol{\xi}$ denote points in $\mathbb{R}^n$ (with $n \geq 2$). Then a typical linear PDE is written as

$$L u(\mathbf{x}) = f(\mathbf{x}), \quad \mathbf{x} \in \Omega \subset \mathbb{R}^n,$$

subject to appropriate boundary conditions on $\partial\Omega$. The corresponding Green's function $G(\mathbf{x}, \boldsymbol{\xi})$ is defined by

$$L G(\mathbf{x}, \boldsymbol{\xi}) = \delta(\mathbf{x} - \boldsymbol{\xi}),$$

where the Dirac delta function in n dimensions satisfies

$$\int_{\mathbb{R}^n} \delta(\mathbf{x} - \boldsymbol{\xi}) \phi(\mathbf{x}) d\mathbf{x} = \phi(\boldsymbol{\xi}).$$

Once $G(\mathbf{x}, \boldsymbol{\xi})$ is determined—using methods such as separation of variables, eigenfunction expansions, or integral transforms—the solution is given by

$$u(\mathbf{x}) = \int_{\Omega} G(\mathbf{x}, \boldsymbol{\xi}) f(\boldsymbol{\xi}) d\boldsymbol{\xi}.$$

This integral sums the contributions of point sources throughout the domain Ω , yielding the complete solution $u(\mathbf{x})$. For instance, in three-dimensional electrostatics, the Poisson equation for the electric potential $\phi(\mathbf{x})$ is

$$-\Delta\phi(\mathbf{x}) = \rho(\mathbf{x}),$$

with appropriate boundary conditions. In free space, the Green's function for the Laplacian is

$$G(\mathbf{x}, \boldsymbol{\xi}) = \frac{1}{4\pi |\mathbf{x} - \boldsymbol{\xi}|},$$

and hence the potential is given by

$$\phi(\mathbf{x}) = \int_{\mathbb{R}^3} \frac{1}{4\pi |\mathbf{x} - \boldsymbol{\xi}|} \rho(\boldsymbol{\xi}) d\boldsymbol{\xi}.$$

In other geometries or with different boundary conditions, the construction of $G(\mathbf{x}, \boldsymbol{\xi})$ may be more involved, but the underlying principle remains the same.

Below is a practical, step-by-step guide for applying the Green's function method to solve linear PDEs in any dimension. We also mention some known Green's functions for common classes of PDEs.

1. Express the PDE in Operator Form.

Write the PDE in the form

$$L u(\mathbf{x}) = f(\mathbf{x}), \quad \mathbf{x} \in \Omega,$$

where L is a linear differential operator (such as the Laplacian $-\Delta$ or the heat operator $\partial_t - \Delta$), and $f(\mathbf{x})$ is the forcing function. Clearly specify the domain $\Omega \subset \mathbb{R}^n$ and the boundary conditions on $\partial\Omega$. For example:

- **Poisson's Equation:** $-\Delta u(\mathbf{x}) = f(\mathbf{x})$ in Ω with, say, $u(\mathbf{x}) = 0$ on $\partial\Omega$.
- **Heat Equation:** $u_t(x, t) - \Delta u(x, t) = f(x, t)$ in $\Omega \times (0, \infty)$ with specified initial and boundary conditions.

2. Represent the Forcing Term via the Delta Function.

Recognize that a point source in $\mathbb{R}^n$ is modeled by the Dirac delta function $\delta(\mathbf{x} - \boldsymbol{\xi})$, which satisfies

$$\int_{\mathbb{R}^n} \delta(\mathbf{x} - \boldsymbol{\xi}) \phi(\mathbf{x}) d\mathbf{x} = \phi(\boldsymbol{\xi})$$

for any test function ϕ . This impulse response is the basic building block for constructing the full solution.

3. Define the Green's Function.

Define $G(\mathbf{x}, \boldsymbol{\xi})$ as the solution to

$$L G(\mathbf{x}, \boldsymbol{\xi}) = \delta(\mathbf{x} - \boldsymbol{\xi}),$$

subject to the same boundary conditions as $u(\mathbf{x})$. Here, $\mathbf{x}$ is the field point (where the solution is observed) and $\boldsymbol{\xi}$ is the source point. Known examples include:

4. Solve for the Green's Function.

Depending on the operator L and the geometry of Ω , solve the equation

$$L G(\mathbf{x}, \boldsymbol{\xi}) = \delta(\mathbf{x} - \boldsymbol{\xi})$$

using techniques such as separation of variables, eigenfunction expansions, or integral transforms. For many standard problems, the Green's function is already known from the literature. In more complex domains, methods like the method of images or numerical techniques might be employed. Below are some common solutions:

Table 6.2: *Comprehensive Table of Common PDE Operators and Their Green's Functions. In these expressions, the notation $G(x, t; \xi, 0)$ (or $G(\mathbf{x}, \boldsymbol{\xi})$) denotes the Green's function where the first set of variables indicates the field point (where the solution is observed) and the second set indicates the source point (where the impulse is applied).*

PDE/Operator	Green's Function
Laplacian in $\mathbb{R}^3$	$G(\mathbf{x}, \boldsymbol{\xi}) = \frac{1}{4\pi \mathbf{x} - \boldsymbol{\xi} }$
Laplacian in $\mathbb{R}^2$	$G(\mathbf{x}, \boldsymbol{\xi}) = -\frac{1}{2\pi} \ln \mathbf{x} - \boldsymbol{\xi} $
Heat Equation in $\mathbb{R}$	$G(x, t; \xi, 0) = \frac{1}{\sqrt{4\pi t}} \exp\left(-\frac{(x - \xi)^2}{4t}\right), \quad t > 0$
Wave Equation in $\mathbb{R}^3$	$G(t, \mathbf{x}; t', \boldsymbol{\xi}) = \frac{\delta\left(t - t' - \frac{ \mathbf{x} - \boldsymbol{\xi} }{c}\right)}{4\pi \mathbf{x} - \boldsymbol{\xi} }$
Free Schrödinger Equation in 1D	$G(x, t; \xi, 0) = \sqrt{\frac{m}{2\pi i \hbar t}} \exp\left(\frac{im(x - \xi)^2}{2\hbar t}\right)$
Helmholtz Equation in $\mathbb{R}^3$	$G(\mathbf{x}, \boldsymbol{\xi}) = \frac{e^{ik \mathbf{x} - \boldsymbol{\xi} }}{4\pi \mathbf{x} - \boldsymbol{\xi} }$

5. Construct the Solution via Superposition.

With $G(\mathbf{x}, \boldsymbol{\xi})$ determined, the solution to the original PDE is obtained by superposing the responses to point sources:

$$u(\mathbf{x}) = \int_{\Omega} G(\mathbf{x}, \boldsymbol{\xi}) f(\boldsymbol{\xi}) d\boldsymbol{\xi}.$$

In time-dependent problems, the integration may include time as well, e.g.,

$$u(x, t) = \int_{\mathbb{R}} G(x, t; \xi, 0) g(\xi) d\xi,$$

where $g(\xi)$ represents the initial condition.

6. Verify the Solution.

Finally, check that the solution $u(\mathbf{x})$ satisfies both the PDE and the boundary conditions. Applying L to $u(\mathbf{x})$ and using the defining property of G , we obtain

$$L u(\mathbf{x}) = \int_{\Omega} [L G(\mathbf{x}, \boldsymbol{\xi})] f(\boldsymbol{\xi}) d\boldsymbol{\xi} = \int_{\Omega} \delta(\mathbf{x} - \boldsymbol{\xi}) f(\boldsymbol{\xi}) d\boldsymbol{\xi} = f(\mathbf{x}).$$

This procedure, while conceptually similar across dimensions, becomes practical when applied to well-studied operators. Many classical PDEs—such as Laplace’s, Poisson’s, and the heat equation—have known Green’s functions, greatly simplifying the solution process. For more complex or irregular domains, the same principles apply, though one might need to resort to numerical or approximate methods to construct $G(\mathbf{x}, \boldsymbol{\xi})$.

▲ Example 6.6.3 Consider the three-dimensional Poisson equation in free space with a Gaussian source:

$$-\Delta u(\mathbf{x}) = e^{-|\mathbf{x}|^2}, \quad \mathbf{x} \in \mathbb{R}^3$$

with the condition that $u(\mathbf{x})$ vanishes at infinity. Here, $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ is the Laplacian operator. Find the solution using the known Green’s function.

Solution Before diving into the mathematics, let’s understand what we’re solving: we have a source that decays exponentially in all directions (like heat spreading from a hot spot), and we want to find how this source influences the surrounding space. We’ll approach this systematically:

1. Identify the Green’s function.

For the 3D Laplacian in free space, the Green’s function is:

$$G(\mathbf{x}, \boldsymbol{\xi}) = \frac{1}{4\pi|\mathbf{x} - \boldsymbol{\xi}|}$$

This function represents the response to a unit point source, where:

- $\mathbf{x} = (x, y, z)$ is the observation point
- $\boldsymbol{\xi} = (\xi, \eta, \zeta)$ is the source point

2. **Write the solution integral.**

By superposition, our solution is:

$$u(\mathbf{x}) = \int_{\mathbb{R}^3} G(\mathbf{x}, \boldsymbol{\xi}) f(\boldsymbol{\xi}) d\boldsymbol{\xi}$$

Substituting our Green's function and source:

$$u(\mathbf{x}) = \frac{1}{4\pi} \int_{\mathbb{R}^3} \frac{e^{-|\boldsymbol{\xi}|^2}}{|\mathbf{x} - \boldsymbol{\xi}|} d\boldsymbol{\xi}$$

3. **Exploit symmetry.**

Notice that our source $e^{-|\boldsymbol{\xi}|^2}$ is radially symmetric. This suggests our solution will also be radially symmetric:

$$u(\mathbf{x}) = u(r), \quad \text{where } r = |\mathbf{x}| = \sqrt{x^2 + y^2 + z^2}$$

4. **Use potential theory result.**

A key result from potential theory states that:

$$\int_{\mathbb{R}^3} \frac{e^{-|\boldsymbol{\xi}|^2}}{|\mathbf{x} - \boldsymbol{\xi}|} d\boldsymbol{\xi} = \pi^{3/2} \frac{\text{erf}(r)}{r}$$

where $\text{erf}(r)$ is the error function:

$$\text{erf}(r) = \frac{2}{\sqrt{\pi}} \int_0^r e^{-t^2} dt$$

5. **Write the final solution.**

Combining our results:

$$u(\mathbf{x}) = \frac{\sqrt{\pi}}{4} \frac{\text{erf}(|\mathbf{x}|)}{|\mathbf{x}|}$$

Several key observations about our solution:

- The solution inherits the radial symmetry of the source
- The error function $\text{erf}(r)$ approaches 1 as $r \rightarrow \infty$

- The $1/|\mathbf{x}|$ decay is characteristic of 3D potential problems
- Near the origin, $\text{erf}(r) \approx \frac{2}{\sqrt{\pi}}r$, so $u(0)$ is finite

The solution reveals how the Gaussian source influences space: near the source, the influence varies smoothly, while far from the source, it decays like $1/|\mathbf{x}|$, characteristic of three-dimensional potential problems.

Finding a Green's Function the Hard Way

Thus far, we've worked with known Green's functions, inserting them into our solution machinery like well-crafted tools. But how are these fundamental objects discovered in the first place? Let's embark on a journey to derive one of quantum mechanics' most important Green's functions from scratch: the propagator for the free Schrödinger equation. This derivation will showcase the interplay between Fourier analysis, differential equations, and the delicate handling of distributions that characterizes modern mathematical physics.

We'll see how the machinery of Fourier transforms can convert our challenging PDE into a manageable ODE, and how careful attention to initial conditions leads us to a result that has profound implications for quantum mechanics. The techniques we develop here extend far beyond this specific example, forming a template for deriving Green's functions in many other contexts.

▲ **Example 6.6.4** Find the Green's function (propagator) for the one-dimensional free Schrödinger equation:

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2}$$

where $\hbar$ is the reduced Planck's constant and m is the particle mass.

Solution Let's solve this systematically using Fourier transform methods:

1. Set up the propagator equation.

The Green's function $K(x, t; x', 0)$ must satisfy:

$$i\hbar \frac{\partial K}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 K}{\partial x^2}$$

with initial condition:

$$K(x, 0; x', 0) = \delta(x - x')$$

This delta function initial condition ensures our Green's function represents the system's response to a perfectly localized disturbance.

2. Transform to Fourier space.

Define the spatial Fourier transform:

$$\tilde{K}(k, t; x', 0) = \int_{-\infty}^{\infty} K(x, t; x', 0) e^{-ikx} dx$$

Under this transform, spatial derivatives become:

$$\frac{\partial^2}{\partial x^2} \rightarrow -k^2$$

Our equation becomes:

$$i\hbar \frac{\partial \tilde{K}}{\partial t} = \frac{\hbar^2 k^2}{2m} \tilde{K}$$

3. Solve the resulting ODE.

For each k , we have:

$$\frac{\partial \tilde{K}}{\partial t} = -i \frac{\hbar k^2}{2m} \tilde{K}$$

with solution:

$$\tilde{K}(k, t; x', 0) = A(k) e^{-i \frac{\hbar k^2}{2m} t}$$

where $A(k)$ will be determined by initial conditions.

4. Apply the initial condition.

The Fourier transform of $\delta(x - x')$ is $e^{-ikx'}$, so:

$$A(k) = e^{-ikx'}$$

Therefore:

$$\tilde{K}(k, t; x', 0) = e^{-ikx'} e^{-i \frac{\hbar k^2}{2m} t}$$

5. Perform the inverse transform.

Our Green's function is:

$$K(x, t; x', 0) = \frac{1}{2\pi} \int_{-\infty}^{\infty} e^{ik(x-x')} e^{-i \frac{\hbar k^2}{2m} t} dk$$

This Gaussian integral evaluates to:

$$K(x, t; x', 0) = \sqrt{\frac{m}{2\pi i \hbar t}} e^{i \frac{m(x-x')^2}{2\hbar t}}$$

Several profound observations about our solution:

- The magnitude $|K|$ spreads out like $1/\sqrt{t}$, reflecting quantum uncertainty
- The phase factor oscillates more rapidly for larger $|x - x'|$

- The solution conserves probability: $\int |K|^2 dx = 1$
- The classical limit ($\hbar \rightarrow 0$) concentrates the propagation along classical paths

This Green's function, known as the free particle propagator, is fundamental to path integral formulations of quantum mechanics and quantum field theory.

Formulas and Definitions

Important Vocabulary

- **Ordinary Differential Equation (ODE):** An equation containing derivatives of an unknown function with respect to a single independent variable.
- **Order:** The highest derivative appearing in the differential equation.
- **Linear ODE:** A differential equation where the unknown function and its derivatives appear only to the first power and are not multiplied together.
- **Homogeneous Equation:** A linear differential equation where there is no forcing term (right-hand side is zero).
- **Characteristic Equation:** The algebraic equation obtained by substituting e^{rx} into a linear differential equation with constant coefficients.
- **Integrating Factor:** A function that, when multiplied through a differential equation, makes it exact or separable.
- **Initial Value Problem (IVP):** A differential equation together with initial conditions specifying the value of the solution and its derivatives at a point.
- **Boundary Value Problem (BVP):** A differential equation with conditions specified at multiple points.
- **Existence and Uniqueness:** Theorems specifying when solutions exist and are unique based on continuity of functions and derivatives.
- **Wronskian:** A determinant that tests linear independence of solutions.
- **Particular Solution:** A specific solution to a non-homogeneous equation.
- **Complementary Function:** The general solution to the homogeneous equation.
- **Laplace Transform:** An integral transform that converts differential equations to algebraic equations.
- **Characteristic Equations:** System of ordinary differential equations derived from a PDE that describe curves along which the solution propagates.
- **Green's Function:** A fundamental solution to an inhomogeneous differential equation for a point source (delta function) input, used to construct solutions for arbitrary forcing functions.
- **First-Order PDE:** A partial differential equation involving only first derivatives of the unknown function.

- **Linear PDE:** A partial differential equation where the unknown function and its derivatives appear to the first power only.
- **Characteristic Curves:** Paths in the domain along which information propagates in the solution of a PDE.
- **Delta Function:** A generalized function $\delta(x)$ that equals infinity at $x = 0$, zero elsewhere, with unit integral.
- **Forcing Function:** The inhomogeneous term in a differential equation representing external influences or sources.
- **Self-Adjoint Operator:** A differential operator L whose Green's function exhibits symmetry: $G(x, \xi) = G(\xi, x)$.

Key Formulas

First-Order Linear ODEs:

$$\frac{dy}{dx} + P(x)y = Q(x)$$

Solution Method:

$$\mu(x) = e^{\int P(x)dx}$$

$$y = \frac{1}{\mu(x)} \left[\int \mu(x)Q(x)dx + C \right]$$

Separable Equations:

$$\frac{dy}{dx} = f(x)g(y)$$

Solution Method:

$$\int \frac{dy}{g(y)} = \int f(x)dx + C$$

Bernoulli Equations:

$$\frac{dy}{dx} + P(x)y = Q(x)y^n$$

Substitution:

$$v = y^{1-n}$$

Second-Order Linear ODEs:

$$ay'' + by' + cy = f(x)$$

Characteristic Equation:

$$ar^2 + br + c = 0$$

Solution Forms: Distinct Real Roots (r_1, r_2) :

$$y = c_1 e^{r_1 x} + c_2 e^{r_2 x}$$

Repeated Root (r_1) :

$$y = c_1 e^{r_1 x} + c_2 x e^{r_1 x}$$

Complex Roots $(\alpha \pm i\beta)$:

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Laplace Transform Properties: Definition:

$$\mathcal{L}\{f(t)\} = \int_0^\infty e^{-st} f(t) dt = F(s)$$

Derivatives:

$$\mathcal{L}\{f'(t)\} = sF(s) - f(0)$$

$$\mathcal{L}\{f''(t)\} = s^2 F(s) - sf(0) - f'(0)$$

Method of Undetermined Coefficients: Solution via Green's Function:

For $f(x)$ of form:

- Polynomial $\rightarrow$ try polynomial
- $e^{ax} \rightarrow$ try Ae^{ax}
- $\sin(ax)$ or $\cos(ax) \rightarrow$ try $A\cos(ax) + B\sin(ax)$

Variation of Parameters: For $y'' + p(x)y' + q(x)y = g(x)$:

$$y_p = -y_1 \int \frac{y_2 g}{W} dx + y_2 \int \frac{y_1 g}{W} dx$$

where W is the Wronskian:

$$W = y_1 y_2' - y_2 y_1'$$

System Properties: Wronskian Test:

$W(y_1, y_2)(x) \neq 0 \implies$ Linear Independence

General Solution:

$$y = y_h + y_p$$

where y_h is homogeneous solution and y_p is particular solution.

First-Order PDE Standard Form:

$$a(x, y) \frac{\partial u}{\partial x} + b(x, y) \frac{\partial u}{\partial y} = c(x, y, u)$$

Characteristic Equations:

$$\frac{dx}{dt} = a(x, y), \quad \frac{dy}{dt} = b(x, y), \quad \frac{du}{dt} = c(x, y, u)$$

Green's Function Definition:

$$L G(x, \xi) = \delta(x - \xi)$$

$$u(x) = \int G(x, \xi) f(\xi) d\xi$$

Delta Function Property:

$$\int_{-\infty}^{\infty} \delta(x - \xi) \phi(x) dx = \phi(\xi)$$

PDE Solution Methods: Method of Characteristics:

$$\frac{dy}{dx} = \frac{b(x, y)}{a(x, y)}$$

Green's Function Solution:

$$u(\mathbf{x}) = \int_{\Omega} G(\mathbf{x}, \boldsymbol{\xi}) f(\boldsymbol{\xi}) d\boldsymbol{\xi}$$

Common PDEs: Poisson's Equation:

$$-\Delta u = f$$

Heat Equation:

$$\frac{\partial u}{\partial t} = \alpha \Delta u + f$$

Wave Equation:

$$\frac{\partial^2 u}{\partial t^2} = c^2 \Delta u + f$$

Schrödinger Equation:

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \Delta \psi + V\psi$$

For tables for Laplace transforms or Green's functions please see earlier in the chapter or Appendix E.

Practice Problems

- Determine whether each of the following equations is an ordinary differential equation (ODE) or a partial differential equation (PDE):
 - $\frac{dy}{dx} = 3x^2$.
 - $\frac{\partial u}{\partial t} = 4 \frac{\partial u}{\partial x}$.
 - $\frac{d^2y}{dx^2} + 5y = \sin(x)$.
- Identify the order of each of the following differential equations:
 - $\frac{d^2y}{dx^2} + 4\frac{dy}{dx} - 5y = 0$.
 - $\frac{d^3y}{dx^3} - 2\frac{d^2y}{dx^2} + y = \cos(x)$.
 - $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$.
- Classify each of the following equations as linear or nonlinear:
 - $\frac{dy}{dx} = y^2 - \sin(x)$.
 - $\frac{dy}{dx} + y \cos(x) = e^x$.
 - $\frac{d^2y}{dx^2} + 3y\frac{dy}{dx} = x$.
- Determine whether each of the following differential equations is homogeneous or non-homogeneous:
 - $\frac{d^2y}{dx^2} - 3\frac{dy}{dx} + 2y = e^x$.
 - $\frac{dy}{dx} + 2y = 0$.
 - $\frac{d^2y}{dx^2} + y = \sin(x)$.
- Identify whether each of the following ODEs has constant coefficients:
 - $x\frac{d^2y}{dx^2} + 2\frac{dy}{dx} - y = 0$.
 - $3\frac{d^2y}{dx^2} + 2\frac{dy}{dx} - y = 0$.
 - $\frac{d^2y}{dx^2} - 4y = 0$.
- Verify that the explicit function $y(x) = e^{2x}$ is a solution of the differential equation $y'' - 4y = 0$.
- Verify that the explicit function $y(x) = 3e^{-x} + 2$ is a solution of the differential equation $y' + y = 2$.
- Verify that the implicit relation $x^2 + y^2 = 1$ is a solution of the differential equation $\frac{dy}{dx} = -\frac{x}{y}$. (Differentiate $x^2 + y^2 = 1$ implicitly with respect to x and solve for $\frac{dy}{dx}$.)
- Verify that the implicit relation $\ln|y| + y = x$ is a solution of the differential equation $\left(\frac{1}{y} + 1\right)y' = 1$.
- Solve the separable ODE

$$\frac{dy}{dx} = \frac{2x}{y}, \quad y(1) = 2.$$
- Solve the separable ODE

$$\frac{dy}{dx} = x e^y.$$
- Solve the linear first-order ODE

$$\frac{dy}{dx} + 2y = \sin(x).$$
- Solve the exact differential equation

$$(2xy + y^2)dx + (x^2 + 2xy)dy = 0.$$
- Solve the Bernoulli equation

$$\frac{dy}{dx} - y = e^x y^2.$$
- Solve the initial value problem

$$\frac{dy}{dx} = \frac{y}{x} + x^2, \quad y(1) = 2.$$
- Solve the separable ODE

$$\frac{dy}{dx} = 3\sqrt{y}, \quad y(4) = 9.$$

17. Solve the linear ODE (for
- $x > 0$
-)

$$\frac{dy}{dx} - \frac{y}{x} = x^2.$$

18. Solve the separable ODE

$$\frac{dy}{dx} = \frac{2x}{1+y^2}.$$

19. Verify that

$$y(x) = e^x - 1$$

is a solution of

$$\frac{dy}{dx} - y = e^x.$$

20. Determine if the ODE

$$(y \cos(x) - \sin(x)) dx + \cos(x) dy = 0$$

is exact; if so, find its general solution.

21. Solve the Bernoulli equation

$$\frac{dy}{dx} + y = x y^2.$$

22. Solve the ODE

$$\frac{dy}{dx} = \frac{xy}{1+x^2}.$$

23. Solve the initial value problem

$$\frac{dy}{dx} = \frac{y}{x} + \ln(x), \quad y(1) = 0.$$

24. Solve the ODE

$$\frac{dy}{dx} + \frac{2}{x}y = \sin(x), \quad x > 0.$$

25. Solve the ODE

$$\frac{dy}{dx} = \frac{y-x}{y+x}.$$

26. Solve the ODE

$$\frac{dy}{dx} = \frac{e^x}{1+y}.$$

27. Find the general solution of

$$y'' - 3y' + 2y = 0.$$

28. Find the general solution of

$$y'' + y = 0.$$

29. Find the general solution of

$$y'' + 4y' + 4y = 0.$$

30. Solve the third-order homogeneous ODE

$$y''' - 3y'' + 3y' - y = 0.$$

31. Find the general solution of the fourth-order homogeneous equation

$$y^{(4)} - 5y''' + 8y'' - 4y' = 0.$$

32. Using the method of undetermined coefficients, find a particular solution of

$$y'' - y = e^x.$$

33. Using undetermined coefficients, find a particular solution of

$$y'' - y = e^{-x}.$$

34. Use variation of parameters to find a particular solution of

$$y'' + 2y' + y = x.$$

35. Use variation of parameters to find a particular solution of

$$y'' - 2y' + y = xe^x.$$

36. Solve the initial value problem

$$y'' + y = \sin x, \quad y(0) = 0, \quad y'(0) = 1.$$

37. Solve the initial value problem

$$y'' + 2y' + 5y = 0, \quad y(0) = 2, \quad y'(0) = 0.$$

38. Find a particular solution by undetermined coefficients for

$$y'' + 4y = \cos(2x).$$

39. Find the general solution of the fourth-order ODE

$$y^{(4)} - 4y'' + 4y = 0.$$

40. Use undetermined coefficients to find a particular solution of

$$y''' - 2y'' + y' = e^x.$$

41. Find the general solution of the resonant equation

$$y'' - 2y' + y = 3e^x.$$

42. Solve the third-order ODE

$$y^{(3)} - 4y'' + 5y' - 2y = 0.$$

43. Find a particular solution of the resonant equation

$$y'' + 2y' + y = e^{-x}$$

using the method of undetermined coefficients.

44. Solve the ODE with complex forcing:

$$y'' + y = e^{ix}.$$

(Then extract the real part as the physical solution.)

45. Find the general solution of

$$y^{(4)} - 2y'' + y = 0.$$

46. Solve the third-order non-homogeneous ODE

$$y''' + 3y'' + 3y' + y = e^{-x}.$$

47. Use variation of parameters to find a particular solution of

$$y'' + 2y' + y = x.$$

48. Use variation of parameters to find a particular solution of

$$y'' - 2y' + y = xe^x.$$

49. Find a particular solution of

$$y'' + y = \tan x$$

using variation of parameters.

50. Use variation of parameters to find a particular solution of

$$y'' + y = \sec^2 x.$$

51. Solve the non-homogeneous integro-differential equation

$$y''(t) + y(t) = \int_0^t y(s) ds$$

given $y(0) = 0, \quad y'(0) = 1.$

52. Solve the ODE with discontinuous forcing using Laplace transforms:

$$y'' + 2y' + 5y = u(t - 3)$$

given $y(0) = 0, \quad y'(0) = 1.$, where $u(t - 3)$ is the Heaviside step function

53. Solve the ODE with impulsive forcing using Laplace transforms:

$$y'' - y = \delta(t - 2), \quad y(0) = 0, \quad y'(0) = 0,$$

where $\delta(t - 2)$ is the Dirac delta function.

54. Solve the ODE using Laplace transforms:

$$y'' + 3y' + 2y = t e^{-t}$$

given $y(0) = 1, \quad y'(0) = 0.$

55. Solve the ODE with a discontinuous forcing function:

$$y'' + y = u(t - \pi), \quad y(0) = 0, \quad y'(0) = 0.$$

56. Using Laplace transforms, solve the integro-differential equation

$$y'(t) + \int_0^t y(s) ds = e^{-t}, \quad y(0) = 0.$$

57. Solve the ODE with a hard forcing function using Laplace transforms:

$$y'' + y = e^t \sin(2t)$$

given $y(0) = 0, \quad y'(0) = 0.$

58. Solve the ODE with impulsive forcing using Laplace transforms:

$$y'' + 2y' + y = \delta(t - 1) + \delta(t - 2)$$

given $y(0) = 0, \quad y'(0) = 0.$

59. Use variation of parameters to find a particular solution of the non-homogeneous ODE

$$y'' + 4y = e^{2x} \cos(3x).$$

60. Solve the heat equation

$$u_t = \alpha u_{xx}, \quad 0 < x < L, \quad t > 0,$$

with homogeneous Dirichlet boundary conditions

$$u(0, t) = u(L, t) = 0,$$

and initial condition

$$u(x, 0) = f(x),$$

using separation of variables and Fourier sine series.

61. Solve Laplace's equation

$$u_{xx} + u_{yy} = 0, \quad 0 < x < L, \quad 0 < y < H,$$

subject to $u(0, y) = u(L, y) = 0$, $u(x, 0) = 0$, $u(x, H) = g(x)$ via separation of variables and Fourier series.

62. Solve the one-dimensional wave equation

$$u_{tt} = c^2 u_{xx}, \quad 0 < x < L, \quad t > 0,$$

with fixed endpoints

$$u(0, t) = u(L, t) = 0,$$

and initial conditions

$$u(x, 0) = f(x), \quad u_t(x, 0) = g(x),$$

using separation of variables.

63. Use the method of characteristics to solve the first-order linear PDE

$$u_x + 2u_y = 3, \quad u(x, 0) = \sin x.$$

64. Use the method of characteristics to solve the nonlinear PDE

$$u u_x + u_y = 0, \quad u(x, 0) = x.$$

65. Solve the heat equation with a discontinuous forcing function:

$$u_t = u_{xx} + H\left(x - \frac{L}{2}\right), \quad 0 < x < L, \quad t > 0,$$

with boundary conditions

$$u(0, t) = u(L, t) = 0, \quad u(x, 0) = 0,$$

where $H(x - \frac{L}{2})$ is the Heaviside function.

66. Solve the inhomogeneous heat equation

$$u_t = u_{xx} + \sin(\pi x), \quad 0 < x < 1, \quad t > 0,$$

with $u(0, t) = u(1, t) = 0$ and $u(x, 0) = 0$ using Fourier series.

67. Solve Laplace's equation in polar coordinates in the disk
- $0 \leq r < a$
- :

$$u_{rr} + \frac{1}{r}u_r + \frac{1}{r^2}u_{\theta\theta} = 0,$$

with boundary condition

$$u(a, \theta) = f(\theta),$$

using separation of variables.

68. Solve the one-dimensional wave equation on
- $\mathbb{R}$

$$u_{tt} = c^2 u_{xx}, \quad -\infty < x < \infty, \quad t > 0,$$

with impulsive initial displacement

$$u(x, 0) = \delta(x), \quad u_t(x, 0) = 0,$$

by applying Fourier transforms.

69. Use the Green's function method to solve Poisson's equation on the interval:

$$-u_{xx} = f(x), \quad 0 < x < L,$$

with boundary conditions $u(0) = u(L) = 0$.

70. Determine the Fourier sine series of

$$f(x) = x(L - x),$$

for $0 < x < L$ to be used in solving the heat equation.

71. Solve the first-order PDE

$$u_t + u_x = 0, \quad u(x, 0) = e^{-x^2},$$

using the method of characteristics.

72. Solve the variable-coefficient PDE

$$x u_x + u_y = 0, \quad u(x, 0) = \ln(1 + x^2),$$

using the method of characteristics.

73. Solve the integro-differential PDE

$$u_t = u_{xx} + \int_0^x u(\xi, t) d\xi$$

given $0 < x < L$, $t > 0$ with $u(0, t) = u(L, t) = 0$ and initial condition $u(x, 0) = f(x)$ via separation of variables.

74. Solve the heat equation with Neumann boundary conditions:

$$u_t = u_{xx}$$

given $0 < x < L$, $t > 0$ with $u_x(0, t) = u_x(L, t) = 0$ and $u(x, 0) = f(x)$ using Fourier cosine series.

75. Solve the advection-diffusion equation

$$u_t + cu_x = D u_{xx}, \quad -\infty < x < \infty, \quad t > 0,$$

with initial condition $u(x, 0) = \phi(x)$ using Fourier transforms or separation of variables.

76. Use Laplace transforms to solve the heat equation on the semi-infinite domain:

$$u_t = u_{xx}, \quad x > 0, \quad t > 0,$$

with boundary condition $u(0, t) = g(t)$ and initial condition $u(x, 0) = 0$.

77. Use the method of characteristics to solve the transport equation with variable coefficients:

$$(1 + x^2)u_x + u_y = 0, \quad u(x, 0) = \frac{1}{1 + x^2}.$$

78. Solve the two-dimensional wave equation

$$u_{tt} = c^2(u_{xx} + u_{yy})$$

given $0 < x < L$, $0 < y < H$, $t > 0$ with homogeneous Dirichlet boundary conditions and initial conditions

$$u(x, y, 0) = f(x, y), \quad u_t(x, y, 0) = g(x, y),$$

using separation of variables and double Fourier series.

Chapter 7

Complex Variables

Introduction

Our journey through calculus has taken us from single-variable functions to multivariable calculus in $\mathbb{R}^2$ and $\mathbb{R}^3$. Now, we turn to complex variables—not an esoteric detour but a natural extension of multidimensional calculus. The complex plane provides a fresh perspective on two-dimensional space, reshaping our understanding of functions, derivatives, and integrals.

The complex plane is isomorphic to $\mathbb{R}^2$ – a fancy way of saying there is a one-to-one correspondence preserving mathematical operations. Writing $z = x + iy$ simply recasts (x, y) in a new framework. Yet, with complex multiplication and division, stunning geometric patterns emerge—multiplying by i corresponds to a 90-degree rotation, blending algebra and geometry in ways real analysis only hints at.

Concepts from vector calculus, such as line integrals and Green’s theorem, reappear in unexpected ways. Complex line integrals, though syntactically similar to real ones, exhibit striking properties—paths can often be deformed without altering the integral, mirroring conservative vector fields. The complex derivative, when it exists, ensures not just differentiability but analyticity, a property far stronger than in real analysis.

This journey promises both challenges and insights. Though the ideas may seem abstract at first, they build naturally on your knowledge, unifying and transforming familiar calculus concepts. Complex analysis reveals profound symmetries and patterns, making it a cornerstone of modern mathematics.

Lecture 7.1: The Complex Number System

Our journey through multivariable calculus has equipped us with tools to analyze functions of multiple real variables. However, there exists another number system—one that extends our familiar real numbers in profound and beautiful ways. This system of complex numbers not only helps us solve seemingly impossible equations but also provides deep insights into geometry, physics, and the very nature of mathematical structure itself.

Historical Development and Motivation

The story of complex numbers begins with a mathematical impossibility—or what seemed impossible at the time. Consider the equation:

$$x^2 + 1 = 0$$

In the real number system, this equation has no solution. After all, any real number squared must be non-negative, so adding 1 can never yield zero. Yet mathematicians of the 16th century, particularly in their work on cubic equations, found themselves forced to work with expressions involving square roots of negative numbers.



Figure 7.1: *Girolamo Cardano, engraving by R. Cooper Wellcome*

The Italian mathematician Girolamo Cardano, in his 1545 work *Ars Magna*, presented a method for solving cubic equations that inevitably led to these “impossible” quantities. Even more remarkably, by manipulating these supposedly impossible numbers according to normal algebraic rules, he obtained correct real number solutions. This suggested something profound: these “imaginary” quantities, while seemingly nonsensical, somehow encoded real mathematical truth.

The modern understanding of complex numbers emerged gradually through the work of mathematicians like Euler, Gauss, and Cauchy, who showed that far from being mere computational tools, complex numbers form a coherent and beautiful mathematical system with

deep connections to geometry, analysis, and physics. The term “imaginary” is somewhat misleading, as complex analysis has tangible physical applications. In my view, it was a missed opportunity not to adopt Gauss’s term “lateral numbers” instead, which more aptly reflects their geometric nature and maintains their tangibility.

Algebraic Form and Fundamental Properties

Let’s begin our formal development. We define the **imaginary unit** i as a quantity satisfying:

$$i^2 = -1$$

This single definition opens up an entirely new number system. A **complex number** is an expression of the form:

$$z = a + bi$$

where a and b are real numbers. We call a the **real part** and b the **imaginary part** of z , denoted:

$$\operatorname{Re}(z) = a \quad \text{and} \quad \operatorname{Im}(z) = b$$

Sometimes you might see $\Re(z)$ and $\Im(z)$ in other texts to represent real and imaginary parts respectively. The set of all complex numbers, denoted $\mathbb{C}$, forms a field—it satisfies all the familiar properties of arithmetic (commutativity, associativity, distributivity, etc.) and every non-zero complex number has a multiplicative inverse. This makes $\mathbb{C}$ a natural extension of $\mathbb{R}$, the real numbers. However, you may notice that one fundamental arithmetic property is absent: order. Unlike real numbers, complex numbers cannot be meaningfully compared using inequalities, as there is no natural way to define $a < b$ when $a, b \in \mathbb{C}$. We will explore this limitation in more detail later.

Indeed, we can view $\mathbb{R}$ as a subset of $\mathbb{C}$ by identifying each real number r with the complex number $r + 0i$. This embedding preserves all algebraic operations, meaning complex arithmetic agrees with real arithmetic when applied to real numbers.

The Complex Plane and Geometric Interpretation

One of the most powerful aspects of complex numbers is their geometric interpretation. Just as real numbers correspond to points on a line, complex numbers naturally correspond to points in a plane. This **complex plane** (also called the **Argand plane** after Jean-Robert Argand) represents the complex number $a + bi$ as the point with coordinates (a, b) .

This geometric view transforms algebraic operations into geometric ones:

- Addition of complex numbers corresponds to vector addition in the plane
- Multiplication by a real number corresponds to scaling
- Multiplication by i corresponds to a 90-degree counterclockwise rotation

To see why multiplication by i represents rotation, consider its effect:

$$i(a + bi) = ai + bi^2 = ai + b(-1) = -b + ai$$

This transforms the point (a, b) to the point $(-b, a)$, which is exactly what a 90-degree counterclockwise rotation does!




Figure 7.2: *Multiplication by i rotates a complex number 90° counterclockwise about the origin on the Argand plane.*

Basic Operations and Their Geometric Meaning

Let's examine how the basic operations on complex numbers manifest geometrically. For two complex numbers $z_1 = a + bi$ and $z_2 = c + di$:

1. Addition:

$$z_1 + z_2 = (a + c) + (b + d)i$$

This corresponds to vector addition in the plane, following the parallelogram law.

2. Subtraction:

$$z_1 - z_2 = (a - c) + (b - d)i$$

This corresponds to vector subtraction.

3. Multiplication:

$$z_1 z_2 = (ac - bd) + (ad + bc)i$$

This formula, while algebraically straightforward, hides a beautiful geometric interpretation we'll uncover when we study polar form.

Complex Conjugates and Absolute Values

For any complex number $z = a + bi$, its **complex conjugate**, denoted $\bar{z}$ or z^* , is defined as:

$$\bar{z} = a - bi$$

Geometrically, the conjugate reflects z across the real axis. This operation has several crucial properties:

- $\overline{(\bar{z})} = z$
- $\overline{z_1 + z_2} = \bar{z}_1 + \bar{z}_2$
- $\overline{z_1 z_2} = \bar{z}_1 \cdot \bar{z}_2$
- $z \cdot \bar{z} = a^2 + b^2$ (always a non-negative real number)

The last property leads us to define the **absolute value** or **modulus** of a complex number:

$$|z| = \sqrt{z \cdot \bar{z}} = \sqrt{a^2 + b^2}$$

This definition generalizes the absolute value of real numbers and has a clear geometric interpretation: $|z|$ is the distance from z to the origin in the complex plane. Now because the complex plane loses the property of number ordering, the modulus will be our primary tool in order to express distance and inequality between values and functions.

A fundamental concept in complex analysis is that an equation of the form $|z - z_0| = r$ describes a circle in the complex plane centered at z_0 with radius r . To understand why, let $z = x + yi$ and $z_0 = a + bi$. Then:

$$|z - z_0| = r \quad |(x + yi) - (a + bi)| = r \quad |(x - a) + (y - b)i| = r \quad \sqrt{(x - a)^2 + (y - b)^2} = r$$

This geometric interpretation makes the modulus an essential tool for describing regions in the complex plane. For instance, $|z| < r$ describes the interior of a circle centered at the origin with radius r , while $|z - z_1| < |z - z_2|$ describes all points closer to z_1 than to z_2 .




Figure 7.3: The equation $|z - z_0| = r$ describes all points z that are exactly distance r from the point z_0 , forming a circle.

▲ **Example 7.1.1** Express the complex fraction $\frac{2-3i}{4+i}$ in the form $a + bi$ where a and b are real numbers.

Solution We'll solve this by multiplying both numerator and denominator by the complex conjugate of the denominator.

1. Multiply numerator and denominator by $4 - i$:

$$\frac{2-3i}{4+i} = \frac{2-3i}{4+i} \cdot \frac{4-i}{4-i}$$

2. Multiply terms in numerator:

$$\begin{aligned} &= \frac{(2-3i)(4-i)}{(4+i)(4-i)} \\ &= \frac{8-2i-12i+3i^2}{16-i^2} \end{aligned}$$

3. Simplify numerator using $i^2 = -1$:

$$\begin{aligned} &= \frac{8-14i-3}{16+1} \\ &= \frac{5-14i}{17} \end{aligned}$$

4. Separate real and imaginary parts:

$$= \frac{5}{17} - \frac{14}{17}i$$

Therefore, $\frac{2-3i}{4+i} = \frac{5}{17} - \frac{14}{17}i$

▲ **Example 7.1.2** Find all complex numbers z that satisfy $|z - 1 - i| = 2$ and $|z + 1 - i| = 2$. Explain the geometric meaning of your answer.

Solution Let's solve this systematically by interpreting the geometric meaning and then finding the algebraic solution.

1. First, interpret each equation geometrically:

$|z - 1 - i| = 2$ describes a circle centered at $1 + i$ with radius 2

$|z + 1 - i| = 2$ describes a circle centered at $-1 + i$ with radius 2

2. Let $z = x + yi$ and expand the first equation:

$$\begin{aligned}|z - 1 - i| &= 2 \\ \sqrt{(x - 1)^2 + (y - 1)^2} &= 2 \\ (x - 1)^2 + (y - 1)^2 &= 4\end{aligned}$$

3. Expand the second equation:

$$\begin{aligned}|z + 1 - i| &= 2 \\ \sqrt{(x + 1)^2 + (y - 1)^2} &= 2 \\ (x + 1)^2 + (y - 1)^2 &= 4\end{aligned}$$

4. Subtract these equations to eliminate the quadratic term in y :

$$\begin{aligned}(x - 1)^2 - (x + 1)^2 &= 0 \\ (x^2 - 2x + 1) - (x^2 + 2x + 1) &= 0 \\ -4x &= 0 \\ x &= 0\end{aligned}$$

5. Substitute $x = 0$ back into either equation:

$$\begin{aligned}(0 - 1)^2 + (y - 1)^2 &= 4 \\ 1 + (y - 1)^2 &= 4 \\ (y - 1)^2 &= 3 \\ y &= 1 \pm \sqrt{3}\end{aligned}$$

6. Therefore, the two solutions are:

$$z = \sqrt{3}i \text{ and } z = (2 - \sqrt{3})i$$

7. Verify these points by checking that each is equidistant from $1 + i$ and $-1 + i$, forming an isosceles triangle with these points as the base vertices.
-

Polar Form and Euler's Formula

While the algebraic form $z = a + bi$ is convenient for addition and subtraction, many properties of complex numbers become clearer in polar form. Given a complex number $z = a + bi$, we can write:

$$z = r(\cos \theta + i \sin \theta)$$

where:

- $r = |z| = \sqrt{a^2 + b^2}$ is the **modulus**
- $\theta = \arg(z) = \arctan\left(\frac{b}{a}\right)$ (with appropriate quadrant adjustments) is the **argument**

The connection between exponentials and trigonometric functions emerges naturally from the Taylor series expansions. Consider the series for e^x :

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots$$

Now substitute $x = i\theta$:

$$\begin{aligned} e^{i\theta} &= 1 + i\theta + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \cdots \\ &= 1 + i\theta - \frac{\theta^2}{2!} - i\frac{\theta^3}{3!} + \frac{\theta^4}{4!} + \cdots \\ &= \left(1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \cdots\right) + i\left(\theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \cdots\right) \\ &= \cos \theta + i \sin \theta \end{aligned}$$

This remarkable derivation leads to **Euler's formula**:

$$e^{i\theta} = \cos \theta + i \sin \theta$$

This formula gives birth to what is often called the most beautiful equation in mathematics. Setting $\theta = \pi$:

$$e^{i\pi} + 1 = 0$$

This single equation unites five fundamental mathematical constants:

- e : The base of natural logarithms
- i : The imaginary unit
- π : The ratio of a circle's circumference to its diameter
- 1 : The multiplicative identity
- 0 : The additive identity

Euler's formula allows us to write any complex number in the elegant form:

$$z = re^{i\theta}$$

This representation transforms complex arithmetic into a more intuitive form:

- Multiplication becomes: $z_1 z_2 = r_1 r_2 e^{i(\theta_1 + \theta_2)}$
- Division becomes: $\frac{z_1}{z_2} = \frac{r_1}{r_2} e^{i(\theta_1 - \theta_2)}$
- Powers become: $(re^{i\theta})^n = r^n e^{in\theta}$

The genius of this representation is that it reveals the geometric nature of complex multiplication: multiplying by $e^{i\theta}$ rotates a complex number by angle θ , while multiplying by r scales its distance from the origin.




Figure 7.4: The polar form $z = re^{i\theta}$ represents a complex number using its distance from the origin (r) and angle from the positive real axis (θ).

▲ **Example 7.1.3** Find all possible values of i^i and express them in the form $a + bi$ where a and b are real numbers.

Solution Let's solve this step by step using the polar form of complex numbers.

1. First, write i in polar form. Since i is located at $(0, 1)$:

$$i = 1 \cdot e^{i\pi/2}$$

2. To compute i^i , we use the complex exponential form:

$$\begin{aligned} i^i &= (e^{i\pi/2})^i \\ &= e^{i^2\pi/2} \\ &= e^{-\pi/2} \end{aligned}$$

3. However, this is only one value. Since $i = e^{i\pi/2+2\pi in}$ for any integer n , we have:

$$\begin{aligned} i^i &= (e^{i\pi/2+2\pi in})^i \\ &= e^{i^2(\pi/2+2\pi n)} \\ &= e^{-\pi/2-2\pi n} \end{aligned}$$

4. The principal value (when $n = 0$) is:

$$\begin{aligned} i^i &= e^{-\pi/2} \\ &\approx 0.2079 \end{aligned}$$

5. Therefore, all values of i^i are real numbers of the form:

$$i^i = e^{-\pi/2-2\pi n}, \quad n \in \mathbb{Z}$$

This surprising result shows that raising an imaginary number to an imaginary power can yield a real number! The result having infinitely many values demonstrates the multi-valued nature of complex exponentiation.

De Moivre's Theorem and Complex Roots

One of the most powerful applications of polar form is **De Moivre's theorem**, which states that for any real number n :

$$(re^{i\theta})^n = r^n e^{in\theta} = r^n (\cos(n\theta) + i \sin(n\theta))$$

This theorem provides a beautiful connection between complex exponentiation and geometric rotation. When $r = 1$, raising $e^{i\theta}$ to the n th power rotates through angle $n\theta$. For example:

- $(e^{i\theta})^2 = e^{2i\theta}$ rotates twice as far
- $(e^{i\theta})^{1/2} = e^{i\theta/2}$ rotates half as far
- $(e^{i\theta})^{-1} = e^{-i\theta}$ rotates in the opposite direction

One of the most fascinating aspects of complex numbers emerges when we study their roots—particularly the **Roots of Unity**. Consider what happens when we try to solve the equation $z^n = 1$. In the real number system, this equation would have at most two solutions (depending on whether n is even or odd). However, in the complex plane, this equation always has exactly n distinct solutions.

To find these solutions, we need to understand why $z^n = 1$ has multiple answers in the first place. When we write a complex number in polar form $z = re^{i\theta}$, we're saying this number

lies r units from the origin at angle θ . If we raise this to the n th power using De Moivre's theorem:

$$z^n = (re^{i\theta})^n = r^n e^{in\theta}$$

For this to equal 1, two things must happen:

- r^n must equal 1, which means $r = 1$ (since r is a positive real number)
- $e^{in\theta}$ must equal 1

Now, Euler's formula tells us that $e^{2\pi i} = 1$, and more generally, $e^{2\pi i k} = 1$ for any integer k . Therefore:

$$e^{in\theta} = 1 \quad \text{if and only if} \quad n\theta = 2\pi k \quad \text{for some integer } k$$

Solving for θ :

$$\theta = \frac{2\pi k}{n}$$

This leads us to the n th roots of unity, given by:

$$\zeta_k = e^{2\pi i k/n}, \quad k = 0, 1, \dots, n-1$$

For example, the fourth roots of unity are:

$$\begin{aligned}\zeta_0 &= 1 = e^{2\pi i \cdot 0/4} = 1 + 0i \\ \zeta_1 &= i = e^{2\pi i \cdot 1/4} = \cos(\pi/2) + i \sin(\pi/2) \\ \zeta_2 &= -1 = e^{2\pi i \cdot 2/4} = \cos(\pi) + i \sin(\pi) \\ \zeta_3 &= -i = e^{2\pi i \cdot 3/4} = \cos(3\pi/2) + i \sin(3\pi/2)\end{aligned}$$




Figure 7.5: The fourth roots of unity form a square in the complex plane, with vertices equally spaced around the unit circle.

More generally, for any complex number $z = re^{i\theta}$, its n th roots are given by:

$$w_k = \sqrt[n]{r} e^{i(\theta + 2\pi k)/n}, \quad k = 0, 1, \dots, n-1$$

This formula has a beautiful geometric interpretation:

- $\sqrt[n]{r}$ determines the distance of all roots from the origin
- The angles $\frac{\theta+2\pi k}{n}$ space the roots evenly around a circle
- The roots form a regular n -gon in the complex plane




Figure 7.6: The sixth roots of unity form a regular hexagon in the complex plane. Here $\omega = e^{2\pi i/6}$ is a primitive sixth root of unity.

The roots of unity play a crucial role in many areas of mathematics:

- They are key to understanding cyclotomic polynomials
- They appear in the Fast Fourier Transform algorithm
- They help solve equations like $x^n = 1$
- They provide symmetries for geometric constructions

A particularly important concept is that of a **primitive root of unity**. A complex number ω is called a primitive n th root of unity if:

- $\omega^n = 1$
- $\omega^k \neq 1$ for all $1 \leq k < n$

For example, if $\omega = e^{2\pi i/6}$, then $\{\omega, \omega^5\}$ are the primitive sixth roots of unity, while $\{1, \omega^2, \omega^3, \omega^4\}$ are not primitive (they are also roots of unity of smaller orders).




Figure 7.7: Powers of a primitive sixth root of unity $\omega = e^{2\pi i/6}$. The arrows show how multiplication by ω cycles through all sixth roots before returning to 1.

De Moivre's theorem and roots of unity demonstrate how deeply intertwined complex numbers are with both algebra and geometry. The regular polygons formed by roots of unity reveal symmetries that would be hidden in purely algebraic manipulations, while the polar form makes clear why these symmetries exist in the first place.

▲ Example 7.1.4 Find all values of $(-8)^{1/3}$ in the complex plane, expressing them both in polar and rectangular form. Which of these is the principal cube root?

Solution Let's solve this step by step using the polar representation of complex numbers.

1. First, write -8 in polar form:

$$-8 = 8e^{i\pi} \quad (\text{since } -8 \text{ lies on the negative real axis})$$

2. To find the cube roots, we use the formula:

$$w_k = \sqrt[3]{r}e^{i(\theta+2\pi k)/3}, \quad k = 0, 1, 2$$

where $r = 8$ and $\theta = \pi$

3. Calculate $\sqrt[3]{r} = 2$ and substitute:

$$\begin{aligned} w_k &= 2e^{i(\pi+2\pi k)/3} \quad k = 0, 1, 2 \\ &= 2e^{i\pi(1+2k)/3} \end{aligned}$$

4. Calculate each root for $k = 0, 1, 2$:

$$w_0 = 2e^{i\pi/3} = 2(\cos(\pi/3) + i\sin(\pi/3)) = 1 + i\sqrt{3}$$

$$w_1 = 2e^{i\pi} = -2$$

$$w_2 = 2e^{5i\pi/3} = 2(\cos(5\pi/3) + i\sin(5\pi/3)) = 1 - i\sqrt{3}$$

5. The principal cube root is the one with argument in $[-\pi/3, \pi/3)$. Here, that's $w_1 = -2$.




Figure 7.8: The three cube roots of -8 form an equilateral triangle when scaled to show their relative positions.

▲ **Example 7.1.5** Find all primitive roots of $z^4 - 1 = 0$, explaining why each root is or is not primitive.

Solution Let's solve this systematically.

1. First, find all fourth roots of unity. Using $\zeta_k = e^{2\pi i k/4}$ for $k = 0, 1, 2, 3$:

$$\zeta_0 = 1 = 1 + 0i$$

$$\zeta_1 = i = 0 + i$$

$$\zeta_2 = -1 = -1 + 0i$$

$$\zeta_3 = -i = 0 - i$$

2. To determine which roots are primitive, we must check the powers of each root:

3. For $\zeta_1 = i$:

$$\begin{aligned} i^1 &= i \\ i^2 &= -1 \\ i^3 &= -i \\ i^4 &= 1 \end{aligned}$$

Since no smaller power equals 1, i is primitive.

4. For $\zeta_2 = -1$:

$$\begin{aligned} (-1)^1 &= -1 \\ (-1)^2 &= 1 \end{aligned}$$

Not primitive as it's a solution to $z^2 = 1$.

5. For $\zeta_3 = -i$:

$$\begin{aligned} (-i)^1 &= -i \\ (-i)^2 &= -1 \\ (-i)^3 &= i \\ (-i)^4 &= 1 \end{aligned}$$

Since no smaller power equals 1, $-i$ is primitive.

6. Therefore, the primitive fourth roots of unity are i and $-i$.




Figure 7.9: The fourth roots of unity. The primitive roots (circled in red) generate all other roots through repeated multiplication.

As shown in the diagram, multiplying by a primitive root (i or $-i$) cycles through all four roots before returning to 1, while multiplying by the non-primitive root -1 only cycles through two values.

Fundamental Theorem of Algebra

Our journey through complex roots and De Moivre's theorem leads us naturally to one of the most profound results in mathematics: the **Fundamental Theorem of Algebra**. This theorem states that every non-constant polynomial with complex coefficients has at least one complex root. While its statement is remarkably simple, its implications are far-reaching.

To appreciate the power of this theorem, let's consider what it tells us. Given any polynomial:

$$p(z) = a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0$$

where $a_n \neq 0$ and the coefficients a_k are complex numbers, there exists at least one complex number c such that $p(c) = 0$.

But the theorem tells us much more. Once we find one root c_1 , we can factor out $(z - c_1)$:

$$p(z) = (z - c_1)(a_n z^{n-1} + b_{n-2} z^{n-2} + \cdots + b_1 z + b_0)$$

The remaining factor is still a polynomial with complex coefficients, so the theorem applies again. This process continues until we have found all n roots (counting multiplicity):

$$p(z) = a_n(z - c_1)(z - c_2) \cdots (z - c_n)$$

This complete factorization has several profound implications:

- Every polynomial of degree n has exactly n complex roots (counting multiplicity)
- There's no need to invent new kinds of numbers to solve polynomial equations
- The complex numbers form an algebraically closed field

The connection to our previous study of roots of unity becomes clear when we consider the equation $z^n - 1 = 0$. The Fundamental Theorem guarantees that this equation has exactly n solutions, and De Moivre's theorem shows us they are:

$$\zeta_k = e^{2\pi i k/n}, \quad k = 0, 1, \dots, n-1$$

More generally, for any complex number w , the equation $z^n = w$ has exactly n solutions:

$$z_k = \sqrt[n]{|w|} e^{i(\arg(w) + 2\pi k)/n}, \quad k = 0, 1, \dots, n-1$$

The completeness of complex numbers stands in stark contrast to the real numbers $\mathbb{R}$. Consider these examples:

- In $\mathbb{R}$: $x^2 + 1 = 0$ has no solutions
- In $\mathbb{C}$: $z^2 + 1 = 0$ has two solutions: i and $-i$
- In $\mathbb{R}$: $x^4 - 2x^2 + 2 = 0$ has no solutions
- In $\mathbb{C}$: $z^4 - 2z^2 + 2 = (z^2 - 1 + i)(z^2 - 1 - i)$ has four solutions

While there are larger number systems that contain $\mathbb{C}$ (such as the quaternions $\mathbb{H}$ or the octonions $\mathbb{O}$), they necessarily sacrifice some of the nice properties of complex arithmetic:

- Quaternions lose commutativity of multiplication ($ij \neq ji$)
- Octonions lose both commutativity and associativity
- Neither system admits an ordering compatible with their operations

This makes $\mathbb{C}$ a sort of “sweet spot” in mathematics—large enough to solve all polynomial equations, yet well-behaved enough to maintain the familiar rules of arithmetic. The Fundamental Theorem of Algebra thus marks complex numbers as both complete and minimal: they are exactly what we need to do algebra, no more and no less.

Lecture 7.2: Complex Functions

Complex functions extend the ideas of real-valued functions into the complex plane, revealing deep connections between algebra and geometry. Unlike real functions, which map points on a line to other points on a line, complex functions transform entire regions of the plane, often producing intricate and elegant geometric structures. These functions play a crucial role in various mathematical fields, offering insights into both theoretical and applied problems.

The Nature of Complex Functions

A complex function f is a rule that assigns to each complex number z in some domain $D \subseteq \mathbb{C}$ another complex number $w = f(z)$. Since both input and output are complex numbers, we can write:

$$f : D \rightarrow \mathbb{C}, \quad z \mapsto w = f(z)$$

Because complex numbers can be written in the form $z = x + yi$, every complex function can be expressed in terms of two real-valued functions $u(x, y)$ and $v(x, y)$:

$$f(z) = f(x + yi) = u(x, y) + iv(x, y)$$

This representation, splitting a complex function into its real and imaginary parts, provides a bridge between complex analysis and multivariable calculus. The functions u and v are called the **component functions** of f .

The behavior of complex functions differs fundamentally from real functions in several ways. When we extend a real function to the complex plane, new phenomena emerge. Consider these key differences:

- A complex function $f(z)$ actually maps points from a plane to another plane, making it inherently two-dimensional in both its domain and range
- The presence of i introduces a rotational aspect to function behavior that has no real-valued analog
- Even simple functions like $f(z) = z^2$ exhibit richer geometric properties than their real counterparts

For example, consider the function $f(z) = z^2$. Writing $z = x + yi$:

$$\begin{aligned} f(z) &= (x + yi)^2 \\ &= x^2 - y^2 + (2xy)i \end{aligned}$$

Here, $u(x, y) = x^2 - y^2$ and $v(x, y) = 2xy$.

The interaction between the real and imaginary parts creates interesting geometric effects. When we square a complex number:

- The function doubles the argument (angle) of the input: $\arg(z^2) = 2\arg(z)$

- The magnitude is squared: $|z^2| = |z|^2$
- Points on the unit circle $|z| = 1$ map to points on the unit circle, but they travel twice as far around it

This geometric behavior contrasts sharply with real functions. When we square a real number, we only see magnitude changes—the geometric aspect of angle doubling is completely hidden in the real case.

Even familiar operations take on new meaning. Consider multiplication by i :

$$\begin{aligned} i(x + yi) &= ix - y \\ &= -y + xi \end{aligned}$$

This simple operation rotates every point in the complex plane by 90 degrees counter-clockwise about the origin. No real-valued function can capture this rotational behavior, highlighting how complex functions incorporate both algebraic and geometric transformations simultaneously.

The real and imaginary component functions u and v are not independent—they are intimately connected through the properties of complex differentiation, as we'll discover when we study the Cauchy-Riemann equations. This interplay between components gives complex functions their distinctive properties and leads to powerful theorems with no real-valued analogues.

Domain and Range Considerations

The domain of a complex function requires careful consideration. While real functions might be restricted by operations like division by zero or taking square roots of negative numbers, complex functions often have broader domains. However, certain functions exhibit multi-valued behavior in the complex plane, necessitating special handling. For instance:

- $\sqrt{z}$ is defined for all complex numbers except $z = 0$, though it requires a branch cut to make it single-valued.
- $\ln(z)$ is defined for all nonzero complex numbers, but since logarithms of complex numbers involve arguments that can wrap around 2π , a branch cut is needed to maintain consistency.
- $\frac{1}{z}$ is defined for all complex numbers except $z = 0$, with no need for a branch cut since it does not exhibit multi-valued behavior.

In cases where a function is naturally multi-valued, a **branch cut** is introduced to restrict its domain to a single-valued region. A branch cut is a deliberate discontinuity in the complex plane that defines a consistent way to evaluate the function. We will discuss this concept in more detail in the following paragraphs.

The concept of **natural domain** extends to complex functions: it's the largest subset of $\mathbb{C}$ where the function can be meaningfully defined. However, we often restrict the domain further to ensure nice properties like continuity or differentiability.

Elementary Complex Functions

The extension of elementary functions to the complex domain reveals surprising depth and beauty. Let's examine each class:

Polynomial and Rational Functions

Complex polynomials greet us as the most approachable extension from real calculus into the vast plane of $\mathbb{C}$, offering a smooth transition from the multivariable mappings we've explored to the intricate world of complex numbers. They take the familiar form:

$$P(z) = a_n z^n + a_{n-1} z^{n-1} + \cdots + a_1 z + a_0,$$

where the coefficients a_k are complex numbers, and $a_n \neq 0$ sets the degree n . These functions are defined everywhere in $\mathbb{C}$, a testament to their well-mannered nature. With no division or roots to restrict them, polynomials operate seamlessly across the plane, free of the gaps or breaks that other functions might encounter. This universality comes from their reliance on addition and multiplication—operations that never falter in the complex realm—making them a steady foundation as we venture deeper.

Their simplicity hides a profound truth: every non-constant polynomial has roots in $\mathbb{C}$, a fact we'll call upon often. For instance, take $P(z) = z^2 - 2z + 2$. Factoring yields:

$$P(z) = (z - 1 - i)(z - 1 + i),$$

with roots at $z = 1 + i$ and $z = 1 - i$, ensuring the degree (2) matches the number of roots when counted appropriately. As z roams far from the origin, the leading term $a_n z^n$ takes charge; for $P(z) = z^2 - 2z + 2$, as $|z| \rightarrow \infty$, $|P(z)| \approx |z|^2 \rightarrow \infty$, stretching magnitudes skyward while angles twist twice around the plane—a geometric flair we'll visualize later.

Rational functions, born from the quotient of polynomials:

$$R(z) = \frac{P(z)}{Q(z)},$$

shift the narrative by introducing points of drama—places where $Q(z) = 0$. These are the **poles**, spots where the function surges to infinity, breaking the calm expanse of polynomials. Consider $R(z) = \frac{z+1}{z-2}$. The pole at $z = 2$ marks where the denominator vanishes, and as z nears 2, say at $z = 2.01$:

$$R(2.01) = \frac{2.01 + 1}{2.01 - 2} = \frac{3.01}{0.01} = 301,$$

a sharp climb that hints at the pole's power. The **order of a pole** at z_0 tells us how fiercely this happens. If $Q(z) = (z - z_0)^m Q'(z)$, where $Q'(z_0) \neq 0$, the pole has order m . A **simple**

pole (order 1) grows linearly in $1/|z - z_0|$, while **multiple poles** (order $m > 1$) amplify this ascent. For $R(z) = \frac{1}{(z-1)^2}$, the pole at $z = 1$ is order 2:

$$R(1.01) = \frac{1}{(0.01)^2} = \frac{1}{0.0001} = 10000,$$

a quadratic leap compared to a simple pole's gentler rise.

This interplay of order shapes the function's behavior near singularities. Picture $R(z) = \frac{z}{(z+i)^3(z-i)}$: it has a triple pole at $z = -i$ and a simple pole at $z = i$. Near $z = -i + 0.01$:

$$R(-i + 0.01) \approx \frac{-i + 0.01}{(0.01)^3(0.01 - i)} \approx \frac{-i}{0.0001 \cdot (-i)} = \frac{1}{0.0001} = 10000,$$

dominated by the triple pole's cubic surge. The following diagram captures this landscape:




Figure 7.10: Poles of a rational function, with simple poles (red dots) at $-1.5 + i$ and $1 - i$, a double pole (blue ring) at $0.5 + 1.5i$, and a triple pole (green rings) at $-2 - 1.5i$.

The diagram paints a vivid scene: simple poles at $-1.5 + i$ and $1 - i$ rise steadily, the double pole at $0.5 + 1.5i$ accelerates quadratically, and the triple pole at $-2 - 1.5i$ soars cubically, each order sculpting the function's climb. Polynomials, in contrast, stretch outward without such interruptions, their roots quietly anchoring the plane while their growth at infinity hints at a rotational twist— $P(z) = z^3$ triples angles as magnitudes cube, a preview of mapping wonders to come.

▲ **Example 7.2.1** Identify the poles and their orders for $R(z) = \frac{z^2+1}{(z-1)^2(z+2)}$, and describe the behavior near $z = 1$.

Solution Set the denominator to zero: $(z - 1)^2(z + 2) = 0$, giving poles at $z = 1$

(order 2) and $z = -2$ (order 1). The numerator $z^2 + 1$ is nonzero at both ($1^2 + 1 = 2$, $(-2)^2 + 1 = 5$), confirming these as poles. Near $z = 1$, let $z = 1 + h$ with small h :

$$R(1+h) = \frac{(1+h)^2 + 1}{h^2(1+h+2)} = \frac{1+2h+h^2+1}{h^2(3+h)} = \frac{2+2h+h^2}{h^2(3+h)}.$$

For $h = 0.01$:

$$R(1.01) = \frac{2 + 0.02 + 0.0001}{0.0001 \cdot 3.01} \approx \frac{2.0201}{0.000301} \approx 6711,$$

a rapid quadratic growth ($|R| \propto 1/h^2$) due to the double pole, far steeper than a simple pole's linear rise.

Polynomials and rational functions thus offer a dual tale: the steady reach of polynomials across $\mathbb{C}$, punctuated by roots and infinite growth, versus the dramatic peaks of rational functions at their poles, each shaping the complex plane in distinct, vivid ways.

Exponential Function

The exponential function, a cornerstone of real calculus, unfurls into a marvel of complexity and elegance when we extend it into the complex plane, revealing a tapestry of growth, rotation, and periodicity that transcends its real origins. In $\mathbb{R}$, e^x is a relentless climber, always positive and surging toward infinity as x grows. In $\mathbb{C}$, it becomes a dancer across the plane, guided by Euler's formula, which breathes life into its complex form:

$$e^z = e^{x+yi} = e^x(\cos y + i \sin y).$$

Here, $z = x + yi$ splits into a real part x that dictates magnitude and an imaginary part y that spins the output around the unit circle, marrying exponential growth with trigonometric oscillation. This definition, elegant in its simplicity, preserves the sacred property of exponents we know so well: $e^{z_1+z_2} = e^{z_1}e^{z_2}$. For $z_1 = 1 + i$ and $z_2 = 2 - i$:

$$e^{(1+i)+(2-i)} = e^3 = e^3(\cos 0 + i \sin 0) = e^3,$$

while separately:

$$e^{1+i}e^{2-i} = e^1(\cos 1 + i \sin 1) \cdot e^2(\cos(-1) + i \sin(-1))$$

$$= e^3(\cos 1 \cos(-1) + i \sin 1 \cos(-1) + i \cos 1 \sin(-1) - \sin 1 \sin(-1)),$$

and since $\cos(-1) = \cos 1$, $\sin(-1) = -\sin 1$, this simplifies to $e^3(\cos^2 1 + \sin^2 1) = e^3$, a harmony unbroken across $\mathbb{C}$.

This extension unveils a function that is **entire**—defined and complex-differentiable for all $z \in \mathbb{C}$ —a trait it shares with polynomials, free of the singularities that haunt rational functions. Compute e^{2+3i} :

$$e^{2+3i} = e^2(\cos 3 + i \sin 3),$$

a point in the plane whose magnitude e^2 reflects $x = 2$, and whose argument 3 radians (about 171.9°) reflects $y = 3$, reachable with any complex input. Unlike its real sibling, which never dips below zero, the complex exponential never rests at zero: $|e^z| = e^x > 0$, and its range is $\mathbb{C} \setminus \{0\}$. Try $e^{0+i\pi} = \cos \pi + i \sin \pi = -1$, a negative real number, impossible in $\mathbb{R}$.

A mesmerizing property emerges in the imaginary direction: periodicity. Since $e^{i2\pi} = \cos 2\pi + i \sin 2\pi = 1$, we find:

$$e^{z+2\pi i} = e^z e^{2\pi i} = e^z \cdot 1 = e^z,$$

a period of $2\pi i$ along the imaginary axis, a cycle absent in real exponentials. This periodicity mirrors the circular motion of $\cos y + i \sin y$, tying the function to the geometry of the unit circle. For $z = x + iy$, as y increases by 2π , the output loops back, tracing a spiral when x varies—a dynamic we'll visualize.

The exponential's mapping properties deepen its allure. Consider $z = x + iy$ as a point in the z -plane. The real part x controls growth ($|e^z| = e^x$), while y dictates rotation ($\arg(e^z) = y$). Horizontal lines $y = c$ map to rays from the origin at angle c , since $e^{x+ic} = e^x e^{ic}$, with magnitude e^x stretching from 0 to ∞ . Vertical lines $x = c$ map to circles $|w| = e^c$, as $e^{c+iy} = e^c(\cos y + i \sin y)$ traces the circle of radius e^c . This transformation from Cartesian strips to polar rays and circles is a hallmark of the complex exponential, connecting to polar coordinates in a way real exponentials cannot.




Figure 7.11: Mapping of the complex exponential $w = e^z$. Horizontal lines in the z -plane ($y = \text{constant}$) map to rays in the w -plane, and vertical lines ($x = \text{constant}$) map to circles of radius e^x .

Its growth rate further distinguishes it. In $\mathbb{R}$, e^x outpaces polynomials; in $\mathbb{C}$, this persists,

but direction matters. Along the real axis ($y = 0$), $|e^x| = e^x \rightarrow \infty$ as $x \rightarrow \infty$. Along the imaginary axis ($x = 0$), $e^{iy} = \cos y + i \sin y$ stays on the unit circle, $|e^{iy}| = 1$. In general directions, $|e^{x+iy}| = e^x$ grows or decays with x , while y spins the phase—a balance of magnitude and motion.

This function's unity with trigonometric functions, via Euler's formula, is profound. Since $e^{iy} = \cos y + i \sin y$ and $e^{-iy} = \cos y - i \sin y$, we derive:

$$\cos y = \frac{e^{iy} + e^{-iy}}{2}, \quad \sin y = \frac{e^{iy} - e^{-iy}}{2i},$$

a bridge we'll cross in the next subsection. For $z = i\pi/4$:

$$e^{i\pi/4} = \cos \frac{\pi}{4} + i \sin \frac{\pi}{4} = \frac{\sqrt{2}}{2} + i \frac{\sqrt{2}}{2},$$

a point at 45° , effortlessly computed.




Figure 7.12: *The complex exponential e^z maps a strip to a spiral, growing with real part x and cycling with imaginary part y , orbiting the unit circle (dashed).*

The exponential's role in visualization is pivotal. Its periodicity and growth make it a lens for periodic phenomena—like waves or rotations—in $\mathbb{C}$, a tool we'll wield in mapping diagrams. For $z = -1 + i2\pi$:

$$e^{-1+i2\pi} = e^{-1}(\cos 2\pi + i \sin 2\pi) = e^{-1} \approx 0.367,$$

a full cycle back to a scaled real value, illustrating its rhythmic nature.

▲ **Example 7.2.2** Find all z such that $e^z = 1 + i$.

Solution Let $e^z = e^{x+iy} = e^x(\cos y + i \sin y) = 1 + i$. Equate magnitudes:

$$|e^z| = e^x = |1 + i| = \sqrt{2}, \quad x = \ln \sqrt{2} = \frac{1}{2} \ln 2.$$

Equate arguments:

$$\tan y = \frac{\sin y}{\cos y} = \frac{1}{1} = 1, \quad y = \frac{\pi}{4} + k\pi,$$

but adjust for periodicity: $e^{x+i(y+2\pi)} = e^{x+iy}$, so principal $y = \pi/4$, with $y = \pi/4 + 2\pi k$.

Thus:

$$z = \frac{1}{2} \ln 2 + i \left(\frac{\pi}{4} + 2\pi k \right), \quad k \in \mathbb{Z}.$$

Trigonometric Functions

In the real world of calculus, sine and cosine weave a rhythmic tale of oscillation, bounded between -1 and 1 , tracing circles and waves with a familiar period of 2π . As we step into the complex plane, these trigonometric functions shed their constraints, unfurling into a broader, more dynamic narrative that ties them intimately to the exponential function we've just explored. Their complex definitions emerge naturally from Euler's formula, revealing a deep kinship:

$$\sin(z) = \frac{e^{iz} - e^{-iz}}{2i}, \quad \cos(z) = \frac{e^{iz} + e^{-iz}}{2}.$$

These expressions, born from the exponential's dance of growth and rotation, extend $\sin x$ and $\cos x$ to all $z = x + yi$ in $\mathbb{C}$, preserving their algebraic essence while unveiling new geometric wonders. Let's test this bridge: for $z = x$ (real),

$$e^{ix} = \cos x + i \sin x, \quad e^{-ix} = \cos x - i \sin x,$$

so:

$$\sin(x) = \frac{(\cos x + i \sin x) - (\cos x - i \sin x)}{2i} = \frac{2i \sin x}{2i} = \sin x,$$

a seamless return to the real case, affirming the extension's harmony.

This connection preserves cherished identities, like $\sin^2(z) + \cos^2(z) = 1$. Compute:

$$\begin{aligned} \sin^2(z) &= \left(\frac{e^{iz} - e^{-iz}}{2i} \right)^2 = \frac{e^{i2z} - 2e^{iz}e^{-iz} + e^{-i2z}}{-4}, \\ \cos^2(z) &= \left(\frac{e^{iz} + e^{-iz}}{2} \right)^2 = \frac{e^{i2z} + 2e^{iz}e^{-iz} + e^{-i2z}}{4}, \end{aligned}$$

and since $e^{iz}e^{-iz} = 1$, sum them:

$$\sin^2(z) + \cos^2(z) = \frac{-e^{i2z} + 2 - e^{-i2z} + e^{i2z} + 2 + e^{-i2z}}{4} = \frac{4}{4} = 1,$$

a truth that holds across $\mathbb{C}$. Yet, the complex plane reshapes their behavior in striking ways. Unlike their real counterparts, $\sin(z)$ and $\cos(z)$ are no longer tethered between -1 and 1 . For $z = i$:

$$\sin(i) = \frac{e^{i \cdot i} - e^{-i \cdot i}}{2i} = \frac{e^{-1} - e}{2i} = i \frac{e - e^{-1}}{2} \approx i \cdot 1.175,$$

a purely imaginary value exceeding 1, reflecting unbounded growth along imaginary paths.

This unboundedness ties to their exponential roots— e^{iz} grows as $\text{Im}(z)$ increases negatively—so $|\sin(z)|$ and $|\cos(z)|$ can soar to infinity. Consider $z = x + iy$:

$$\sin(x + iy) = \frac{e^{i(x+iy)} - e^{-i(x+iy)}}{2i} = \frac{e^{-y+ix} - e^{y-ix}}{2i} = \sin x \cosh y + i \cos x \sinh y,$$

where $\cosh y = (e^y + e^{-y})/2$ and $\sinh y = (e^y - e^{-y})/2$ explode as $|y| \rightarrow \infty$. For $z = \pi/2 + 2i$:

$$\sin\left(\frac{\pi}{2} + 2i\right) = \sin \frac{\pi}{2} \cosh 2 + i \cos \frac{\pi}{2} \sinh 2 = 1 \cdot \frac{e^2 + e^{-2}}{2} + i \cdot 0 \cdot \frac{e^2 - e^{-2}}{2} \approx 3.762,$$

far beyond the real bounds.

Their zeros multiply in $\mathbb{C}$. For $\sin(z) = 0$:

$$\frac{e^{iz} - e^{-iz}}{2i} = 0 \implies e^{iz} = e^{-iz} \implies e^{i2z} = 1 \implies i2z = 2\pi ik,$$

so $z = k\pi$, $k \in \mathbb{Z}$, an infinite parade along the real axis, unlike the finite zeros of polynomials of fixed degree. Similarly, $\cos(z) = 0$ at $z = \pi/2 + k\pi$, echoing this boundless pattern.

Periodicity doubles in complexity. In $\mathbb{R}$, $\sin(x + 2\pi) = \sin x$. In $\mathbb{C}$, this holds, but the imaginary direction adds nuance:

$$\sin(z + 2\pi) = \frac{e^{i(z+2\pi)} - e^{-i(z+2\pi)}}{2i} = \frac{e^{iz}e^{i2\pi} - e^{-iz}e^{-i2\pi}}{2i} = \sin z,$$

since $e^{i2\pi} = 1$. Along the imaginary axis, no new period emerges directly from a simple shift as it does along the real axis, but the exponential terms within $\sin(z)$ weave a rhythmic interplay that unfolds in a striking and dramatic fashion, which we can explore through careful computation. To see this, let's set $z = iy$, where y is a real number, and examine $\sin(z)$ along the imaginary axis:

$$\sin(iy) = \frac{e^{i(iy)} - e^{-i(iy)}}{2i} = \frac{e^{-y} - e^y}{2i} = i \frac{e^y - e^{-y}}{2} = i \sinh y.$$

Here, $\sinh y = (e^y - e^{-y})/2$ —the hyperbolic sine—grows exponentially as $|y|$ increases. For small y , say $y = 0.1$, we find $\sinh 0.1 \approx 0.1002$, so $\sin(i \cdot 0.1) \approx i \cdot 0.1002$, a modest imaginary value. But as y climbs to 2, $\sinh 2 \approx 3.627$, making $\sin(i \cdot 2) \approx i \cdot 3.627$, and by $y = 3$, $\sinh 3 \approx 10.018$, yielding $\sin(i \cdot 3) \approx i \cdot 10.018$ —a value soaring far beyond the real sine's bounds of -1 to 1 . This exponential surge, driven by the e^y term overpowering e^{-y} for large

$|y|$, marks a stark departure from the gentle oscillation we know, revealing an unbounded reach tied to the exponential's influence. Meanwhile, the zeros of $\sin(z)$ march steadily along the real axis at $z = k\pi$, reflecting the 2π period inherited from $e^{i2\pi} = 1$. Together, these behaviors—periodic repetition in one direction and explosive growth in another—craft a vivid contrast, painting $\sin(z)$ as a function of both rhythm and unrestrained expansion across the complex plane.

▲ **Example 7.2.3** Find all z where $\cos(z) = 2$.

Solution Set $\cos(z) = (e^{iz} + e^{-iz})/2 = 2$. Let $w = e^{iz}$:

$$\frac{w + 1/w}{2} = 2 \implies w + \frac{1}{w} = 4 \implies w^2 - 4w + 1 = 0.$$

Solve: $w = 2 \pm \sqrt{3}$. Then $e^{iz} = 2 + \sqrt{3}$ or $2 - \sqrt{3}$, so:

$$iz = \ln(2 \pm \sqrt{3}) + 2\pi ik, \quad z = -i \left(\ln(2 \pm \sqrt{3}) + 2\pi ik \right), \quad k \in \mathbb{Z}.$$

For $k = 0$, $z \approx \pm 1.316i$ (using $\ln(2 + \sqrt{3}) \approx 1.316$), values purely imaginary, far from real possibilities.

These functions weave a tale of boundless reach and infinite repetition, their exponential heritage unlocking a complex rhythm we'll soon contrast with the logarithm's subtleties.

Logarithmic Function

The logarithm, a steadfast companion in real calculus for unwinding exponentials, ventures into the complex plane with a flourish, extending $\ln(x)$ into a domain of depth and subtlety that challenges our single-valued instincts. For $z \neq 0$, we define:

$$\ln(z) = \ln|z| + i \arg(z),$$

where $|z| = \sqrt{x^2 + y^2}$ measures the distance from the origin, and $\arg(z)$ is the angle θ such that $z = |z|e^{i\theta}$. In the real realm, $\ln(x)$ demands $x > 0$, delivering a unique real output with quiet assurance. In $\mathbb{C}$, the condition $z \neq 0$ opens nearly the entire plane—barring the origin, where the logarithm falters—but $\arg(z)$ unveils a profound twist. Since θ is only defined up to multiples of 2π (if $z = e^{i\theta}$, then $z = e^{i(\theta+2\pi k)}$ for any integer k), the complex logarithm emerges as **multi-valued**:

$$\ln(z) = \ln|z| + i(\theta + 2\pi k), \quad k \in \mathbb{Z}.$$

Consider $z = -1 = e^{i\pi}$:

$$\ln(-1) = \ln 1 + i(\pi + 2\pi k) = i\pi(1 + 2k),$$

an infinite cascade of values— $i\pi$, $i3\pi$, $i(-\pi)$, and beyond—whereas real $\ln(-1)$ simply shrugs and steps aside.

This multiplicity is no mere quirk; it mirrors the exponential's periodicity, $e^{z+2\pi i} = e^z$, demanding that the logarithm, as its inverse, embrace every cycle of the plane's spiral. Take $z = 1 + i = \sqrt{2}e^{i\pi/4}$:

$$\ln(1 + i) = \ln \sqrt{2} + i \left(\frac{\pi}{4} + 2\pi k \right),$$

yielding a base value at $k = 0$ of $\frac{1}{2} \ln 2 + i\frac{\pi}{4}$, then stacking others like $\frac{1}{2} \ln 2 + i(2\pi + \frac{\pi}{4})$ or $\frac{1}{2} \ln 2 + i(-2\pi + \frac{\pi}{4})$. Each integer k lifts the imaginary part by $2\pi i$, forming a tower of possibilities that reflects the circular heartbeat of $\mathbb{C}$.

To harness this multiplicity, we often select a **principal branch**, restricting $\arg(z)$ to a fixed range, such as $(-\pi, \pi]$. This **principal logarithm**, denoted $\text{Log}(z)$, becomes:

$$\text{Log}(z) = \ln |z| + i \text{Arg}(z), \quad \text{Arg}(z) \in (-\pi, \pi],$$

slicing a single, consistent layer from the infinite stack. For $z = -2$:

$$\text{Log}(-2) = \ln 2 + i\pi,$$

a clear choice, but approach from just above, at $z = -2 + 0.01i$, and $\text{Arg}(z) \approx 3.14$, or just below, at $-2 - 0.01i$, and $\text{Arg}(z) \approx -3.14$. This jump along the negative real axis hints at a seam—a boundary we'll soon see as a mere edge of a grander structure.

Geometrically, $\ln(z)$ performs a remarkable inversion. Where e^z spins the plane into a polar spiral, mapping $z = x + iy$ to a point with magnitude e^x and angle y , the logarithm unwinds this spiral into a layered expanse. For $z = re^{i\theta}$, we get $w = \ln(z) = \ln r + i(\theta + 2\pi k)$, with real part $u = \ln r$ stretching magnitudes logarithmically and imaginary part $v = \theta + 2\pi k$ stacking angles vertically. Test this with $z = e^{2+i3}$:

$$\ln(e^{2+i3}) = 2 + i3 + 2\pi ik,$$

retrieving the exponent plus periodic shifts, each k placing the result on a different level of an infinite tower. This multi-valued mapping doesn't fit neatly into a flat plane—it cries out for a richer canvas, one that captures its spiraling, layered essence.

To truly grasp this, we turn to the Riemann surface of $\text{Log}(z)$, a three-dimensional marvel that unfurls the logarithm's many values into a continuous, helical structure. Each branch, indexed by k , becomes a sheet, stacked and connected along a seam at the negative real axis, where the argument's 2π jumps weave the surface into a single, flowing form. For $z = -1$, the values $i\pi$, $i3\pi$, $i(-\pi)$ lie on successive sheets, and as z circles the origin, the surface spirals upward or downward, revealing the logarithm's full complexity in a way a flat plot cannot.




Figure 7.13: The Riemann surface of $\text{Log}(z)$, a helical structure where each sheet represents a branch of $\ln(z)$ for $k \in \mathbb{Z}$, connected along the branch cut at the negative real axis. As z circles the origin, the imaginary part ascends or descends through layers, with $\text{Log}(z)$ as the principal sheet ($-\pi < \text{Arg}(z) \leq \pi$).

▲ **Example 7.2.4** Compute all values of $\ln(i)$ and the principal value on the Riemann surface.

Solution Write $i = e^{i\pi/2}$:

$$\ln(i) = \ln 1 + i \left(\frac{\pi}{2} + 2\pi k \right) = i \frac{\pi}{2} (1 + 4k), \quad k \in \mathbb{Z},$$

giving values like $i\pi/2$, $i5\pi/2$, $i(-\pi/2)$, each on a distinct sheet of the Riemann surface. The principal branch ($-\pi < \text{Arg}(z) \leq \pi$) selects:

$$\text{Log}(i) = i \frac{\pi}{2},$$

since $\pi/2$ fits within $(-\pi, \pi]$, anchoring this value to the base sheet.

The logarithm's layered nature, beautifully captured by its Riemann surface, primes us for the next subsection on branch cuts, where we'll slice through this helical expanse to navigate its many dimensions.

Multiple-Valued Functions and Branch Cuts

The complex logarithm's multi-valued nature, stacking outputs like $\ln(-1) = i\pi(1 + 2k)$, hints at a broader phenomenon in $\mathbb{C}$: many functions assign multiple values to a single input. Take the square root: for $z = 4$, $w = 2$ and $w = -2$ both satisfy $w^2 = 4$, doubling the real case. For $z = re^{i\theta}$, $z^{1/2} = \sqrt{r}e^{i\theta/2}$ or $\sqrt{r}e^{i(\theta/2+\pi)}$, a pair tied to $e^{i\pi} = -1$. Similarly, $z^{1/n}$ yields n roots, $r^{1/n}e^{i(\theta+2\pi k)/n}$, $k = 0, 1, \dots, n-1$, cycling around the origin.

To pin these down, we use **branch cuts**—lines where we restrict the function to a single **branch**. For $\ln(z)$, the negative real axis cut gives $\text{Log}(z) = \ln|z| + i\text{Arg}(z)$, $\text{Arg}(z) \in (-\pi, \pi]$. Power functions like $z^{1/3}$ often adopt the same cut, selecting one of three roots. Inverse trigonometric functions, say $\arcsin(z)$, cut from $-\infty$ to -1 and 1 to ∞ , picking a principal path from infinite possibilities. These cuts, arbitrary yet purposeful, echo the Riemann surface's layers, slicing a single sheet from the spiral stack.

▲ **Example 7.2.5** Find the principal value of $(-8)^{1/3}$, cut along the negative real axis, $\text{Arg}(z) \in (-\pi, \pi]$.

Solution Write $-8 = 8e^{i\pi}$: $(-8)^{1/3} = 8^{1/3}e^{i(\pi+2\pi k)/3} = 2e^{i(\pi/3+2\pi k/3)}$. For $k = 0, 1, 2$:

- $k = 0$: $2e^{i\pi/3} = 1 + i\sqrt{3}$ ($\text{Arg} = \pi/3$),
- $k = 1$: $2e^{i\pi} = -2$ ($\text{Arg} = \pi$),
- $k = 2$: $2e^{i5\pi/3} = 1 - i\sqrt{3}$ ($\text{Arg} = -\pi/3$). All fit $(-\pi, \pi]$, but $k = 0$ is typically principal: $1 + i\sqrt{3}$.

Branch cuts thus refine multi-valued functions into single-valued tools, their placement shaping our journey through $\mathbb{C}$'s layered expanse.

Computer-Aided Visualization

As we've journeyed through the complex plane, from polynomials to logarithms, static sketches have hinted at the richness of these functions. Yet, in an era of digital prowess, software unfurls a new dimension of exploration, breathing life into complex mappings with tools that twist, spin, and zoom at our command. These programs don't merely

draw—they invite us to interact with the mathematics, revealing dynamics that paper alone can't capture. A direct plot of a complex function, mapping one two-dimensional plane to another, demands four dimensions—a feat beyond a simple page—but computational tools rise to the challenge with elegant solutions like domain coloring.

Domain coloring paints the complex plane with hues and shades to unveil a function's hidden dance, assigning colors to the range based on its properties. We let **hue** reflect the argument of $f(z)$, the angle it makes in the plane—picture a color wheel where red marks 0, yellow $\pi/2$, green π , cycling through the spectrum over 2π —while **brightness** mirrors the magnitude $|f(z)|$, bright where it soars, dark where it dips. What emerges is a vivid tapestry: **zeros** appear as dark points, colors converging since $|f(z)|$ vanishes; **poles** blaze bright, hues spiraling tightly; **branch cuts** slice as sharp discontinuities—think of $\ln(z)$'s jump from π to $-\pi$ —and **periodicity** unfurls in repeating bands. Consider $f(z) = z^2$: software reveals its zero at the origin as a dark center, with hues doubling around it, a visual echo of $\arg(z^2) = 2 \arg(z)$.




Figure 7.14: Domain coloring of $f(z) = z^2$, where hue encodes the argument of $f(z)$, ranging from $-\pi$ to π to span all directions in the complex plane once, and brightness reflects the magnitude $|f(z)|$ on a logarithmic scale. The doubled cycling of hues around the dark center reveals the angle-doubling property, $\arg(z^2) = 2 \arg(z)$, a hallmark of this quadratic mapping.

```
1 import numpy as np
2 import matplotlib.pyplot as plt
```

```

3 from matplotlib.colors import hsv_to_rgb
4 from matplotlib.cm import ScalarMappable
5
6 # Define the complex function  $f(z) = z^2$ 
7 f = lambda z: z**2
8
9 # Set up the grid in the complex plane
10 x = np.linspace(-2, 2, 400)
11 y = np.linspace(-2, 2, 400)
12 X, Y = np.meshgrid(x, y)
13 Z = X + 1j * Y
14
15 # Compute  $f(z)$ 
16 W = f(Z)
17
18 # Calculate magnitude and argument
19 magnitude = np.abs(W)
20 argument = np.angle(W)
21
22 # Normalize magnitude for brightness (log scale for better contrast)
23 magnitude = np.log1p(magnitude)
24 magnitude = (magnitude - np.min(magnitude)) / (np.max(magnitude) - np.min(magnitude))
25
26 # Create HSV image: hue from argument, full saturation, value from magnitude
27 h = (argument + np.pi) / (2 * np.pi) # Hue: 0 to 1
28 s = np.ones_like(h) # Saturation: full
29 v = magnitude # Value: brightness
30 hsv = np.dstack((h, s, v))
31
32 # Convert HSV to RGB for plotting
33 rgb = hsv_to_rgb(hsv)
34
35 # Generate the plot
36 plt.imshow(rgb, extent=(-2, 2, -2, 2), origin='lower')
37 plt.title(r'Domain Coloring of  $f(z) = z^2$ ')
38 plt.xlabel('Re(z)')
39 plt.ylabel('Im(z)')
40
41 # Add colorbar for argument
42 sm = ScalarMappable(cmap='hsv', norm=plt.Normalize(-np.pi, np.pi))
43 sm.set_array([]) # Required for colorbar to work with ScalarMappable
44 cbar = plt.colorbar(sm, label='Argument of  $f(z)$  (radians)')
45 cbar.set_ticks([-np.pi, -np.pi/2, 0, np.pi/2, np.pi])
46 cbar.set_ticklabels([r'$-\pi$', r'$-\pi/2$', r'$0$', r'$\pi/2$', r'$\pi$'])
47
48 plt.show()

```

Beyond domain coloring, software offers more wonders: **interactive plots** let you click a domain point and see its range image, or drag to trace preimages; **real-time parameter tweaks** shift $f(z) = z^2 + c$ instantly; **animations** layer transformations like e^z atop z^2 ; and **surface plots of $|f(z)|$** offer a topographic view of magnitude. These tools sharpen

our grasp of **geometric effects**—addition nudges, multiplication rotates, as with iz at 90 degrees—while zooming near **singular points** reveals poles flaring or essential singularities writhing. For **multi-valued functions**, software peels apart branches, stacking $\ln(z)$'s layers, and previews **conformal mappings**' angle preservation, as grids bend yet hold shape.

Take $f(z) = e^z$: software maps a grid—horizontal lines to rays, verticals to circles—letting you tilt the view or animate z 's path, the output spiraling in rhythm. This builds on the static depiction in Figure 7.11, amplifying what such figures suggest and opening a window into the function's soul.

Software also lifts us into three dimensions, offering vivid perspectives on complex functions beyond the flat plane. One approach plots the **magnitude** $|f(z)|$ as a surface over the domain, with height revealing growth—zeros dip to the plane, poles spike skyward, and the function's scale unfurls in a topographic sweep. For $f(z) = z^2$, this surface rises parabolically from its zero at the origin, a 3D testament to its quadratic stretch across the plane.




Figure 7.15: Surface plot of $|f(z)| = |z|^2$ over the complex plane, with height showing magnitude rising quadratically from the zero at the origin.

Alternatively, software can surface the **real or imaginary parts**, plotting $u(x, y)$ or $v(x, y)$ to trace their undulations over the complex plane. Take $f(z) = e^z$: its real part $e^x \cos y$ blends exponential growth along the real axis with cosine waves along the imaginary, a 3D ripple that echoes its oscillatory nature. Then there's the **Riemann surface**, stacking multi-valued outputs like $\ln(z)$'s spiral sheets, as seen in Figure 7.13, where branches connect across cuts in a helical dance. These 3D views—crafted by tools like Python or Mathematica—bring the invisible into focus, marrying computation with intuition.




Figure 7.16: Surface plot of the real part of $f(z) = e^z$, where $u(x, y) = e^x \cos y$ blends exponential growth along the real axis with cosine oscillations along the imaginary.

These visualizations, from domain coloring to 3D surfaces, are ingenious compromises. A complex function $f : \mathbb{C} \rightarrow \mathbb{C}$ demands four dimensions—two for the input z , two for the output $f(z)$ —but our tools are bound to three or two. Domain coloring squeezes both magnitude and argument into a 2D image via color, while surface plots of $|f(z)|$, $\text{Re}(f(z))$, or $\text{Im}(f(z))$ reduce the output to one real number, plotted over the complex plane in 3D. Riemann surfaces stack multi-valued outputs, hinting at the function's layers. Each method sacrifices a dimension to make the invisible visible, offering a glimpse into the function's soul at the cost of completeness.

Lecture 7.3: Basic Limits in Complex Analysis

In Chapter 3, Lecture 2, we developed the epsilon–delta definition of limits for functions in $\mathbb{R}^n$. When we move into complex analysis, the formal definition remains identical—using the modulus $|z - z_0|$ as our measure of distance—but the two-dimensional nature of $\mathbb{C}$ and its algebraic structure open up additional techniques for evaluating limits. In this lecture, we expand on the basics of complex limits, highlight methods unique to the complex setting, and explain in detail how to handle expressions involving complex conjugates.

Basic Definition and Similarities

Let $f : D \rightarrow \mathbb{C}$ be a function defined on a domain $D \subseteq \mathbb{C}$, and let z_0 be a limit point of D . We say that

$$\lim_{z \rightarrow z_0} f(z) = L$$

if for every $\epsilon > 0$ there exists a $\delta > 0$ such that for all $z \in D$ with $0 < |z - z_0| < \delta$ we have

$$|f(z) - L| < \epsilon.$$

This is exactly the epsilon–delta definition used in $\mathbb{R}^n$, with the key difference that here distance is measured by the modulus $|z - z_0|$. Because $\mathbb{C}$ is essentially $\mathbb{R}^2$, many aspects of higher-dimensional limits carry over directly. However, the two-dimensional nature of $\mathbb{C}$ introduces an important nuance: there are infinitely many directions from which z can approach z_0 .

In $\mathbb{R}^n$, we typically consider sequences or paths converging to a point and verify that the limit is independent of the chosen path. In $\mathbb{C}$, this requirement is even more critical. For the limit L to exist, the condition $|f(z) - L| < \epsilon$ must hold regardless of the angle or path along which z approaches z_0 . This path-independence guarantees the uniqueness of the limit and is automatically satisfied in one-dimensional real analysis, but must be explicitly checked in the complex setting. Thus, while the definition is formally the same as in $\mathbb{R}^n$, the complex case forces us to account for a richer set of approaches to a limit point.




Figure 7.17: An open disk $D(z_0, \epsilon)$ centered at z_0 with radius ϵ in the complex plane.

New Techniques in Evaluating Complex Limits

Although the basic epsilon–delta definition remains unchanged, several additional techniques are especially useful in complex analysis:

- **Use of Polar Form:** Expressing a complex number z in polar coordinates as

$$z = re^{i\theta},$$

where $r = |z|$ and $\theta = \arg(z)$, can simplify the evaluation of limits. In many cases, limits reduce to a function of r (the distance from the origin) and θ (the angle of approach). This method also helps us check if the limit is path-independent; if the limit depends on θ , it does not exist.

- **Series Expansion:** Similar to real analysis, Taylor series expansions for functions such as $\sin(z)$, $\cos(z)$, or e^z allow us to approximate and evaluate limits. For example, expanding $\sin(z)$ as

$$\sin(z) = z - \frac{z^3}{3!} + \frac{z^5}{5!} - \cdots,$$

simplifies the evaluation of $\frac{\sin(z)}{z}$ as z approaches 0.

- **Handling Complex Conjugates:** Complex conjugates play a special role in $\mathbb{C}$. If $z = re^{i\theta}$, then the conjugate is given by

$$\bar{z} = re^{-i\theta}.$$

This relationship is extremely useful in simplifying expressions where both z and $\bar{z}$ appear. For example, note that $|z|^2 = z\bar{z}$ and that ratios like $\frac{\bar{z}}{z}$ can be expressed in terms of the argument θ , as we will see.

- **Modulus and Argument Techniques:** By decomposing a complex number into its modulus and argument, we can often determine if a limit is well-defined. If the limit expression depends on the angle θ (i.e., the direction of approach), then the limit might not exist. This method is particularly useful when dealing with functions that do not behave uniformly in every direction.

A common scenario in complex limits is the appearance of the complex conjugate. Consider the ratio:

$$\frac{\bar{z}}{z}.$$

Express z in polar form as $z = re^{i\theta}$. Then the conjugate is $\bar{z} = re^{-i\theta}$, and the ratio simplifies to

$$\frac{\bar{z}}{z} = \frac{re^{-i\theta}}{re^{i\theta}} = e^{-2i\theta}.$$

Here, the modulus of the expression is always 1, but its value depends entirely on the angle θ . As $z \rightarrow 0$, although r tends to 0, the value of θ may vary depending on the path taken toward the origin. For example:

- Along the positive real axis ($\theta = 0$), $e^{-2i\theta} = 1$.
- Along the positive imaginary axis ($\theta = \pi/2$), $e^{-2i\theta} = e^{-i\pi} = -1$.

Since the limit depends on the direction of approach, the limit of $\frac{\bar{z}}{z}$ as $z \rightarrow 0$ does not exist.

Properties of Limits in $\mathbb{C}$

The algebraic properties of limits that you are familiar with from real analysis extend naturally to complex functions. Suppose that

$$\lim_{z \rightarrow z_0} f(z) = L \quad \text{and} \quad \lim_{z \rightarrow z_0} g(z) = M.$$

Then the following properties hold:

- $\lim_{z \rightarrow z_0} (f(z) + g(z)) = L + M$.
- $\lim_{z \rightarrow z_0} (f(z) \cdot g(z)) = L \cdot M$.
- For any constant $c \in \mathbb{C}$, $\lim_{z \rightarrow z_0} c f(z) = c L$.
- If $M \neq 0$, $\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = \frac{L}{M}$.

These properties rely on the fact that the modulus $|\cdot|$ in $\mathbb{C}$ obeys the triangle inequality and product rule, much like the Euclidean norm in $\mathbb{R}^2$. The proofs of these properties follow along the same lines as in Chapter 3, Lecture 2, and serve as a useful bridge between real and complex analysis.

▲ **Example 7.3.1** Evaluate the following limits:

1. $\lim_{z \rightarrow 0} z^2$
2. $\lim_{z \rightarrow 0} \frac{\sin(z)}{z}$
3. $\lim_{z \rightarrow 0} \frac{|z|^2}{z}$
4. $\lim_{z \rightarrow 0} \frac{\bar{z}}{z}$

Solution

1. **Limit of z^2 :** Since z^2 is a polynomial, it is continuous everywhere in $\mathbb{C}$. Therefore, by direct substitution:

$$\lim_{z \rightarrow 0} z^2 = 0^2 = 0.$$

2. **Limit of $\frac{\sin(z)}{z}$:** The Taylor series for $\sin(z)$ is:

$$\sin(z) = z - \frac{z^3}{3!} + \frac{z^5}{5!} - \cdots.$$

Dividing by z yields:

$$\frac{\sin(z)}{z} = 1 - \frac{z^2}{3!} + \frac{z^4}{5!} - \cdots.$$

As $z \rightarrow 0$, all terms involving z vanish, hence:

$$\lim_{z \rightarrow 0} \frac{\sin(z)}{z} = 1.$$

3. **Limit of $\frac{|z|^2}{z}$:** Recall that $|z|^2 = z\bar{z}$. Thus, for $z \neq 0$:

$$\frac{|z|^2}{z} = \frac{z\bar{z}}{z} = \bar{z}.$$

As $z \rightarrow 0$, $\bar{z} \rightarrow 0$ (since complex conjugation is continuous), so:

$$\lim_{z \rightarrow 0} \frac{|z|^2}{z} = 0.$$

4. **Limit of $\frac{\bar{z}}{z}$:** Write $z = re^{i\theta}$ so that $\bar{z} = re^{-i\theta}$. Then,

$$\frac{\bar{z}}{z} = e^{-2i\theta}.$$

Since this expression depends on the angle θ and not on the magnitude r , different paths to 0 (with different θ) yield different values. For example:

- If z approaches 0 along the positive real axis ($\theta = 0$), then $\frac{\bar{z}}{z} = 1$.
- If z approaches 0 along the positive imaginary axis ($\theta = \pi/2$), then $\frac{\bar{z}}{z} = e^{-i\pi} = -1$.

Therefore, the limit does not exist.

In summary, while the fundamental definition of limits in $\mathbb{C}$ mirrors that in $\mathbb{R}^n$, the additional structure of complex numbers allows us to employ techniques such as polar representation and series expansions. These methods, along with careful treatment of expressions involving complex conjugates, are essential tools in evaluating complex limits and in detecting path-dependent behavior.

Continuity and Topological Foundations in $\mathbb{C}$

In complex analysis, the notion of continuity is defined in exactly the same way as in real analysis using the epsilon–delta formulation. However, the two-dimensional structure of $\mathbb{C}$ and its rich topology provide additional tools and perspectives that are crucial for a deeper understanding of limits and continuity. In this section, we discuss these foundational ideas and illustrate their usefulness with detailed examples.

Continuity in $\mathbb{C}$

Let $f : D \rightarrow \mathbb{C}$ be a function on a domain $D \subseteq \mathbb{C}$ and let z_0 be a limit point of D . We say that

$$\lim_{z \rightarrow z_0} f(z) = L$$

if for every $\epsilon > 0$ there exists a $\delta > 0$ such that whenever $z \in D$ with $0 < |z - z_0| < \delta$ it follows that

$$|f(z) - L| < \epsilon.$$

This definition uses the modulus $|z - z_0|$ as the distance measure, leveraging the fact that $\mathbb{C}$ is isometric to $\mathbb{R}^2$.

An equivalent formulation employs open sets: f is continuous at z_0 if for every open neighborhood V of $f(z_0)$ there is an open neighborhood U of z_0 such that

$$f(U) \subseteq V.$$

In $\mathbb{C}$, open neighborhoods are typically open disks of the form

$$D(z_0, \epsilon) = \{z \in \mathbb{C} : |z - z_0| < \epsilon\}.$$

▲ Example 7.3.2 Consider the function $f(z) = |z|^2 = z\bar{z}$. Show that f is continuous at any $z_0 \in \mathbb{C}$.

Solution Let $\epsilon > 0$ be given. We wish to find a $\delta > 0$ such that for all z with $|z - z_0| < \delta$,

$$||z|^2 - |z_0|^2| = |z\bar{z} - z_0\bar{z}_0| < \epsilon.$$

Notice that

$$|z\bar{z} - z_0\bar{z}_0| = |z(\bar{z} - \bar{z}_0) + \bar{z}_0(z - z_0)| \leq |z||z - z_0| + |z_0||z - z_0|.$$

Since $|z - z_0| < 1$ implies $|z| \leq |z_0| + 1$, we have

$$|f(z) - f(z_0)| < ((|z_0| + 1) + |z_0|)|z - z_0| = (1 + 2|z_0|)|z - z_0|.$$

Thus, choosing

$$\delta = \min \left\{ 1, \frac{\epsilon}{1 + 2|z_0|} \right\},$$

ensures that $|f(z) - f(z_0)| < \epsilon$ whenever $|z - z_0| < \delta$. Hence, f is continuous at z_0 .

Topological Properties of $\mathbb{C}$ and Their Usefulness

The rich topological structure of $\mathbb{C}$ – inherited from its homeomorphism with $\mathbb{R}^2$ – provides us with powerful tools such as open sets, connectedness, and compactness, which are essential for rigorously understanding limits and continuity. In the following sections, we will develop these technical foundations further to deepen our understanding of continuity in complex analysis.

Open Sets: One of the cornerstones of topology—and by extension, of continuity and limit theory—is the concept of an open set. In Chapter 3, Lecture 2, we covered the basics of open sets in $\mathbb{R}^n$. Since $\mathbb{C}$ is very similar to $\mathbb{R}^2$, these ideas translate directly to the complex plane with an added geometric elegance.

In $\mathbb{C}$, every open set can be expressed as a union of open disks. An open disk centered at z_0 with radius ϵ is defined by

$$D(z_0, \epsilon) = \{z \in \mathbb{C} : |z - z_0| < \epsilon\}.$$

This disk is the basic building block or “neighborhood” of z_0 in the complex plane. It provides a precise way to describe what it means for points to be “close” to z_0 using the modulus as a distance measure.

The usefulness of open disks becomes evident when we prove local properties like continuity. Specifically, to show that a function f is continuous at a point z_0 , one must show that for every open disk $D(f(z_0), \eta)$ around $f(z_0)$ in the codomain, there exists an open disk $D(z_0, \epsilon)$ around z_0 in the domain such that

$$f(D(z_0, \epsilon)) \subseteq D(f(z_0), \eta).$$

This topological formulation is entirely equivalent to the epsilon–delta definition but emphasizes the spatial, geometric intuition behind continuity.

▲ **Example 7.3.3** Consider again $f(z) = z^2$. Demonstrate that for any open disk $D(f(z_0), \eta)$ centered at $f(z_0)$, there exists an open disk $D(z_0, \epsilon)$ such that

$$f(D(z_0, \epsilon)) \subseteq D(f(z_0), \eta).$$

Solution Since $f(z) = z^2$ is continuous, for any $\eta > 0$ there exists $\delta > 0$ so that if $|z - z_0| < \delta$ then

$$|z^2 - z_0^2| = |z - z_0||z + z_0| < \eta.$$

By choosing δ sufficiently small (ensuring, for example, that $|z - z_0| < 1$ so that $|z + z_0|$ is bounded), we guarantee that the entire disk $D(z_0, \delta)$ is mapped inside $D(z_0^2, \eta)$.

Connectedness: The complex plane $\mathbb{C}$ is a prime example of a connected space—it is not only connected but also path-connected. This means that for any two points $z_1, z_2 \in \mathbb{C}$, there exists a continuous path $\gamma : [0, 1] \rightarrow \mathbb{C}$ such that $\gamma(0) = z_1$ and $\gamma(1) = z_2$. In practical terms, this property implies that the complex plane cannot be split into two disjoint nonempty open subsets.

Connectedness has profound implications in analysis. One of the key theorems states that if a function $f : D \rightarrow \mathbb{C}$ is continuous on a connected domain D , then the image $f(D)$ is also connected. This result is critical because it ensures that continuous functions cannot “break apart” the domain into separate pieces. As a consequence, if a continuous function takes two distinct values on a connected set, it must take every value in between (this is an extension of the intermediate value property to connected sets in $\mathbb{C}$).

The equivalence of connectedness and path-connectedness in $\mathbb{C}$ (a feature of all metric spaces like $\mathbb{R}^2$) further simplifies many arguments. When dealing with limits and continuity, knowing that any two points are joined by some continuous path allows us to analyze function behavior along various paths and conclude that the limit (or image) must be the same regardless of the chosen path.

▲ **Example 7.3.4** Let $f(z) = e^z$. Prove that $e^{\mathbb{C}}$ is connected.

Solution The function $f(z) = e^z$ is continuous and maps $\mathbb{C}$ onto $\mathbb{C} \setminus \{0\}$. Since $\mathbb{C}$ is path-connected, any continuous image of $\mathbb{C}$ must be connected. Therefore, $e^{\mathbb{C}} = \mathbb{C} \setminus \{0\}$ is connected.




Figure 7.18: Any two points in $\mathbb{C}$ can be connected by a continuous path.

Compactness: In the setting of $\mathbb{C}$ (which, as we know, is essentially $\mathbb{R}^2$), a set is called compact if it is both closed and bounded. Here, a set is *closed* if it contains all its boundary points, and it is *bounded* if it can be entirely contained within a disk of some finite radius. This result is known as the **Heine–Borel theorem**. Compact sets are very useful in analysis because any continuous function defined on a compact set is guaranteed to reach both a maximum and a minimum value—a fact learned earlier as the Extreme Value Theorem. This property helps ensure that functions behave in a controlled manner, which is critical when dealing with limits.

▲ **Example 7.3.5** Let

$$A = \{z \in \mathbb{C} : 1 \leq |z| \leq 2\}$$

be the closed annulus in $\mathbb{C}$. First, show that A is compact. Then, consider the function

$$f(z) = z^2.$$

Show that f is continuous on A and deduce that $f(A)$ is compact. Finally, determine the range of $|f(z)|$ on A and verify that f attains its maximum and minimum values.

Solution

1. **Prove A is Compact.** The set A is defined by the condition $1 \leq |z| \leq 2$. It is bounded because every $z \in A$ satisfies $|z| \leq 2$, and it is closed because it includes all points on its boundary (those with $|z| = 1$ and $|z| = 2$). By the Heine–Borel theorem, every closed and bounded subset of $\mathbb{R}^2$ is compact. Since $\mathbb{C}$ is homeomorphic to $\mathbb{R}^2$, A is compact.
2. **Show $f(z) = z^2$ is Continuous on A and $f(A)$ is Compact.** Since $f(z) = z^2$ is a polynomial, it is continuous on the entire complex plane, and in particular on the compact set A . The continuous image of a compact set is compact, so $f(A)$ is compact.
3. **Determine the Range of $|f(z)|$ on A .** Notice that for any $z \in \mathbb{C}$,

$$|f(z)| = |z^2| = |z|^2.$$

Since $z \in A$ satisfies $1 \leq |z| \leq 2$, we have:

$$1^2 \leq |z|^2 \leq 2^2 \implies 1 \leq |f(z)| \leq 4.$$

Thus, the minimum modulus of $f(z)$ on A is 1, and the maximum modulus is 4. These extreme values are attained when $|z| = 1$ (yielding $|f(z)| = 1$) and when $|z| = 2$ (yielding $|f(z)| = 4$), respectively.

4. **Conclusion:** We have shown that the closed annulus

$$A = \{z \in \mathbb{C} : 1 \leq |z| \leq 2\}$$

is compact. The function $f(z) = z^2$ is continuous on A , so its image $f(A)$ is also compact. Moreover, since $|f(z)| = |z|^2$, the range of $|f(z)|$ on A is the interval $[1, 4]$, meaning that f attains a minimum modulus of 1 and a maximum modulus of 4 on A .

Extended Complex Plane: To study limits that involve behavior “at infinity,” it is useful to introduce the extended complex plane, denoted

$$\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}.$$

This construction “adds” a single point at infinity to $\mathbb{C}$, thereby compactifying the plane and making it homeomorphic to the Riemann sphere. In other words, by including ∞ , we transform $\mathbb{C}$ from an unbounded space into a compact space, which has many desirable properties in analysis.

One major advantage of working in the extended complex plane is that it allows us to handle limits at infinity in a manner analogous to limits at finite points. Specifically, we define

$$\lim_{z \rightarrow \infty} f(z) = L$$

if, for every $\epsilon > 0$, there exists an $R > 0$ such that whenever $|z| > R$ (that is, z lies outside a disk of radius R), it follows that

$$|f(z) - L| < \epsilon.$$

This definition mirrors the standard epsilon–delta definition of a limit, but with the roles reversed: instead of z approaching a specific finite point, we require the values of $f(z)$ to approach L as z “escapes” to infinity.

The compactification of $\mathbb{C}$ via the addition of ∞ is not merely a technical trick—it has significant consequences. For example, many theorems that hold on compact sets (such as the Extreme Value Theorem) can now be extended to functions defined on $\hat{\mathbb{C}}$. Moreover, considering limits at infinity in this framework unifies our approach to finite and infinite limits.

To visualize the extended complex plane, one often uses the Riemann sphere. Imagine the complex plane as the equatorial plane of a sphere. Each point in $\mathbb{C}$ corresponds to a point on the sphere via stereographic projection, and the “north pole” of the sphere represents the point at infinity. This geometric picture helps us understand how sequences that “escape” to infinity in $\mathbb{C}$ are seen as converging to the north pole on the Riemann sphere.




Figure 7.19: A schematic of the Riemann sphere representing the extended complex plane $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$. The north pole corresponds to the point at infinity, while the equator (solid and dashed) is a cross-sectional view of the complex plane. The points 1, -1 , i , and $-i$ are placed at approximate locations on the equator for illustrative purposes.

▲ **Example 7.3.6** Show that

$$\lim_{z \rightarrow 0} \frac{1}{z} = \infty$$

in the extended complex plane.

Solution For $z \neq 0$,

$$\left| \frac{1}{z} \right| = \frac{1}{|z|}.$$

As $z \rightarrow 0$, $|z|$ becomes arbitrarily small, and consequently, $\frac{1}{|z|}$ becomes arbitrarily large. Formally, for any $M > 0$, if we choose $\delta = \frac{1}{M}$ then whenever $0 < |z| < \delta$, we have

$$\left| \frac{1}{z} \right| > M.$$

Thus, in $\hat{\mathbb{C}}$, we write

$$\lim_{z \rightarrow 0} \frac{1}{z} = \infty.$$

This concept is best visualized via the Riemann sphere, where the point at infinity “completes” the compact space.

In summary, the topological properties of $\mathbb{C}$ —including the structure of open sets, connectedness, and compactness, as well as the notion of the extended complex plane—provide a

rigorous framework for understanding limits. These properties are essential not only in establishing basic limit results but also in preparing us for more advanced topics in the theory of generalized change.

Sequential Continuity in $\mathbb{C}$

In addition to the epsilon–delta definition, continuity in the complex plane can also be characterized via sequences. Specifically, a function $f : D \rightarrow \mathbb{C}$ is **sequentially continuous** at a point z_0 if, for every sequence $\{z_n\}$ in D converging to z_0 , the sequence $\{f(z_n)\}$ converges to $f(z_0)$. This viewpoint can be especially intuitive in $\mathbb{C}$ because there are infinitely many directions from which a point can approach z_0 . By considering *all* sequences that converge to z_0 , we ensure that the limit of $f(z)$ is path-independent. If different sequences converging to z_0 yield different limits for $\{f(z_n)\}$, then f cannot be continuous at z_0 . This sequential perspective thus confirms that continuity demands uniformity of behavior in every direction in the complex plane.

▲ **Example 7.3.7** Consider the function

$$f(z) = \frac{\bar{z}}{z}, \quad z \neq 0.$$

Express z in polar form as $z = re^{i\theta}$, so that $\bar{z} = re^{-i\theta}$. Then,

$$f(z) = \frac{re^{-i\theta}}{re^{i\theta}} = e^{-2i\theta}.$$

Now, observe that if we take different sequences approaching 0 with different fixed angles θ , the value of $e^{-2i\theta}$ will vary. For instance:

- If z approaches 0 along the positive real axis ($\theta = 0$), then $f(z) = e^0 = 1$.
- If z approaches 0 along the positive imaginary axis ($\theta = \pi/2$), then $f(z) = e^{-i\pi} = -1$.

Because the limit depends on the direction of approach, f is not sequentially continuous at $z = 0$.

Solution Since different sequences approaching 0 yield different limit values, by the sequential criterion, $f(z) = \frac{\bar{z}}{z}$ does not have a unique limit as $z \rightarrow 0$. Hence, f is not continuous at 0.

In conclusion, the interplay between the topology of $\mathbb{C}$ and the study of limits is central to complex analysis. By understanding open sets, connectedness, and compactness—and by applying both epsilon–delta and sequential definitions of continuity—we establish a robust

framework for analyzing how functions behave in the complex plane. This foundational work is indispensable not only for basic limit proofs but also for the more advanced topics that follow.

In particular, these topological and continuity concepts pave the way for *complex differentiability* (also known as *holomorphy*), where a function has a complex derivative at every point in a domain. Holomorphic functions, in turn, exhibit remarkable properties such as *analyticity*, meaning they can be locally represented by power series and have infinitely many derivatives. These deeper results in complex analysis—such as the Cauchy integral theorems, the Maximum Modulus Principle, and analytic continuation—all rely heavily on the topological underpinnings we have established here.

Thus, by thoroughly understanding limits and continuity in the context of open sets, connected domains, and compact subsets of $\mathbb{C}$, we equip ourselves with the essential “toolkit” needed to explore the elegant and far-reaching consequences of holomorphic functions in the broader theory of generalized change.

Lecture 7.4: Complex Differentiation

In calculus and real analysis, the derivative is defined as the limit of the difference quotient—a tool that provides the best linear approximation to a function at a given point. But what does it mean to differentiate a function on the complex plane, where every point has infinitely many directions of approach? The idea is similar: we wish for the limit

$$f'(z_0) = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}$$

to exist. However, because the complex plane is two-dimensional, this limit must be the same regardless of the path along which Δz (or equivalently, $z - z_0$) approaches zero. In other words, $f'(z_0)$ must be independent of the direction of approach. This extra requirement is what makes complex differentiability a much stronger condition than in the real setting.

Motivation and Definition

Recall that every complex number z can be expressed as

$$z = x + iy,$$

and accordingly a function f defined on a domain $D \subset \mathbb{C}$ can be written as

$$f(z) = u(x, y) + iv(x, y).$$

We say that f is **complex differentiable** (or **holomorphic**) at a point $z_0 \in D$ if

$$f'(z_0) = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}$$

exists and is independent of the manner in which Δz approaches 0. This condition ensures that, locally, the function can be well-approximated by the linear map

$$f(z) \approx f(z_0) + f'(z_0)(z - z_0),$$

where $f'(z_0)$ acts as both a scaling factor (its modulus $|f'(z_0)|$) and a rotation (its angle $\arg(f'(z_0))$). In the one-dimensional real setting the function has only two possible directions (from the left or the right), but in $\mathbb{C}$ there is a continuum of directions. This stronger requirement forces the real and imaginary parts of f to satisfy interrelated conditions, later formalized as the Cauchy-Riemann equations.

To illustrate the concept, consider the following simplified diagram. The left side shows an ϵ -neighborhood (a circle) in the z -plane centered at z_0 . The requirement is that, no matter which point z is chosen on this circle, the difference quotient

$$\frac{f(z) - f(z_0)}{z - z_0} \quad (\text{or equivalently, } \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z})$$

tends to the same complex number $f'(z_0)$. The right side of the diagram depicts the corresponding linear approximation in the w -plane.




Figure 7.20: Complex differentiation requires the same limit $f'(z_0)$ regardless of the direction of approach. Here, two different approaches, with increments h_1 and h_2 (denoted as Δz), demonstrate that f transforms these vectors via its derivative $f'(z_0)$, preserving both scaling and rotation.

In summary, complex differentiability insists that the difference quotient

$$\frac{f(z) - f(z_0)}{z - z_0} = \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}$$

converges to the same limit $f'(z_0)$ as $\Delta z \rightarrow 0$, regardless of the direction in which Δz approaches 0. This uniformity not only provides a robust linear approximation to f near z_0 but also lays the foundation for many of the elegant and powerful results in complex analysis, such as power series representations and the preservation of angles (conformality). With this understanding, we will next explore how these stringent requirements lead naturally to the Cauchy-Riemann equations and other properties of holomorphic functions.

▲ **Example 7.4.1** Using the limit definition of the derivative, compute the derivatives of:

1. $f(z) = e^z$.
2. $g(z) = \text{Log}(z)$, where

$$\text{Log}(z) = \ln |z| + i \text{Arg}(z) \quad \text{with} \quad -\pi < \text{Arg}(z) \leq \pi,$$

denotes the principal branch of the logarithm.

Solution We will compute the derivatives in two parts.

1. Differentiation of $f(z) = e^z$

(a) By definition, the derivative at z_0 is given by

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{e^z - e^{z_0}}{z - z_0}.$$

Factor the numerator as

$$e^z - e^{z_0} = e^{z_0} (e^{z-z_0} - 1),$$

so that

$$\frac{e^z - e^{z_0}}{z - z_0} = e^{z_0} \cdot \frac{e^{z-z_0} - 1}{z - z_0}.$$

Let $w = z - z_0$; then as $z \rightarrow z_0$ we have

$$e^w = 1 + w + \frac{w^2}{2!} + \frac{w^3}{3!} + \cdots,$$

so that

$$\frac{e^w - 1}{w} = 1 + \frac{w}{2!} + \frac{w^2}{3!} + \cdots \longrightarrow 1 \quad \text{as } w \rightarrow 0.$$

Therefore,

$$f'(z_0) = e^{z_0} \cdot 1 = e^{z_0}.$$

Since z_0 was arbitrary, we conclude that

$$(e^z)' = e^z.$$

2. Differentiation of $g(z) = \text{Log}(z)$ (Principal Branch)

(a) Recall that the principal logarithm is defined by

$$\text{Log}(z) = \ln |z| + i \text{Arg}(z) \quad \text{with } -\pi < \text{Arg}(z) \leq \pi,$$

so that z avoids the negative real axis.

(b) The derivative at z_0 is given by

$$g'(z_0) = \lim_{z \rightarrow z_0} \frac{\text{Log}(z) - \text{Log}(z_0)}{z - z_0}.$$

Using the logarithm identity,

$$\operatorname{Log}(z) - \operatorname{Log}(z_0) = \operatorname{Log}\left(\frac{z}{z_0}\right),$$

we rewrite the difference quotient as

$$\frac{\operatorname{Log}(z) - \operatorname{Log}(z_0)}{z - z_0} = \frac{\operatorname{Log}\left(\frac{z}{z_0}\right)}{z - z_0}.$$

- (c) Express z in terms of z_0 by writing $z = z_0 w$, so that $z - z_0 = z_0(w - 1)$ and $\operatorname{Log}\left(\frac{z}{z_0}\right) = \operatorname{Log}(w)$ with $w \rightarrow 1$ as $z \rightarrow z_0$. Then,

$$\frac{\operatorname{Log}(z) - \operatorname{Log}(z_0)}{z - z_0} = \frac{1}{z_0} \cdot \frac{\operatorname{Log}(w)}{w - 1}.$$

- (d) Write $w = 1 + t$ with $t \rightarrow 0$. Then the Taylor series for the natural logarithm gives

$$\operatorname{Log}(1 + t) = \ln(1 + t) = t - \frac{t^2}{2} + \frac{t^3}{3} - \cdots,$$

so that

$$\frac{\operatorname{Log}(1 + t)}{t} = 1 - \frac{t}{2} + \frac{t^2}{3} - \cdots \longrightarrow 1 \quad \text{as } t \rightarrow 0.$$

- (e) Hence,

$$\lim_{w \rightarrow 1} \frac{\operatorname{Log}(w)}{w - 1} = 1,$$

and it follows that

$$g'(z_0) = \frac{1}{z_0} \cdot 1 = \frac{1}{z_0}.$$

Therefore, for the principal branch,

$$(\operatorname{Log}(z))' = \frac{1}{z}.$$

In conclusion, using the limit definition we have derived:

$$(e^z)' = e^z \quad \text{and} \quad (\operatorname{Log}(z))' = \frac{1}{z}.$$

Cauchy-Riemann Equations

One remarkable consequence of demanding that the difference quotient

$$\frac{f(z) - f(z_0)}{z - z_0}$$

converges to the same limit along every possible direction is that the real and imaginary parts of a holomorphic function cannot vary independently. Writing

$$f(z) = u(x, y) + iv(x, y) \quad \text{for } z = x + iy,$$

the existence of the complex derivative forces the partial derivatives of u and v to satisfy the celebrated **Cauchy-Riemann equations**:

$$u_x(x, y) = v_y(x, y) \quad \text{and} \quad u_y(x, y) = -v_x(x, y).$$

To understand how these equations arise, consider the derivative of f along two natural directions. First, let h be a real increment. Then approaching $z_0 = (x, y)$ along the real axis (i.e. from $z = x + h + iy$) yields

$$\frac{f(x + h, y) - f(x, y)}{h} = u_x(x, y) + iv_x(x, y).$$

Next, consider an imaginary increment ih (i.e. approaching from $z = x + i(y + h)$). In this case, we have

$$\frac{f(x, y + h) - f(x, y)}{ih} = \frac{u(x, y + h) - u(x, y)}{ih} + i \frac{v(x, y + h) - v(x, y)}{ih} = v_y(x, y) - iu_y(x, y).$$

For the limit defining the derivative $f'(z_0)$ to exist and be unique—that is, independent of the direction of approach—these two expressions must be equal. Equating the real parts gives

$$u_x(x, y) = v_y(x, y),$$

and equating the imaginary parts yields

$$v_x(x, y) = -u_y(x, y).$$

Thus, we obtain the Cauchy-Riemann equations:

$$u_x(x, y) = v_y(x, y) \quad \text{and} \quad u_y(x, y) = -v_x(x, y).$$

These equations not only serve as a necessary condition for complex differentiability but, under suitable continuity assumptions on the first partial derivatives, they also imply that f is holomorphic. Moreover, a further consequence is that both u and v are harmonic functions; that is, they satisfy Laplace's equation:

$$u_{xx} + u_{yy} = 0 \quad \text{and} \quad v_{xx} + v_{yy} = 0.$$

This connection to partial differential equations illuminates the deep interplay between analysis and geometry in complex function theory.

▲ **Example 7.4.2** Verify that the following functions satisfy the Cauchy-Riemann equations in their natural coordinate systems:

1. $f(z) = z^2$ (expressed in Cartesian coordinates).
2. $g(z) = z^{3/2}$ (expressed in polar coordinates on its principal branch).

Solution We provide a two-part treatment.

1. Verification for $f(z) = z^2$ in Cartesian Coordinates

- (a) Write $z = x + iy$, so that

$$f(z) = z^2 = (x + iy)^2 = (x^2 - y^2) + i(2xy).$$

Define

$$u(x, y) = x^2 - y^2, \quad v(x, y) = 2xy.$$

- (b) Compute the necessary partial derivatives:

$$u_x = 2x, \quad u_y = -2y, \quad v_x = 2y, \quad v_y = 2x.$$

- (c) Verify the Cauchy-Riemann equations:

$$u_x = 2x = v_y \quad \text{and} \quad u_y = -2y = -v_x.$$

- (d) Since these equations hold for all (x, y) , the function $f(z) = z^2$ is holomorphic.

2. Verification for $g(z) = z^{3/2}$ in Polar Coordinates

- (a) Express z in polar form as

$$z = re^{i\theta}, \quad -\pi < \theta \leq \pi,$$

and define the principal branch by

$$g(z) = z^{3/2} = r^{3/2} e^{\frac{3i\theta}{2}}.$$

Thus, the real and imaginary parts are

$$u(r, \theta) = r^{3/2} \cos\left(\frac{3\theta}{2}\right), \quad v(r, \theta) = r^{3/2} \sin\left(\frac{3\theta}{2}\right).$$

- (b) **Derivation of the polar Cauchy-Riemann equations:** Starting with $z = x + iy$ where $x = r \cos \theta$ and $y = r \sin \theta$, the chain rule gives

$$u_r = u_x \cos \theta + u_y \sin \theta, \quad u_\theta = -r u_x \sin \theta + r u_y \cos \theta,$$

and similarly for v . A brief calculation then leads to the polar Cauchy-Riemann equations:

$$u_r = \frac{1}{r} v_\theta \quad \text{and} \quad v_r = -\frac{1}{r} u_\theta.$$

- (c) Compute the partial derivatives:

- With respect to r :

$$u_r = \frac{3}{2} r^{1/2} \cos \left(\frac{3\theta}{2} \right), \quad v_r = \frac{3}{2} r^{1/2} \sin \left(\frac{3\theta}{2} \right).$$

- With respect to θ :

$$u_\theta = -\frac{3}{2} r^{3/2} \sin \left(\frac{3\theta}{2} \right), \quad v_\theta = \frac{3}{2} r^{3/2} \cos \left(\frac{3\theta}{2} \right).$$

- (d) Verify the polar Cauchy-Riemann equations:

$$u_r = \frac{1}{r} v_\theta \quad \text{and} \quad v_r = -\frac{1}{r} u_\theta.$$

Substituting the derivatives:

$$\frac{1}{r} v_\theta = \frac{1}{r} \cdot \frac{3}{2} r^{3/2} \cos \left(\frac{3\theta}{2} \right) = \frac{3}{2} r^{1/2} \cos \left(\frac{3\theta}{2} \right) = u_r,$$

and

$$-\frac{1}{r} u_\theta = -\frac{1}{r} \left(-\frac{3}{2} r^{3/2} \sin \left(\frac{3\theta}{2} \right) \right) = \frac{3}{2} r^{1/2} \sin \left(\frac{3\theta}{2} \right) = v_r.$$

- (e) Both equations are satisfied, which implies that $g(z) = z^{3/2}$ is holomorphic on its domain (excluding the branch cut).

Rules of Complex Differentiation

The familiar rules of differentiation from real analysis extend to the complex setting. If f and g are holomorphic, then:

- $(f + g)'(z) = f'(z) + g'(z),$

- $(f \cdot g)'(z) = f'(z)g(z) + f(z)g'(z),$
- $\left(\frac{f}{g}\right)'(z) = \frac{f'(z)g(z) - f(z)g'(z)}{[g(z)]^2}$ (provided $g(z) \neq 0$),
- $(f \circ g)'(z) = f'(g(z))g'(z).$

These standard rules serve as the backbone for calculations in complex analysis. In addition, L'Hôpital's rule holds in the complex setting: if

$$\lim_{z \rightarrow z_0} f(z) = 0 \quad \text{and} \quad \lim_{z \rightarrow z_0} g(z) = 0,$$

(or both limits are infinite), and if f and g are holomorphic in a neighborhood of z_0 (except possibly at z_0) with $g'(z) \neq 0$, then

$$\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = \lim_{z \rightarrow z_0} \frac{f'(z)}{g'(z)},$$

provided the limit on the right exists.

▲ **Example 7.4.3** We illustrate these rules with two examples.

1. Application of L'Hôpital's Rule:

Evaluate the limit

$$\lim_{z \rightarrow 0} \frac{e^z - 1}{z}.$$

(a) As $z \rightarrow 0$, the numerator and denominator both tend to 0.

(b) Differentiate the numerator and the denominator with respect to z :

$$\frac{d}{dz}(e^z - 1) = e^z, \quad \frac{d}{dz}z = 1.$$

(c) Thus, by L'Hôpital's rule,

$$\lim_{z \rightarrow 0} \frac{e^z - 1}{z} = \lim_{z \rightarrow 0} \frac{e^z}{1} = e^0 = 1.$$

2. Differentiation of a Composite Function:

Differentiate the function

$$f(z) = \sin(e^z).$$

(a) Recognize that $f(z) = \sin(e^z)$ is a composition of the functions $h(w) = \sin(w)$ and $w(z) = e^z$.

(b) By the chain rule,

$$f'(z) = h'(e^z) \cdot (e^z)'.$$

(c) Since $h'(w) = \cos(w)$ and $(e^z)' = e^z$, we obtain

$$f'(z) = \cos(e^z) \cdot e^z.$$

Analytic Functions and Harmonic Functions

Once a function f is established to be holomorphic—that is, complex differentiable throughout an open domain—it enjoys a suite of properties that extend and deepen the ideas of real analysis. In particular, a holomorphic function is automatically **analytic**. This means that in a neighborhood of any point z_0 in its domain, the function can be expressed as a convergent power series:

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n,$$

so that the behavior of f is completely determined by its derivatives at z_0 . This power series representation not only shows that f is infinitely differentiable, but it also provides practical tools for approximating, differentiating, and integrating f term-by-term within its radius of convergence.

Moreover, if we write

$$f(z) = u(x, y) + iv(x, y) \quad \text{with } z = x + iy,$$

then the holomorphicity of f forces the real function $u(x, y)$ and the imaginary function $v(x, y)$ to satisfy the Cauchy-Riemann equations. A significant consequence of these equations is that both u and v are **harmonic functions**; that is, they satisfy the Laplace equation:

$$u_{xx} + u_{yy} = 0 \quad \text{and} \quad v_{xx} + v_{yy} = 0.$$

Harmonic functions frequently arise in physical contexts—such as in potential theory for gravitational and electrostatic fields, in steady-state heat conduction, and in fluid dynamics—highlighting the practical importance of this concept.

A particularly important property of harmonic functions is the *mean value property*. This property states that the value of a harmonic function at any point equals the average of its values over any circle centered at that point. More precisely, if $u(x, y)$ is harmonic in a domain D and the closed disk

$$\overline{D(z_0, r)} = \{z \in \mathbb{C} : |z - z_0| \leq r\}$$


is contained in D , then

$$u(z_0) = \frac{1}{2\pi} \int_0^{2\pi} u(z_0 + re^{i\theta}) d\theta.$$

This mean value property is a powerful consequence of the Laplace equation and has critical implications for uniqueness and stability in various physical and mathematical problems.

Another striking feature of holomorphic functions is the geometric relationship between the level curves of u and v . In many applications—such as fluid flow or electrostatics—the level curves of u (often interpreted as equipotential lines) and the level curves of v (which can represent streamlines) intersect at right angles. This orthogonality reflects the deep connection between analytic functions and the Laplace equation, and it is a direct consequence of the Cauchy-Riemann equations.

Here the equipotential lines (level curves of $u(x, y) = \ln \sqrt{x^2 + y^2}$) are circles centered at the origin, and the orthogonal trajectories (level curves of $v(x, y) = \arctan(y/x)$) are rays emanating from the origin.



▲ **Example 7.4.4** For the function $f(z) = \text{Log}(z)$ (with the principal branch),

$$\text{Log}(z) = \ln |z| + i \text{Arg}(z) \quad \text{with} \quad -\pi < \text{Arg}(z) \leq \pi,$$

verify that the real part

$$u(x, y) = \ln |z| = \frac{1}{2} \ln(x^2 + y^2)$$

is harmonic.

Solution We proceed as follows:

1. Express the real part of $\text{Log}(z)$ in terms of x and y :

$$u(x, y) = \frac{1}{2} \ln(x^2 + y^2).$$

2. Compute the first partial derivatives:

$$u_x = \frac{1}{2} \cdot \frac{1}{x^2 + y^2} \cdot 2x = \frac{x}{x^2 + y^2}, \quad u_y = \frac{1}{2} \cdot \frac{1}{x^2 + y^2} \cdot 2y = \frac{y}{x^2 + y^2}.$$

3. Compute the second partial derivative with respect to x :

$$u_{xx} = \frac{\partial}{\partial x} \left(\frac{x}{x^2 + y^2} \right) = \frac{(x^2 + y^2) \cdot 1 - x \cdot 2x}{(x^2 + y^2)^2} = \frac{x^2 + y^2 - 2x^2}{(x^2 + y^2)^2} = \frac{y^2 - x^2}{(x^2 + y^2)^2}.$$

4. Compute the second partial derivative with respect to y :

$$u_{yy} = \frac{\partial}{\partial y} \left(\frac{y}{x^2 + y^2} \right) = \frac{(x^2 + y^2) \cdot 1 - y \cdot 2y}{(x^2 + y^2)^2} = \frac{x^2 + y^2 - 2y^2}{(x^2 + y^2)^2} = \frac{x^2 - y^2}{(x^2 + y^2)^2}.$$

5. Add the second derivatives:

$$u_{xx} + u_{yy} = \frac{y^2 - x^2}{(x^2 + y^2)^2} + \frac{x^2 - y^2}{(x^2 + y^2)^2} = 0.$$

6. Since the Laplace equation $u_{xx} + u_{yy} = 0$ holds, $u(x, y) = \ln |z|$ is harmonic.

Power Series and Analyticity

A defining feature of holomorphic functions is their analyticity. That is, if a function f is holomorphic at a point z_0 , then it is equal to its Taylor series in some open disk centered at z_0 . More precisely, there exists a radius $R > 0$ such that for all z with $|z - z_0| < R$,

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^n,$$

where the coefficients are given by the usual Taylor formula:

$$a_n = \frac{f^{(n)}(z_0)}{n!}.$$

This result shows that every holomorphic function is automatically infinitely differentiable, and its entire local behavior is captured by its derivatives at a single point. Within the disk of convergence, this series can be differentiated and integrated term by term, providing a

powerful computational framework.

The analytic structure of holomorphic functions is more than a technical convenience; it reflects the rigid and highly regular nature of complex differentiability. The infinite differentiability is not just formal but constructive—convergence of the Taylor series is guaranteed in a disk where the function is holomorphic.

▲ **Example 7.4.5** Find the power series expansion of the function $f(z) = \frac{1}{z(1-z)}$ centered at $z_0 = 0$, and determine its radius of convergence.

Solution We rewrite $f(z)$ in a form suitable for expansion. Note that

$$f(z) = \frac{1}{z(1-z)} = \frac{1}{z} \cdot \frac{1}{1-z}.$$

1. Recall the geometric series expansion

$$\frac{1}{1-z} = \sum_{n=0}^{\infty} z^n \quad \text{for } |z| < 1.$$

2. Multiply term by term:

$$f(z) = \frac{1}{z} \cdot \sum_{n=0}^{\infty} z^n = \sum_{n=0}^{\infty} z^{n-1}.$$

3. Reindex the series by letting $m = n - 1$, so that $n = m + 1$. Then the sum becomes

$$f(z) = \sum_{m=-1}^{\infty} z^m.$$

4. Thus, the power series expansion of $f(z)$ about $z_0 = 0$ is

$$f(z) = \frac{1}{z} + 1 + z + z^2 + \cdots,$$

valid for $0 < |z| < 1$.

5. The radius of convergence is $R = 1$, but since the term $\frac{1}{z}$ appears, the origin $z = 0$ is a simple pole. So the domain of analyticity is the punctured open disk $0 < |z| < 1$, not the entire open disk.

Liouville's Theorem

Among the most surprising and powerful results in complex analysis (and one that seems counterintuitive when you try to visualize at first) is Liouville's Theorem. This theorem reveals a profound constraint on the behavior of entire functions—those that are holomorphic throughout the entire complex plane. An **entire function** is a function that is holomorphic (complex differentiable) at every point in the complex plane $\mathbb{C}$. Examples include polynomials, the exponential function e^z , and trigonometric functions like $\sin z$ and $\cos z$. Such functions are defined and analytic everywhere, with no singularities at any finite point.

Liouville's Theorem: If f is an entire function and f is bounded (i.e., there exists a constant M such that $|f(z)| \leq M$ for all $z \in \mathbb{C}$), then f must be constant.

This result stands in stark contrast to real analysis, where there are many non-constant functions that are both bounded and differentiable on the entire real line, such as $\sin x$, $\cos x$, $\arctan x$, and e^{-x^2} . In the complex domain, holomorphicity imposes a far stronger constraint—any bounded entire function must reduce to a mere constant.

Liouville's Theorem has profound consequences throughout complex analysis. It serves as the cornerstone for proving the Fundamental Theorem of Algebra, which guarantees that every non-constant polynomial has at least one root. It also reveals the remarkable rigidity of holomorphic functions, showing that they must either grow unboundedly as $|z| \rightarrow \infty$ or remain constant—a stark departure from the more flexible behavior permitted in real analysis.

▲ **Example 7.4.6** Use Liouville's Theorem to prove that there is no entire function $f(z)$ satisfying $|f(z)| = e^{-|z|^2}$ for all $z \in \mathbb{C}$.

Solution We proceed by contradiction. Suppose there exists an entire function $f(z)$ such that $|f(z)| = e^{-|z|^2}$ for all $z \in \mathbb{C}$. First, observe that $e^{-|z|^2}$ is bounded for all $z \in \mathbb{C}$. In fact: $0 < e^{-|z|^2} \leq 1$ for all $z \in \mathbb{C}$ with the maximum value of 1 occurring at $z = 0$. Since $|f(z)| = e^{-|z|^2}$, we know that f is also bounded, with:

$$|f(z)| \leq 1 \quad \text{for all } z \in \mathbb{C}$$

By Liouville's Theorem, any bounded entire function must be constant. Therefore, $f(z) = C$ for some constant C .

But if f is constant, then $|f(z)| = |C|$ for all $z \in \mathbb{C}$. This contradicts our original assumption that $|f(z)| = e^{-|z|^2}$, because $e^{-|z|^2}$ is not constant (it varies with $|z|$). Therefore, no such entire function exists.

Connection to Conformal Mapping

A fundamental consequence of complex differentiability is that, wherever the derivative $f'(z_0) \neq 0$, the function f acts as a local **conformal map**. In other words, not only does f possess a linear approximation

$$f(z) \approx f(z_0) + f'(z_0)(z - z_0),$$

but this approximation has a very special geometric interpretation: the complex number $f'(z_0)$ acts as a scaling and rotation factor. Specifically, the modulus $|f'(z_0)|$ scales infinitesimal distances (i.e., stretches or compresses small figures) and the argument $\arg(f'(z_0))$ rotates them. Consequently, if two curves intersect at z_0 with an angle θ , then their images under f will intersect at $f(z_0)$ with the same angle θ .

The following diagram illustrates this property. In the z -plane (left), two curves intersect at z_0 ; dashed lines represent the tangent directions at z_0 and a small arc between these lines marks the angle θ between them. Under the mapping f , these curves are transformed in the w -plane (right) with the same relative angle preserved.




Figure 7.21: A conformal mapping f (e.g. $f(z) = z + z^2$ near z_0) transforms the original curves in the z -plane (left) into modified curves in the w -plane (right). The red angle arc in the z -plane and the blue angle arc in the w -plane both measure θ , illustrating that f preserves the angle between intersecting curves.

Lecture 7.5: Elementary Complex Integration

Complex integration extends the familiar concept of integration from calculus to the complex plane, allowing us to evaluate integrals along paths rather than just intervals. In Chapter 5, Lecture 2, we studied line integrals in the context of vector fields in $\mathbb{R}^n$; complex integration builds upon these concepts but adds powerful new structures unique to complex analysis. This seemingly straightforward extension leads to surprisingly powerful results that form the cornerstone of complex analysis. In this lecture, we will develop the theory of complex integration, explore its geometric meaning, and establish key properties that will serve as the foundation for the elegant theorems to follow.

Path Integrals in the Complex Plane

In real calculus, we integrate functions over intervals. In complex analysis, we integrate functions along paths or curves in the complex plane, similar to the line integrals we studied earlier, but with the added richness of complex function theory. As we'll see, the requirement of holomorphicity imposes much stronger conditions on complex integrals than we observed with real vector fields, leading to remarkable consequences.

Parametrization of Curves

To precisely define integration along a curve, we first need to formalize the concept of a curve in the complex plane. A **parametrized curve** (or **path**) γ is a continuous function that maps a real interval $[a, b]$ to points in the complex plane:

$$\gamma : [a, b] \rightarrow \mathbb{C}$$

We often express the curve parametrically as $\gamma(t) = x(t) + iy(t)$ for $t \in [a, b]$, where $x(t)$ and $y(t)$ are real-valued continuous functions representing the real and imaginary parts, respectively. This should be familiar from our work with parametrized curves in vector calculus.

A curve γ is said to be **piecewise smooth** if its component functions $x(t)$ and $y(t)$ have continuous derivatives except possibly at finitely many points. A **smooth curve** has continuous derivatives throughout its domain. These conditions ensure that the curve behaves well enough for integration purposes.

The derivative of a curve is denoted by $\gamma'(t) = x'(t) + iy'(t)$, and it represents the velocity vector along the curve at time t . This derivative plays a crucial role in defining the complex line integral, as it provides the direction and rate of traversal along the path.

A curve γ is **simple** if it does not intersect itself, except possibly at its endpoints. When $\gamma(a) = \gamma(b)$, we say the curve is **closed**.

The **length** of a piecewise smooth curve γ over $[a, b]$ is given by:

$$L(\gamma) = \int_a^b |\gamma'(t)| dt = \int_a^b \sqrt{[x'(t)]^2 + [y'(t)]^2} dt$$

Different parametrizations can represent the same geometric curve. For instance, if $\phi : [c, d] \rightarrow [a, b]$ is a strictly increasing, continuously differentiable function with $\phi(c) = a$ and $\phi(d) = b$, then $\gamma \circ \phi$ is a reparametrization of γ . The physical curve is the same, but the "speed" at which we traverse it may differ.

Definition of the Complex Line Integral

Given a complex-valued function $f(z) = u(x, y) + iv(x, y)$ defined on a region containing a piecewise smooth curve $\gamma : [a, b] \rightarrow \mathbb{C}$, the **complex line integral** of f along γ is defined as:

$$\int_{\gamma} f(z) dz = \int_a^b f(\gamma(t)) \gamma'(t) dt$$

This definition requires both f and γ to be well-behaved, with f being continuous along the curve and γ being piecewise smooth. The integral combines the function value at each point with the derivative of the curve, creating a sort of "weighted average" of the function along the path.




Figure 7.22: A geometric representation of a complex line integral. The function $f(z)$ (green vector) is evaluated at each point along the path γ (blue curve) and multiplied by the tangent vector $\gamma'(t)$ (red) before integration.

To understand this definition more clearly, let's express it in terms of real and imaginary parts. Let $f(z) = u(x, y) + iv(x, y)$ and $\gamma(t) = x(t) + iy(t)$. Then:

$$\begin{aligned} \int_{\gamma} f(z) dz &= \int_a^b f(\gamma(t)) \gamma'(t) dt \\ &= \int_a^b [u(x(t), y(t)) + iv(x(t), y(t))] [x'(t) + iy'(t)] dt \\ &= \int_a^b [u \cdot x' - v \cdot y' + i(u \cdot y' + v \cdot x')] dt \end{aligned}$$

This expression highlights that the complex line integral combines the real and imaginary parts of the function in a specific way that accounts for both the function's values and the curve's direction.

For computational purposes, we can write:

$$\begin{aligned} \int_{\gamma} f(z) dz &= \int_a^b [u(x(t), y(t)) \cdot x'(t) - v(x(t), y(t)) \cdot y'(t)] dt \\ &\quad + i \int_a^b [u(x(t), y(t)) \cdot y'(t) + v(x(t), y(t)) \cdot x'(t)] dt \end{aligned}$$

When γ is a segment along the real axis from a to b , parametrized as $\gamma(t) = t$ for $t \in [a, b]$, the complex line integral reduces to the familiar real integral:

$$\int_{\gamma} f(z) dz = \int_a^b f(t) dt$$

which connects complex integration to the real integrals we've studied previously. You can use this list of common parameterizations to aid you. Some of the parameterizations like a circle or arc are a little bit different compared to their real counterpart.

Table 7.1: Common Parameterizations for Complex Line Integrals

Curve Type	Complex Parameterization	Parameter Range	Derivative $\gamma'(t)$
Line Segment	$\gamma(t) = z_1 + t(z_2 - z_1)$	$0 \leq t \leq 1$	$\gamma'(t) = z_2 - z_1$
Circle/Arc	$\gamma(t) = z_0 + re^{it}$	$\alpha \leq t \leq \beta$	$\gamma'(t) = ire^{it}$
Ellipse	$\gamma(t) = z_0 + a \cos t + ib \sin t$	$0 \leq t \leq 2\pi$	$\gamma'(t) = -a \sin t + ib \cos t$
Ray/Radial Line	$\gamma(t) = z_0 + te^{i\theta}$	$0 \leq t \leq R$	$\gamma'(t) = e^{i\theta}$
Rectangular Contour	$\gamma_1 \cup \gamma_2 \cup \gamma_3 \cup \gamma_4$	Piecewise definition	Piecewise constant

▲ **Example 7.5.1** Evaluate the complex line integral $\int_{\gamma} z^2 dz$ where γ is the straight line segment from $z_1 = 1 + i$ to $z_2 = 3 + 4i$.

Solution We proceed step by step to evaluate this integral:

1. First, we parametrize the straight line from z_1 to z_2 . A natural parametrization is:

$$\gamma(t) = (1 - t)(1 + i) + t(3 + 4i) = 1 + i + t(2 + 3i), \quad t \in [0, 1]$$

Simplifying:

$$\gamma(t) = (1 + 2t) + i(1 + 3t), \quad t \in [0, 1]$$

2. The derivative of the path is:

$$\gamma'(t) = 2 + 3i$$

3. We evaluate $f(\gamma(t))$ where $f(z) = z^2$:

$$\begin{aligned} f(\gamma(t)) &= \left[(1 + 2t) + i(1 + 3t) \right]^2 \\ &= (1 + 2t)^2 - (1 + 3t)^2 + 2i(1 + 2t)(1 + 3t) \\ &= \left[(1 + 4t + 4t^2) - (1 + 6t + 9t^2) \right] + 2i(1 + 5t + 6t^2) \\ &= (-2t - 5t^2) + 2i(1 + 5t + 6t^2) \end{aligned}$$

4. Now we compute the integral:

$$\begin{aligned} \int_{\gamma} z^2 dz &= \int_0^1 f(\gamma(t)) \gamma'(t) dt \\ &= \int_0^1 \left[(-2t - 5t^2) + 2i(1 + 5t + 6t^2) \right] (2 + 3i) dt. \end{aligned}$$

Multiplying:

$$\begin{aligned} &\left[(-2t - 5t^2) + 2i(1 + 5t + 6t^2) \right] (2 + 3i) \\ &= (-2t - 5t^2)(2 + 3i) + 2i(1 + 5t + 6t^2)(2 + 3i) \\ &= \left[-4t - 10t^2 - 6it - 15it^2 \right] + \left[-6 - 30t - 36t^2 + 4i + 20it + 24it^2 \right] \\ &= \left[-6 - 34t - 46t^2 \right] + i \left[4 + 14t + 9t^2 \right] \end{aligned}$$

5. Finally, we integrate with respect to t :

$$\int_{\gamma} z^2 dz = \int_0^1 \left[-6 - 34t - 46t^2 + i(4 + 14t + 9t^2) \right] dt$$

$$\begin{aligned}
&= \left[-6t - 17t^2 - \frac{46}{3}t^3 + i(4t + 7t^2 + 3t^3) \right]_0^1 \\
&= -6 - 17 - \frac{46}{3} + i(4 + 7 + 3) \\
&= -\frac{115}{3} + 14i.
\end{aligned}$$

Therefore,

$$\int_{\gamma} z^2 dz = -\frac{115}{3} + 14i.$$

Alternative Forms and Notation

There are several common notations for complex line integrals. If γ is a curve from point z_1 to z_2 , we might write:

$$\int_{\gamma} f(z) dz \quad \text{or} \quad \int_{z_1}^{z_2} f(z) dz \quad \text{or} \quad \int_C f(z) dz$$

When the curve is closed, we sometimes use the notation:

$$\oint_{\gamma} f(z) dz$$

where the circle in the integral sign emphasizes that we're integrating around a closed contour.




Figure 7.23: Various types of paths in the complex plane: a simple curve γ_1 (blue), a closed contour γ_2 (red), and a piecewise curve γ_3 (green). Each path requires appropriate notation and treatment when computing complex line integrals.

For a curve parametrized as $z = z(t)$, we sometimes write:

$$\int_{\gamma} f(z) dz = \int_a^b f(z(t)) z'(t) dt$$

In vector calculus terms, if we denote $\mathbf{r}(t) = (x(t), y(t))$ as the position vector in the complex plane, and $f(z) = P(x, y) + iQ(x, y)$, then:

$$\int_{\gamma} f(z) dz = \int_{\gamma} (P + iQ) d(x + iy) = \int_{\gamma} P dx - Q dy + i \int_{\gamma} Q dx + P dy$$

This relates the complex line integral to the line integrals of vector fields we studied in vector calculus.

▲ **Example 7.5.2** Evaluate the complex line integral $\int_{\gamma} \bar{z} dz$ where γ is the unit circle $|z| = 1$ traversed once counterclockwise.

Solution We need to parametrize the unit circle and apply the definition of the complex line integral.

1. Parametrize the unit circle:

$$\gamma(t) = e^{it}, \quad t \in [0, 2\pi]$$

The derivative is:

$$\gamma'(t) = ie^{it}$$

2. For $z = e^{it}$, the complex conjugate is:

$$\bar{z} = \overline{e^{it}} = e^{-it}$$

3. Now we compute the integrand:

$$f(\gamma(t)) \cdot \gamma'(t) = e^{-it} \cdot ie^{it} = i$$

Notice that this simplifies to a constant.

4. Evaluating the integral:

$$\int_{\gamma} \bar{z} dz = \int_0^{2\pi} e^{-it} \cdot ie^{it} dt$$

$$\begin{aligned}
&= \int_0^{2\pi} i \, dt \\
&= i \cdot 2\pi \\
&= 2\pi i
\end{aligned}$$

Therefore, $\int_{\gamma} \bar{z} \, dz = 2\pi i$ when γ is the unit circle traversed counterclockwise.

This example illustrates an important point: even for a function like $\bar{z}$ that is not holomorphic, the integral around a closed curve need not be zero. As we'll see later in the chapter, this contrasts with holomorphic functions, for which integrals around closed curves in appropriate domains vanish.

Basic Properties of Complex Line Integrals

Complex line integrals inherit many properties from real integrals while also exhibiting some distinctive features due to their path-dependent nature.

Linearity

For any complex-valued functions f and g that are continuous along a piecewise smooth curve γ , and for any complex constants α and β :

$$\int_{\gamma} [\alpha f(z) + \beta g(z)] \, dz = \alpha \int_{\gamma} f(z) \, dz + \beta \int_{\gamma} g(z) \, dz$$

This property follows directly from the linearity of real integrals and allows us to break down complex integrals into simpler components.

Additivity with Respect to Paths

If a curve γ is composed of two curves γ_1 and γ_2 joined end-to-end (i.e., the endpoint of γ_1 is the starting point of γ_2), then:

$$\int_{\gamma} f(z) \, dz = \int_{\gamma_1} f(z) \, dz + \int_{\gamma_2} f(z) \, dz$$

This property allows us to evaluate complex integrals by breaking the path into simpler segments.

Reverse Orientation

If $-\gamma$ denotes the curve γ traversed in the opposite direction, then:

$$\int_{-\gamma} f(z) \, dz = - \int_{\gamma} f(z) \, dz$$

This property reflects the fact that reversing the path direction negates the line integral. If γ is parametrized by $\gamma(t)$ for $t \in [a, b]$, then $-\gamma$ can be parametrized as $(-\gamma)(t) = \gamma(a + b - t)$ for $t \in [a, b]$.

ML Inequality

If f is continuous on a piecewise smooth curve γ , then we have the following estimate:

$$\left| \int_{\gamma} f(z) dz \right| \leq L(\gamma) \cdot \max_{z \in \gamma} |f(z)|$$

where $L(\gamma)$ is the length of the curve. This inequality, often called the ML inequality (where M stands for the maximum value of $|f|$ and L for the path length), provides an upper bound for the magnitude of the complex line integral.

The ML inequality is particularly useful for estimating the size of complex integrals without explicitly evaluating them, which becomes important when studying convergence of series of complex integrals or when establishing bounds for error terms.

To prove this inequality, we use the definition of the complex line integral and basic properties of real integrals:

$$\begin{aligned} \left| \int_{\gamma} f(z) dz \right| &= \left| \int_a^b f(\gamma(t)) \gamma'(t) dt \right| \\ &\leq \int_a^b |f(\gamma(t))| |\gamma'(t)| dt \\ &\leq \max_{t \in [a, b]} |f(\gamma(t))| \cdot \int_a^b |\gamma'(t)| dt \\ &= \max_{z \in \gamma} |f(z)| \cdot L(\gamma) \end{aligned}$$

Integration of Specific Types of Functions

Certain categories of complex functions have special integration properties that are worth highlighting.

Integration of Powers and Polynomials

For simple curves like line segments or circular arcs, we can often compute integrals of powers of z directly. For instance, on a line segment from z_1 to z_2 parametrized as $\gamma(t) = (1-t)z_1 + tz_2$ for $t \in [0, 1]$, we have:

$$\int_{\gamma} z^n dz = \frac{z_2^{n+1} - z_1^{n+1}}{n+1} \quad \text{for } n \neq -1$$

For $n = -1$, the integral of $\frac{1}{z}$ depends on the path and will be discussed in the context of logarithmic functions and multi-valued functions. For polynomials $P(z) = a_0 + a_1z + a_2z^2 +$

$\cdots + a_n z^n$, we can integrate term by term using linearity:

$$\int_{\gamma} P(z) dz = a_0 \int_{\gamma} dz + a_1 \int_{\gamma} z dz + \cdots + a_n \int_{\gamma} z^n dz$$

Integration of Rational Functions

Rational functions, which are quotients of polynomials, are integrated using techniques like partial fraction decomposition, similar to real calculus. The approach breaks down the rational function into simpler terms that can be integrated individually.

For instance, a rational function of the form $\frac{P(z)}{(z-a)^n}$ where $P(z)$ is a polynomial and $(z-a)^n$ is a factor in the denominator, contributes terms involving logarithms (for $n = 1$) or negative powers of $(z-a)$ (for $n > 1$) in the antiderivative.

Integration of Trigonometric and Exponential Functions

The complex exponential function e^z integrates to itself, similar to the real case:

$$\int_{\gamma} e^z dz = e^z \Big|_{\gamma_{\text{start}}}^{\gamma_{\text{end}}} = e^{\gamma_{\text{end}}} - e^{\gamma_{\text{start}}}$$

assuming the integral is taken along a simple curve.

Since the complex trigonometric functions are defined in terms of complex exponentials, their integrals follow familiar patterns from real calculus:

$$\begin{aligned} \int_{\gamma} \sin z dz &= -\cos z \Big|_{\gamma_{\text{start}}}^{\gamma_{\text{end}}} \\ \int_{\gamma} \cos z dz &= \sin z \Big|_{\gamma_{\text{start}}}^{\gamma_{\text{end}}} \end{aligned}$$

▲ Example 7.5.3 Let $k > 0$ be given, and for each $R > 0$ define the closed contour $\Gamma_R = \gamma_1 \cup \gamma_2$, where

1. γ_1 is the line segment along the real axis from $-R$ to R ,
2. γ_2 is the upper semicircular arc from R to $-R$, parameterized by $z(t) = Re^{it}$, $t \in [0, \pi]$.

Show that $\lim_{R \rightarrow \infty} \int_{\gamma_2} e^{ikz} dz = 0$, using the ML inequality and the additivity and linearity of the integral, and explain the role of the γ_1 part.

Solution We first parameterize γ_2 by $z(t) = Re^{it}$, $dz = iRe^{it} dt$, $t \in [0, \pi]$. Then,

$$\int_{\gamma_2} e^{ikz} dz = \int_0^{\pi} e^{ikRe^{it}} iRe^{it} dt.$$

Taking absolute values and applying the ML inequality,

$$\left| \int_{\gamma_2} e^{ikz} dz \right| \leq R \int_0^\pi \left| e^{ikRe^{it}} \right| dt.$$

Since $e^{ikRe^{it}} = e^{ikR(\cos t + i \sin t)} = e^{ikR \cos t} e^{-kR \sin t}$ and hence $\left| e^{ikRe^{it}} \right| = e^{-kR \sin t}$, we obtain

$$\left| \int_{\gamma_2} e^{ikz} dz \right| \leq R \int_0^\pi e^{-kR \sin t} dt.$$

Letting $J(R) = \int_0^\pi e^{-kR \sin t} dt$, we estimate $J(R)$ as follows: For any fixed $\epsilon \in (0, \frac{\pi}{2})$,

- For $t \in [\epsilon, \pi - \epsilon]$, $\sin t \geq \sin \epsilon$ so that

$$\int_\epsilon^{\pi-\epsilon} e^{-kR \sin t} dt \leq (\pi - 2\epsilon) e^{-kR \sin \epsilon}.$$

- For $t \in [0, \epsilon]$ (and similarly near π), using $\sin t \geq t/2$ gives

$$\int_0^\epsilon e^{-kR \sin t} dt \leq \int_0^\epsilon e^{-kRt/2} dt = \frac{2}{kR} (1 - e^{-kR\epsilon/2}).$$

Thus, $J(R)$ decays at least as $O(1/R)$ and exponentially on the middle interval, so that $R \cdot J(R) \rightarrow 0$ as $R \rightarrow \infty$. Hence,

$$\lim_{R \rightarrow \infty} \int_{\gamma_2} e^{ikz} dz = 0.$$

Now, consider γ_1 . Parameterizing by $z = x$ with $x \in [-R, R]$ (so $dz = dx$), we have

$$\int_{\gamma_1} e^{ikz} dz = \int_{-R}^R e^{ikx} dx.$$

Since e^{ikz} is entire, Cauchy's theorem tells us that $\int_{\Gamma_R} e^{ikz} dz = 0$. Thus, by additivity,

$$\int_{-R}^R e^{ikx} dx = - \int_{\gamma_2} e^{ikz} dz.$$

Taking the limit as $R \rightarrow \infty$ and noting that the arc contribution vanishes, we obtain

$$\lim_{R \rightarrow \infty} \int_{-R}^R e^{ikx} dx = 0.$$

This analysis of both γ_2 and γ_1 is a key step in many contour integration techniques used to evaluate real integrals via complex methods.

Antiderivatives and Path Independence

One of the most significant results in complex integration is the fact that under certain conditions, a complex line integral becomes independent of the specific path taken between two points, depending only on the endpoints. This property leads to the concept of antiderivatives in the complex plane.

Antiderivatives in the Complex Plane

A function $F(z)$ is called an **antiderivative** (or **primitive**) of $f(z)$ on a domain D if $F'(z) = f(z)$ for all $z \in D$. When an antiderivative exists on a suitable domain, it allows us to compute complex line integrals using a formula analogous to the Fundamental Theorem of Calculus. Specifically, if $f(z)$ has an antiderivative $F(z)$ on a domain containing a piecewise smooth curve γ from z_1 to z_2 , then:

$$\int_{\gamma} f(z) dz = F(z_2) - F(z_1)$$

This powerful result means that when an antiderivative exists, the complex line integral depends only on the endpoints of the path, not on the specific path taken between them.

Path Independence and Simply Connected Domains

A complex line integral $\int_{\gamma} f(z) dz$ is **path-independent** in a domain D if its value depends only on the endpoints of the path, not on the specific path taken. Path independence is equivalent to the existence of an antiderivative: The integral $\int_{\gamma} f(z) dz$ is path-independent in D if and only if $f(z)$ has an antiderivative in D .

Path independence is closely linked to **simply connected domains** – domains where every simple closed curve can be continuously shrunk to a point while staying within the domain (intuitively, domains without "holes"). For a continuous function $f(z)$ on a simply connected domain D , the following are equivalent:

1. $f(z)$ has an antiderivative in D .
2. $\int_{\gamma} f(z) dz$ is path-independent in D .
3. For every simple closed curve γ in D , $\oint_{\gamma} f(z) dz = 0$.

Multiply Connected Domains and Winding Numbers

A domain that is not simply connected is called **multiply connected**. Such domains contain "holes" that can create path-dependent integrals, even for holomorphic functions.

The **winding number** of a closed curve γ with respect to a point a (not on γ), denoted $n(\gamma, a)$, counts how many times γ encircles the point a in the counterclockwise direction. For a simple closed counterclockwise curve, the winding number is 1 for points inside the curve and 0 for points outside.

For example, in the punctured plane $\mathbb{C} \setminus \{0\}$, the function $f(z) = \frac{1}{z}$ is holomorphic, but the integral $\int_{\gamma} \frac{1}{z} dz$ depends on the winding number of γ around the origin:

$$\oint_{\gamma} \frac{1}{z} dz = 2\pi i \cdot n(\gamma, 0)$$

This formula reveals that the integral equals $2\pi i$ multiplied by the number of counterclockwise revolutions around the origin (negative if clockwise). These quantities, called **periods**, are fundamental in the study of Riemann surfaces and multi-valued functions like the logarithm.

▲ **Example 7.5.4** Evaluate $\oint_{\gamma} \frac{1}{z} dz$ where γ is a closed path that winds twice around the origin counterclockwise.

Solution

1. The winding number of γ around the origin is $n(\gamma, 0) = 2$.




Figure 7.24: A freehand closed contour γ with winding number 2 around the origin. The spiral dial indicator shows how the path winds twice around the origin before closing.

2. Using the winding number formula:

$$\oint_{\gamma} \frac{1}{z} dz = 2\pi i \cdot n(\gamma, 0) = 2\pi i \cdot 2 = 4\pi i$$

3. This confirms that the integral value is directly proportional to the winding number, demonstrating the path dependence in the multiply connected domain $\mathbb{C} \setminus \{0\}$.

Fundamental Theorem for Line Integrals (Revisited)

In vector calculus, we learned that if $\mathbf{F} = \nabla\phi$ is a conservative vector field (i.e., the gradient of a scalar potential ϕ), then the line integral of $\mathbf{F}$ along any curve depends only on the endpoints:

$$\int_{\gamma} \mathbf{F} \cdot d\mathbf{r} = \phi(P_2) - \phi(P_1)$$

where P_1 and P_2 are the endpoints of the curve. The complex analog of this result states that if $f(z)$ has an antiderivative $F(z)$ in a domain D , then for any piecewise smooth curve γ in D with endpoints z_1 and z_2 :

$$\int_{\gamma} f(z) dz = F(z_2) - F(z_1)$$

As in vector calculus, this result is known as the Fundamental Theorem for Line Integrals. It provides a powerful computational tool, allowing us to evaluate complex line integrals without performing the actual integration along the path. The Fundamental Theorem also has a special case for closed curves (i.e., curves where the starting and ending points coincide). If γ is a closed curve in a domain where $f(z)$ has an antiderivative, then:

$$\oint_{\gamma} f(z) dz = 0$$

This vanishing of the integral around closed curves in domains where antiderivatives exist is a precursor to Cauchy's Integral Theorem, which we'll explore in more detail.

▲ **Example 7.5.5** Show that $f(z) = \frac{z}{(z^2+1)^2}$ has an antiderivative in $\mathbb{C} \setminus \{i, -i\}$, and use the Fundamental Theorem for Line Integrals to evaluate $\int_{\gamma} f(z) dz$ along any path from $z_1 = 2$ to $z_2 = 3$ avoiding $\pm i$.

Solution To find an antiderivative for $f(z) = \frac{z}{(z^2+1)^2}$, we make a strategic substitution. Let $u = z^2 + 1$, then $du = 2z dz$, which gives:

$$\begin{aligned} f(z) dz &= \frac{z}{(z^2+1)^2} dz \\ &= \frac{1}{2} \cdot \frac{2z}{(z^2+1)^2} dz \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \cdot \frac{du}{u^2} \\
&= \frac{1}{2} \cdot \left(-\frac{1}{u}\right) + C \\
&= -\frac{1}{2(z^2 + 1)} + C
\end{aligned}$$

We can verify by differentiation:

$$\begin{aligned}
\frac{d}{dz} \left(-\frac{1}{2(z^2 + 1)} \right) &= -\frac{1}{2} \cdot \frac{d}{dz} \left(\frac{1}{z^2 + 1} \right) \\
&= -\frac{1}{2} \cdot \frac{-2z}{(z^2 + 1)^2} \\
&= \frac{z}{(z^2 + 1)^2} = f(z)
\end{aligned}$$

Therefore, $F(z) = -\frac{1}{2(z^2 + 1)}$ is an antiderivative of $f(z)$ in $\mathbb{C} \setminus \{i, -i\}$.
By the Fundamental Theorem for Line Integrals:

$$\begin{aligned}
\int_{\gamma} f(z) dz &= F(z_2) - F(z_1) \\
&= -\frac{1}{2(3^2 + 1)} - \left(-\frac{1}{2(2^2 + 1)} \right) \\
&= -\frac{1}{2(10)} + \frac{1}{2(5)} \\
&= -\frac{1}{20} + \frac{1}{10} \\
&= -\frac{1}{20} + \frac{2}{20} \\
&= \frac{1}{20}
\end{aligned}$$

Thus, $\int_{\gamma} f(z) dz = \frac{1}{20}$ for any path from $z_1 = 2$ to $z_2 = 3$ avoiding $\pm i$.

Lecture 7.6: Cauchy's Integral Theorem and Residues

In our previous lecture, we developed the theory of complex integration along paths in the complex plane. We discovered that the behavior of complex line integrals depends fundamentally on the domain of integration and the analytic properties of the integrand. Now we turn to one of the most profound results in complex analysis: Cauchy's Integral Theorem, which reveals the deep connection between complex differentiability and integration.

This final lecture brings together the key concepts we've developed throughout our exploration of complex analysis and in the study of multivariate calculus. The theorems we'll examine—Cauchy's Integral Theorem, Cauchy's Integral Formula, and the Residue Theorem—provide both theoretical insight and powerful computational tools. These results showcase how the restrictive condition of holomorphicity yields remarkable rewards, allowing us to solve problems that would be difficult or impossible using only real analysis techniques.

Cauchy's Integral Theorem

Cauchy's Integral Theorem represents one of the cornerstones of complex analysis, fundamentally differentiating the behavior of complex functions from their real counterparts. Augustin-Louis Cauchy (1789-1857) first published this result in 1825, though earlier versions appeared in his work from 1814. This theorem establishes a profound connection between the local property of holomorphicity and the global behavior of complex integrals.

Cauchy's Integral Theorem: Let $f(z)$ be holomorphic in a simply connected domain D (sometimes written $f \in \mathcal{O}(D)$). Then for any simple closed curve γ in D :

$$\oint_{\gamma} f(z) dz = 0$$

This remarkable result tells us that when integrating a holomorphic function around a closed loop in a domain where the function has no singularities, the result is always zero. This is true regardless of the specific shape of the loop, as long as it can be continuously deformed within the domain without crossing any points where $f(z)$ fails to be holomorphic.

The power of this theorem becomes evident when contrasted with real analysis. For real-valued functions, no similar general result exists—the integral of a differentiable function around a closed path need not vanish. This difference arises because holomorphicity imposes much stronger conditions than mere differentiability. The Cauchy-Riemann equations couple the behavior of the real and imaginary parts of a function in a way that forces this path-independence property.




Figure 7.25: *Cauchy's Integral Theorem: The integral of a holomorphic function around any closed curve in a simply connected domain equals zero.*

Proof of Cauchy's Integral Theorem

We'll prove the theorem for a special case and then extend it. Let's assume that $f(z)$ is holomorphic in a simply connected domain D and that γ is a simple closed curve forming the boundary of a rectangle in D .

First, we'll show that the integral vanishes for a small rectangle. Let R be a rectangle with vertices at z_0 , $z_0 + h$, $z_0 + h + ik$, and $z_0 + ik$, oriented counterclockwise.




Figure 7.26: *A rectangular contour for proving Cauchy's Integral Theorem.*

Computing the integral along each edge:

- Along γ_1 (from z_0 to $z_0 + h$):

$$\int_{\gamma_1} f(z) dz = \int_0^h f(z_0 + t) dt$$

- Along γ_2 (from $z_0 + h$ to $z_0 + h + ik$):

$$\int_{\gamma_2} f(z) dz = \int_0^k f(z_0 + h + it) i dt$$

- Along γ_3 (from $z_0 + h + ik$ to $z_0 + ik$):

$$\int_{\gamma_3} f(z) dz = \int_h^0 f(z_0 + t + ik) dt = - \int_0^h f(z_0 + t + ik) dt$$

- Along γ_4 (from $z_0 + ik$ to z_0):

$$\int_{\gamma_4} f(z) dz = \int_k^0 f(z_0 + it) i dt = - \int_0^k f(z_0 + it) i dt$$

Adding these four integrals:

$$\begin{aligned} \oint_{\gamma} f(z) dz &= \int_0^h f(z_0 + t) dt + \int_0^k f(z_0 + h + it) i dt \\ &\quad - \int_0^h f(z_0 + t + ik) dt - \int_0^k f(z_0 + it) i dt \\ &= \int_0^h [f(z_0 + t) - f(z_0 + t + ik)] dt \\ &\quad + i \int_0^k [f(z_0 + h + it) - f(z_0 + it)] dt \end{aligned}$$

Now, as h and k become small, we can use the Mean Value Theorem and the Cauchy-Riemann equations:

$$\begin{aligned} f(z_0 + t + ik) - f(z_0 + t) &\approx ik \frac{\partial f}{\partial z}(z_0 + t) = ik f'(z_0 + t) \\ f(z_0 + h + it) - f(z_0 + it) &\approx h \frac{\partial f}{\partial z}(z_0 + it) = h f'(z_0 + it) \end{aligned}$$

Substituting into our integral:

$$\begin{aligned} \oint_{\gamma} f(z) dz &\approx -ik \int_0^h f'(z_0 + t) dt + ih \int_0^k f'(z_0 + it) dt \\ &= -ik[f(z_0 + h) - f(z_0)] + ih[f(z_0 + ik) - f(z_0)] \end{aligned}$$

As $h, k \rightarrow 0$, the error in our approximation approaches zero faster than the area of the rectangle, showing that:

$$\oint_{\gamma} f(z) dz = O(h^2k + hk^2)$$

This means that for sufficiently small rectangles, the integral approaches zero. For a general region, we can subdivide it into a grid of small rectangles. By the additivity property of line integrals, the integrals along internal edges cancel out (each is traversed twice in opposite directions), leaving only the integral around the exterior boundary. This extends the result to any simple closed curve in a simply connected domain.

This proof reveals an important connection: the vanishing of the integral is a direct consequence of the Cauchy-Riemann equations, which define holomorphicity. The theorem thus establishes a profound link between the differential and integral properties of complex functions.

Intuition and Significance

To understand why Cauchy's theorem is so significant, consider the following:

1. In real analysis, the integral $\int_a^b f(x) dx$ can be non-zero even if $f(x)$ is differentiable everywhere. For instance, $\int_0^1 1 dx = 1$.
2. In complex analysis, Cauchy's theorem tells us that the integral of a holomorphic function around any closed loop is always zero, revealing a deep connection between complex differentiability and integration.
3. Holomorphicity is a much stronger condition than real differentiability. This coupling forces the integral around closed curves to vanish.
4. Geometrically, we can think of the complex line integral as measuring the “net circulation” of the function around the curve. Cauchy's theorem tells us that holomorphic functions have zero circulation.

Cauchy's theorem is not merely a computational tool; it reveals a deep structure in the theory of holomorphic functions that has no parallel in real analysis.

▲ **Example 7.6.1** Use Cauchy's Integral Theorem to evaluate

$$\oint_{\gamma} \Gamma(z) dz$$

where γ is a simple closed curve lying entirely in the right half-plane $\operatorname{Re}(z) > 0$.

Solution We begin by recalling the integral definition of the Gamma function for $\operatorname{Re}(z) > 0$:

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt$$

To apply Cauchy's Integral Theorem, we must verify that $\Gamma(z)$ is holomorphic on the domain $\operatorname{Re}(z) > 0$. Let $z = x + iy$, and define

$$\Gamma(z) = u(x, y) + iv(x, y) \quad \text{with} \quad t^{z-1} = t^{x-1} e^{iy \log t}$$

Then the integrand is:

$$t^{z-1} e^{-t} = t^{x-1} e^{-t} [\cos(y \log t) + i \sin(y \log t)]$$

So we have:

$$u(x, y) = \int_0^{\infty} t^{x-1} e^{-t} \cos(y \log t) dt, \quad v(x, y) = \int_0^{\infty} t^{x-1} e^{-t} \sin(y \log t) dt$$

Since the integrand and its partial derivatives with respect to x and y are continuous (for $x > 0$), we may differentiate under the integral sign. Then we compute:

$$\frac{\partial u}{\partial x} = \int_0^\infty t^{x-1} e^{-t} \log t \cdot \cos(y \log t) dt, \quad \frac{\partial v}{\partial y} = \int_0^\infty t^{x-1} e^{-t} \log t \cdot \cos(y \log t) dt$$

$$\frac{\partial u}{\partial y} = - \int_0^\infty t^{x-1} e^{-t} \log t \cdot \sin(y \log t) dt, \quad \frac{\partial v}{\partial x} = \int_0^\infty t^{x-1} e^{-t} \log t \cdot \sin(y \log t) dt$$

So we conclude:

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

Thus, $\Gamma(z)$ satisfies the Cauchy-Riemann equations on the domain $\operatorname{Re}(z) > 0$, and both u and v are continuously differentiable. Therefore, $\Gamma(z)$ is holomorphic on the right half-plane. Since γ lies entirely within this domain and is a simple closed curve, the hypotheses of Cauchy's Integral Theorem are satisfied. Therefore:

$$\oint_{\gamma} \Gamma(z) dz = 0$$

Morera's Theorem

While Cauchy's Integral Theorem tells us that holomorphic functions have vanishing integrals around closed curves, a natural question arises: Is the converse true? That is, if a function's integral vanishes around every closed curve, must the function be holomorphic? Giacinto Morera (1856-1909) answered this question affirmatively in 1886, providing an elegant characterization of holomorphicity through integration rather than differentiation.

Morera's Theorem: Let $f(z)$ be a continuous function in a simply connected domain D . If for every simple closed curve γ in D ,

$$\oint_{\gamma} f(z) dz = 0$$

then $f(z)$ is holomorphic in D .

This remarkable result provides an alternative path to proving holomorphicity. Rather than verifying the Cauchy-Riemann equations—which can be computationally intensive—we can sometimes more easily demonstrate that integrals vanish around closed contours. Morera's Theorem has proven particularly valuable for establishing that limits of holomorphic functions remain holomorphic, and for proving holomorphicity across boundary regions where direct verification via the Cauchy-Riemann equations might be unwieldy.

▲ **Example 7.6.2** Use Morera's Theorem to prove that the function $f(z) = \bar{z}^2$ is not holomorphic in any domain containing the origin.

Solution By Morera's Theorem, if a continuous function f is holomorphic in a domain, then the integral of f over any closed contour in that domain must vanish. To show that $f(z) = \bar{z}^2$ is not holomorphic, it suffices to find a closed contour on which

$$\oint_{\gamma} \bar{z}^2 dz \neq 0.$$

Consider the circle γ centered at $z_0 = 1$ with radius $r = \frac{1}{2}$, parameterized by

$$z(t) = 1 + \frac{1}{2}e^{it}, \quad t \in [0, 2\pi].$$

Then,

$$\overline{z(t)} = 1 + \frac{1}{2}e^{-it} \quad \text{and} \quad dz = \frac{1}{2}i e^{it} dt.$$

Thus,

$$\oint_{\gamma} \bar{z}^2 dz = \int_0^{2\pi} \left(1 + \frac{1}{2}e^{-it}\right)^2 \left(\frac{1}{2}i e^{it}\right) dt.$$

Expanding the square,

$$\left(1 + \frac{1}{2}e^{-it}\right)^2 = 1 + e^{-it} + \frac{1}{4}e^{-2it}.$$

Substitute this into the integral:

$$\oint_{\gamma} \bar{z}^2 dz = \frac{i}{2} \int_0^{2\pi} \left(e^{it} + 1 + \frac{1}{4}e^{-it}\right) dt.$$

Since

$$\int_0^{2\pi} e^{it} dt = 0, \quad \int_0^{2\pi} 1 dt = 2\pi, \quad \int_0^{2\pi} e^{-it} dt = 0,$$

we obtain

$$\oint_{\gamma} \bar{z}^2 dz = \frac{i}{2}(2\pi) = \pi i \neq 0.$$

Therefore, by Morera's Theorem, $f(z) = \bar{z}^2$ is not holomorphic in any domain that contains the origin.

Cauchy's Integral Formula

Cauchy's Integral Theorem leads to one of the most powerful tools in complex analysis: Cauchy's Integral Formula. This formula allows us to express the value of a holomorphic function at any point in terms of its values on a surrounding contour. The formula also reveals a profound connection between holomorphic functions and singularities, particularly poles, which we introduced in Chapter 7, Lecture 2 as points where a function grows unbounded.

Cauchy's Integral Formula: Let $f(z)$ be holomorphic in a simply connected domain D , and let γ be a simple closed positively oriented contour in D . If a is a point inside γ , then:

$$f(a) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{z-a} dz$$

This remarkable formula shows that if we know the values of a holomorphic function on a closed contour, we can determine its value at any point inside that contour. It's worth noting that the integrand $\frac{f(z)}{z-a}$ has a simple pole at $z = a$, but the formula works because a is inside the contour. When functions have poles inside a contour, the corresponding integrals no longer vanish, which gives rise to the theory of residues we'll explore later.

Proof of Cauchy's Integral Formula

Let f be holomorphic on a domain D and let γ be a positively oriented, simple closed contour contained in D , with a a point inside γ . We aim to prove that:

$$f(a) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{z-a} dz$$

Consider a small positively oriented circle γ_{ϵ} of radius $\epsilon > 0$ centered at a , fully contained within γ and D . Define a new function

$$g(z) = \begin{cases} \frac{f(z)-f(a)}{z-a} & z \neq a \\ f'(a) & z = a \end{cases}$$

Since f is holomorphic, so is g . Then,

$$\frac{f(z)}{z-a} = \frac{f(a)}{z-a} + g(z)$$

Now apply Cauchy's Theorem to the annular region between γ and γ_{ϵ} . The integrals over γ and γ_{ϵ} cancel for $g(z)$ since g is holomorphic on this entire region:

$$\begin{aligned} \oint_{\gamma} g(z) dz &= \oint_{\gamma_{\epsilon}} g(z) dz \\ \Rightarrow \oint_{\gamma} \frac{f(z)}{z-a} dz &= f(a) \oint_{\gamma} \frac{1}{z-a} dz + \oint_{\gamma} g(z) dz \end{aligned}$$

By Cauchy's Theorem, $\oint_{\gamma} g(z) dz = \oint_{\gamma_{\epsilon}} g(z) dz$, and since γ_{ϵ} becomes arbitrarily small, this integral vanishes as $\epsilon \rightarrow 0$ due to the continuity of g at a . Thus:

$$\oint_{\gamma} \frac{f(z)}{z-a} dz = f(a) \oint_{\gamma} \frac{1}{z-a} dz$$

But $\oint_{\gamma} \frac{1}{z-a} dz = 2\pi i$ by a standard result of contour integration, so:

$$\oint_{\gamma} \frac{f(z)}{z-a} dz = 2\pi i f(a) \quad \Rightarrow \quad f(a) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{z-a} dz$$

This completes the proof.




Figure 7.27: *Cauchy's Integral Formula: The value of a holomorphic function at a point is determined by its values on surrounding contours.*

This formula has profound implications. It shows that holomorphic functions possess a remarkable "averaging property" where their values at interior points are weighted averages of their boundary values. It also demonstrates that a holomorphic function is completely determined by its values on the boundary of its domain, reflecting the rigidity imposed by the Cauchy-Riemann equations.

Cauchy's Integral Formula has many consequential results that spin off from it. Below are a collection of useful formulas and results that arise in complex analysis:

- **Cauchy's Integral Formula:**

$$f(a) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{z-a} dz$$

where f is holomorphic inside and on γ , and a is inside γ .

• **Cauchy's Formula for Derivatives:**

$$f^{(n)}(a) = \frac{n!}{2\pi i} \oint_{\gamma} \frac{f(z)}{(z-a)^{n+1}} dz$$

for any non-negative integer n . This shows that holomorphic functions are infinitely differentiable.

• **Cauchy's Inequality:**

$$|f^{(n)}(a)| \leq \frac{n! \cdot M}{R^n}$$

where $M = \max_{|z-a|=R} |f(z)|$ on the circle $|z-a|=R$. This provides bounds on the magnitudes of derivatives.

• **Integral Formula for Taylor Coefficients:** If $f(z) = \sum_{n=0}^{\infty} c_n(z-a)^n$, then:

$$c_n = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{(z-a)^{n+1}} dz = \frac{f^{(n)}(a)}{n!}$$

This connects complex integration with power series representations.

▲ **Example 7.6.3** Use Cauchy's Integral Formula to evaluate $\oint_{\gamma} \frac{e^z}{z-2} dz$ where γ is the circle $|z|=3$ oriented counterclockwise.

Solution The function $f(z) = e^z$ is holomorphic everywhere, and the point $a = 2$ lies inside the contour γ . By Cauchy's Integral Formula:

$$f(a) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{z-a} dz$$

Rearranging to solve for the integral:

$$\oint_{\gamma} \frac{f(z)}{z-a} dz = 2\pi i \cdot f(a)$$

In our case:

$$\oint_{|z|=3} \frac{e^z}{z-2} dz = 2\pi i \cdot e^2 = 2\pi i e^2$$

This demonstrates the power of Cauchy's Integral Formula: it allows us to evaluate complex integrals without direct computation, simply by knowing the value of the function at specific points.

▲ **Example 7.6.4** Evaluate the integral

$$\oint_{\gamma} \frac{e^{z^2}}{(z-i)^3} dz$$

where γ is the positively oriented circle $|z| = 2$. What does this tell us about the value of the second derivative of $f(z) = e^{z^2}$ at $z = i$?

Solution We observe that the integrand has a third-order pole at $z = i$, and $f(z) = e^{z^2}$ is entire, so it's holomorphic everywhere. Since i lies inside γ , we apply Cauchy's Integral Formula for derivatives:

$$f^{(n)}(a) = \frac{n!}{2\pi i} \oint_{\gamma} \frac{f(z)}{(z-a)^{n+1}} dz$$

Here $a = i$ and $n = 2$, so the exponent is $n + 1 = 3$. Thus,

$$\oint_{\gamma} \frac{e^{z^2}}{(z-i)^3} dz = 2\pi i \cdot \frac{f^{(2)}(i)}{2!}$$

Now compute $f^{(2)}(z)$ for $f(z) = e^{z^2}$:

$$f'(z) = 2ze^{z^2}, \quad f''(z) = 2e^{z^2} + 4z^2e^{z^2} = (2 + 4z^2)e^{z^2}$$

Then:

$$f^{(2)}(i) = (2 + 4i^2)e^{i^2} = (2 - 4)e^{-1} = -2e^{-1}$$

So:

$$\oint_{\gamma} \frac{e^{z^2}}{(z-i)^3} dz = 2\pi i \cdot \frac{-2e^{-1}}{2} = -2\pi i \cdot \frac{e^{-1}}{1} = -2\pi i e^{-1}$$

Conclusion: The value of this contour integral reveals the second derivative of e^{z^2} at $z = i$ via pure Cauchy theory, demonstrating how the analytic structure of entire functions can be interrogated using higher-order poles.

Laurent Series and Singular Points

While Cauchy's Integral Theorem provides powerful insights about holomorphic functions, in practical applications we frequently encounter functions with points where holomorphicity fails. What happens when we integrate around such points? This question is central to physical applications in fluid dynamics, electromagnetism, and quantum mechanics, where singularities often have physical significance representing sources, vortices, or particles. Cauchy's theorem, by itself, cannot address these cases directly, as it applies only to holomorphic functions integrated along contours in domains where they remain holomorphic.

To bridge this gap, we need to develop tools that characterize the behavior of functions near their singularities and provide a systematic approach to computing integrals around them. This necessity leads us to Laurent series and residue theory, which extend the scope of complex integration beyond purely holomorphic functions. When a function $f(z)$ is not holomorphic at a point z_0 but is holomorphic in some punctured disk $0 < |z - z_0| < R$ around this point, we call z_0 an isolated singularity of f . To understand the behavior of f near such points, we need to generalize power series to include negative powers of $(z - z_0)$.

Laurent Series Expansion

Laurent Series: Let $f(z)$ be holomorphic in an annular region $r < |z - z_0| < R$ for some $0 \leq r < R \leq \infty$. Then $f(z)$ can be represented by a series of the form:

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n = \sum_{n=0}^{\infty} a_n(z - z_0)^n + \sum_{n=1}^{\infty} \frac{a_{-n}}{(z - z_0)^n}$$

which converges absolutely and uniformly on any compact subset of the annulus.

The coefficients a_n in this expansion are given by:

$$a_n = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{(z - z_0)^{n+1}} dz$$

where γ is any positively oriented simple closed contour in the annulus and encircling z_0 once.




Figure 7.28: *Laurent series expansion in an annular region around an isolated singularity. The coefficients of negative powers in the expansion determine the singular behavior of the function at z_0 .*

The Laurent series consists of two parts:

- The **analytic part** (or regular part): $\sum_{n=0}^{\infty} a_n(z - z_0)^n$, which is holomorphic at z_0 .
- The **principal part**: $\sum_{n=1}^{\infty} \frac{a_{-n}}{(z - z_0)^n}$, which captures the singular behavior near z_0 .

The existence and uniqueness of Laurent series can be established using Cauchy's Integral Formula applied to appropriately chosen contours in the annular region. The Laurent expansion provides a complete characterization of the behavior of a function near an isolated singularity.

Classification of Isolated Singularities

Based on the Laurent series expansion, we can classify isolated singularities into three types:

1. **Removable Singularity:** A point z_0 is a removable singularity of $f(z)$ if $a_{-n} = 0$ for all $n \geq 1$ in the Laurent expansion around z_0 . This means the principal part vanishes, and f can be defined (or redefined) at z_0 to be holomorphic there.
2. **Pole of Order m :** A point z_0 is a pole of order m if $a_{-m} \neq 0$ but $a_{-n} = 0$ for all $n > m$. Near such a point, $f(z)$ behaves like $\frac{a_{-m}}{(z - z_0)^m}$ as $z \rightarrow z_0$. We say f has a simple pole at z_0 when $m = 1$.
3. **Essential Singularity:** A point z_0 is an essential singularity if infinitely many coefficients a_{-n} are non-zero, meaning the principal part has infinitely many terms. The behavior of $f(z)$ near an essential singularity can be extremely complicated.

This classification is fundamental in complex analysis and provides a systematic framework for understanding the behavior of meromorphic functions—those that are holomorphic except at isolated poles.




Figure 7.29: Left: A removable singularity where $\lim_{z \rightarrow z_0} f(z)$ exists, as indicated by the dashed circle. Center: A pole where $|f(z)| \rightarrow \infty$, visualized by diverging arrows. Right: An essential singularity; by Picard's theorem, in any neighborhood of z_0 , $f(z)$ assumes every complex value (with at most one exception), illustrated here by overlaid spiral curves...don't stare too long!

▲ **Example 7.6.5** Find the Laurent series expansion of $f(z) = \frac{z^2+1}{z(z-1)(z-2)}$ around $z = 0$ and classify the singularity at this point.

Solution First, we express $f(z)$ using partial fractions:

$$f(z) = \frac{z^2 + 1}{z(z-1)(z-2)} = \frac{A}{z} + \frac{B}{z-1} + \frac{C}{z-2}$$

Multiplying both sides by $z(z-1)(z-2)$:

$$z^2 + 1 = A(z-1)(z-2) + Bz(z-2) + Cz(z-1)$$

- Setting $z = 0$ gives: $1 = A \cdot (-1) \cdot (-2) \Rightarrow A = \frac{1}{2}$
- Setting $z = 1$ gives: $2 = B \cdot 1 \cdot (-1) \Rightarrow B = -2$
- Setting $z = 2$ gives: $5 = C \cdot 2 \cdot 1 \Rightarrow C = \frac{5}{2}$

Therefore:

$$f(z) = \frac{1/2}{z} + \frac{-2}{z-1} + \frac{5/2}{z-2} = \frac{1}{2z} - \frac{2}{z-1} + \frac{5/2}{z-2}$$

To obtain the Laurent series around $z = 0$, we need to expand the terms with $(z-1)$ and $(z-2)$ in the denominator:

$$\frac{1}{z-1} = -\frac{1}{1-z} = -\sum_{n=0}^{\infty} z^n \quad \text{for } |z| < 1$$

$$\frac{1}{z-2} = -\frac{1}{2} \cdot \frac{1}{1-\frac{z}{2}} = -\frac{1}{2} \sum_{n=0}^{\infty} \left(\frac{z}{2}\right)^n = -\frac{1}{2} \sum_{n=0}^{\infty} \frac{z^n}{2^n} \quad \text{for } |z| < 2$$

Thus, for $0 < |z| < 1$:

$$\begin{aligned} f(z) &= \frac{1}{2z} - \frac{2}{z-1} + \frac{5/2}{z-2} \\ &= \frac{1}{2z} + 2 \sum_{n=0}^{\infty} z^n - \frac{5}{4} \sum_{n=0}^{\infty} \frac{z^n}{2^n} \\ &= \frac{1}{2z} + 2 + 2z + 2z^2 + \cdots - \frac{5}{4} - \frac{5}{8}z - \frac{5}{16}z^2 - \cdots \\ &= \frac{1}{2z} + \left(2 - \frac{5}{4}\right) + \left(2 - \frac{5}{8}\right)z + \left(2 - \frac{5}{16}\right)z^2 + \cdots \\ &= \frac{1}{2z} + \frac{3}{4} + \frac{11}{8}z + \frac{27}{16}z^2 + \cdots \end{aligned}$$

Since there is only a $\frac{1}{z}$ term in the principal part (with $a_{-1} = \frac{1}{2}$), and no higher negative powers, the function has a simple pole at $z = 0$.

Residue Theory

When integrating a function with isolated singularities along a closed contour, Cauchy's Integral Theorem no longer applies directly. However, a remarkably elegant theory—residue theory—allows us to evaluate such integrals by examining the behavior of the function near each enclosed singularity.

Definition and Basic Properties of Residues

Residue: Given a function $f(z)$ with an isolated singularity at z_0 and Laurent expansion $f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n$ valid in some punctured disk around z_0 , the residue of f at z_0 is the coefficient a_{-1} and is denoted:

$$\text{Res}(f, z_0) = a_{-1}$$

The fundamental importance of residues is revealed by the following theorem:

Residue Theorem: Let $f(z)$ be analytic in a simply connected domain D except for isolated singularities at $z_1, z_2, \dots, z_n$. If γ is a simple closed positively oriented contour in D that doesn't pass through any of the singularities but encloses them, then:

$$\oint_{\gamma} f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f, z_k)$$

This theorem reduces the evaluation of complex contour integrals to the calculation of residues at the enclosed singularities, providing a powerful computational tool.

$$f(z) \notin \mathcal{O}(\Omega), \Omega \subset D$$




Figure 7.30: *The Residue Theorem: The integral equals $2\pi i$ times the sum of residues at enclosed singularities.*

Computing Residues

Several techniques exist for calculating residues, particularly for poles:

1. **From Laurent Series:** If we can find the Laurent expansion of $f(z)$ around z_0 , then $\text{Res}(f, z_0) = a_{-1}$.
2. **For Simple Poles:** If $f(z) = \frac{g(z)}{h(z)}$ has a simple pole at z_0 (meaning $h(z_0) = 0$ but $h'(z_0) \neq 0$ and $g(z_0) \neq 0$), then:

$$\text{Res}(f, z_0) = \frac{g(z_0)}{h'(z_0)}$$

3. **For Poles of Order m :** If $f(z)$ has a pole of order m at z_0 , then:

$$\text{Res}(f, z_0) = \frac{1}{(m-1)!} \lim_{z \rightarrow z_0} \frac{d^{m-1}}{dz^{m-1}} [(z - z_0)^m f(z)]$$

4. **For Simple Poles with Special Form:** If $f(z) = \frac{p(z)}{q(z)}$ where p and q are holomorphic at z_0 , $q(z_0) = 0$, $q'(z_0) \neq 0$, and $p(z_0) \neq 0$, then:

$$\text{Res}(f, z_0) = \frac{p(z_0)}{q'(z_0)}$$

▲ **Example 7.6.6** Evaluate $\oint_{\gamma} \frac{e^z}{z^2(z-1)^3} dz$ where γ is the circle $|z| = 2$ oriented counterclockwise.

Solution The function $f(z) = \frac{e^z}{z^2(z-1)^3}$ has singularities at $z = 0$ (a pole of order 2) and $z = 1$ (a pole of order 3). Both are inside the contour $\gamma : |z| = 2$. By the Residue Theorem:

$$\oint_{\gamma} f(z) dz = 2\pi i [\text{Res}(f, 0) + \text{Res}(f, 1)]$$

Let's compute each residue:

For $z = 0$ (pole of order 2):

$$\begin{aligned} \text{Res}(f, 0) &= \lim_{z \rightarrow 0} \frac{d}{dz} [z^2 f(z)] \\ &= \lim_{z \rightarrow 0} \frac{d}{dz} \left[\frac{e^z}{(z-1)^3} \right] \\ &= \lim_{z \rightarrow 0} \left[\frac{e^z}{(z-1)^3} - \frac{3e^z}{(z-1)^4} \right] \\ &= \frac{e^0}{(-1)^3} - \frac{3e^0}{(-1)^4} = -1 - 3 = -4 \end{aligned}$$

For $z = 1$ (pole of order 3):

$$\begin{aligned}\operatorname{Res}(f, 1) &= \frac{1}{2!} \lim_{z \rightarrow 1} \frac{d^2}{dz^2} [(z-1)^3 f(z)] \\ &= \frac{1}{2} \lim_{z \rightarrow 1} \frac{d^2}{dz^2} \left[\frac{e^z}{z^2} \right]\end{aligned}$$

Computing the derivatives:

$$\begin{aligned}\frac{d}{dz} [e^z z^{-2}] &= e^z z^{-2} - 2e^z z^{-3} \\ \frac{d^2}{dz^2} [e^z z^{-2}] &= e^z z^{-2} - 2e^z z^{-3} - 2e^z z^{-3} + 6e^z z^{-4} \\ &= e^z z^{-2} - 4e^z z^{-3} + 6e^z z^{-4}\end{aligned}$$

Evaluating at $z = 1$:

$$\begin{aligned}\operatorname{Res}(f, 1) &= \frac{1}{2} [e^1 \cdot 1^{-2} - 4e^1 \cdot 1^{-3} + 6e^1 \cdot 1^{-4}] \\ &= \frac{1}{2} [e - 4e + 6e] \\ &= \frac{3e}{2}\end{aligned}$$

Therefore:

$$\begin{aligned}\oint_{\gamma} \frac{e^z}{z^2(z-1)^3} dz &= 2\pi i \left[-4 + \frac{3e}{2} \right] \\ &= \pi i (3e - 8)\end{aligned}$$

▲ **Example 7.6.7** Evaluate

$$I = \oint_{\gamma} \frac{e^{1/z}}{1+z} dz,$$

where γ is the circle $|z| = 2$ oriented counterclockwise.

Solution The function

$$f(z) = \frac{e^{1/z}}{1+z}$$

has two singularities inside γ :

- An essential singularity at $z = 0$ (from $e^{1/z}$),

- A simple pole at $z = -1$ (since $1 + z = 0$).

By the Residue Theorem:

$$I = 2\pi i [\text{Res}(f, 0) + \text{Res}(f, -1)].$$

For the residue at $z = -1$, which is a simple pole:

$$\begin{aligned} \text{Res}(f, -1) &= \lim_{z \rightarrow -1} (z + 1)f(z) \\ &= \lim_{z \rightarrow -1} \frac{(z + 1) \cdot e^{1/z}}{1 + z} \\ &= \lim_{z \rightarrow -1} e^{1/z} \\ &= e^{-1} \end{aligned}$$

For the residue at $z = 0$, which is an essential singularity, we need to find the coefficient of z^{-1} in the Laurent series expansion. We expand each factor:

$$e^{1/z} = \sum_{n=0}^{\infty} \frac{1}{n!} \frac{1}{z^n}$$

and

$$\frac{1}{1+z} = \sum_{m=0}^{\infty} (-1)^m z^m \quad \text{for } |z| < 1$$

Multiplying these series:

$$\frac{e^{1/z}}{1+z} = \sum_{n=0}^{\infty} \sum_{m=0}^{\infty} \frac{(-1)^m}{n!} z^{m-n}$$

The z^{-1} term appears when $m - n = -1$, or $m = n - 1$. Therefore:

$$\begin{aligned} \text{Res}(f, 0) &= \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n!} \\ &= - \sum_{n=1}^{\infty} \frac{(-1)^n}{n!} \\ &= - \left(\sum_{n=0}^{\infty} \frac{(-1)^n}{n!} - 1 \right) \\ &= - (e^{-1} - 1) \\ &= 1 - e^{-1} \end{aligned}$$

Adding the residues:

$$\text{Res}(f, 0) + \text{Res}(f, -1) = (1 - e^{-1}) + e^{-1}$$

$$= 1$$

Therefore, by the Residue Theorem:

$$I = 2\pi i \cdot 1 = 2\pi i$$

Applications to Real Definite Integrals

One of the most powerful applications of residue theory is the evaluation of real definite integrals that would be difficult or impossible to compute using elementary calculus techniques. The strategy involves transforming a real integral into a complex contour integral, applying the residue theorem, and then extracting the value of the original integral.

This approach is particularly effective for three classes of integrals:

1. **Rational Functions on the Entire Real Line:** Integrals of the form $\int_{-\infty}^{\infty} \frac{P(x)}{Q(x)} dx$ where P and Q are polynomials with $\deg(P) < \deg(Q)$.
2. **Trigonometric Integrals:** Integrals of the form $\int_0^{2\pi} R(\cos \theta, \sin \theta) d\theta$ where R is a rational function.
3. **Integrals with Exponential Factors:** Integrals like $\int_{-\infty}^{\infty} \frac{e^{iax}}{P(x)} dx$ or $\int_{-\infty}^{\infty} \frac{\cos(ax)}{P(x)} dx$.

Key Transformations and Techniques

Several transformations and identities facilitate the conversion between real and complex integrals:

- **Substitution $z = e^{i\theta}$:** This transforms trigonometric functions:

$$\begin{aligned}\cos \theta &= \frac{z + z^{-1}}{2} \\ \sin \theta &= \frac{z - z^{-1}}{2i} \\ d\theta &= \frac{dz}{iz}\end{aligned}$$

With this substitution, $\int_0^{2\pi} f(\cos \theta, \sin \theta) d\theta$ becomes $\oint_{|z|=1} f\left(\frac{z+z^{-1}}{2}, \frac{z-z^{-1}}{2i}\right) \frac{dz}{iz}$.

- **Contour Selection:** For integrals over $(-\infty, \infty)$, we often use a semicircular contour in the upper half-plane, consisting of the real axis from $-R$ to R and a semicircle Γ_R of radius R in the upper half-plane.

- **Jordan's Lemma:** For integrals of the form $\int_{-\infty}^{\infty} f(x)e^{iax} dx$ with $a > 0$, the contribution from the upper semicircular contour Γ_R vanishes as $R \rightarrow \infty$ if $f(z) \rightarrow 0$ as $|z| \rightarrow \infty$ in the upper half-plane.
- **Evaluation of Real Integrals Using Complex Methods:**

$$\begin{aligned}\int_{-\infty}^{\infty} \frac{P(x)}{Q(x)} dx &= 2\pi i \sum_{\text{Im}(z_k) > 0} \text{Res} \left(\frac{P(z)}{Q(z)}, z_k \right) \\ \int_0^{\infty} f(x) dx &= \int_0^{\infty} f(x) dx + i \int_0^{\infty} f(ix) dx \\ &= 2\pi i \sum_{\text{Re}(z_k) > 0, \text{Im}(z_k) > 0} \text{Res}(f(z), z_k)\end{aligned}$$

where z_k are the poles in the specified regions.

Important Relations for Trigonometric Integrals

When dealing with trigonometric integrals, these identities are particularly useful:

$$\begin{aligned}\cos^2 \theta &= \frac{1 + \cos 2\theta}{2} = \frac{1}{2} + \frac{z^2 + z^{-2}}{4} \\ \sin^2 \theta &= \frac{1 - \cos 2\theta}{2} = \frac{1}{2} - \frac{z^2 + z^{-2}}{4} \\ \cos^n \theta &= \frac{1}{2^n} \sum_{k=0}^n \binom{n}{k} \cos((n-2k)\theta) \\ \sin^n \theta &= \frac{1}{2^n} \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \binom{n}{k} \sin((n-2k)\theta)\end{aligned}$$

After substituting $z = e^{i\theta}$, a rational function of trigonometric functions transforms into a rational function of z , allowing us to apply residue theory effectively.

▲ **Example 7.6.8** Evaluate the challenging trigonometric integral:

$$I = \int_0^{2\pi} \frac{d\theta}{(2 + \cos \theta)^2}$$

Solution This integral appears formidable using standard techniques. We'll solve it using residue theory with the substitution $z = e^{i\theta}$.

First, let's express $\cos \theta$ in terms of z :

$$\cos \theta = \frac{z + z^{-1}}{2}$$

and note that:

$$d\theta = \frac{dz}{iz}$$

The integral becomes:

$$\begin{aligned} I &= \int_0^{2\pi} \frac{d\theta}{(2 + \cos \theta)^2} \\ &= \oint_{|z|=1} \frac{1}{\left(2 + \frac{z+z^{-1}}{2}\right)^2} \cdot \frac{dz}{iz} \\ &= \oint_{|z|=1} \frac{1}{\left(\frac{4z+z^2+1}{2z}\right)^2} \cdot \frac{dz}{iz} \\ &= \frac{1}{i} \oint_{|z|=1} \frac{4z}{(z^2 + 4z + 1)^2} dz \end{aligned}$$

We need to find the poles of the integrand by solving $z^2 + 4z + 1 = 0$:

$$\begin{aligned} z &= \frac{-4 \pm \sqrt{16 - 4}}{2} \\ &= \frac{-4 \pm \sqrt{12}}{2} \\ &= -2 \pm \sqrt{3} \end{aligned}$$

This gives $z_1 = -2 + \sqrt{3} \approx -0.268$ and $z_2 = -2 - \sqrt{3} \approx -3.732$. Since $|z_1| < 1$ and $|z_2| > 1$, only z_1 lies inside the unit circle. The pole at z_1 is of order 2, so we use the formula for a second-order pole:

$$\begin{aligned} \text{Res}(f, z_1) &= \lim_{z \rightarrow z_1} \frac{d}{dz} \left[(z - z_1)^2 \cdot \frac{4z}{(z^2 + 4z + 1)^2} \right] \\ &= \lim_{z \rightarrow z_1} \frac{d}{dz} \left[\frac{4z}{(z - z_1)^2 \cdot (z - z_2)^2} \cdot (z - z_1)^2 \right] \\ &= \lim_{z \rightarrow z_1} \frac{d}{dz} \left[\frac{4z}{(z - z_2)^2} \right] \end{aligned}$$

Computing the derivative:

$$\begin{aligned} \frac{d}{dz} \left[\frac{4z}{(z - z_2)^2} \right] &= \frac{4 \cdot (z - z_2)^2 - 4z \cdot 2(z - z_2)}{(z - z_2)^4} \\ &= \frac{4(z - z_2)^2 - 8z(z - z_2)}{(z - z_2)^4} \\ &= \frac{-4(z + z_2)}{(z - z_2)^3} \end{aligned}$$

Evaluating at $z = z_1$:

$$\begin{aligned}\operatorname{Res}(f, z_1) &= \frac{-4(z_1 + z_2)}{(z_1 - z_2)^3} \\ &= \frac{-4(-2 + \sqrt{3} - 2 - \sqrt{3})}{((-2 + \sqrt{3}) - (-2 - \sqrt{3}))^3} \\ &= \frac{2\sqrt{3}}{9}\end{aligned}$$

By the Residue Theorem:

$$\begin{aligned}I &= \frac{1}{i} \cdot 2\pi i \cdot \operatorname{Res}(f, z_1) \\ &= \frac{4\pi\sqrt{3}}{9}\end{aligned}$$

Therefore:

$$\int_0^{2\pi} \frac{d\theta}{(2 + \cos \theta)^2} = \frac{4\pi\sqrt{3}}{9}$$

This example demonstrates the power of residue theory in evaluating trigonometric integrals that would be extremely challenging using elementary techniques.

Formulas and Definitions

Important Vocabulary

- **Complex Number:** A number of the form $z = a + bi$ where a and b are real numbers and i is the imaginary unit satisfying $i^2 = -1$.
- **Real Part:** For $z = a + bi$, the real part $\operatorname{Re}(z) = a$.
- **Imaginary Part:** For $z = a + bi$, the imaginary part $\operatorname{Im}(z) = b$.
- **Complex Conjugate:** For $z = a + bi$, the complex conjugate $\bar{z} = a - bi$.
- **Modulus/Absolute Value:** For $z = a + bi$, the modulus $|z| = \sqrt{a^2 + b^2}$.
- **Argument:** The angle $\theta = \arg(z)$ such that $z = |z|e^{i\theta}$.
- **Holomorphic Function:** A complex function that is complex differentiable in a neighborhood of every point in its domain.
- **Analytic Function:** A function that can be represented by a convergent power series in a neighborhood of each point in its domain.
- **Singularity:** A point where a function is not holomorphic.
- **Pole:** A singularity where a function becomes unbounded.
- **Essential Singularity:** A non-removable singularity that is not a pole.
- **Residue:** The coefficient of $(z - z_0)^{-1}$ in the Laurent series expansion of a function around z_0 .
- **Branch Cut:** A curve in the complex plane across which a multi-valued function is discontinuous.
- **Conformal Mapping:** A transformation that locally preserves angles.
- **Simply Connected Domain:** A domain where every simple closed curve can be continuously shrunk to a point within the domain.

Key Formulas

Basic Operations:

$$(a + bi) + (c + di) = (a + c) + (b + d)i$$

$$(a + bi)(c + di) = (ac - bd) + (ad + bc)i$$

$$\frac{a + bi}{c + di} = \frac{(a + bi)(c - di)}{(c + di)(c - di)} = \frac{ac + bd}{c^2 + d^2} + \frac{bc - ad}{c^2 + d^2}i$$

Complex Conjugate:

$$\bar{z} = \overline{a + bi} = a - bi$$

$$\overline{z_1 + z_2} = \bar{z}_1 + \bar{z}_2$$

$$\overline{z_1 \cdot z_2} = \bar{z}_1 \cdot \bar{z}_2$$

$$z \cdot \bar{z} = |z|^2 = a^2 + b^2$$

Modulus/Absolute Value:

$$|z| = \sqrt{a^2 + b^2} = \sqrt{z\bar{z}}$$

$$|z_1 \cdot z_2| = |z_1| \cdot |z_2|$$

$$\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|}$$

Polar Form:

$$z = a + bi = r(\cos \theta + i \sin \theta) = re^{i\theta}$$

$$r = |z| = \sqrt{a^2 + b^2}$$

$$\theta = \arg(z) = \arctan\left(\frac{b}{a}\right)$$

Euler's Formula:

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$e^z = e^x(\cos y + i \sin y) \quad \text{for } z = x + iy$$

De Moivre's Theorem:

$$(re^{i\theta})^n = r^n e^{in\theta} = r^n(\cos(n\theta) + i \sin(n\theta))$$

 n th Roots of Complex Numbers:

$$w_k = \sqrt[n]{r} e^{i(\theta + 2\pi k)/n}, \quad k = 0, 1, \dots, n-1$$

Roots of Unity:

$$\zeta_k = e^{2\pi i k/n}, \quad k = 0, 1, \dots, n-1$$

Exponential Function:

$$e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!}$$

Trigonometric Functions:

$$\sin(z) = \frac{e^{iz} - e^{-iz}}{2i}$$

$$\cos(z) = \frac{e^{iz} + e^{-iz}}{2}$$

Complex Logarithm:

$$\ln(z) = \ln|z| + i \arg(z)$$

$$\text{Log}(z) = \ln|z| + i \text{Arg}(z) \quad (\text{principal branch})$$

Complex Derivatives:

$$f'(z_0) = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}$$

Cauchy-Riemann Equations: For $f(z) = u(x, y) + iv(x, y)$, where $z = x + iy$:

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

Polar Cauchy-Riemann Equations:

$$\frac{\partial u}{\partial r} = \frac{1}{r} \frac{\partial v}{\partial \theta}, \quad \frac{\partial v}{\partial r} = -\frac{1}{r} \frac{\partial u}{\partial \theta}$$

Rules of Differentiation:

$$(f + g)'(z) = f'(z) + g'(z)$$

$$(f \cdot g)'(z) = f'(z)g(z) + f(z)g'(z)$$

$$\left(\frac{f}{g}\right)'(z) = \frac{f'(z)g(z) - f(z)g'(z)}{[g(z)]^2}$$

$$(f \circ g)'(z) = f'(g(z))g'(z)$$

Power Series:

$$f(z) = \sum_{n=0}^{\infty} a_n(z - z_0)^n$$

$$a_n = \frac{f^{(n)}(z_0)}{n!}$$

Complex Line Integral:

$$\int_{\gamma} f(z) dz = \int_a^b f(\gamma(t)) \gamma'(t) dt$$

Complex Line Integral in Component Form:

$$\int_{\gamma} f(z) dz = \int_a^b [u \cdot x' - v \cdot y' + i(u \cdot y' + v \cdot x')] dt$$

ML Inequality:

$$\left| \int_{\gamma} f(z) dz \right| \leq L(\gamma) \cdot \max_{z \in \gamma} |f(z)|$$

Fundamental Theorem for Line Integrals:

$$\int_{\gamma} f(z) dz = F(z_2) - F(z_1)$$

where $F'(z) = f(z)$ and γ is a path from z_1 to z_2 .

Cauchy's Integral Theorem:

$$\oint_{\gamma} f(z) dz = 0$$

for f holomorphic within and on a simple closed contour γ .

Cauchy's Integral Formula:

$$f(a) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{z-a} dz$$

where γ is a simple closed contour containing a .

Cauchy's Formula for Derivatives:

$$f^{(n)}(a) = \frac{n!}{2\pi i} \oint_{\gamma} \frac{f(z)}{(z-a)^{n+1}} dz$$

Cauchy's Inequality:

$$|f^{(n)}(a)| \leq \frac{n! \cdot M}{R^n}$$

where $M = \max_{|z-a|=R} |f(z)|$.

Laurent Series:

$$\begin{aligned} f(z) &= \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n \\ &= \sum_{n=0}^{\infty} a_n (z - z_0)^n + \sum_{n=1}^{\infty} \frac{a_{-n}}{(z - z_0)^n} \end{aligned}$$

Laurent Series Coefficients:

$$a_n = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{(z - z_0)^{n+1}} dz$$

Residue Theorem:

$$\oint_{\gamma} f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f, z_k)$$

where z_k are the poles of f inside γ .

Residue Computation (Simple Pole):

$$\text{Res}(f, z_0) = \lim_{z \rightarrow z_0} (z - z_0) f(z)$$

Residue for Simple Pole with Special Form:

$$\text{Res}(f, z_0) = \frac{p(z_0)}{q'(z_0)}$$

for $f(z) = \frac{p(z)}{q(z)}$ where $q(z_0) = 0$ and $q'(z_0) \neq 0$

Residue Computation (Pole of Order m):

$$\text{Res}(f, z_0) = \frac{1}{(m-1)!} \lim_{z \rightarrow z_0} \frac{d^{m-1}}{dz^{m-1}} [(z - z_0)^m f(z)]$$

Winding Number:

$$n(\gamma, a) = \frac{1}{2\pi i} \oint_{\gamma} \frac{dz}{z - a}$$

Integration Along Unit Circle:

$$\oint_{|z|=1} z^n dz = \begin{cases} 0 & \text{if } n \neq -1 \\ 2\pi i & \text{if } n = -1 \end{cases}$$

Practice Problems

1. Compute the modulus and complex conjugate of the number $3 - 4i$. Verify that

$$(3 - 4i)(3 + 4i) = |3 - 4i|^2.$$

2. Given $z_1 = 2 + 3i$ and $z_2 = -1 + 4i$, find:

- (a) $z_1 + z_2$

- (b) $z_1 - z_2$

- (c) $z_1 z_2$

- (d) $\frac{z_1}{z_2}$

Express all answers in the form $a + bi$.

3. Express the complex number $-1 - i$ in polar form by determining its modulus and argument.
4. Use Euler's formula to show that

$$\cos \theta + i \sin \theta = e^{i\theta},$$

and compare the first few terms of the Taylor series on both sides to verify the equality.

5. Use De Moivre's theorem to compute

$$\left(\frac{\sqrt{2}}{2} + i \frac{\sqrt{2}}{2} \right)^5,$$

and then express your answer in rectangular form.

6. Find all cube roots of $8i$. Express your answers both in polar form and in the form $a + bi$.
7. Solve the equation

$$z^2 + 4 = 0$$

using polar representation, and write the solutions in $a + bi$ form.

8. Describe the geometric transformation on the complex plane corresponding to multiplication by $-i$. Choose a nonzero complex number to illustrate how its position changes.

9. Find all complex numbers z satisfying

$$|z - (1 + 2i)| = 3.$$

Describe the geometric locus of these points in the Argand plane.

10. Determine all the 5th roots of unity, and sketch their approximate positions on the complex plane.
11. Verify that for $z = 5 - 2i$, the product $z\bar{z}$ equals $|z|^2$ by direct computation.
12. Determine the locus of points z in the complex plane that satisfy

$$|z - 2| = |z + 2|.$$

Explain the geometric significance of this locus.

13. Factorize the polynomial

$$z^3 - 8$$

completely over $\mathbb{C}$, and find all its roots.

14. Solve the equation

$$z^4 + 16 = 0$$

by expressing the solutions in polar form and then converting them to rectangular form.

15. Solve the quadratic equation

$$z^2 + (3 - 2i)z + (5 - i) = 0$$

using the quadratic formula for complex coefficients, and express the solutions in the form $a + bi$.

16. For
- $f(z) = z^2$
- with
- $z = x + iy$
- , write
- $f(z) = u(x, y) + iv(x, y)$
- by expanding
- $(x + iy)^2$
- . Identify
- $u(x, y)$
- and
- $v(x, y)$
- . Then, show that if
- z
- lies on the real axis, its image under
- f
- is also real.

17. Express
- $f(z) = \frac{1}{z}$
- in terms of
- x
- and
- y
- for
- $z = x + iy$
- (by rationalizing the denominator) and identify the real function
- $u(x, y)$
- and the imaginary function
- $v(x, y)$
- . Also, state the set of points where
- $f(z)$
- is undefined.

18. For
- $f(z) = e^z$
- , write
- $f(z) = e^x(\cos y + i \sin y)$
- when
- $z = x + iy$
- . Describe the image of the vertical line
- $x = 0$
- (i.e.
- $z = iy$
-) in the
- w
- plane.

19. Determine the natural domain of
- $f(z) = \ln(z)$
- in the complex plane and explain how and why a branch cut (e.g. along the negative real axis) is introduced to obtain a single-valued principal branch.

20. Starting from the definition

$$\sin(z) = \frac{e^{iz} - e^{-iz}}{2i},$$

derive the form

$$\sin(z) = \sin x \cosh y + i \cos x \sinh y,$$

for $z = x + iy$.

21. Solve
- $e^z = -2$
- for
- z
- by writing
- $z = x + iy$
- and expressing the answer in the form
- $x + iy$
- . List all solutions.

22. Let
- $f(z) = z^3$
- . Show that if
- z
- lies on the unit circle (i.e.
- $|z| = 1$
-), then
- $f(z)$
- also lies on the unit circle and its argument is tripled. Verify this with
- $z = e^{i\pi/3}$
- .

23. Consider
- $f(z) = \frac{z^2+1}{z-i}$
- . Determine its domain in
- $\mathbb{C}$
- and identify any singularities. Specify the type and order (if applicable) of each singularity.

24. Define
- $f(z) = \sqrt{z}$
- using the principal branch with a branch cut along the negative real axis. Compute
- $f(4e^{i\pi})$
- and express your answer in the form
- $a + bi$
- .

25. For
- $f(z) = \ln(z)$
- , compute all values of
- $\ln(-1 - i)$
- in terms of its modulus and argument. Then, specify the principal value using the conventional range
- $(-\pi, \pi]$
- for the argument.

26. Determine the image of the horizontal line
- $y = 1$
- (i.e.
- $z = x + i$
-) under the mapping
- $f(z) = e^z$
- . Describe the geometric nature of the resulting curve in the
- w
- plane.

27. Solve the equation
- $\cos(z) = 0$
- for all
- $z \in \mathbb{C}$
- . (Hint: Use the definition
- $\cos(z) = \frac{e^{iz} + e^{-iz}}{2}$
- and set up an exponential equation.)

28. For
- $f(z) = z + \frac{1}{z}$
- , determine the condition(s) on
- z
- so that
- $f(z)$
- is a real number. Express your answer in terms of the real and imaginary parts of
- z
- .

29. Consider
- $f(z) = \frac{z-1}{z+1}$
- . Prove that
- f
- maps the imaginary axis (i.e.
- $z = iy$
- with
- $y \in \mathbb{R}$
-) into the unit circle.

30. Let
- $f(z) = e^{z^2}$
- . Express
- $f(z)$
- in terms of
- x
- and
- y
- for
- $z = x + iy$
- .

31. Evaluate the following limits:

$$(a) \lim_{z \rightarrow 0} z^3.$$

- (b) $\lim_{z \rightarrow 0} \frac{\sin z}{z}$.
- (c) $\lim_{z \rightarrow 0} \frac{e^z - 1}{z}$.
- (d) $\lim_{z \rightarrow 0} \frac{1 - \cos z}{z^2}$.
32. Using the Taylor series expansion for $\sin z$, evaluate
- $$\lim_{z \rightarrow 0} \frac{\sin z - z}{z^3}.$$
33. Compute
- $$\lim_{z \rightarrow 1+i} \frac{z^3 - (1+i)^3}{z - (1+i)}.$$
- (Hint: Factor the numerator or interpret the limit as the derivative of z^3 at $z = 1 + i$.)
34. Prove, using the epsilon–delta definition, that for any integer $n \geq 1$
- $$\lim_{z \rightarrow 0} z^n = 0.$$
35. Using polar form, prove that $\lim_{r \rightarrow 0} r e^{2i\theta} = 0$ (the limit is independent of θ).
36. Let $f(z) = \frac{z^2 - \bar{z}^2}{z - \bar{z}}$ (with $z \notin \mathbb{R}$). Evaluate $\lim_{z \rightarrow 2+i} f(z)$.
37. Show that by writing $z = r e^{i\theta}$
- $$\lim_{r \rightarrow 0} r e^{2i\theta} = 0,$$
- and explain why this limit does not depend on the angle θ .
38. Prove that
- $$\lim_{z \rightarrow 0} \frac{\bar{z}}{z}$$
- does not exist by showing that the value depends on the direction (i.e. on θ) along which z approaches 0.
39. (a) Evaluate $\lim_{|z| \rightarrow \infty} \frac{1}{z}$ in the extended complex plane.
- (b) Determine whether the limit
- $$\lim_{z \rightarrow \infty} \frac{z}{|z|}$$
- exists; justify your answer.
40. Evaluate
- $$\lim_{z \rightarrow 0} \frac{e^z - e^{\bar{z}}}{z - \bar{z}}.$$
- (Hint: Express $z = x + iy$ so that $z - \bar{z} = 2iy$, then simplify using the series for e^z .)
41. Write down the epsilon–delta definition of the limit
- $$\lim_{z \rightarrow z_0} f(z) = L$$
- for a complex function f , and explain why path independence (i.e. the fact that the limit must be the same along every approach to z_0) is crucial in $\mathbb{C}$.
42. Describe the extended complex plane $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ and, using the epsilon–delta idea adapted for infinity, prove that
- $$\lim_{z \rightarrow 0} \frac{1}{z} = \infty.$$
43. Using the definition of the derivative, compute
- $$f'(z) \quad \text{for} \quad f(z) = z^2.$$
44. Using the limit definition, show that
- $$(e^z)' = e^z.$$
45. Using the limit definition (and the power series of the logarithm), prove that on its principal branch
- $$(\text{Log } z)' = \frac{1}{z} \quad \text{for } z \notin (-\infty, 0].$$

46. Compute the derivative of

$$f(z) = \sin z$$

by directly applying the definition.

47. Use the limit definition to differentiate

$$f(z) = \frac{1}{z},$$

for $z \neq 0$.

48. Show that the function

$$f(z) = |z|^2$$

is not complex differentiable at any point $z \neq 0$ (by considering different paths of approach).

49. Prove that the function

$$f(z) = \operatorname{Re}(z)$$

is nowhere differentiable in $\mathbb{C}$.

50. Write
- $f(z) = z^3$
- in Cartesian form and verify that its real and imaginary components satisfy the Cauchy-Riemann equations.

51. Check that the function

$$f(z) = \bar{z}$$

fails the Cauchy-Riemann equations, and conclude that it is not holomorphic.

52. Prove (outline a proof) the chain rule for complex functions: that if
- g
- is holomorphic at
- z_0
- and
- f
- is holomorphic at
- $g(z_0)$
- , then

$$(f \circ g)'(z_0) = f'(g(z_0)) \cdot g'(z_0).$$

53. Differentiate the functions

$$(a) \quad f(z) = z^2 \cdot e^z$$

$$(b) \quad f(z) = \sin(e^z).$$

$$(c) \quad f(z) = z^3 + 2z^2 - 3z + 1$$

54. For the function defined on the principal branch by

$$f(z) = z^{3/2},$$

express z in polar form and compute $f'(z)$.

55. Prove that if
- $f'(z) = 0$
- for all
- z
- in a connected domain, then
- f
- is constant. (Hint: Use the uniqueness of antiderivatives.)

56. Express the function

$$f(z) = z^2$$

in polar coordinates. Derive the polar form of the Cauchy-Riemann equations

$$u_r = \frac{1}{r} v_\theta \quad \text{and} \quad v_r = -\frac{1}{r} u_\theta,$$

and verify that these hold for $f(z)$.

57. Show that

$$u(x, y) = \frac{1}{2} \ln(x^2 + y^2)$$

is harmonic on $\mathbb{R}^2 \setminus \{(0, 0)\}$; that is, verify that

$$u_{xx}(x, y) + u_{yy}(x, y) = 0$$

for all $(x, y) \neq (0, 0)$.

58. Prove that

$$u(x, y) = e^x \cos y$$

is a harmonic function on $\mathbb{R}^2$; in other words, show that it satisfies Laplace's equation

$$u_{xx} + u_{yy} = 0.$$

59. Verify that the real part of
- z^3
- , namely

$$u(x, y) = x^3 - 3xy^2,$$

is harmonic in $\mathbb{R}^2$ by showing that

$$u_{xx}(x, y) + u_{yy}(x, y) = 0.$$

60. Find the power series expansion of

$$f(z) = \frac{1}{1-z}$$

about $z = 0$ and determine its radius of convergence.

61. Expand

$$f(z) = \text{Log}(1+z)$$

in a Taylor series around $z = 0$ (using the principal branch) and state its radius of convergence.

62. Let
- γ
- be the straight line segment from
- $z = 1 + 0i$
- to
- $z = 1 + 2i$
- . Compute the integral

$$\int_{\gamma} dz.$$

63. Let
- γ
- be the straight line from
- $z = 0$
- to
- $z = 1 + i$
- . Compute

$$\int_{\gamma} e^z dz$$

by parametrizing γ appropriately.

64. Compute

$$\int_{\gamma} z^2 dz,$$

where γ is the circle $z = 2e^{it}$ with $t \in [0, 2\pi]$. (Hint: Use the fact that z^2 is entire and explain why the integral vanishes.)

65. Let
- γ_R
- be the semicircular arc defined by

$$z = Re^{it}, \quad t \in [0, \pi].$$

Use the ML inequality to show that, for $k > 0$,

$$\left| \int_{\gamma_R} e^{ikz} dz \right|$$

tends to zero as $R \rightarrow \infty$.

66. Evaluate

$$\int_{\gamma} \frac{dz}{z},$$

where γ is the unit circle $z = e^{it}$, $t \in [0, 2\pi]$. Explain briefly why the answer reflects the winding number.

67. Find an antiderivative
- $F(z)$
- of
- $f(z) = z^3$
- and use the Fundamental Theorem for Line Integrals to evaluate

$$\int_{\gamma} z^3 dz,$$

where γ is any piecewise smooth curve joining $z = 1$ and $z = 2$.

68. Let
- γ
- be parametrized by
- $\gamma(t) = t + it$
- for
- $t \in [0, 1]$
- . Compute

$$\int_{\gamma} \sin z dz.$$

69. Evaluate

$$\int_{\gamma} \frac{dz}{z-1},$$

where γ is the circle $|z| = 2$ traversed counterclockwise. (Hint: Since $z = 1$ lies inside γ , the answer will involve $2\pi i$.)

70. Prove that if
- $f(z)$
- is holomorphic on a domain
- D
- and has an antiderivative
- $F(z)$
- in
- D
- , then

$$\int_{\gamma} f(z) dz = F(z_2) - F(z_1)$$

for any piecewise smooth curve γ from z_1 to z_2 in D . Briefly explain why this result implies that the integral is path-independent.

71. Evaluate

$$\int_{\gamma} \frac{z}{z^2 + 1} dz,$$

where γ is the line segment from $z = 0$ to $z = 2$ (assuming the segment avoids any singularities). (Hint: Find an antiderivative and apply the Fundamental Theorem.)

72. Show that reparametrizing a piecewise smooth curve does not change the value of the complex line integral. Briefly describe the invariance property.

73. Let γ be a closed contour that winds n times counterclockwise around the origin. Prove that for $f(z) = 1/z$ the integral satisfies

$$\oint_{\gamma} \frac{dz}{z} = 2\pi i n.$$

74. Let $f(z) = e^z$. Prove that for any closed, piecewise smooth curve γ in $\mathbb{C}$,

$$\oint_{\gamma} e^z dz = 0.$$

75. Let γ be the unit circle $z = e^{it}$ with $t \in [0, 2\pi]$ (oriented counterclockwise). Show that

$$\oint_{\gamma} \frac{dz}{z} = 2\pi i.$$

76. Use Cauchy's Integral Formula to evaluate

$$\oint_{|z|=3} \frac{e^z}{z-2} dz.$$

77. Apply the generalized Cauchy Integral Formula to compute the second derivative $f^{(2)}(0)$ for $f(z) = \sin z$ via

$$f^{(2)}(0) = \frac{2!}{2\pi i} \oint_{|z|=R} \frac{\sin z}{z^3} dz.$$

78. Evaluate

$$\oint_{|z|=2} \frac{dz}{z-1},$$

and explain why the answer depends on the fact that $z = 1$ lies inside the contour.

79. Let γ be the circle centered at $z = 1$ of radius $r > 0$. Compute

$$\oint_{\gamma} \frac{dz}{(z-1)^2},$$

and explain how the order of the pole affects the result.

80. Evaluate

$$\oint_{|z|=2} \frac{dz}{z(z-1)},$$

indicating which singularities are enclosed by the contour.

81. Find the Laurent series expansion of

$$f(z) = \frac{1}{z(z-2)}$$

valid in the annulus $0 < |z| < 2$ and classify the singularity at $z = 0$.

82. Compute the residue of

$$f(z) = \frac{e^{1/z}}{1+z}$$

at $z = 0$ by finding the coefficient a_{-1} in its Laurent expansion.

83. Find the Laurent series for

$$f(z) = \frac{z^2 + 1}{z(z-1)(z-2)}$$

about $z = 0$ for $0 < |z| < 1$ and determine the type of singularity at $z = 0$.

84. Use the Residue Theorem to evaluate

$$\oint_{|z|=3} \frac{e^z}{z^2 + 1} dz.$$

85. Evaluate the real integral

$$\int_{-\infty}^{\infty} \frac{dx}{x^2 + 1}$$

by interpreting it as a contour integral and applying residue theory.

86. Prove Liouville's Theorem: Show that if $f(z)$ is entire and there exists an $M > 0$ such that $|f(z)| \leq M$ for all

$z \in \mathbb{C}$, then $f(z)$ is constant. **Hint:** Consider the power series expansion of f and use Cauchy's estimates.

87. Evaluate

$$\oint_{|z-i|=R} \frac{e^{z^2}}{(z-i)^3} dz,$$

where $R > 0$ is sufficiently small so that the circle is contained in a region where e^{z^2} is holomorphic. Express your answer in terms of $f^{(2)}(i)$ for $f(z) = e^{z^2}$ and then compute $f^{(2)}(i)$.

Appendices

Introduction

These appendices review all of the formulas for this book. I highly emphasize that teachers let their students use these during quizzes and exams. Upon completion of basic differential equations proficiency, it is reasonable to assume that a student should be able to look up what they need quickly in order to solve a question on an exam, and hence a question that they may face in real life. Students in this day and age will always have something to reference, either Google, ChatGPT, or this book even. In my opinion, if one is able to quickly produce an answer and be confident that it is correct, then all resources are therefore valid to use.

Appendix A: Derivative and Integral Identities

Derivative Identities

Basic Rules

- $\frac{d}{dx}(c) = 0$, where c is a constant
- $\frac{d}{dx}(x^n) = nx^{n-1}$
- $\frac{d}{dx}(e^x) = e^x$
- $\frac{d}{dx}(\ln|x|) = \frac{1}{x}$, $x \neq 0$
- $\frac{d}{dx}(a^x) = a^x \ln(a)$, $a > 0$
- $\frac{d}{dx}(\sin x) = \cos x$
- $\frac{d}{dx}(\cos x) = -\sin x$
- $\frac{d}{dx}(\tan x) = \sec^2 x$
- $\frac{d}{dx}(\cot x) = -\csc^2 x$
- $\frac{d}{dx}(\sec x) = \sec x \tan x$
- $\frac{d}{dx}(\csc x) = -\csc x \cot x$

Product Rule

- $\frac{d}{dx}(f(x)g(x)) = f'(x)g(x) + f(x)g'(x)$

Quotient Rule

- $\frac{d}{dx}\left(\frac{f(x)}{g(x)}\right) = \frac{f'(x)g(x) - f(x)g'(x)}{[g(x)]^2}$

Chain Rule

- $\frac{d}{dx}(f(g(x))) = f'(g(x))g'(x)$

Inverse Function Rule

- $\frac{d}{dx}(f^{-1}(x)) = \frac{1}{f'(f^{-1}(x))}$

Inverse Trigonometric Functions

- $\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$, $|x| < 1$
- $\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$, $|x| < 1$
- $\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}$
- $\frac{d}{dx}(\operatorname{arccot} x) = -\frac{1}{1+x^2}$
- $\frac{d}{dx}(\operatorname{arcsec} x) = \frac{1}{|x|\sqrt{x^2-1}}$, $|x| > 1$
- $\frac{d}{dx}(\operatorname{arccsc} x) = -\frac{1}{|x|\sqrt{x^2-1}}$, $|x| > 1$

Hyperbolic Functions

- $\frac{d}{dx}(\sinh x) = \cosh x$
- $\frac{d}{dx}(\cosh x) = \sinh x$
- $\frac{d}{dx}(\tanh x) = \operatorname{sech}^2 x$
- $\frac{d}{dx}(\coth x) = -\operatorname{csch}^2 x$
- $\frac{d}{dx}(\operatorname{sech} x) = -\operatorname{sech} x \tanh x$
- $\frac{d}{dx}(\operatorname{csch} x) = -\operatorname{csch} x \coth x$

Integral Identities

Basic Integrals

- $\int x^n dx = \frac{x^{n+1}}{n+1} + C, n \neq -1$
- $\int \frac{1}{x} dx = \ln |x| + C$
- $\int e^x dx = e^x + C$
- $\int a^x dx = \frac{a^x}{\ln a} + C, a > 0, a \neq 1$
- $\int \sin x dx = -\cos x + C$
- $\int \cos x dx = \sin x + C$
- $\int \tan x dx = -\ln |\cos x| + C$
- $\int \cot x dx = \ln |\sin x| + C$
- $\int \sec x dx = \ln |\sec x + \tan x| + C$
- $\int \csc x dx = -\ln |\csc x + \cot x| + C$

Trigonometric Integrals

- $\int \sin^2 x dx = \frac{x}{2} - \frac{\sin(2x)}{4} + C$
- $\int \cos^2 x dx = \frac{x}{2} + \frac{\sin(2x)}{4} + C$
- $\int \tan^2 x dx = \tan x - x + C$
- $\int \sec^2 x dx = \tan x + C$
- $\int \csc^2 x dx = -\cot x + C$

Integration by Parts

- $\int u dv = uv - \int v du$

Integration by Substitution

- $\int f(g(x))g'(x)dx = \int f(u)du$, where $u = g(x)$

Partial Fractions

For $\frac{P(x)}{Q(x)}$ where degree of $P <$ degree of Q :

- Linear factors: $\frac{A}{(x-a)} + \frac{B}{(x-b)} + \dots$
- Repeated linear factors: $\frac{A}{(x-a)} + \frac{B}{(x-a)^2} + \dots$
- Quadratic factors: $\frac{Ax+B}{x^2+px+q} + \dots$

Trigonometric Substitutions

- $\sqrt{a^2 - x^2}$: use $x = a \sin \theta$
- $\sqrt{a^2 + x^2}$: use $x = a \tan \theta$
- $\sqrt{x^2 - a^2}$: use $x = a \sec \theta$

Appendix B: Vector Properties and Formulas

Vector Algebra

- Addition: $\vec{u} + \vec{v} = \langle u_1 + v_1, u_2 + v_2, u_3 + v_3 \rangle$
- Scalar Multiplication: $c\vec{u} = \langle cu_1, cu_2, cu_3 \rangle$
- Dot Product: $\vec{u} \cdot \vec{v} = u_1v_1 + u_2v_2 + u_3v_3 = |\vec{u}||\vec{v}|\cos\theta$
- Cross Product: $\vec{u} \times \vec{v} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$
- Magnitude: $|\vec{u}| = \sqrt{u_1^2 + u_2^2 + u_3^2}$
- Unit Vector: $\hat{u} = \frac{\vec{u}}{|\vec{u}|}$

Equations of a Line

- Parametric Form: $\vec{r}(t) = \vec{r}_0 + t\vec{v}$, where $\vec{v}$ is the direction vector
- Symmetric Form: $\frac{x-x_0}{v_1} = \frac{y-y_0}{v_2} = \frac{z-z_0}{v_3}$

Equations of a Plane

- General Form: $ax + by + cz + d = 0$
- Normal Form: $\vec{n} \cdot (\vec{r} - \vec{r}_0) = 0$
- Parametric Form: $\vec{r}(s, t) = \vec{r}_0 + s\vec{v}_1 + t\vec{v}_2$

Curvature and Torsion

- Curvature: $\kappa = \frac{|\vec{T}'(s)|}{|\vec{r}'(t)|} = \frac{|\vec{r}'(t) \times \vec{r}''(t)|}{|\vec{r}'(t)|^3}$
- Torsion: $\tau = \frac{\vec{B} \cdot (\vec{T}' \times \vec{N})}{|\vec{r}'(t)|}$
- TNB Frame:
 - Tangent: $\vec{T} = \frac{\vec{r}'(t)}{|\vec{r}'(t)|}$
 - Normal: $\vec{N} = \frac{\vec{T}'(t)}{|\vec{T}'(t)|}$
 - Binormal: $\vec{B} = \vec{T} \times \vec{N}$

Equations of Motion in Physics

- Position: $\vec{r}(t) = \vec{r}_0 + \vec{v}_0t + \frac{1}{2}\vec{a}t^2$
- Velocity: $\vec{v}(t) = \vec{v}_0 + \vec{a}t$
- Acceleration: $\vec{a}(t) = \frac{d\vec{v}}{dt}$

Common Parameterizations

- Circle: $\vec{r}(t) = \langle R \cos t, R \sin t \rangle$
- Helix: $\vec{r}(t) = \langle R \cos t, R \sin t, ht \rangle$
- Ellipse: $\vec{r}(t) = \langle a \cos t, b \sin t \rangle$
- Parabola: $\vec{r}(t) = \langle t, at^2 \rangle$
- Sphere:

$$\vec{r}(\theta, \phi) = \langle R \sin \phi \cos \theta, R \sin \phi \sin \theta, R \cos \phi \rangle$$

Vector Calculus

- Gradient: $\nabla f = \langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \rangle$
- Divergence: $\nabla \cdot \vec{F} = \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} + \frac{\partial F_z}{\partial z}$
- Curl: $\nabla \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_x & F_y & F_z \end{vmatrix}$
- Laplacian: $\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}$

Vector Projections

- Projection of $\vec{u}$ onto $\vec{v}$: $\text{proj}_{\vec{v}} \vec{u} = \frac{\vec{u} \cdot \vec{v}}{\vec{v} \cdot \vec{v}} \vec{v}$
- Component of $\vec{u}$ along $\vec{v}$: $\text{comp}_{\vec{v}} \vec{u} = \frac{\vec{u} \cdot \vec{v}}{|\vec{v}|}$

Distance Formulas

- Point to Line: $d = \frac{|\vec{n} \cdot (\vec{r} - \vec{r}_0)|}{|\vec{n}|}$
- Point to Plane: $d = \frac{|ax_0 + by_0 + cz_0 + d|}{\sqrt{a^2 + b^2 + c^2}}$
- Line to Line (Skew): $d = \frac{|(\vec{r}_2 - \vec{r}_1) \cdot (\vec{v}_1 \times \vec{v}_2)|}{|\vec{v}_1 \times \vec{v}_2|}$

Appendix C: Calculus Theorems

Calculus I (Single Variable)

Limits and Continuity

- **Squeeze Theorem:** If $f(x) \leq g(x) \leq h(x)$ for all x near a and $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} h(x) = L$, then $\lim_{x \rightarrow a} g(x) = L$.
- **Intermediate Value Theorem (IVT):** If f is continuous on $[a, b]$ and N is between $f(a)$ and $f(b)$, then there exists $c \in [a, b]$ such that $f(c) = N$.

Differentiation

- **Mean Value Theorem (MVT):** If f is continuous on $[a, b]$ and differentiable on (a, b) , then there exists $c \in (a, b)$ such that $f'(c) = \frac{f(b)-f(a)}{b-a}$.
- **Rolle's Theorem:** If f is continuous on $[a, b]$, differentiable on (a, b) , and $f(a) = f(b)$, then there exists $c \in (a, b)$ such that $f'(c) = 0$.
- **L'Hôpital's Rule:** If $\lim_{x \rightarrow c} \frac{f(x)}{g(x)}$ is indeterminate, and f and g are differentiable, then $\lim_{x \rightarrow c} \frac{f(x)}{g(x)} = \lim_{x \rightarrow c} \frac{f'(x)}{g'(x)}$.

Integration

- **Fundamental Theorem of Calculus (FTC):**
 1. Part 1: If $F(x) = \int_a^x f(t) dt$, then $F'(x) = f(x)$.

2. Part 2: $\int_a^b f(x) dx = F(b) - F(a)$, where F is any antiderivative of f .

- **Integration by Substitution:** If $u = g(x)$, then $\int f(g(x))g'(x)dx = \int f(u)du$.

Calculus II (Techniques and Series)

Sequences and Series

- **Convergence of Series:** $\sum a_n$ converges if $\lim_{n \rightarrow \infty} S_n = S$ (finite).
- **Comparison Test:** If $0 \leq a_n \leq b_n$ for all n and $\sum b_n$ converges, then $\sum a_n$ converges.
- **Ratio Test:** $\sum a_n$ converges absolutely if $\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$.
- **Alternating Series Test:** An alternating series $\sum (-1)^n a_n$ converges if a_n is decreasing and $\lim_{n \rightarrow \infty} a_n = 0$.

Parametric and Polar Calculus

- **Arc Length (Parametric):** $L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$.
- **Arc Length (Polar):** $L = \int_\alpha^\beta \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$.
- **Area (Polar):** $A = \frac{1}{2} \int_\alpha^\beta r^2 d\theta$.

Calculus III (Multivariable Calculus)

Vector Calculus

- **Gradient:** $\nabla f = \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right\rangle$
- **Divergence:** $\nabla \cdot \vec{F} = \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} + \frac{\partial F_z}{\partial z}$
- **Curl:** $\nabla \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_x & F_y & F_z \end{vmatrix}$
- **Laplacian:** $\nabla^2 f = \frac{\partial^2 f}{\partial x^2} + \frac{\partial^2 f}{\partial y^2} + \frac{\partial^2 f}{\partial z^2}$

Theorems of Vector Calculus

- **Green's Theorem:** $\int_C \vec{F} \cdot d\vec{r} = \iint_R \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA$

- **Stokes' Theorem:** $\int_S (\nabla \times \vec{F}) \cdot \vec{n} dS = \int_C \vec{F} \cdot d\vec{r}$

- **Divergence Theorem:** $\iiint_V (\nabla \cdot \vec{F}) dV = \int_S \vec{F} \cdot \vec{n} dS$

Extrema and Optimization

- **Critical Points:** $\nabla f = \vec{0}$
- **Second Derivative Test:** Classify critical points using the Hessian matrix H .
- **Lagrange Multipliers:** $\nabla f = \lambda \nabla g$ to find extrema subject to $g(x, y, z) = 0$.

Appendix D: Quick Reference for First-Order ODEs

Basic Recipes

1. **Separable Equations:** $\frac{dy}{dx} = g(x)h(y)$
 - Rewrite as $\frac{dy}{h(y)} = g(x)dx$
 - Integrate both sides: $\int \frac{dy}{h(y)} = \int g(x)dx + C$
2. **Linear Equations:** $\frac{dy}{dx} + P(x)y = Q(x)$
 - Find integrating factor: $\mu(x) = e^{\int P(x)dx}$
 - Multiply both sides by $\mu(x)$
 - Rewrite as $\frac{d}{dx}[\mu(x)y] = \mu(x)Q(x)$
 - Integrate: $y = \frac{1}{\mu(x)}[\int \mu(x)Q(x)dx + C]$
3. **Exact Equations:** $M(x, y)dx + N(x, y)dy = 0$
 - Check if $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$
 - Find $\psi(x, y)$ such that $\frac{\partial \psi}{\partial x} = M$ and $\frac{\partial \psi}{\partial y} = N$
 - Solution is $\psi(x, y) = C$
4. **Bernoulli Equations:** $\frac{dy}{dx} + P(x)y = Q(x)y^n$
 - Substitute $v = y^{1-n}$
 - Solve resulting linear equation in v
 - Substitute back $y = v^{\frac{1}{1-n}}$

Appendix E: Useful Tables for Differential Equations

Table 2: *Comprehensive Table of Laplace Transform Pairs: These pairs enable both forward transformation for problem-solving and inverse transformation for solution recovery.*

$f(t)$	$\mathcal{L}\{f(t)\}$	$f(t)$	$\mathcal{L}\{f(t)\}$
1	$\frac{1}{s}$	$t \sin(at)$	$\frac{2a}{(s^2+a^2)^2}$
t	$\frac{1}{s^2}$	$t \cos(at)$	$\frac{s^2-a^2}{(s^2+a^2)^2}$
t^n	$\frac{n!}{s^{n+1}}$	$t^n e^{at}$	$\frac{n!}{(s-a)^{n+1}}$
e^{at}	$\frac{1}{s-a}$	$\frac{t^n}{n!}$	$\frac{1}{s^{n+1}}$
$\sin(at)$	$\frac{a}{s^2+a^2}$	$t^n \cos(at)$	$\frac{n!s}{(s^2+a^2)^{n+1}}$
$\cos(at)$	$\frac{s}{s^2+a^2}$	$t^n \sin(at)$	$\frac{n!a}{(s^2+a^2)^{n+1}}$
$\sinh(at)$	$\frac{a}{s^2-a^2}$	$\frac{\sin(at)}{t}$	$\tan^{-1}\left(\frac{a}{s}\right)$
$\cosh(at)$	$\frac{s}{s^2-a^2}$	$J_0(at)$	$\frac{1}{\sqrt{s^2+a^2}}$
$e^{at} \sin(bt)$	$\frac{b}{(s-a)^2+b^2}$	$u(t-c)$	$\frac{e^{-cs}}{s}$
$e^{at} \cos(bt)$	$\frac{s-a}{(s-a)^2+b^2}$	$\delta(t-c)$	e^{-cs}
$e^{at} f(t)$	$F(s-a)$	$u(t-a)$	$\frac{e^{-as}}{s}$
te^{at}	$\frac{1}{(s-a)^2}$	$\int_0^t f(\tau)g(t-\tau) d\tau$	$F(s)G(s)$
e^{-at^2}	$\frac{\sqrt{\pi}}{2\sqrt{a}} e^{\frac{s^2}{4a}} \operatorname{erfc}\left(\frac{s}{2\sqrt{a}}\right)$	$t^n u(t)$	$\frac{n!}{s^{n+1}}$

Table 3: *Comprehensive Table of Common PDE Operators and Their Green's Functions. In these expressions, the notation $G(x, t; \xi, 0)$ (or $G(\mathbf{x}, \boldsymbol{\xi})$) denotes the Green's function where the first set of variables indicates the field point (where the solution is observed) and the second set indicates the source point (where the impulse is applied).*

PDE/Operator	Green's Function
Laplacian in $\mathbb{R}^3$	$G(\mathbf{x}, \boldsymbol{\xi}) = \frac{1}{4\pi \mathbf{x} - \boldsymbol{\xi} }$
Laplacian in $\mathbb{R}^2$	$G(\mathbf{x}, \boldsymbol{\xi}) = -\frac{1}{2\pi} \ln \mathbf{x} - \boldsymbol{\xi} $
Heat Equation in $\mathbb{R}$	$G(x, t; \xi, 0) = \frac{1}{\sqrt{4\pi t}} \exp\left(-\frac{(x - \xi)^2}{4t}\right), \quad t > 0$
Wave Equation in $\mathbb{R}^3$	$G(t, \mathbf{x}; t', \boldsymbol{\xi}) = \frac{\delta\left(t - t' - \frac{ \mathbf{x} - \boldsymbol{\xi} }{c}\right)}{4\pi \mathbf{x} - \boldsymbol{\xi} }$
Free Schrödinger Equation in 1D	$G(x, t; \xi, 0) = \sqrt{\frac{m}{2\pi i \hbar t}} \exp\left(\frac{im(x - \xi)^2}{2\hbar t}\right)$
Helmholtz Equation in $\mathbb{R}^3$	$G(\mathbf{x}, \boldsymbol{\xi}) = \frac{e^{ik \mathbf{x} - \boldsymbol{\xi} }}{4\pi \mathbf{x} - \boldsymbol{\xi} }$

Table 4: *Comprehensive Table of Fourier Transform Pairs: These pairs enable both forward and inverse transformations for analysis and signal processing.*

$f(t)$	$F(\omega)$	$f(t)$	$F(\omega)$
$\delta(t)$	1	1	$2\pi \delta(\omega)$
e^{-at^2}	$\sqrt{\frac{\pi}{a}} e^{-\frac{\omega^2}{4a}}$	$e^{i\omega_0 t}$	$2\pi \delta(\omega - \omega_0)$
$\cos(\omega_0 t)$	$\pi [\delta(\omega - \omega_0) + \delta(\omega + \omega_0)]$	$\sin(\omega_0 t)$	$i\pi [\delta(\omega + \omega_0) - \delta(\omega - \omega_0)]$
$e^{-at}u(t)$	$\frac{1}{a+i\omega}$	$t e^{-at}u(t)$	$\frac{1}{(a+i\omega)^2}$
$u(t)$	$\pi \delta(\omega) + \frac{1}{i\omega}$	$e^{-a t }$	$\frac{2a}{a^2 + \omega^2}$

Appendix F: Complex Analysis Identities

Basic Identities

Euler's Formula

- $e^{ix} = \cos x + i \sin x$
- $e^{i\pi} + 1 = 0$
- $\cos x = \frac{e^{ix} + e^{-ix}}{2}$
- $\sin x = \frac{e^{ix} - e^{-ix}}{2i}$

Complex Conjugate

- $\overline{z_1 + z_2} = \overline{z_1} + \overline{z_2}$
- $\overline{z_1 \cdot z_2} = \overline{z_1} \cdot \overline{z_2}$
- $\overline{z_1/z_2} = \overline{z_1}/\overline{z_2}$
- $\overline{\overline{z}} = z$
- $z \cdot \overline{z} = |z|^2$
- $\overline{z^n} = \overline{z}^n$
- $\overline{e^z} = e^{\overline{z}}$

Modulus Properties

- $|z_1 \cdot z_2| = |z_1| \cdot |z_2|$
- $|z_1/z_2| = |z_1|/|z_2|$
- $|z_1 + z_2| \leq |z_1| + |z_2|$ (Triangle Inequality)

- $|z_1 - z_2| \geq ||z_1| - |z_2||$
- $|z^n| = |z|^n$
- $|e^z| = e^{\operatorname{Re}(z)}$
- $|i^n| = 1$ for all integers n

Trigonometric Connections

- $\cos(iz) = \cosh(z)$
- $\sin(iz) = i \sinh(z)$
- $\cosh(iz) = \cos(z)$
- $\sinh(iz) = i \sin(z)$
- $\tan(iz) = i \tanh(z)$
- $\tanh(iz) = i \tan(z)$

Hyperbolic Trigonometric Identities

- $\sinh(z) = \frac{e^z - e^{-z}}{2}$
- $\cosh(z) = \frac{e^z + e^{-z}}{2}$
- $\tanh(z) = \frac{\sinh(z)}{\cosh(z)} = \frac{e^z - e^{-z}}{e^z + e^{-z}}$
- $\coth(z) = \frac{\cosh(z)}{\sinh(z)} = \frac{e^z + e^{-z}}{e^z - e^{-z}}$
- $\operatorname{sech}(z) = \frac{1}{\cosh(z)} = \frac{2}{e^z + e^{-z}}$
- $\operatorname{csch}(z) = \frac{1}{\sinh(z)} = \frac{2}{e^z - e^{-z}}$

- $\cosh^2(z) - \sinh^2(z) = 1$
- $1 - \tanh^2(z) = \operatorname{sech}^2(z)$
- $\coth^2(z) - 1 = \operatorname{csch}^2(z)$
- $\sinh(z_1 \pm z_2) = \sinh(z_1) \cosh(z_2) \pm \cosh(z_1) \sinh(z_2)$
- $\cosh(z_1 \pm z_2) = \cosh(z_1) \cosh(z_2) \pm \sinh(z_1) \sinh(z_2)$
- $\sinh(2z) = 2 \sinh(z) \cosh(z)$
- $\cosh(2z) = \cosh^2(z) + \sinh^2(z) = 2 \cosh^2(z) - 1 = 1 + 2 \sinh^2(z)$
- $\tanh(z_1 \pm z_2) = \frac{\tanh(z_1) \pm \tanh(z_2)}{1 \pm \tanh(z_1) \tanh(z_2)}$

Complex Functions

Logarithmic Identities

- $\log(z) = \ln |z| + i \arg(z)$
- $\log(z_1 z_2) = \log(z_1) + \log(z_2) + 2\pi i k$ for some integer k
- $\log(z^n) = n \log(z) + 2\pi i k$ for some integer k
- $\log(1/z) = -\log(z) + 2\pi i k$ for some integer k
- $\log(e^z) = z + 2\pi i k$ for some integer k
- $\log(-z) = \log(z) + i\pi$ up to multiples of $2\pi i$

Power Identities

- $z^{1/n} = |z|^{1/n} e^{i(\arg(z) + 2\pi k)/n}$ for $k = 0, 1, \dots, n-1$
- $z^w = e^{w \log(z)}$
- $(e^z)^w = e^{zw}$
- $(-1)^n = e^{in\pi} = \cos(n\pi) + i \sin(n\pi)$
- $(\cos \theta + i \sin \theta)^n = \cos(n\theta) + i \sin(n\theta)$

Analytic Functions

Cauchy-Riemann Equations

If $f(z) = u(x, y) + iv(x, y)$ is analytic, then:

- $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$
- $\frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$
- $\nabla^2 u = \nabla^2 v = 0$ (Laplace's equation)

Conformal Mappings

Transformation Properties

- $w = z + a$ (Translation)
- $w = az$ (Rotation and scaling)
- $w = 1/z$ (Inversion)
- $w = (az + b)/(cz + d)$ (Möbius transformation)
- $w = z^n$ (Power mapping)
- $w = e^z$ (Exponential mapping)
- $w = \log(z)$ (Logarithmic mapping)

References

- OpenStax. *Calculus Volume 1*. OpenStax, 2016. Accessed August 16, 2024. <https://openstax.org/details/books/calculus-volume-1>
- OpenStax. *Calculus Volume 2*. OpenStax, 2016. Accessed August 16, 2024. <https://openstax.org/details/books/calculus-volume-2>
- OpenStax. *Calculus Volume 3*. OpenStax, 2016. Accessed August 16, 2024. <https://openstax.org/details/books/calculus-volume-3>
- Strang, Gilbert. *Introduction to Linear Algebra*. Fifth Edition. Wellesley-Cambridge Press, 2016. OpenCourseWare: <https://ocw.mit.edu/courses/mathematics/18-06-linear-algebra-spring-2010/>
- University of British Columbia. “CLP Calculus Textbooks.” Creative Commons. Accessed August 16, 2024. <https://www.math.ubc.ca/~CLP/>
- Feynman, Richard P. *The Feynman Lectures on Physics*. Open Access Edition. Accessed August 16, 2024. <http://www.feynmanlectures.caltech.edu/>
- MIT OpenCourseWare. “Single Variable Calculus.” Massachusetts Institute of Technology. Accessed August 16, 2024. <https://ocw.mit.edu/courses/mathematics/18-01sc-single-variable-calculus-fall-2010/>
- MIT OpenCourseWare. “Multivariable Calculus.” Massachusetts Institute of Technology. Accessed August 16, 2024. <https://ocw.mit.edu/courses/mathematics/18-02sc-multivariable-calculus-fall-2010/>
- Wikipedia Contributors. “Gradient.” Wikipedia, The Free Encyclopedia. Accessed January 10, 2025. <https://en.wikipedia.org/wiki/Gradient>
- Wikipedia Contributors. “Divergence.” Wikipedia, The Free Encyclopedia. Accessed January 10, 2025. <https://en.wikipedia.org/wiki/Divergence>
- Wikipedia Contributors. “Curl (mathematics).” Wikipedia, The Free Encyclopedia. Accessed January 10, 2025. [https://en.wikipedia.org/wiki/Curl_\(mathematics\)](https://en.wikipedia.org/wiki/Curl_(mathematics))
- Wikipedia Contributors. “Line Integral.” Wikipedia, The Free Encyclopedia. Accessed January 10, 2025. https://en.wikipedia.org/wiki/Line_integral

- Wikipedia Contributors. “Surface Integral.” Wikipedia, The Free Encyclopedia. Accessed January 10, 2025. https://en.wikipedia.org/wiki/Surface_integral
- Wikipedia Contributors. “Green’s Theorem.” Wikipedia, The Free Encyclopedia. Accessed January 10, 2025. https://en.wikipedia.org/wiki/Green's_theorem
- Wikipedia Contributors. “Stokes’ Theorem.” Wikipedia, The Free Encyclopedia. Accessed January 10, 2025. https://en.wikipedia.org/wiki/Stokes'_theorem
- Wikipedia Contributors. “Divergence Theorem.” Wikipedia, The Free Encyclopedia. Accessed January 10, 2025. https://en.wikipedia.org/wiki/Divergence_theorem
- Wikipedia Contributors. “Laplacian.” Wikipedia, The Free Encyclopedia. Accessed January 10, 2025. <https://en.wikipedia.org/wiki/Laplacian>
- Wikipedia Contributors. “Parametric Surface.” Wikipedia, The Free Encyclopedia. Accessed January 10, 2025. https://en.wikipedia.org/wiki/Parametric_surface
- Wikipedia Contributors. “Tangent, Normal, and Binormal Vectors.” Wikipedia, The Free Encyclopedia. Accessed January 10, 2025. https://en.wikipedia.org/wiki/FrenetSerret_formulas

Index

- Ampère's Law, [462](#)
- Analytic Functions, [631](#)
- Analyticity, [693](#)
- Angular Momentum, [316](#), [350](#)
- Antiderivatives, [647](#)
- Approximations in Riemann Sums, [12](#)
- Arc Length, [133](#)
- Argand Plane, [579](#)
- Average Distribution, [318](#), [355](#)
- Average Value, [318](#), [355](#)

- Basic Integral Formulas, [15](#)
- Basic Terminology, [v](#)
- Bernoulli Differential Equations, [512](#)
- Boundary Conditions, [496](#)
- Boundary Value Problem, [497](#)
- Branch Cuts, [606](#)

- Calculus, [2](#)
- Cauchy's Integral Formula, [657](#)
- Cauchy's Integral Theorem, [651](#)
- Cauchy-Riemann Equations, [627](#)
- Center of Mass, [310](#), [346](#)
- Chain Rule, [10](#)
- Change in Variables, [336](#)
- Characteristic Equation Method, [518](#)
- Clairaut's Theorem, [190](#), [423](#)
- Closed Sets, [229](#)
- Compact Sets, [229](#)
- Complex Conjugate, [581](#)
- Complex Differentiation, [623](#)
- Complex Domain, [595](#)
- Complex Exponential, [598](#)
- Complex Function Definition, [594](#)
- Complex Functions, [594](#), [693](#)
- Complex Integration, [637](#)
- Complex Line Integral, [638](#)
- Complex Logarithm, [603](#)
- Complex Numbers, [578](#), [692](#)
- Complex Trigonometric Functions, [601](#)
- Complex Visualization Software, [606](#)
- Conformal Mapping, [636](#)
- Conformal Mappings, [693](#)
- Conservation Laws, [474](#)
- Conservative Fields, [382](#)
- Constrained Optimization, [253](#), [261](#)
- Continuous Functions, [180](#)
- Coordinate Systems, [25](#)
- Critical Points, [226](#)
- Cross Product, [49](#)
- Curl, [417](#)
- Curvature, [120](#), [122](#)
- Cylindrical Coordinates, [31](#)

- De Moivre's Theorem, [586](#)
- Definite Integrals, [11](#)
- Derivative Rules, [9](#)
- Derivatives, [7](#), [683](#)
- Differential Equations, [491](#)
 - Classifications, [495](#)
 - Order, [495](#)
- Differential Form, [465](#)
- Direction Cosines, [40](#), [92](#)
- Directional Derivatives, [203](#)
- Discriminant, [237](#)
- Divergence, [417](#)
- Divergence Theorem, [468](#)
- Gauss's Theorem, [468](#)
- Dot Product, [48](#)
- Double Integral Regions, [291](#)

Type I Regions, [292](#)
 Type II Regions, [295](#)
 Double Integrals, [276](#)

 Eigenvalues, [244](#)
 Elementary Complex Functions, [596](#)
 Epsilon-Delta Definition of Limits, [5](#)
 Euler's Formula, [584](#)
 Exact Differential Equations, [508](#)
 Existence and Uniqueness Theorems, [513](#)
 Existence Theorem, [513](#)
 Extrema in Several Variables, [228](#)
 Extreme Value Theorem, [22](#), [182](#), [232](#)

 Faraday's Law, [462](#)
 First Derivative Test, [233](#)
 First-Order Approximation, [224](#)
 First-Order ODEs, [502](#)
 Flux, [406](#), [447](#)
 Flux Integral, [425](#)
 Fubini's Theorem, [281](#)
 Functions of Three or More Variables, [165](#)
 Functions of Two Variables, [152](#)
 Fundamental Theorem for Line Integrals, [649](#)
 Fundamental Theorem of Algebra, [592](#)
 Fundamental Theorem of Calculus, [22](#)
 Fundamental Theorem of Line Integrals, [400](#)

 General Regions, [290](#)
 Geometric Optimization, [251](#)
 Global Extrema, [230](#), [239](#)
 Global Maxima, [230](#)
 Global Minima, [230](#)
 Gradient Ascent, [209](#)
 Gradient Descent, [209](#)
 Gradient Vector, [204](#)
 Green's Functions
 Advanced Methods, [566](#)
 Green's Theorem, [406](#)

 Harmonic Functions, [631](#)
 Heine-Borel Theorem, [618](#)
 Hessian Matrix, [237](#), [243](#)
 High-Dimensional Arc Length, [138](#)

 Higher-Order ODEs, [515](#), [528](#)
 Fundamentals, [515](#)
 Second Order, [517](#)
 Superposition, [516](#)
 Higher-Order Partial Derivatives, [188](#)
 Holomorphic Functions, [623](#)
 Homogeneous Differential Equations, [495](#)
 Hyperplanes, [220](#), [221](#)

 Implicit Differentiation, [197](#)
 Indefinite Double Integration, [287](#)
 Indefinite Integrals, [11](#)
 Indefiniteness, [244](#)
 Infinite Limits, [5](#)
 Initial Conditions, [496](#)
 Initial Value Problem, [497](#)
 Integral Approximations, [12](#)
 Integrals, [11](#), [684](#)
 Integrating Factor, [505](#)
 Integration by Parts, [16](#)
 Integration Rules, [17](#)
 Intermediate Value Theorem, [22](#), [181](#)
 Iterated Integrals, [282](#)

 Jacobian, [336](#)

 Karush-Kuhn-Tucker Conditions, [263](#)

 L'Hôpital's Rule, [6](#), [22](#)
 Lagrange Condition, [254](#)
 Lagrange Function, [254](#)
 Lagrange Multiplier, [253](#)
 Laplace Transform, [528](#)
 Laplacian, [429](#)
 Laurent Series, [660](#)
 Definition, [661](#)
 Leading Principal Minors, [245](#)
 Leibniz Integral Rule, [202](#)
 Leibniz Rule, [22](#)
 Level Curves, [159](#), [210](#)
 Level Surfaces, [166](#)
 Limit Rules, [4](#)
 Limits, [4](#), [611](#)
 Limits and Continuity, [169](#)
 Line Integrals, [387](#)
 Linear Approximation, [222](#)

Linear Classifier, [221](#)
 Linear Differential Equations, [495](#)
 Linear First-Order ODEs, [504](#)
 Linear Independence, [518](#)
 Lines, [59](#)
 Liouville's Theorem, [635](#)
 Local Extrema, [228](#), [238](#)
 Local Maxima, [228](#)
 Local Minima, [228](#)

 Machine Learning, [167](#)
 Maclaurin Series, [19](#)
 Mass and Density, [310](#), [346](#)
 Mathematical Symbols, [ix](#)
 Matrices, [336](#)
 Matrix Definiteness, [244](#)
 Matrix Operations, [351](#)
 Maxwell's Equations, [462](#)
 Mean Value Theorem, [21](#), [182](#)
 Method of Undetermined Coefficients, [519](#)
 Metric Formulas, [27](#)
 ML Inequality, [644](#)
 Modulus, [581](#)
 Moment of Inertia, [314](#), [349](#)
 Morera's Theorem, [655](#)
 Motion, [110](#)
 Multiply Connected Domains, [647](#)
 Multivariable Limits, [170](#)

 Negative Definiteness, [244](#)
 Non-Conservative Fields, [383](#)
 Non-homogeneous Differential Equations, [496](#)
 Nonlinear Differential Equations, [495](#)
 Numerical Arc Length, [142](#)
 Numerical Integration, [13](#)

 Open Sets, [229](#)
 Optimization with Inequality Constraints, [263](#)
 Ordinary Differential Equations, [495](#)
 Osculating Circles, [124](#)

 Parametric Equations, [99](#)
 Parametric Line, [92](#)
 Parametric Surfaces, [434](#)

 Partial Derivatives, [184](#)
 Partial Differential Equations, [495](#)
 Classification, [539](#)
 Fourier Series, [549](#)
 Green's Functions, [560](#)
 Method of Characteristics, [553](#)
 Methods, [539](#), [553](#)
 Separation of Variables, [541](#)
 Partial Fractions, [18](#)
 Path Independence, [647](#)
 Path Integrals, [637](#)
 Planes, [59](#)
 Poisson's Equation, [551](#)
 Polar Coordinate Arc Length, [135](#)
 Polar Coordinates, [30](#), [297](#)
 Polar Regions, [297](#)
 Positive Definiteness, [244](#)
 Power Series, [633](#)
 Primitive Root, [588](#)
 Product of Inertia, [349](#)
 Product Rule, [10](#)
 Projectile Motion, [112](#)
 Properties of Complex Integrals, [643](#)
 Properties of Complex Numbers, [579](#)
 Python, [57](#), [125](#), [214](#), [377](#), [538](#), [607](#)

 Quadric Surfaces, [80](#)
 Quotient Rule, [10](#)

 Rate of Change, [208](#)
 Rectangular Coordinates, [25](#)
 Rectangular Regions, [276](#)
 Reduction Formulas, [18](#)
 Reparameterization, [133](#)
 Residue Theorem, [664](#)
 Residue Theory, [651](#), [664](#)
 Real Integrals, [668](#)
 Residues
 Computation, [665](#)
 Definition, [664](#)
 Riemann Sum, [11](#), [277](#)
 Riemann Sum Types, [280](#)
 Rolle's Theorem, [21](#)
 Roots of Unity, [586](#)
 Rotational Motion, [314](#), [350](#)

Saddle Points, [242](#)
Scalar Functions, [99](#)
Second Derivative Test, [237](#)
Second-Order Approximation, [225](#)
Semidefiniteness, [244](#)
Separation of Variables, [502](#)
Simpson's Rule, [14](#)
Single-Sided Limits, [4](#)
Singular Points, [660](#)
Singularities
 Classification, [662](#)
Spherical Coordinates, [33](#)
Squeeze Theorem, [182](#)
Stokes' Theorem, [450](#)
Substitution Method, [17](#)
Support Vector Machines, [264](#)
Supremum and Infimum, [13](#)
Surface Area, [319](#)
Surface Integrals, [434](#)
Sylvester's Criterion, [246](#)

Tabular Method, [16](#)
Taco-Related Examples, [153](#)
Tangent Planes, [217](#)
Taylor Series, [19](#)
Taylor's Theorem, [23](#)
Tensors, [350](#)
The Chain Rule, [194](#)
TNB Frame, [120](#)

Toroidal Coordinates, [339](#)
Torque, [316](#), [351](#)
Torsion, [120](#), [128](#)
Total Differentials, [200](#)
Transformations, [336](#)
Trapezoidal Rule, [14](#)
Trigonometric Integrals, [18](#)
Trigonometric Substitution, [18](#)
Triple Integral Regions, [323](#)
 Type I Regions, [323](#)
 Type II Regions, [324](#)
 Type III Regions, [324](#)

Unconstrained Optimization, [248](#)
Uniqueness Theorem, [513](#)

Variation of Parameters, [524](#)
Vector Angle, [92](#)
Vector Calculus, [103](#)
Vector Fields, [371](#)
Vector Operations, [44](#)
Vector Operators, [428](#)
Vector Projection, [72](#)
Vector-Valued Functions, [99](#)
Vectors, [38](#)
Verification of Solutions, [499](#)
Vibrating Membrane, [547](#)

Winding Number, [647](#)
Wronskian, [518](#)