

## Lecture 8: Partial Derivatives

### Background & Motivation

Now we differentiate. The idea is simple: hold all variables constant except one, and differentiate with respect to that one. This gives us partial derivatives—the multivariate analog of  $f'(x)$ . Today we define them, compute them, and see how they measure the slope of a surface in each coordinate direction.

### 1. Partial Derivatives

#### Definition 1: Partial Derivative (Limit Definition)

The **partial derivative**  $f_x(a, b)$  is the instantaneous rate of change of  $f(x, y)$  in the  $x$ -direction at the point  $(a, b)$ , with  $y$  held fixed at  $b$ . Geometrically, slice the graph of  $f$  with the vertical plane  $y = b$ ; the resulting cross-section is a single-variable curve  $z = f(x, b)$ , and  $f_x(a, b)$  is the **slope of the tangent line** to that curve at  $x = a$ . In other words, partial derivatives are ordinary single-variable slopes taken along axis-aligned slices of the surface.

$$f_x(x, y) = \frac{\partial f}{\partial x} = \underline{\hspace{2cm}}$$


With respect to  $y$ :

$$f_y(x, y) = \frac{\partial f}{\partial y} = \underline{\hspace{2cm}}$$

**Practical rule:**  $\frac{\partial}{\partial x}$  means differentiate with respect to  $x$ , treating  $y$  as a constant.

#### Geometric Interpretation

- $f_x(a, b)$  is the slope of the tangent line to  $z = f(x, b)$  at  $x = a$
- Measures rate of change of  $f$  in the  $x$ -direction while  $y$  is held fixed
- Similarly,  $f_y(a, b)$  is the slope of  $z = f(a, y)$  at  $y = b$



Tangent line (red) at  $(1, 1, 2)$  on  $z = x^2 + y^2$ . Blue dashed:  $xz$ -projection (slope  $= f_x = 2$ ). Green dashed:  $yz$ -projection (slope  $= f_y = 2$ ).

### Example 1: Using the Limit Definition

Use the limit definition to find  $f_x$  for  $f(x, y) = x^2y + 3xy^2$ .

### Example 2: Computing Partial Derivatives

Find all first partial derivatives:

(a)  $f(x, y) = x^3e^y + \sin(xy)$

(b)  $g(x, y) = \frac{x^2}{x + y}$

(c)  $h(x, y, z) = xye^z + z \ln(x + y)$

## 2. Derivative Rules for Partialials

### Rules (Hold Other Variables Constant)

- **Linearity:**  $\partial_x[af + bg] = af_x + bg_x$
- **Product:**  $\partial_x[fg] = \underline{\hspace{2cm}}$
- **Quotient:**  $\partial_x[f/g] = \underline{\hspace{2cm}}$
- **Chain:**  $\partial_x[f(g(x, y))]$  =  $\underline{\hspace{2cm}}$
- **Zero partial:** if  $f_x = 0$  everywhere  $\implies f$  does not depend on  $x$
- **Constants:**  $\partial_x[c] = 0$ ; treat other variables as constants

## 3. Higher-Order Partial Derivatives

### Definition 2: Second-Order Partial Derivatives

A **second-order partial derivative** is a partial derivative of a partial derivative. The two *pure* second partials  $f_{xx}$  and  $f_{yy}$  measure **concavity along coordinate directions** — how fast the slope  $f_x$  changes as you move further in  $x$ , and similarly for  $y$  — exactly analogous to the second derivative of a single-variable function. The two *mixed* partials  $f_{xy}$  and  $f_{yx}$  measure something new: how the slope in *one* direction changes as you move in the *other* direction, capturing the graph's twist or saddle-like behavior at a point.

$$f_{xx} = \frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right) \quad (\text{pure, in } x)$$

$$f_{yy} = \frac{\partial^2 f}{\partial y^2} \quad (\text{pure, in } y)$$

$$f_{xy} = \frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial y} \left( \frac{\partial f}{\partial x} \right) \quad (\text{mixed: } x \text{ first, then } y)$$

$$f_{yx} = \frac{\partial^2 f}{\partial x \partial y} \quad (\text{mixed: } y \text{ first, then } x)$$

### Theorem 1: Clairaut's Theorem (Equality of Mixed Partialials)

If  $f_{xy}$  and  $f_{yx}$  are both continuous on an open region, then:

\_\_\_\_\_

The order of differentiation does not matter (under mild continuity conditions).

### Why Clairaut's Theorem Works (Proof Outline)

- Consider the second-order difference quotient over a small rectangle  $[x, x + h] \times [y, y + k]$
- Define  $\Delta = f(x + h, y + k) - f(x + h, y) - f(x, y + k) + f(x, y)$
- Apply the Mean Value Theorem twice: first in  $x$ , then in  $y \implies \Delta = f_{xy}(c_1, d_1) \cdot hk$

- Apply MVT in the other order: first in  $y$ , then in  $x \implies \Delta = f_{yx}(c_2, d_2) \cdot hk$
- As  $h, k \rightarrow 0$ , continuity gives  $f_{xy}(x, y) = f_{yx}(x, y)$

### When Clairaut Fails – A Counterexample

- Consider  $f(x, y) = \begin{cases} \frac{xy(x^2 - y^2)}{x^2 + y^2} & (x, y) \neq (0, 0) \\ 0 & (x, y) = (0, 0) \end{cases}$
- One can show  $f_{xy}(0, 0) = -1$  and  $f_{yx}(0, 0) = 1$  – the mixed partials are *not equal* at the origin because they are not continuous there

#### Example 3: Clairaut Verification

Let  $f(x, y) = x^3y^2 + e^{xy}$ . Find all four second-order partials and verify  $f_{xy} = f_{yx}$ .

#### Example 4: Application: Container Design

A cylindrical container has volume  $V(r, h) = \pi r^2 h$  and surface area  $S(r, h) = 2\pi r h + 2\pi r^2$ . If  $r = 5$  cm and  $h = 10$  cm, how does  $V$  change if we increase  $r$  by 0.1 cm? If we increase  $h$  by 0.1 cm?

## Practice Problems

---

**Problem 1.** Find  $f_x$  and  $f_y$  for  $f(x, y) = x^2 \cos y + y \ln x$ .

---

**Problem 2.** Find all first partials of  $f(x, y, z) = xyz + z^2 \sin(x)$ .

---

**Problem 3.** For  $f(x, y) = \arctan(y/x)$ , show that  $f_{xx} + f_{yy} = 0$  (i.e.,  $f$  is **harmonic**).

---

**Problem 4.** Use the limit definition to compute  $f_x(0, 0)$  for  $f(x, y) = \begin{cases} \frac{xy}{x^2+y^2} & (x, y) \neq (0, 0) \\ 0 & (x, y) = (0, 0) \end{cases}$

---

**Problem 5.** Find  $f_{xxy}$  for  $f(x, y) = e^x \sin(xy)$ .