

Lecture 10: Partial Derivatives

Background & Motivation

Now we differentiate. The idea is simple: hold all variables constant except one, and differentiate with respect to that one. This gives us partial derivatives—the multivariate analog of $f'(x)$. Today we define them, compute them, and see how they measure the slope of a surface in each coordinate direction.

1. Partial Derivatives

Definition 1: Partial Derivative (Limit Definition)

The **partial derivative** $f_x(a, b)$ is the instantaneous rate of change of $f(x, y)$ in the x -direction at the point (a, b) , with y held fixed at b . Geometrically, slice the graph of f with the vertical plane $y = b$; the resulting cross-section is a single-variable curve $z = f(x, b)$, and $f_x(a, b)$ is the **slope of the tangent line** to that curve at $x = a$. In other words, partial derivatives are ordinary single-variable slopes taken along axis-aligned slices of the surface.

$$f_x(x, y) = \frac{\partial f}{\partial x} = \underline{\hspace{2cm}}$$


With respect to y :

$$f_y(x, y) = \frac{\partial f}{\partial y} = \underline{\hspace{2cm}}$$

Practical rule: $\underline{\hspace{2cm}}$

Geometric Interpretation

- $f_x(a, b)$ is the slope of the tangent line to $z = f(x, b)$ at $x = a$
- Measures rate of change of f in the x -direction while y is held fixed
- Similarly, $f_y(a, b)$ is the slope of $z = f(a, y)$ at $y = b$



Tangent line (red) at $(1, 1, 2)$ on $z = x^2 + y^2$. Blue dashed: xz -projection (slope $= f_x = 2$). Green dashed: yz -projection (slope $= f_y = 2$).

Example 1: Using the Limit Definition

Use the limit definition to find f_x for $f(x, y) = x^2y + 3xy^2$.

Example 2: Computing Partial Derivatives

Find all first partial derivatives:

(a) $f(x, y) = x^3e^y + \sin(xy)$

(b) $g(x, y) = \frac{x^2}{x + y}$

(c) $h(x, y, z) = xye^z + z \ln(x + y)$

2. Derivative Rules for Partialials

Rules (Hold Other Variables Constant)

- **Linearity:** $\partial_x[af + bg] = af_x + bg_x$
- **Product:** $\partial_x[fg] = \underline{\hspace{2cm}}$
- **Quotient:** $\partial_x[f/g] = \underline{\hspace{2cm}}$
- **Chain:** $\partial_x[f(g(x, y))]$ = $\underline{\hspace{2cm}}$
- **Zero partial:** if $f_x = 0$ everywhere $\implies f$ does not depend on x
- **Constants:** $\partial_x[c] = 0$; treat other variables as constants

3. Higher-Order Partial Derivatives

Definition 2: Second-Order Partial Derivatives

A **second-order partial derivative** is a partial derivative of a partial derivative. The two *pure* second partials f_{xx} and f_{yy} measure **concavity along coordinate directions** — how fast the slope f_x changes as you move further in x , and similarly for y — exactly analogous to the second derivative of a single-variable function. The two *mixed* partials f_{xy} and f_{yx} measure something new: how the slope in *one* direction changes as you move in the *other* direction, capturing the graph's twist or saddle-like behavior at a point.

$$f_{xx} = \frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right) \quad (\text{pure, in } x)$$

$$f_{yy} = \frac{\partial^2 f}{\partial y^2} \quad (\text{pure, in } y)$$

$$f_{xy} = \frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right) \quad (\text{mixed: } x \text{ first, then } y)$$

$$f_{yx} = \frac{\partial^2 f}{\partial x \partial y} \quad (\text{mixed: } y \text{ first, then } x)$$

Theorem 1: Clairaut's Theorem (Equality of Mixed Partialials)

If f_{xy} and f_{yx} are both continuous on an open region, then:

The order of differentiation does not matter (under mild continuity conditions).

Why Clairaut's Theorem Works (Proof Outline)

- Consider the second-order difference quotient over a small rectangle $[x, x + h] \times [y, y + k]$
- Define $\Delta = f(x + h, y + k) - f(x + h, y) - f(x, y + k) + f(x, y)$
- Apply the Mean Value Theorem twice: first in x , then in $y \implies \Delta = f_{xy}(c_1, d_1) \cdot hk$

- Apply MVT in the other order: first in y , then in $x \implies \Delta = f_{yx}(c_2, d_2) \cdot hk$
- As $h, k \rightarrow 0$, continuity gives $f_{xy}(x, y) = f_{yx}(x, y)$

When Clairaut Fails – A Counterexample

- Consider $f(x, y) = \begin{cases} \frac{xy(x^2 - y^2)}{x^2 + y^2} & (x, y) \neq (0, 0) \\ 0 & (x, y) = (0, 0) \end{cases}$
- One can show $f_{xy}(0, 0) = -1$ and $f_{yx}(0, 0) = 1$ – the mixed partials are *not equal* at the origin because they are not continuous there

Example 3: Clairaut Verification

Let $f(x, y) = x^3y^2 + e^{xy}$. Find all four second-order partials and verify $f_{xy} = f_{yx}$.

Example 4: Application: Container Design

A cylindrical container has volume $V(r, h) = \pi r^2 h$ and surface area $S(r, h) = 2\pi r h + 2\pi r^2$. If $r = 5$ cm and $h = 10$ cm, how does V change if we increase r by 0.1 cm? If we increase h by 0.1 cm?

Practice Problems

Problem 1. Find f_x and f_y for $f(x, y) = x^2 \cos y + y \ln x$.

Problem 2. Find all first partials of $f(x, y, z) = xyz + z^2 \sin(x)$.

Problem 3. For $f(x, y) = \arctan(y/x)$, show that $f_{xx} + f_{yy} = 0$ (i.e., f is **harmonic**).

Problem 4. Use the limit definition to compute $f_x(0, 0)$ for $f(x, y) = \begin{cases} \frac{xy}{x^2+y^2} & (x, y) \neq (0, 0) \\ 0 & (x, y) = (0, 0) \end{cases}$

Problem 5. Find f_{xxy} for $f(x, y) = e^x \sin(xy)$.