

A Tutorial Overview of Ordinal Notations

Jacques Bailhache (jacques.bailhache@gmail.com)

January-February 2018

1 Interest of transfinite ordinal numbers

The domain of transfinite ordinals has the particularity of being the only mathematical domain that cannot be automated. In all other domains of mathematics, it is at least theoretically possible to deduce the theorems automatically from a formal system consisting of a finite set of axioms and rules. But Gödel proved that given any formal system of a theory sufficiently powerful to contain arithmetics, it is possible to build a proposition that expresses its own unprovability in this formal system. This proposition, which is very huge, has also a meaning as an ordinary arithmetic proposition, but is very useless in ordinary arithmetics. If the formal system is consistent, then this proposition is undecidable.

At first sight one could think that we just have to add this proposition to the system as an axiom, but this augmented system also have its own Gödelian proposition. By adding successively Gödelian propositions, we obtain an infinite sequence of systems, and the system defined as the union of all these systems also has its Gödelian proposition, and so on. But according to Solomon Feferman in "Penrose's Gödelian argument" <http://math.stanford.edu/~feferman/papers/penrose.pdf> p.9 :

"one obtains completeness for all arithmetical sentences in a progression based on the transfinite iteration of the so-called global or uniform reflection principle"

The uniform reflection principle, which is something similar to adding the Gödelian proposition as an axiom, is described for example in John Harrison's paper "Metatheory and Reflection in Theorem Proving: A Survey and Critique"

<http://www.cl.cam.ac.uk/~jrh13/papers/reflect.ps.gz> p.18 :

$$\vdash \forall n. Pr(\lceil \phi[n] \rceil) \Rightarrow \phi[n]$$

Harrison also says p.19 :

"Feferman showed that a transfinite iteration based on it proves all true sentences of number theory".

So we can say that the construction of transfinite ordinals can store all the creative part of mathematics.

2 Mathematical reminders

2.1 Combinatory logic and lambda calculus

These theories are formalization of the notion of information processing.

Everything is represented by information processing or functions, even data. An elementary piece of data, like a boolean data (true or false), can be represented by a function with two variables, which gives the first one if the value is true, or the second one if the value is false. A structured information, for example a couple of value, is represented by a function that, when applied to the boolean true value, gives the first value of the couple, and when applied to the boolean false, gives the second value of the couple.

A function with two variables is represented by a function that, when applied to the first variable, gives another function which, when applied to the second variable, gives the final result, and so on. A function that gives several result can be represented by a function that gives a structured containing all the results. So we have to consider only functions that, when applied to one variable, also called "argument" or sometimes "parameter", give one result.

The application of a function f to a variable x is written " $f\ x$ ".

" $(f\ x)\ y$ " may be written simply " $f\ x\ y$ ".

The following cases can be distinguished according to the relationship between the variable to which the function is applied and the result of the application of this function to this variable :

- The result is the variable itself : the function is called identity, written " I ". For any x , we have $I\ x = x$.
- The result is y which does not depend on the variable. The function is a constant function which always gives y as result. It is written " $K\ y$ ". For any x and y , we have $K\ y\ x = y$.

- The result is the result of the application of a to b, where both a and b may depend on the variable. In this case, the function is written "S f g" where f is a function that gives a when applied to the variable and g is a function that gives b when applied to the variable. For any f, g and x, $S\ f\ g\ x = f\ x\ (g\ x)$.

Any function can be represented by applications of I, K and S, or even only K and S, because $I = S\ K\ K$. This is called "combinatory logic".

But with this representation we obtain huge expressions difficult to read and understand. So we will introduce a notation to represent the function that, when applied to a variable x, gives a result M, where M represents an expression that may contain one or several occurrences of x. Different notations are used, for example :

- $[x].M$
- $\hat{x}.M$
- $\wedge x.M$
- $\backslash x.M$
- $\lambda x.M$
- $(\lambda x M)$
- $\lambda x[M]$
- $x \mapsto M$
- $\lambda x \rightarrow M$ or $\backslash x \rightarrow M$ (in Haskell)

$\lambda x.\lambda y.\lambda z.M$ may be written $\lambda xyz.M$.

This is the lambda calculus notation.

The combinatory logic representation of a function can be retrieved from its lambda calculus representation using the following correspondence rules :

- $\lambda x.x = I$
- $\lambda x.y = Ky$ if y does not contain x
- $\lambda x.(ab) = S(\lambda x.a)(\lambda x.b)$

The lambda calculus notation has a disadvantage for example $\lambda x.x$ and *lambday.y* represent the same function although they are different expressions. To avoid this disadvantage, we can use De Bruijn index. With this notation, this function is written $\lambda 1$. Each occurrence of a variable is replaced by a natural number n which means the variable corresponding to the n-th *lambda* in which it is nested, starting from the innermost.

With this notation we have :

- $I = \lambda 1$
- $K = \lambda \lambda 2$
- $S = \lambda \lambda \lambda 31(21)$

I will sometime use the notation $[\dots * \dots]$ or $[\dots \bullet \dots]$ for $\lambda \dots 1 \dots$.

See also <https://ryanflannery.net/research/logic-notes/Church-CalculiOfLambdaConversion.pdf> for more information about combinatory logic and lambda calculus.

2.2 Natural numbers

Natural numbers are defined by Peano axioms :

- 0 is a natural number.
- Every natural number has a successor.
- 0 is not the successor of any natural number.
- If two natural numbers have the same successor, they are equal.
- If 0 has a property, and if the fact that some natural number has this property implies that its successor also has this property, then every natural number has this property.

Arithmetical operations are defined as follow, where $\text{suc}(n)$ represents the successor of the natural number n :

- addition : $a + 0 = a$; $a + \text{suc}(b) = \text{suc}(a + b)$

- multiplication : $a \cdot 0 = 0; a \cdot \text{suc}(b) = (a \cdot b) + a$
- exponentiation : $a^0 = 1; a^{\text{suc}(b)} = a^b \cdot a$

For natural numbers, the addition and the multiplication are commutative : $a + b = b + a; a \cdot b = b \cdot a$, but not the exponentiation : generally $a^b \neq b^a$.

We shall see later that the addition and the multiplication of transfinite ordinal numbers are not commutative.

2.3 Composition and iteration of functions

The composition of two functions f and g , written $B f g$ or $f \circ g$ is a function satisfying $(f \circ g)x = f(gx)$.

The composition of a function with itself $f \circ f$ can be written f^2 .

More generally, the n -th iterate of the function f is defined by :

- $f^0 = I$
- $f^{\text{suc}(n)} = f \circ f^n$

and has the following properties :

- $f^{a+b} = f^b \circ f^a$
- $f^{a \cdot b} = (f^a)^b$

3 Defining transfinite ordinal numbers

Natural numbers can be represented by sets. Each number is represented by the set of all numbers smaller than it.

- $0 = \{\}$ (the empty set)
- $1 = \{0\} = \{\{\}\}$
- $2 = \{0, 1\} = \{\{\}, \{\{\}\}\}$
- $3 = \{0, 1, 2\} = \{\{\}, \{\{\}\}, \{\{\}, \{\{\}\}\}\}$
- ...

The successor of a natural number can be defined by $\text{suc}(n) = n + 1 = n \cup \{n\}$.

We have $n \leq p$ if and only if $n \subseteq p$.

\mathbb{N} is the set of all natural numbers : $\mathbb{N} = \{0, 1, 2, 3, \dots\}$ The natural numbers can be generalized to what is called "transfinite ordinal numbers", or simply "ordinal numbers" or "ordinals", by considering that infinite sets represent ordinal numbers. \mathbb{N} considered as an ordinal number is written ω . ω is the least ordinal which is greater than all the numbers $0, 1, 2, 3, \dots$ We say that ω is a limit ordinal and $0, 1, 2, 3, \dots$ is a fundamental sequence of ω . This is written : $\omega = \sup\{0, 1, 2, 3, \dots\}$ or $\omega = \lim(n \mapsto n)$ because the n -th element (starting with 0) of the sequence is n . An ordinal does not have a unique fundamental sequence, for example $1, 2, 3, 4, \dots$ is also a fundamental sequence of ω , because the least ordinal that is greater than all ordinals of this sequence is also ω (more generally the limit ordinal is the same if any number of the least items of a sequence are removed), and the same stands for the sequence $0, 2, 4, 6, \dots$

Any ordinal can be defined as the least ordinal strictly greater than all ordinals of a set : the empty set for 0, $\{\alpha\}$ for the successor of α , $\{\alpha_0, \alpha_1, \alpha_2, \dots\}$ for an ordinal with fundamental sequence $\alpha_0, \alpha_1, \alpha_2, \dots$

The successor can be generalized to transfinite ordinal numbers : $\text{suc}(\omega) = \omega + 1 = \omega \cup \{\omega\} = \{0, 1, 2, 3, \dots, \omega\}$; $\text{suc}(\text{suc}(\omega)) = \omega + 2 = \{0, 1, 2, 3, \dots, \omega, \omega + 1\}$ and so on.

Then we can consider the set $\{0, 1, 2, 3, \dots, \omega, \omega + 1, \omega + 2, \omega + 3, \dots\}$ which is a limit ordinal, and $\omega, \omega + 1, \omega + 2, \omega + 3, \dots$ is a fundamental sequence of this ordinal. This ordinal is $\omega + \omega = \omega \cdot 2$ or $\omega \times 2$ or $\omega 2$.

Then we can go on building greater and greater ordinals : $\omega \cdot 3, \dots, \omega \cdot \omega = \omega^2, \omega^3, \dots, \omega^\omega, \omega^{\omega^\omega}, \dots$

For natural numbers, arithmetical operations are defined as follows :

- addition : $a + 0 = a; a + \text{suc}(b) = \text{suc}(a + b)$
- multiplication : $a \cdot 0 = 0; a \cdot \text{suc}(b) = (a \cdot b) + a$
- exponentiation : $a^0 = 1; a^{\text{suc}(b)} = a^b \cdot a$

The definitions of arithmetical operations can be generalized to ordinals by adding canonical rules for limit ordinals :

- addition : $\alpha + 0 = \alpha; \alpha + \text{suc}(\beta) = \text{suc}(\alpha + \beta); \alpha + \lim(f) = \lim(n \mapsto \alpha + f(n))$

Now suppose we have $\omega^{\omega^{\omega^{\varepsilon_0+1}}} = \varepsilon_0^{\omega^{\omega^{\varepsilon_0+1}}}$.

We must prove the equality for $n+1$, which can be written $\omega^{\omega^{\omega^{\omega^{\varepsilon_0+1}}}} = \varepsilon_0^{\omega^{\omega^{\omega^{\varepsilon_0+1}}}}$.

We have $\omega^{\omega^{\omega^{\omega^{\varepsilon_0+1}}}} = \omega^{\varepsilon_0^{\omega^{\omega^{\varepsilon_0+1}}}}$ (by our hypothesis) $= \omega^{\varepsilon_0^{1+\varepsilon_0}}$ (for the same reason than $1 + \omega = \omega$, see above) $= \omega^{\varepsilon_0 \cdot \varepsilon_0^{\omega^{\varepsilon_0+1}}}$ $= (\omega^{\varepsilon_0})^{\varepsilon_0^{\omega^{\varepsilon_0+1}}}$ $= \varepsilon_0^{\varepsilon_0^{\omega^{\varepsilon_0+1}}}$. QED.

In a similar way, the limit of $\varepsilon_1, \varepsilon_1^{\varepsilon_1}, \varepsilon_1^{\varepsilon_1^{\varepsilon_1}}, \dots$ is called ε_2 and is also the limit of $\varepsilon_1 + 1, \omega^{\varepsilon_1+1}, \omega^{\omega^{\varepsilon_1+1}}, \dots$

We can define the same way ε_n for any natural number n . Then ε_ω is defined as the limit of $\varepsilon_0, \varepsilon_1, \varepsilon_2, \varepsilon_3, \dots$, and $\varepsilon_{\omega+1}$ as the limit of $\varepsilon_\omega, \varepsilon_\omega^{\varepsilon_\omega}, \varepsilon_\omega^{\varepsilon_\omega^{\varepsilon_\omega}}, \dots$ or $\varepsilon_\omega + 1, \omega^{\varepsilon_\omega+1}, \omega^{\omega^{\varepsilon_\omega+1}}, \dots$

After comes $\varepsilon_{\varepsilon_0}$, and the limit of $\varepsilon_0, \varepsilon_{\varepsilon_0}, \varepsilon_{\varepsilon_{\varepsilon_0}}, \dots$ which is called ζ_0 . This is the least fixed point of $\alpha \mapsto \varepsilon_\alpha$. The next one is ζ_1 which is the limit of $\zeta_0 + 1, \varepsilon_{\zeta_0+1}, \varepsilon_{\varepsilon_{\zeta_0+1}}, \dots$. Then we get $\zeta_2, \zeta_3, \dots, \zeta_\omega, \zeta_{\omega+1}, \dots, \zeta_{\varepsilon_0}, \dots, \zeta_{\zeta_0}, \dots, \zeta_{\zeta_{\zeta_0}}, \dots$. The limit of $0, \zeta_0, \zeta_{\zeta_0}, \zeta_{\zeta_{\zeta_0}}, \dots$ is called η_0 .

We can go on using successively different greek letters, or we can use functions indexed by numbers

- $\varphi_0(\alpha) = \omega^\alpha$
- $\varphi_1(\alpha) = \varepsilon_\alpha$
- $\varphi_2(\alpha) = \zeta_\alpha$
- $\varphi_3(\alpha) = \eta_\alpha$
- $\varphi_{\alpha+1}(\beta)$ is the $(1 + \beta)$ -th fixed point of $\xi \mapsto \varphi_\alpha(\xi)$.

or a function with two variables :

- $\varphi(0, \alpha) = \omega^\alpha$
- $\varphi(1, \alpha) = \varepsilon_\alpha$
- $\varphi(2, \alpha) = \zeta_\alpha$
- $\varphi(3, \alpha) = \eta_\alpha$
- $\varphi(\alpha + 1, \beta)$ is the $(1 + \beta)$ -th fixed point of $\xi \mapsto \varphi(\alpha, \xi)$.

Then we can enumerate the fixed points of the function $\alpha \mapsto \varphi(\alpha, 0)$ and define Γ_α as the $(1 + \alpha)$ -th fixed point of this function, or add another variable to the φ function and define $\varphi_{1,0}(\alpha)$ or $\varphi(1, 0, \alpha)$ as the $(1 + \alpha)$ -th fixed point of this function. So we have $\Gamma_\alpha = \varphi_{1,0}(\alpha) = \varphi(1, 0, \alpha)$.

More generally, we can define a function with any (finite) number of variables $\varphi_{\alpha_n, \alpha_{n-1}, \dots, \alpha_1, \alpha_0}(\beta) = \varphi(\alpha_n, \alpha_{n-1}, \dots, \alpha_1, \alpha_0, \beta)$, with $\varphi(\alpha) = \varphi_0(\alpha) = \varphi(0, \alpha) = \omega^\alpha$.

The notation $\varphi_{\alpha_n, \alpha_{n-1}, \dots, \alpha_1, \alpha_0}(\beta)$ has the advantage of highlighting the different role played by the last variable β .

For a complete definition of this Veblen function with finitely many variables, see for example :

https://en.wikipedia.org/wiki/Veblen_function :

"Let z be an empty string or a string consisting of one or more comma-separated zeros $0, 0, \dots, 0$ and s be an empty string or a string consisting of one or more comma-separated ordinals $\alpha_1, \alpha_2, \dots, \alpha_n$ with $\alpha_1 > 0$. The binary function $\varphi(\beta, \gamma)$ can be written as $\varphi(s, \beta, z, \gamma)$ where both s and z are empty strings.

The finitary Veblen functions are defined as follows:

- $\varphi(\gamma) = \omega^\gamma$
- $\varphi(z, s, \gamma) = \varphi(s, \gamma)$
- if $\beta > 0$, then $\varphi(s, \beta, z, \gamma)$ denotes the $(1 + \gamma)$ -th common fixed point of the functions $\xi \mapsto \varphi(s, \delta, \xi, z)$ for each $\delta < \beta$

(...)

The limit of the $\varphi(1, 0, \dots, 0)$ where the number of zeroes ranges over ω , is sometimes known as the "small" Veblen ordinal.

Every non-zero ordinal α less than the small Veblen ordinal (SVO) can be uniquely written in normal form for the finitary Veblen function:

$$\alpha = \varphi(s_1) + \varphi(s_2) + \dots + \varphi(s_k)$$

where

- k is a positive integer
- $\varphi(s_1) \geq \varphi(s_2) \geq \dots \geq \varphi(s_k)$
- s_m is a string consisting of one or more comma-separated ordinals $\alpha_{m,1}, \alpha_{m,2}, \dots, \alpha_{m,n_m}$ where $\alpha_{m,1} > 0$ and each $\alpha_{m,i} < \varphi(s_m)$

For limit ordinals $\alpha < SVO$, written in normal form for the finitary Veblen function:

- $(\varphi(s_1) + \varphi(s_2) + \dots + \varphi(s_k))[n] = \varphi(s_1) + \varphi(s_2) + \dots + \varphi(s_k)[n]$,
- $\varphi(\gamma)[n] =$
 - n if $\gamma = 1$
 - $\varphi(\gamma - 1) \cdot n$ if γ is a successor ordinal
 - $\varphi(\gamma[n])$ if γ is a limit ordinal
- $\varphi(s, \beta, z, \gamma)[0] = 0$ and $\varphi(s, \beta, z, \gamma)[n + 1] = \varphi(s, \beta - 1, \varphi(s, \beta, z, \gamma)[n], z)$ if $\gamma = 0$ and β is a successor ordinal,
- $\varphi(s, \beta, z, \gamma)[0] = \varphi(s, \beta, z, \gamma - 1) + 1$ and $\varphi(s, \beta, z, \gamma)[n + 1] = \varphi(s, \beta - 1, \varphi(s, \beta, z, \gamma)[n], z)$ if γ and β are successor ordinals,
- $\varphi(s, \beta, z, \gamma)[n] = \varphi(s, \beta, z, \gamma[n])$ if γ is a limit ordinal,
- $\varphi(s, \beta, z, \gamma)[n] = \varphi(s, \beta[n], z, \gamma)$ if $\gamma = 0$ and β is a limit ordinal,
- $\varphi(s, \beta, z, \gamma)[n] = \varphi(s, \beta[n], \varphi(s, \beta, z, \gamma - 1) + 1, z)$ if γ is a successor ordinal and β is a limit ordinal. "

The Veblen function can be generalized to transfinitely many variables with a finite number different from 0. Instead of writing the list of all the variable of the Veblen function, we can write only the non zero variables with position as indice, for example $\varphi(\alpha, 0, \beta, \gamma) = \varphi(\alpha_3, \beta_1, \gamma_0)$. We can then generalize the Veblen function by allowing any ordinal as indices, writing for example $SVO = \varphi(1_\omega)$. The limit of the ordinals that can be written with this notation is called the large Veblen ordinal (LVO).

According to Wikipedia, "The definition can be given as follows: let α be a transfinite sequence of ordinals (i.e., an ordinal function with finite support) which ends in zero (i.e., such that $\alpha_0=0$), and let $\alpha[0 \mapsto \gamma]$ denote the same function where the final 0 has been replaced by γ . Then $\gamma \mapsto \varphi(\alpha[0 \mapsto \gamma])$ is defined as the function enumerating the common fixed points of all functions $\xi \mapsto \varphi(\beta)$ where β ranges over all sequences which are obtained by decreasing the smallest-indexed nonzero value of α and replacing some smaller-indexed value with the indeterminate ξ (i.e., $\beta = \alpha[\iota_0 \mapsto \zeta, \iota \mapsto \xi]$ meaning that for the smallest index ι_0 such that α_{ι_0} is nonzero the latter has been replaced by some value $\zeta < \alpha_{\iota_0}$ and that for some smaller index $\iota < \iota_0$, the value $\alpha_\iota = 0$ has been replaced with ξ)."

Schütte brackets or Klammersymbols are another way to write Veblen fuctions with transfinitely many variables. A Schütte bracket consists in a matrix with two lines, with the positions of the variables in the second line in increasing order, and the corresponding values in the first line. This matrix is preceded by the function $\xi \mapsto \varphi(\xi)$. If we take $\xi \mapsto \omega^\xi$, we get the equivalent of the Veblen function. With this notation, the previous example is written :

$$(\xi \mapsto \omega^\xi) \begin{pmatrix} \gamma & \beta & \alpha \\ 0 & 1 & 3 \end{pmatrix}$$

In some of his papers, Harold Simmons puts the function after the matrix, which is more logical, the matrix being considered as a function which, when applied to a function, gives an ordinal :

$$\begin{pmatrix} \gamma & \beta & \alpha \\ 0 & 1 & 3 \end{pmatrix} (\xi \mapsto \omega^\xi)$$

When the function at the left of the matrix is $\xi \mapsto \omega^\xi$, it is sometimes omitted. Example :

$$\begin{pmatrix} \gamma & \beta & \alpha \\ 0 & 1 & 3 \end{pmatrix}$$

The corresponding fundamental sequences can be found in <https://sites.google.com/site/travelingtotheinfinity/fundamental-sequences-for-extended-veblen-function> .

Another possible notation is to represent the parameters of the φ function by a polynom of variable Ω where the exponent corresponds to the position of the variable, for example $\varphi(\alpha, 0, \beta, \gamma) = \varphi(\gamma_0, \beta_1, \alpha_3) = (\xi \mapsto \omega^\xi) \begin{pmatrix} \gamma & \beta & \alpha \\ 0 & 1 & 3 \end{pmatrix} = \varphi(\Omega^3 \cdot \alpha + \Omega \cdot \beta + \gamma)$. For Ω , we can choose an ordinal which is greater than all the ordinals we want to produce. Since they all are countable, we can

take for example $\Omega = \omega_1$ which is the least uncountable ordinal. The method consisting in using uncountable ordinals to define countable ordinals is called "collapsing". We will see later other examples of notations using this method.

Note that $\varphi(1, 0) = \varphi(\Omega)$ is the least α such that $\alpha = \varphi(\alpha) = \omega^\alpha$ (the least fixed point of $\alpha \mapsto \omega^\alpha$); $\varphi(1, 0, 0) = \varphi(\Omega^2) = \varphi(\Omega \cdot \Omega)$ is the least α such that $\alpha = \varphi(\alpha, 0) = \varphi(\Omega \cdot \alpha)$. Generally speaking, we can see that $f(\Omega)$ is the least fixed point of f . We shall see other examples of this equality later concerning ordinal collapsing functions. Note also that " Ω " can be replaced by " $1, 0$ " in the formulas.

If we want to distinguish the last variable, we can also use collapsing with the notation $\varphi_{\alpha_n, \dots, \alpha_0}(\beta)$, writing for example $\varphi_{\alpha, \beta, \gamma}(\delta) = \varphi_{\Omega^2 \cdot \alpha + \Omega \cdot \beta + \gamma}(\delta)$, or $\varphi(\alpha, \beta, \gamma, \delta) = \varphi(\Omega^2 \cdot \alpha + \Omega \cdot \beta + \gamma, \delta)$.

See Veblen's article "Continuous Increasing Functions of Finite and Transfinite Ordinals" (<http://www.ams.org/journals/tran/1908-009-03/S0002-9947-1908-1500814-9/S0002-9947-1908-1500814-9.pdf>) for more information.

Here is an Agda implementation of the Veblen function with transfinitely many variables :

```
{-
  A definition of the large Veblen ordinal in Agda
  by Jacques Bailhache, March 2016

  See https://en.wikipedia.org/wiki/Veblen\_function

  (1) phi(a)=w**a for a single variable,

  (2) phi(0,an-1,...,a0)=phi(an-1,...,a0), and

  (3) for a>0, c->phi(an,...,ai+1,a,0,...,0,c) is the function enumerating the common fixed points of the
      functions x->phi(an,...,ai+1,b,x,0,...,0) for all b<a.

  (4) Let a be a transfinite sequence of ordinals (i.e., an ordinal function with finite support) which ends in
      zero (i.e., such that a0=0), and let a[0->c] denote the same function where the final 0 has been replaced
      by c.
      Then c->phi(a[0->c]) is defined as the function enumerating the common fixed points of all functions
      x->phi(b) where b ranges over all sequences which are obtained by decreasing the smallest-indexed nonzero
      value of a and replacing some smaller-indexed value with the indeterminate x (i.e., b=a[i0->z,i->x]
      meaning that for the smallest index i0 such that ai0 is nonzero the latter has been replaced by some value
      z<ai0 and that for some smaller index i<i0, the value ai=0 has been replaced with x).

-}

module LargeVeblen where

data Nat : Set where
  0 : Nat
  1+ : Nat -> Nat

data Ord : Set where
  Zero : Ord
  Suc : Ord -> Ord
  Lim : (Nat -> Ord) -> Ord

-- rpt n f x = f^n(x)
rpt : {t : Set} -> Nat -> (t -> t) -> t -> t
rpt 0 f x = x
rpt (1+ n) f x = rpt n f (f x)
```

```

-- smallest fixed point of f greater than x, limit of x, f x, f (f x), ...
fix : (Ord -> Ord) -> Ord -> Ord
fix f x = Lim (\n -> rpt n f x)

w = fix Suc Zero -- not a fixed point in this case !

-- cantor a b = b + w^a
cantor : Ord -> Ord -> Ord
cantor Zero a = Suc a
cantor (Suc b) a = fix (cantor b) a
cantor (Lim f) a = Lim (\n -> cantor (f n) a)

-- phi0 a = w^a
phi0 : Ord -> Ord
phi0 a = cantor a Zero

-- Another possibility is to use phi'0 instead of phi0 in the definition of phi,
-- this gives a phi function which grows slower
phi'0 : Ord -> Ord
phi'0 Zero = Suc Zero
phi'0 (Suc a) = Suc (phi'0 a)
phi'0 (Lim f) = Lim (\n -> phi'0 (f n))

-- Associative list of ordinals
infixr 40 _=>_&_
data OrdAList : Set where
  Zeros : OrdAList
  _=>_&_ : Ord -> Ord -> OrdAList -> OrdAList

-- Usage : phi al, where al is the associative list of couples index => value ordered by increasing values,
-- absent indexes corresponding to Zero values

phi : OrdAList -> Ord
phi
  Zeros = phi0 Zero -- (1) phi(0) = w**0 = 1
phi (Zero => a & Zeros) = phi0 a -- (1) phi(a) = w**a
phi (
  k => Zero & al) = phi al -- eliminate unnecessary Zero value
phi (Zero => a & k => Zero & al) = phi (Zero => a & al) -- idem
phi (Zero => a & Zero => b & al) = phi (Zero => a & al) -- should not appear but necessary for completeness
phi (Zero => Lim f & al) = Lim (\n -> phi (Zero => f n & al)) -- canonical treatment of limit
phi (
  Suc k => Suc b & al) = fix (\x -> phi (k => x & Suc k => b & al)) Zero
  -- (3) least fixed point
phi (Zero => Suc a & Suc k => Suc b & al) = fix (\x -> phi (k => x & Suc k => b & al)) (Suc (phi (Zero => a &
Suc k => Suc b & al))) -- (3) following fixed points
phi (
  Suc k => Lim f & al) = Lim (\n -> phi (Suc k => f n & al)) -- idem
phi (Zero => Suc a & Suc k => Lim f & al) = Lim (\n -> phi (k => Suc (phi (Zero => a & Suc k => Lim f & al)) &
Suc k => f n & al)) -- idem
phi (
  Lim f => Suc b & al) = Lim (\n -> phi (f n => (Suc Zero) & Lim f => b & al))
phi (Zero => Suc a & Lim f => Suc b & al) = Lim (\n -> phi (f n => phi (Zero => a & Lim f => Suc b & al) & Lim
f => b & al))
phi (
  Lim f => Lim g & al) = Lim (\n -> phi (Lim f => g n & al))
phi (Zero => Suc a & Lim f => Lim g & al) = Lim (\n -> phi (f n => phi (Zero => a & Lim f => Lim g & al) & Lim
f => g n & al))

SmallVeblen = phi (w => Suc Zero & Zeros)

LargeVeblen = fix (\x -> phi (x => Suc Zero & Zeros)) (Suc Zero)

```


Normally it should terminate because the parameter of `phi` lexicographically decreases, but Agda is not clever enough so it must be called with no termination check option :

[illegible]

Main>

5 Going beyond Veblen function with transfinitely many variables

...

There is another way to express this construction.

There are different conventions for $\varphi_0(x)$, like ω^x or ε_x . We can write explicitly the convention chosen for φ_0 by writing " $\varphi_f(\alpha, \beta)$ " for " $\varphi_\alpha(\beta)$ with function f used for φ_0 ". With this notation we have:

- $\varphi_f(0, \beta) = f(\beta)$
- $\varphi_f(\alpha + 1, \beta) = (1 + \beta)$ th fixed point of the function $\beta \mapsto \varphi_f(\alpha, \beta)$
- $\varphi_f(\lambda, \beta) = (1 + \beta)$ th common fixed point of the function $\beta \mapsto \varphi_f(\alpha, \beta)$ for all $\alpha < \lambda$, if λ is a limit ordinal.

(See <http://www.cs.man.ac.uk/~hsimmons/TEMP/OrdNotes.pdf>)

Then we generalize the binary function $\varphi_f(\alpha, \beta)$ to finitely many variables: for example $\varphi_f(1, 0, \alpha) = (1 + \alpha)$ th common fixed point of the function $\xi \mapsto \varphi_f(\xi, 0)$ (see https://en.wikipedia.org/wiki/Veblen_function) and to infinitely many variables with a finite number of them different from 0, for example $\varphi_f(1_\omega)$.

Then we can define new φ functions by taking for φ_0 the function $\xi \mapsto \varphi_f(1_\xi)$ and define functions $\varphi_{\xi \mapsto \varphi_f(1_\xi)}$ with 2 variables, with finitely many variables and with transfinitely many variables.

To make a correspondence with my previous construction, if f is the function $\xi \mapsto \omega^\xi$, then $\varphi_f(\alpha, \beta)$ corresponds to what I wrote $\varphi_\alpha(\beta)$, and $\varphi_{\xi \mapsto \varphi_f(1_\xi)}(\alpha, \beta)$ to $\varphi_\alpha^+(\beta)$.

If we define the function S by $S(f)(\xi) = \varphi_f(1_\xi)$, then $\varphi_{\xi \mapsto \varphi_f(1_\xi)}$ can be written $\varphi_{S(f)}$. We can then consider $\varphi_{S(S(f))}$ and so on. Given an ordinal α , we can iterate transfinitely " α times" the application of S to an initial function f_0 , for example $f_0(\xi) = \omega^\xi$, to obtain a function which I will write $S^\alpha(f_0)$. We can use this function to define a function $\varphi_{S^\alpha(f_0)}$ which permits to construct big ordinals.

6 Simmons notation

6.1 Presentation

Harold Simmons defined a notation (see <http://www.cs.man.ac.uk/~hsimmons/ORDINAL-NOTATIONS/ordinal-notations.html>) based on fixed points enumeration which "contains" Veblen functions and permits to go further.

He uses the lambda calculus formalism, in which $f \ x$ represents the application of function f to x , and $f \ x \ y = (f \ x) \ y$ the application of function f to x which gives another function which is applied to y giving the final result. He uses the notation $x \mapsto y$ to represent the function which, when applied to x , gives y (instead of the traditional lambda calculus notation $\lambda x. y$).

He also uses the notation ω^\bullet for $\alpha \mapsto \omega^\alpha$.

$f \circ g$ represents the composition of functions f and g : $(f \circ g)\alpha = f(g\alpha)$.

f^α is a canonical generalization of of exponentiation of a function to an ordinal power : f^n represents $f \circ f \circ \dots \circ f$ with f repeated n times, $f^\omega \zeta$ is the limit of $\zeta, f \ \zeta, f(f \ \zeta), \dots, f^{\omega+1} \zeta = f(f^\omega \zeta)$ and so on.

More precisely, Simmons gives the following definitions in <http://www.cs.man.ac.uk/~hsimmons/TEMP/OrdNotes.pdf> page 11 :

- $g^0 \zeta = \zeta$
- $g^{\alpha+1} \zeta = g(g^\alpha \zeta)$
- $g^\lambda \zeta = V\{g^\alpha \zeta \mid \alpha < \lambda\}$ (if λ is a limit ordinal, where V denotes the pointwise supremum)

and the following equivalent definitions in <http://www.cs.man.ac.uk/~hsimmons/ORDINAL-NOTATIONS/Fruitful.pdf> page 4 :

- $g^0 = id$
- $g^{\alpha+1} = g \circ g^\alpha$
- $g^\lambda = V\{g^\alpha \mid \alpha < \lambda\}$

and he generalizes these definitions to higher order functions.

Then Simmons defines the following functions :

Fix $f \zeta = f^\omega(\zeta + 1) = \text{limit of } \zeta + 1, f(\zeta + 1), f(f(\zeta + 1)), \dots$ is the least fixed point of the function f which is strictly greater than ζ , which means the least ordinal ν satisfying $f \ \nu = \nu$ and $\nu > \zeta$.

Next $= \text{Fix } \omega^\bullet = \text{Fix}(\alpha \mapsto \omega^\alpha)$; *Next* ζ is the next ε_α after ζ .

$[0]h = \text{Fix}(\alpha \mapsto h^\alpha 0)$

$[1]hg = \text{Fix}(\alpha \mapsto h^\alpha g 0)$

$[2]hgf = \text{Fix}(\alpha \mapsto h^\alpha g f 0)$

... and so on ...

In <http://www.cs.man.ac.uk/~hsimmons/ORDINAL-NOTATIONS/OrdSlides.pdf> Simmons gives another equivalent definition :

$$\begin{aligned} [0]h &= \text{Fix}(\alpha \mapsto h^\alpha \omega) \\ [1]hg &= \text{Fix}(\alpha \mapsto h^\alpha g \omega) \\ [2]hgf &= \text{Fix}(\alpha \mapsto h^\alpha g f \omega) \end{aligned}$$

Simmons also defines :

$$\begin{aligned} \text{Veb } f \zeta &= (\text{Fix } f)^{1+\zeta} 0 \text{ is the } (1 + \zeta)\text{-th fixed point of } f \\ \text{Enm } h \alpha &= h^{1+\alpha} 0 \\ \text{Veb} &= \text{Enm} \circ \text{Fix} \\ [0] &= \text{Fix} \circ \text{Enm} \\ \text{Fix} \circ \text{Veb} &= \text{Fix} \circ \text{Enm} \circ \text{Fix} = [0] \circ \text{Fix} \\ \text{Fix} \circ \text{Veb}^\alpha &= [0]^\alpha \circ \text{Fix} \\ \Delta[0] &= \omega \\ \Delta[1] &= \text{Next } \omega = \varepsilon_0 \\ \Delta[2] &= [0] \text{Next } \omega = \text{least } \nu \text{ with } \nu = \text{Next}^\nu \omega = \zeta_0 \\ \Delta[3] &= [1][0] \text{Next } \omega = \text{least } \nu \text{ with } \nu = [0]^\nu \text{Next } \omega = \Gamma_0 \\ \Delta[4] &= [2][1][0] \text{Next } \omega = \text{least } \nu \text{ with } \nu = [1]^\nu [0] \text{Next } \omega = LVO \text{ (large Veblen ordinal)} \\ \dots &\text{ and so on } \dots \end{aligned}$$

6.2 Implementation

Here is an implementation of the Simmons hierarchy in Haskell :

```
module Simmons where

-- Natural numbers
data Nat
  = ZeroN
  | SucN Nat

-- Ordinals
data Ord
  = Zero
  | Suc Ord
  | Lim (Nat -> Ord)

-- Ordinal corresponding to a given natural
ordOfNat ZeroN = Zero
ordOfNat (SucN n) = Suc (ordOfNat n)

-- omega
w = Lim ordOfNat

lim0 s = Lim s
lim1 f x = lim0 (\n -> f n x)
lim2 f x = lim1 (\n -> f n x)

-- this does not work :
-- lim ZeroN s = Lim s
-- lim (SucN p) f = \x -> lim p (\n -> f n x)

-- f^a(x)
fpower0 f Zero x = x
fpower0 f (Suc a) x = f (fpower0 f a x)
fpower0 f (Lim s) x = Lim (\n -> fpower0 f (s n) x)
```

```

fpower 1 f Zero x = x
fpower 1 f (Suc a) x = f (fpower 1 f a x)
fpower 1 f (Lim s) x = 1 (\n -> fpower 1 f (s n) x)

-- fix f z = least fixed point of f which is > z
fix f z = fpower lim0 f w (Suc z) -- Lim (\n -> fpower0 f (ordOfNat n) (Suc z))

-- cantor b a = a + w^b
cantor Zero a = Suc a
cantor (Suc b) a = fix (cantor b) a
cantor (Lim s) a = Lim (\n -> cantor (s n) a)

-- expw a = w^a
expw a = cantor a Zero

-- next a = least epsilon_b > a
next = fix expw

-- [0]
simmons0 h = fix (\a -> fpower lim0 h a Zero)

-- [1]
simmons1 h1 h0 = fix (\a -> fpower lim1 h1 a h0 Zero)

-- [2]
simmons2 h2 h1 h0 = fix (\a -> fpower lim2 h2 a h1 h0 Zero)

-- Large Veblen ordinal
lvo = simmons2 simmons1 simmons0 next w

```

\$ hugs

```

-- -- -- -- --
||  ||  ||  ||  ||  ||  ||__  Hugs 98: Based on the Haskell 98 standard
||__||  ||__||  ||__||  __||  Copyright (c) 1994-2005
||---||          ---||  World Wide Web: http://haskell.org/hugs
||  ||          Bugs: http://hackage.haskell.org/trac/hugs
||  || Version: September 2006 -----

```

Haskell 98 mode: Restart with command line option -98 to enable extensions

```

Type :? for help
Hugs> :load simmons
Simmons> lvo
ERROR - Cannot find "show" function for:
*** Expression : lvo
*** Of type    : Ord

Simmons>

```

6.3 Correspondence with Veblen functions

ε_0 is the next ε_α after 0 (or after ω , or after any ordinal less than ε_0 , so we have $\varepsilon_0 = \text{Next } 0 = \text{Next } \omega$).

ε_1 is the next ε_α after ε_0 , so we have $\varepsilon_1 = \text{Next } \varepsilon_0 = \text{Next } (\text{Next } 0) = \text{Next}^2 0 = \text{Next } (\text{Next } \omega) = \text{Next}^2 \omega$.
 ε_2 is the next ε_α after ε_1 , so we have $\varepsilon_2 = \text{Next } \varepsilon_1 = \text{Next } (\text{Next } (\text{Next } 0)) = \text{Next}^3 0 = \text{Next } (\text{Next } (\text{Next } \omega)) = \text{Next}^3 \omega$.
 \dots
 ε_ω is the limit of $\varepsilon_0, \varepsilon_1, \varepsilon_2, \dots$. It is the limit of $\text{Next}^1 0, \text{Next}^2 0, \text{Next}^3 0, \dots$ which is $\text{Next}^\omega 0$.
More generally, we have $\varepsilon_\alpha = \varphi(1, \alpha) = \text{Next}^{1+\alpha} 0 = \text{Next}^{1+\alpha} \omega$.

$\zeta_0 = \varphi(2, 0)$ is the least fixed point of $\alpha \mapsto \varepsilon_\alpha$ (greater than 0), so $\zeta_0 = \text{Fix}(\alpha \mapsto \varepsilon_\alpha) 0 = \text{Fix}(\alpha \mapsto \text{Next}^{1+\alpha} 0) 0 = \text{Fix}(\alpha \mapsto \text{Next}^\alpha 0) 0$ (because the "1+" is "absorbed" after a few iterations) $= [0] \text{Next } 0$. Since ζ_0 is also greater than ω , it is also $[0] \text{Next } \omega$ according to a similar computation.

$\zeta_1 = \varphi(2, 1)$ is the next fixed point of $\alpha \mapsto \varepsilon_\alpha$, the least one which is strictly greater than ζ_0 , so $\zeta_1 = \text{Fix}(\alpha \mapsto \varepsilon_\alpha) \zeta_0 = \text{Fix}(\alpha \mapsto \text{Next}^\alpha 0) \zeta_0 = [0] \text{Next } \zeta_0 = [0] \text{Next}([0] \text{Next } 0) = ([0] \text{Next})^2 0 = [0] \text{Next}([0] \text{Next } \omega) = ([0] \text{Next})^2 \omega$.

More generally, $\zeta_\alpha = ([0] \text{Next})^{1+\alpha} 0$.

Similar computations give $\eta_0 = \varphi(3, 0) = [0]^2 \text{Next } 0$ and $\eta_\alpha = ([0]^2 \text{Next})^{1+\alpha} 0$.

More generally, $\varphi(1 + \beta, \alpha) = ([0]^\beta \text{Next})^{1+\alpha} 0$ or $([0]^\beta \text{Next})^{1+\alpha} \omega$.

$\Gamma_0 = \varphi(1, 0, 0)$ is the least fixed point (greater than 0) of the function $\alpha \mapsto \varphi(\alpha, 0)$ or $\alpha \mapsto \varphi(1 + \alpha, 0)$ (for the same reason of "absorption" of "1+" than previously), so $\Gamma_0 = \text{Fix}(\alpha \mapsto \varphi(1 + \alpha, 0)) 0 = \text{Fix}(\alpha \mapsto ([0]^\alpha \text{Next})^{1+0} 0) 0 = \text{Fix}(\alpha \mapsto [0]^\alpha \text{Next } 0) 0 = [1][0] \text{Next } 0$.

$\Gamma_1 = \varphi(1, 0, 1)$ is the next fixed point : $\Gamma_1 = \text{Fix}(\alpha \mapsto [0]^\alpha \text{Next } 0) \Gamma_0 = [1][0] \text{Next } \Gamma_0 = [1][0] \text{Next } ([1][0] \text{Next } 0) = ([1][0] \text{Next})^2 0$.

More generally, we have $\varphi(1, 0, \alpha) = ([1][0] \text{Next})^{1+\alpha} 0$.

$\varphi(1, 1, 0)$ is the least fixed point (greater than 0) of the function $\alpha \mapsto \varphi(1, 0, \alpha)$, so it is $\text{Fix}(\alpha \mapsto \varphi(1, 0, \alpha)) 0 = \text{Fix}(\alpha \mapsto ([1][0] \text{Next})^{1+\alpha} 0) 0 = \text{Fix}(\alpha \mapsto ([1][0] \text{Next})^\alpha 0) 0$ (absorption of 1+) $= [0]([1][0] \text{Next}) 0$.

$\varphi(1, 1, 1)$ is the next fixed point $\text{Fix}(\alpha \mapsto ([1][0] \text{Next})^\alpha 0) \varphi(1, 1, 0) = ([0]([1][0] \text{Next})([1][0] \text{Next}) 0) = ([0]([1][0] \text{Next}))^2 0$.

More generally, $\varphi(1, 1, \alpha) = ([0]([1][0] \text{Next}))^{1+\alpha} 0$.

$\varphi(1, 2, 0)$ is the least fixed point (greater than 0) of the function $\alpha \mapsto \varphi(1, 1, \alpha)$, $\text{Fix}(\alpha \mapsto \varphi(1, 1, \alpha)) 0 = \text{Fix}([0]([1][0] \text{Next})^{1+\alpha} 0) 0 = \text{Fix}(\alpha \mapsto ([0]([1][0] \text{Next}))^\alpha 0) 0 = [0]([0]([1][0] \text{Next})) 0 = [0]^2([1][0] \text{Next}) 0$.

Like previously, $\varphi(1, 2, \alpha)$ is the $(1 + \alpha)$ -th fixed point of the previous function, which is $([0]^2([1][0] \text{Next}))^{1+\alpha} 0$.

More generally, $\varphi(1, \beta, \alpha) = ([0]^\beta([1][0] \text{Next}))^{1+\alpha} 0$.

$\varphi(2, 0, 0)$ is the least fixed point (greater than 0) of the function $\beta \mapsto \varphi(1, \beta, 0)$, which is $\text{Fix}(\alpha \mapsto \varphi(1, \beta, 0)) 0 = \text{Fix}(\beta \mapsto ([0]^\beta([1][0] \text{Next}))^{1+0} 0) 0 = \text{Fix}(\beta \mapsto [0]^\beta([1][0] \text{Next}) 0) 0 = [1][0]([1][0] \text{Next}) 0 = ([1][0])^2 \text{Next } 0$.

The $(1 + \alpha)$ -th fixed point of the previous function is $\varphi(2, 0, \alpha) = (([1][0])^2 \text{Next})^{1+\alpha} 0$.

The least fixed point of the function $\alpha \mapsto \varphi(2, 0, \alpha)$ is $\varphi(2, 1, 0) = \text{Fix}(\alpha \mapsto \varphi(2, 0, \alpha)) 0 = \text{Fix}(\alpha \mapsto (([1][0])^2 \text{Next})^{1+\alpha} 0) 0 = \text{Fix}(\alpha \mapsto ([1][0])^2 \text{Next}^\alpha 0) 0 = [0]([1][0])^2 \text{Next } 0$ and its $(1 + \alpha)$ -th fixed point is $\varphi(2, 1, \alpha) = ([0]([1][0])^2 \text{Next})^{1+\alpha} 0$.

More generally, we have $\varphi(2, \beta, \alpha) = ([0]^\beta([1][0])^2 \text{Next})^{1+\alpha} 0$.

The general formula with three variables (with $\gamma \neq 0$) is $\varphi(\gamma, \beta, \alpha) = ([0]^\beta([1][0])^\gamma \text{Next})^{1+\alpha} 0$.

In particular, we have $\varphi(\gamma, 0, 0) = ([1][0])^\gamma \text{Next } 0$.

Using collapsing, we can write $\varphi(\gamma, \beta, \alpha) = \varphi_{\gamma, \beta}(\alpha) = \varphi_{\Omega \cdot \gamma + \beta}(\alpha) = \varphi(\Omega \cdot \gamma + \beta, \alpha) = \varphi(1 + \Omega \cdot \gamma + \beta, \alpha) = ([0]^{\Omega \cdot \gamma + \beta} \text{Next})^{1+\alpha} 0 = ([0]^\beta([0]^\Omega)^\gamma \text{Next})^{1+\alpha} 0 = ([0]^\beta([1][0])^\gamma \text{Next})^{1+\alpha} 0$ if we consider that $[0]^\Omega = [1][0]$.

$\varphi(1, 0, 0, 0)$ is the least fixed point of the function $\gamma \mapsto \varphi(\gamma, 0, 0)$, $\text{Fix}(\gamma \mapsto \varphi(\gamma, 0, 0)) 0 = \text{Fix}(\gamma \mapsto ([1][0])^\gamma \text{Next } 0) 0 = [1]([1][0] \text{Next } 0) 0 = [1]^2[0] \text{Next } 0$.

All of these computations could be done with ω instead of 0 at the end of the formulas so we also have $\varphi(\gamma, \beta, \alpha) = ([0]^\beta([1][0])^\gamma \text{Next})^{1+\alpha} \omega$.

In a similar way, we can obtain the formula with 4 variables :

$\varphi(1, 0, 0, \alpha) = ([1]^2[0] \text{Next})^{1+\alpha} 0$

$\varphi(1, 0, 1, 0) = \text{Fix}(\alpha \mapsto ([1]^2[0] \text{Next})^\alpha 0) 0 = [0]([1]^2[0]) 0$

$\varphi(1, 0, 1, \alpha) = ([0]([1]^2[0] \text{Next}))^{1+\alpha} 0$

$\varphi(1, 0, \beta, \alpha) = ([0]^\beta([1]^2[0] \text{Next}))^{1+\alpha} 0$

$\varphi(1, 1, 0, 0) = \text{Fix}(\alpha \mapsto \varphi(1, 0, \alpha, 0)) 0 = \text{Fix}(\alpha \mapsto [0]^\alpha([1]^2[0] \text{Next}) 0) 0 = [1][0]([1]^2[0] \text{Next}) 0$

$\varphi(1, 1, 0, \alpha) = ([1][0]([1]^2[0] \text{Next}))^{1+\alpha} 0$

$\varphi(1, 1, 1, 0) = \text{Fix}(\alpha \mapsto \varphi(1, 1, 0, \alpha)) 0 = \text{Fix}(\alpha \mapsto ([1][0]([1]^2[0] \text{Next}))^\alpha 0) 0 = [0]([1][0]([1]^2[0] \text{Next})) 0$

$\varphi(1, 1, 1, \alpha) = ([0]([1][0]([1]^2[0] \text{Next})))^{1+\alpha} 0$

$\varphi(1, 1, \beta, \alpha) = ([0]^\beta([1][0]([1]^2[0] \text{Next})))^{1+\alpha} 0$

$\varphi(1, 2, 0, 0) = \text{Fix}(\alpha \mapsto \varphi(1, 1, \alpha, 0)) 0 = \text{Fix}(\alpha \mapsto [0]^\alpha([1][0]([1]^2[0] \text{next})) 0) 0 = [1][0]([1][0]([1]^2[0] \text{Next})) 0 = ([1][0])^2([1]^2[0] \text{Next}) 0$

$\varphi(1, 0, 0, 0) = [1]^2[0] \text{Next } 0$

$\varphi(1, 1, 0, 0) = [1][0]([1]^2[0]Next)0$
 $\varphi(1, 2, 0, 0) = ([1][0])^2([1]^2[0]Next)0$
 $\varphi(1, \gamma, 0, 0) = ([1][0])^\gamma([1]^2[0]Next)0$
 $\varphi(1, \gamma, \beta, \alpha) = ([0]^\beta((([1][0])^\gamma([1]^2[0]Next))))^{1+\alpha}0$
 $\varphi(2, 0, 0, 0) = Fix(\alpha \mapsto \varphi(1, \alpha, 0, 0))0 = Fix(\alpha \mapsto ([1][0])^\alpha([1]^2[0]Next)0)0 = [1]([1][0])([1]^2[0]Next)0 = [1]^2[0]([1]^2[0]Next)0 = ([1]^2[0])^2Next0$
 $\varphi(\delta, 0, 0, 0) = ([1]^2[0])^\delta Next 0$
 The general formula with four variables is :
 $\varphi(\delta, \gamma, \beta, \alpha) = ([0]^\beta((([1][0])^\gamma((([1]^2[0])^\delta Next))))^{1+\alpha}0 = ([0]^\beta((([1][0])^\gamma((([1]^2[0])^\delta Next))))^{1+\alpha}\omega$
 and so on.

Using collapsing, we can write $\varphi(\delta, \gamma, \beta, \alpha) = \varphi_{\delta, \gamma, \beta}(\alpha) = \varphi_{\Omega^2 \cdot \delta + \Omega \cdot \gamma + \beta}(\alpha) = \varphi(\Omega^2 \cdot \delta + \Omega \cdot \gamma + \beta, \alpha) = \varphi(1 + \Omega^2 \cdot \delta + \Omega \cdot \gamma + \beta, \alpha) = ([0]^{\Omega^2 \cdot \delta + \Omega \cdot \gamma + \beta} Next)^{1+\alpha}0 = ([0]^\beta((([0]^\Omega)^\gamma((([0]^{\Omega^2})^\delta Next))))^{1+\alpha}0 = ([0]^\beta((([1][0])^\gamma((([1]^2[0])^\delta Next))))^{1+\alpha}0$ if we consider that $[0]^\Omega = [1][0]$ and $[0]^{\Omega^2} = ([0]^\Omega)^\Omega = ([1][0])^\Omega = [1]([1][0]) = [1]^2[0]$.

The small Veblen ordinal is the limit of :

$\varphi(1) = \omega, \varphi(1, 0) = Next \omega, \varphi(1, 0, 0) = [1][0]Next \omega, \varphi(1, 0, 0, 0) = [1]^2[0]Next \omega, \varphi(1, 0, 0, 0, 0) = [1]^3[0]Next \omega, \dots$
 This limit is $[1]^\omega[0]Next \omega = [1]^\omega[0]Next 0$.

Allowing variables at any finite or transfinite positions (which is equivalent to Schütte brackets or Klammersymbols) gives ordinals smaller than the large Veblen ordinal which is the least fixed point of the function $\alpha \mapsto \varphi(1_\alpha)$. It is $Fix(\alpha \mapsto \varphi(1_\alpha))0 = Fix(\alpha \mapsto [1]^\alpha[0]Next 0)0 = [2][1][0]Next 0$
 $[2][1][0]Next 0$ or $[2][1][0]Next \omega$.

The conversion rule from Schütte Klammersymbol to Simmons notation are described by Simmons in his paper : <http://www.cs.man.ac.uk/NOTATIONS/FromBelow.pdf> (Simmons also wrote other papers but it seems to me that they contain inaccuracies and maybe even errors).

In summary :

$Fix f\zeta = f^\omega(\zeta + 1)$

$Enm h \alpha = h^{1+\alpha}0$

$Next = Fix(\alpha \mapsto \omega^\alpha)$

$[0]h = Fix(\alpha \mapsto h^\alpha 0)$

$[1]hg = Fix(\alpha \mapsto h^\alpha g 0)$

$\nabla \begin{bmatrix} \alpha + 1 \\ i + 1 \end{bmatrix} = ([1]^i[0])^{1+\alpha}$ if $i \neq 0$; $[0]^\alpha$ if $i = 0$

$\nabla \begin{bmatrix} \alpha_1 + 1 & \dots & \alpha_s + 1 \\ i_1 + 1 & \dots & i_s + 1 \end{bmatrix} = \nabla \begin{bmatrix} \alpha_1 + 1 \\ i_1 + 1 \end{bmatrix} \circ [0] \circ \dots \circ [0] \circ \nabla \begin{bmatrix} \alpha_s + 1 \\ i_s + 1 \end{bmatrix}$

where $f \circ g$ is the composition of functions f and g : $(f \circ g)x = f(g x)$

$Sch \begin{bmatrix} 1 + \alpha_1 & \dots & 1 + \alpha_s \\ 1 + i_1 & \dots & 1 + i_s \end{bmatrix} = Enm \circ \nabla \begin{bmatrix} 1 + \alpha_1 & \dots & 1 + \alpha_s \\ 1 + i_1 & \dots & 1 + i_s \end{bmatrix} \circ Fix$

f may be any function but it is usually $\alpha \mapsto \omega^\alpha$.

$f \begin{pmatrix} \zeta & 1 + \alpha_1 & \dots & 1 + \alpha_s \\ 0 & 1 + i_1 & \dots & 1 + i_s \end{pmatrix}$

$= Sch \begin{bmatrix} 1 + \alpha_1 & \dots & 1 + \alpha_s \\ 1 + i_1 & \dots & 1 + i_s \end{bmatrix} f\zeta$

$= (Enm \circ \nabla \begin{bmatrix} 1 + \alpha_1 & \dots & 1 + \alpha_s \\ 1 + i_1 & \dots & 1 + i_s \end{bmatrix} \circ Fix)f\zeta$

$= (Enm \circ \nabla \begin{bmatrix} \alpha_1 + 1 \\ i_1 + 1 \end{bmatrix} \circ [0] \circ \dots \circ [0] \circ \nabla \begin{bmatrix} \alpha_s + 1 \\ i_s + 1 \end{bmatrix} \circ Fix)f\zeta$

$= Enm((\nabla \begin{bmatrix} \alpha_1 + 1 \\ i_1 + 1 \end{bmatrix} \circ [0] \circ \dots \circ [0] \circ \nabla \begin{bmatrix} \alpha_s + 1 \\ i_s + 1 \end{bmatrix})(Fix f))\zeta$

$= (\nabla \begin{bmatrix} \alpha_1 + 1 \\ i_1 + 1 \end{bmatrix} \circ [0] \circ \dots \circ [0] \circ \nabla \begin{bmatrix} \alpha_s + 1 \\ i_s + 1 \end{bmatrix})(Fix f)^{1+\zeta}0$

If $f = \alpha \mapsto \omega^\alpha$, then $Fix f = Next$ and

$f \begin{pmatrix} \zeta & 1 + \alpha_1 & \dots & 1 + \alpha_s \\ 0 & 1 + i_1 & \dots & 1 + i_s \end{pmatrix} = (\nabla \begin{bmatrix} \alpha_1 + 1 \\ i_1 + 1 \end{bmatrix} \circ [0] \circ \dots \circ [0] \circ \nabla \begin{bmatrix} \alpha_s + 1 \\ i_s + 1 \end{bmatrix})Next^{1+\zeta}0$

Examples :

$$\begin{aligned}
& \varphi(1 + \beta, \alpha) \\
&= (\xi \mapsto \omega^\xi) \begin{pmatrix} \alpha & 1 + \beta \\ 0 & 1 \end{pmatrix} \\
&= ((\nabla \begin{bmatrix} \beta + 1 \\ 1 \end{bmatrix})(\text{Fix}(\xi \mapsto \omega^\xi)))^{1+\alpha} 0 \\
&= ((\nabla \begin{bmatrix} \beta + 1 \\ 1 \end{bmatrix})\text{Next})^{1+\alpha} 0 \\
&= ([0]^\beta \text{Next})^{1+\alpha} 0
\end{aligned}$$

$$\begin{aligned}
& \varphi(1 + \gamma, 1 + \beta, \alpha) \\
&= (\xi \mapsto \omega^\xi) \begin{pmatrix} \alpha & 1 + \beta & 1 + \gamma \\ 0 & 1 & 2 \end{pmatrix} \\
&= ((\nabla \begin{bmatrix} \beta + 1 \\ 1 \end{bmatrix} \circ [0] \circ \nabla \begin{bmatrix} \gamma + 1 \\ 2 \end{bmatrix})(\text{Fix}(\xi \mapsto \omega^\xi)))^{1+\alpha} 0 \\
&= ((\nabla \begin{bmatrix} \beta + 1 \\ 1 \end{bmatrix} \circ [0] \circ \nabla \begin{bmatrix} \gamma + 1 \\ 2 \end{bmatrix})\text{Next})^{1+\alpha} 0 \\
&= (([0]^\beta \circ [0] \circ ([1][0])^{1+\gamma})\text{Next})^{1+\alpha} 0 \\
&= ([0]^{1+\beta}([1][0])^{1+\gamma}\text{Next})^{1+\alpha} 0
\end{aligned}$$

Compare with the previously found formula :

if $\gamma > 0$, $\varphi(\gamma, \beta, \alpha) = ([0]^\beta([1][0])^\gamma \text{Next})^{1+\alpha} 0$

and note the "round trip" $1 + \gamma \rightarrow \gamma + 1 \rightarrow 1 + \gamma$.

$$\begin{aligned}
& \varphi(1 + \delta, 1 + \gamma, 1 + \beta, \alpha) \\
&= (\xi \mapsto \omega^\xi) \begin{pmatrix} \alpha & 1 + \beta & 1 + \gamma & 1 + \delta \\ 0 & 1 & 2 & 3 \end{pmatrix} \\
&= ((\nabla \begin{bmatrix} \beta + 1 \\ 1 \end{bmatrix} \circ [0] \circ \nabla \begin{bmatrix} \gamma + 1 \\ 2 \end{bmatrix} \circ [0] \circ \nabla \begin{bmatrix} \delta + 1 \\ 3 \end{bmatrix})(\text{Fix}(\xi \mapsto \omega^\xi)))^{1+\alpha} 0 \\
&= ((\nabla \begin{bmatrix} \beta + 1 \\ 1 \end{bmatrix} \circ [0] \circ \nabla \begin{bmatrix} \gamma + 1 \\ 2 \end{bmatrix} \circ [0] \circ \nabla \begin{bmatrix} \delta + 1 \\ 3 \end{bmatrix})\text{Next})^{1+\alpha} 0 \\
&= (([0]^\beta \circ [0] \circ ([1][0])^{1+\gamma} \circ [0] \circ ([1]^2[0])^{1+\delta})\text{Next})^{1+\alpha} 0 \\
&= ([0]^{1+\beta}([1][0])^{1+\gamma}([0]([1]^2[0])^{1+\delta}\text{Next})))^{1+\alpha} 0 \\
&= ([0]^{1+\beta}([1][0])^{1+\gamma}([1]^2[0])^{1+\delta}\text{Next}))^{1+\alpha} 0
\end{aligned}$$

because $[0]$ is absorbed by the following operator (see <http://www.cs.man.ac.uk/~hsimmons/ORDINAL-NOTATIONS/FromBelow.pdf> p 33, 6.7)

Compare with the previously mentioned formula :

$\varphi(\delta, \gamma, \beta, \alpha) = ([0]^\beta([1][0])^\gamma([1]^2[0])^\delta \text{Next}))^{1+\alpha} 0$

The equality

$$(\xi \mapsto \omega^\xi) \begin{pmatrix} \zeta & 1 + \alpha_1 & \dots & 1 + \alpha_s \\ 0 & 1 + i_1 & \dots & 1 + i_s \end{pmatrix} = (\nabla \begin{bmatrix} \alpha_1 + 1 \\ i_1 + 1 \end{bmatrix} \circ [0] \circ \dots \circ [0] \circ \nabla \begin{bmatrix} \alpha_s + 1 \\ i_s + 1 \end{bmatrix})\text{Next})^{1+\zeta} 0$$

can be reformulated, distinguishing four cases :

- $(\xi \mapsto \omega^\xi) \begin{pmatrix} \zeta \\ 0 \end{pmatrix} = \varphi(0, \zeta) = \omega^\zeta$
- $(\xi \mapsto \omega^\xi) \begin{pmatrix} \zeta & 1 + \alpha \\ 0 & 1 \end{pmatrix} = \varphi(1 + \alpha, \zeta) = (\nabla \begin{bmatrix} \alpha + 1 \\ 1 \end{bmatrix} \text{Next})^{1+\zeta} 0 = ([0]^\alpha \text{Next})^{1+\zeta} 0$
- $(\xi \mapsto \omega^\xi) \begin{pmatrix} \zeta & 1 + \alpha_1 & 1 + \alpha_2 & \dots & 1 + \alpha_s \\ 0 & 1 & 1 + i_2 & \dots & 1 + i_s \end{pmatrix}$
 $= ((\nabla \begin{bmatrix} \alpha_1 + 1 \\ 1 \end{bmatrix} \circ [0] \circ \nabla \begin{bmatrix} \alpha_2 + 1 \\ i_2 + 1 \end{bmatrix} \circ [0] \circ \dots \circ [0] \circ \nabla \begin{bmatrix} \alpha_s + 1 \\ i_s + 1 \end{bmatrix})\text{Next})^{1+\zeta} 0$
 $= (([0]^{\alpha_1} \circ [0] \circ ([1]^{i_2}[0])^{1+\alpha_2} \circ [0] \circ \dots \circ [0] \circ ([1]^{i_s}[0])^{1+\alpha_s})\text{Next})^{1+\zeta} 0$
 $= (([0]^{1+\alpha_1} \circ ([1]^{i_2}[0])^{1+\alpha_2} \circ [0] \circ \dots \circ [0] \circ ([1]^{i_s}[0])^{1+\alpha_s})\text{Next})^{1+\zeta} 0$

$$= ((([0]^{1+\alpha_1} \circ ([1]^{i_2}[0])^{1+\alpha_2} \circ \dots \circ ([1]^{i_s}[0])^{1+\alpha_s})Next)^{1+\zeta}0$$

The first separating $[0]$ is combined with $[0]^{\alpha_1}$ giving $[0]^{1+\alpha_1}$ and the other are absorbed.

$$\begin{aligned} & \bullet (\xi \mapsto \omega^\xi) \left(\begin{pmatrix} \zeta & 1+\alpha_1 & \dots & 1+\alpha_s \\ 0 & 1+i_1 & \dots & 1+i_s \end{pmatrix} \text{ with } i_1 \neq 0 \right) \\ &= ((\nabla \begin{bmatrix} \alpha_1+1 \\ i_1+1 \end{bmatrix} \circ [0] \circ \dots \circ [0] \circ \nabla \begin{bmatrix} \alpha_s+1 \\ i_s+1 \end{bmatrix})Next)^{1+\zeta}0 \\ &= ((([1]^{i_1}[0])^{1+\alpha_1} \circ [0] \circ \dots \circ [0] \circ ([1]^{i_s}[0])^{1+\alpha_s})Next)^{1+\zeta}0 \\ &= ((([1]^{i_1}[0])^{1+\alpha_1} \circ \dots \circ ([1]^{i_s}[0])^{1+\alpha_s})Next)^{1+\zeta}0 \end{aligned}$$

The separating $[0]$ are absorbed.

We can see that the third case is contained in the fourth one if we remove the restriction $i_1 \neq 0$ because if $i_1 = 0$ we have $([1]^{i_1}[0])^{1+\alpha_1} = [0]^{1+\alpha_1}$ like in the third case.

For more information concerning the correspondence between Simmons notation and Schütte Klammersymbols, see : <http://www.cs.man.ac.uk/~hsimmons/ORDINAL-NOTATIONS/FromBelow.pdf> pages 28 - 34.

The Simmons notation can also be used to represent the notation going beyond Veblen functions that we saw previously.

As we saw previously, the large Veblen ordinal is the least fixed point of the function $\alpha \mapsto \varphi(1_\alpha)$ or $\alpha \mapsto (\xi \mapsto \omega^\xi) \left(\frac{1}{\alpha} \right)$. It is $Fix(\alpha \mapsto \varphi(1_\alpha))0 = Fix(\alpha \mapsto [1]^\alpha[0]Next\ 0)0 = [2][1][0]Next\ 0$.

Using collapsing, we can write it $\varphi(1_\Omega) = [1]^\Omega[0]Next\ 0$. Compare with the previously obtained equality $[0]^{\Omega^2} = ([0]^\Omega)^\Omega = ([1][0])^\Omega = [1]([1][0]) = [1]^2[0]$ which can be generalized to $[0]^{\Omega^\alpha} = [1]^\alpha[0]$. We can also write $LVO = \varphi_{\Omega^\Omega}(0) = [0]^{\Omega^\Omega}Next\ 0 = [1]^\Omega[0]Next\ 0 = [2][1][0]Next\ 0$ with $[1]^\Omega = [2][1]$.

The fixed points of this function $\alpha \mapsto \varphi(1_\alpha)$ are enumerated by the function F , so we have $LVO = F(0)$. More generally, the $(1 + \alpha)$ -th fixed point of $\alpha \mapsto \varphi(1_\alpha)$ is $F(\alpha) = \varphi_1^+(\alpha) = ([2][1][0]Next)^{1+\alpha}0$.

Then the fixed points of $F = \varphi_1^+$ are enumerated by $G = \varphi_2^+$. The least fixed point of F is $G(0) = \varphi_2^+(0) = Fix(\alpha \mapsto ([2][1][0]Next)^{1+\alpha}0)0 = [0]([2][1][0]Next)0$ (because of the absorption of "1+") and its $(1 + \alpha)$ -th fixed point is $G(\alpha) = \varphi_2^+(\alpha) = ([0]([2][1][0]Next))^{1+\alpha}0$.

Then the fixed points of $G = \varphi_2^+$ are enumerated by $H = \varphi_3^+$. The least fixed point of H is $H(0) = \varphi_3^+(0) = Fix(\alpha \mapsto ([0]([2][1][0]Next)^{1+\alpha}0)0 = [0]([0]([2][1][0]Next))0 = [0]^2([2][1][0]Next)0$ and its $(1 + \alpha)$ -th fixed point is $H(\alpha) = \varphi_3^+(\alpha) = ([0]^2([2][1][0]Next))^{1+\alpha}0$.

More generally, we have $\varphi_{1+\alpha}^+(0) = [0]^\alpha([2][1][0]Next)0$ and $\varphi_{1+\alpha}^+(\beta) = ([0]^\alpha([2][1][0]Next))^{1+\beta}0$.

Then we generalize the function φ^+ to any number of variables :

$$\varphi^+(\alpha, \beta) = \varphi_\alpha^+(\beta)$$

$\varphi^+(1, 0, 0)$ is the least fixed point of the function $\alpha \mapsto \varphi^+(\alpha, 0) = \alpha \mapsto [0]^\alpha([2][1][0]Next)0$. It is $Fix(\alpha \mapsto [0]^\alpha([2][1][0]Next)0)0 = [1][0]([2][1][0]Next)0$.

Compare with $\varphi(1, 0, 0) = [1][0]Next0$.

More generally, like we found $\varphi(\gamma, \beta, \alpha) = ([0]^\beta([1][0]^\gamma Next))^{1+\alpha}0$, we have $\varphi^+(\gamma, \beta, \alpha) = ([0]^\beta([1][0]^\gamma([2][1][0]Next)))^{1+\alpha}0$.

Like we generalized the φ function to transfinitely many variables reaching all ordinals less than $LVO = [2][1][0]Next\ 0$, we can generalize the φ^+ function to transfinitely many variables and reach all ordinals less than a new limit which we will call $LVO^+ = [2][1][0]([2][1][0]Next)0$ which is the least fixed point of $\alpha \mapsto [1]^\alpha[0]([2][1][0]Next)0$.

Then we can do the same with $\varphi^{++} = \Phi_2$ and we shall get similar results with $([2][1][0])^2Next$, and generally with Φ_α , getting formulas with $([2][1][0])^\alpha Next$.

The limit of $Next\ 0, [0]Next\ 0, [1][0]Next\ 0, [2][1][0]Next\ 0, [3][2][1][0]Next\ 0, \dots$

or $Next\ \omega, [0]Next\ \omega, [1][0]Next\ \omega, [2][1][0]Next\ \omega, [3][2][1][0]Next\ \omega, \dots$

is called the Bachmann-Howard ordinal (BHO).

It could be written $[\omega \dots 0]Next\ 0$ or $[\omega \dots 0]Next\ \omega$.

7 Rationalization of the Veblen functions

When we have defined the different notations, we have arbitrarily chosen some conventions, for example the limit of $\omega, \omega^\omega, \omega^{\omega^\omega}, \dots$ have been called ε_0 . We could have called it ε_1 . In this case, ε_α would have been the α -th fixed point of $\xi \mapsto \omega^\xi$ instead of the $(1 + \alpha)$ -th one. Also we chose to define $\varphi(0, \alpha) = \omega^\alpha$. We could have chosen to define $\varphi(0, \alpha) = \varepsilon_\alpha$. The "1+" which appear in the correspondence between Simmons and Veblen notations may be due to the fact that the choices that have been made are not the most logical.

We will define a rationalized variant of the Veblen notations which simplifies the correspondence with the Simmons notation :

- $\varepsilon_\alpha = \varphi(1, \alpha) = \varepsilon'_{1+\alpha} = \varphi'(0, 1 + \alpha)$
- $\zeta_\alpha = \varphi(2, \alpha) = \zeta'_{1+\alpha} = \varphi'(1, 1 + \alpha)$
- $\eta_\alpha = \varphi(3, \alpha) = \eta'_{1+\alpha} = \varphi'(2, 1 + \alpha)$
- Generally, $\varphi(1 + \beta, \alpha) = \varphi'(\beta, 1 + \alpha)$
- $\Gamma_0 = \varphi(1, 0, 0) = \varphi'(1, 0, 1)$
- Generally, if $\gamma \neq 0$, $\varphi(\gamma, \beta, \alpha) = \varphi'(\gamma, \beta, 1 + \alpha)$
- In a similar way, if $\gamma \neq 0$ or $\delta \neq 0$, $\varphi(\delta, \gamma, \beta, \alpha) = \varphi'(\delta, \gamma, \beta, 1 + \alpha)$ and so on.

With this notation, the correspondence with Simmons notation becomes simpler, for example we have :

- $\varepsilon'_\alpha = \text{Next}^\alpha 0$ instead of $\varepsilon_\alpha = \text{Next}^{1+\alpha} 0$
- $\varphi'(\beta, \alpha) = ([0]^\beta \text{Next})^\alpha 0$ instead of $\varphi(1 + \beta, \alpha) = ([0]^\beta \text{Next})^{1+\alpha} 0$
- $\varphi'(\gamma, \beta, \alpha) = ([0]^\beta ([1][0])^\gamma \text{Next})^\alpha 0$ instead of $\varphi(\gamma, \beta, \alpha) = ([0]^\beta ([1][0])^\gamma \text{Next})^{1+\alpha} 0$
- $\varphi'(\delta, \gamma, \beta, \alpha) = ([0]^\beta ([1][0])^\gamma ([1]^2[0])^\delta \text{Next})^{1+\alpha} 0$ instead of $\varphi(\delta, \gamma, \beta, \alpha) = ([0]^\beta ([1][0])^\gamma ([1]^2[0])^\delta \text{Next})^{1+\alpha} 0$

It appears that the last variable (α in the previous examples) plays a different role from the other variables, so it could be more logical to write for example $\varphi'_{\delta, \gamma, \beta}(\alpha)$ instead of $\varphi'(\delta, \gamma, \beta, \alpha)$ and to consider that β is at position 0, γ at position 1 and δ at position 2. In this case, we see that the position corresponds to the exponent of $[1]$ in the Simmons representation.

We can also use collapsing to represent the index list, writing for example :

- $\Gamma_0 = \varphi'_{1,0}(1) = \varphi'_\Omega(1)$
- Ackermann ordinal $= \varphi'_{1,0,0}(1) = \varphi'_{\Omega^2}(1)$
- $SVO = \varphi'_{\Omega^\omega}(1)$

This notation even permits writing ordinals that are out of range of Veblen notation like :

- $LVO = \varphi'_{\Omega^\Omega}(1)$
- $\varphi'_{\Omega^{\Omega^\Omega}}(1)$
- ...

8 RHS0 notation

8.1 Basic principles

Like Simmons notation, the RHS0 notation uses lambda calculus formalism.

The basic method consists in :

- Start from 0
- If we don't see any regularity, take the successor (add 1)
- If we see a regularity and we don't have a notation for it, invent it and jump to the limit
- If we see a regularity and we already have a notation for it, use it and jump to the limit.

The difficulty, which requires intelligence, is to see the regularities. It gives the following sequence :

- 0 : no regularity, take the successor
- $suc\ 0$: no regularity, take the successor
- $suc(suc\ 0)$: regularity : suc repeatedly applied to 0 . No notation, invent it : $H\ f\ x = \text{limit of } x, f\ x, f\ (f\ x), \dots$
- $Hsuc\ 0$: no regularity, take the successor
- $suc(Hsuc\ 0)$: no regularity, take the successor
- $suc(suc(Hsuc\ 0))$: regularity : suc repeatedly applied to $Hsuc\ 0$, notation exists
- $Hsuc(Hsuc\ 0)$: regularity : $Hsuc$ repeatedly applied to 0 , notation exists
- $H(Hsuc)0$: regularity : H repeatedly applied to suc , notation exists
- $HHsuc\ 0$: regularity ($suc\ 0, \dots, Hsuc\ 0, \dots, HHsuc\ 0, \dots, HHHsuc\ 0, \dots$), invent notation $R_1Hsuc\ 0$ for the limit of this sequence
- $R_1Hsuc\ 0$: no regularity, take the successor
- $suc(R_1Hsuc\ 0)$
- $suc(suc(R_1Hsuc\ 0))$
- $Hsuc(R_1Hsuc\ 0)$
- $suc(Hsuc(R_1Hsuc\ 0))$
- $suc(suc(Hsuc(R_1Hsuc\ 0)))$
- $Hsuc(Hsuc(R_1Hsuc\ 0))$
- $H(Hsuc)(R_1Hsuc\ 0)$
- $HHsuc(R_1Hsuc\ 0)$
- $R_1Hsuc(R_1Hsuc\ 0)$
- $H(R_1Hsuc)0$
- $suc(H(R_1Hsuc)0)$
- $suc(suc(H(R_1Hsuc)0))$
- $Hsuc(H(R_1Hsuc)0)$
- $suc(suc(Hsuc(H(R_1Hsuc)0)))$
- $Hsuc(Hsuc(H(R_1Hsuc)0))$
- $H(Hsuc)(H(R_1Hsuc)0)$
- $HHsuc(H(R_1Hsuc)0)$
- $R_1Hsuc(H(R_1Hsuc)0)$
- $suc(R_1Hsuc(H(R_1Hsuc)0))$
- $suc(suc(R_1Hsuc(H(R_1Hsuc)0)))$
- $Hsuc(R_1Hsuc(H(R_1Hsuc)0))$
- $suc(Hsuc(R_1Hsuc(H(R_1Hsuc)0)))$
- $suc(suc(Hsuc(R_1Hsuc(H(R_1Hsuc)0))))$
- $Hsuc(Hsuc(R_1Hsuc(H(R_1Hsuc)0)))$
- $H(Hsuc)(R_1Hsuc(H(R_1Hsuc)0))$
- $HHsuc(R_1Hsuc(H(R_1Hsuc)0))$
- $R_1Hsuc(R_1Hsuc(H(R_1Hsuc)0))$
- $H(R_1Hsuc)(H(R_1Hsuc)0)$
- $H(H(R_1Hsuc))0$
- $HH(R_1Hsuc)0$
- $R_1H(R_1Hsuc)0$
- $H(R_1H)suc\ 0$
- \dots
- $R_1H(R_1H)suc\ 0$
- $R_1(R_1H)suc\ 0$
- $HR_1Hsuc\ 0$
- \dots
- $R_1HR_1Hsuc\ 0$: invent notation $R_2R_1Hsuc\ 0 = \text{limit of } suc\ 0, R_1Hsuc\ 0, R_1HR_1Hsuc\ 0, \dots$
- \dots
- $R_3R_2R_1Hsuc\ 0$: invent notation $R_{3\dots 1}Hsuc\ 0$ and jump to limit
- $R_{\omega\dots 1}Hsuc\ 0$

- ...
- $R_2 R_{\omega+1} Hsuc\ 0$: invent notation $R_{\omega+1} Hsuc\ 0$
- ...

To progress faster, we can use the following rule :

If we have found an ordinal α , and later another ordinal β of the form $f(s(sz))$, we may produce an ordinal $\gamma = f([suc \rightarrow s, 0 \rightarrow z]\alpha)$ where $[suc \rightarrow s, 0 \rightarrow z]\alpha$ means the expression obtained by replacing suc by s and 0 by z in α .

For example :

- $\alpha = R_1 Hsuc\ 0$
- $\beta = R_1 H(R_1 Hsuc)0$
- $s = R_1 H$
- $z = suc$
- $fx = x\ 0$
- $[suc \rightarrow R_1 H, 0 \rightarrow suc]\alpha = R_1 H(R_1 H)suc$
- $\gamma = f([suc \rightarrow R_1 H, 0 \rightarrow suc]\alpha) = R_1 H(R_1 H)suc\ 0$

With the following rules :

- $0 : \rightarrow 0$
- $suc : x \rightarrow suc\ x$
- $H : f(fx) - > Hfx$
- $R_1 : ff - > R_1 f$
- $R_2 : fgfg - > R_2 fg$
- $R_3 : fghfgh - > R_3 fgh$
- ...
- $Repl : \alpha, f(s(sz)) \rightarrow f([suc \rightarrow s, 0 \rightarrow z]\alpha)$

we can produce the following sequence of ordinals :

- $0 : 0 : 0$
- $1 : suc\ 0 : suc\ 0$
- $2 : suc\ 1 : suc\ (suc\ 0)$
- $3 : H2 : Hsuc\ 0$
- $4 : suc\ 3 : suc\ (Hsuc\ 0)$
- $5 : suc\ 4 : suc\ (suc\ (Hsuc\ 0))$
- $6 : H5 : Hsuc\ (Hsuc\ 0)$
- $7 : H6 : H(Hsuc)0$
- $8 : H7 : HHsuc\ 0$
- $9 : R_1 8 : R_1 Hsuc\ 0$
- $10 : suc\ 9 : suc\ (R_1 Hsuc\ 0)$
- $11 : suc\ 10 : suc\ (suc\ (R_1 Hsuc\ 0))$
- $12 : Repl\ 9\ 11[suc - > suc, 0 - > R_1 Hsuc\ 0] : R_1 Hsuc\ (R_1 Hsuc\ 0)$
- $13 : Repl\ 9\ 12[suc - > R_1 Hsuc, 0 - > 0] : R_1 H(R_1 Hsuc)0$
- $14 : Repl\ 9\ 13[suc - > R_1 H, 0 - > suc] : R_1 H(R_1 H)suc\ 0$
- $15 : R_1 14 : R_1 (R_1 H)suc\ 0$
- $16 : Repl\ 9\ 15[suc - > R_1, 0 - > H] : R_1 H R_1 Hsuc\ 0$
- $17 : R_2 16 : R_2 R_1 Hsuc\ 0$

The rules R_1, R_2, R_3, \dots may be replaced by H or $Repl$ if $f_1 \dots f_n \dots f_1 \dots f_n$ is reformulated in $\langle f_1, \dots, f_n \rangle (\dots (\langle f_1, \dots, f_n \rangle I) \dots)$ with $\langle f_1, \dots, f_n \rangle g = g\ f_1 \dots f_n$:

- $0 : 0 : 0$
- $1 : suc\ 0 : suc\ 0$
- $2 : suc\ 1 : suc\ (suc\ 0)$

- 3 : H 2 : H suc 0
- 4 : suc 3 : suc (H suc 0)
- 5 : suc 4 : suc (suc (H suc 0))
- 6 : Repl 3 5 [suc->suc,0->H suc 0] : H suc (H suc 0)
- 7 : Repl 3 6 [suc->H suc,0->0] : H (H suc) 0
- 8 : Repl 3 7 [suc->H,0->suc] : H H suc 0 = <H> (<H> I) suc 0
- 9 : Repl 3 8 [suc-><H>,0->I] : H <H> I suc 0
- 10 : suc 9 : suc (H <H> I suc 0)
- 11 : suc 10 : suc (suc (H <H> I suc 0))
- 12 : Repl 9 10 [suc->suc,0->H <H> I suc 0] : H <H> I suc (H <H> I suc 0)
- 13 : Repl 9 12 [suc->H <H>: I suc,0->0] : H <H> I (H <H> I suc) 0
- 14 : Repl 9 13 [suc->H <H> I,0->suc] : H <H> I (H <H> I) suc 0 = <H <H> I> (<H <H> I> I) suc 0
- 15 : Repl 3 14 [suc-><H <H> I>,0->I] : H <H <H> I> I suc 0 = [H <*> I] ([H <*> I] H) suc 0
- 16 : Repl 9 15 [suc->[H <*> I],0->H] : H <H> I [H <*> I] H suc 0
- = [H <*> I] H [H <*> I] H suc 0 = <[H <*> I],H> (<[H <*> I], H> I) suc 0
- 17 : Repl 3 16 [suc-><[H <*> I],H>,0->I] : H <[H <*> I], H> I suc 0

More formally, the RHS0 notation uses lambda calculus with De Bruijn indexes. $\lambda.x$ is written $[x]$ and variables are written $*$, $**$, $***$, ..., or $\bullet, \bullet\bullet, \bullet\bullet\bullet, \dots$ for example $[\dots * \dots] = [\dots \bullet \dots] = \lambda x(\dots x \dots)$

CI = C I is defined by CI $x f = f x$.
 CI $x = \langle x \rangle$
 $\langle x_1, \dots, x_n \rangle f = f x_1 \dots x_n$
 $\text{tuple } n f x_1 \dots x_n = f \langle x_1, \dots, x_n \rangle$
 $\text{tuple } 0 = \langle I \rangle$
 $\text{tuple } (n+1) f x = \text{tuple } n [f (\text{insert } x *)]$
 with $\text{insert } x a f = a (f x)$

$r \ 0 f x = x$
 $r \ (n+1) f x = f (r \ n f x)$
 $r \ (\text{lim } g) f x = \text{lim } [r * f x]$

H $f x$ represents the limit of $x, f x, f (f x), \dots$
 $H f x = r \ w f x$

$R_1 = [H \langle \bullet \rangle I] = \text{tuple } 1 [H \bullet I]$
 $R_2 = [[H \langle \bullet\bullet, \bullet \rangle I]] = \text{tuple } 2 [H \bullet\bullet I]$
 $R_3 = [[[H \langle \bullet\bullet\bullet, \bullet\bullet, \bullet \rangle I]]] = \text{tuple } 3 [H \bullet\bullet\bullet I]$
 $R_n = \text{tuple } n [H \bullet^n I]$
 $R_{n\dots 1} = R_n \dots R_1$
 $S_{n\dots 1} = [S_{\bullet\dots 1}]_n = \langle R_n, \dots, R_1 \rangle$
 $R_{n\dots 1} = S_{n\dots 1} I$
 $[S_{\bullet\dots 1}]_0 = I$
 $[S_{\bullet\dots 1}](n+1) = \text{insert}(\text{tuple}(n+1)[H \bullet^n I])([S_{\bullet\dots 1}]_n)$

$L f = \text{lim } f \ 0, f \ 1, \dots$
 $L f x = L [f * x]$
 $H = [[L [r * *** **]]]$
 or
 $L0 = \text{lim } f \ 0, f \ 1, \dots$
 $L \ n f = \text{tuple } n [L0 [** (f *)]]$
 $L \ n = [\text{tuple } n [L0 [** (***) *]]]$
 $L = [[\text{tuple } ** [L0 [** (***) *]]]]$

= \n \f (tuple n \a (L0 \i (a (f i))))

To represent the replacement $[suc \rightarrow s, 0 \rightarrow z]$ we can represent ordinals by ordinal functions which, when applied to suc and 0 , give the considered ordinal. For example, $R_1Hsuc\ 0$ is represented by the ordinal function $s \mapsto z \mapsto R_1Hsz, R_1H(R_1Hsuc)0$ by $s \mapsto z \mapsto R_1H(R_1Hs)z$. From these ordinals, with the replacement $[suc \rightarrow R_1H, 0 \rightarrow suc]$ we can produce a new ordinal represented by $s \mapsto z \mapsto ((s \mapsto z \mapsto R_1Hsz)(R_1H)sz) = s \mapsto z \mapsto R_1H(R_1H)sz$ which, when applied to suc and 0 , gives $R_1H(R_1H)suc\ 0$.

Operations can be represented with replacements :

- $\alpha + \beta = [0 \rightarrow \alpha]\beta$
- $\alpha \cdot \beta = [suc \rightarrow [\bullet + \alpha]]\beta = [suc \rightarrow > [[0 \rightarrow \bullet\bullet]\alpha]]\beta$
- $\alpha^\beta = [suc \rightarrow [\bullet \cdot \alpha], 0 \rightarrow 1]\beta = [suc \rightarrow [[suc \rightarrow [[0 \rightarrow \bullet\bullet] \bullet \bullet \bullet]]\alpha], 0 \rightarrow suc\ 0]\beta$
- $\omega^\alpha = [suc \rightarrow [suc \rightarrow Hsuc], 0 \rightarrow suc\ 0]\alpha = [suc \rightarrow H, 0 \rightarrow suc]\alpha\ 0$
- $\varepsilon_0^\alpha = [suc \rightarrow R_1H, 0 \rightarrow suc]\alpha\ 0$
- $\varepsilon_\alpha = [suc \rightarrow R_1, 0 \rightarrow H](1 + \alpha)suc\ 0; 1 + \alpha = [0 \rightarrow suc\ 0]\alpha$

8.2 Correspondence with other notations

- $suc\ 0 = 0 + 1 = 1$
- $suc\ (suc\ 0) = 1 + 1 = 2$
- $Hsuc\ 0 = \omega$
- $suc\ (Hsuc\ 0) = \omega + 1$
- $Hsuc\ (Hsuc\ 0) = \omega + \omega = \omega \cdot 2$
- $H(Hsuc)\ 0 = \omega \cdot \omega = \omega^2$
- $HHsuc\ 0 = \omega^\omega$
- $R_1Hsuc\ 0 = \text{limit of } suc\ 0, Hsuc\ 0, HHsuc\ 0, HHHsuc\ 0, \dots = \varepsilon_0 = \varphi(1, 0) = \varphi'(0, 1) = \text{Next } \omega$
- $suc(R_1Hsuc\ 0) = \varepsilon_0 + 1$
- $R_1Hsuc(R_1Hsuc\ 0) = \varepsilon_0 + \varepsilon_0 = \varepsilon_0 \cdot 2$
- $R_1H(R_1Hsuc)0 = \varepsilon_0 \cdot \varepsilon_0 = \varepsilon_0^2$
- $R_1H(R_1H)suc\ 0 = \varepsilon_0^{\varepsilon_0}$
- $R_1H(R_1H)(R_1H)suc\ 0 = \varepsilon_0^{\varepsilon_0^{\varepsilon_0}}$
- $R_1(R_1H)suc\ 0 = \varepsilon_1 = \varphi(1, 1) = \varphi'(0, 2) = \text{Next}(\text{Next } \omega)$ (note again that the correspondence is clearer with the rationalized function φ'

We have previously seen that ε_1 is the limit of $\varepsilon_0, \varepsilon_0^{\varepsilon_0}, \varepsilon_0^{\varepsilon_0^{\varepsilon_0}}, \dots$ and is also the limit of $\varepsilon_0 + 1, \omega^{\varepsilon_0+1}, \omega^{\omega^{\varepsilon_0+1}}, \dots$ and we have proved the equivalence of these two fundamental sequences. We have seen that the first fundamental sequence is equivalent to $\omega, \varepsilon_0^\omega, \varepsilon_0^{\varepsilon_0^\omega}, \dots$, so we proved the equivalence of the two fundamental sequences by proving that for any n , we have :

$$\begin{array}{c} \omega \\ \vdots \\ \omega^{\omega^{\varepsilon_0+1}} \end{array} = \begin{array}{c} \varepsilon_0^\omega \\ \vdots \\ \varepsilon_0^{\varepsilon_0^\omega} \end{array}$$

We will now see how we can prove it using RHS0 notation.

First we will write the two sides of this equality using RHS0 notation :

We will use the notation $X \dots X$ for X repeated n times.

- $\varepsilon_0 = R_1Hsuc\ 0$
- $\varepsilon_0 + 1 = suc(R_1Hsuc\ 0)$
- $\omega^{\varepsilon_0+1} = [suc \rightarrow H, 0 \rightarrow suc](suc(R_1Hsuc)0)0 = H(R_1HHsuc)0 = H(R_1Hsuc)0$
- $\omega^{\omega^{\varepsilon_0+1}} = H(R_1HH)suc\ 0 = H(R_1H)suc\ 0$
- $\omega^{\omega^{\omega^{\varepsilon_0+1}}} = H(R_1H)Hsuc\ 0$
- $\omega^{\omega^{\omega^{\omega^{\varepsilon_0+1}}}} = H(R_1H)HHsuc\ 0$
- \dots
- $\omega^{\omega^{\omega^{\omega^{\omega^{\varepsilon_0+1}}}}}} = H(R_1H)H \dots Hsuc\ 0$
- $\omega = Hsuc\ 0$

- $\varepsilon_0^\omega = [suc \rightarrow R_1H, 0 \rightarrow suc]\omega 0 = H(R_1H)suc\ 0$
- $\varepsilon_0^\omega = H(R_1H)(R_1H)suc\ 0$
- $\varepsilon_0^{\varepsilon_0^\omega} = H(R_1H)(R_1H)(R_1H)suc\ 0$
- \dots
- $\varepsilon_0^{\varepsilon_0^\omega} = H(R_1H) \dots (R_1H)(R_1H)suc\ 0$

We will now prove the equality $H(R_1H)H \dots Hsuc\ 0 = H(R_1H) \dots (R_1H)(R_1H)suc\ 0$ for any n by induction.

For $n = 0$, the equality is trivial : $H(R_1H)suc\ 0 = H(R_1H)suc\ 0$.

We will now suppose $H(R_1H)H \dots Hsuc\ 0 = H(R_1H) \dots (R_1H)(R_1H)suc\ 0$ for a given n and prove it for $n + 1$:

By elevating ω at the power of each side of this equality, we get :
 $[suc \rightarrow H, 0 \rightarrow suc](H(R_1H)H \dots Hsuc\ 0)0 = [suc \rightarrow H, 0 \rightarrow suc](H(R_1H) \dots (R_1H)(R_1H)suc\ 0)0$
(1) $H(R_1H)H \dots HHsuc\ 0 = H(R_1H) \dots (R_1H)(R_1H)suc\ 0$

We also have :

$$H(R_1H) \dots (R_1H)suc\ 0 = H(R_1H) \dots (R_1H)suc(suc\ 0)$$

which corresponds to the RHS0 notation for :

$$\varepsilon_0^\omega = 1 + \varepsilon_0^\omega$$

by absorpsion of "1+" and "suc".

Now we elevate ε_0 to the power of each side of this equality, which gives :

$$[suc \rightarrow R_1H, 0 \rightarrow suc](H(R_1H) \dots (R_1H)suc\ 0)0 = [suc \rightarrow R_1H, 0 \rightarrow suc](H(R_1H) \dots (R_1H)suc(suc\ 0))0$$

$$H(R_1H) \dots (R_1H)(R_1H)suc\ 0 = H(R_1H) \dots (R_1H)(R_1H)(R_1H)suc\ 0$$

Then we elevate ω to the power of each side of this equality :

$$[suc \rightarrow H, 0 \rightarrow suc](H(R_1H) \dots (R_1H)(R_1H)suc\ 0)0 = [suc \rightarrow H, 0 \rightarrow suc](H(R_1H) \dots (R_1H)(R_1H)(R_1H)suc\ 0)0$$

$$H(R_1H) \dots (R_1H)(R_1H)suc\ 0 = H(R_1H) \dots (R_1H)(R_1H)(R_1H)suc\ 0$$

which can be simplified to :

$$H(R_1H) \dots (R_1H)(R_1H)suc\ 0 = H(R_1H) \dots (R_1H)(R_1H)(R_1H)suc\ 0$$

Noting that the left side of this last equality is the same as the right side of (1), we get by transitivity of equality, equating the left side of (1) with the right side of the last equality :

$$H(R_1H)H \dots HHsuc\ 0 = H(R_1H) \dots (R_1H)(R_1H)(R_1H)suc\ 0$$

which corresponds to the equality we wanted to prove for $n+1$.

Then the correspondence continues with :

- $R_1(R_1(R_1H))suc\ 0 = \varepsilon_2 = \varphi(1, 2) = \varphi'(0, 3) = Next(Next(Next\ \omega))$
- $HR_1Hsuc\ 0 = \varepsilon_\omega = \varphi(1, \omega) = \varphi'(0, \omega) = Next^\omega\omega$
- $R_1HR_1Hsuc\ 0 = \varepsilon_{\varepsilon_0}$
- $R_1HR_1HR_1Hsuc\ 0 = \varepsilon_{\varepsilon_{\varepsilon_0}}$
- $R_2R_1Hsuc\ 0 = \zeta_0 = \varphi(2, 0) = \varphi'(1, 1) = [0]Next\ \omega$

The next step is ζ_1 which is the next fixed point of the function $\alpha \mapsto \varepsilon_\alpha$, the limit of $\zeta_0 + 1, \varepsilon_{\zeta_0+1}, \varepsilon_{\varepsilon_{\zeta_0+1}}, \dots, \varepsilon_\alpha$ is $[suc \rightarrow R_1, 0 \rightarrow H](1 + \alpha)suc\ 0$, or $[suc \rightarrow R_1, 0 \rightarrow H]\alpha\ suc\ 0$ if $\alpha \geq \omega$ by absorpsion of "1+". This is the result of replacing suc by R_1 and 0 by H in α and applying the result to suc and 0. So by iterating this transformation we get that ζ_1 is the limit of :

- $\zeta_0 + 1 = suc(R_2R_1Hsuc\ 0)$
- $R_1(R_2R_1HR_1H)suc\ 0 = R_1(R_2R_1H)suc\ 0$
- $R_1(R_2R_1H)R_1Hsuc\ 0$
- $R_1(R_2R_1H)R_1HR_1Hsuc\ 0$
- \dots

In the previous correnspondence formulas, we can see a correspondence between RHS0 and Simmons notations :

- $R_2 \leftrightarrow [0]$
- $R_1 \leftrightarrow Next$
- $H \leftrightarrow \omega$
- $suc\ 0$ at the end of the RHS0 notation

If we apply this correspondence to $\zeta_1 = [0]Next([0]Next\ \omega)$ (see "Simmons notation / Correspondence with Veblen functions") we get $\zeta_1 = R_2R_1(R_2R_1H)suc\ 0$.

This is the limit of :

- $R_1(R_2R_1H)suc\ 0$
- $R_1(R_2R_1H)R_1(R_2R_1H)suc\ 0$
- $R_1(R_2R_1H)R_1(R_2R_1H)R_1(R_2R_1H)suc\ 0$
- ...

Compare with what we found previously :

- $\zeta_0 + 1 = suc(R_2R_1Hsuc\ 0)$
- $R_1(R_2R_1HR_1H)suc\ 0 = R_1(R_2R_1H)suc\ 0$
- $R_1(R_2R_1H)R_1Hsuc\ 0$
- $R_1(R_2R_1H)R_1HR_1Hsuc\ 0$
- ...

and with the previously proven equality :

$$H(R_1H)H \dots Hsuc\ 0 = H(R_1H) \dots (R_1H)(R_1H)suc\ 0$$

which could also be written :

$$H(R_1H)H \dots Hsuc\ 0 = H(R_1H)(R_1H) \dots (R_1H)suc\ 0$$

There is probably a similar equality :

$$R_1(R_2R_2H)R_1H \dots R_1Hsuc\ 0 = R_1(R_2R_1H)R_1(R_2R_1H) \dots R_1(R_2R_1H)suc\ 0$$

which would prove the equivalence of the two fundamental sequences.

So we can go on with our correspondences :

- $R_2R_1(R_2R_1H)suc\ 0 = \zeta_1 = \varphi(2, 1) = \varphi'(1, 2) = [0]Next([0]Next\ \omega)$
- $H(R_2R_1)Hsuc\ 0 = \zeta_\omega$
- $R_2R_1H(R_2R_1)Hsuc\ 0 = \zeta_{\zeta_0}$
- $R_2(R_2R_1)Hsuc\ 0 = \eta_0 = \varphi(3, 0) = \varphi'(2, 1) = [0]([0]Next)\omega$
- $HR_2R_1Hsuc\ 0 = \varphi(\omega, 0) = \varphi'(\omega, 1)$
- $R_1HR_2R_1Hsuc\ 0 = \varphi(\varepsilon_0, 0) = \varphi(\varphi(1, 0), 0) = \varphi'(\varepsilon_0, 1) = \varphi'(\varphi'(0, 1), 1)$
- $R_2R_1HR_2R_1Hsuc\ 0 = \varphi(\zeta_0, 0) = \varphi(\varphi(2, 0), 0) = \varphi'(\zeta_0, 1) = \varphi'(\varphi'(1, 1), 1)$
- $R_3R_2R_1Hsuc\ 0 = \Gamma_0 = \varphi(1, 0, 0) = \varphi'(1, 0, 1) = [1][0]Next\ \omega$

We may then extend our correspondence rule :

- $R_3 \leftrightarrow [1]$
- $R_2 \leftrightarrow [0]$
- $R_1 \leftrightarrow Next$
- $H \leftrightarrow \omega$
- $suc\ 0$ at the end of the RHS0 notation

It is likely that this correspondence can be generalized in a simple and logical way, and it seems to me that the simpler generalization is :

- $R_{n+2} \leftrightarrow [n]$
- $R_1 \leftrightarrow Next$
- $H \leftrightarrow \omega$
- $suc\ 0$ at the end of the RHS0 notation

I will call it the "Simmons - RHS0 correspondence conjecture".

Then, if the correspondence conjecture is true, the correspondence goes on with :

- $R_3 R_2 R_1 Hsuc\ 0 = \Gamma_0 = \varphi(1, 0, 0) = \varphi'(1, 0, 1) = [1][0]Next\ \omega$
- $R_3(R_3 R_2)R_1 Hsuc\ 0 = \varphi(1, 0, 0, 0) = \varphi'(1, 0, 0, 1) = [1]([1][0])Next\ 0$ (Note that in the φ and φ' functions, the last variable plays a different role than the others, as mentioned previously, so the most logical representation should probably be $\varphi'_{1,0,0}(1)$ where the first 1 should be considered at position 2 and not 3, in this case its position corresponds to the number of occurrences (or the exponent) of R_3 and $[1]$)
- $HR_3 R_2 R_1 Hsuc\ 0 = SVO = [1]^\omega[0]Next\ \omega$
- $R_4 R_3 R_2 R_1 Hsuc\ 0 = LVO = [2][1][0]Next\ \omega$
- $R_{\omega \dots 1} Hsuc\ 0 = BHO$
- ...

Note the importance of using logical notations to make correct conjectures : if, instead of φ' , we use the less logical function φ , we have the correspondence :

- $R_1 Hsuc\ 0 = \varepsilon_0 = \varphi(1, 0)$
- $R_2 R_1 Hsuc\ 0 = \zeta_0 = \varphi(2, 0)$

and we could think that it continues with :

- $R_3 R_2 R_1 Hsuc\ 0 = \eta_0 = \varphi(3, 0)$
- $R_{\omega \dots 1} Hsuc\ 0 = \varphi(\omega, 0)$

Like with the Veblen functions, we can use collapsing with RHS0 notation, writing for example :

$$\Gamma_0 = \varphi_{1,0}(0) = \varphi'_{1,0}(1) = \varphi'_\Omega(0) = \varphi'_\Omega(1) = [0]^\Omega Next\ \omega = [1][0]Next\ \omega = (R_2)^\Omega R_1 Hsuc\ 0 = R_3 R_2 R_1 Hsuc\ 0$$

which gives $(R_2)^\Omega = R_3 R_2$

to be compared with $[0]^\Omega = [1][0]$.

Γ_0 is also the limit of the following sequence :

- $\zeta_0 = \varphi'_1(1) = [0]Next\ \omega = R_2 R_1 Hsuc\ 0$
- $\varphi'_{\zeta_0}(1) = \varphi'_{\varphi'_1(1)}(1) = [0]^{[0]Next\ \omega} Next\ \omega = (R_2)^{R_2 R_1 Hsuc\ 0} R_1 Hsuc\ 0 = R_2 R_1 H R_2 R_1 Hsuc\ 0$
- $\varphi'_{\varphi'_{\varphi'_1(1)}(1)}(1) = [0]^{[0]^{[0]Next\ \omega} Next\ \omega} Next\ \omega = R_2 R_1 H R_2 R_1 H R_2 R_1 Hsuc\ 0$

This limit is $R_3 R_2 R_1 Hsuc\ 0$.

8.3 Going further with RHS0 notation and collapsing

The Bachmann-Howard ordinal (BHO) is the limit of $R_1 Hsuc\ 0, R_2 R_1 Hsuc\ 0, R_3 R_2 R_1 Hsuc\ 0, \dots$ which we will write $R_{\omega \dots 1} Hsuc\ 0$. We can go on ascending ordinals after BHO :

- $BHO = R_{\omega \dots 1} Hsuc\ 0$
- $suc(R_{\omega \dots 1} Hsuc\ 0)$
- $R_{\omega \dots 1} Hsuc\ (R_{\omega \dots 1} Hsuc\ 0)$
- $R_{\omega \dots 1} H(R_{\omega \dots 1} Hsuc\ 0)$
- $R_{\omega \dots 1} H(R_{\omega \dots 1} H)suc\ 0$
- $R_1(R_{\omega \dots 1} H)suc\ 0$
- $R_{\omega \dots 1}(R_{\omega \dots 1} H)suc\ 0$
- $HR_{\omega \dots 1} Hsuc\ 0$
- $R_2 R_{\omega \dots 1} Hsuc\ 0$
- $R_3 R_2 R_{\omega \dots 1} Hsuc\ 0$
- $R_{\omega \dots 2} R_{\omega \dots 1} Hsuc\ 0$ which we will write $R_{\omega \cdot 2 \dots 1} Hsuc\ 0$
- $R_{\omega \dots 3} R_{\omega \dots 2} R_{\omega \dots 1} Hsuc\ 0$ which we will write $R_{\omega \cdot 3 \dots 1} Hsuc\ 0$
- $R_{\omega^2 \dots 1} Hsuc\ 0$
- $R_{\varepsilon_0 \dots 1} Hsuc\ 0 = R_{R_1 Hsuc\ 0 \dots 1} Hsuc\ 0$

Then we can take the least fixed point of the function $\alpha \mapsto R_{\alpha \dots 1} Hsuc\ 0$ which we can also write $[R_{\bullet \dots 1} Hsuc\ 0]$. This fixed point is $H[R_{\bullet \dots 1} Hsuc\ 0]0$ which we may also write $R_1^1 Hsuc\ 0$ if we define $R_1^1 x_1 x_2 x_3 = H[R_{\bullet \dots 1} x_1 x_2 x_3]0$. Then the ascension goes on with :

- $H[R_{\bullet \dots 1} Hsuc\ 0]0 = R_1^1 Hsuc\ 0$
- $R_2 R_1^1 Hsuc\ 0$
- $R_3 R_2 R_1^1 Hsuc\ 0$
- $R_{\omega \dots 2} R_1^1 Hsuc\ 0$
- $H[R_{\bullet \dots 2} R_1^1 Hsuc\ 0]0 = R_2^1 R_1^1 Hsuc\ 0 = R_{2 \dots 1}^1 Hsuc\ 0$ with $R_2^1 x_1 x_2 x_3 x_4 = H[R_{\bullet \dots 2} x_1 x_2 x_3 x_4]0$
- $R_3^1 R_2^1 R_1^1 Hsuc\ 0 = R_{3 \dots 1}^1 Hsuc\ 0$
- $R_{\omega \dots 1}^1 Hsuc\ 0$
- $H[R_{\bullet \dots 1}^1 Hsuc\ 0]0 = R_1^2 Hsuc\ 0$ with $R_1^2 x_1 x_2 x_3 = H[R_{\bullet \dots 1}^1 x_1 x_2 x_3]0$
- $R_1^3 Hsuc\ 0$
- $R_1^\omega Hsuc\ 0$
- $H[R_1^\bullet Hsuc\ 0]0 = R_1^{1,0} Hsuc\ 0$ with $R_1^{1,0} x_1 x_2 x_3 = H[R_1^\bullet x_1 x_2 x_3]0$
- $R_1^{1,0,0} Hsuc\ 0$

We can number the positions in the list of upper indices of R or introduce collapsing to write

- $R_1^1 = R_1^{1_0} = R_1^{0:1}$
- $R_1^{1,0} = R_1^{1_1} = R_1^{1:1} = R_1^\Omega$
- $R_1^{1,0,0} = R_1^{1_2} = R_1^{2:1} = R_1^{\Omega^2}$
- ...

We also need a notation for uncountable ordinals. We can take $\Omega = \omega_1$ the least uncountable ordinal and use a notation similar to the one we used for countable ordinals, replacing H by H_1 when ω is replaced by $\Omega = \omega_1$, writing for example :

- $\Omega = \omega_1 = H_1 suc\ 0$
- $\Omega^2 = H_1(H_1 suc)0$
- $\Omega^\omega = HH_1 suc\ 0$
- $\Omega^\Omega = H_1 H_1 suc\ 0$
- $\Omega^{\Omega^\Omega} = H_1 H_1 H_1 suc\ 0$
- ...

Then we can go on ascending ordinals by using greater and greater uncountable ordinals as upper indices of R, for example :
 $R_1^{H[R_1^\bullet H_1 suc\ 0]0} Hsuc\ 0$

9 Extending Simmons notation

The limit of the Simmons notation is the limit of :

- $Next\ \omega = \varepsilon_0$
- $[0]Next\ \omega = \zeta_0$
- $[1][0]Next\ \omega = \Gamma_0$
- $[2][1][0]Next\ \omega = LVO$
- $[3][2][1][0]Next\ \omega$
- ...

which is BHO, the Bachmann-Howard ordinal.

Using RHS0 notation, it corresponds to :

- $R_1 Hsuc\ 0 = \varepsilon_0$
- $R_2 R_1 Hsuc\ 0 = \zeta_0$
- $R_3 R_2 R_1 Hsuc\ 0 = \Gamma_0$
- $R_4 R_3 R_2 R_1 Hsuc\ 0 = LVO$

- $R_5 R_4 R_3 R_2 R_1 Hsuc\ 0$
- ...

which can be written $R_{\omega \dots 1} Hsuc\ 0$ in RHS0 notation.

And we just saw that the RHS0 notation goes much further.

So, using the correspondence, we can extend the Simmons notation in a similar way the RHS0 extends beyond BHO.

Using similar notations, we can write $[\omega \dots 0]Next\omega$ for the BHO.

Then we can go on :

- $R_{\omega \dots 1} Hsuc\ 0 = R_{\omega \dots 2} R_1 Hsuc\ 0 = [\omega \dots 0]Next\omega$
- $R_2 R_{\omega \dots 1} Hsuc\ 0 = R_2 (R_{\omega \dots 2} R_1) Hsuc\ 0 = [0]([\omega \dots 0]Next)\omega$
- $R_3 R_2 R_{\omega \dots 1} Hsuc\ 0 = R_3 R_2 (R_{\omega \dots 2} R_1) Hsuc\ 0 = [1][0]([\omega \dots 0]Next)\omega$
- $R_{\omega \cdot 2 \dots 1} Hsuc\ 0 = R_{\omega \dots 2} R_{\omega \dots 1} Hsuc\ 0 = R_{\omega \dots 2} (R_{\omega \dots 2} R_1) Hsuc\ 0 = [\omega \dots 0]([\omega \dots 0]Next)\omega$
- $HR_{\omega \dots 2} R_1 Hsuc\ 0 = [\omega \dots 0]^\omega Next\ \omega$
- $R_1 HR_{\omega \dots 2} R_1 Hsuc\ 0 = [\omega \dots 0]^{\varepsilon_0} Next\ \omega$
- $R_3 R_{\omega \dots 2} R_1 Hsuc\ 0 = Fix(\alpha \mapsto [\omega \dots 0]^\alpha Next\ \omega)\omega = [1][\omega \dots 0]Next\ \omega = [\omega + 1 \dots 0]Next\ \omega$
- $R_{\omega \cdot 2 + 1 \dots 1} Hsuc\ 0$
 $= R_3 R_{\omega \dots 2} R_{\omega \dots 1} Hsuc\ 0$
 $= R_3 R_{\omega \dots 2} (R_{\omega \dots 2} R_1) Hsuc\ 0$
 $= [1][\omega \dots 0]([\omega \dots 0]Next)\omega$
 $= Fix(\alpha \mapsto [\omega \dots 0]^\alpha ([\omega \dots 0]Next)\omega)\omega$
 $= Fix(\alpha \mapsto [\omega \dots 0]^\alpha Next\omega)\omega$ (absorbsion of $[\omega \dots 0]$)
 $= [1][\omega \dots 0]Next\ \omega$
 $= [\omega + 1 \dots 0]Next\ \omega$
- ...

10 Ordinal collapsing functions

Remember that collapsing consists in using uncountable ordinals to define countable ordinals.

We already saw how collapsing can be used to define ordinals, for example :

- $\varphi(\alpha, 0, \beta, \gamma) = \varphi(\gamma_0, \beta_1, \alpha_3) = (\xi \mapsto \omega^\xi) \begin{pmatrix} \gamma & \beta & \alpha \\ 0 & 1 & 3 \end{pmatrix} = \varphi(\Omega^3 \cdot \alpha + \Omega \cdot \beta + \gamma).$
- $\varphi_{\alpha, \beta, \gamma}(\delta) = \varphi_{\Omega^2 \cdot \alpha + \Omega \cdot \beta + \gamma}(\delta)$
- $\varphi(\gamma, \beta, \alpha) = \varphi_{\gamma, \beta}(\alpha) = \varphi_{\Omega \cdot \gamma + \beta}(\alpha) = \varphi(\Omega \cdot \gamma + \beta, \alpha) = \varphi(1 + \Omega \cdot \gamma + \beta, \alpha) = ([0]^{\Omega \cdot \gamma + \beta} Next)^{1+\alpha} 0 = ([0]^\beta ([0]^\Omega)^\gamma Next)^{1+\alpha} 0 = ([0]^\beta ([1][0])^\gamma Next)^{1+\alpha} 0$ with $[0]^\Omega = [1][0]$
- $\varphi(\delta, \gamma, \beta, \alpha) = \varphi_{\delta, \gamma, \beta}(\alpha) = \varphi_{\Omega^2 \cdot \delta + \Omega \cdot \gamma + \beta}(\alpha) = \varphi(\Omega^2 \cdot \delta + \Omega \cdot \gamma + \beta, \alpha) = \varphi(1 + \Omega^2 \cdot \delta + \Omega \cdot \gamma + \beta, \alpha) = ([0]^{\Omega^2 \cdot \delta + \Omega \cdot \gamma + \beta} Next)^{1+\alpha} 0 = ([0]^\beta ([0]^\Omega)^\gamma ([0]^\Omega)^\delta Next)^{1+\alpha} 0 = ([0]^\beta ([1][0])^\gamma ([1]^2[0])^\delta Next)^{1+\alpha} 0$ with $[0]^\Omega = [1][0]$ and $[0]^{\Omega^2} = ([0]^\Omega)^\Omega = ([1][0])^\Omega = [1]([1][0]) = [1]^2[0]$
- $\varphi(1_\Omega) = [1]^\Omega [0] Next\ 0.$
- $LVO = \varphi_{\Omega^\Omega}(0) = [0]^{\Omega^\Omega} Next\ 0 = [1]^\Omega [0] Next\ 0 = [2][1][0] Next\ 0$ with $[1]^\Omega = [2][1]$
- $\Gamma_0 = \varphi'_{1,0}(1) = \varphi'_\Omega(1)$
- Ackermann ordinal $= \varphi'_{1,0,0}(1) = \varphi'_{\Omega^2}(1)$
- $SVO = \varphi'_{\Omega^\omega}(1)$
- $LVO = \varphi'_{\Omega^\Omega}(1)$
- $\varphi'_{\Omega^{\Omega^\Omega}}(1)$
- $\Gamma_0 = \varphi_{1,0}(0) = \varphi'_{1,0}(1) = \varphi_\Omega(0) = \varphi'_\Omega(1) = [0]^\Omega Next\ \omega = [1][0] Next\ \omega = (R_2)^\Omega R_1 Hsuc\ 0 = R_3 R_2 R_1 Hsuc\ 0$
- $(R_2)^\Omega = R_3 R_2$
- $[0]^\Omega = [1][0].$

We will now see another way to use collapsing : ordinal collapsing functions.

An ordinal collapsing function is a function which, when applied to an uncountable ordinal, gives a countable ordinal.

The general idea is to define a set of ordinals $C(a)$ or $C(a,b)$ where a and b are ordinals, which contains all ordinals that can be built using an initial set of ordinals and some operations or functions, and then define $\psi(a)$ or $\psi(a,b)$ as the smallest ordinal which is not in $C(a)$ or $C(a,b)$, or the least ordinal which is greater than all countable ordinals of $C(a)$ or $C(a,b)$.

Different ordinal collapsing functions are described in http://googology.wikia.com/wiki/Ordinal_notation .

These functions are extensions of functions on countable ordinals, whose fixed points can be reached by applying them to an uncountable ordinal.

Here is a correspondence between basic notation systems and their collapsing extensions based on formula : least fixed point of $f = f(\Omega)$:

Basic notation	Formula	Limit	Extension	Correspondence	Crossing
Cantor	$cantor(\alpha, \beta)$ $= \beta + \omega^\alpha$	least $\alpha = cantor(\alpha, 0)$ $= \omega^\alpha = \varepsilon_0$	Taranovsky's C	$C(\alpha, \beta) = \beta + \omega^\alpha$ iff $C(\alpha, \beta) \geq \alpha$	$C(\Omega, 0) = \varepsilon_0$
Epsilon	ε_α	least $\alpha = \varepsilon_\alpha$ $= \zeta_0$	Madore's ψ	$\psi(\alpha) = \varepsilon_\alpha$ for all $\alpha < \zeta_0$	$\psi(\Omega) = \zeta_0$
Binary Veblen	$\varphi_\alpha(\beta)$ or $\varphi(\alpha, \beta)$	least $\alpha = \varphi(\alpha, 0)$ $= \Gamma_0$	θ	$\theta(\alpha, \beta) = \varphi(\alpha, \beta)$ below Γ_0	$\theta(\Omega, 0) = \Gamma_0$

10.1 Madore's ψ

This ordinal collapsing function is described in https://en.wikipedia.org/wiki/Ordinal_collapsing_function and <http://quibb.blogspot.fr/impredicative-ordinals.html> .

The definition of this function uses the ordinal Ω which is the least uncountable ordinal.

$C(\alpha)$ is the set of all ordinals constructible using only $0, 1, \omega, \Omega$ and addition, multiplication, exponentiation, and the function ψ (which will be defined later) restricted to ordinals smaller than α .

$\psi(\alpha)$ is the smallest ordinal not in $C(\alpha)$.

The smallest ordinal not in $C(0)$ is the limit of $\omega, \omega^\omega, \omega^{\omega^\omega}, \dots$ which is ε_0 , so $\psi(0) = \varepsilon_0$. More generally, $\psi(\alpha) = \varepsilon_\alpha$ for all $\alpha < \zeta_0$, $\psi(\alpha) = \zeta_0$ for $\zeta_0 \leq \alpha \leq \Omega$, and $\psi(\Omega + \alpha) = \varepsilon(\zeta_0 + \alpha)$ for $\alpha < \Omega$.

Note that $\psi(\Omega) = \zeta_0$ is the least fixed point of $\alpha \mapsto \varepsilon_\alpha$; we already saw such an equality when we introduced collapsing in the Veblen function.

The limit $\psi(\varepsilon_\Omega + 1)$ of $\psi(\Omega), \psi(\Omega^\Omega), \psi(\Omega^{\Omega^\Omega}), \dots$ is the Bachmann-Howard ordinal.

Some examples of fundamental sequences (FS) are :

A FS of ω is $0, 1, 2, 3, \dots$

A FS of $\psi(0)$ is $\omega, \omega^\omega, \omega^{\omega^\omega}, \dots$

A FS of $\psi(\alpha + 1)$ is $\psi(\alpha), \psi(\alpha)^{\psi(\alpha)}, \psi(\alpha)^{\psi(\alpha)^{\psi(\alpha)}}, \dots$

A FS of $\psi(f(\Omega))$ is $\psi(0), \psi(f(\psi(0))), \psi(f(\psi(f(\psi(0))))) , \dots$

For example :

A FS of $\psi(\Omega)$ is $\psi(0), \psi(\psi(0)), \psi(\psi(\psi(0))), \dots$

A FS of $\psi(\Omega \cdot 2)$ is $\psi(0), \psi(\Omega + \psi(0)), \psi(\Omega + \psi(\Omega + \psi(0))), \dots$

A FS of $\psi(\Omega^\Omega \cdot 3)$ is $\psi(0), \psi(\Omega^\Omega \cdot 2 + \Omega^{\psi(0)}), \psi(\Omega^\Omega \cdot 2 + \Omega^{\Omega^\Omega \cdot 2 + \Omega^{\psi(0)}}), \dots$

10.2 Correspondence between Veblen functions and Madore's ψ

To distinguish between the different Veblen functions, let us call φ_F the Veblen function with finitely many variables, and φ_T the Veblen function with transfinitely many variables.

φ_F is a function which, when applied to a list of countable ordinals, gives a countable ordinal. A list of countable ordinals can be seen as a function which, when applied to a natural number, gives a countable ordinal, with the restriction that the result differs from 0 for finitely many integers. If we denote ω the set of natural numbers and Ω the set of countable ordinals, then this can be written : $\varphi_F : (\omega \rightarrow \Omega) \rightarrow \Omega$. If we replace $\alpha \rightarrow \beta$ by β^α , we get Ω^{Ω^ω} , and if we apply ψ to it, we get $\psi(\Omega^{\Omega^\omega})$, which is the small Veblen ordinal, the least ordinal that cannot be reached using φ_F .

For φ_T , the position of a variable is represented by a countable ordinal instead of a natural number, also with the restriction that finitely many variables differ from 0, so we have $\varphi_T : (\Omega \rightarrow \Omega) \rightarrow \Omega$. If we replace $\alpha \rightarrow \beta$ by β^α , we get Ω^{Ω^Ω} , and if we apply ψ to it, we get $\psi(\Omega^{\Omega^\Omega})$, which is the large Veblen ordinal, the least ordinal that cannot be reached using φ_T .

10.3 θ function

θ function is a binary function. It's defined as follows:

- $C_0(\alpha, \beta) = \{\gamma \mid \gamma < \beta\} \cup \{0\}$.
- $C_{n+1}(\alpha, \beta) = \{\gamma + \delta \mid \gamma, \delta \in C_n(\alpha, \beta)\} \cup \{\theta(\gamma, \delta) \mid \gamma < \alpha \& \gamma, \delta \in C_n(\alpha, \beta)\} \cup \{\Omega_c \mid c \in C_n(\alpha, \beta)\}$.

- $C(\alpha, \beta) = \cup_{n < \omega} C_n(\alpha, \beta)$
- $\theta(\alpha, \beta) = \min\{c \mid (c \in C(\alpha, \gamma) \& (\forall \delta < \beta : \gamma > \theta(\alpha, \delta)))\}$
where $\Omega_0 = 0$ and Ω_a represents the a-th uncountable ordinal.

It means that $\theta(\alpha, \beta)$ is the $(1+\beta)$ -th ordinal such that it cannot be built from ordinals less than it by addition, applying $\theta(\delta, \dots)$ where $\delta < \alpha$ and getting an uncountable cardinal.

It seems that $\theta(\alpha, \beta) = \varphi(\alpha, \beta)$ below Γ_0 , making θ function an extension of φ function. Even $\theta(\Gamma_0, \beta) = \varphi(\Gamma_0, \beta)$ is true.

Other important values are :

- $\theta(\Omega, \alpha) = \Gamma_\alpha$
- $\theta(\Omega^\omega, 0) = \text{small Veblen ordinal}$
- $\theta(\Omega^\Omega, 0) = \text{large Veblen ordinal}$
- $\theta(\varepsilon_{\Omega+1}, 0) = \text{Bachmann Howard ordinal}$

See <https://stepstowardinfinity.wordpress.com/2015/05/04/ordinal2/> for more information.

10.4 Taranovsky's C

$C(a, b)$ is the least element above b that has degree a .

Definition: A degree for a well-ordered set S is a binary relation on S such that :

- Every element $c \in S$ has degree 0_S (the least element of S). 0_S only has degree 0_S .
- For a limit a , c has degree a iff it has every degree less than a .
- For a successor $a' = a+1$, either of the following holds:
 - An element has degree a' iff it is a limit of elements of degree a .
 - There is a limit element $d \models a$ such that for every c in S , c has degree a' iff it has degree a and either $c \models d$ or c is a limit of elements of degree a (or both).

Note: The third condition can be equivalently written as $\forall a (C_{a+1} = \lim(C_a) \vee \exists d \in \lim(S) \cap (a+1) C_{a+1} = \lim(C_a) \cup (C_a \cap (d+1)))$, where S is identified with an ordinal (so $a+1$ consists of ordinals $\leq a$), C_a is the set of elements that have degree a , and \lim is limit points.

In other terms : Let η be an ordinal, and let 0_S and let $Ld(a, b)$ be the statement that a is a limit of ordinals c such that $(c, b) \in D$. Let D be the following binary relation over η :

- $\forall a < \eta : (a, 0) \in D$
- $\forall a < \eta : a \neq 0 \Rightarrow (0, a) \notin D$
- $\forall b \in \text{Lim} \cup \eta : (a, b) \in D \Leftrightarrow \forall c < b : (a, c) \in D$
- $\forall b : (a, b) \in D \Leftrightarrow Ld(a, b+1) \forall b : (a, b) \in D \Leftrightarrow Ld(a, b+1)$
- $\forall b : \exists d \in \text{Lim} \cup \eta : d \leq b \Rightarrow \forall c : (c, a+1) \in D \Leftrightarrow (c \leq d \vee Ld(c, b))$

Then $C(a, b) = \min c : c \in \eta \wedge c > b \wedge (c, a) \in D$.

$C(a, b) = b + \omega^a$ iff $C(a, b) \geq a$.

For ordinals in the standard representation written in the postfix form, the comparison is done in the lexicographical order where ' C ' < ' 0 ' < ' Ω ': For example, $C(C(0, 0), 0) < C(\Omega, 0)$ because $000CC < 0\omega C$. (This does not hold for non-standard representations of ordinals.)

The fundamental sequences of Taranovsky's notation can be easily defined.

Let $L(\alpha)$ be the amount of C 's in standard representation of α , then $\alpha[n] = \max\{\beta \mid \beta < \alpha \wedge L(\beta) \leq L(\alpha) + n\}$.

Here is a summary of the system by Taranovsky (see <https://cs.nyu.edu/pipermail/fom/2012-March/016349.html>) :

I discovered a conjectured ordinal notation system that I conjecture reaches full second order arithmetic. I implemented the system in a python module/program:

<http://web.mit.edu/dmytro/www/other/OrdinalArithmetic.py>

along with ordinal arithmetic operations (addition, multiplication, exponentiation, etc.) and other functions. The ordinal arithmetic functionality is useful even if you are only interested in ordinals below ϵ_0 .

The notation system is simple enough to be defined in full here.

Definition: An ordinal a is 0-built from below from b iff $a \leq b$
 a is $n+1$ -built from below from b iff the standard representation of a does not use ordinals above a except in the scope of an ordinal n -built from below from b .

(Note: "in the scope of" means "as a subterm of".)

The n th (n is a positive integer) ordinal notation system is defined as follows.

Syntax: Two constants (0 , W_n) and a binary function C .

Comparison: For ordinals in the standard representation written in the postfix form, the comparison is done in the lexicographical order where ' C ' < ' 0 ' < ' W_n ': For example, $C(C(0,0),0) < C(W_n, 0)$ because $0\ 0\ 0\ C$
 $C < 0\ W_n\ C$.

Standard Form:

0 , W_n are standard

" $C(a, b)$ " is standard iff

1. " a " and " b " are standard,
2. b is 0 or W_n or " $C(c, d)$ " with $a \leq c$, and
3. a in n -built from below from b .

I conjecture that the strength of the n th ordinal notation system is between $\Pi^1_{n-1}\text{-CA}$ and $\Pi^1_n\text{-CA}_0$, and thus the sum of the order types of these ordinal notation systems is the proof-theoretical ordinal of second order arithmetic.

The full notation system is obtained by combining these notation systems as follows:

Constants 0 and W_i (for every positive integer i), and a binary function C .

$W_i = C(W_{i+1}, 0)$ and the standard form always uses W_i instead of $C(W_{i+1}, 0)$.

To check for standard form and compare ordinals use $W_i = C(W_{i+1}, 0)$ to convert each W to W_n for a single positive integer n (it does not matter which n) and then use the n th ordinal notation system.

To make C a total function for a and b in the notation system (this is not required for standard forms), let $C(a, b)$ be the least ordinal (in the notation system) of degree $\geq a$ above b , where the degree of W_i is W_{i+1} and the degree of $C(c,d)$ is c if " $C(c,d)$ " is the standard form. A polynomial time computation of $C(a, b)$ (that I believe is correct) is included in the program.

To complete ordinal analysis of second order arithmetic, one would need:

- * A canonical assignment of notations to formulas that provably in second order arithmetic denote an ordinal, and such that for every two ordinals/formulas, comparison is provable in second order arithmetic. The idea is that the notation system captures not only provably recursive ordinals of second order arithmetic but all ordinals that have a provable canonical definition in second order arithmetic. For example, W_1 is best assigned to the least admissible ordinal above ω . A partial assignment is in my paper. (It is because of such assignment that I believe that the system reaches full second order arithmetic.)
- * Proof that the system is well-founded and that it has the right strength, etc. (If you do not fully understand the notation system, or if you think that it is not well-founded, let me know.)

Historical Note: In 2005, I discovered the right general form of C , defined a notation system at the level of alpha-recursively inaccessible ordinals (FOM postings in August 2005), and had an idea for reaching second order arithmetic. In January 2006 (or possibly late 2005), I defined the notation system with W_2 and in 2009 (June 29, 2009 FOM posting) implemented it is a computer program. This year I defined the key concept -- n -built from below -- that allowed me to complete the full notation system.

Details about the ordinal notation system and its initial segments are in my paper:

<http://web.mit.edu/dmytro/www/other/OrdinalNotation.htm>

Sincerely,
Dmytro Taranovsky

Here are some examples of representations of some ordinals :

- $0 = 0$
- $1 = 0 + \omega^0 = C(0, 0)$
- $2 = 1 + \omega^0 = C(0, 1) = C(0, C(0, 0))$
- $\omega = 0 + \omega^1 = C(1, 0)$
- $\omega + 1 = \omega + \omega^0 = C(0, \omega) = C(0, C(1, 0))$
- $\omega \cdot 2 = \omega + \omega^1 = C(1, \omega) = C(1, C(1, 0))$
- $\omega^2 = 0 + \omega^2 = C(2, 0)$
- $\omega^\omega = 0 + \omega^\omega = C(\omega, 0) = C(C(1, 0), 0)$
- $\omega^{\omega^\omega} = 0 + \omega^{\omega^\omega} = C(\omega^\omega, 0) = C(C(C(1, 0), 0), 0)$
- $\varepsilon_0 = \varphi(1, 0) = \varphi'(0, 1) = C(\Omega_1, 0)$
- $\varepsilon_1 = \varphi(1, 1) = \varphi'(0, 2) = C(W, C(W, 0))$ (note that the correspondence with φ' is simpler than with φ)
- $\zeta_0 = \varphi(2, 0) = \varphi'(1, 1) = C(C(\Omega_1, \Omega_1), 0) = C(\Omega_1 \cdot 2, 0)$ with $\Omega_1 \cdot 2 = C(\Omega_1, \Omega_1)$
- $\zeta_1 = \varphi(2, 1) = \varphi'(1, 2) = C(\Omega_1 \cdot 2, C(\Omega_1 \cdot 2, 0))$
- $\eta_0 = \varphi(3, 0) = \varphi'(2, 1) = C(\Omega_1 \cdot 3, 0)$ with $\Omega_1 \cdot 3 = C(\Omega_1, C(\Omega_1, \Omega_1))$

- $\Gamma_0 = \varphi(1, 0, 0) = \varphi'(1, 0, 1) = C(C(\Omega_1 \cdot 2, \Omega_1), 0) = C(\Omega_1^2, 0)$ with $\Omega_1^2 = C(\Omega_1 \cdot 2, \Omega_1)$
- $\Gamma_1 = C(\Omega_1^2, C(\Omega_1^2, 0))$
- $\Gamma_\omega = C(\Omega_1^2 + 1, 0)$
- Small Veblen ordinal $= C(\Omega_1^\omega, 0)$
- Large Veblen ordinal $= C(\Omega_1^{\Omega_1}, 0)$
- Bachmann Howard ordinal $= C(C(\Omega_2, \Omega_1), 0)$

See <http://web.mit.edu/dmytro/www/other/OrdinalNotation.htm> and <https://stepstowardinfinity.wordpress.com/2015/06/22/ordinal3/> for more information.

11 Ordinal trees

Ordinal can also be represented by trees. An example of such a representation is given in : <http://www.madore.org/~david/math/ordtrees.pdf> .

In this representation, the order on finite rooted trees is recursively defined as follows : $A < B$ if and only if one of this conditions is true :

- There is some immediate subtree B' of B such that $A \leq B'$.
- Every child A' of A satisfies $A' < B$ and the list of children of A is lexicographically less than the list of children of B for the order $<$ with the leftmost children having the most weight.

Trees can also be represented by parenthesized expressions, for example :

- $0 = ()$
- $1 = (())$
- $2 = ((()))$
- $\omega = (()())$
- $\omega + 1 = (((())))$
- $\omega \cdot 2 = (()())()$
- $\omega \cdot 3 = (()((())))$
- $\omega^2 = (()())()$
- $\omega^2 + \omega = (()((()())))$
- $\omega^2 \cdot 2 = (()()())()$
- $\omega^2 \cdot 3 = (()()((())))$
- $\omega^3 = (()()()())$
- $\omega^\omega = (((()))())$
- $\omega^{\omega^\omega} = (((()())())())$
- $\epsilon_0 = \varphi_1(0) = (()()())$
- $\epsilon_1 = \varphi_1(1) = (()()())()$
- $\epsilon_2 = \varphi_1(2) = (()()())()$
- $\epsilon_\omega = \varphi_1(\omega) = (()()()())$
- $\epsilon_{\epsilon_0} = (()()()())()$
- $\zeta_0 = \varphi_2(0) = (()()())()$
- $\Gamma_0 = \varphi(1, 0, 0) = (((()))())$
- $\varphi(1, 0, 0, 0) = (()()()())$
- ...

Another example of tree representation is Takeuti ordinal diagrams, see : https://projecteuclid.org/download/pdf_1/euclid.jmsj/1261153819 .

12 Summary

Any ordinal can be defined as the least ordinal strictly greater than all ordinals of a set : the empty set for 0, $\{\alpha\}$ for the successor of α , $\{\alpha_0, \alpha_1, \alpha_2, \dots\}$ for an ordinal with fundamental sequence $\alpha_0, \alpha_1, \alpha_2, \dots$

12.1 Algebraic notation

We define the following operations on ordinals :

- addition : $\alpha + 0 = \alpha$; $\alpha + \text{suc}(\beta) = \text{suc}(\alpha + \beta)$; $\alpha + \text{lim}(f) = \text{lim}(n \mapsto \alpha + f(n))$
- multiplication : $\alpha \times 0 = 0$; $\alpha \times \text{suc}(\beta) = (\alpha \times \beta) + \alpha$; $\alpha \times \text{lim}(f) = \text{lim}(n \mapsto \alpha \times f(n))$
- exponentiation : $\alpha^0 = 1$; $\alpha^{\text{suc}(\beta)} = \alpha^\beta \times \alpha$; $\alpha^{\text{lim}(f)} = \text{lim}(n \mapsto \alpha^{f(n)})$

12.2 Veblen functions

These functions use fixed points enumeration : $\varphi(\dots, \beta, 0, \dots, 0, \gamma)$ represents the $(1 + \gamma)^{\text{th}}$ common fixed point of the functions $\xi \mapsto \varphi(\dots, \delta, \xi, 0, \dots, 0)$ for all $\delta < \beta$.

12.3 Simmons notation

$\text{Fix}fz = f^w(z + 1) =$ least fixed point of f strictly greater than z .

$\text{Next} = \text{Fix}(\alpha \mapsto \omega^\alpha)$

$[0]h = \text{Fix}(\alpha \mapsto h^\alpha \omega)$; $[1]hg = \text{Fix}(\alpha \mapsto h^\alpha g\omega)$; $[2]hgf = \text{Fix}(\alpha \mapsto h^\alpha g f\omega)$; etc...

Correspondence with Veblen's ϕ : $\phi(1 + \alpha, \beta) = ([0]^\alpha \text{Next})^{1+\beta} \omega$; $\phi(\alpha, \beta, \gamma) = ([0]^\beta (([1][0])^\alpha \text{Next}))^{1+\gamma} \omega$

12.4 RHS0 notation

We start from 0, if we don't see any regularity we take the successor, if we see a regularity, if we have a notation for this regularity, we use it, else we invent it, then we jump to the limit.

$Hfx = \text{lim } x, fx, f(fx), \dots$; $R_1fgx = \text{lim } gx, fgx, ffgx, \dots$; $R_2fghx = \text{lim } hx, fghx, fgfghx, \dots$

Correspondence with Simmons notation : $\dots, [3] \rightarrow R5, [2] \rightarrow R4, [1] \rightarrow R3, [0] \rightarrow R2, \text{Next} \rightarrow R1, \omega \rightarrow H\text{suc } 0$

12.5 Ordinal collapsing functions

These functions use uncountable ordinals to define countable ordinals.

We define sets of ordinals that can be built from given ordinals and operations, then we take the least ordinal which is not in this set, or the least ordinal which is greater than all countable ordinals of this set.

These functions are extensions of functions on countable ordinals, whose fixed points can be reached by applying them to an uncountable ordinal.

Examples :

- Madore's ψ : $\psi(\alpha) = \varepsilon_\alpha$ if $\alpha < \zeta_0$; $\psi(\Omega) = \zeta_0$ which is the least fixed point of $\alpha \mapsto \varepsilon_\alpha$.
- Feferman's θ : $\theta(\alpha, \beta) = \varphi(\alpha, \beta)$ if $\alpha < \Gamma_0$ and $\beta < \Gamma_0$; $\theta(\Omega, 0) = \Gamma_0$ which is the least fixed point of $\alpha \mapsto \varphi(\alpha, 0)$.
- Taranovsky's C : $C(\alpha, \beta) = \beta + \omega^\alpha$ if α is countable; $C(\Omega_1, 0) = \varepsilon_0$ which is the least fixed point of $\alpha \mapsto \omega^\alpha$.

13 Comparison table

Name	Symbol	Algebraic	Veblen	Simmons	RHS0	Madore	Taranovsky
Zero	0	0			0		0
One	1	1	$\varphi(0, 0)$		suc 0		$C(0, 0)$
Two	2	2			suc (suc 0)		$C(0, C(0, 0))$
Omega	ω	ω	$\varphi(0, 1)$	ω	H suc 0		$C(1, 0)$
		$\omega + 1$			suc (H suc 0)		$C(0, C(1, 0))$
		$\omega \times 2$			H suc (H suc 0)		$C(1, C(1, 0))$
		ω^2	$\varphi(0, 2)$		H (H suc) 0		$C(C(0, C(0, 0)), 0)$
		ω^ω	$\varphi(0, \omega)$		H H suc 0		$C(C(1, 0), 0)$
		ω^{ω^ω}	$\varphi(0, \omega^\omega)$		H H H suc 0		$C(C(C(1, 0), 0), 0)$
Epsilon zero	ε_0	ε_0	$\varphi(1, 0)$	$Next \ \omega$	$R_1 H suc \ 0$	$\psi(0)$	$C(\Omega_1, 0)$
		ε_1	$\varphi(1, 1)$	$Next^2 \omega$	$R_1(R_1 H) suc \ 0$	$\psi(1)$	$C(\Omega_1, C(\Omega_1, 0))$
		ε_ω	$\varphi(1, \omega)$	$Next^\omega \omega$	$H R_1 H suc \ 0$	$\psi(\omega)$	$C(C(0, \Omega_1), 0)$
		$\varepsilon_{\varepsilon_0}$	$\varphi(1, \varphi(1, 0))$	$Next^{Next^\omega \omega}$	$R_1 H R_1 H suc \ 0$	$\psi(\psi(0))$	$C(C(C(\Omega_1, 0), \Omega_1), 0)$
Zeta zero	ζ_0	ζ_0	$\varphi(2, 0)$	$[0] Next \ \omega$	$R_2 R_1 H suc \ 0$	$\psi(\Omega)$	$C(C(\Omega_1, \Omega_1), 0)$
Eta zero	η_0	η_0	$\varphi(3, 0)$	$[0]^2 Next \ \omega$	$R_2(R_2 R_1) H suc \ 0$		$C(C(\Omega, C(\Omega, \Omega)), 0)$
			$\varphi(\omega, 0)$	$[0]^\omega Next \ \omega$	$H R_2 R_1 H suc \ 0$		$C(C(C(0, \Omega_1), \Omega_1), 0)$
Feferman -Schütte	Γ_0	Γ_0	$\varphi(1, 0, 0)$ $= \varphi(2 \mapsto 1)$	$[1][0] Next \ \omega$	$R_3 R_2 R_1 H suc \ 0$ $= R_{3 \dots 1} H suc \ 0$	$\psi(\Omega^\Omega)$	$C(C(C(\Omega_1, \Omega_1), \Omega_1), 0)$
Ackermann			$\varphi(1, 0, 0, 0)$ $= \varphi(3 \mapsto 1)$	$[1]^2 [0] Next \ \omega$	$R_3(R_3 R_2) R_1 H suc \ 0$	$\psi(\Omega^{\Omega^2})$	
Small Veblen ordinal			$\varphi(\omega \mapsto 1)$	$[1]^\omega [0] Next \ \omega$	$H R_3 R_2 R_1 H suc \ 0$	$\psi(\Omega^{\Omega^\omega})$	$C(\Omega_1^\omega, 0)$ $= C(C(C(C(0, \Omega_1), \Omega_1), \Omega_1), 0)$
Large Veblen ordinal			least ord. not rep.	$[2][1][0] Next \ \omega$	$R_4 R_3 R_2 R_1 H suc \ 0$ $= R_{4 \dots 1} H suc \ 0$	$\psi(\Omega^{\Omega^\Omega})$	$C(\Omega_1^{\Omega^\Omega}, 0)$ $= C(C(C(C(C(\Omega_1, \Omega_1), \Omega_1), \Omega_1), \Omega_1), 0)$
Bachmann- Howard ordinal				least ord. not rep.	$R_{\omega \dots 1} H suc \ 0$	$\psi(\varepsilon_{\Omega+1})$	$C(C(\Omega_2, \Omega_1), 0)$

14 Links

- <http://www.madore.org/%7Edavid/weblog/2011-09-18-nombres-ordinaux-intro.html> : Tutorial introduction to ordinal numbers in French
- <https://sites.google.com/site/pointlesslargenumberstuff/home/1/pglin?tmpl=%2Fsystem%2Fapp%2Ftemplates%2Fprint%2F> : Pointless Gigantic List of Infinite Numbers
- https://sites.google.com/site/largenumber/home/appendix/a/infinite_numbers : Sbiis Saibian's !!! FORBIDDEN LIST !!! of Infinite Numbers
- <http://quibb.blogspot.fr/p/infinity-series-portal.html> : Professor Quibb's Infinity Series Portal
- http://googology.wikia.com/wiki/Ordinal_notation : Ordinal notation
- <https://sites.google.com/site/travelingtotheinfinity/> : Traveling to the infinity
- <http://www.cs.man.ac.uk/~hsimmons/TEMP/OrdNotes.pdf> : A short introduction to Ordinal Notations by Harold Simmons
- http://www.mathematik.uni-muenchen.de/~buchholz/articles/jaegerfestschr_buchholz3.pdf : A survey on ordinal notations around the Bachmann-Howard ordinal by Wilfried Buchholz
- <http://web.mit.edu/dmytro/www/other/OrdinalNotation.htm> : Ordinal Notation by Dmytro Taranovsky
- <http://arxiv.org/html/1203.2270> : Higher Order Set Theory with Reflective Cardinals by Dmytro Taranovsky
- <https://www1.maths.leeds.ac.uk/~rathjen/realms.pdf> : The Realm of Ordinal Analysis by Michael Rathjen
- https://cage.ugent.be/~jvdm/Site/Research_files/DissertationJeroenVanderMeerenPrinted.pdf : Connecting the Two Worlds: Well-partial-orders and Ordinal Notation Systems by Jeroen Van der Meeren
- https://en.wikipedia.org/wiki/Veblen_function : Veblen function on Wikipedia

- <http://www.ams.org/journals/tran/1908-009-03/S0002-9947-1908-1500814-9/S0002-9947-1908-1500814-9.pdf> : Continuous increasing functions of finite and transfinite ordinals by Oswald Veblen
- http://en.wikipedia.org/wiki/Ordinal_collapsing_function : Ordinal collapsing function on Wikipedia
- https://en.wikipedia.org/wiki/Buchholz_psi_functions : Buchholz psi functions on Wikipedia
- <http://www.madore.org/%7Edavid/math/ordtrees.pdf> : Ordinal trees
- <https://johncarlosbaez.wordpress.com/2016/06/29/large-countable-ordinals-part-1/> : Large Countable Ordinals by John Baez, Part 1
- <https://johncarlosbaez.wordpress.com/2016/07/04/large-countable-ordinals-part-2/> : Large Countable Ordinals by John Baez, Part 2
- <https://johncarlosbaez.wordpress.com/2016/07/07/large-countable-ordinals-part-3/> : Large Countable Ordinals by John Baez, Part 3
- <https://medium.com/@joshkerr/mind-blown-the-fast-growing-hierarchy-for-laymen-aka-enormous-numbers-d9a865c6443b> : Mind blown: the fast growing hierarchy for laymen — aka enormous numbers
- <https://sites.google.com/site/largenumbers/home> : Sbiis Saibian's Large Number Site
- <http://forums.xkcd.com/viewtopic.php?f=14&t=7469> : My number is bigger !
- <http://www.cl.cam.ac.uk/%7Ejrh13/papers/reflect.html> : Metatheory and Reflection in Theorem Proving: A Survey and Critique by John Harrison
- <http://math.stanford.edu/%7Efeferman/papers/penrose.pdf> : Penrose's Gödelian argument by Solomon Feferman
- <http://www.turingarchive.org/browse.php/B/15> : Systems of logic based on ordinals by Alan Turing
- <https://coq.inria.fr/documentation> : Coq documentation
- <http://wiki.portal.chalmers.se/agda/pmwiki.php?n=Main.Documentation> : Agda documentation