

The EMC Effect for Tritium and Helium-3 from JLab's MARATHON Experiment using DIS.

Jason Bane

University of Tennessee

jbane1@vols.utk.edu

Thomas Jefferson National Accelerator Facility is a U.S.

Department of Energy Office of Science national laboratory.

Jefferson Lab's unique and exciting mission is to expand humankind's knowledge of the universe by **studying the fundamental building blocks of matter** within the nucleus: subatomic particles known as **quarks and gluons**.




More than 1,500 nuclear physicists worldwide come to Jefferson Lab to conduct and collaborate on research.

Outline


- Deep Inelastic Scattering
- The EMC Effect
- The MARATHON Experiment
- Data Analysis
- EMC Effect of $A=3$

Deep Inelastic Scattering (DIS)




- $Q^2 \equiv 4EE' \sin^2 \frac{\theta}{2}$
- $X_{Bj} = \frac{Q^2}{2\nu M}$
- $\nu \equiv E - E'$
- $W^2 = 2M\nu + M^2 - Q^2$
- $W^2 > 4 \rightarrow \text{DIS}$


DIS ??????




Idealized spectra of high-energy electron scattering as a function of energy transfer [G. T. Garvey, et al., 2015].





- Elastic scattering





- Elastic scattering
- Quasielastic scattering






- Elastic scattering

- Quasieastic scattering

- DIS




Why DIS?

$$\sigma_{eN} = \frac{\alpha^2}{4E^2 \sin^4\left(\frac{\theta}{2}\right)} \left[\frac{F_2(x)}{\nu} \cos^2\left(\frac{\theta}{2}\right) + \frac{2F_1(x)}{M} \sin^2\left(\frac{\theta}{2}\right) \right]$$


Quark parton model

- $F_2(x) = x \cdot \sum_i z_i^2 f_i(x)$
- $F_1(x) = 1/2 \cdot \sum_i z_i^2 f_i(x)$
- **Spin 1/2 quarks**
 $F_2(x) = F_1(x) \cdot 2x$



Ratio of $2x \cdot F_1(x)$ and $F_2(x)$ vs. x .
 [Povh, 1995]

The EMC Effect




[D. Higinbotham, 2013]

European Muon Collaboration

- X = fraction of momentum carried by quark
- Expected Unity \rightarrow Binding $<$ interaction
- $F^A = Z \cdot F^p + (A - Z) \cdot F^n$
- Unexpected Relative Decrease
- Missing high-momentum quarks in $A > 2$
- EMC Effect \equiv structure of the A/D Ratio


EMC Effect

SLAC experiment E139 [J. Gomez et al., 1994] .



EMC Effect


JLab experiment E03103
[J.Seely, A. Daniel et al, 2009]



- Studied for last 30+ years
- CERN, SLAC, HERMES, BCDMS, and JLab
- Slope of A/D Ratios from 0.3 - 0.7 in x
- Models have difficulty matching data for all criteria
- $\approx \log$ dependence in A

EMC Effect

JLab experiment E03103
 [J.Seely, A. Daniel et al, 2009]



- Studied for last 30+ years
- CERN, SLAC, HERMES, BCDMS, and JLab
- Slope of A/D Ratios from 0.3 - 0.7 in x
- Models have difficulty matching data for all criteria
- $\approx \log$ dependence in A
- sensitivity to "local" effects



EMC Effect



- Studied for last 30+ years
- CERN, SLAC, HERMES, BCDMS, and JLab
- Slope of A/D Ratios from 0.3 - 0.7 in x
- Models have difficulty matching data for all criteria
- ~~≈ log dependence in A~~
- sensitivity to "local" effects


MARATHON

MeAsurement of F_2^n/F_2^p , d/u RAtios and $A = 3$ EMC Effect in Deep Inelastic Electron Scattering off the Tritium and Helium MirOr Nuclei.



- Lightest and simplest mirror system
 - ▶ Number of protons in 3H = neutrons in 3He
- Differences in the nuclear effects are small

The Continuous Electron Beam Accelerator Facility (CEBAF) at Thomas Jefferson Accelerator Facility.



- ≈ 2.2 GeV per revolution
- 12 GeV for Hall D
- Superconducting RF cavities
- RF separators split the beam to each Hall

MARATHON's proposal

- 11 GeV Beam
- Bigbite spectrometer
- Hall A's high resolution spectrometers (HRS)
- Tritium


Uses of Tritium




Tritium Target Cell

First tritium target at JLab

- Thin Al entrance and exit windows 0.01 inches
- 1090Ci of Tritium (0.1 g)
- 25 cm long
- Tritium Cell was filled in Savannah River
- 40 kelvin Helium is used to cool an attached heat sink



Bigbite Spectrometer



Large Acceptance Spectrometer


- Solid angle = 96 mrad
- Large Momentum Acceptance: $> 500 \text{ MeV}/c$

Preparing Bigbite


- Signal quality checks
- PMT performance studies
- Designing trigger logic
- Supply detector HV

After about 2 years of work:
Removed from then run plan!

Hall A




High Resolution Spectrometers (HRSs)






The Run Period



Rey Torres

Kinematic Coverage



Kinematic coverage between Q^2 , x , and Θ . The band around the points represents the approximate spectrometer acceptance in the y axis.

Preparing Data for Analysis

Calibration

ADC calibration

- Calorimeters, Scintillators, and Cherenkov


TDC calibration

- Scintillators and VDC


Detector calibrations

- BPMs, BCMS, Optics


Cherenkov Calibration



Beam Position Monitor(BPM) Calibration




Intrusive absolute position measurement




Relative position measurement

$$\begin{pmatrix} X_{position} \\ Y_{position} \end{pmatrix} = \begin{pmatrix} C(0,0) & C(0,1) \\ C(0,0) & C(0,1) \end{pmatrix} * \begin{pmatrix} X_{BPM} \\ Y_{BPM} \end{pmatrix} + \begin{pmatrix} X_{offset} \\ Y_{offset} \end{pmatrix}$$

Beam position from BPM and harp for a collection of runs



Before Calibration



After Calibration

Experimentally Measured Cross Section

$$\frac{d\sigma}{dE'd\Omega} = \frac{(N - BG)}{L \cdot \epsilon \cdot \Delta E' \Delta \Omega \cdot A(E', \theta)}$$




- N = Number of electrons
- BG = Background
- L = Luminosity
 - ▶ Density Correction
- ϵ = Efficiency
- $\Delta E' \Delta \Omega$ = Bin Size
- $A(E', \theta)$ = Acceptance probability


Charge Symmetric Background

- γ decay into an e^+e^- pairs
- Pair produced e^- by detecting e^+
- Extraction based on fit to Exponential function


Pair Production



Tritium positron contamination. Credit: Tong Su




Aluminum Endcap Background



- Extract ratio of the normalized yield from the gas cell to that of the empty cell

3H Decay




$$\tau(^3H) = 4500 \pm 8 \text{ days}$$

$$c = \frac{\eta_{^3He}}{\eta_{tot}}$$





$$\sigma_{^3H} = \left(\frac{\sigma_{tot}}{\sigma_{^3He}} \right) \left(\frac{1}{1 - c} \right) - \left(\frac{1}{1 - c} \right)$$

Beta Decay Helium Fraction



Tyler Kutz

Density Fluctuations


(a) ${}^3\text{H}$ Density Analysis.(b) ${}^3\text{He}$ Density Analysis.(c) ${}^2\text{H}$ Density Analysis.(d) ${}^1\text{H}$ Density Analysis.

[S.N.Santiesteban et. al (2019)]

Efficiencies (ϵ)

- Particle Identification(PID)
 - ▶ Cherenkov
 - ▶ Calorimeters
- Trigger
 - ▶ Scintillators
- Tracking
 - ▶ Vertical Drift Chambers(VDCs)
- Electronic Deadtime

Particle ID Efficiency



- Use well defined samples
- Apply PID cuts and count
- PID efficiency $\approx 98\%$ for all kinematics

Monte Carlo Ratio Method

$$Y_{MC}(E', \theta) = L \cdot \sigma^{model} \cdot (\Delta E' \Delta \Omega) \cdot A(E', \theta)$$

$$\sigma_{data} = \frac{Y_{data}(E', \theta)}{L \cdot (\Delta E' \Delta \Omega) \cdot A(E', \theta)}$$

Use a Monte Carlo simulation


- $(\Delta E' \Delta \Omega)_{Data} = (\Delta E' \Delta \Omega)_{MC}$
- $A(E', \theta)_{Data} = A(E', \theta)_{MC}$

$$\sigma_{Data} = \sigma_{model} \cdot \frac{Y_{Data}}{Y_{MC}}$$


Monte Carlo

- Generate events → Pass through Magnetic apertures
- Tune Simulation offsets to match detector response
- Use model to weight events
 - ▶ Deep Inelastic and resonance region from Ari Bodek Fit from E139

[A. Bodek and U.K. Yang, 2002]



DIS Cross Section




Normalization uncertainty due to target thickness uncertainty
He3 - 1.12% • H3 - 0.97% • D - 0.56%

Relative Error Contributions in % for Cross Section for a selection of bins



X bin	0.215	0.455	0.705
Statistical Error	0.512	0.889	1.106
Positron Correction Error	0.036	0.016	0.005
Efficiency Error*	0.665	1.477	2.951
Density Correction Error	0.002	0.002	0.002
Monte Carlo Error	0.193	0.217	0.209
Total Error	0.95	1.931	3.316

* Contains contributions from PID, tracking, trigger, and livetime

Per Nucleon Cross Section Ratio





Isoscalar Corrections




- Correct for the unpaired nucleon in the A/D ratio.

$$\text{Cor.} = \frac{\left(0.5 * (1.0 + \frac{F_2^n}{F_2^p})\right)}{\left(\frac{1}{A} \cdot (Z + (A - Z) \cdot \frac{F_2^n}{F_2^p})\right)}$$


Isoscalar Corrections



$$\text{Cor.} = \frac{\left(0.5 * (1.0 + \frac{F_2^n}{F_2^p})\right)}{\left(\frac{1}{A} \cdot (Z + (A - Z) \cdot \frac{F_2^n}{F_2^p})\right)}$$



EMC Effect




Future Plans

- Model dependence on cross section extraction
- Study the edge of the HRS acceptance for a long target
- Model Dependence on Isoscalar correction
- Determine the magnitude of the EMC effect for Tritium and Helium-3



Detailed Studies of the nuclear dependence of F2 in light nuclei [E12-100-008: J. Arrington, A Daniel, NF, D. Gaskell]


Target Choice motivated by physics impact



Coming soon* in Hall C

What is soon?


Slide credit Nadia Fomin



The EMC effect in spin structure functions

[E12-14-001: Will Brooks and Sebastian Kuhn]

[ICC, W. Bentz and A. W. Thomas, Phys. Lett. B **642**, 210 (2006)]



- A polarized EMC effect arises because in-medium quarks are more relativistic
 - Lower components of quark wave functions are enhanced
 - Quark Spin is converted to orbital angular momentum
- Spin Dependent cross-section is suppressed by $1/A$
- Experiment to measure spin structure functions of ${}^7\text{Li}$

Slide credit Nadia Fomin

Summary

- The puzzling EMC Effect - decrease in high momentum quarks $A > 2$
- DIS - probe the quarks
- Used JLab's 12GeV electron beam & Hall A's HRS
- Use the ${}^3\text{He}$ & ${}^3\text{H}$ EMC Effect to add a puzzle piece to the EMC puzzle.

Special Thanks

- Nadia Fomin and Douglas Higinbotham
- The Tritium Students
- DOE and JSA
- Rex Tayloe for the invite.



The JLab MARATHON Tritium Collaboration

D. Abrams, H. Albataineh, **S. Alsalmi**, D. Androic, K. Aniol, W. Armstrong, J. Arrington, H. Atac, T. Averett, C. Ayerbe Gayoso, X. Bai, **J. Bane***, **S. Barcus**, A. Beck, V. Bellini, H. Bhatt, D. Bhetuwal, D. Biswas, D. Blyth, W. Boeglin, D. Bulumulla, A. Camsonne, **M. Carmignotto**, **J. Castellanos**, J-P. Chen, C. Ciofi degli Atti, E. O. Cohen, S. Covrig, K. Craycraft, **R. Cruz-Torres**, B. Dongwi, M. Duer, B. Duran, D. Dutta, N. Fomin, E. Fuchey, C. Gal, T. N. Gautam, S. Gilad, K. Gnanvo, T. Gogami, J. Gomez, C. Gu, A. Habarakada, **T. Hague***, O. Hansen, M. Hattawy, **F. Hauenstein**, O. Hen, D. W. Higinbotham, R. Holt, E. Hughes, C. Hyde, H. Ibrahim, S. Jian, S. Joosten, A. Karki, B. Karki, A. T. Katramatou, C. Keppel, M. Khachatryan, V. Khachatryan, A. Khanal, D. King, P. King, I. Korover, S. A. Kulagin, **T. Kutz***, N. Lashley-Colthirst, G. Laskaris, **S. Li**, W. Li, **H. Liu***, S. Liuti, N. Liyanage, D. Lonardoni, R. Machleidt, L.E. Marcucci, P. Markowitz, **E. McClellan**, D. Meekins, W. Melnitchouk, S. Mey-Tal Beck, Z-E. Meziani, R. Michaels, M. Mihovilović, V. Nelyubin, **D. Nguyen**, N. Nuruzzaman, **M. Nycz***, R. Obrecht, M. Olson, L. Ou, V. Owen, E. Pace, **B. Pandey**, V. Pandey, A. Papadopoulou, M. Paolone, S. Park, M. Patsyuk, S. Paul, G. G. Petratos, R. Pettit, E. Piasetzky, R. Pomatsalyuk, S. Premathilake, A. J. R. Puckett, V. Punjabi, R. Ransome, M. N. H. Rashad, P. E. Reimer, S. Riordan, J. Roche, F. Sammarruca, G. Salmè, **N. Santiesteban**, B. Sawatzky, J. Segal, E. P. Segarra, B. Schmookler, A. Schmidt, S. Scopetta, A. Shahinyan, S. Sirca, N. Sparveris, **T. Su***, R. Suleiman, H. Szumila-Vance, A. S. Tadepalli, L. Tang, W. Tireman, F. Tortorici, G. Urciuoli, M. Viviani, L. B. Weinstein, B. Wojtsekowski, S. Wood, **Z. H. Ye**, Z. Y. Ye, and J. Zhang.

More than 140 Collaborators

Red-Boldfaced Names: Tritium Program grad students; **starred**: MARATHON Ph.D. students

Blue-Boldfaced Names: Tritium Program postdoctoral associates



The JLab MARATHON Tritium Collaboration

Forty Five Institutions (in no particular order): University of Virginia; Texas A & M University; Kent State University; University of Zagreb; California State University, Los Angeles; Argonne National Laboratory; Temple University; The College of William and Mary; University of Tennessee; Massachusetts Institute of Technology; INFN Sezione di Catania; INFN Sezione di Roma, INFN Sezione di Pisa; Mississippi State University; Hampton University; Florida International University; Old Dominion University; Jefferson Lab; University of Perugia; Tel Aviv University; University of Connecticut; Tohoku University; Columbia University; Cairo University; Ohio University; Stony Brook, State University of New York; Syracuse University; Nuclear Research Center-Negev, Beer-Sheva; Institute for Nuclear Research of the Russian Academy of Sciences; University of New Hampshire; University of Regina; Columbia University; Facility for Rare Isotope Beams, Michigan State University; Los Alamos National Laboratory; University of Idaho; University of Pisa; Jožef Stefan Institute, University of Ljubljana; Johannes Gutenberg-Universität Mainz; Saint Norbert College; Center for Neutrino Physics, Virginia Tech; University of South Carolina; Kharkov Institute of Physics and Technology; Norfolk State University; Rutgers University; Artem Alikhanian National Laboratory; Tel Aviv University; Northern Michigan University; University of Illinois, Chicago.

Twelve Countries: Armenia, Canada, Croatia, Egypt, Germany, Israel, Italy, Japan, Russia, Slovenia, Ukraine, United States.



References I



A. Bodek and U.K. Yang

Nuclear Physics B, Procc. Suppl. 112 (2002) 70-76



Garvey, G. T. et. al.

Phys. Rept., 580 (2015) 1-45



J. Gomez et al.

Phys. Rev D 49 (1994) 4348



Douglas Higinbotham (2013)

The EMC effect still puzzles after 30 years. *Cern Courier* April 2013.



K. Nakamura et. al.

Review of Particle Physics, 37 (2010)



S.N.Santiestebana et. al

Nucl. Instrum. Meth. A 940 (2019) 351-358



J.Seely, A. Daniel et al

Phys. Rev. Lett. 103, 202301 (2009).



Bogdan Povh,

Particles and Nuclei: An Introduction to the Physical Concepts, (1995)