

Short corn 2025

yield comparison

Jesse Bealsburg

2025-10-31

Date = 2025-10-31

Site = Rosemount, R22 and R34 Field

Methods = Harvested 43 ft of 2 rows at 30" row spacing in 6 adjacent corn rows. Moisture and test weight were estimated two ways...

1. using the HarvestMaster system on combine and
2. using a Perten 5200 on a subsample immediately afterwards

Does rosemount combine accurately determine moisture and test weight compared to benchtop equipment?

Data


Testweight = good. with 0.5 lbs per bushel.

Moisture = ok, within 1.5 - 2 moisture %


Discussion

The corn was bagged and driven back (1hr) in bed of truck during a light drizzle which could possibly have slightly increased moisture. When brought back to campus, they were put in dryer for 30 min, then analyzed.

moisture



test weight



data

raw

site	position	method	moisture	testweight
r22	middle	Perten 5200	22.8	52.5
r22	middle	HarvestMaster on combine	21.1	52.0
r22	north	Perten 5200	23.2	50.9
r22	north	HarvestMaster on combine	20.6	50.1
r22	south	Perten 5200	23.3	50.2
r22	south	HarvestMaster on combine	21.2	49.6
r34	east	Perten 5200	20.3	52.5
r34	east	HarvestMaster on combine	18.3	51.6
r34	middle	Perten 5200	19.9	52.6
r34	middle	HarvestMaster on combine	18.6	53.7
r34	west	Perten 5200	19.6	55.2
r34	west	HarvestMaster on combine	18.1	52.7

summarized

Table 2: moisture


site	Perten 5200	HarvestMaster on combine	difference	%difference
r22	23.10000	20.96667	2.133333	9.235209
r34	19.93333	18.33333	1.600000	8.026756

Table 3: test weight

site	Perten 5200	HarvestMaster on combine	difference	%difference
r22	51.20000	50.56667	0.633333	1.236979
r34	53.43333	52.66667	0.766667	1.434810


more in depth figures

moisture



- Trend is consistent across site*position

test weight



- Trend is consistent across site*position except R34 middle

photos



Figure 1: Short corn strip (2 rows at 43 feet long) harvested with wintersteiger quantum on 31Oct2025