



‘Recent Progress in Collisionless Models,’ Imperial College 8 Sept 2015



Phase mixing vs. nonlinear advection in drift-kinetic plasma turbulence

Alex Schekochihin

w i t h

Joseph Parker, Paul Dellar (Oxford Maths),

Edmund Highcock (Oxford Th. Phys.),

Bill Dorland (Maryland), Greg Hammett (Princeton)




*with thanks to the Wolfgang Pauli Institute, Vienna,
where much of this work was conceived and done
(and where maths meet physics as if they'd always been together)*

arXiv:1508.05988



CAMBRIDGE
UNIVERSITY PRESS




Norbert and Claude are coming for you...

<http://journals.cambridge.org/pla>



‘Recent Progress in Collisionless Models,’ Imperial College 8 Sept 2015



Phase mixing vs. nonlinear advection in drift-kinetic plasma turbulence

Alex Schekochihin

w i t h

Joseph Parker, Paul Dellar (Oxford Maths),

Edmund Highcock (Oxford Th. Phys.),

Bill Dorland (Maryland), Greg Hammett (Princeton)



*with thanks to the Wolfgang Pauli Institute, Vienna,
where much of this work was conceived and done
(and where maths meet physics as if they'd always been together)*

arXiv:1508.05988



‘Recent Progress in Collisionless Models,’ Imperial College 8 Sept 2015



Phase mixing vs. nonlinear advection in drift-kinetic plasma turbulence

Alex Schekochihin

w i t h


Joseph Parker, Paul Dellar (Oxford Maths),
Edmund Highcock (Oxford Th. Phys.),
Bill Dorland (Maryland), Greg Hammett (Princeton)



*with thanks to the Wolfgang Pauli Institute, Vienna,
where much of this work was conceived and done
(and where maths meet physics as if they'd always been together)*

arXiv:1508.05988

You put a tokamak on your web page at your peril...



I will assume you really want to know
what sort of problems fusion physicists think about

*(actually, what I will give you is my take on what's interesting, to me, right now;
disclaimer: I am not really a hard-core fusion physicist,
most of them actually think of much more practical things...)*



A Prototypical Kinetic Problem

Plasma near Maxwellian equilibrium:

$$f = F_0 + \delta f$$

Strong (uniform) magnetic field:

$$\omega \ll \Omega_i, \quad k_{\parallel} \ll k_{\perp}$$

Electrostatic:

$$\delta \mathbf{E} = -\nabla \phi, \quad \delta \mathbf{B} = 0$$

Long wavelength:

$$k_{\perp} \rho_i \ll 1$$

\Downarrow

Drift-kinetic equation for

$$g(t, \mathbf{r}, v_{\parallel}) = \frac{1}{n_{0i}} \int d^2 \mathbf{v}_{\perp} \delta f_i$$

$$\frac{\partial g}{\partial t} + v_{\parallel} \nabla_{\parallel} (g + \varphi F_0) + \mathbf{u}_{\perp} \cdot \nabla_{\perp} g = C[g] + \chi$$

\uparrow parallel \uparrow $\mathbf{E} \times \mathbf{B}$ drift \uparrow coll. \uparrow energy
 el. field nonlinearity phase injection
 $\mathbf{u}_{\perp} = \frac{\rho_i v_{th}}{2} \hat{\mathbf{z}} \times \nabla_{\perp} \varphi$ space cutoff

Boltzmann electrons:

$$\varphi \equiv \frac{Ze\phi}{T_i} = \alpha \int dv_{\parallel} g, \quad \alpha = \frac{ZT_e}{T_i}$$

ITG/drift-wave problem:

$$\chi = -\mathbf{u}_{\perp} \cdot \nabla F_0 = -\frac{\rho_i v_{th}}{2} \frac{\partial \varphi}{\partial y} \left[\frac{1}{L_n} + \left(\frac{v_{\parallel}^2}{v_{th}^2} - \frac{1}{2} \right) \frac{1}{L_T} \right]$$



A Prototypical Kinetic Problem

Plasma near Maxwellian equilibrium:

$$f = F_0 + \delta f$$

Strong (uniform) magnetic field:

$$\omega \ll \Omega_i, \quad k_{\parallel} \ll k_{\perp}$$

Electrostatic:

$$\delta \mathbf{E} = -\nabla \phi, \quad \delta \mathbf{B} = 0$$

Long wavelength:

$$k_{\perp} \rho_i \ll 1$$

\Downarrow

Drift-kinetic equation for

$$g(t, \mathbf{r}, v_{\parallel}) = \frac{1}{n_{0i}} \int d^2 \mathbf{v}_{\perp} \delta f_i$$

$$\frac{\partial g}{\partial t} + v_{\parallel} \nabla_{\parallel} (g + \varphi F_0) + \mathbf{u}_{\perp} \cdot \nabla_{\perp} g = C[g] + \chi$$

\uparrow parallel \uparrow $\mathbf{E} \times \mathbf{B}$ drift \uparrow coll. \uparrow energy
 el. field nonlinearity phase injection
 $\mathbf{u}_{\perp} = \frac{\rho_i v_{th}}{2} \hat{\mathbf{z}} \times \nabla_{\perp} \varphi$ space cutoff

Boltzmann electrons:

$$\varphi \equiv \frac{Ze\phi}{T_i} = \alpha \int dv_{\parallel} g, \quad \alpha = \frac{ZT_e}{T_i}$$

ITG/drift-wave problem:

$$\chi = -\mathbf{u}_{\perp} \cdot \nabla F_0 = -\frac{\rho_i v_{th}}{2} \frac{\partial \varphi}{\partial y} \left[-\frac{1}{n_i} \frac{dn_i}{dx} - \left(\frac{v_{\parallel}^2}{v_{th}^2} - \frac{1}{2} \right) \frac{1}{T_i} \frac{dT_i}{dx} \right]$$



The Plasma Turbulence Problem

Energy injected into perturbations can be thermalised:

EITHER by **phase mixing** (=Landau damping), producing fine scales in v_{\parallel} and thus making $C[g]$ finite even if the collisionality is small:

$$C[g] \sim \nu v_{\text{th}}^2 \frac{\partial^2 g}{\partial v_{\parallel}^2} \sim \omega g \quad \text{if} \quad \frac{\delta v_{\parallel}}{v_{\text{th}}} \sim \left(\frac{\nu}{\omega}\right)^{1/2}$$

$$\frac{\partial g}{\partial t} + \underbrace{v_{\parallel} \nabla_{\parallel} (g)}_{\text{phase mixing}} + \varphi F_0 + \underbrace{\mathbf{u}_{\perp} \cdot \nabla_{\perp} g}_{\text{turbulent mixing}} = C[g] + \chi$$

AND/OR by **turbulent mixing**, producing fine scales in real space, eventually accessing various dissipation mechanisms at $k_{\perp} \rho_i \lesssim 1$ (which are an interesting but separate story, for another talk)

So what does the system choose to do?

The Plasma Turbulence Problem



“Idle” theory questions:


- Which **thermalisation route** does the system favour?
- Therefore, what is the **structure of turbulence** at scales between injection and dissipation (in phase space, so φ, g vs. $k_{\perp}, k_{\parallel}, v_{\perp}$)



‘Which here is the “+”
and which is the “-”?’

“Pragmatic” modeling questions:

- At what **rate** is the injected energy removed to small scales?
- Therefore, what is the typical **amplitude** of the fluctuations?
(get that by balancing injection rate with removal rate)
- Therefore, what is the typical “**turbulent diffusivity**” relaxing large-scale gradients?



$$D_T \sim \underbrace{\langle u_{\perp}^2 \rangle}_{\text{amplitude}} \underbrace{\tau_c}_{\text{correlation time}}$$



Free Energy

“Energy” in δf kinetics is in fact the **free energy** of the fluctuations:

$$\mathcal{F} = - \sum_s T_s \delta S_s = - \sum_s T_s \delta \int d^3 \mathbf{v} \langle f_s \ln f_s \rangle = \sum_s \int d^3 \mathbf{v} \frac{T_s \langle \delta f_s^2 \rangle}{2F_{0s}} = n_i T_i W$$

where $W = \int dv_{\parallel} \frac{\langle g^2 \rangle}{2F_0} + \frac{\langle \varphi^2 \rangle}{2\alpha}$ is conserved by our equations

$$\frac{\partial g}{\partial t} + v_{\parallel} \nabla_{\parallel} (g + \varphi F_0) + \mathbf{u}_{\perp} \cdot \nabla_{\perp} g = C[g] + \chi$$

$$\varphi \equiv \frac{Ze\phi}{T_i} = \alpha \int dv_{\parallel} g, \quad \alpha = \frac{ZT_e}{T_i}$$

$$\frac{dW}{dt} = \int dv_{\parallel} \frac{\langle g\chi \rangle}{F_0} + \int dv_{\parallel} \frac{\langle gC[g] \rangle}{F_0}$$

↑
injection
(instabilities,
forcing...)

↑
dissipation
(collisions)

Kruskal & Oberman 1958

Bernstein 1958

Fowler 1963, 68

Krommes & Hu 1994

Krommes 1999

Sugama et al. 1996

Hallatschek 2004

Howes et al. 2006

Candy & Waltz 2006

Schekochihin et al. 2007-09

Scott 2010

Banon, Teaca, Hatch, Morel,

Jenko et al. 2011-14

Plunk et al 2012

Abel et al. 2013

Kunz et al. 2015

...



Landau Damping = Phase Mixing

Landau damping/phase mixing is the transfer of free energy from φ to g via refinement of velocity-space structure of the perturbed distribution

$$W = \int dv_{\parallel} \frac{\langle g^2 \rangle}{2F_0} + \frac{\langle \varphi^2 \rangle}{2\alpha} \text{ is conserved by our equations}$$

$$\frac{\partial g}{\partial t} + v_{\parallel} \nabla_{\parallel} (g + \varphi F_0) + \mathbf{u}_{\perp} \cdot \nabla_{\perp} g = C[g] + \chi$$

$$\varphi \equiv \frac{Ze\phi}{T_i} = \alpha \int dv_{\parallel} g, \quad \alpha = \frac{ZT_e}{T_i}$$

$$\frac{dW}{dt} = \int dv_{\parallel} \frac{\langle g\chi \rangle}{F_0} + \int dv_{\parallel} \frac{\langle gC[g] \rangle}{F_0}$$

↑
↑
injection
dissipation
(instabilities,
(collisions)
forcing...)

Hammett,
Perkins,
Dorland,
Beer,
Smith,
Snyder 1990-2001:

*development
of Landau fluid models
based on understanding
of phase mixing
as energy removal
into phase space
and eventual
collisional thermalisation*



Life in Hermite Space

$$H_m(x) = (-1)^m e^{x^2} \frac{d^m}{dx^m} e^{-x^2} \Rightarrow g_m = \int dv_{\parallel} \frac{H_m(v_{\parallel}/v_{\text{th}})}{\sqrt{2^m m!}} g(v_{\parallel})$$

$$H_0 = 1 \Rightarrow g_0 = \frac{\delta n}{n} = \frac{\varphi}{\alpha}$$

$$H_1 = 2x \Rightarrow g_1 = \sqrt{2} \frac{u_{\parallel}}{v_{\text{th}}}$$

$$H_2 = 4 \left(x^2 - \frac{1}{2} \right) \Rightarrow g_2 = \frac{1}{\sqrt{2}} \frac{\delta T_{\parallel}}{T}, \text{ etc.}$$

“fluid” moments
 $m = 0, 1, 2$

$$\frac{\partial g}{\partial t} + v_{\parallel} \nabla_{\parallel} (g + \varphi F_0) + \mathbf{u}_{\perp} \cdot \nabla_{\perp} g = C[g] + \chi$$

$$\varphi \equiv \frac{Ze\phi}{T_i} = \alpha \int dv_{\parallel} g, \quad \alpha = \frac{ZT_e}{T_i}$$

$$\chi = -\mathbf{u}_{\perp} \cdot \nabla F_0 = -\frac{\rho_i v_{\text{th}}}{2} \frac{\partial \varphi}{\partial y} \left[\frac{1}{L_n} + \left(\frac{v_{\parallel}^2}{v_{\text{th}}^2} - \frac{1}{2} \right) \frac{1}{L_T} \right]$$

H_0 H_2 energy
injection
only at low m



Life in Hermite Space

$$\underline{\frac{\partial}{\partial t} \frac{\varphi}{\alpha}} + \underline{v_{th} \nabla_{\parallel} \frac{u_{\parallel}}{v_{th}}} = -\frac{v_{th}}{2L_n} \rho_i \frac{\partial \varphi}{\partial y}$$

Standard “fluid”

ITG equations

(e.g., Cowley et al. 1992)

$$\left(\underline{\frac{\partial}{\partial t}} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{u_{\parallel}}{v_{th}} + v_{th} \nabla_{\parallel} \left(\underline{\frac{1}{2} \frac{\delta T_{\parallel}}{T}} + \frac{1+\alpha}{\alpha} \varphi \right) = 0$$

$$\omega^3 \approx \frac{\alpha}{2} (k_{\parallel} v_{th})^2 \omega_{*T}$$

$$\left(\underline{\frac{\partial}{\partial t}} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{\delta T_{\parallel}}{T} + v_{th} \nabla_{\parallel} \left(\sqrt{3} g_3 + 2 \frac{u_{\parallel}}{v_{th}} \right) = -\underline{\underline{\frac{v_{th}}{2L_T} \rho_i \frac{\partial \varphi}{\partial y}}}$$

$$\omega_{*T} = \frac{k_y \rho_i v_{th}}{2L_T}$$

From $m=3$ on, the equations are universal:

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) g_m + v_{th} \nabla_{\parallel} \left(\sqrt{\frac{m+1}{2}} g_{m+1} + \sqrt{\frac{m}{2}} g_{m-1} \right) = -\nu m g_m$$

↑

all moments are
advected by the same
velocity

$$\mathbf{u}_{\perp} = \frac{\rho_i v_{th}}{2} \hat{\mathbf{z}} \times \nabla_{\perp} \varphi$$

↑

higher moments couple to lower ones,
so **even though free energy is injected**
at low m , it gets to large m

↑

at large enough m ,
free energy
is removed
by collisions

NB: we use the LB operator

$$C[g] = \nu \frac{\partial}{\partial v_{\parallel}} \left(\frac{1}{2} \frac{\partial}{\partial v_{\parallel}} + v_{\parallel} \right) g$$



Life in Hermite Space

$$\frac{\partial}{\partial t} \frac{\varphi}{\alpha} + v_{th} \nabla_{\parallel} \frac{u_{\parallel}}{v_{th}} = -\frac{v_{th}}{2L_n} \rho_i \frac{\partial \varphi}{\partial y}$$

Standard “fluid”

ITG equations

(e.g., Cowley et al. 1992)

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{u_{\parallel}}{v_{th}} + v_{th} \nabla_{\parallel} \left(\frac{1}{2} \frac{\delta T_{\parallel}}{T} + \frac{1+\alpha}{\alpha} \varphi \right) = 0$$

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{\delta T_{\parallel}}{T} + v_{th} \nabla_{\parallel} \left(\sqrt{3} g_3 + 2 \frac{u_{\parallel}}{v_{th}} \right) = -\frac{v_{th}}{2L_T} \rho_i \frac{\partial \varphi}{\partial y}$$

The **free energy** is (via Parseval’s theorem for H_m ’s)

$$W = \sum_{m=3}^{\infty} \frac{\langle g_m^2 \rangle}{2} + \frac{1}{4} \frac{\langle \delta T_{\parallel}^2 \rangle}{T^2} + \frac{\langle u_{\parallel}^2 \rangle}{v_{th}^2} + \frac{1+\alpha}{2\alpha^2} \langle \varphi^2 \rangle$$



Life in Hermite Space

$$\frac{\partial}{\partial t} \frac{\varphi}{\alpha} + v_{th} \nabla_{\parallel} \frac{u_{\parallel}}{v_{th}} = -\frac{v_{th}}{2L_n} \rho_i \frac{\partial \varphi}{\partial y}$$

Standard “fluid”

ITG equations

(e.g., Cowley et al. 1992)

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{u_{\parallel}}{v_{th}} + v_{th} \nabla_{\parallel} \left(\frac{1}{2} \frac{\delta T_{\parallel}}{T} + \frac{1+\alpha}{\alpha} \varphi \right) = 0$$

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{\delta T_{\parallel}}{T} + v_{th} \nabla_{\parallel} \left(\sqrt{3} g_3 + 2 \frac{u_{\parallel}}{v_{th}} \right) = -\frac{v_{th}}{2L_T} \rho_i \frac{\partial \varphi}{\partial y}$$

The **free energy** is (via Parseval’s theorem for H_m ’s)

Landau damping/phase mixing is the transfer of free energy
From low moments ($\varphi, u_{\parallel}, \delta T_{\parallel}$) into higher ones ($g_{m \geq 3}$).

$$W = \sum_{m=3}^{\infty} \frac{\langle g_m^2 \rangle}{2} + \frac{1}{4} \frac{\langle \delta T_{\parallel}^2 \rangle}{T^2} + \frac{\langle u_{\parallel}^2 \rangle}{v_{th}^2} + \frac{1+\alpha}{2\alpha^2} \langle \varphi^2 \rangle$$

Turbulence (in the usual sense) is the mixing of $\varphi, u_{\parallel}, \delta T_{\parallel}$
by \mathbf{u}_{\perp} transferring their energy to small scales (large k_{\perp}).



“Fluid” Turbulence Theory

$$\frac{\partial}{\partial t} \frac{\varphi}{\alpha} + v_{th} \nabla_{\parallel} \frac{u_{\parallel}}{v_{th}} = -\frac{v_{th}}{2L_n} \rho_i \frac{\partial \varphi}{\partial y}$$

Standard “fluid”

ITG equations

(e.g., Cowley et al. 1992)

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{u_{\parallel}}{v_{th}} + v_{th} \nabla_{\parallel} \left(\frac{1}{2} \frac{\delta T_{\parallel}}{T} + \frac{1+\alpha}{\alpha} \varphi \right) = 0$$

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{\delta T_{\parallel}}{T} + v_{th} \nabla_{\parallel} \left(\cancel{\sqrt{3} g_3} + 2 \frac{u_{\parallel}}{v_{th}} \right) = -\frac{v_{th}}{2L_T} \rho_i \frac{\partial \varphi}{\partial y}$$

Let us construct a turbulence theory for ITG ignoring coupling to phase space...

$$W = \cancel{\sum_{m=3}^{\infty} \frac{\langle g_m^2 \rangle}{2}} + \frac{1}{4} \frac{\langle \delta T_{\parallel}^2 \rangle}{T^2} + \frac{\langle u_{\parallel}^2 \rangle}{v_{th}^2} + \frac{1+\alpha}{2\alpha^2} \langle \varphi^2 \rangle$$

Turbulence (in the usual sense) is the mixing of $\varphi, u_{\parallel}, \delta T_{\parallel}$ by \mathbf{u}_{\perp} transferring their energy to small scales (large k_{\perp}).



“Fluid” Turbulence Theory: Outer Scale

$$\frac{\partial}{\partial t} \frac{\varphi}{\alpha} + v_{th} \nabla_{\parallel} \frac{u_{\parallel}}{v_{th}} = - \frac{v_{th}}{2L_n} \rho_i \frac{\partial \varphi}{\partial y}$$

Standard “fluid”

ITG equations

(e.g., Cowley et al. 1992)

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{u_{\parallel}}{v_{th}} + v_{th} \nabla_{\parallel} \left(\frac{1}{2} \frac{\delta T_{\parallel}}{T} + \frac{1+\alpha}{\alpha} \varphi \right) = 0$$

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{\delta T_{\parallel}}{T} + v_{th} \nabla_{\parallel} \left(\cancel{\sqrt{3} g_3} + 2 \frac{u_{\parallel}}{v_{th}} \right) = - \frac{v_{th}}{2L_T} \rho_i \frac{\partial \varphi}{\partial y}$$

Let us construct a turbulence theory for ITG ignoring coupling to phase space...

Where is energy injected?

➤ “Linear balance”: $k_{\parallel 0} v_{th} \sim \omega_{*T} = k_{y0} \rho_i \frac{v_{th}}{L_T}$

(for ITG injection to work)

➤ Largest possible scale: $k_{\parallel 0} \sim \frac{1}{L_{\parallel}} \left(= \frac{1}{qR} \right)$

➤ Isotropy: $k_{x0} \sim k_{y0} \sim k_{\perp 0}$

($k_{x0} \sim S_{ZF} k_{y0} \tau_c \sim k_{y0}$ if $S_{ZF} \sim \tau_c^{-1}$)

↑
zonal flow shear

$$k_{\perp 0} \rho_i \sim \frac{L_T}{L_{\parallel}}$$

[Barnes, Parra & AAS PRL **107**, 115003 (2011)]



“Fluid” Turbulence Theory: Outer Scale

$$\frac{\partial}{\partial t} \frac{\varphi}{\alpha} + v_{th} \nabla_{\parallel} \frac{u_{\parallel}}{v_{th}} = -\frac{v_{th}}{2L_n} \rho_i \frac{\partial \varphi}{\partial y}$$

Standard “fluid”
ITG equations
(e.g., Cowley et al. 1992)

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{u_{\parallel}}{v_{th}} + v_{th} \nabla_{\parallel} \left(\frac{1}{2} \frac{\delta T_{\parallel}}{T} + \frac{1+\alpha}{\alpha} \varphi \right) = 0$$

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{\delta T_{\parallel}}{T} + v_{th} \nabla_{\parallel} \left(\cancel{\sqrt{3} g_3} + 2 \frac{u_{\parallel}}{v_{th}} \right) = -\frac{v_{th}}{2L_T} \rho_i \frac{\partial \varphi}{\partial y}$$

Let us construct a turbulence theory for ITG ignoring coupling to phase space...

At what rate is energy removed from this scale?

$$\omega_{*T} \sim k_{\perp 0} \rho_i \frac{v_{th}}{L_T} \sim k_{\perp 0} u_{\perp 0} \sim \rho_i v_{th} k_{\perp 0}^2 \varphi_0$$

injection
nonlinear removal to smaller
scales, $k_{\perp} > k_{\perp 0}$

$$\Downarrow$$

$$\varphi_0 \sim \frac{1}{k_{\perp 0} L_T}$$

$$k_{\perp 0} \rho_i \sim \frac{L_T}{L_{\parallel}}$$

$$\varphi_0 \sim \frac{\rho_i L_{\parallel}}{L_T^2}$$



“Fluid” Turbulence Theory: Outer Scale

$$\frac{\partial}{\partial t} \frac{\varphi}{\alpha} + v_{th} \nabla_{\parallel} \frac{u_{\parallel}}{v_{th}} = - \frac{v_{th}}{2L_n} \rho_i \frac{\partial \varphi}{\partial y}$$

Standard “fluid”
ITG equations
(e.g., Cowley et al. 1992)

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{u_{\parallel}}{v_{th}} + v_{th} \nabla_{\parallel} \left(\frac{1}{2} \frac{\delta T_{\parallel}}{T} + \frac{1+\alpha}{\alpha} \varphi \right) = 0$$

$$\left(\frac{\partial}{\partial t} + \mathbf{u}_{\perp} \cdot \nabla_{\perp} \right) \frac{\delta T_{\parallel}}{T} + v_{th} \nabla_{\parallel} \left(\cancel{\sqrt{3} g_3} + 2 \frac{u_{\parallel}}{v_{th}} \right) = - \frac{v_{th}}{2L_T} \rho_i \frac{\partial \varphi}{\partial y}$$

Let us construct a turbulence theory for ITG ignoring coupling to phase space...

What is the turbulent diffusivity?

$$D_T \sim u_{\perp 0}^2 \tau_c \sim \frac{u_{\perp 0}}{k_{\perp 0}} \sim \rho_i v_{th} \varphi_0 \sim \frac{\rho_i^2 v_{th}}{L_{\parallel}} \left(\frac{L_{\parallel}}{L_T} \right)^2$$

$$k_{\perp 0} \rho_i \sim \frac{L_T}{L_{\parallel}}$$

$$\varphi_0 \sim \frac{\rho_i L_{\parallel}}{L_T^2}$$

and so the **heat flux** is

More formally,

$Q = \langle u_x \delta T_{\parallel} \rangle$
and you want to bound
it from above by some
function of L_T .


$$Q \sim \frac{n D_T T}{L_T} \sim \frac{n \rho_i^2 v_{th}}{L_{\parallel}^2} \left(\frac{L_{\parallel}}{L_T} \right)^3$$

gyro-Bohm stiff

[Barnes, Parra & AAS PRL **107**, 115003 (2011)]



“Fluid” Turbulence Theory: Outer Scale



Let us construct a turbulence theory for ITG ignoring coupling to phase space...

These scalings basically work.

$$k_{\perp 0}\rho_i \sim \frac{L_T}{L_{\parallel}}$$


$$\varphi_0 \sim \frac{\rho_i L_{\parallel}}{L_T^2}$$

$$Q \sim \frac{nD_T T}{L_T} \sim \frac{n\rho_i^2 v_{th}}{L_{\parallel}^2} \left(\frac{L_{\parallel}}{L_T} \right)^3$$

gyro-Bohm stiff

[Barnes, Parra & AAS PRL **107**, 115003 (2011)]

“Fluid” Turbulence Theory: Inertial Range



Let us construct a turbulence theory for ITG ignoring coupling to phase space...

These scalings basically work.

So what? All this says is that leakage rate into phase space, $\sim k_{\parallel 0} v_{th}$, is at most same order as $k_{\perp 0} u_{\perp 0}$, as we indeed assumed!

A more sensitive (if less interesting to modelers) question is how free energy cascades to smaller scales...



“Fluid” Turbulence Theory: Inertial Range

Kolmogorov-style argument: constant flux of free energy to small scales


NB: assuming no damping, i.e., energy stays in “fluid” ($m=0, 1, 2$) moments.

Then, at $k_{\perp} > k_{\perp 0}$,

$$\frac{\varphi^2}{\tau_c} \sim k_{\perp} u_{\perp} \varphi^2 \sim k_{\perp}^2 \varphi^3 = \text{const} \Rightarrow \varphi \propto k_{\perp}^{-2/3}$$

The “1D spectrum”:

$$E(k_{\perp}) = 2\pi k_{\perp} \int dk_{\parallel} \langle |\varphi_{\mathbf{k}}|^2 \rangle \sim \frac{\varphi^2}{k_{\perp}} \propto k_{\perp}^{-7/3}$$



$$W = \cancel{\sum_{m=3}^{\infty} \frac{\langle g_m^2 \rangle}{2}} + \frac{1}{4} \frac{\langle \delta T_{\parallel}^2 \rangle}{T^2} + \frac{\langle u_{\parallel}^2 \rangle}{v_{\text{th}}^2} + \frac{1 + \alpha}{2\alpha^2} \langle \varphi^2 \rangle$$



“Fluid” Turbulence Theory: Inertial Range

Kolmogorov-style argument: constant flux of free energy to small scales


NB: assuming no damping, i.e., energy stays in “fluid” ($m=0, 1, 2$) moments.

Then, at $k_{\perp} > k_{\perp 0}$,

$$\frac{\varphi^2}{\tau_c} \sim k_{\perp} u_{\perp} \varphi^2 \sim k_{\perp}^2 \varphi^3 = \text{const} \Rightarrow \varphi \propto k_{\perp}^{-2/3}$$


The “1D spectrum”:

$$E(k_{\perp}) = 2\pi k_{\perp} \int dk_{\parallel} \langle |\varphi_{\mathbf{k}}|^2 \rangle \sim \frac{\varphi^2}{k_{\perp}} \propto k_{\perp}^{-7/3}$$



Critical balance: by **causality**, turbulence cannot stay correlated at parallel scales larger than those over which linear communication happens faster than nonlinear decorrelation:
so, no correlation if

$$k_{\parallel} v_{th} < k_{\perp} u_{\perp} \propto k_{\perp}^{4/3} \Rightarrow k_{\parallel} L_{\parallel} < \left(\frac{k_{\perp}}{k_{\perp 0}} \right)^{4/3}$$



[Barnes, Parra & AAS PRL **107**, 115003 (2011)]



“Fluid” Turbulence Theory: Inertial Range

Kolmogorov-style argument: constant flux of free energy to small scales

NB: assuming no damping, i.e., energy stays in “fluid” ($m=0, 1, 2$) moments.

Then, at $k_{\perp} > k_{\perp 0}$,

$$\frac{\varphi^2}{\tau_c} \sim k_{\perp} u_{\perp} \varphi^2 \sim k_{\perp}^2 \varphi^3 = \text{const} \Rightarrow \varphi \propto k_{\perp}^{-2/3}$$

The “1D spectrum”:

$$E(k_{\perp}) = 2\pi k_{\perp} \int dk_{\parallel} \langle |\varphi_{\mathbf{k}}|^2 \rangle \sim \frac{\varphi^2}{k_{\perp}} \propto k_{\perp}^{-7/3}$$

That this works suggests
there is no phase mixing
in the inertial range

Critical balance: by causality, turbulence cannot stay correlated at parallel scales larger than those over which linear communication happens faster than nonlinear decorrelation:
so, no correlation if

$$k_{\parallel} v_{\text{th}} < k_{\perp} u_{\perp} \propto k_{\perp}^{4/3} \Rightarrow k_{\parallel} L_{\parallel} < \left(\frac{k_{\perp}}{k_{\perp 0}} \right)^{4/3}$$

That this works suggests
the notional phase mixing
rate $\sim k_{\parallel} v_{\text{th}}$
is nevertheless same order
as $k_{\perp} u_{\perp}$ at all scales.

*So why is there no exponential
cutoff of the spectrum?*



Hermite “Cascade”

Let us go back to our kinetic equation and now ask how transfer of free energy to high m 's occurs linearly:

$$\left(\frac{\partial}{\partial t} + \cancel{\mathbf{u}_{\perp} \cdot \nabla_{\perp}} \right) g_m + v_{th} \nabla_{\parallel} \left(\sqrt{\frac{m+1}{2}} g_{m+1} + \sqrt{\frac{m}{2}} g_{m-1} \right) = -\nu m g_m$$

In Fourier space: $\nabla_{\parallel} \rightarrow i k_{\parallel}$, $\tilde{g}_m(k_{\parallel}) = (i \operatorname{sgn} k_{\parallel})^m g_m(k_{\parallel})$

$$\frac{\partial \tilde{g}_m}{\partial t} + \frac{|k_{\parallel}| v_{th}}{\sqrt{2}} (\sqrt{m+1} \tilde{g}_{m+1} - \sqrt{m} \tilde{g}_{m-1}) = -\nu m \tilde{g}_m$$



Hermite “Cascade”

Let us go back to our kinetic equation and now ask how transfer of free energy to high m 's occurs linearly:

$$\left(\frac{\partial}{\partial t} + \cancel{\mathbf{u}_\perp \cdot \nabla_\perp} \right) g_m + v_{th} \nabla_\parallel \left(\sqrt{\frac{m+1}{2}} g_{m+1} + \sqrt{\frac{m}{2}} g_{m-1} \right) = -\nu m g_m$$

In Fourier space: $\nabla_\parallel \rightarrow i k_\parallel$, $\tilde{g}_m(k_\parallel) = (i \operatorname{sgn} k_\parallel)^m g_m(k_\parallel)$

$$\frac{\partial \tilde{g}_m}{\partial t} + \frac{|k_\parallel| v_{th}}{\sqrt{2}} \left(\sqrt{m+1} \tilde{g}_{m+1} - \sqrt{m} \tilde{g}_{m-1} \right) = -\nu m \tilde{g}_m$$

↑
this looks like a derivative: indeed,

$$\begin{aligned} &= \sqrt{m} \left(\sqrt{1 + \frac{1}{m}} \tilde{g}_{m+1} - \tilde{g}_{m-1} \right) \\ &\approx \sqrt{m} \left(\tilde{g}_m + \frac{1}{2m} + \frac{\partial \tilde{g}_m}{\partial m} - \tilde{g}_m + \frac{\partial \tilde{g}_m}{\partial m} \right) \\ &= 2m^{1/4} \frac{\partial}{\partial m} m^{1/4} \tilde{g}_m \quad \text{this propagates perturbations} \\ &\quad \text{towards higher } m \end{aligned}$$



Hermite “Cascade”

Let us go back to our kinetic equation and now ask how transfer of free energy to high m 's occurs linearly:

$$\left(\frac{\partial}{\partial t} + \cancel{\mathbf{u}_\perp \cdot \nabla_\perp} \right) g_m + v_{th} \nabla_\parallel \left(\sqrt{\frac{m+1}{2}} g_{m+1} + \sqrt{\frac{m}{2}} g_{m-1} \right) = -\nu m g_m$$

In Fourier space: $\nabla_\parallel \rightarrow i k_\parallel$, $\tilde{g}_m(k_\parallel) = (i \operatorname{sgn} k_\parallel)^m g_m(k_\parallel)$

$$\frac{\partial \tilde{g}_m}{\partial t} + \frac{|k_\parallel| v_{th}}{\sqrt{2}} m^{1/4} \frac{\partial}{\partial m} m^{1/4} \tilde{g}_m = -\nu m \tilde{g}_m$$


Hermite spectrum $C_m = \frac{1}{2} \langle |g_m|^2 \rangle$ satisfies

$$\cancel{\frac{\partial C_m}{\partial t}} + |k_\parallel| v_{th} \frac{\partial}{\partial m} \sqrt{2m} C_m = -2\nu m C_m$$

stationary

Hermite flux,
constant for $m \ll m_c$.

$$C_m \propto \frac{1}{\sqrt{m}} e^{-(m/m_c)^{3/2}}$$





Hermite “Cascade”

So this is what Landau damping looks like in a system with some persistent energy source at low m [see detailed tutorial in Kanekar et al. JPP **81**, 305810104 (2015)]

It will dissipate collisionally all the energy that is injected, at the rate $\sim |k_{\parallel}| v_{th}$, independent of collisionality (because the m spectrum is shallow):

$$\frac{dW}{dt} = (\text{injection}) - \sum_{k_{\parallel}} \sum_m 2\nu m C_m(k_{\parallel})$$

NB: W diverges as $\sim \nu^{-1/3}$

$$\sim \nu \int_0^{m_c} dm \sqrt{m} \sim \nu m_c^{3/2} \sim |k_{\parallel}| v_{th}$$


Hermite spectrum $C_m = \frac{1}{2} \langle |g_m|^2 \rangle$ satisfies

$$\cancel{\frac{\partial C_m}{\partial t}} + |k_{\parallel}| v_{th} \frac{\partial}{\partial m} \sqrt{2m} C_m = -2\nu m C_m$$

stationary

↑
Hermite flux,
constant for $m \ll m_c$.

$$C_m \propto \frac{1}{\sqrt{m}} e^{-(m/m_c)^{3/2}}$$



$$m_c = \left(\frac{3|k_{\parallel}| v_{th}}{2\sqrt{2}\nu} \right)^{2/3}$$

[Zocco & AAS PoP **18**, 102309 (2011)]




Hermite “Cascade”

So this is what Landau damping looks like in a system with some persistent energy source at low m [see detailed tutorial in Kanekar et al. JPP **81**, 305810104 (2015)]

It will dissipate collisionally all the energy that is injected, at the rate $\sim |k_{\parallel}| v_{th}$, independent of collisionality (because the m spectrum is shallow):

$$\frac{dW}{dt} = (\text{injection}) - \sum_{k_{\parallel}} \sum_m 2\nu m C_m(k_{\parallel})$$

$$\uparrow \sim \nu \int_0^{m_c} dm \sqrt{m} \sim \nu m_c^{3/2} \sim |k_{\parallel}| v_{th}$$



“Un-phase-mixing”



The crucial step that gave us robust phase mixing was assuming continuity in m space:

$$\frac{\partial \tilde{g}_m}{\partial t} + \frac{|k_{\parallel}| v_{th}}{\sqrt{2}} (\sqrt{m+1} \tilde{g}_{m+1} - \sqrt{m} \tilde{g}_{m-1}) = -\nu m \tilde{g}_m$$
$$\Downarrow$$
$$\frac{\partial \tilde{g}_m}{\partial t} + \frac{|k_{\parallel}| v_{th}}{\sqrt{2}} m^{1/4} \frac{\partial}{\partial m} m^{1/4} \tilde{g}_m = -\nu m \tilde{g}_m$$



“Un-phase-mixing”

The crucial step that gave us robust phase mixing was assuming continuity in m space:

$$\cancel{\frac{\partial \tilde{g}_m}{\partial t}} + \frac{|k_{\parallel}| v_{th}}{\sqrt{2}} (\sqrt{m+1} \tilde{g}_{m+1} - \sqrt{m} \tilde{g}_{m-1}) = \cancel{-\nu m \tilde{g}_m}$$

$$\Downarrow$$

$$\frac{\partial \tilde{g}_m}{\partial t} + \frac{|k_{\parallel}| v_{th}}{\sqrt{2}} m^{1/4} \frac{\partial}{\partial m} m^{1/4} \tilde{g}_m = -\nu m \tilde{g}_m$$

For $1 \ll m \ll m_c$, to lowest order,

$$\sqrt{m+1} \tilde{g}_{m+1} - \sqrt{m} \tilde{g}_{m-1} = 0 \Rightarrow \tilde{g}_{m+1} \approx \tilde{g}_{m-1}$$

This allows two solutions: $\tilde{g}_{m+1} \approx \pm \tilde{g}_m$, so either \tilde{g}_m or $(-1)^m \tilde{g}_m$ is continuous.

This can be encoded in the following decomposition:

$$\tilde{g}_m = \tilde{g}_m^+ + (-1)^m \tilde{g}_m^-$$

where $\tilde{g}_m^+ = \frac{\tilde{g}_m + \tilde{g}_{m+1}}{2}$ and $\tilde{g}_m^- = (-1)^m \frac{\tilde{g}_m - \tilde{g}_{m+1}}{2}$ are continuous in m .



“Un-phase-mixing”

$$\frac{\partial \tilde{g}_m}{\partial t} + \frac{|k_{\parallel}| v_{th}}{\sqrt{2}} (\sqrt{m+1} \tilde{g}_{m+1} - \sqrt{m} \tilde{g}_{m-1}) = -\nu m \tilde{g}_m$$

\Downarrow

$$\boxed{\frac{\partial \tilde{g}_m^{\pm}}{\partial t} \pm \sqrt{2} |k_{\parallel}| v_{th} m^{1/4} \frac{\partial}{\partial m} m^{1/4} \tilde{g}_m^{\pm} = -\nu m \tilde{g}_m^{\pm}}$$

For $1 \ll m \ll m_c$, to lowest order,

$$\sqrt{m+1} \tilde{g}_{m+1} - \sqrt{m} \tilde{g}_{m-1} = 0 \Rightarrow \tilde{g}_{m+1} \approx \tilde{g}_{m-1}$$

This allows two solutions: $\tilde{g}_{m+1} \approx \pm \tilde{g}_m$, so either \tilde{g}_m or $(-1)^m \tilde{g}_m$ is continuous.

This can be encoded in the following decomposition:

$$\boxed{\tilde{g}_m = \tilde{g}_m^+ + (-1)^m \tilde{g}_m^-}$$

where $\tilde{g}_m^+ = \frac{\tilde{g}_m + \tilde{g}_{m+1}}{2}$ and $\tilde{g}_m^- = (-1)^m \frac{\tilde{g}_m - \tilde{g}_{m+1}}{2}$ are continuous in m .

propagates
from low to high m
(phase mixing)

propagates
from high to low m
(un-phase-mixing!)

[AAS et al., arXiv:1508.05988]



“Un-phase-mixing”

$$\frac{\partial \tilde{g}_m}{\partial t} + \frac{|k_{\parallel}| v_{th}}{\sqrt{2}} (\sqrt{m+1} \tilde{g}_{m+1} - \sqrt{m} \tilde{g}_{m-1}) = -\nu m \tilde{g}_m$$

↓

$$\boxed{\frac{\partial \tilde{g}_m^{\pm}}{\partial t} \pm \sqrt{2} |k_{\parallel}| v_{th} m^{1/4} \frac{\partial}{\partial m} m^{1/4} \tilde{g}_m^{\pm} = -\nu m \tilde{g}_m^{\pm}}$$

In energy terms: $C_m = C_m^+ + C_m^-$ satisfies

$$\frac{\partial C_m}{\partial t} + \frac{\partial}{\partial m} |k_{\parallel}| v_{th} \sqrt{2m} (C_m^+ - C_m^-) = -2\nu m C_m$$

Hermite flux to high m can be
cancelled (on average) by the ‘-’ modes

$$\boxed{\tilde{g}_m = \tilde{g}_m^+ + (-1)^m \tilde{g}_m^-}$$

where $\tilde{g}_m^+ = \frac{\tilde{g}_m + \tilde{g}_{m+1}}{2}$ and $\tilde{g}_m^- = (-1)^m \frac{\tilde{g}_m - \tilde{g}_{m+1}}{2}$ are continuous in m .

propagates
from low to high m
(phase mixing)

propagates
from high to low m
(un-phase-mixing!)

[AAS et al., arXiv:1508.05988]



“Un-phase-mixing”

$$\frac{\partial \tilde{g}_m}{\partial t} + \frac{|k_{\parallel}| v_{th}}{\sqrt{2}} (\sqrt{m+1} \tilde{g}_{m+1} - \sqrt{m} \tilde{g}_{m-1}) = -\nu m \tilde{g}_m$$

↓

$$\frac{\partial \tilde{g}_m^{\pm}}{\partial t} \pm \sqrt{2} |k_{\parallel}| v_{th} m^{1/4} \frac{\partial}{\partial m} m^{1/4} \tilde{g}_m^{\pm} = -\nu m \tilde{g}_m^{\pm}$$

In energy terms: $C_m = C_m^+ + C_m^-$ satisfies

$$\frac{\partial C_m}{\partial t} + \frac{\partial}{\partial m} |k_{\parallel}| v_{th} \sqrt{2m} (C_m^+ - C_m^-) = -2\nu m C_m$$

Hermite flux to high m can be cancelled (on average) by the ‘-’ modes

$$\tilde{g}_m = \tilde{g}_m^+ + (-1)^m \tilde{g}_m^-$$

where $\tilde{g}_m^+ = \frac{\tilde{g}_m + \tilde{g}_{m+1}}{2}$ and $\tilde{g}_m^- = (-1)^m \frac{\tilde{g}_m - \tilde{g}_{m+1}}{2}$ are continuous in m .

propagates
from low to high m
(phase mixing)

propagates
from high to low m
(un-phase-mixing!)

Linearly, none of this happens because there are no energy sources at high m and to satisfy

$$\tilde{g}_{m \rightarrow \infty} \rightarrow 0,$$

we must have

$$\tilde{g}_m^- = 0$$

Time to bring back nonlinearity...



“Un-phase-mixing”

Restore nonlinearity:

$$\left(\frac{\partial g_m}{\partial t}\right)_{\text{nl}} = -[\mathbf{u}_{\perp} \cdot \nabla_{\perp} g_m](k_{\parallel}) = - \sum_{p_{\parallel} + q_{\parallel} = k_{\parallel}} \mathbf{u}_{\perp}(p_{\parallel}) \cdot \nabla_{\perp} g_m(q_{\parallel})$$

For $\tilde{g}_m = (i \operatorname{sgn} k_{\parallel})^m g_m$, the nonlinearity becomes

$$\left(\frac{\partial \tilde{g}_m}{\partial t}\right)_{\text{nl}} = -(i \operatorname{sgn} k_{\parallel})^m [\mathbf{u}_{\perp} \cdot \nabla_{\perp} g_m](k_{\parallel}) = - \sum_{p_{\parallel} + q_{\parallel} = k_{\parallel}} \mathbf{u}_{\perp}(p_{\parallel}) \cdot \nabla_{\perp} \frac{(i \operatorname{sgn} k_{\parallel})^m}{(i \operatorname{sgn} q_{\parallel})^m} \tilde{g}_m(q_{\parallel})$$

And for the ‘+’ and ‘−’ modes (add/subtract the above for m and $m+1$),

$$\left(\frac{\partial \tilde{g}_m^{\pm}}{\partial t}\right)_{\text{nl}} = - \sum_{p_{\parallel} + q_{\parallel} = k_{\parallel}} \mathbf{u}_{\perp}(p_{\parallel}) \cdot \nabla_{\perp} \left[\underset{\substack{\uparrow \\ 1 \text{ if } k_{\parallel} \text{ and } q_{\parallel} \\ \text{have same sign} \\ 0 \text{ otherwise}}}{\delta_{k_{\parallel}, q_{\parallel}}^{+}} \tilde{g}_m^{\pm}(q_{\parallel}) + \delta_{k_{\parallel}, q_{\parallel}}^{-} \tilde{g}_m^{\mp}(q_{\parallel}) \right]$$

$\underset{\substack{\uparrow \\ 1 \text{ if } k_{\parallel} \text{ and } q_{\parallel} \\ \text{have opposite sign} \\ 0 \text{ otherwise}}}{\delta_{k_{\parallel}, q_{\parallel}}^{-}}$

Plasma Echo



$$\frac{\partial \tilde{g}_m^\pm}{\partial t} \pm \sqrt{2}|k_\parallel|v_{th}m^{1/4}\frac{\partial}{\partial m}m^{1/4}\tilde{g}_m^\pm = -\nu m\tilde{g}_m^\pm$$

$$- \sum_{p_\parallel + q_\parallel = k_\parallel} \mathbf{u}_\perp(p_\parallel) \cdot \nabla_\perp \left[\delta_{k_\parallel, q_\parallel}^+ \tilde{g}_m^\pm(q_\parallel) + \delta_{k_\parallel, q_\parallel}^- \tilde{g}_m^\mp(q_\parallel) \right]$$

\uparrow
 1 if k_\parallel and q_\parallel
 have same sign
 0 otherwise

\uparrow
 1 if k_\parallel and q_\parallel
 have opposite sign
 0 otherwise

\uparrow
‘+’ and ‘-’ modes couple!

$\tilde{g}_m^- = 0$ is no longer a solution.

Free energy can come back from phase space!

[AAS et al., arXiv:1508.05988]

[cf. Hammett et al. 1993]



Plasma Echo

A very compact form of the Hermite-space equation is achieved by defining

$$f = m^{1/4} \begin{cases} \tilde{g}_m^+ & \text{for } k_{\parallel} \geq 0, \\ \tilde{g}_m^- & \text{for } k_{\parallel} < 0 \end{cases} \quad \text{and} \quad s = \sqrt{m}$$

This is a bit like a Fourier transform, with $iv_{\parallel} \sim \partial_s$

$$\frac{\partial f}{\partial t} + \frac{k_{\parallel} v_{th}}{\sqrt{2}} \frac{\partial f}{\partial s} + \nu s^2 f = - \sum_{p_{\parallel}} \mathbf{u}_{\perp}(p_{\parallel}) \cdot \nabla_{\perp} f(k_{\parallel} - p_{\parallel})$$

$$\frac{\partial \tilde{g}_m^{\pm}}{\partial t} \pm \sqrt{2} |k_{\parallel}| v_{th} m^{1/4} \frac{\partial}{\partial m} m^{1/4} \tilde{g}_m^{\pm} = -\nu m \tilde{g}_m^{\pm}$$

$$- \sum_{p_{\parallel} + q_{\parallel} = k_{\parallel}} \mathbf{u}_{\perp}(p_{\parallel}) \cdot \nabla_{\perp} \left[\delta_{k_{\parallel}, q_{\parallel}}^+ \tilde{g}_m^{\pm}(q_{\parallel}) + \delta_{k_{\parallel}, q_{\parallel}}^- \tilde{g}_m^{\mp}(q_{\parallel}) \right]$$

\uparrow \uparrow
 1 if k_{\parallel} and q_{\parallel} have same sign 1 if k_{\parallel} and q_{\parallel} have opposite sign
 0 otherwise 0 otherwise

‘+’ and ‘-’ modes couple!

$\tilde{g}_m^- = 0$ is no longer a solution.

Free energy can come back from phase space!

[AAS et al., arXiv:1508.05988]

[cf. Hammett et al. 1993]



Plasma Echo

A very compact form of the Hermite-space equation is achieved by defining

$$f = m^{1/4} \begin{cases} \tilde{g}_m^+ & \text{for } k_{\parallel} \geq 0, \\ \tilde{g}_m^- & \text{for } k_{\parallel} < 0 \end{cases} \quad \text{and} \quad s = \sqrt{m}$$

This is a bit like a Fourier transform, with $iv_{\parallel} \sim \partial_s$

$$\frac{\partial f}{\partial t} + \frac{k_{\parallel} v_{th}}{\sqrt{2}} \frac{\partial f}{\partial s} + \nu s^2 f = - \sum_{p_{\parallel}} \mathbf{u}_{\perp}(p_{\parallel}) \cdot \nabla_{\perp} f(k_{\parallel} - p_{\parallel})$$

A phase-mixing perturbation can turn around and come back (un-phase-mix) if the advecting velocity couples it to a parallel wave number of opposite sign – **plasma echo** effect.




Plasma Echo

A very compact form of the Hermite-space equation is achieved by defining

$$f = m^{1/4} \begin{cases} \tilde{g}_m^+ & \text{for } k_{\parallel} \geq 0, \\ \tilde{g}_m^- & \text{for } k_{\parallel} < 0 \end{cases} \quad \text{and} \quad s = \sqrt{m}$$

This is a bit like a Fourier transform, with $iv_{\parallel} \sim \partial_s$

$$\frac{\partial f}{\partial t} + \frac{k_{\parallel} v_{th}}{\sqrt{2}} \frac{\partial f}{\partial s} + \nu s^2 f = - \sum_{p_{\parallel}} \mathbf{u}_{\perp}(p_{\parallel}) \cdot \nabla_{\perp} f(k_{\parallel} - p_{\parallel})$$



A phase-mixing perturbation can turn around and come back (un-phase-mix) if the advecting velocity couples it to a parallel wave number of opposite sign – **plasma echo** effect.

...And it does indeed do it! Here are some plots of the relative Hermite flux

$$\frac{C_m^+ - C_m^-}{C_m^+ + C_m^-}$$

from a drift-kinetic slab ITG simulation by **J. Parker & E. Highcock**



Phase Mixing vs. Turbulence

A very compact form of the Hermite-space equation is achieved by defining

$$f = m^{1/4} \begin{cases} \tilde{g}_m^+ & \text{for } k_{\parallel} \geq 0, \\ \tilde{g}_m^- & \text{for } k_{\parallel} < 0 \end{cases} \quad \text{and} \quad s = \sqrt{m}$$

This is a bit like a Fourier transform, with $iv_{\parallel} \sim \partial_s$

$$\frac{\partial f}{\partial t} + \frac{k_{\parallel} v_{th}}{\sqrt{2}} \frac{\partial f}{\partial s} + \nu s^2 f = - \sum_{p_{\parallel}} \mathbf{u}_{\perp}(p_{\parallel}) \cdot \nabla_{\perp} f(k_{\parallel} - p_{\parallel})$$

The perturbation at low s , $f(s \sim 1) \sim \varphi$, and at some fixed k_{\perp} and k_{\parallel} , will propagate to higher s along the characteristic:

$$s \sim k_{\parallel} v_{th} t,$$


until it is swept by nonlinear advection (\mathbf{u}_{\perp}) to higher k_{\perp} in one nonlinear time,

$$t \sim (k_{\perp} u_{\perp})^{-1} \propto k_{\perp}^{-4/3}.$$

Thus, $f(s) \sim \varphi$ for $s \lesssim \frac{k_{\parallel} v_{th}}{k_{\perp} u_{\perp}}$, or, equivalently, for the phase-space spectrum:

$$E_m(k_{\perp}, k_{\parallel}) = 2\pi k_{\perp} \langle |g_m|^2 \rangle \sim \frac{E_{\varphi}(k_{\perp}, k_{\parallel})}{\sqrt{m}} \quad \text{for } k_{\parallel} \gtrsim \frac{k_{\perp} u_{\perp}}{v_{th}} \sqrt{m} \propto k_{\perp}^{4/3} \sqrt{m}$$

Phase-Space Spectra



Phase-Space Spectra



Phase-mixing region:
everything is linear, **no echo**

$$E_m^+ \sim \frac{k_\perp^3 k_\parallel^{-5}}{\sqrt{m}}$$

phase-mixing threshold $k_\parallel v_{th} \sim \sqrt{m} k_\perp^{4/3}$

✓ By pure kinematics of
correlation functions, in 2D,

$$E_m \sim \text{const } k_\perp^3 + \dots \text{ as } k_\perp \rightarrow 0$$

✓ Parallel exponent fixed by **matching**
at the phase-mixing threshold

Phase-mixing term
dominant

$$\frac{\partial f}{\partial t} + \frac{k_\parallel v_{th}}{\sqrt{2}} \frac{\partial f}{\partial s} + \cancel{\nu s^2} f = - \sum_{p_\parallel} \mathbf{u}_\perp(p_\parallel) \cdot \nabla_\perp f(k_\parallel - p_\parallel)$$

Phase-Space Spectra

Phase-mixing region:
everything is linear, **no echo**


$$E_m^+ \sim \frac{k_\perp^3 k_\parallel^{-5}}{\sqrt{m}}$$

phase-mixing threshold $k_\parallel v_{th} \sim \sqrt{m} k_\perp^{4/3}$

✓ By pure kinematics of
correlation functions, in 2D,

$$E_m \sim \text{const } k_\perp^3 + \dots \text{ as } k_\perp \rightarrow 0$$

✓ Parallel exponent fixed by **matching**
at the phase-mixing threshold



Phase-mixing term
dominant


$$\frac{\partial f}{\partial t} + \frac{k_\parallel v_{th}}{\sqrt{2}} \frac{\partial f}{\partial s} + \cancel{\nu s^2} f = - \sum_{p_\parallel} \mathbf{u}_\perp(p_\parallel) \cdot \nabla_\perp f(k_\parallel - p_\parallel)$$

“phase-space critical balance”


found numerically

by Hatch et al. JPP **80**, 531 (2014)


Phase-Space Spectra




Phase-Space Spectra




Phase-Space Spectra






Phase-Space Spectra





Phase-Space Spectra





Phase-Space Spectra






Phase-Space Spectra



Energy Flows



Energy Flows



Conclusions


- ✓ At $k_{\parallel} \gtrsim k_{\perp}^{4/3} \sqrt{m}$, linear phase mixing dominates, $E_m \propto \frac{1}{\sqrt{m}}$, but there is very little energy ($\sim k_{\parallel}^{-5}$)
- ✓ At $k_{\parallel} \lesssim k_{\perp}^{4/3} \sqrt{m}$, nonlinear mixing (turbulence) dominates, $E_m \propto \frac{1}{m^{5/2}}$, most energy is there, but collisional dissipation $\rightarrow 0$ as $\nu \rightarrow 0$;
total free energy stored in phase space is finite and independent of collisionality

$$\sum_m E_m \rightarrow \text{const}, \quad \nu \sum_m m E_m \rightarrow 0 \text{ as } \nu \rightarrow +0$$

*This means spatial mixing
("turbulence")
wins over phase mixing*


In contrast, in the linear problem,

$$\sum_m E_m \sim \nu^{-1/3}, \quad \nu \sum_m m E_m \rightarrow \text{const as } \nu \rightarrow +0$$



Conclusions


- ✓ At $k_{\parallel} \gtrsim k_{\perp}^{4/3} \sqrt{m}$, linear phase mixing dominates, $E_m \propto \frac{1}{\sqrt{m}}$, but there is very little energy ($\sim k_{\parallel}^{-5}$)
- ✓ At $k_{\parallel} \lesssim k_{\perp}^{4/3} \sqrt{m}$, nonlinear mixing (turbulence) dominates, $E_m \propto \frac{1}{m^{5/2}}$, most energy is there, but collisional dissipation $\rightarrow 0$ as $\nu \rightarrow 0$;
total free energy stored in phase space is finite and independent of collisionality



Conclusions


- ✓ At $k_{\parallel} \gtrsim k_{\perp}^{4/3} \sqrt{m}$, linear phase mixing dominates, $E_m \propto \frac{1}{\sqrt{m}}$, but there is very little energy ($\sim k_{\parallel}^{-5}$)
- ✓ At $k_{\parallel} \lesssim k_{\perp}^{4/3} \sqrt{m}$, nonlinear mixing (turbulence) dominates, $E_m \propto \frac{1}{m^{5/2}}$, most energy is there, but collisional dissipation $\rightarrow 0$ as $\nu \rightarrow 0$;
total free energy stored in phase space is finite and independent of collisionality

from a drift-kinetic
slab ITG simulation
by **J. Parker**
& **E. Highcock:**



Conclusions

- ✓ At $k_{\parallel} \gtrsim k_{\perp}^{4/3} \sqrt{m}$, linear phase mixing dominates, $E_m \propto \frac{1}{\sqrt{m}}$, but there is very little energy ($\sim k_{\parallel}^{-5}$)
- ✓ At $k_{\parallel} \lesssim k_{\perp}^{4/3} \sqrt{m}$, nonlinear mixing (turbulence) dominates, $E_m \propto \frac{1}{m^{5/2}}$, most energy is there, but collisional dissipation $\rightarrow 0$ as $\nu \rightarrow 0$; total free energy stored in phase space is finite and independent of collisionality



- ✓ Return echo flux cancels phase-mixing flux at $k_{\parallel} \lesssim k_{\perp}^{4/3}$ (below critical balance); turbulent cascade of low Hermite moments is effectively fluid... we might say that, for the purposes of free-energy accounting in turbulence, “Landau damping is suppressed”

[AAS et al., arXiv:1508.05988]

All of the arguments presented above rely on the approximation of

$$m \gg 1 \text{ and, indeed, } m^{1/4} \gg 1,$$

i.e., truly asymptotically small collisionality (= a lot of velocity-space structure).

In reality (experimental and certainly numerical), the collisionality or effective collisionality (in codes) is rarely truly small. When it is moderate and only relatively little Hermite space is available to the free energy, processes that require such space – most notably the echo flux – are likely to be less pronounced.

This probably accounts for how well Landau fluid closures have tended to capture quantitative behaviour of turbulence in tokamaks.

So perhaps (perhaps!) the scenario is

$\nu \gg \omega$ – collisional system, fluid

$\nu \lesssim \omega$ – weakly collisional system, “Landau-fluid”

$\nu \ll \omega$ – “collisionless” system, like fluid again?

In any event, this is quite a cute problem to think about...