Draft version October 27, 2026
Typeset using LaTeX **preprint** style in AASTeX7.0.2




# Physically Motivated Gaussian Processes for Stellar Variability I: A Starspot Kernel with Differential Rotation

Jessica Birky [1,2] Eric Agol [1] Rory Barnes [1] James Davenport [1] and Sean Matt [2]

[1] *Department of Astronomy, University of Washington, 3910 15th Avenue NE, Seattle, WA 98195, USA*
[2] *Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman, OK 73019, USA*

## ABSTRACT

We present a novel, physically motivated Gaussian process (GP) kernel designed to model starspot variability from photometric lightcurves. Using an analytic flux model, we derive an analytic kernel function that describes the temporal correlations induced by rotating starspots as a function of spot properties and stellar rotation: equitorial rotation period ($P_{\rm eq}$), differential rotation ($\kappa$), inclination angle ($I$), spot lifetimes ($\ell_{\rm spot}$), spot emergence and decay timescales ($\tau_{\rm spot}$), maximum spot size ($\alpha_{\rm max}$), spot contrast ($f_{\rm spot}$), and spot emergence rate ($\dot{N}_{\rm spot}$). We thoroughly validate our derivation using `SymPy` and compare our analytic solutions against to numerical simulations. Furthermore, we interpret how each parameter affects the kernel structure, showing that $\alpha_{\rm max}$, $f_{\rm spot}$, and $\dot{N}_{\rm spot}$ are degenerate parameters controlling the amplitude of the correlations (which can be re-parameterized into a single parameter $\sigma_k$), while $P_{\rm eq}$, $\kappa$, and $I$ control the periodicity and harmonic structure, and $\ell_{\rm spot}$, $\tau_{\rm em}$, and $\tau_{\rm dec}$ control the decay of the correlations over time. Our implementation is available in the package `spotgp`, which uses `JAX` for efficient automatic differentiation and GPU acceleration. Furthermore, our implementation makes use of a sparse banded matrix solver (with bandwidth $b$), reducing the computational complexity of GP inference from $\mathcal{O}(N^3)$ to $\mathcal{O}(Nb^2)$ for $N$ data points. Using simulated lightcurves, we use Markov chain Monte Carlo (MCMC) and Cramer-Rao bound analysis to systematically explore how inference performs for different parameter regimes, and observational conditions (cadence, baseline, and noise level). Finally, our simulated inference demonstrates that `spotgp` is a highly promising tool for extracting differential rotation and inclinations just from *individual* lightcurves of stars.

## 1. INTRODUCTION

High-precision photometric surveys such as *Kepler* (Borucki et al. 2010), *K2* (Howell et al. 2014), and *TESS* (Ricker et al. 2014) have been transforming our understanding of fundamental stellar physics, revealing the effects of stellar variability and dynamics for stars across a wide range of masses, ages, and Galactic environments. In particular, precise time-domain data have enabled constraints

Corresponding author: Jess Birky
jess@birky.net

on the rotation periods of thousands of stars (McQuillan et al. 2014; Gordon et al. 2021; Boyle et al. 2026), which has been crucial for understanding the evolution of stellar angular momentum (Weber & Davis 1967; Skumanich 1972), the surface variability and magnetic activity of stars (Santos et al. 2017; Forgács-Dajka et al. 2021), and the stellar dynamo process (Brun et al. 2017; Käpylä et al. 2023; Noraz et al. 2022b).

Over time, stars lose angular momentum through magnetized stellar winds, causing them to spin down as they age. The rate of this spin-down depends on the star's mass, age, and magnetic field strength, and is driven by the torque from stellar winds flowing along open magnetic field lines, a process known as *magnetic braking*. Early work established that $P_{\rm rot} \propto t^{-1/2}$ (Skumanich 1972), opening up the field of gyrochronology for inferring stellar ages from rotation periods (Barnes 2003, 2007). Further work has developed empirical relations for how magnetic braking torque depends on stellar properties (Kawaler 1988; Sills et al. 2000; Reiners & Mohanty 2012; van Saders & Pinsonneault 2013; Matt et al. 2015; Amard & Matt 2020; Ireland et al. 2021). However open questions remain: recent studies have discovered evidence for weakened magnetic braking in rapidly rotating old field stars (van Saders et al. 2016; Saunders et al. 2024), suggesting that the braking process may be more complex than originally thought.

Much of our knowledge of stellar rotation comes from photometric variability driven by starspots. Low-mass stars ($M \lesssim 1.5\,M_\odot$) possess outer convective envelopes in which turbulent motions interact with rotation and magnetic fields to drive a stellar dynamo (Brun et al. 2017). Differential rotation (the latitudinal variation in angular velocity) plays a fundamental role in this process: it shears poloidal magnetic field lines into a toroidal configuration (the $\Omega$-effect), which is a key mechanism driving the activity cycle (Noraz et al. 2022b). Numerical simulations of rotating convection predict that the magnitude and geometry of differential rotation depend on the stellar mass, rotation rate, and convective structure (Augustson et al. 2012; Guerrero et al. 2013; Käpylä et al. 2023). In particular, these simulations predict a transition from solar-like differential rotation (fast equator, slow poles) to anti-solar differential rotation (fast poles, slow equator) as the Rossby number $Ro = P_{\rm rot}/\tau_{\rm conv}$ increases past $Ro \sim 1$. Many starspot properties, including spot lifetimes (Giles et al. 2017), latitude distributions, and emergence patterns, are also governed by $Ro$, making it a highly important dimensionless parameter for characterizing the stellar dynamo regime (Gastine et al. 2013; Noraz et al. 2022a). However, this predicted transition in differential rotation remains largely untested observationally, because measuring differential rotation for large samples of stars has been extremely challenging.

Existing techniques for measuring stellar differential rotation include: Doppler imaging (Collier Cameron 2007), the Fourier transform method for spectral line profiles (Reiners & Mohanty 2012), asteroseismology (Benomar et al. 2018), and transit spot-crossing (Morris et al. 2017). However, each have each been limited to samples of ∼tens of stars, and are restricted to specific stellar types (e.g., fast rotators, bright stars, or transiting systems). On the other hand, the high-precision, long-baseline photometric datasets from *Kepler* and *TESS* encode information on the spot variability for hundreds of thousands of stars, which has yet to be fully leveraged. However, extracting differential rotation from disk-integrated photometry is notoriously difficult: Aigrain et al. (2015) conducted a blind hare-and-hounds exercise on 1,000 simulated *Kepler*-like lightcurves and found *little correlation between the reported and simulated values of the differential rotation shear*. Morris et al. (2019) reinforced this conclusion using the Sun itself as a benchmark, finding no detectable

differential rotation signal even in a noise-free solar lightcurve reconstructed from archival sunspot records. These results suggest that the stochastic nature of spot emergence at a broad range of latitudes causes the measured period to reflect an average over active latitudes, rather than revealing the latitude-dependent rotation law.

Modeling the magnetohydrodynamical processes that accurately reproduce the convection, magnetic fields, and dynamo process of low-mass stars is inherently complex and computationally expensive (Miesch et al. 2000, 2008; Charbonneau 2020). As an alternative, analytic spot models (Kipping 2012; Luger et al. 2019) treat the stellar surface as hosting an arbitrary number of circular spots that emerge and decay at different latitudes, each modulating the disk-integrated flux according to its projected area and contrast. These models have been successful in matching the statistical properties of observed lightcurves (Aigrain et al. 2015), while providing a physically interpretable framework for the spot geometry.

Although there are approaches for forward modeling surfaces with spots, inferring information about *individual* star spots is an ill-posed problem with many degeneracies (Luger et al. 2019; Basri & Shah 2020). Basri & Shah (2020) showed that the single-dip ratio (a metric for whether a lightcurve shows one or two dips per rotation) is only weakly sensitive to differential rotation shear and is strongly confounded by spot lifetime, concluding that differential rotation may be "nearly impossible to accurately infer from light curves alone unless spots live for many rotations." A Gaussian process (GP; Rasmussen & Williams 2006) offers an alternative approach that treats starspot variability as a stochastic process and models the *distribution* of spot properties rather than individual spots. GPs have become a standard tool for modeling stellar rotation (Angus et al. 2018; Foreman-Mackey et al. 2017; Gordon et al. 2021), but the most widely used quasi-periodic (QP) kernel has no parameter that encodes differential rotation, meaning that any differential rotation signal is absorbed into the other hyperparameters in an uncontrolled way. Nicholson & Aigrain 2022 systematically tested the mapping between QP kernel hyperparameters and physical spot properties, finding that while the rotation period and evolution timescale are reliably recovered, the harmonic complexity parameter has no intuitive physical interpretation. These results collectively motivate the development of a physically grounded GP kernel: one whose hyperparameters map directly to measurable stellar properties including differential rotation.

In this paper, we derive a physically motivated GP kernel for stellar rotation from first principles. Starting from an analytic starspot model (Kipping 2012), we obtain a closed-form kernel function that naturally encodes the effects of spot geometry, evolution, and differential rotation. We validate the analytic kernel against numerical Monte Carlo simulations, demonstrate that it captures harmonic structure and non-sinusoidal periodicity absent in standard quasi-periodic and simple harmonic oscillator kernels, and show that it provides significant computational speedups over numerical approaches. The paper is organized as follows: Section 2 describes the lightcurve model, Section 4 derives the GP kernel, Section 8 presents the `JAX` implementation of the kernel and MCMC inference framework, Section 9 validates the analytic kernel against numerical results, Section 11 discusses the physical interpretability of our results with intuitive visualizations, and the Appendix Section provides detailed derivations and `sympy` validation notebooks. A companion paper (J. Birky et al. 2026, in preparation; hereafter Paper II) extends the kernel with additive composite terms for several coexisting spot populations and for non-spot variability, and applies the full framework to the four-year *Kepler* lightcurve of Kepler-63.

## 2. LIGHTCURVE MODEL

We use the analytic starspot model defined in Kipping (2012). The flux model for $N_{\rm spot}$ spots with contrast $f_k$ is:

$$F(t) = 1 - \frac{1}{\pi}\sum_{k=1}^{N_{\rm spot}} A_k(t)\cdot(1-f_{\rm spot,k}), \tag{1}$$

where $A_k(t)$ is the projected area of spot $k$ at time $t$ and $f_{\rm spot,k}$ is the contrast of the spot (where $f_{\rm spot}=0$ represents a totally black spot, $0<f_{\rm spot}<1$ represents a partially dark spot, $f_{\rm spot}=1$ represents no contrast, and $f_{\rm spot}>1$ represents a bright spot or stellar faculae). The flux deficit at a given time is the sum of the contributions from all visible spots on the surface of the star, where each contribution is determined by the projected area of the spot and its contrast relative to the stellar photosphere.

### 2.1. *Single-Spot Projected Area*

For each individual spot, the flux deficit is determined by the projected area of the spot on the visible hemisphere of the star. For ease of notation we drop the spot index $k$ and write the projected area of a single spot as $A(t)$, however these equations are applied to each spot $k$ with its own time-dependent area $A_k(t)$ and angle $\beta_k(t)$. This projected area of starspot $k$ at time $t$ is given by:

$$A_k(t) = \mathbb{R}\Big[\cos^{-1}[\cos\alpha_k\csc\beta_k] + \cos\beta_k\sin\alpha_k\Xi_k - \cos\alpha_k\sin\beta_k\Psi_k\Big] \tag{2}$$

where the intermediate terms are

$$\Xi_k = \sin\alpha_k\cos^{-1}[-\cot\alpha_k\cot\beta_k], \tag{3}$$

$$\Psi_k = \sqrt{1-\cos^2\alpha_k\csc^2\beta_k} \tag{4}$$

Expanded out, $A(t)$ becomes a piecewise expression with three cases:

$$A_k(t) = \begin{cases} \pi\sin^2\alpha_k\cos\beta_k & (0<\beta_k<\pi/2-\alpha_k) \\ \cos^{-1}[\cos\alpha_k\csc\beta_k] + \cos\beta_k\sin^2\alpha_k\cos^{-1}[-\cot\alpha_k\cot\beta_k] & \\ \qquad -\cos\alpha_k\sin\beta_k\sqrt{1-\cos^2\alpha_k\csc^2\beta_k} & (\pi/2-\alpha_k<\beta_k<\pi/2+\alpha_k) \\ 0 & (\pi/2+\alpha_k<\beta_k<\pi) \end{cases} \tag{5}$$

where $\alpha_k$ is the angular radius of the spot and $\beta_k$ is the angle between the spot normal and the line of sight (Kipping 2012). The three cases correspond to distinct geometric configurations of the spot on the stellar disk: fully visible, partiay occulted by the limb, and fully hidden on the far side, as shown in Figure 1. Paper II presents a self-contained derivation of Eq. (5) as a surface integral over the spot cap.

The angle between spot normal and the line of sight for a single spot at a given time is:

$$\cos\beta(t) = \cos I\sin\Phi + \sin I\cos\Phi\cos\Lambda(t), \tag{6}$$

where $I$ is the inclination of the stellar rotation axis, $\Phi$ is the latitude of the spot, and $\Lambda(t)$ is the longitude of the spot as a function of time. We emphasize the distinction between the two frames





**Figure 1.** Illustration of the three geometric regimes for the projected area of a circular starspot on a rotating star. Each panel shows the stellar disk as seen by the observer, with latitude and longitude grid lines, the spot (dark red), the angular radius $\alpha_k$, and the angle $\beta_k$ between the spot normal and the line of sight. **Left:** The spot is fully visible on the near hemisphere. **Center:** The spot straddles the stellar limb and is partially occulted. **Right:** The spot is on the far side and hidden from view.

related by Eq. (6): $\Phi$ is a *stellar* coordinate, measured from the stellar equator in the rotation frame, with $\Phi \in [-\pi/2, \pi/2]$ and $\Phi = +\pi/2$ the pole inclined toward the observer; $I \in [0, \pi/2]$ is the angle between the rotation axis and the line of sight, so that the sub-observer point lies at latitude $\pi/2 - I$. Defining $\Phi$ in the rotation frame is what makes the differential rotation law $\omega_0(\Phi)$ (Eq. 59) well posed, since latitudinal shear is an intrinsic property of the star and cannot depend on the viewing geometry. Consequently the inclination enters the model not through the coordinate $\Phi$ itself, but through the observer-dependent visibility mask it imposes on that coordinate: since $\cos\beta$ ranges over $[\sin(\Phi - I), \sin(\Phi + I)]$ as the star rotates, spots at $\Phi \geq I$ are circumpolar and permanently visible, spots at $\Phi \leq -I$ never rotate into view, and only the band $|\Phi| < I$ produces rotationally modulated flux. The time dependence of $\beta(t)$ arises from the rotation of the star, which causes the spot to move in longitude and therefore change its angle relative to the line of sight. The projected area $A(t)$ is then determined by both $\alpha(t)$ and $\beta(t)$ according to the piecewise function in Eqn 5.

When the spot angular radius $\alpha_k$ is small compared to 1 radian (typical for Sun-like stars where spots subtend $\lesssim 6°$), we can make the approximation $\sin\alpha_k \approx \alpha_k$, so $\sin^2\alpha_k \approx \alpha_k^2$. In this limit:

- The **fully visible** case simplifies to $A_k \approx \pi\alpha_k^2 \cos\beta_k$.
- The **partial-visibility window** has width $2\alpha_k$ and becomes vanishingly narrow, so the spot transitions nearly instantaneously from fully visible to fully hidden as it crosses the limb.
- The **geometric projection effect** is purely $\cos\beta$: the same projection factor as a flat disk viewed at angle $\beta$. Higher-order corrections from spherical curvature (the $\Xi_k$, $\Psi_k$ terms) vanish at leading order in $\alpha_k$.

This allows us to replace the entire three-case piecewise function with a single expression:

$$A(t) \approx \pi\,\alpha^2(t)\ \cdot\ \max\{\cos\beta(t),\ 0\} \tag{7}$$

where $\alpha(t)$ is the angular radius of the spot (whose time evolution is described in Section 2.2). Figure 2 illustrates this approximation. The left panel shows the size of spots with three different

angular radii $\alpha_{\rm max}$ on the stellar disk at $\beta = 0$ (disk center), providing a visual reference for the spot-to-star size ratio. The right panel compares the exact projected area $A(\beta)$ from Eq. (5) (solid lines) with the small-spot approximation $\max\{\cos\beta, 0\}$ (dashed line). For small spots ($\alpha_{\rm max} \lesssim 15°$), the exact and approximate curves are nearly indistinguishable, validating the factorization. For larger spots ($\alpha_{\rm max} \sim 30°$), the transition at the limb ($\beta = \pi/2$) becomes more gradual in the exact solution, but the approximation remains reasonable.





**Figure 2.** *Left:* Stellar disk viewed face-on, with circles indicating the projected size of spots with angular radii $\alpha_{\rm max} = 5°$, 15°, and 30° centered at $\beta = 0$ (disk center). *Right:* Normalized projected area $A(\beta)/A(0)$ as a function of the angle $\beta$ between the spot normal and the line of sight. Solid colored lines show the exact piecewise expression (Eq. 5) for each $\alpha_{\rm max}$; the dashed black line shows the small-spot approximation $\max\{\cos\beta, 0\}$ (Eq. 7). The two converge for $\alpha_{\rm max} \lesssim 15°$.

The key consequence for the kernel derivation is that the projected area now *factorizes* into a product of two functions:

$$A(t) \propto \Gamma(t) \cdot \mathcal{V}(t), \tag{8}$$

where we define $\Gamma(t) \equiv \alpha^2(t)/\alpha^2_{\rm max}$ as the normalized squared spot-size envelope (with unit peak), and $\mathcal{V}(t) \equiv \max\{\cos\beta(t), 0\}$ as the spot visibility function. Figure 3 illustrates this decomposition. In this paper we proceed to derive the GP kernel under the simplifying assumptions of small spot approximation and negligible limb darkening. These assumptions allow us to solve for an analytic, closed-form solution for the kernel function, however we will refer the reader to Paper II for the numerical approach for how to compute the full visibility function including these terms.

### 2.2. *The Emergence and Decay Profile of Spots*

In this work, we assume that the angular size of the spot, $\alpha(t)$, evolves over time as the spot emerges and decays. Here we test three different parameterizations of the spot evolution: a piecewise linear function, an exponential growth/decay function, and a skew-normal function. In each case, the spot reaches its maximum angular size $\alpha_{\rm max}$ near time $t_{\rm ref}$, which is the reference time for the spot. In the first two, the spot emerges over a timescale $\tau_{\rm em}$ and decays over a timescale $\tau_{\rm dec}$; in the skew-normal profile, the asymmetry between emergence and decay is instead set by a single skewness parameter.

**Trapezoidal emergence/decay profile:** Following Kipping (2012), we define a piecewise linear function with an emergence timescale $\tau_{\rm em}$, a plateau duration $\ell_{\rm spot}$, and a decay timescale $\tau_{\rm dec}$. The spot holds its maximum angular size $\alpha_{\rm max}$ over a plateau centered on $t_{\rm ref}$, which is the reference time for the spot. The Heaviside step function $\mathsf{H}(x)$ is used to ensure that the spot only contributes to the flux deficit during its emergence and decay phases. The spot angular radius is given by:

$$\frac{\alpha_{\rm trap}(t)}{\alpha_{\rm max}} = \mathcal{I}^{-1}[\Delta t_1 \mathsf{H}(\Delta t_1) - \Delta t_2 \mathsf{H}(\Delta t_2)] - \mathcal{E}^{-1}[\Delta t_3 \mathsf{H}(\Delta t_3) - \Delta t_4 \mathsf{H}(\Delta t_4)], \tag{9}$$

where $\Delta t_1 = t - (t_{\rm ref} - \ell_{\rm spot}/2 - \tau_{\rm em})$, $\Delta t_2 = t - (t_{\rm ref} - \ell_{\rm spot}/2)$, $\Delta t_3 = t - (t_{\rm ref} + \ell_{\rm spot}/2)$, and $\Delta t_4 = t - (t_{\rm ref} + \ell_{\rm spot}/2 + \tau_{\rm dec})$. The normalization factors $\mathcal{I} = \tau_{\rm em}$ and $\mathcal{E} = \tau_{\rm dec}$ are the maximum values of the two bracketed ramp functions, which ensures that the peak of the trapezoidal function is normalized to 1, so that $\alpha_{\rm trap}(t)$ reaches $\alpha_{\rm max}$ at its maximum. The trapezoidal profile is a simple, piecewise linear model that captures the general shape of spot emergence and decay, with a linear rise to maximum size, a plateau, and a linear decay back to zero. For equal emergence and decay timescales, $\tau_{\rm em} = \tau_{\rm dec} \equiv \tau_{\rm spot}$, it reduces to the symmetric trapezoid of Eq. (C11) (with $t_{\rm ref} = 0$).

**Exponential emergence/decay profile:** An alternative, model for the spot-size evolution (similar to Aigrain et al. 2015) uses exponential emergence and decay rather than linear ramps. The spot angular radius is:

$$\alpha_{\rm exp}(t) = \alpha_{\rm max} \times \begin{cases} e^{(t-t_1)/\tau_{\rm em}} & t < t_1 \quad \text{(exponential rise)} \\ 1 & t_1 \le t \le t_2 \quad \text{(plateau)} \\ e^{-(t-t_2)/\tau_{\rm dec}} & t > t_2 \quad \text{(exponential decay)} \end{cases} \tag{10}$$

where $t_1$ and $t_2 = t_1 + \tau_{\rm plat}$ mark the start and end of the plateau phase. The three parameters are: $\tau_{\rm em}$ (emergence $e$-folding time), $\tau_{\rm plat}$ (plateau duration), and $\tau_{\rm dec}$ (decay $e$-folding time). Without loss of generality we center the envelope by setting $t_1 = 0$, $t_2 = \tau_{\rm plat}$.

**Skew-normal emergence/decay profile:** Forgács-Dajka et al. (2021) modeled sunspot group area evolution using a skew-normal function (their Eq. 1), which captures the observed asymmetry between rapid emergence and slower decay with a single smooth profile. Since the projected spot area scales as $\alpha^2$ in the small-spot limit, we apply the skew-normal to the squared angular radius:

$$\alpha_{\rm sn}(t) = \alpha_{\rm max}\sqrt{\frac{S(t - t_{\rm ref})}{S_{\rm max}}}, \qquad S(t) = \exp\left(-\frac{t^2}{2\sigma_{\rm sn}^2}\right)\left[1 + \mathrm{erf}\left(\frac{n_{\rm sn}\, t}{\sigma_{\rm sn}\sqrt{2}}\right)\right], \tag{11}$$

where $S_{\rm max}$ is the peak value of $S(t)$, so that $\alpha_{\rm sn}(t)$ reaches $\alpha_{\rm max}$ at its maximum. The two parameters are $\sigma_{\rm sn}$ (the scale parameter in days, controlling the overall spot lifetime) and $n_{\rm sn}$ (the dimensionless skewness parameter). When $n_{\rm sn} = 0$ the profile reduces to a symmetric Gaussian; $n_{\rm sn} > 0$ gives rapid emergence and slow decay (typical of sunspots), while $n_{\rm sn} < 0$ gives the opposite asymmetry. Unlike the trapezoidal and exponential profiles, the skew-normal profile has no plateau, and its peak is offset from $t_{\rm ref}$ when $n_{\rm sn} \neq 0$.

### 2.3. *Lattitudinal and Longitudinal Evolution of Spots*

The position of a starspot on the surface is described by its latitude ($\Phi$) and longitude ($\Lambda$). Similarly to the Sun, the latitude of the spot is assumed to be constant over time, while the longitude evolves





**Figure 3.** Decomposition of the single-spot projected area signal $A(t)$ into the normalized squared spot-size envelope $\Gamma(t) = \alpha^2(t)/\alpha^2_{\max}$ (top) and the visibility function $\mathcal{V}(t) = \max\{\cos\beta(t), 0\}$ (middle), with their product $A(t) \propto \Gamma(t) \cdot \mathcal{V}(t)$ (bottom). *Left:* trapezoidal envelope (Eq. 9). *Middle:* exponential envelope (Eq. 10). *Right:* skew-normal envelope (Eq. 11) with $n_{\rm sn} > 0$ (rapid emergence, slow decay). All three use the same rotation period $P = 3\,\mathrm{d}$, spot latitude $\Phi = 30°$, and edge-on inclination $I = 90°$.

due to the rotation of the star:

$$\Phi = \Phi_{\rm ref}, \tag{12}$$

$$\Lambda(t) = \Lambda_{\rm ref} + \frac{2\pi(t - t_{\rm ref})}{P(\Phi)}, \tag{13}$$

We assume a Sun-like differential rotation profile, where the rotation period $P$ at latitude $\Phi_k$ depends on the latitudinal shear $\kappa$ and the rotation period at the equator of the star $P_{\rm eq}$:

$$P(\Phi_k) = \frac{P_{\rm eq}}{1 - \kappa \sin^2 \Phi_k}, \tag{14}$$

where $\kappa = 0$ corresponds to solid body rotation and $\kappa > 0$ corresponds to solar-like differential rotation (equator rotates faster than poles), while $\kappa < 0$ corresponds to anti-solar differential rotation (poles rotate faster than equator). Or in other words, $\kappa$ quantifies the relative difference in rotation rate between the equator and the poles, $\kappa = (\Omega_{\rm eq} - \Omega_{\rm pole})/\Omega_{\rm eq}$ for equatorial rotation rate $\Omega_{\rm eq} = 2\pi/P_{\rm eq}$ and polar rotation rate $\Omega_{\rm pole} = 2\pi/P_{\rm pole}$. Figure 4 illustrates the effect of differential rotation on the rotational velocity at different latitudes on the stellar surface.





**Figure 4.** Illustration of differential rotation on a stellar sphere viewed at inclination $I = 70°$. Arrows show the rotational velocity at each latitude, with length proportional to the angular velocity $\omega(\Phi) = 2\pi/P(\Phi)$. *Left:* Solar-like differential rotation ($\kappa = 1.0$), where the equator rotates faster than the poles. *Right:* Anti-solar differential rotation ($\kappa = -1.0$), where the poles rotate faster than the equator.

### 2.4. *Summary of Parameters*

Tables 1 and 2 summarize the physical parameters of the lightcurve model. Table 1 lists the stellar parameters that are held fixed for all spots on the star, while Table 2 lists the parameters that can vary for each individual spot.

| *Star variables* | |
|---|---|
| $P_{\rm eq}$ | rotation period at equator |
| $\kappa$ | latitudinal differential rotation shear |
| $I$ | inclination angle between the rotation axis and the line of sight $(0, \pi/2)$ |
| $N_{\rm spot}$ | total number of spots observed within the simulated time $T_{\rm obs}$ |
| $l_{\rm spot}$ | lifetime of each spot |
| $\tau_{\rm spot}$ | emergence and decay timescale of each spot |

**Table 1**. Stellar parameters and their definitions.

| *Individual spot variables* | |
|---|---|
| $k$ | index number for each spot |
| $A_k(t)$ | projected area of star spot |
| $\beta_k(t)$ | angle between spot normal and line of sight |
| $\alpha_k(t)$ | angular size of spot (emerges and decays) |

| | |
|---|---|
| $\Lambda_k(t)$ | starspot longitude (0, $2\pi$) |
| $\Lambda_{\rm ref,k}$ | reference longitude of spot (0, $2\pi$) |
| $\Phi_{\rm ref,k}$ | reference latitude of spot ($-\pi/2$, $\pi/2$) |
| $t_{\rm ref,k}$ | time at which spot reaches maximum size |
| $\alpha_{\rm max,k}$ | maximum angular size of spot |
| $f_{\rm spot,k}$ | spot-star contrast (0 = black spot, 1 = no contrast) |

**Table 2**. Individual spot parameters and their definitions.

Figure 5 shows an example lightcurve generated from the starspot model, with contributions from each individual spot shown in color and the cumulative flux deficit shown in black.





**Figure 5.** Examples of simulated lightcurves generated from the starspot model from the Kipping (2012) model. The top panel shows the case with solar-like differential rotation ($\kappa = 0.5$), while the bottom panel shows the case with anti-solar differential rotation ($\kappa = -0.5$). The contributions from each individual spot are shown in color, while the cumulative flux deficit is shown in black. The lightcurve exhibits quasi-periodic variability due to the rotation of the star and the emergence and decay of spots.

## 3. GAUSSIAN PROCESS

### 3.1. *Motivation for Gaussian Processes*

In practice, extracting the physical parameters of a star from its lightcurve is challenging due to the complex and stochastic nature of starspot-induced variability. The observed lightcurve is a superposition of signals from multiple spots, each emerging at different times and locations on the stellar surface. Even with very high quality data, performing model inversion to infer the underlying spot properties is inherently challenging for at least several reasons:

- **Degeneracies:** Multiple combinations of spot properties can produce similar lightcurves, leading to degeneracies in the inferred parameters (Figure 6).
- **Non-uniqueness:** The same lightcurve can be explained by different spot configurations, making it difficult to uniquely determine the spot properties (Figure 7).
- **Noise and systematics:** Real lightcurves are affected by instrumental noise, stellar granulation, and other sources of variability that can obscure the starspot signal and further complicate the inference of spot properties.





**Figure 6.** Illustration of degeneracies in starspot lightcurve modeling.

Given these challenges, a more practical approach is to not just model the properties of *individual* spots, but to instead model the *statistical properties* of the starspot population as a whole. This is where Gaussian processes (GPs) come in: they provide a flexible and powerful framework for modeling the covariance structure of the lightcurve without needing to explicitly model each individual spot. By deriving a physically motivated kernel function that captures the key features of starspot variability, we can use GPs to infer the underlying physical parameters of the star and its spots from the observed lightcurve, while marginalizing over the complex and degenerate spot configurations. This allows us to extract meaningful information about the star's rotation period, differential rotation, spot lifetimes, and emergence/decay timescales as a distribution over the entire spot population, rather than trying to pin down the properties of each individual spot.





**Figure 7.** Illustration of non-uniqueness in starspot lightcurve modeling. All lightcurves have the same parameters: $P_{\rm eq} = 5\,{\rm d}$, $\kappa = 0$, $I = 80°$, $N_{\rm spot} = 10$, $l_{\rm spot} = 8\,{\rm d}$, $\tau_{\rm spot} = 3\,{\rm d}$, yet the resulting lightcurves differ due to the random positions and emergence times of individual spots.

### 3.2. *Mathematical Formulation of Gaussian Processes*

Formally, a Gaussian process is defined by a multivariate Gaussian:

$$\mathcal{GP}(\mathbf{x}) \sim \mathcal{N}(m(\mathbf{x}), k(\mathbf{x}, \mathbf{x}')) \tag{15}$$

where the mean function, $m(\mathbf{x})$, represents the expected value of the function at each input point $\mathbf{x}$, while the covariance function, $k(\mathbf{x}, \mathbf{x}')$ models the correlation between function values at different input points $\mathbf{x}$ and $\mathbf{x}'$ and quantifies the smoothness and spatial properties of the underlying function.

This GP, defined by a mean function and a covariance function, represents a probability distribution over functions and can be used to model the lightcurve data. Given some lightcurve data $(\mathbf{x}, \mathbf{y})$, the log marginal likelihood (or evidence) of a GP is given by:

$$\log P(\mathbf{y}|\mathbf{x}, \theta) = -\frac{1}{2}\mathbf{y}^T K^{-1}\mathbf{y} - \frac{1}{2}\log|K| - \frac{n}{2}\log 2\pi \tag{16}$$

This distribution allows us to make predictions at new test points, where the predictive mean and variance at a given test point $x^*$ can be computed as:

$$\mu = k(x_*, \mathbf{x})k(\mathbf{x}, \mathbf{x})^{-1}\mathbf{y} \tag{17}$$

$$\sigma^2 = k(x_*, x_*) - k(x_*, \mathbf{x})k(\mathbf{x}, \mathbf{x})^{-1}k(\mathbf{x}, x_*) \tag{18}$$

For this particular application, the mean function $m(t)$ is determined by the expected flux deficit from the average spot coverage, which we derive analytically in Section 6.1. In practice, we subtract the mean from the observed data and set $m(t) = 0$. The covariance function $k(\mathbf{x}, \mathbf{x}')$ is derived in Section 4, and encodes the physical properties of the star and its spots. By maximizing the log marginal likelihood with respect to the hyperparameters $\theta$ of the kernel, we can infer the underlying physical parameters of the star and its spots from the observed lightcurve data. For simplification we assume that the distribution of starspot properties does not evolve with time. In other words, the kernel is *stationary* and depends only the time lag $\tau$ between two points. With these basic assumptions, we move on to how to compute the kernel function numerical and analytically. However, for a more comprehensive introduction to Gaussian processes and their properties, see Rasmussen & Williams (2006).

### 3.3. *The Covariance Matrix and Kernel Function*

A brute-force approach to modeling the covariance structure of the lightcurve is to directly compute the covariance matrix $K$ from a large sample of simulated lightcurves generated from the starspot model. Assuming we have 1-dimensional training data, where $\mathbf{x} = t_1, \dots t_n$ represent the times and $\mathbf{y} = F_1, \dots F_n$ represent the fluxes of the lightcurve, then we want to solve for covariance matrix:

$$K = \begin{bmatrix} \mathrm{Var}(F_1) & \cdots & \mathrm{Cov}(F_1, F_n) \\ \vdots & \ddots & \vdots \\ \mathrm{Cov}(F_n, F_1) & \cdots & \mathrm{Var}(F_n) \end{bmatrix} \tag{19}$$

Each element of the covariance matrix $K$ is the equivalent to the covariance between $F_i$ and $F_j$

$$K_{ij} = \mathrm{Cov}(F_i, F_j) \tag{20}$$

The first term is the expectation of the flux product at two different times $t$ and $t+\tau$. Flux at a given time is given by the Kipping (2012) lightcurve model (Eqn. 1). The random variables of this process are the reference time, latitude, and longitude of each spot ($t_{\mathrm{ref,k}}$, $\Lambda_{\mathrm{ref,k}}$, and $\Phi_{\mathrm{ref,k}}$ respectively).

The covariance matrix can then be estimated over a large sample of simulations with random variables (e.g., a uniform distribution with $t_{\mathrm{ref,k}} \in [0, T]$, $\Lambda_{\mathrm{ref,k}} \in [0, 2\pi]$, and $\Phi_{\mathrm{ref,k}} \in [-\pi/2, \pi/2]$). The covariance can then be estimated as the sample covariance of the fluxes at two different times $t$ and $t + \tau$:

$$\mathcal{K}_{\mathrm{num}}(\tau) = \mathrm{Cov}[F(t), F(t+\tau)] \tag{21}$$

$$= \frac{1}{N-1} \sum_{i=1}^{N} (F(t_i) - \mathbb{E}[F])(F(t_i + \tau) - \mathbb{E}[F]). \tag{22}$$

In practice, the numerical approach is computationally expensive, since it requires generating a large number of lightcurves and computing the covariance for each pair of time points. Moreover, it does not provide a closed-form expression for the kernel function, which limits its interpretability and makes it difficult to use in a Gaussian process regression framework.

## 4. DERIVING THE ANALYTICAL KERNEL FOR STARSPOT VARIABILITY

The goal of this section is to derive an analytical expression for the kernel function $\mathcal{K}(\tau)$, which describes the covariance of the flux at two different times $t$ and $t+\tau$ due to starspot variability. We first start with the analytic flux model defined in Equation 1 and then derive the kernel function by computing the autocorrelation of the flux signal. In this section we will show the derivation for the simplified case assuming the small spot approximation and no limb darkening, however the derivation can be extended to these more complex cases, which we derive in Paper II.

The result is a physically motivated kernel function that captures the key features of starspot-induced variability, such as the rotation period, spot lifetime, and emergence/decay timescales. The kernel function depends on the underlying physical properties of the star and its spots, which are encapsulated in the hyperparameters:

$$\theta = \{P_{\rm eq}, \kappa, I, \dot{N}_{\rm spot}, l_{\rm spot}, \tau_{\rm spot}, \alpha_{\rm max}\}.$$

This means that the kernel can be used in a Gaussian process regression framework to model and predict stellar variability, while also allowing for inference on the physical properties of the star and its spots through the hyperparameters. In the following sections, we give an overview of the Fourier transform notation and identities used in the derivation, and then we present the step-by-step derivation of the kernel function $\mathcal{K}(\tau)$ from the flux model.

### 4.1. *Fourier Transform Notation and Identities*

We use the Fourier transform convention $\hat{f}(\omega) = \int_{-\infty}^{\infty} f(t)\, e^{-i\omega t}\, dt$ with inverse $f(t) = (2\pi)^{-1} \int \hat{f}(\omega)\, e^{i\omega t}\, d\omega$ throughout. Table 3 summarizes the notation.

**Table 3.** Notation used in Section 4.

| Symbol | Description |
|---|---|
| $t$ | Time |
| $\tau$ | Time lag |
| $\omega$ | Angular frequency |
| $\omega_0$ | Fundamental angular frequency, $2\pi/T$ |
| $T$ | Period of a periodic function |
| $n$ | Harmonic index (integer) |
| $f(t)$, $g(t)$ | Functions in the time domain |
| $\hat{f}(\omega)$, $\hat{g}(\omega)$ | Fourier transforms of $f$ and $g$ |
| $c_n$ | Fourier series coefficients |
| $\delta(\cdot)$ | Dirac delta function |
| $(f * g)(t)$ | Convolution of $f$ and $g$ |
| $R_f(\tau)$ | Autocorrelation function of $f$ |
| $\varepsilon(\omega)$ | Energy spectral density, $\lvert\hat{f}(\omega)\rvert^2$ |
| $k(\tau)$ | Autocovariance kernel |
| $A(t)$ | Single-spot flux signal |
| $t_0$ | Fixed time shift |

The derivation relies on three standard results. First, a periodic function with period $T$ and Fourier coefficients $c_n = T^{-1}\int_0^T f(t)\,e^{-in\omega_0 t}\,dt$ (where $\omega_0 = 2\pi/T$) has a Fourier transform given by the weighted Dirac comb $\hat{f}(\omega) = 2\pi\sum_n c_n\,\delta(\omega - n\omega_0)$. Second, the convolution theorem: convolution in the time domain corresponds to multiplication in the frequency domain (FT4), and a product in the time domain becomes a convolution in the frequency domain (FT5). Third, the Wiener–Khinchin theorem: the autocorrelation $R_f(\tau) = \int f(t)\,f(t+\tau)\,dt$ is the inverse Fourier transform of the power spectral density $S_f(\omega) = |\hat{f}(\omega)|^2$:

$$R_f(\tau) = \frac{1}{2\pi}\int_{-\infty}^{\infty} |\hat{f}(\omega)|^2\,e^{i\omega\tau}\,d\omega. \tag{23}$$

Together, these allow us to obtain the autocovariance kernel $\mathcal{K}(\tau)$ from the Fourier transform of the single-spot signal $A(t)$. In the Appendix, Table 13 lists the Fourier transform properties and Table 14 lists the specific transform pairs used in this paper.

### 4.2. *Approach to Derivation*

By Eq. (7) and the flux model (Eq. 1), a single dark spot dims the star by $A(t)/\pi = \alpha_{\max}^2\,\Gamma(t)\,\mathcal{V}(t)$. The constant $\alpha_{\max}^2$ passes unchanged through every Fourier transform below, so in this section we write $A(t) \equiv \Gamma(t)\cdot\mathcal{V}(t)$ for the unit-amplitude single-spot signal; the kernel $\mathcal{K}(\tau)$ of the flux is $\alpha_{\max}^4$ times its autocorrelation. The autocorrelation function of function $A(t)$ is equivalent to taking the convolution between $A(t)$ and a lagged version of the function. By the convolution property (FT4), convolution in the time domain is multiplication in the frequency domain. Here $\mathcal{F}\{A(t)\} = \hat{A}(\omega)$ denotes the Fourier transform of a function $A(t)$, and $\mathcal{F}^{-1}\{\hat{A}(\omega)\} = A(t)$ denotes the inverse Fourier transform. So the Fourier transform of the kernel function is:

$$\begin{aligned}
\mathcal{F}\{\mathcal{K}(\tau)\} &= \alpha_{\max}^4\,A(t) * A(t-\tau) && (24)\\
&= \alpha_{\max}^4\,\hat{A}(\omega)\cdot\hat{A}(\omega) && (25)\\
&= \alpha_{\max}^4\,|\hat{A}(\omega)|^2 && (26)\\
&= \alpha_{\max}^4\,S_A(\omega) && (27)
\end{aligned}$$

The term $S_A(\omega)$ is the *power spectral density* of the single-spot signal. By the Wiener–Khinchin theorem (Eq. 23), the autocorrelation function is the inverse Fourier transform of the power spectral density:

$$\mathcal{K}(\tau) \overset{\mathcal{F}}{\longleftrightarrow} \alpha_{\max}^4\,S_A(\omega) \tag{28}$$

$$\mathcal{K}(\tau) = \frac{\alpha_{\max}^4}{2\pi}\int_{-\infty}^{\infty} S_A(\omega)e^{i\omega t}\,d\omega. \tag{29}$$

Hence to solve for $\mathcal{K}(\tau)$, we need to first solve for the Fourier transform of a single spot, $\hat{A}(\omega) = \mathcal{F}\{A(t)\}$. Referring back to Equation 7, ignoring the edge cases (for small spot sizes), the unit-amplitude signal $A(t)$ is the product of the normalized squared spot-size envelope $\Gamma(t) = \alpha^2(t)/\alpha_{\max}^2$ and the visibility function $\mathcal{V}(t) = \max\{\cos\beta(t), 0\}$. The visibility function is a truncated cosine, which can be written as the product of a square wave (P4) and cosine wave (P2): $\mathcal{V}(t) = \sqcap(t)\cdot C(t)$.

By the multiplication property (FT5), a product in the time domain becomes a convolution in the frequency domain:

$$\mathcal{F}\{A(t)\} = \mathcal{F}\{\Gamma(t) \cdot \mathcal{V}(t)\} \tag{30}$$
$$= \mathcal{F}\{\Gamma(t)\} * \mathcal{F}\{\mathcal{V}(t)\} \tag{31}$$
$$\hat{A}(\omega) = \hat{\Gamma}(\omega) * \hat{\mathcal{V}}(\omega) \tag{32}$$

As we show in the next section, the Fourier transform of the spot evolution $\hat{A}(\omega)$ can be solved analytically for both of the components $\hat{\Gamma}(\omega)$ and $\hat{\mathcal{V}}(\omega)$. In Section 4.3 we derive the Fourier transform of the truncated cosine $\hat{\mathcal{V}}(\omega)$ (representing the variability as the star rotates), and in Section 4.4 we derive the Fourier transform of the squared spot-size envelope $\hat{\Gamma}(\omega)$ (representing the growth and decay of the spot). Finally, in Section 4.7.2 we combine these results to obtain the final expression for the kernel $\mathcal{K}(\tau)$.





**Figure 8.** Overview of the kernel derivation strategy. *Top row (blue):* The analytic path derives $k(\tau)$ from the single-spot flux model $A(t) = \Gamma(t) \cdot \mathcal{V}(t)$ by (1) computing Fourier transforms of the envelope and visibility terms, (2) averaging the energy spectral density over random spot parameters ($t_{\rm ref}$, $\Lambda_{\rm ref}$, $\Phi_{\rm ref}$), and (3) applying the Wiener–Khinchin theorem. *Bottom row (red):* The numerical path simulates $N$ lightcurves from the same flux model with randomly drawn spot parameters and estimates the kernel as the sample-averaged autocorrelation. *Center (green):* The two paths converge, verifying the analytic kernel against the numerical estimate.

Figure 8 provides an overview of the derivation strategy. The analytic path (top row) derives $\mathcal{K}(\tau)$ through Fourier analysis, while a parallel numerical path (bottom row) estimates the kernel via Monte Carlo simulation. Finally, we compare the numerical and analytical kernels as a validation test of the derivation.

### 4.3. *Fourier Transform: The Rotation Term $\mathcal{V}(t)$*

As given in Equation 6, the drop in flux due to a spot rotating into and out of view is determined by the angle $\beta$ between the spot normal and the line of sight.[3] The spot is visible when $\cos\beta > 0$, and the visibility function is $\mathcal{V}(t) = \max\{\cos\beta(t), 0\}$. The angle $\beta$ depends on the spot latitude $\Phi$, the stellar inclination $I$, and the rotation period $P$ of the star.

Figure 9 shows how the signal varies as a function of spot latitude $\Phi$ and stellar inclination $I$. For a star viewed equator-on ($I = 90°$, left panel of Figure 9), the visibility function looks like a half-truncated cosine wave, where the spot rotates into view for half a rotation period and then rotates out of view for the other half. The different colors show the relative change in flux for spots at different latitudes $\Phi$, with spots at higher latitudes having smaller amplitudes due to their smaller projected areas.

Figure 9 also shows how the signal varies as a function of stellar inclination $I$ for a fixed spot latitude $\Phi = 60°$ (right panel). For a star viewed equator-on ($I = 90°$), the visibility function is a half-truncated cosine wave (red). For a star viewed at intermediate inclination ($60° < I < 90°$), the visibility function is a semi-truncated cosine wave (blue). For a star viewed closer to pole-on ($I = 30°$), the visibility function is a full cosine wave which never rotates out of view, but varies in amplitude due to the changing projected area of the spot (orange). Finally, for a star viewed pole-on ($I = 0°$), the visibility function is a constant (blue) since the spot is always visible with the same projected area.





**Figure 9.** Visibility function $\mathcal{V}(t) = \max\{\cos\beta(t), 0\}$ over three rotation periods. *Left:* varying spot latitude $\Phi$ at fixed edge-on inclination $I = 90°$. *Right:* varying stellar inclination $I$ at fixed latitude $\Phi = 60°$. The amplitude and duty cycle of $\mathcal{V}(t)$ set the Fourier coefficients $|c_n|^2$ that enter the kernel.

The visibility function $\mathcal{V}(t) = \max\{\cos\beta(t), 0\}$ is periodic with the rotation period $P = 2\pi/\omega_0$, so it has a discrete Fourier series (Section 4):

$$\mathcal{V}(t) = \sum_{n=-\infty}^{\infty} c_n \, e^{in\omega_0 t}, \tag{33}$$

[3] Symbolic and numerical check: `derive_ft_spot_rotation.ipynb`

and its Fourier transform is a summation over Dirac functions (P3):

$$\hat{\mathcal{V}}(\omega) = 2\pi \sum_{n=-\infty}^{\infty} c_n \, \delta(\omega - n\omega_0). \tag{34}$$

**Solution for the edge-on case**: For the edge-on case ($I = 90°$), $\cos\beta(t) = \cos\Phi\cos(\omega_0 t)$, and the visibility function reduces to $\mathcal{V}(t) = \cos\Phi \cdot \max\{\cos(\omega_0 t), 0\}$. The Fourier coefficients of the half-rectified cosine $\max\{\cos\theta, 0\}$ are (see Appendix B for the full derivation):

$$c_n = \cos\Phi \begin{cases} 1/\pi & n = 0 \\ 1/4 & n = \pm 1 \\ \dfrac{(-1)^{|n|/2+1}}{\pi(n^2-1)} & n = \pm 2, \pm 4, \pm 6, \ldots \\ 0 & n = \pm 3, \pm 5, \pm 7, \ldots \end{cases} \tag{35}$$

All coefficients are real and satisfy $c_{-n} = c_n$, as expected for a real, even function.

**Solution for general inclinations**: For general inclination, $\cos\beta(t) = b_0 + b_1\cos(\omega_0 t)$ where $b_0 = \cos I \sin\Phi$ and $b_1 = \sin I \cos\Phi$. The spot is visible when $|\omega_0 t| < \theta_{\rm vis} = \arccos(-b_0/b_1)$, and the general Fourier coefficients (derived in Appendix B) are:

$$c_n = \frac{1}{2\pi}\int_{-\theta_{\rm vis}}^{\theta_{\rm vis}} \left(b_0 + b_1\cos\theta\right) e^{-in\theta}\, d\theta = \frac{1}{\pi}\left[\frac{b_0\sin(n\,\theta_{\rm vis})}{n} + \frac{b_1}{2}\left(\frac{\sin\left((n-1)\theta_{\rm vis}\right)}{n-1} + \frac{\sin\left((n+1)\theta_{\rm vis}\right)}{n+1}\right)\right] \tag{36}$$

with $c_0 = (b_0\,\theta_{\rm vis} + b_1\sin\theta_{\rm vis})/\pi$ and $c_{\pm 1} = (b_0\sin\theta_{\rm vis} + b_1[\theta_{\rm vis} + \sin\theta_{\rm vis}\cos\theta_{\rm vis}]/2)/\pi$ obtained by taking the appropriate limits. These are not separate cases but removable singularities of Eq. (36), recovered using $\sin(n\theta_{\rm vis})/n \to \theta_{\rm vis}$ as $n \to 0$ and $\sin[(n-1)\theta_{\rm vis}]/(n-1) \to \theta_{\rm vis}$ as $n \to 1$.

In general, it is difficult to solve for a closed-form solution to the integral of the full kernel function in Equation 59 due to the complexity of the latitude-dependent terms. However, for specialized cases (e.g., a star with edge-on inclination, solid body rotation, and uniform latitude spots), the kernel reduces to a simple, closed-form solution. Truncating to second harmonic order, the Fourier transform of the visibility function is:

$$\hat{\mathcal{V}}(\omega) \approx 2\pi\left[c_0\,\delta(\omega) + c_1\left[\delta(\omega-\omega_0) + \delta(\omega+\omega_0)\right] + c_2\left[\delta(\omega-2\omega_0) + \delta(\omega+2\omega_0)\right]\right]. \tag{37}$$

As we will show in Section 4.5, the kernel $\mathcal{K}(\tau)$ is dominated by the first two harmonics, so this approximation is sufficient for our purposes (though in our `spotgp` implementation, this Fourier series can be expanded to an arbitrary number of terms as specified by the user).

### 4.4. *Fourier Transform: Spot Evolution Envelope Term* $\Gamma(t)$

In this section we show how to derive the Fourier transform of the spot evolution envelope, which captures the growth and decay of the spot over time.[4] Here we present several common spot evolution

[4] Symbolic and numerical checks: `derive_ft_squared_trapezoid.ipynb` for the squared trapezoid $\Gamma(t)$, and `derive_ft_trapezoid.ipynb` for the trapezoid $\alpha(t)$ itself.

profiles, but the same approach can be applied to any arbitrary profile $\alpha(t)$ by performing the appropriate integral. These profiles include: (1) squared trapezoidal (Kipping 2012), (2) exponential (Aigrain et al. 2015), and (3) skewed normal (Forgács-Dajka et al. 2021). The trapezoidal and exponential profiles are simple functions which produce an analytic kernel function. On the other hand, the skewed normal profile is a more flexible profile that can capture asymmetric growth and decay timescales and reproduce the observed spot lifetimes of the Sun (Forgács-Dajka et al. 2021).

In the Results and Discussion (Sections 9–11) we show a detailed analysis of how the properties of the trapezoidal profile translate to the structure of the kernel function, as its analytic solution is easily interpretable. While perhaps not the most realistic model, the trapezoidal parameterization provides an intuitive insight into how the kernel behaves in different limits. We include the implementations for all three kernels in our `spotgp` implementation (as well as the flexibility to define custom envelope functions) so that users can experiment with different assumptions for spot lifetimes and decay timescales.

**(1) Squared trapezoidal profile for $\Gamma(t)$:** In the first example, we consider the trapezoidal profile (Equation 9) from Kipping (2012). Appendix C, shows how we solve for the Fourier transform of the normalized squared envelope $\Gamma(t) = \alpha^2(t)/\alpha^2_{\rm max}$: Since $\alpha(t)$ is a trapezoid with peak $\alpha_{\rm max}$, the normalized squared envelope $\Gamma(t) = \alpha^2(t)/\alpha^2_{\rm max}$ has unit peak height, the same support $\ell_{\rm spot} + 2\tau_{\rm spot}$, the same plateau duration $\ell_{\rm spot}$, but parabolic (rather than linear) ramps. Since $\Gamma(t)$ is real and even, its Fourier transform is purely real and can be computed via the cosine transform $\hat{\Gamma}(\omega) = 2\int_0^\infty \Gamma(t)\cos(\omega t)\,dt$, splitting into a plateau integral and a parabolic-ramp integral with two integrations by parts (see Appendix C):

$$\hat{\Gamma}(\omega) = \frac{4}{\tau^2_{\rm spot}\,\omega^3}\left[\tau_{\rm spot}\,\omega\cos\left(\frac{\omega\,\ell_{\rm spot}}{2}\right) + \sin\left(\frac{\omega\,\ell_{\rm spot}}{2}\right) - \sin\left(\frac{\omega\,\ell_{\rm spot}}{2} + \omega\,\tau_{\rm spot}\right)\right]. \tag{38}$$

A visualization of this profile is shown in the left panel of Figure 10.

**(2) Exponential profile for $\Gamma(t)$:** For the piecewise exponential envelope (Eq. 10), setting $t_1 = 0$ and $t_2 = \tau_{\rm plat}$, the normalized squared envelope $\Gamma(t) = \alpha^2_{\rm exp}(t)/\alpha^2_{\rm max}$ is

$$\Gamma(t) = \begin{cases} e^{2t/\tau_{\rm em}} & t < 0 \\ 1 & 0 \le t \le \tau_{\rm plat} \\ e^{-2(t-\tau_{\rm plat})/\tau_{\rm dec}} & t > \tau_{\rm plat} \end{cases} \tag{39}$$

with doubled exponential rates compared to $\alpha(t)$. Because $\Gamma(t)$ is not symmetric when $\tau_{\rm em} \neq \tau_{\rm dec}$, the Fourier transform is complex. Splitting $\hat{\Gamma}(\omega) = \hat{\Gamma}_R(\omega) - i\,\hat{\Gamma}_I(\omega)$ and evaluating each piece integral analytically:

$$\hat{\Gamma}_R(\omega) = \frac{2\tau_{\rm em}}{4+\omega^2\tau^2_{\rm em}} + \frac{\sin(\omega\tau_{\rm plat})}{\omega} + \frac{2\tau_{\rm dec}\cos(\omega\tau_{\rm plat})}{4+\omega^2\tau^2_{\rm dec}} - \frac{\omega\tau^2_{\rm dec}\sin(\omega\tau_{\rm plat})}{4+\omega^2\tau^2_{\rm dec}} \tag{40}$$

$$\hat{\Gamma}_I(\omega) = -\frac{\omega\tau^2_{\rm em}}{4+\omega^2\tau^2_{\rm em}} + \frac{1-\cos(\omega\tau_{\rm plat})}{\omega} + \frac{\omega\tau^2_{\rm dec}\cos(\omega\tau_{\rm plat})}{4+\omega^2\tau^2_{\rm dec}} + \frac{2\tau_{\rm dec}\sin(\omega\tau_{\rm plat})}{4+\omega^2\tau^2_{\rm dec}} \tag{41}$$

The full derivation, including the direct convolution for $R_\Gamma(\tau)$, is analogous to the trapezoidal case (Appendix C), and a figure of the function is shown in the middle panel of Figure 10.

**(3) Skewed Normal profile for $\Gamma(t)$:** Forgács-Dajka et al. (2021) modeled sunspot group area evolution using a skew-normal function (their Eq. 1), which naturally captures the observed asymmetry between rapid emergence and slower decay phases of sunspot lifetimes. We adopt this as a third envelope option. The normalized squared envelope is

$$\Gamma(t) \;\propto\; \exp\!\left(-\frac{t^2}{2\sigma_{\rm sn}^2}\right)\left[1+\operatorname{erf}\!\left(\frac{n_{\rm sn}\,t}{\sigma_{\rm sn}\sqrt{2}}\right)\right], \tag{42}$$

rescaled so that the peak value equals unity. The two free parameters are $\sigma_{\rm sn}$ (the scale parameter in days, controlling the overall spot lifetime) and $n_{\rm sn}$ (the dimensionless skewness parameter). When $n_{\rm sn} = 0$ the envelope reduces to a symmetric Gaussian; $n_{\rm sn} > 0$ gives rapid rise and slow decay (typical of solar spots), while $n_{\rm sn} < 0$ gives the opposite asymmetry. Unlike the trapezoidal and exponential profiles, the skew-normal $\Gamma(t)$ does not yield a closed-form autocorrelation $R_\Gamma(\tau)$ or Fourier transform $\widehat{\Gamma}(\omega)$, so we therefore compute both numerically via FFT-based methods in our `spotgp` implementation. This function is shown in the right panel of Figure 10.





**Figure 10.** Comparison of the three spot envelope models: trapezoidal (blue), exponential (green), and skew-normal (red). *Left:* Normalized squared envelopes $\Gamma(t)$. *Middle:* Energy spectral densities $S_\Gamma(\omega)$ (normalized, log scale), showing the different high-frequency decay rates set by envelope smoothness. *Right:* Normalized envelope autocorrelation $R_\Gamma(\tau)/R_\Gamma(0)$. All three implementations are included in our `spotgp` package, along with the functionality to define custom envelope profiles.

### 4.5. *Combined Kernel for a Single Spot*

Substituting the delta train (34) into Eq. (32) and using the sifting property of the Dirac delta (P5, FT3), $f(\omega) * \delta(\omega - a) = f(\omega - a)$:

$$\hat{A}(\omega) = \sum_n c_n\,\hat{\Gamma}(\omega - n\omega_0) \tag{43}$$

Explicitly to second harmonic order, using linearity (FT1):

$$\boxed{\hat{A}(\omega) = c_0\,\hat{\Gamma}(\omega) + c_1\big[\hat{\Gamma}(\omega-\omega_0)+\hat{\Gamma}(\omega+\omega_0)\big] + c_2\big[\hat{\Gamma}(\omega-2\omega_0)+\hat{\Gamma}(\omega+2\omega_0)\big]} \tag{44}$$

where $c_0$, $c_1$, $c_2$ are the Fourier coefficients of the visibility function given by Eq. (36) (all real, with $c_{-n} = c_n$). Each term has a clear physical interpretation in the time domain:

- The $n = 0$ (DC "direct current") term, $c_0\,\Gamma(t)$, is the non-oscillatory component: the mean flux deficit that tracks the spot-size envelope. It is largest when the spot is visible for most of the rotation (e.g., a high-latitude spot on a low-inclination star), contributing a smooth, aperiodic dimming.
- The $n = 1$ (fundamental) term, $2c_1\,\Gamma(t)\cos(\omega_0 t)$, is the dominant periodic component at the stellar rotation frequency. It produces the familiar once-per-rotation brightness dip as the spot rotates in and out of view.
- The $n = 2$ (second harmonic) term, $2c_2\,\Gamma(t)\cos(2\omega_0 t)$, captures the non-sinusoidal shape of the visibility function (the flattened profile near disk center and sharp limb transitions), which a pure cosine cannot reproduce.

Figure 11 shows the Fourier decomposition of the single-spot signal $A(t) = \Gamma(t)\,\mathcal{V}(t)$ for three stellar inclinations at a fixed latitude $\Phi = 30°$, with the same spot decay envelope $\Gamma(t)$. Here we see that terms up to second harmonic order are sufficient for capturing the shape of the signal for varying $I$.





**Figure 11.** Fourier decomposition of the single-spot signal $A(t) = \Gamma(t) \cdot \mathcal{V}(t)$ for a spot at latitude $\Phi = 30°$. *Left:* Time-domain signal for three stellar inclinations, with the normalized squared envelope $\Gamma(t) = \alpha^2(t)/\alpha^2_{\max}$ shown as a dashed gray line. *Right:* Normalized squared Fourier coefficients $|c_n|^2/\sum|c_n|^2$ of the visibility function for each inclination.

### 4.6. *Averaging Over Random Spot Parameters*

The expression for $\hat{A}(\omega)$ in Eq. (43) describes a *single* spot with a particular reference time $t_{\rm ref}$, reference longitude $\Lambda_{\rm ref}$, and latitude $\Phi_{\rm ref}$. To obtain the kernel for a *population* of spots, we must average the power spectral density $|\hat{A}(\omega)|^2$ over these random variables.[5] We treat $t_{\rm ref}$ and $\Lambda_{\rm ref}$ as uniformly distributed, and $\Phi_{\rm ref}$ as drawn from some latitude distribution $p(\Phi)$.

#### 4.6.1. *Effect of $t_{\rm ref}$: time shift of the envelope*

Shifting the envelope by $t_{\rm ref}$ gives $\Gamma(t - t_{\rm ref})$, whose Fourier transform picks up a phase by the time-shifting property (FT2, P5):

$$\hat{\Gamma}(\omega;\, t_{\rm ref}) = \hat{\Gamma}(\omega)\, e^{-i\omega\, t_{\rm ref}}. \tag{45}$$

[5] Symbolic and numerical check: `derive_averaging_random_params.ipynb`

This phase propagates to $\hat{A}(\omega)$, but since $|e^{-i\omega\, t_{\rm ref}}|^2 = 1$, it **drops out** of the power spectral density $|\hat{A}(\omega)|^2$. The kernel is therefore independent of $t_{\rm ref}$: any spot reference time yields the same autocorrelation.

#### 4.6.2. *Effect of $\Lambda_{\rm ref}$: longitude phase*

Shifting the initial longitude $\Lambda_{\rm ref}$ is equivalent to a time shift of $\Lambda_{\rm ref}/\omega_0$ in the periodic visibility function. By the time-shifting property (FT2), the Fourier series coefficients acquire a harmonic-dependent phase:

$$c_n(\Lambda_{\rm ref}) = c_n\, e^{in\Lambda_{\rm ref}}, \tag{46}$$

where $c_n \equiv c_n(\Lambda_{\rm ref} = 0)$. Substituting into Eq. (43), the full single-spot spectrum is:

$$\hat{A}(\omega) = e^{-i\omega t_{\rm ref}} \sum_n c_n\, e^{in\Lambda_{\rm ref}}\, \hat{\Gamma}(\omega - n\omega_0). \tag{47}$$

The power spectral density is:

$$|\hat{A}(\omega)|^2 = \sum_n \sum_m c_n\, c_m^*\, e^{i(n-m)\Lambda_{\rm ref}}\, \hat{\Gamma}(\omega - n\omega_0)\, \hat{\Gamma}^*(\omega - m\omega_0). \tag{48}$$

Averaging over $\Lambda_{\rm ref}$ uniformly distributed on $[0, 2\pi]$:

$$\left\langle e^{i(n-m)\Lambda_{\rm ref}} \right\rangle_{\Lambda_{\rm ref}} = \frac{1}{2\pi} \int_0^{2\pi} e^{i(n-m)\Lambda_{\rm ref}}\, d\Lambda_{\rm ref} = \delta_{nm}. \tag{49}$$

This **cancels out all cross terms** between different harmonics. Only the diagonal ($n = m$) terms survive:

$$\boxed{\left\langle |\hat{A}(\omega)|^2 \right\rangle_{t_{\rm ref},\, \Lambda_{\rm ref}} = \sum_{n=-\infty}^{\infty} |c_n|^2\, S_\Gamma(\omega - n\omega_0),} \tag{50}$$

where $S_\Gamma(\omega) \equiv |\hat{\Gamma}(\omega)|^2$ is the power spectral density of the squared envelope, and each harmonic of the visibility function contributes independently to the power spectrum, with weight $|c_n|^2$.

#### 4.6.3. *From averaged PSD to the kernel*

Applying the Wiener–Khinchin theorem (Eq. 23) to the averaged power spectral density:

$$\mathcal{K}(\tau) = \frac{\alpha_{\rm max}^4}{2\pi} \int_{-\infty}^{\infty} \left\langle |\hat{A}(\omega)|^2 \right\rangle\, e^{i\omega\tau}\, d\omega = \alpha_{\rm max}^4 \sum_n |c_n|^2 \cdot \frac{1}{2\pi} \int_{-\infty}^{\infty} S_\Gamma(\omega - n\omega_0)\, e^{i\omega\tau}\, d\omega. \tag{51}$$

Substituting $u = \omega - n\omega_0$:

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} S_\Gamma(u)\, e^{i(u+n\omega_0)\tau}\, du = e^{in\omega_0\tau} \cdot \underbrace{\frac{1}{2\pi} \int_{-\infty}^{\infty} S_\Gamma(u)\, e^{iu\tau}\, du}_{R_\Gamma(\tau)}, \tag{52}$$

where $R_\Gamma(\tau)$ is the autocorrelation of the normalized squared envelope $\Gamma(t) = \alpha^2(t)/\alpha_{\rm max}^2$. Therefore:

$$\mathcal{K}(\tau) = \alpha_{\rm max}^4\, R_\Gamma(\tau) \sum_{n=-\infty}^{\infty} |c_n|^2\, e^{in\omega_0\tau}. \tag{53}$$

Since the $c_n$ are real with $c_{-n} = c_n$, the complex exponentials combine into cosines:

$$\boxed{\mathcal{K}(\tau) = \alpha_{\max}^4 \, R_\Gamma(\tau) \left[ |c_0|^2 + 2 \sum_{n=1}^{\infty} |c_n|^2 \, \cos(n\omega_0 \tau) \right]} \tag{54}$$

The kernel factorizes into two interpretable components:

- $R_\Gamma(\tau)$: the **normalized envelope autocorrelation** of $\Gamma(t) = \alpha^2(t)/\alpha_{\max}^2$, which sets the overall decay timescale of correlations (governed by spot lifetime $\ell_{\rm spot}$ and rise/decay time $\tau_{\rm spot}$).
- The **cosine sum**: a quasi-periodic modulation at the rotation frequency and its harmonics, with amplitudes set by the squared Fourier coefficients $|c_n|^2$ of the visibility function.

Truncating to second harmonic order gets:

$$\mathcal{K}(\tau) \approx \alpha_{\max}^4 \, R_\Gamma(\tau) \left[ |c_0|^2 + 2|c_1|^2 \cos(\omega_0 \tau) + 2|c_2|^2 \cos(2\omega_0 \tau) \right], \tag{55}$$

which represents the kernel function for a *single* spot.

### 4.7. *Multi-Spot Kernel*

The observed flux deficit is the sum over $N_{\rm spot}$ independent spots (Eq. 1). With the small-spot projected area $A_k(t) = \pi \, \alpha_k^2(t) \, \mathcal{V}_k(t)$ (Eq. 7), the factor of $\pi$ cancels the $1/\pi$ of the flux model:

$$\delta F(t) = \frac{1}{\pi} \sum_{k=1}^{N_{\rm spot}} A_k(t) \, (1 - f_{{\rm spot},k}) = \sum_{k=1}^{N_{\rm spot}} (1 - f_{{\rm spot},k}) \, \alpha_k^2(t) \, \mathcal{V}_k(t). \tag{56}$$

Each term is the single-spot flux deficit $\alpha^2(t) \, \mathcal{V}(t) = \alpha_{\max}^2 A(t)$, whose autocorrelation, averaged over the spot's random parameters, is the single-spot kernel of Eq. (54). That autocorrelation is integrated over time, so it has units of time. Spots emerge at random reference times $t_{{\rm ref},k}$, which we model as a Poisson process with rate $\dot{N}_{\rm spot}$ [spots/day]. For such a sum of independent pulses at random times (Campbell's theorem), the covariance is the emergence rate times the time-integrated autocorrelation of a single pulse. For spots with identical contrast $f_{\rm spot}$:

$$\mathcal{K}(\tau) \equiv {\rm Cov}[\delta F(t), \, \delta F(t+\tau)] = \dot{N}_{\rm spot} \, (1 - f_{\rm spot})^2 \, \langle \mathcal{K}(\tau) \rangle_\Phi \, , \tag{57}$$

where $\langle \mathcal{K}(\tau) \rangle_\Phi$ is the single-spot kernel (Eq. 54) averaged over the latitude distribution $p(\Phi)$. Equivalently, $N_{\rm spot}$ spots with reference times spread uniformly over a long baseline $T_{\rm obs}$ each contribute $\langle \mathcal{K}(\tau) \rangle_\Phi / T_{\rm obs}$, so that $\dot{N}_{\rm spot} = N_{\rm spot}/T_{\rm obs}$.

#### 4.7.1. *Averaging over latitude*

The latitude $\Phi_{\rm ref}$ affects both the Fourier coefficients $c_n$ (through $b_0 = \cos I \sin \Phi$ and $b_1 = \sin I \cos \Phi$) and, in the presence of differential rotation, the rotation frequency $\omega_0(\Phi) = 2\pi(1 - \kappa \sin^2 \Phi)/P_{\rm eq}$. Averaging Eq. (54) over $p(\Phi)$:

$$\langle \mathcal{K}(\tau) \rangle_\Phi = \alpha_{\max}^4 \int_{-\pi/2}^{\pi/2} R_\Gamma(\tau) \left[ |c_0(\Phi)|^2 + 2 \sum_{n=1}^{N} |c_n(\Phi)|^2 \, \cos\big(n\omega_0(\Phi) \, \tau\big) \right] p(\Phi) \, d\Phi. \tag{58}$$

In general, this integral must be evaluated numerically.

#### 4.7.2. *Full kernel*

Combining Eqs. (57) and (58), the complete GP kernel for the flux variability due to a population of independent spots is:

$$\boxed{\mathcal{K}(\tau) = \dot{N}_{\rm spot}\,(1-f_{\rm spot})^2\,\alpha_{\rm max}^4 \int_{-\pi/2}^{\pi/2} R_\Gamma(\tau)\left[\,|c_0(I,\Phi)|^2 + 2\sum_{n=1}^{N}|c_n(I,\Phi)|^2\,\cos\!\big(n\,\omega_0(\Phi)\,\tau\big)\right] p(\Phi)\,d\Phi} \tag{59}$$

where $R_\Gamma(\tau)$ is the normalized autocorrelation of the squared spot-size envelope $\Gamma(t) = \alpha^2(t)/\alpha_{\rm max}^2$, $c_n(I,\Phi)$ are the Fourier coefficients of the visibility function (Eq. 36), $\omega_0(\Phi) = 2\pi(1-\kappa\sin^2\Phi)/P_{\rm eq}$ is the latitude-dependent rotation frequency, and $p(\Phi)$ is the spot latitude distribution.

The stellar inclination enters the kernel *only* through the visibility coefficients $c_n$; the rotation frequency $\omega_0(\Phi)$ is independent of $I$. Writing $b_0 = \cos I \sin\Phi$ and $b_1 = \sin I\cos\Phi$ out explicitly, the half-width of the visibility window depends on $I$ and $\Phi$ through the single combination $\cot I\tan\Phi$,

$$\theta_{\rm vis}(I,\Phi) = \arccos\!\big(-\cot I\,\tan\Phi\big), \tag{60}$$

and the Fourier coefficients of Eq. (36) become

$$c_n(I,\Phi) = \frac{1}{\pi}\left[\frac{\cos I\sin\Phi}{n}\,\sin\!\big(n\,\theta_{\rm vis}\big) + \frac{\sin I\cos\Phi}{2}\left(\frac{\sin\!\big((n-1)\theta_{\rm vis}\big)}{n-1} + \frac{\sin\!\big((n+1)\theta_{\rm vis}\big)}{n+1}\right)\right]. \tag{61}$$

Equation (60) partitions the stellar surface at the latitudes $\Phi = \pm I$, as described following Eq. (6): spots at $\Phi \geq I$ are circumpolar ($\theta_{\rm vis} = \pi$), spots at $\Phi \leq -I$ never rotate into view ($\theta_{\rm vis} = 0$, $c_n \equiv 0$), and only the band $|\Phi| < I$ contributes rotationally modulated power, with duty cycle $\Delta = \theta_{\rm vis}/\pi$. We stress that the integral in Eq. (59) runs over the full physical range $[-\pi/2, \pi/2]$ and not the visible range $(-I, \pi/2]$: the two are equivalent because $c_n$ vanishes identically on the hidden cap, and retaining the full range is what keeps $p(\Phi)$ normalized over the whole star. Restricting the bounds and renormalizing $p(\Phi)$ over them would rescale the kernel amplitude by $\pi/(\pi/2 + I)$.

Two limits are instructive. For edge-on viewing ($I = \pi/2$) the $\cot I$ term vanishes, so $\theta_{\rm vis} = \pi/2$ at every latitude and $c_n(\pi/2,\Phi) = g_n\cos\Phi$, recovering the closed-form kernel of Eq. (66) below. For pole-on viewing ($I \to 0$) we have $b_1 \to 0$, the visibility function reduces to the constant $\sin\Phi$ over the northern hemisphere, all $c_{n\geq 1} \to 0$, and the kernel collapses to pure envelope decay with no rotational signal, $\mathcal{K}(\tau) \to \sigma_k^2 R_\Gamma(\tau)/4$ for $p(\Phi) = 1/\pi$. Decreasing the inclination therefore does not reduce the total variance $\mathcal{K}(0)$; it transfers power out of the rotation harmonics and into the aperiodic $n = 0$ term.

The full hyperparameters of this kernel are:

$$\theta = \{P_{\rm eq},\ \kappa,\ I,\ \dot{N}_{\rm spot},\ \ell_{\rm spot},\ \tau_{\rm spot},\ \alpha_{\rm max},\ f_{\rm spot}\}. \tag{62}$$

The amplitude in Eq. (59) is set by the spot emergence rate $\dot{N}_{\rm spot}$ [spots/day], not by the total number of spots. For a time series of duration $T_{\rm obs}$ the expected number of spots is $N_{\rm spot} = \dot{N}_{\rm spot}\,T_{\rm obs}$, but a longer baseline adds spots without changing the variance of the flux. The amplitude prefactor $\dot{N}_{\rm spot}\,(1-f_{\rm spot})^2\,\alpha_{\rm max}^4$ is therefore independent of the observing baseline, an intrinsic stellar property. It sets only the spot part of the zero-lag variance $\mathcal{K}(0)$ (granulation, stellar oscillations and instrumental noise add variance of their own, and when they are not modeled the fitted spot amplitude absorbs

part of it and is biased high; Section 11.3). In our `spotgp` implementation the amplitude is specified through `nspot_rate` (emergence rate in spots/day), which is the recommended parameterization for inference.

The amplitude prefactor is degenerate in practice (only the product can be constrained from the lightcurve variance) so it can be absorbed into a single amplitude hyperparameter $\sigma_k^2$. Defining the spot contrast-area product as $a_{\rm spot} = (1 - f_{\rm spot})\,\alpha_{\rm max}^2$, then:

$$\sigma_k^2 \equiv \dot{N}_{\rm spot}\, a_{\rm spot}^2, \tag{63}$$

and the reduced hyperparameter set becomes:

$$\theta = \{P_{\rm eq},\ \kappa,\ I,\ \ell_{\rm spot},\ \tau_{\rm spot},\ \sigma_k\}. \tag{64}$$

#### 4.7.3. *Analytic solution for $\kappa = 0$ and $I = 90$*

In some cases (e.g. a solid-body rotator viewed at 90 deg inclination), there is a closed-form analytic solution for the kernel.[6] Truncating to second harmonic order ($n \le 2$) and specializing to solid-body rotation ($\kappa = 0$), the cosines factor out of the latitude integral and the kernel becomes:

$$\mathcal{K}(\tau) = \sigma_k^2\, R_\Gamma(\tau)\left[\langle|c_0|^2\rangle_\Phi + 2\langle|c_1|^2\rangle_\Phi \cos(\omega_0\tau) + 2\langle|c_2|^2\rangle_\Phi \cos(2\omega_0\tau)\right], \tag{65}$$

where $\omega_0 = 2\pi/P_{\rm eq}$ and the latitude-averaged squared coefficients are $\langle|c_n|^2\rangle_\Phi = \int_{-\pi/2}^{\pi/2} |c_n(\Phi)|^2\, p(\Phi)\, d\Phi$, with $c_n(\Phi)$ given by Eq. (36).

For the special case of edge-on viewing ($I = \pi/2$) and a uniform latitude distribution $p(\Phi) = 1/\pi$ on $[-\pi/2, \pi/2]$, all Fourier coefficients are proportional to $\cos\Phi$: $c_n(\Phi) = g_n \cos\Phi$, where $g_0 = 1/\pi$, $g_1 = 1/4$, and $g_2 = 1/(3\pi)$. The latitude integrals evaluate to $\langle|c_n|^2\rangle_\Phi = g_n^2/2$, giving the fully closed-form kernel:

$$\mathcal{K}(\tau) = \sigma_k^2\, R_\Gamma(\tau)\left[\frac{1}{2\pi^2} + \frac{1}{16}\cos(\omega_0\tau) + \frac{1}{9\pi^2}\cos(2\omega_0\tau)\right] \quad (I = \pi/2,\ \kappa = 0). \tag{66}$$

For general inclination $I$, the latitude integrals $\langle|c_n|^2\rangle_\Phi$ must be evaluated numerically, since $c_0$ and $c_1$ depend on $\theta_{\rm vis}(\Phi) = \arccos(-\cot I \tan\Phi)$, which prevents elementary integration. However, the numerical quadrature over $\Phi$ is inexpensive (a single 1D integral per harmonic). This simplified form (66) is applicable when a system is viewed edge-on ($I = \pi/2$) with no differential rotation ($\kappa = 0$) and under the assumption that spots are uniformly distributed across all latitudes, however it is flexible enough to work for different envelope terms $R_\Gamma(\tau)$.

### 4.8. *Power Spectral Density of the Full Kernel*

By the Wiener–Khinchin theorem (Eq. 23), the kernel $\mathcal{K}(\tau)$ and its power spectral density (PSD) $S_\mathcal{K}(\omega)$ form a Fourier transform pair:

$$S_\mathcal{K}(\omega) = \int_{-\infty}^{\infty} \mathcal{K}(\tau)\, e^{-i\omega\tau}\, d\tau. \tag{67}$$

[6] Symbolic and numerical checks: `derive_closed_form_kernel_n2.ipynb` for $n \le 2$ and general $I$, and `derive_closed_form_kernel.ipynb` for all harmonics.

Since $\mathcal{K}(\tau)$ is real and even, $S_{\mathcal{K}}(\omega)$ is real, non-negative, and even in $\omega$. To evaluate the transform, we use the frequency-shift property: for a real, even function $R_\Gamma(\tau)$ with transform $S_\Gamma(\omega) = \mathcal{F}\{R_\Gamma\}$,

$$\mathcal{F}\{R_\Gamma(\tau)\cos(n\omega_0\tau)\} = \tfrac{1}{2}\big[S_\Gamma(\omega - n\omega_0) + S_\Gamma(\omega + n\omega_0)\big], \tag{68}$$

which follows from writing $\cos(n\omega_0\tau) = \frac{1}{2}(e^{in\omega_0\tau} + e^{-in\omega_0\tau})$ and applying the frequency-shift property (FT3).

Applying this to the solid-body rotation kernel (Eq. 65) yields:

$$S_{\mathcal{K}}(\omega) = \sigma_k^2 \left[\langle|c_0|^2\rangle_\Phi\, S_\Gamma(\omega) + \sum_{n=1}^{N}\langle|c_n|^2\rangle_\Phi \Big[S_\Gamma(\omega - n\omega_0) + S_\Gamma(\omega + n\omega_0)\Big]\right], \quad (\kappa = 0), \tag{69}$$

where $S_\Gamma(\omega) = |\hat{\Gamma}(\omega)|^2$ is the energy spectral density of the squared envelope and $\omega_0 = 2\pi/P_{\rm eq}$. The PSD is a sum of shifted copies of the envelope spectrum $S_\Gamma(\omega)$, each centered on a harmonic $n\omega_0$ and weighted by the corresponding latitude-averaged squared Fourier coefficient $\langle|c_n|^2\rangle_\Phi$. The $n = 0$ term produces a low-frequency component centered at $\omega = 0$, while each $n \geq 1$ term contributes a pair of symmetric peaks at $\omega = \pm n\omega_0$, whose width is set by the envelope spectral width (inversely proportional to the spot lifetime).

For the general case with differential rotation ($\kappa \neq 0$), the latitude-dependent rotation frequency $\omega_0(\Phi)$ prevents factoring the cosine out of the latitude integral. Taking the Fourier transform of the full kernel (Eq. 59):

$$\boxed{S_{\mathcal{K}}(\omega) = \sigma_k^2 \int_{-\pi/2}^{\pi/2} \left[|c_0(\Phi)|^2\, S_\Gamma(\omega) + \sum_{n=1}^{N} |c_n(\Phi)|^2 \Big[S_\Gamma\big(\omega - n\omega_0(\Phi)\big) + S_\Gamma\big(\omega + n\omega_0(\Phi)\big)\Big]\right] p(\Phi)\, d\Phi.} \tag{70}$$

Differential rotation causes the harmonic peaks to broaden: spots at different latitudes contribute power at slightly different frequencies $n\omega_0(\Phi)$, smearing each peak over the range $n\omega_0(0)$ to $n\omega_0(\pi/2) = n\omega_0(0)(1 - \kappa)$.

## 5. DERIVING THE MEAN FUNCTION

A Gaussian process is fully specified by its mean function $m(t) = \mathbb{E}[F(t)]$ and covariance function $\mathcal{K}(\tau)$. Having derived the kernel in Section 4, we now compute the mean function analytically from the flux model.

As in Section 4.7, we model spot emergence as a stationary process with rate $\dot{N}_{\rm spot}$ [spots day$^{-1}$], so that each spot contributes a finite-duration flux pulse. For identical spots with contrast $f_{\rm spot}$, the expected flux deficit at any instant is the emergence rate times the time-integrated flux deficit of a single spot:

$$\mathbb{E}[\delta F] = \frac{\dot{N}_{\rm spot}\,(1 - f_{\rm spot})}{\pi} \int_{-\infty}^{\infty} \mathbb{E}_{\Lambda,\,\Phi}[A(t)]\, dt. \tag{71}$$

The integral converges because each spot has finite support through $\Gamma(t)$, and $t_{\rm ref}$ does not appear: a shift in spot emergence time is equivalent to a phase rotation of the longitude, which is already uniformly distributed.

Using the small-spot factorization (Eq. 7), $A(t) = \pi\,\alpha_{\max}^2\,\Gamma(t)\,\mathcal{V}(t;\,\Lambda_{\rm ref},\,\Phi)$, we average over $\Lambda_{\rm ref}$ inside the time integral. Since $\Lambda_{\rm ref}$ is uniform on $[0, 2\pi]$, the longitude-averaged visibility is constant in time:

$$\bar{\mathcal{V}}(\Phi, I) \equiv \langle \mathcal{V}(t) \rangle_\Lambda = \frac{1}{2\pi} \int_0^{2\pi} \max\!\big\{ b_0 + b_1 \cos(\omega_0 t + \Lambda_{\rm ref}),\; 0 \big\}\, d\Lambda_{\rm ref} = \frac{b_0\,\theta_{\rm vis} + b_1 \sin\theta_{\rm vis}}{\pi}, \tag{72}$$

where $b_0 = \cos I \sin\Phi$, $b_1 = \sin I \cos\Phi$, and $\theta_{\rm vis} = \arccos(-b_0/b_1)$. This is exact: averaging over the initial longitude phase is equivalent to averaging over one full rotation period, so $\bar{\mathcal{V}}$ is independent of $t$ regardless of the spot lifetime. This is exactly the zeroth Fourier coefficient $c_0(\Phi)$ of the visibility function derived in Section 4.3.

Since $\bar{\mathcal{V}}$ is constant, it factors out of the time integral, and the remaining integral acts only on the spot-size envelope:

$$\int_{-\infty}^{\infty} \langle A(t) \rangle_\Lambda\, dt = \pi\,\alpha_{\max}^2\,\bar{\mathcal{V}}(\Phi, I) \int_{-\infty}^{\infty} \Gamma(t)\, dt. \tag{73}$$

Averaging over the latitude distribution $p(\Phi)$ and combining, the expected flux for a general envelope $\Gamma(t)$ is:

$$\boxed{\mathbb{E}[F] = 1 - \dot{N}_{\rm spot}\,(1 - f_{\rm spot})\,\alpha_{\max}^2 \left( \int_{-\infty}^{\infty} \Gamma(t)\, dt \right) \int_{-\pi/2}^{\pi/2} \bar{\mathcal{V}}(\Phi, I)\, p(\Phi)\, d\Phi} \tag{74}$$

This is an intrinsic stellar property, independent of both the observing baseline $T_{\rm obs}$ and the rotation parameters $P_{\rm eq}$ and $\kappa$, since the time-averaged visibility $\bar{\mathcal{V}}$ depends only on the inclination and latitude geometry. The only envelope-dependent quantity is $\int \Gamma(t)\, dt$, which has a closed-form expression for each profile considered in this work. Note the contrast with the kernel amplitude (Section 4.5), which involves $R_\Gamma(0) = \int \Gamma^2(t)\, dt$: the mean depends on the first moment of the envelope, while the variance depends on the second. In practice, we set the GP mean function to $m(t) = 0$ and model the mean-subtracted residuals, however we include this derivation for completeness and to show how the mean depends on the physical parameters of the starspot model.

## 6. PHYSICAL INTERPRETATION OF KERNEL HYPERPARAMETERS

### 6.1. *Information Content of the Mean and Covariance*

Before discussing the individual hyperparameters, we illustrate how the single-spot signal connects to the two statistical quantities modeled by the GP: the mean function (first moment) and the kernel (second moment). Figure 12 shows 80 realizations of the single-spot signal $A(t) = \Gamma(t)\,\mathcal{V}(t)$ for random initial longitudes $\Lambda_{\rm ref}$, at fixed inclination $I = 60°$ and latitude $\Phi = 30°$. The ensemble mean over all longitudes is the smooth, aperiodic curve $\langle A(t) \rangle_\Lambda = c_0\,\Gamma(t)$: the spot-size envelope scaled by the zeroth Fourier coefficient of the visibility function. This is the first moment of the process: the quantity that enters the GP mean function (Section 6.1). It depends linearly on $c_0$ and on the first moment of the envelope, $\int \Gamma\, dt$.

The lower panel shows the residuals after subtracting this mean. These oscillatory residuals (produced by the periodic rotation of the spot in and out of view) are what the kernel (second moment) describes. Their variance depends quadratically on the Fourier coefficients ($|c_n|^2$) and on the autocorrelation of the envelope, $R_\Gamma(\tau) = \int \Gamma(t)\,\Gamma(t+\tau)\, dt$. The distinction between these two moments

(linear versus quadratic dependence on the visibility coefficients, and first versus second functional of the envelope) is what allows the mean and kernel to constrain different combinations of the amplitude parameters.





**Figure 12.** Decomposition of the single-spot signal into first and second moments of the GP, for a spot at latitude $\Phi = 30°$ with $I = 60°$, $P_{\rm eq} = 5\,\mathrm{d}$, $\ell_{\rm spot} = 15\,\mathrm{d}$, and $\tau_{\rm spot} = 3\,\mathrm{d}$. *Top:* 80 realizations of $A(t) = \Gamma(t)\,\mathcal{V}(t)$ for random longitudes (blue), with the ensemble mean $\langle A\rangle_\Lambda = c_0\,\Gamma(t)$ (red). The mean depends linearly on $c_0$ and on $\int \Gamma\, dt$ (first moment of the envelope). *Bottom:* Residuals after subtracting the mean, with $\pm\sigma(t)$ envelope (orange). This spread is what the kernel describes; it depends quadratically on $|c_n|^2$ and on the envelope autocorrelation $R_\Gamma(\tau) = \int \Gamma(t)\,\Gamma(t+\tau)\, dt$ (second moment of the envelope).

### 6.2. *Kernel Components and Hyperparameters*

The full kernel (Eq. 59) is a sum over rotation harmonics, and the cleanest way to understand what each hyperparameter does is to keep it that way. Writing

$$\mathcal{K}(\tau) = \sum_{n=0}^{N} \mathcal{K}_n(\tau), \qquad \mathcal{K}_n(\tau) = \sigma_k^2\, s_n \int_{-\pi/2}^{\pi/2} R_\Gamma(\tau)\; |c_n(I,\Phi)|^2 \cos\!\big(n\,\omega_0(\Phi)\,\tau\big)\; p(\Phi)\, d\Phi, \tag{75}$$

with $s_0 = 1$ and $s_n = 2$ for $n \geq 1$, each harmonic term is the product of exactly three factors, and every physical hyperparameter enters through exactly one of them:

- **The envelope autocorrelation**: $R_\Gamma(\tau)$ encodes how long spots live. It carries $\ell_{\rm spot}$ and $\tau_{\rm spot}$, and it multiplies every order identically.
- **The geometry weight**: $|c_n(I,\Phi)|^2$ sets how much variance lands in order $n$. It carries the inclination $I$ and the latitude distribution $p(\Phi)$, and nothing else.
- **The rotation phase**: $\cos(n\,\omega_0(\Phi)\tau)$ carries $P_{\rm eq}$ and $\kappa$ and sets the positions of the kernel peaks.

The amplitude $\sigma_k^2$ multiplies all orders equally. Because the three harmonics are sensitive to disjoint pieces of the physics, they are not three views of the same measurement: which parameters a fit can recover is decided by which orders the data constrain. The subsections below examine each order in turn; all quoted numbers are computed with `spotgp` (`AnalyticKernel.kernel_components`, Gauss–Legendre latitude quadrature) for a fiducial model with $\ell_{\rm spot} = 15\,$d, $\tau_{\rm spot} = 3\,$d, $I = 60°$, $P_{\rm eq} = 5\,$d, uniform $p(\Phi)$, varying one parameter at a time.[7]

Each $\mathcal{K}_n$ is separately a valid (positive semi-definite) kernel: the $n = 0$ term has spectrum $|\hat{\Gamma}(\omega)|^2\langle|c_0|^2\rangle_\Phi \geq 0$, the $n \geq 1$ terms have spectra $[S_\Gamma(\omega - n\omega_0) + S_\Gamma(\omega + n\omega_0)]$ that are shifted copies of the same non-negative function, and a positive mixture over latitude preserves both statements. The harmonics are therefore *independently samplable* stochastic processes whose sum is a draw from the full GP. Figure 13 shows this decomposition in the time domain: the $n = 0$ component is an aperiodic random walk–like dimming set by spot emergence and decay, the $n = 1$ component is the familiar rotational modulation, and the $n = 2$ component is a weaker signal at half the rotation period; their sum is statistically identical to a draw from the full kernel. Figure 14 shows the same decomposition of the kernel and its PSD: at the fiducial inclination the $n = 0$, 1, 2 terms contribute 59%, 37%, and 4% of the variance.

#### 6.2.1. *The $n = 0$ Term: A Direct Readout of the Spot Lifetime*

With no cosine factor, the $n = 0$ term is the envelope autocorrelation times a constant, $\mathcal{K}_0(\tau) = \sigma_k^2\,\langle|c_0|^2\rangle_\Phi\,R_\Gamma(\tau)$. Its shape is a direct readout of how long spots live, uncontaminated by anything else: for the trapezoidal envelope it decays through the piecewise-polynomial breakpoints at $\tau = \tau_{\rm spot}$, $\ell_{\rm spot}$, $\ell_{\rm spot}+\tau_{\rm spot}$ and reaches zero *exactly* at $\tau = \ell_{\rm spot}+2\tau_{\rm spot}$ (21 d for the fiducial model; Appendix D). The plateau carries most of the leverage: doubling $\ell_{\rm spot}$ from 15 to 30 d moves the half-height lag from 8.9 to 16.4 d, while $\tau_{\rm spot}$ enters at roughly half that weight.

Because $P_{\rm eq}$ and $\kappa$ enter only through the cosine factor, the $n = 0$ term is completely blind to both: the normalized $\mathcal{K}_0$ is identical for any rotation period or shear. This invariance is what makes it useful: rotation cannot leak into it, so it is the fixed reference against which the decay of the $n \geq 1$ orders is measured (Section 6.2.4). Its amplitude, by contrast, is pure geometry: $\langle|c_0|^2\rangle_\Phi$ grows by a factor $\sim$5 from edge-on to $I = 10°$ as more of the spot population becomes permanently visible.

The compact support of $\mathcal{K}_0$ also provides a clean model-checking diagnostic. Unlike the exponential or Gaussian decays of phenomenological kernels, which are strictly positive at all lags, the starspot kernel vanishes identically beyond $\ell_{\rm spot}+2\tau_{\rm spot}$; any observed correlation at longer lags must come from other sources (activity cycles, instrument systematics), or indicates a longer spot lifetime. Combined with the rotational recurrences of the $n \geq 1$ terms, this sets the number of visible ACF peaks,

$$N_{\rm peaks} \approx \left\lfloor \frac{\ell_{\rm spot} + 2\tau_{\rm spot}}{P_{\rm eq}} \right\rfloor, \tag{76}$$

[7] The accompanying notebook is `src/notebooks/harmonics_plots.ipynb`.





**Figure 13.** Harmonic decomposition of GP lightcurve samples ($I = 60°$, $P_{\rm eq} = 5\,$d, $\kappa = 0$, $\ell_{\rm spot} = 15\,$d, $\tau_{\rm spot} = 3\,$d). Each of the top three panels is an independent draw from a single harmonic term $\mathcal{K}_n$ of Eq. (75): the aperiodic $n = 0$ (spot emergence and decay), the fundamental $n = 1$ (rotational modulation at $P_{\rm eq}$), and the second harmonic $n = 2$ (at $P_{\rm eq}/2$). Annotations give each order's share of the variance, $\mathcal{K}_n(0)/\mathcal{K}(0)$. Because the terms are independently samplable, their sum (fourth panel) is a draw from the full kernel.





**Figure 14.** Harmonic decomposition of the kernel (left) and its power spectral density (right) for the fiducial model ($P_{\rm eq} = 5\,$d, $\kappa = 0.2$, $I = 60°$, $\ell_{\rm spot} = 15\,$d, $\tau_{\rm spot} = 3\,$d). The $n = 0$ term (blue) is the pure envelope autocorrelation: aperiodic, spanning the low-frequency lobe of the PSD; the $n = 1$ term (orange) carries the rotation peak at $1/P_{\rm eq}$; the $n = 2$ term (pink) carries the first-harmonic peak at $2/P_{\rm eq}$. Annotations give the fractional contribution of each order to the variance $\mathcal{K}(0)$.

the starspot analog of the quality factor of a damped oscillator kernel (in practice the last peak sits on the tail of $R_\Gamma$ and is barely visible).

Finally, the $n = 0$ harmonic is the point of contact between the kernel and the mean function: $c_0 = \bar{\mathcal{V}}$ enters the mean linearly (Section 6.1) and the kernel quadratically. The $|c_0|^2$ floor keeps the kernel non-negative at all lags: each spot only ever dims the star, and the autocorrelation of a one-signed pulse is non-negative, whereas sample ACFs of real lightcurves dip negative at half-period lags. The difference is an estimator effect: subtracting the finite-window time mean biases the sample ACF downward by roughly $\bar{\mathcal{K}} = T^{-1}\int \mathcal{K}\, ds$, which is dominated by the $n = 0$ power. The DC power fraction $f_0 = \mathcal{K}_0(0)/\mathcal{K}(0)$ quantifies how large this effect can be, growing from 0.41 at $I = 90°$ toward unity pole-on. Since the $n = 0$ fluctuations are genuinely stochastic (the number of visible spots varies), the term belongs in the kernel even for mean-subtracted data.

In practice, a second and typically larger effect compounds the estimator bias above. That bias is a statistical artifact of not knowing the true baseline flux level: it would persist even for a perfectly instrumented, unfiltered lightcurve, simply because the sample mean used in its place is itself contaminated by the same low-frequency power it is meant to remove. The second effect is different in kind: the lightcurve detrending applied upstream of the GP fit (e.g., a Savitzky–Golay filter with a window of tens of days) is a real spectral filter, applied by the reduction pipeline before the GP ever sees the data, that removes power at exactly the frequencies the DC bump occupies. The bump has half-width $\sim 1/(\ell_{\rm spot} + 2\tau_{\rm spot})$, comparable to or narrower than the typical detrending window, so the filter does not merely attenuate the $n = 0$ term uniformly; it can carve out most of it. Because $\ell_{\rm spot}$ and $\tau_{\rm spot}$ are shared with the $n \geq 1$ harmonics through $R_\Gamma(\tau)$, forcing the kernel to match its full geometric DC prediction against a lightcurve with that power removed pulls the fitted envelope

timescale away from its true value to compensate; and because the ratio $\langle|c_0|^2\rangle_\Phi/\langle|c_1|^2\rangle_\Phi$ is a principal lever for constraining the inclination (Section 6.2), that constraint degrades along with it.

We therefore do not treat the $n = 0$ amplitude as fixed by the geometry alone. Instead we introduce a free multiplicative scale $s_0$ on the DC term,

$$\mathcal{K}(\tau) = \sigma_k^2\, R_\Gamma(\tau) \left[ s_0 \,\langle|c_0|^2\rangle_\Phi + 2\sum_{n\geq 1}\langle|c_n|^2\rangle_\Phi \,\cos(n\,\omega_0(\Phi)\,\tau)\right], \tag{77}$$

with $s_0 \geq 0$ the only requirement for positive semi-definiteness, since it multiplies an already positive-definite sub-kernel. $s_0 = 1$ recovers the standard kernel (Eq. 75) exactly, and $s_0 = 0$ is equivalent to dropping the $n = 0$ term and retaining only $n \geq 1$ harmonics; intermediate $s_0 < 1$ absorbs whatever fraction of the geometric DC power is missing. We do not attempt to separate the two suppression mechanisms above: the estimator bias and the detrending filter both act on the same term with the same sign, so their combined effect is what a single scale factor can hope to capture, not either one individually.

### 6.2.2. *The $n = 1$ Term: Rotation Phase and the Differential-Rotation Gap*

The fundamental multiplies the same envelope by $\cos(\omega_0\tau)$. Under solid-body rotation this adds no new timescale: at integer multiples of the period the cosine is unity, so the recurrence heights of the $n = 1$ term land exactly on the $n = 0$ envelope,

$$\frac{\mathcal{K}(mP_{\rm eq})}{\mathcal{K}(0)} = \frac{R_\Gamma(mP_{\rm eq})}{R_\Gamma(0)} \qquad (\kappa = 0), \tag{78}$$

which evaluates to 0.741 and 0.432 at one and two periods for the fiducial model. When $mP_{\rm eq}$ falls in the linear regime of $R_\Gamma$ ($\tau_{\rm spot} < mP_{\rm eq} < \ell_{\rm spot}$), the peak heights decay linearly,

$$\frac{\mathcal{K}(mP_{\rm eq})}{\mathcal{K}(0)} = \frac{\ell_{\rm spot} + 2\tau_{\rm spot}/3 - mP_{\rm eq}}{\ell_{\rm spot} + 2\tau_{\rm spot}/5}, \tag{79}$$

a distinctive prediction of the trapezoidal profile (exponential profiles give exponentially decaying peaks). Under solid-body rotation, then, all of the *new* information in the fundamental is in where its peaks sit, not how tall they are: peak positions measure $P_{\rm eq}$, peak heights re-measure the envelope.

Differential rotation is what breaks this tie. Spots at latitude $\Phi$ accumulate phase $\omega_0(\Phi)\tau$, with a characteristic spread across the latitude distribution of

$$\Delta\phi(\tau) = \frac{2\pi\,|\kappa|\,\tau}{P_{\rm eq}}\,\langle\sin^2\Phi\rangle_\Phi \tag{80}$$

(the full range across latitudes is $2\pi|\kappa|\tau/P_{\rm eq}$). Because $\cos(\omega_0(\Phi)\tau) = 1$ for every latitude at $\tau = 0$, the shear leaves $\mathcal{K}_1(0)$ and the variance shares untouched at any $\kappa$; what it does instead is pull the recurrence heights *below* the $n = 0$ envelope and skew them toward longer lags (high-latitude spots recur later for $\kappa > 0$). For the fiducial geometry the first recurrence retains only 0.58 of its solid-body height at $\kappa = 0.4$. **The gap between the $n = 1$ recurrence heights and the $n = 0$ envelope is the shear; nothing else opens it.** Setting $\Delta\phi \approx \pi$ gives the coherence lag

$$\tau_{\rm coh} \approx \frac{P_{\rm eq}}{2\,|\kappa|\,\langle\sin^2\Phi\rangle_\Phi}, \tag{81}$$

beyond which the rotational recurrences are strongly suppressed (roughly halved, for the fiducial latitude distribution) even where $R_\Gamma$ has not yet decayed: differential rotation imposes a second coherence timescale competing with the spot lifetime. The dephasing depends only on $|\kappa|$; the sign of $\kappa$ appears solely in the direction of the peak skew. Figure 15 shows this decoherence in both the lightcurves and the ACF: as $|\kappa|$ increases, GP samples lose their periodic pattern and the ACF recurrences broaden and wash out, while the $R_\Gamma$ envelope (the $n = 0$ reference) remains unchanged.





**Figure 15.** Decoherence from differential rotation. *Left:* Three GP lightcurve samples drawn from the kernel for each $\kappa$ value. *Right:* Normalized kernel ACF (solid) with the envelope autocorrelation $R_\Gamma(\tau)/R_\Gamma(0)$ (dashed). For $\kappa = 0$ (solid-body rotation), the lightcurves are highly periodic and the ACF recurrences ride the $n = 0$ envelope. As $|\kappa|$ increases, spots at different latitudes rotate at different rates and the periodic signal decoheres: at $|\kappa| = 0.5$ the signal is noticeably aperiodic beyond the first two recurrences ($\tau_{\rm coh} \approx 10\,$d from Eq. 81), and at $|\kappa| = 1$ only the first survives ($\tau_{\rm coh} \approx 5\,$d). The envelope is unchanged throughout, since it depends only on the spot lifetime parameters; the widening gap between the recurrence heights and the envelope is the shear signal. The decoherence is symmetric in $\pm\kappa$. Parameters: $P_{\rm eq} = 5\,$d, $I = 90°$, $\ell_{\rm spot} = 30\,$d, $\tau_{\rm spot} = 3\,$d.

Two practical caveats follow directly from the factorization. First, short-lived spots bias the recovered period low: at $\ell_{\rm spot} = 5\,$d the first kernel recurrence sits at 4.73 d for a true 5.0 d period, because the envelope decays across the peak. Second, when the spot lifetime falls below the rotation period the recurrence lands where the envelope has already died (at $P_{\rm eq} = 20\,$d with a 21 d lifetime, $\mathcal{K}_1(P_{\rm eq})$ is zero to three decimal places), and no rotational information survives at any inclination. The amplitude of the fundamental is the mirror image of the $n = 0$ term: half the variance equator-on, collapsing to 1.4% by $I = 10°$.

### 6.2.3. *The $n = 2$ Term: Evidence That Spots Set*

A spot that never sets has projected area exactly $b_0 + b_1\cos(\omega_0 t)$: a pure sinusoid, with no second harmonic to give. Only the clipping of the visibility function at the limb generates power at $n \geq 2$ (Appendix B). The second harmonic is therefore not a smaller copy of the fundamental; it is a statement that spots are disappearing behind the star. The kernel is unambiguous about this: restricting spots to circumpolar latitudes ($\Phi \geq I$; e.g. a 60°–90° band at $I = 60°$) drives $\mathcal{K}_2(0)$ to zero exactly while the fundamental retains 12% of the variance.

The share of variance at $n = 2$ is small everywhere: 9% of the total at best (equator-on), 3.7% at the fiducial $I = 60°$, and $\sim 10^{-4}$ pole-on, so detecting it is a signal-to-noise question before it is a physics question. When it is detected, it adds two things the fundamental cannot. Its recurrence at $P_{\rm eq}/2$ survives envelopes too short for the fundamental (in the $P_{\rm eq} = 20\,$d example above, where $\mathcal{K}_1$ has died, $\mathcal{K}_2$ still retains 0.43 of its zero-lag value at half a period). And the ratio of the harmonic weights,

$$\mathcal{H} \equiv \frac{\langle|c_2|^2\rangle_\Phi}{\langle|c_1|^2\rangle_\Phi}, \tag{82}$$

is a clean inclination diagnostic: $\mathcal{H} = (4/(3\pi))^2 \approx 0.18$ for edge-on viewing with uniform latitudes, falling by about 5× the magnitude as $I = 30°$. In the lightcurve, a significant $n = 2$ share manifests as the familiar "double-dip" shape: two flux minima per rotation.

The same weights control how deeply the kernel oscillates.[8] Writing the extrema of the cosine sum explicitly, the kernel is bounded by $k_{\rm upper}(\tau) = R_\Gamma(\tau)\,[\langle|c_0|^2\rangle + 2\langle|c_1|^2\rangle + 2\langle|c_2|^2\rangle]$ at integer periods and $k_{\rm lower}(\tau) = R_\Gamma(\tau)\,[\langle|c_0|^2\rangle - 2\langle|c_1|^2\rangle + 2\langle|c_2|^2\rangle]$ at half periods, so the fractional oscillation depth

$$\frac{k_{\rm lower}}{k_{\rm upper}} = \frac{\langle|c_0|^2\rangle - 2\langle|c_1|^2\rangle + 2\langle|c_2|^2\rangle}{\langle|c_0|^2\rangle + 2\langle|c_1|^2\rangle + 2\langle|c_2|^2\rangle} \tag{83}$$

is set entirely by the geometry weights: $\approx -0.005$ edge-on (the kernel oscillates essentially to zero at the troughs), +0.25 at $I = 60°$, and +0.77 at $I = 30°$, where the DC floor dominates and the oscillation is a shallow ripple on an aperiodic decay.

### 6.2.4. *Differential Rotation Across the Harmonics*

At fixed latitude, the PSD sideband of order $n$ has width proportional to $n\kappa$, and the textbook expectation is that higher orders decohere proportionally faster. The kernel says otherwise, because each order samples latitude through its own weight $|c_n(I,\Phi)|^2$, and the shear $\kappa\sin^2\Phi$ doing the smearing is itself latitude-dependent. Higher orders need spots that set, so $|c_2|^2$ concentrates toward

[8] Numerical check: `lower_envelope_executed.ipynb`

the equator, precisely where the shear is weakest. The two effects run in opposite directions, and which one wins depends on the inclination. Quantifying the dephasing rate of order $n$ by $n\,\sigma_\omega^{(n)}$, where $\sigma_\omega^{(n)}$ is the $|c_n|^2$-weighted spread of $\omega_0(\Phi)$:

**Table 4.** Latitude sampling and dephasing rate of the $n = 1$ and $n = 2$ harmonics for $\kappa = 0.4$ and a uniform latitude distribution. $\langle|\Phi|\rangle_n$ is the $|c_n|^2$-weighted mean absolute latitude; $n\,\sigma_\omega^{(n)}$ [rad d$^{-1}$] is the dephasing rate.

| Inclination | $\langle\|\Phi\|\rangle_1$ / $\langle\|\Phi\|\rangle_2$ | $n\,\sigma_\omega^{(n)}$, $n=1$ / $n=2$ | Outcome |
|---|---|---|---|
| $I = 90°$ | 26.8° / 26.8° | 0.126 / 0.251 | factor 2, as expected |
| $I = 60°$ | 31.0° / 18.0° | 0.139 / 0.140 | cancels almost exactly |
| $I = 30°$ | 30.7° / 8.4° | 0.127 / 0.037 | $n = 2$ far more coherent |

Only equator-on does the naïve $n$-scaling survive (Table 4). At the fiducial $I = 60°$ the latitude weighting cancels the $n$-scaling almost exactly, and by $I = 30°$ the second harmonic is the *more* coherent of the two: at $\tau = P_{\rm eq}$ with $\kappa = 0.4$, the $n = 1$ recurrence retains 0.58 of the $n = 0$ envelope while $n = 2$ retains 0.67. The same latitude weighting shifts the apparent rates: each order reports the mean rotation rate of the latitudes it samples, $\langle\omega_0\rangle_n = 0.975,\ 1.101,\ 1.192\,\mathrm{rad\,d^{-1}}$ for $n = 0, 1, 2$ at $I = 60°$, $\kappa = 0.4$ (the $n = 0$ weighting samples the highest latitudes, the $n = 2$ weighting the lowest). The spread of apparent periods *between* the orders is therefore itself a measurement of the shear, complementary to the peak damping of Section 6.2.2.

### 6.2.5. *Which Parameter Moves Which Harmonic*

**Table 5.** Which hyperparameter affects which harmonic term, and how. "Amplitude" entries change only the term's overall scale; "shape" entries change its dependence on $\tau$.

| Parameter | $n = 0$ | $n = 1, 2$ | Effect |
|---|---|---|---|
| $\sigma_k$ | amplitude | amplitude | rescales all orders; shares and shapes unchanged |
| $\ell_{\rm spot}$ | shape + amp. | shape + amp. | common envelope; support ends at $\ell_{\rm spot} + 2\tau_{\rm spot}$ |
| $\tau_{\rm spot}$ | shape + amp. | shape + amp. | as $\ell_{\rm spot}$, at roughly half the leverage |
| $I$ | amplitude | amplitude | redistributes variance between orders; no shape change |
| $p(\Phi)$ | amplitude | amplitude$^\dagger$ | same lever as $I$; degenerate with it |
| $P_{\rm eq}$ | – | shape | recurrences at $P_{\rm eq}$ and $P_{\rm eq}/2$ |
| $\kappa$ | – | shape | damps recurrences below the $n = 0$ envelope; skews them later |

NOTE—$^\dagger$For $\kappa \neq 0$ the latitude distribution also enters the $n \geq 1$ shapes through the sampled range of $\omega_0(\Phi)$ (Section 6.2.4).

Four consequences for inference follow directly:

1. **Spot lifetime and shear are separable, but only jointly.** Both damp the ACF, but only the shear makes the $n \geq 1$ terms decay faster than the $n = 0$ term. The comparison between the orders (not either curve alone) carries $\kappa$.
2. **Inclination and the latitude distribution are one lever, not two.** Both act purely through $|c_n|^2$: an equatorial band viewed at $I = 60°$ and a full-latitude star viewed edge-on

put nearly identical power in the fundamental ($n = 1$ share $\approx 0.50$ in both cases). Without a latitude prior, a fitted inclination is really a fitted geometry weight (cf. the Kepler-63 analysis of Paper II, where transit-mapped spot latitudes provide exactly such a prior).

3. $\kappa$ **is never constrained alone.** Dephasing needs a lever arm in latitude: at $\kappa = 0.4$ an equatorial band barely damps the first recurrence at all (0.712 versus 0.741 for solid body), while a polar band moves it from 5.0 to 7.4 d. What the data constrain is $\kappa$ times the occupied latitude spread: the same degeneracy noted for individual-spot modeling by Basri & Shah (2020).
4. **Rotation is measurable only while the envelope survives it.** When $\ell_{\rm spot} + 2\tau_{\rm spot} \lesssim P_{\rm eq}$, the first recurrence lands beyond the envelope's support and $P_{\rm eq}$, $\kappa$, and $I$ all lose their handles at once, leaving only the weak half-period recurrence of $n = 2$. This is the regime, identified independently by the Cramér–Rao analysis of Section 10.1, in which the kernel degrades gracefully into a purely aperiodic model.

### 6.3. *Assumed Distributions of Spot Parameters*

Both the numerical and the analytic kernel are averages over the random parameters of the spot population, so the kernel we derive is specific to the distributions assumed for those parameters. Table 6 lists the distribution we adopt in this paper for each parameter of Tables 1 and 2. Only the reference time, longitude, and latitude of each spot are random. Spots emerge independently of one another, and these three variables are drawn independently for each spot. All other spot properties take the same value for every spot.

**Table 6.** Distributions assumed for the spot and stellar parameters in this paper.

| Parameter | Assumed distribution |
|---|---|
| | *Random for each spot* |
| $t_{{\rm ref},k}$ | Poisson process with uniform random spot emergence rate |
| $\Lambda_{{\rm ref},k}$ | uniform on $[0, 2\pi)$ |
| $\Phi_{{\rm ref},k}$ | uniform in latitude, $p(\Phi) = 1/\pi$ on $[-\pi/2, \pi/2]$ |
| | *Identical for every spot* |
| $\ell_{\rm spot}$, $\tau_{\rm spot}$ | fixed, with $\tau_{\rm em} = \tau_{\rm dec} = \tau_{\rm spot}$ |
| $\alpha_{\rm max}$, $f_{\rm spot}$ | fixed |
| $P_{\rm eq}$, $\kappa$, $I$ | fixed (properties of the star) |

Each of these parameters enters the derivation of Section 4 at a different step. Averaging over the emergence times $t_{{\rm ref},k}$ removes the phase of each spot's spectrum, so the kernel depends on the spot envelope only through its autocorrelation $R_\Gamma(\tau)$, whose shape is set by $\ell_{\rm spot}$ and $\tau_{\rm spot}$ (Section 4.6). Averaging over the longitudes $\Lambda_{{\rm ref},k}$ cancels the cross terms between different harmonics of the visibility function (Eq. 49). The latitude distribution $p(\Phi)$ is the weight in the remaining integral over latitude (Eq. 59), and the stellar parameters set the integrand: $P_{\rm eq}$ and $\kappa$ give the rotation frequency $\omega_0(\Phi)$ of a spot at each latitude, and $I$ together with $\Phi$ gives the Fourier coefficients $c_n(I, \Phi)$ of its visibility function. The spot size $\alpha_{\rm max}$ and contrast $f_{\rm spot}$ only scale the kernel amplitude $\sigma_k$ (Eq. 63).

Of these, the emergence-time and longitude distributions are built into the form of the kernel. A constant emergence rate makes the kernel stationary, and the cross terms between harmonics cancel

only if the longitudes are uniform. An activity cycle (a time-varying emergence rate) or active longitudes that persist in the rotating frame would make the process non-stationary, which calls for a different kernel rather than a different parameter distribution. The latitude distribution is a modeling choice: in the analysis of this paper we assume a uniform latitude destribution, however this assumption can easily be modified for distributions (e.g. to assume active spot latitudes as oserved on the Sun).

Each of these assumptions is a separate component of our `spotgp` implementation. Users can change the latitude distribution, the envelope, or the visibility function by defining their own subclass of `LatitudeDistributionFunction`, `EnvelopeFunction`, or `VisibilityFunction`, without modifying the kernel or solver code. Birky et al. (2026) describes these interfaces.

## 7. GP SOLVER IMPLEMENTATION

In order to solve for the GP posterior and perform inference on the hyperparameters $\theta$, we need to evaluate the GP marginal log-likelihood (Eq. 84), which requires computing the kernel matrix inversion. In this section, we show how the structure of our kernel allows for efficient computation using a banded Cholesky solver, which reduces the computational cost from $\mathcal{O}(N^3)$ to $\mathcal{O}(Nb^2)$, where $b$ is the bandwidth determined by the spot lifetime and observing cadence. We implement this solver in JAX, enabling fast automatic differentiation for MCMC sampling of the hyperparameters.

### 7.1. *Naive Cholesky Solver*

We implement a JAX-accelerated GP solver that uses the analytic kernel for Bayesian inference on stellar lightcurves. The GP marginal log-likelihood for $N$ observations $\mathbf{y}$ at times $\mathbf{x}$ is

$$\ln p(\mathbf{y} \mid \theta) = -\frac{1}{2}\left[(\mathbf{y}-\boldsymbol{\mu})^\top K_\theta^{-1}(\mathbf{y}-\boldsymbol{\mu}) + \ln|K_\theta| + N\ln 2\pi\right], \tag{84}$$

where $K_\theta$ is the $N \times N$ covariance matrix with entries $[K_\theta]_{ij} = k(|x_i - x_j|;\, \theta) + (\sigma_{\mathrm{err},i}^2 + \sigma_n^2)\,\delta_{ij}$, $\boldsymbol{\mu}$ is the mean vector, and $\sigma_n$ is an optional white-noise amplitude.

Evaluating Eq. (84) involves two computationally distinct steps:

1. *Kernel evaluation*: computing $k(|x_i - x_j|;\, \theta)$ for all pairs, which requires the latitude integral in Eq. (59). We perform this integral using Gauss-Legendre quadrature with $n_{\mathrm{lat}} = 64$ nodes. To avoid materializing all latitude contributions simultaneously (which would require an $n_{\mathrm{lat}} \times M$ intermediate tensor, where $M$ is the number of unique lags), we accumulate the weighted latitude contributions sequentially, reducing peak memory from $\mathcal{O}(n_{\mathrm{lat}} \times M)$ to $\mathcal{O}(M)$. We further reduce memory by exploiting the symmetry of the covariance matrix, evaluating the kernel only on the $N(N+1)/2$ upper-triangular lags rather than the full $N^2$ matrix.
2. *Cholesky factorization and solve*: decomposing $K_\theta = LL^\top$ and solving for $L^{-1}(\mathbf{y} - \boldsymbol{\mu})$, which dominates at $\mathcal{O}(N^3)$.

Our implementation provides two interfaces. The `GPSolver` class pre-computes and caches the Cholesky factorization at construction time, enabling fast repeated queries (predictions, log-likelihood evaluation) for fixed hyperparameters. For MCMC sampling, where hyperparameters change at every step, we provide a pure-functional JIT-compiled log-likelihood function that recomputes the full pipeline (kernel evaluation, Cholesky, and solve) from scratch on each call. JAX's just-in-time compilation and automatic differentiation provide efficient gradient computation for Hamiltonian Monte Carlo (HMC) and related samplers.

**Table 7.** GPSolver wall-clock time and peak memory as a function of the number of data points $N$, irregularly spaced over 100 d, with the full (dense) Cholesky solver. *Build*: covariance matrix + Cholesky factorization. *Log-likelihood*: JIT-compiled evaluation (kernel + Cholesky + solve) as called during MCMC. *Predict*: predictive mean at 200 test points. Times are the median of 5 runs after JIT warmup, on one core of an AMD EPYC 7742 64-Core Processor. Memory is the peak RSS increase during each operation, measured in an isolated subprocess.

| | Build | | Log-likelihood | | Predict | |
|---|---|---|---|---|---|---|
| $N$ | Time | Memory | Time | Memory | Time | Memory |
| 10 | 8.61 ms | 0.4 MB | 208 $\mu$s | 0.1 MB | 17.0 ms | 4 MB |
| 100 | 33.4 ms | 12 MB | 20.1 ms | 0.2 MB | 195 ms | 82 MB |
| 1,000 | 4.49 s | 2.10 GB | 1.95 s | 284 MB | 1.81 s | 824 MB |
| 5,000 | 111 s | 51.57 GB | 45.7 s | 674 MB | 8.93 s | 4.02 GB |
| 10,000 | 251 s | 131.26 GB | 183 s | 707 MB | 19.2 s | 8.03 GB |

Table 7 benchmarks the wall-clock time and peak memory usage of the GP solver as a function of the number of data points $N$. The "Build" column measures the `GPSolver` construction (kernel evaluation + Cholesky), while the "Log-likelihood" column measures the JIT-compiled pure-functional evaluation used during MCMC. After JIT warmup, the log-likelihood evaluation is faster than the class-based build because XLA reuses pre-allocated buffers. At $N = 1000$, the full log-likelihood evaluation takes 1.95 s on a single CPU core, making MCMC chains of $\sim 10^4$ samples feasible within hours. The $\mathcal{O}(N^2)$ memory scaling of the covariance matrix is the primary limitation for larger datasets.

### 7.2. *Banded Cholesky Solver*

The compact support of the spot-lifetime envelope $R_\Gamma(\tau)$ (Section 6) guarantees that the kernel vanishes exactly for lags $|\tau| > \tau_{\max} \equiv \ell_{\rm spot} + 2\tau_{\rm spot}$. For uniformly sampled data with cadence $\Delta t$, this means the covariance matrix $K_\theta$ is banded with bandwidth

$$b = \left\lceil \frac{\ell_{\rm spot} + 2\tau_{\rm spot}}{\Delta t} \right\rceil , \tag{85}$$

i.e., $[K_\theta]_{ij} = 0$ whenever $|i - j| > b$. For typical stellar monitoring campaigns where the observing baseline greatly exceeds the spot lifetime ($T_{\rm obs} \gg \tau_{\max}$), the bandwidth satisfies $b \ll N$, and the covariance matrix is sparse.

We exploit this banded structure to reduce the cost of the Cholesky factorization from $\mathcal{O}(N^3)$ to $\mathcal{O}(Nb^2)$ and the memory from $\mathcal{O}(N^2)$ to $\mathcal{O}(Nb)$ (Golub & Van Loan 2013, Section 4.3.5). The lower Cholesky factor $L$ inherits the banded structure of $K_\theta$ (i.e., $L_{ij} = 0$ for $i - j > b$), so both the factorization and the subsequent forward/backward substitution can be restricted to within the band.

We store $L$ in compact banded form as a $(b + 1) \times N$ array $\tilde{L}$, where

$$\tilde{L}_{d,\,j} = L_{j+d,\,j}, \qquad d = 0, 1, \ldots, b, \tag{86}$$

so that the diagonal occupies row $d = 0$ and the $d$-th sub-diagonal occupies row $d$. The kernel is evaluated directly into this compact storage, computing only the $\mathcal{O}(Nb)$ entries within the band rather than the full $N^2$ matrix. The log-determinant is then $\ln|K_\theta| = 2\sum_j \ln \tilde{L}_{0,j}$.





**Figure 16.** Scaling of the GP solver with bandwidth $b$ for fixed $N = 1000$. The banded Cholesky solver scales as $\mathcal{O}(Nb^2)$, while the full Cholesky scales as $\mathcal{O}(N^3)$. For typical spot lifetimes and observing cadences, the bandwidth is small ($b \ll N$), leading to significant speedups.

**Table 8.** GPSolver wall-clock time: `cholesky_full` vs. `cholesky_banded`, for data sampled uniformly at $\Delta t = 0.1$ d over a baseline $x_{\max}$. *Build*: kernel evaluation + Cholesky factorization. *Predict*: GP predictive mean at 200 test points. *Speedup*: full over banded build time. Times are the median of 3 runs after JIT warmup, on an NVIDIA GeForce RTX 5070 Ti GPU. The full Cholesky solver scales as $\mathcal{O}(N^3)$ and was not run above $N = 10{,}000$, where its covariance matrix no longer fits in GPU memory; the banded solver scales as $\mathcal{O}(Nb^2)$ and builds the $N = 100{,}000$ solver in 615 ms. OOM: the run exceeded the GPU's memory.

| | | Full Cholesky | | Banded Cholesky | | |
|---|---|---|---|---|---|---|
| $x_{\max}$ [d] | $N$ | Build | Predict | Build | Predict | Speedup |
| 10 | 100 | 16.1 ms | 25.9 ms | 17.2 ms | 24.4 ms | 0.9× |
| 50 | 500 | 17.4 ms | 23.8 ms | 21.6 ms | 25.4 ms | 0.8× |
| 100 | 1,000 | 19.8 ms | 28.6 ms | 24.8 ms | 29.5 ms | 0.8× |
| 500 | 5,000 | 112 ms | 97.4 ms | 52.6 ms | 72.9 ms | 2.1× |
| 1,000 | 10,000 | OOM | OOM | 90.3 ms | 126 ms | — |
| 5,000 | 50,000 | — | — | 324 ms | OOM | — |
| 10,000 | 100,000 | — | — | 615 ms | OOM | — |

We implement the banded Cholesky factorization using `jax.lax.scan` over columns: at each column $j$, the diagonal and the $b$ sub-diagonal entries are updated using only the $(b \times b)$ block of previously computed entries within the band. The entire factorization is JIT-compiled and end-to-end differentiable via JAX's automatic differentiation, enabling gradient-based sampling with no

additional implementation effort. The forward and backward triangular solves ($L\mathbf{y} = \mathbf{r}$ and $L^\top \mathbf{x} = \mathbf{y}$) similarly operate on the compact storage, restricting each dot product to at most $b$ terms.

We implement the kernel derivation and GP inference framework described above in the open-source Python package `spotgp`. The package is designed with a modular architecture so that each physical assumption (envelope shape, amplitude parameterization, latitude distribution) can be swapped independently without modifying the kernel or solver code. For a detailed description of the code modules and how to implement custom kernel components, we refer the reader to our software paper (Birky et al. 2026).





**Figure 17.** Probabilistic graphical model for the photometric starspot GP. Free parameters (open circles) determine the photometric kernel $\mathcal{K}_{\rm spot}(\tau)$ (Equation 59), which governs the temporal covariance of the flux observations $F_i$ across $N_t$ epochs. The rotation parameters ($P_{\rm eq}$, $\kappa$, $I$) control the harmonic structure, the evolution parameters ($\ell_{\rm sp}$, $\tau_{\rm sp}$) control the temporal envelope $R_\Gamma(\tau)$, and the amplitude parameter $\sigma_k$ controls the overall variability level. The spot latitude distribution $p(\Phi)$ (filled dot) enters the kernel as a fixed input from stellar models.

## 8. MCMC SAMPLING THE KERNEL HYPERPARAMETERS

Figure 17 summarizes the probabilistic structure of the model, showing how the physical parameters determine the photometric kernel and how the flux observations derive from it.

Given a set of observations $\{t_i, y_i, \sigma_i\}_{i=1}^N$, we sample the posterior distribution of the kernel hyperparameters $\boldsymbol{\theta} = (P_{\rm eq}, \kappa, I, \ell_{\rm spot}, \tau_{\rm spot}, \sigma_k)$ using the No-U-Turn Sampler (NUTS; Hoffman & Gelman 2014), a variant of Hamiltonian Monte Carlo (HMC) that automatically tunes the trajectory length. We implement this using the `BlackJAX` library (Cabezas et al. 2024), which provides a JAX-native NUTS kernel that is fully compatible with the JIT-compiled, autodifferentiable log-posterior described below.

### 8.1. *Log-Posterior and Gradients*

The GP marginal log-likelihood is

$$\ln \mathcal{L}(\boldsymbol{\theta}) = -\frac{1}{2}\left[\mathbf{r}^\top K_\theta^{-1}\mathbf{r} + \ln|K_\theta| + N\ln(2\pi)\right], \tag{87}$$





**Figure 18.** Soft uniform prior for a single parameter $\theta$ with bounds $[a, b]$. The sigmoid-based formulation provides a smooth transition at the boundaries while maintaining an approximately uniform density within the bounds.

where $\mathbf{r} = \mathbf{y} - \boldsymbol{\mu}$ is the residual vector and $K_\theta = K(\boldsymbol{\theta}) + \mathrm{diag}(\boldsymbol{\sigma}^2)$ is the noise-augmented covariance matrix, with $\boldsymbol{\sigma}$ the per-point measurement uncertainties. No fixed jitter is added to the diagonal; white noise beyond $\boldsymbol{\sigma}$ is instead modeled as an additive kernel term (Paper II). In our fits to simulated lightcurves the noise level is known, so it enters only through $\boldsymbol{\sigma}$, with $\sigma_i = \sigma_n$. The Cholesky factorization $K_\theta = LL^\top$ is used for efficient computation of both the data-fit term and the log-determinant: $\ln |K_\theta| = 2 \sum_i \ln L_{ii}$.

The entire computational graph, from hyperparameters $\boldsymbol{\theta}$ through the kernel evaluation (Eq. 59), covariance matrix assembly, Cholesky decomposition, and log-likelihood, is implemented in JAX and is end-to-end differentiable. The gradient $\nabla_{\boldsymbol{\theta}} \ln p(\boldsymbol{\theta}|\mathbf{y})$ is computed via reverse-mode automatic differentiation, providing exact gradients at roughly the cost of a single likelihood evaluation.

### 8.2. *Prior*

We adopt a soft uniform prior that enforces parameter bounds while maintaining differentiability. For each parameter $\theta_i$ with bounds $[a_i, b_i]$:

$$\ln p(\boldsymbol{\theta}) = \sum_i \left[ \ln \sigma(k(\theta_i - a_i)) + \ln \sigma(k(b_i - \theta_i)) - \ln(b_i - a_i) \right], \tag{88}$$

where $\sigma(\cdot)$ is the sigmoid function and $k = 100$ controls the steepness of the boundary (Figure 18). This provides effectively uniform support within the bounds while keeping the gradient finite everywhere, which is critical for HMC-based samplers.

### 8.3. *Mass Matrix Estimation*

The efficiency of NUTS depends on the choice of mass matrix $M$, which preconditions the Hamiltonian dynamics to account for differences in the scale and correlation of the parameters. An identity mass matrix performs poorly when the hyperparameters span vastly different scales (e.g., $P_{\rm eq} \sim 5$ d vs. $\alpha_{\rm max} \sim 0.1$ rad). We provide three methods for estimating the inverse mass matrix $M^{-1}$:

1. **Hessian at the MAP.** We first find the maximum a posteriori estimate $\boldsymbol{\theta}_{\rm MAP}$ via L-BFGS-B using JAX-computed gradients, then compute the Hessian of the negative log-likelihood at that

point using `jax.hessian`:

$$M^{-1} = \left[ \nabla^2_{\boldsymbol{\theta}} (-\ln \mathcal{L}) \Big|_{\boldsymbol{\theta}_{\mathrm{MAP}}} \right]^{-1}. \tag{89}$$

The Hessian is regularized by clamping its eigenvalues to be at least $10^{-6}$, ensuring positive definiteness.

2. **Fisher information.** The expected Fisher information matrix for a GP with covariance $K_\theta$ is

$$\mathcal{I}_{ij} = \frac{1}{2} \mathrm{tr} \left( K_\theta^{-1} \frac{\partial K_\theta}{\partial \theta_i} K_\theta^{-1} \frac{\partial K_\theta}{\partial \theta_j} \right). \tag{90}$$

The kernel derivatives $\partial K_\theta / \partial \theta_i$ are computed via forward-mode autodiff (`jax.jacfwd`), which is memory-efficient for the case of few parameters mapping to a large covariance matrix. The inverse mass matrix is then $M^{-1} = \mathcal{I}^{-1}$.

3. **Laplace approximation.** This is equivalent to the Hessian method but additionally provides a Gaussian approximation to the posterior, $p(\boldsymbol{\theta}|\mathbf{y}) \approx \mathcal{N}(\boldsymbol{\theta}_{\mathrm{MAP}}, M^{-1})$, which can be used for rapid approximate inference or as a proposal distribution.

In practice, for NUTS we extract the diagonal of the full inverse mass matrix and use a diagonal mass matrix, which is more numerically robust. Extreme values are clamped to within a factor of $10^4$ of the median diagonal element to prevent instability.

### 8.4. *Sampler Backends*

Because the `spotgp` likelihood and its gradients are implemented as pure JAX functions, they can be readily coupled to any sampler that accepts a callable log-probability (and, optionally, its gradient). We currently provide built-in interfaces to two complementary backends:

- **BlackJAX NUTS.** The default sampler is the NUTS kernel from `BlackJAX` (Cabezas et al. 2024), which exploits the exact gradients provided by JAX autodifferentiation to efficiently explore the posterior. This is the recommended choice when the posterior is unimodal and the mass matrix estimation described in Section 8.3 provides adequate preconditioning.

- **Dynesty nested sampling.** For problems where the posterior may be multimodal or where one additionally requires a Bayesian evidence estimate, `spotgp` also supports the `dynesty` dynamic nested sampler (Speagle 2020). Because nested sampling does not require gradient information, this backend uses only the log-likelihood and prior transform, making it a useful complement to NUTS when gradient-based sampling encounters difficulties such as highly correlated or multi-modal posteriors.

Users may select the sampler backend at runtime, and the interface is designed so that additional samplers can be integrated with minimal effort by supplying the appropriate log-probability and prior functions.

# 9. NUMERICAL RESULTS

## 9.1. *Analytic vs. Numerical Kernel*

In this Section, we compare the analytic kernel derived in Section 4 to a numerical kernel computed by simulating many random spot realizations and empirically estimating their autocorrelation (Eq. 22). For the numerical kernel, we simulate $N_{\rm sim} = 500$ realizations of the flux deficit $\delta F(t)$, each with $N_{\rm spot} = 10$ spots drawn from the same parameter distribution, computed with a cadence of 0.05 days for a duration of 200 days. We then compute the sample autocorrelation function across all simulations to obtain an empirical estimate of the kernel. For the analytic kernel, we evaluate Eq. (59) numerically for the same set of parameters.

Table 9 summarizes the parameter combinations tested and the mean squared error (MSE) between the analytic and numerical autocorrelation functions. For each row, a single parameter is varied from its baseline value ($P_{\rm eq} = 5$ d, $\kappa = 0$, $I = 90°$, $\ell_{\rm spot} = 15$ d, $\tau_{\rm spot} = 3$ d, $\alpha_{\rm max} = 0.1$ rad), with $N_{\rm spot} = 10$ held fixed throughout.

**Table 9.** Mean squared error (MSE) of the analytic kernel ACF compared to numerical Monte Carlo simulations, including stellar limb darkening.

| Varied param. | Value | MSE |
|---|---|---|
| $P_{\rm eq}$ | 3 | $3.59 \times 10^{-4}$ |
| | 5 | $7.76 \times 10^{-4}$ |
| | 10 | $9.59 \times 10^{-4}$ |
| $\kappa$ | $-1.0$ | $8.07 \times 10^{-4}$ |
| | 0.0 | $5.77 \times 10^{-4}$ |
| | 1.0 | $7.32 \times 10^{-4}$ |
| $I$ | 30° | $1.18 \times 10^{-3}$ |
| | 60° | $1.05 \times 10^{-3}$ |
| | 90° | $7.92 \times 10^{-4}$ |
| $\ell_{\rm spot}$ | 5 | $1.47 \times 10^{-4}$ |
| | 15 | $5.68 \times 10^{-4}$ |
| | 30 | $1.62 \times 10^{-3}$ |
| $\tau_{\rm spot}$ | 1 | $6.42 \times 10^{-4}$ |
| | 3 | $7.34 \times 10^{-4}$ |
| | 8 | $9.58 \times 10^{-4}$ |

Figures 19 shows the analytic vs. numerical kernel comparison for each parameter variation. In all cases, the analytic kernel closely matches the numerical kernel, with MSE values on the order of $10^{-3}$ or smaller. Overall, these results validate our analytic kernel against numerical simulations across a range of physically relevant parameters.

Table 10 benchmarks the computational speed of evaluating the analytic kernel vs. computing the numerical kernel. The analytic kernel is at least an order of magnitude faster to evaluate than the numerical kernel, which requires simulating many spot realizations and computing their autocorrelation. This speed advantage makes the analytic kernel practical for use in GP regression on large datasets, where repeated kernel evaluations are required.

**Table 10.** A comparison showing the kernel computation time using the analytic and numerical approaches. The analytic kernel is evaluated at varying lag-array sizes (with $N$ simulated datapoints); the numerical (Monte Carlo) kernel uses 200 and 500 simulations over 100 d. Analytic times are the median of 5 runs after JIT warmup and numerical times are single runs, on one core of an AMD EPYC 7742 64-Core Processor.

| | | $N_{\rm sim} = 200$ | | $N_{\rm sim} = 500$ | |
|---|---|---|---|---|---|
| $N$ | Analytic | Numerical | Speedup | Numerical | Speedup |
| 50 | 5.25 ms | 350 ms | 67× | 463 ms | 88× |
| 100 | 5.48 ms | 343 ms | 63× | 484 ms | 88× |
| 500 | 6.68 ms | 418 ms | 63× | 663 ms | 99× |
| 1,000 | 8.50 ms | 501 ms | 59× | 873 ms | 103× |
| 5,000 | 24.8 ms | 1.31 s | 53× | 2.66 s | 107× |





**Figure 19.** Analytic (dashed) vs. numerical Monte Carlo (solid) kernel comparison, parameter sweeps with varying $P_{\rm eq}$ (upper left), $\kappa$ (upper right), inclination (middle left), $\ell_{\rm spot}$ (middle right), and $\tau_{\rm spot}$ (lower left). The numerical kernel is computed using $N_{\rm sim} = 500$ lightcurve simulations with a baseline of 200d, and cadence of 0.05d. Vertical dotted lines mark the rotation frequency and its first harmonic.

### 9.2. *Comparison to Kernels from Literature*

In this Section, we compare our physically motivated kernel to commonly used kernels in the literature, such as the quasi-periodic (QP) kernel and the simple harmonic oscillator (SHO) kernel. We show that our kernel can reproduce the behavior of these kernels in certain limits, but also captures additional features such as harmonic structure and non-sinusoidal periodicity that are not present in the QP or SHO kernels.

Quasi-periodic kernels typically have the form:

$$k_{\rm QP}(\tau) = A^2 \exp\left(-\frac{\tau^2}{2\lambda^2}\right) \exp\left(-\frac{2\sin^2(\pi\tau/P)}{\omega^2}\right), \tag{91}$$

where $P$ is the period, $\lambda$ is the decay timescale, and $\omega$ controls the smoothness of the periodic component. In the limit of large $\omega$, the periodic component approaches a pure cosine, and the kernel resembles a product of an exponential decay and a cosine.

The simple harmonic oscillator kernel has the form:

$$k_{\rm SHO}(\tau) = A^2 \exp\left(-\frac{\tau}{\lambda}\right) \cos\left(\frac{2\pi\tau}{P}\right), \tag{92}$$

which is a damped cosine with a single period and an exponential decay.

Figure 20 compares the normalized autocorrelation function (left) and power spectral density (right) of our starspot kernel against the QP and SHO kernels for three different parameter regimes. The top panel shows a case with short-lived spots and edge-on geometry. Here, the starspot kernel exhibits a rapidly decaying ACF with prominent harmonic peaks in the PSD, while the QP and SHO kernels show a more sinusoidal ACF and a single dominant peak at the rotation frequency. The second panel shows a case with long-lived spots and inclined geometry, where the starspot kernel has a more complex ACF shape and multiple harmonic peaks in the PSD, which are not captured by the QP or SHO kernels.

The third panel shows a case with strong solar-like differential rotation ($\kappa = 1$). Here, the starspot kernel is able to capture the "skewed" behavior around the ACF peaks (i.e., slower moving spots at higher latitudes have a longer period, thus skewing the ACF around the equitorial period peaks to slightly longer lags). For the QP and SHO kernels, shifts in the ACF peaks can only be made by adjusting the single period parameter, which cannot account for the range of rotation periods due to differential rotation. Differential rotation in the starspot kernel also produces a broadened peak in the PSD around the rotation frequency, reflecting the range of rotation frequencies across different latitudes. In contrast, the QP and SHO kernels cannot capture these features, as they assume a single period and do not account for differential rotation.

Overall, this comparison demonstrates that our physically motivated starspot kernel can capture a richer variety of behaviors in the autocorrelation and power spectrum than commonly used kernels, making it a more flexible, interpretable (and potentially more accurate) choice for modeling stellar variability due to starspots.

### 9.3. *Comparison to Other GP Packages*

While our physically motivated kernel provides richer structure than standard QP or SHO kernels, practical adoption depends on computational performance. We benchmark the `spotgp` GP solver





**Figure 20.** Comparison of the starspot kernel (this work) with the quasi-periodic (QP) and simple harmonic oscillator (SHO) kernels. **Left:** Normalized autocorrelation functions. **Right:** Power spectral densities. **Top row:** Edge-on geometry with short-lived spots ($P_{\rm eq} = 5\,{\rm d}$, $I = 90°$, $\ell = 10\,{\rm d}$, $\tau = 3\,{\rm d}$). **Bottom row:** Inclined geometry with long-lived spots ($P_{\rm eq} = 5\,{\rm d}$, $I = 60°$, $\ell = 25\,{\rm d}$, $\tau = 5\,{\rm d}$). Vertical dotted lines in the PSD panels mark integer harmonics of the rotation frequency. The starspot kernel naturally produces harmonic peaks and non-sinusoidal ACF shapes that the QP and SHO kernels cannot capture.

against two widely used packages: `tinygp` (Foreman-Mackey 2024), which implements the `celerite` $\mathcal{O}(N)$ semi-separable solver (Foreman-Mackey et al. 2017), and `george` (Ambikasaran et al. 2015), which uses a standard $\mathcal{O}(N^3)$ Cholesky factorization.

For a fair comparison, we use a kernel that is representable in all three frameworks: the one-sided exponential envelope $\Gamma(t) = e^{-t/\tau_{\rm spot}}\, H(t)$, which gives $R_\Gamma(\tau) = (2a)^{-1}\, e^{-a|\tau|}$ with $a = 1/\tau_{\rm spot}$. With $N_{\rm harm} = 2$ harmonics, this decomposes into $J = 3$ `celerite` terms (one real + two complex), making it compatible with the $\mathcal{O}(N)$ solver while also being evaluable by the standard dense and banded solvers.

Figure 21 shows the wall-clock time for a single log-likelihood evaluation as a function of the number of datapoints $N$, with fixed cadence $\Delta t = 1\,$d. We compare four solver configurations: (1) `tinygp` with the `celerite` $\mathcal{O}(N)$ solver, (2) `george` with a dense $\mathcal{O}(N^3)$ Cholesky, (3) `spotgp` with a banded $\mathcal{O}(Nb^2)$ Cholesky, and (4) `spotgp` with a full $\mathcal{O}(N^3)$ Cholesky. All JAX-based solvers (`tinygp` and `spotgp`) are JIT-compiled and run on the GPU, while `george` uses compiled C/LAPACK and can only run on the CPU.

The top panel of Figure 21 shows that `tinygp`'s `celerite` solver scales linearly with $N$ and has the lowest overhead at small dataset sizes. The `george` solver scales as $N^3$ but benefits from highly optimized C/LAPACK routines, making it competitive for $N \lesssim 500$. The `spotgp` banded solver carries higher per-evaluation overhead than `tinygp` at small $N$, but because the bandwidth $b$ is set by the kernel correlation length and does not grow with $N$, its $\mathcal{O}(Nb^2)$ scaling is effectively linear for long baselines. At $N = 10{,}000$, the banded solver reaches wall-clock times comparable to `tinygp` ($\sim$0.5 s), while `george` exceeds 30 s. The banded solver overtakes the full $\mathcal{O}(N^3)$ Cholesky around $N \sim 500$.

The middle panel shows matrix storage requirements. The `celerite` solver stores only $\mathcal{O}(NJ)$ elements (where $J = 2N_{\rm harm} + 1$ is the number of semi-separable terms), keeping memory below a few MB even at $N = 10{,}000$. The banded solver stores the $(2b + 1) \times N$ band, which grows linearly with $N$ for fixed $b$. In contrast, the dense solvers (`george` and `spotgp` full) store the full $N \times N$ covariance matrix, reaching $\sim$800 MB at $N = 10{,}000$. For the full `spotgp` solver run on the GPU, the max matrix storage size is limited to the memory capacity of the GPU, which is typically 16–32 GB for consumer hardware (in this case, an Nvidia RTX 5070 TI with 16 GB of memory), thus limiting the maximum $N$ to a few thousand. Similarly for the full `george` solver, the max matrix size is limited by the RAM capacity of the CPU, which is typically 16–64 GB for consumer hardware, thus also limiting the maximum $N$ to a few thousand. In comparison, the banded solver's linear memory scaling allows it to handle much larger datasets without hitting memory limits.

The bottom panel shows the sparsity fraction $(2b+1)/N$ of the banded covariance matrix. Because the bandwidth is determined by the kernel's correlation length (a few rotation periods), it remains fixed as the observation baseline grows. The sparsity fraction therefore drops as $1/N$, falling below 10% for $N \gtrsim 1000$, which is the regime where the banded solver's advantage is most pronounced.

The `celerite` solver's $\mathcal{O}(N)$ scaling is only available for kernels that decompose into a sum of (damped) exponentials and cosines. Our trapezoidal and skew-normal envelopes do not admit this decomposition, so the banded Cholesky solver is the only option for the general kernel. For applications where speed is critical and the one-sided exponential envelope is a sufficient approximation (e.g., for noise modeling, or applications like lightcurve detrending where the user does not care about interpreting the hyperparameters), the `celerite` decomposition provides a practical fast path.





**Figure 21.** Benchmark of GP solver performance as a function of the number of datapoints $N$, using a one-sided exponential envelope with $N_{\rm harm} = 2$ harmonics and cadence $\Delta t = 1\,\mathrm{d}$. *Top:* Wall-clock time for a single log-likelihood evaluation. `tinygp` (célerite, green) scales as $\mathcal{O}(N)$; `george` (red) and `spotgp` full Cholesky (teal) scale as $\mathcal{O}(N^3)$; `spotgp` banded (dark blue) scales as $\mathcal{O}(Nb^2)$. Horizontal gray lines mark 1 s and 1 min. *Middle:* Matrix storage in MB. The `celerite` and banded solvers scale linearly with $N$, while the dense solvers scale as $N^2$. *Bottom:* Sparsity fraction $(2b+1)/N$ of the banded covariance matrix, showing that the bandwidth $b$ remains fixed as $N$ grows. All JAX solvers are JIT-compiled and warmed up before timing.

### 9.4. *Comparison with* `starry_process`

An alternative GP framework for stellar variability is the `starry_process` of Luger et al. (2021b,b), which models the stellar surface as a linear combination of spherical harmonics $Y_{\ell m}$ and derives the covariance of the disk-integrated flux analytically. Our approach differs in several fundamental respects, which we summarize here.

*Surface representation:* `starry_process` represents the stellar surface intensity as a truncated spherical harmonic expansion, $I(\theta, \phi) = \sum_{\ell,m} y_{\ell m} Y_{\ell m}(\theta, \phi)$, with the flux computed via the linear design-matrix relation $\mathbf{f} = \mathbf{A}\,\mathbf{y}$, where $\mathbf{A}$ encodes the rotational geometry and limb darkening (Luger et al. 2019). This yields a non-parametric surface model: the coefficients $y_{\ell m}$ are treated as Gaussian random variables, and the GP covariance of the flux follows from the prior covariance of the harmonic coefficients, $\mathbf{\Sigma}_f = \mathbf{A}\,\mathbf{\Sigma}_y\,\mathbf{A}^\top$. By contrast, our model parameterizes the surface in terms of individual circular spots (each described by a latitude $\Phi$, longitude evolution $\lambda(t)$, and trapezoidal size envelope $\alpha(t)$), and derives the kernel by taking the autocorrelation of the resulting flux analytically.

*Temporal evolution:* A key distinction is the treatment of time dependence. The `starry_process` covariance is *non-stationary*: it depends on absolute time through the rotation matrix $\mathbf{A}(t)$, since the surface pattern is treated as fixed (or drawn once) and the star merely rotates. There is no built-in mechanism for spot emergence, evolution, or decay; the surface is effectively a static random field viewed from different angles. Our kernel, by contrast, is *stationary* (depending only on the time lag $\tau = |t_2 - t_1|$) and explicitly encodes spot lifetimes through the envelope autocorrelation $R_\Gamma(\tau)$, which models the birth, plateau, and decay of each spot. This stationarity arises because we marginalize over the random arrival times of spots (Section 4.7), treating the spot process as a temporal Poisson process. As a consequence, our kernel naturally captures the finite coherence time of the variability signal: the autocorrelation decays to zero at $\tau = \ell_{\rm spot} + 2\tau_{\rm spot}$, a feature that has no analog in `starry_process`.

*Differential rotation:* Our model includes differential rotation through the shear parameter $\kappa$, which allows spots at different latitudes to rotate at different angular velocities (Section 6). This produces observable effects including peak broadening, decoherence, and the coherence lag $\tau_{\rm coh}$ (Figure 15). The `starry_process` framework assumes solid-body rotation.

*Spot parameterization:* In `starry_process`, the surface statistics are controlled by the angular power spectrum of the spherical harmonic coefficients (parameterized by a characteristic angular scale and latitude-dependent modulation). These parameters describe the *spatial* statistics of the surface but do not map directly onto physical spot properties. In our model, the hyperparameters have direct physical interpretations: $P_{\rm eq}$ is the equatorial rotation period, $\ell_{\rm spot}$ is the spot plateau lifetime, $\tau_{\rm spot}$ is the emergence/decay timescale, $\alpha_{\rm max}$ is the peak angular spot radius, and $\kappa$ is the differential rotation shear.

*Computational structure:* The `starry_process` covariance matrix $\mathbf{\Sigma}_f$ is dense and generally $O(N^3)$ to factorize, as it is for any GP. The computational advantage of that framework lies in the analytic evaluation of the matrix entries (avoiding numerical integration over the stellar surface). Our kernel shares this advantage – its evaluation reduces to sums of cosines weighted by tabulated Fourier coefficients – but additionally benefits from sparsity: $K(\tau) = 0$ for $|\tau| > \ell_{\rm spot} + 2\tau_{\rm spot}$, which produces a banded covariance matrix with bandwidth $b = (\ell_{\rm spot} + 2\tau_{\rm spot})/\Delta t$. This enables $O(Nb^2)$ Cholesky factorization via banded linear algebra, a significant speedup for long time series with fine cadence.

*Degeneracies and information content:* Luger et al. (2021b) showed that there exist fundamental degeneracies in mapping stellar surfaces from disk-integrated photometry alone: infinitely many surface maps can produce identical lightcurves. These degeneracies persist in both frameworks. In `starry_process`, they manifest as a null space of the design matrix $\mathbf{A}$; in our model, they appear as the well-known degeneracy between $N_{\rm spot}$, $f_{\rm spot}$, and $\alpha_{\rm max}$ (which are absorbed into the single amplitude parameter $\sigma_k^2$), and the fact that the kernel constrains only the latitude-averaged squared Fourier coefficients $\langle |c_n|^2 \rangle_\Phi$, not the surface map itself. However, our kernel provides direct constraints on *temporal* properties (spot lifetime, evolution timescale, differential rotation) that are not accessible from the static surface model of `starry_process`.

## 10. INFERENCE CONDITIONS

Before applying our GP model to real data, we first test its ability to recover known parameters from synthetic datasets. We generate mock lightcurves by simulating the flux using the same spot model (Equation 1) that underlies our kernel, with known hyperparameters $\boldsymbol{\theta}_{\rm true}$. We then add Gaussian noise (with a standard deviation $\sigma_n$) to the simulated fluxes to create realistic observations.

### 10.0.1. *Kernel Convergence and Data Requirements*

As a first test, we estimate the minimum observation duration ($T_{\rm obs}$) and sampling cadence ($\Delta t$) required for the ACF to converge within a certain error of the true kernel function. To perform this test, we generate a grid of synthetic timeseries with varying $T_{\rm obs}$, $\Delta t$, and $\sigma_n$, and numerically compute the ACF for each individual lightcurve. We then compare the empirical ACF to the true kernel function (computed using the known parameters) and compute the mean squared error (MSE) as a function of $T_{\rm obs}$, $\Delta t$, and $\sigma_n$. This analysis allows us to identify the regimes in which the ACF provides a reliable estimate of the kernel, which is critical for informing the design of observing campaigns and setting expectations for parameter recovery.

Figures 22 shows the results of this test for two sampling cadences: $\Delta t = 0.5\,$d (representative of space-based photometry such as *Kepler* or *TESS*) and $\Delta t = 5.0\,$d (representative of ground-based monitoring campaigns). In each panel, the $x$-axis gives the number of datapoints in the lightcurve $N = T_{\rm obs}/\Delta t$, and the $y$-axis gives the MSE between the sample ACF (computed from a single simulated lightcurve) and the analytic kernel, averaged over 10 random realizations with $1\sigma$ errorbars. Different colored lines correspond to different Gaussian noise levels $\sigma_{\rm noise}$, spanning from $10^{-4}$ (photon-noise-limited space photometry) to $10^{-1}$ (heavily contaminated data). The spot emergence rate is held fixed at $\dot{N}_{\rm spot} = 0.25\,{\rm spots\,day^{-1}}$, so that longer baselines contain proportionally more spots and the kernel amplitude is independent of $T_{\rm obs}$ (Section 4.7).

Several trends are apparent. First, for low noise levels ($\sigma_{\rm noise} \leq 10^{-3}$), the MSE decreases steadily with increasing $N$, roughly as a power law, reflecting the improved statistical convergence of the sample ACF as more independent spot events contribute to the estimate. At the highest noise levels ($\sigma_{\rm noise} = 10^{-1}$), the noise dominates the signal and the MSE saturates at $\sim 10^{-1}$ regardless of the number of datapoints, indicating that the spot-induced correlations are buried in the noise floor.

Second, comparing the two cadences, the coarser sampling ($\Delta t = 5.0\,$d) yields systematically higher MSE at fixed $N$ for the intermediate noise levels ($\sigma_{\rm noise} = 10^{-2}$), because fewer samples per rotation period degrade the ACF estimate at short lags. For the fine cadence ($\Delta t = 0.5\,$d), the low-noise curves ($\sigma_{\rm noise} \leq 10^{-3}$) converge to MSE $\lesssim 10^{-3}$ with $N \gtrsim 10^3$ datapoints (corresponding to $T_{\rm obs} \gtrsim 500\,$d),

while the coarse cadence requires $N \gtrsim 10^2$–$10^3$ points ($T_{\rm obs} \gtrsim 500$–$5000\,{\rm d}$) to reach comparable accuracy.

These results suggest that for typical spot parameters, reliable kernel estimation from a single lightcurve requires observation baselines spanning at least ∼10–20 spot lifetimes ($\ell_{\rm spot} + 2\tau_{\rm spot} = 21\,{\rm d}$ in our test case), with noise levels $\sigma_{\rm noise} \lesssim 10^{-2}$ in relative flux units.





**Figure 22.** MSE between the analytic kernel and the sample ACF computed from individual simulated lightcurves, as a function of the number of datapoints $N$, for a sampling cadence of $\Delta t = 0.5\,{\rm d}$ (left) and $\Delta t = 5.0\,{\rm d}$ (right). Each line corresponds to a different Gaussian noise level $\sigma_{\rm noise}$ added to the simulated flux. Points show the mean over 10 random realizations; errorbars indicate the $1\sigma$ scatter. The spot emergence rate is $\dot{N}_{\rm spot} = 0.25\,{\rm spots\,day^{-1}}$, so $N_{\rm spot}$ scales with the observation duration.

### 10.1. *Testing the Cranmer–Rao Bound of Different Parameter Regimes*

Even though we are only modeling a six parameter kernel model, there are many possible combinations of parameters that can produce similar kernel shapes, leading to degeneracies and non-identifiability. For example, if the spot lifetime is very short compared to the rotation period, the kernel will be dominated by the envelope autocorrelation and may not contain enough information about the periodic structure to constrain $P_{\rm eq}$ or $\kappa$. Conversely, if the spot lifetime is relatively long compared to the rotation period or observational baseline, the kernel may be dominated by the periodic structure and have a very slowly decaying ACF, which could also make it difficult to constrain the envelope parameters $\ell_{\rm spot}$ and $\tau_{\rm spot}$.

Before performing MCMC sampling (which is computationally intensive to do across a wide parameter space), we can first test the theoretical identifiability of the parameters by computing the Fisher information matrix (Eq. 90) across a grid of parameters. The inverse of the Fisher information matrix gives the Cramér–Rao lower bound on the covariance of any unbiased estimator of the parameters, which provides a theoretical limit on how well we can constrain the parameters from data generated by our model. By analyzing the eigenvalues and eigenvectors of the Fisher information matrix, we can identify which combinations of parameters are most and least identifiable, and how this depends on the underlying spot parameters. This analysis will inform our expectations for MCMC sampling and help us interpret the results in terms of parameter degeneracies and uncertainties.

To systematically explore parameter space, we draw $N = 4096$ parameter combinations from a scrambled Sobol quasi-random sequence spanning the bounds listed in Table 11. Sampling $\ell_{\rm spot}$ and $\tau_{\rm spot}$ independently would leave their sum sparse at both ends of its range, so we instead draw the total spot lifetime $\ell_{\rm spot}+2\tau_{\rm spot}$ uniformly over $[3, 300]$ d, draw $\ell_{\rm spot}$ uniformly from the values that keep both timescales within their bounds, and set $\tau_{\rm spot}$ from the remainder. For each sample $\boldsymbol{\theta}_k$, we construct a synthetic lightcurve of duration $T_{\rm sim} = 365$ d sampled at a cadence of $\Delta t = 0.5$ d (i.e., $N_t = 730$ observations) with white noise $\sigma_n = 10^{-4}$ and a mean spot emergence rate of $\dot{N}_{\rm spot} = 0.25$ d$^{-1}$. We then evaluate the expected Fisher information matrix $\mathcal{I}(\boldsymbol{\theta}_k)$ via Eq. (90) and invert it to obtain the CRB covariance matrix $\Sigma_{\rm CRB} = \mathcal{I}^{-1}$. Cases where any eigenvalue of $\mathcal{I}$ falls below a regularization floor of $10^{-6}$ are flagged as numerically singular and excluded from the analysis, as the corresponding CRB is unreliable. From the CRB covariance we extract the marginal standard deviations $\sigma_{\rm CRB}(\theta_i) = \sqrt{\Sigma_{ii}}$ and the pairwise Pearson correlation coefficients $r_{ij} = \Sigma_{ij}/(\sigma_i\sigma_j)$ for all parameter pairs.

### 10.2. *CRB results*

To visualize the CRB parameter sweep, we compute two dimensional projections between each combination of parameters. Each panel of the corner plots projects the six-dimensional sweep onto a pair of parameters, with the two envelope timescales combined into the total spot lifetime $\ell_{\rm spot}+2\tau_{\rm spot}$. The samples are divided into a $20 \times 20$ grid of equal-width bins, about 10 samples per bin, and the color shows the mean of $\log_{10}\sigma_{\rm CRB}$ in each bin. The other four parameters vary freely within a bin, so this mean marginalizes over them, weighted uniformly across their sweep ranges. A panel therefore shows the typical bound at each pair of values, not the bound with the remaining parameters held fixed. The binned means are smoothed with a Gaussian kernel of width 1.5 bins, normalized by the smoothed bin occupancy so that only occupied bins contribute, and any empty bin is masked and drawn in gray.

**Solar versus anti-solar shear:** At long periods the bound on $\kappa$ depends on its sign (the $\kappa$–$P_{\rm eq}$ panel of Fig. 23). For $|\kappa| > 0.4$, the median $\sigma_{\rm CRB}(\kappa)$ for solar-like shear exceeds that for anti-solar shear of the same magnitude. The spread of rotation frequencies does not explain it. The frequency $\omega_0(\Phi) = 2\pi(1 - \kappa\sin^2\Phi)/P_{\rm eq}$ is linear in $\kappa$, so both the sensitivity $\partial\omega_0/\partial\kappa = -2\pi\sin^2\Phi/P_{\rm eq}$ and the dephasing rate (Eq. 80) depend only on $|\kappa|$. What does depend on the sign is where in frequency the high-latitude spots sit, and these spots carry all of the leverage on $\kappa$. For $\kappa > 0$ they rotate more slowly than the equator, down to $\omega_0 = (1-\kappa)\,2\pi/P_{\rm eq}$ at the pole. For $\kappa < 0$ they rotate faster, up to $(1+|\kappa|)\,2\pi/P_{\rm eq}$.

A spot that completes fewer than one or two rotations during its lifetime contributes a term $\cos(\omega_0\tau)R_\Gamma(\tau) \approx R_\Gamma(\tau)(1 - \omega_0^2\tau^2/2)$. This term is nearly indistinguishable from the aperiodic $n = 0$ term, and its derivative with respect to $\omega_0$ vanishes as $\omega_0 \to 0$. In the power spectrum, the fundamental peak from those latitudes merges with the zero-frequency peak from spot evolution, whose width is $\sim 1/\ell_{\rm spot}$.

The controlling quantity is the number of polar rotations per spot lifetime, $(\ell_{\rm spot}+2\tau_{\rm spot})(1-\kappa)/P_{\rm eq}$. At $P_{\rm eq} = 40$ d the polar period is 100 d for $\kappa = 0.6$ and 25 d for $\kappa = -0.6$. At $P_{\rm eq} \lesssim 10$ d the polar spots of most samples still complete many rotations per lifetime. There the asymmetry is confined to $\kappa \gtrsim 0.6$, where the polar period $P_{\rm eq}/(1-\kappa)$ diverges.

**Table 11.** Parameter bounds used for the Sobol quasi-random sweep of the CRB analysis.

| Parameter | Lower | Upper |
|---|---|---|
| $P_{\rm eq}$ [d] | 0.1 | 50 |
| $\kappa$ | $-1.0$ | 1.0 |
| $I$ [deg] | 0 | 90 |
| $\ell_{\rm spot}$ [d] | 1.0 | 100 |
| $\tau_{\rm spot}$ [d] | 1.0 | 100 |
| $\sigma_k$ | $10^{-3}$ | 1 |





**Figure 23.** Cramér–Rao lower bound on the differential rotation shear, $\sigma_{\rm CRB}(\kappa)$, across the 4096 Sobol samples drawn within the bounds of Table 11. The color scale gives the bin-averaged $\log_{10}\sigma_{\rm CRB}(\kappa)$, with dark blue indicating tighter constraints and dark red indicating looser constraints. For short periods ($P_{\rm eq} \lesssim 20\,$d), long spot lifetimes ($\ell_{\rm spot} + 2\tau_{\rm spot} \gtrsim 10 P_{\rm eq}$), and high inclinations ($I \gtrsim 45°$), the bound is typically $\sigma_{\rm CRB}(\kappa) \approx 0.01$, falling to $\approx 0.002$ for $P_{\rm eq} \lesssim 5\,$d; for $P_{\rm eq} \gtrsim 40\,$d it rises to $\gtrsim 0.3$.





**Figure 24.** Same as Fig. 23, for the bound on the inclination, $\sigma_{\rm CRB}(I)$ in degrees. The bound is $\approx 1°$ for $P_{\rm eq} \lesssim 10\,$d and loosens toward long periods and short spot lifetimes. It grows toward edge-on viewing, from $\approx 2°$ at $I \lesssim 25°$ to $\approx 4°$ at $I \approx 45°$–$65°$ and $\gtrsim 10°$ above $80°$: for a latitude distribution symmetric about the equator the kernel is unchanged under $I \to 180° - I$, so its derivative with respect to $I$ vanishes at $I = 90°$. For $I \gtrsim 80°$ the bound exceeds the distance to edge-on, $90° - I$, in most samples, so there the local Gaussian bound is not meaningful and the posterior is instead limited by the $I = 90°$ boundary.





**Figure 25.** Same as Fig. 23, for the bound on the equatorial period, $\sigma_{\rm CRB}(P_{\rm eq})$ in days. $P_{\rm eq}$ is by far the easiest of the parameters to constrain. The bound grows steeply with the period itself, from $\approx 0.002\,$d at $P_{\rm eq} \lesssim 5\,$d to $\approx 0.7\,$d at $P_{\rm eq} \approx 20$–$30\,$d and $\approx 4\,$d at $P_{\rm eq} \gtrsim 40\,$d, and is nearly independent of $\kappa$ and $\sigma_k$. It loosens most at pole-on inclinations ($I \lesssim 10°$), where the rotational harmonics carry little power, and for spots that do not survive a full rotation ($\ell_{\rm spot} + 2\tau_{\rm spot} \lesssim P_{\rm eq}$), where it exceeds the 50 d width of the sweep and the period is effectively unconstrained.





**Figure 26.** Same as Fig. 23, for the bound on the kernel amplitude, $\sigma_{\rm CRB}(\sigma_k)$. As expected for an amplitude, the bound scales nearly in proportion to $\sigma_k$ itself, from $\approx 0.002$ for $\sigma_k \lesssim 0.1$ to $\approx 0.035$ for $\sigma_k \gtrsim 0.8$, so the bottom row is dominated by the amplitude. At fixed $\sigma_k$ the bound depends only weakly on the other parameters, loosening for short spot lifetimes, especially at long periods.

### 10.3. *MCMC Inference*

While Section 10 gives a sense of the minimum observation requirements for reliable ACF estimation, it doesn't directly test the ability of our GP model to recover the underlying parameters $\boldsymbol{\theta}$ and their uncertainties from noisy data. To assess this, we perform MCMC sampling of the posterior distribution $p(\boldsymbol{\theta}|\mathbf{y})$ using the sampling methods described in Section 8. More specifically, we investigate how the noise level and observation duration affect the ability to recover the true parameters, and how this compares to the theoretical limits set by the Cramér–Rao bound. We also analyze the posterior distributions to identify any degeneracies or correlations between parameters, and to understand how well the model can constrain each parameter under different conditions.

As in the CRB analysis (Section 10.1), we simulate lightcurve data with a white noise level of $\sigma_n = 10^{-4}$ ($\sim$ 100 ppm), but using a sampling cadence of $\Delta t = 1.0$ d (i.e., $N_t = 365$ and 1460 observations for $T_{\rm sim} = 365$ and 1460 d). Given that *Kepler* has a finer cadence (1 minute for the short cadence or 30 min for the long cadence data), and typical instrumental noise levels of less than 50 ppm Gilliland et al. (2010), we assume that these simulated values are reasonable for *Kepler*-like data.

For each synthetic lightcurve, we ran both NUTS (with `BlackJAX`) and nested sampling (with `dynesty`) to obtain posterior samples of the parameters $\boldsymbol{\theta}$ and compare the results to the true parameters $\boldsymbol{\theta}_{\rm true}$ used to generate the data. In practice, we found that the NUTS sampler tended to get stuck in small local modes of the posterior and was less effective at exploring the full parameter space, and so we primarily report results from the nested sampling runs, which were more robust to multimodality and provided better coverage of the posterior.

To see how the parameter uncertainties carry through to the covariance structure, Figure 28 shows the kernel and PSD evaluated at random draws from the posteriors in Figure 27. At $I = 90°$, both baselines recover the true kernel and PSD. At $I = 30°$, the 1-year posterior underestimates the correlation at short lags, which matches its inclination posterior lying above the truth ($I = 36 \pm 3°$): a higher inclination deepens the rotational modulation. With 4 years of data, the posterior kernel follows the true kernel.

Figure 29 shows the GP prediction at the posterior mean of each fit: the predictive mean and standard deviation of the noise-free flux, conditioned on all of the data. The predictive standard deviation falls to about the white noise level at each observation and rises between observations, but stays two orders of magnitude below the amplitude of the rotational modulation, so the prediction follows the lightcurve closely between the daily samples. Over the whole baseline of each fit, the noise-free flux lies within the $1\sigma$ and $2\sigma$ predictive intervals at close to the nominal 68% and 95% rates, so the predictive variance is well calibrated at the posterior mean.

**Table 12.** Posterior summary statistics from nested sampling (`dynesty`) on simulated lightcurves. For each combination of observation duration ($T_{\rm sim}$) and true inclination ($I_{\rm true}$), we report the posterior mean and standard deviation alongside the true (injected) parameter value and the bounds of its uniform prior. The inclination is converted from radians to degrees for readability. All simulations use a sampling cadence of $\Delta t = 1\,\mathrm{d}$.

| $I_{\rm true}$ | Parameter | True | Prior | $T_{\rm sim} = 365\,\mathrm{d}$ | | $T_{\rm sim} = 1460\,\mathrm{d}$ | |
|---|---|---|---|---|---|---|---|
| | | Value | (min, max) | Mean | Std | Mean | Std |
| **30°** | $P_{\rm eq}$ [d] | 5 | (4.5, 5.5) | 4.969 | 0.019 | 4.980 | 0.013 |
| | $\kappa$ | 0.3 | (−1, 1) | 0.309 | 0.035 | 0.305 | 0.019 |
| | $I$ [deg] | 30 | (0, 60) | 33.5 | 2.0 | 32.6 | 1.1 |
| | $\ell_{\rm spot}$ [d] | 20 | (10, 30) | 20.10 | 0.44 | 20.03 | 0.13 |
| | $\tau_{\rm spot}$ [d] | 5 | (0, 10) | 5.24 | 0.63 | 4.63 | 0.27 |
| | $\sigma_k$ | 0.0032 | (0.001, 0.01) | 0.00319 | 0.00022 | 0.003046 | 0.000095 |
| **60°** | $P_{\rm eq}$ [d] | 5 | (4.5, 5.5) | 4.945 | 0.037 | 5.005 | 0.018 |
| | $\kappa$ | 0.3 | (−1, 1) | 0.349 | 0.032 | 0.287 | 0.015 |
| | $I$ [deg] | 60 | (30, 90) | 57.2 | 3.4 | 60.7 | 1.5 |
| | $\ell_{\rm spot}$ [d] | 20 | (10, 30) | 22.46 | 0.49 | 22.39 | 0.34 |
| | $\tau_{\rm spot}$ [d] | 5 | (0, 10) | 1.02 | 0.42 | 1.40 | 0.22 |
| | $\sigma_k$ | 0.0032 | (0.001, 0.01) | 0.00253 | 0.00013 | 0.002984 | 0.000074 |
| **90°** | $P_{\rm eq}$ [d] | 5 | (4.5, 5.5) | 4.982 | 0.044 | 5.019 | 0.014 |
| | $\kappa$ | 0.3 | (−1, 1) | 0.330 | 0.033 | 0.287 | 0.012 |
| | $I$ [deg] | 90 | (60, 120) | 90.0 | 5.2 | 90.0 | 3.4 |
| | $\ell_{\rm spot}$ [d] | 20 | (10, 30) | 19.88 | 0.83 | 19.50 | 0.34 |
| | $\tau_{\rm spot}$ [d] | 5 | (0, 10) | 5.2 | 1.2 | 5.69 | 0.44 |
| | $\sigma_k$ | 0.0032 | (0.001, 0.01) | 0.00310 | 0.00017 | 0.003341 | 0.000089 |





**Figure 27.** Posterior distributions for different inclinations: $I = 90$ (bottom left) and 30 (upper right) degrees based on nested sampling runs on synthetic lightcurves. The blue posterior outlines the constraints using 1 year of data, while the green posterior outlines the constraints using 4 years of data (comparable to the Kepler mission duration), and the true parameters are indicated by the dashed black lines. The posterior distributions show that the model can recover the true parameters with tight precision given a typical noise level of $\sigma_n = 10^{-4}$ and a spot emergence rate of $\dot{N}_{\rm spot} = 0.25\,{\rm d}^{-1}$.





**Figure 28.** Kernel and PSD evaluated at 2000 random draws from the nested sampling posteriors in Figure 27, for $I = 90°$ (top) and $I = 30°$ (bottom). The left panels show the normalized kernel $K(\tau)/K(0)$, with each draw normalized by its own $K(0)$. The right panels show the PSD up to the Nyquist frequency of the $\Delta t = 1\,\mathrm{d}$ sampling. Solid lines show the posterior median, and the dark and light shaded regions show the pointwise 68% and 95% intervals for 1 year (blue) and 4 years (green) of data. The dashed black lines show the kernel and PSD at the true parameters. The lower panels show the residuals from the truth: $\Delta K/K(0)$ for the kernel and $P/P_{\rm true} - 1$ for the PSD. The fractional PSD residual spikes near the nulls of the PSD, where $P_{\rm true} \to 0$.





**Figure 29.** GP prediction at the posterior mean of the nested sampling fits in Figure 27, for $I = 90°$ (top) and $I = 30°$ (bottom) with 1 year (left, blue) and 4 years (right, green) of data, over days 200–260 of each simulated lightcurve. The 1-year and 4-year fits are to separate simulations. In each panel, the upper row shows the predictive mean of the noise-free flux (solid line) with its $\pm1\sigma$ and $\pm2\sigma$ intervals (shaded, narrower than the line at this scale), and the observations with their $1\sigma$ errors (circles). The lower row shows the residual of the predictive mean from the true, noise-free simulated flux in units of $\sigma$.

# 11. DISCUSSION

## 11.1. *Summary of Results*

In this work, we derived a physically motivated GP kernel for stellar photometric variability caused by rotating starspots and systematically characterized the performance and limitations of this methodology using simulations; the application to real data is the subject of Paper II. We summarize the main results below.

**Analytic kernel derivation (Section 4).** Starting from a forward model of the flux deficit produced by circular spots on a differentially rotating star, we decomposed the single-spot signal into a product of an envelope term $\Gamma(t)$ (capturing spot emergence and decay) and a visibility term $\mathcal{V}(t)$ (capturing rotational modulation and limb geometry). By taking the Fourier transform of each component and applying the Wiener–Khinchin theorem, we obtained a closed-form expression for the autocovariance kernel $\mathcal{K}(\tau)$ as a sum over rotation harmonics weighted by latitude-averaged visibility coefficients $\langle|c_n|^2\rangle_\Phi$ and modulated by the envelope autocorrelation $R_\Gamma(\tau)$ (Eq. 59). We provided analytic $R_\Gamma(\tau)$ expressions for trapezoidal, exponential, and skew-normal spot profiles, and validated all derivations symbolically with `SymPy` and numerically against brute-force simulations.

**Physical interpretability of hyperparameters (Section 6).** The six free kernel hyperparameters, $P_{\rm eq}$, $\kappa$, $I$, $\ell_{\rm spot}$, $\tau_{\rm spot}$, and $\sigma_k$, map directly onto stellar and spot properties, unlike the phenomenological parameters of standard quasi-periodic or SHO kernels. We showed that $P_{\rm eq}$ and $\kappa$ control the periodicity and harmonic structure of the kernel; $I$ controls the balance between the DC and oscillating components via the visibility coefficients; $\ell_{\rm spot}$ and $\tau_{\rm spot}$ set the correlation lifetime; and the amplitude parameters ($\alpha_{\rm max}$, $f_{\rm spot}$, $N_{\rm spot}$) are degenerate and collapse into a single scale parameter $\sigma_k^2$. We further demonstrated that our kernel captures signatures, such as PSD broadening from differential rotation and harmonic ratio variations with inclination, that standard kernels cannot reproduce (Section 9.2).

**Efficient GP solver (Section 7).** Because the kernel $\mathcal{K}(\tau)$ has compact support (set by the spot lifetime $\ell_{\rm spot} + 2\tau_{\rm spot}$), the covariance matrix is banded with bandwidth $b \propto (\ell_{\rm spot} + 2\tau_{\rm spot})/\Delta t$. Exploiting this structure, we implemented a banded Cholesky solver that reduces the GP inference cost from $\mathcal{O}(N^3)$ to $\mathcal{O}(Nb^2)$. The solver is fully JIT-compiled and autodifferentiable via JAX, enabling gradient-based sampling with no additional implementation effort.

**Inference conditions and Cramér–Rao bounds (Sections 10–10.1).** Using synthetic lightcurves, we showed that reliable kernel estimation requires observation baselines spanning at least ∼10–20 spot lifetimes with noise levels $\sigma_{\rm noise} \lesssim 10^{-2}$ in relative flux. We computed the Cramér–Rao lower bound across a grid of 4096 parameter combinations and found that the differential rotation coefficient $\kappa$ is best constrained for short-period stars ($P_{\rm eq} \lesssim 20\,{\rm d}$) with long-lived spots and moderate-to-high inclinations, where the bound is $\sigma_{\rm CRB}(\kappa) \approx 0.01$ (and $\approx 0.002$ for $P_{\rm eq} \lesssim 5\,{\rm d}$). The bound on the inclination is a few degrees over most of the sweep but diverges toward edge-on viewing, where the kernel is insensitive to $I$.

**Application to real data (Paper II).** The companion paper extends the kernel into an additive composite model, with several spot populations sharing one stellar geometry and with granulation and white-noise terms, and applies the framework to the four-year *Kepler* lightcurve of Kepler-63, comparing the inferred inclination, differential rotation, and spot lifetimes with independent constraints from spectroscopy and transit spot mapping.

## 11.2. *Applications and Future Work*

### 11.2.1. *Direct Applications for this model*

There are numerous potential applications for this starspot kernel. Firstly, it can be used as a tool for inferring star spot parameters using only photometric lightcurves, without the need for spectroscopic or transit data. This is particularly useful for stars that do not have transiting planets or for which high-resolution spectroscopy is not available. By fitting the kernel to the lightcurve, one can extract information about the star's rotation period, differential rotation, inclination, and spot lifetimes, which are important for understanding stellar magnetic activity and its impact on exoplanet habitability.

Additionally, the kernel can be used to model and mitigate stellar variability in the context of exoplanet transit searches and radial velocity measurements, improving the precision of planet detection and characterization. The physically motivated nature of the kernel also allows for more interpretable modeling of stellar variability compared to phenomenological kernels, enabling insights into the underlying stellar physics driving the observed lightcurve features.

### 11.2.2. *Extensions to Spectroscopy*

This model can also be used in combination with spectroscopic data to jointly constrain stellar and spot properties. For example, our GP kernel can be used to infer the inclination angle of a star just from the photometric variability. This means that we can combine this information with spectroscopic measurements of $v_* \sin I$, allowing us to infer the equatorial rotation velocity $v_{\rm eq}$ and thus the stellar radius $R_* = v_{\rm eq} P_{\rm eq}/2\pi$. This methodology would allow us to constrain stellar radii independently of stellar evolution/isochrone models.

Another natural extension of this framework is to high-resolution spectroscopic time series, where starspots induce wavelength-dependent distortions in absorption line profiles as they rotate across the stellar disk. In the Doppler imaging framework (e.g., Luger et al. 2021a), a spot at latitude $\Phi$ and longitude $\Lambda(t)$ perturbs the line profile at a radial velocity $v(t) = v_{\rm eq} \sin I \cos\Phi \sin\Lambda(t)$, which is directly related to the same visibility geometry (Eq. 6) that underpins our photometric kernel. The Fourier decomposition and Wiener–Khinchin machinery developed in Section 4 should therefore generalize to a two-dimensional spectral-temporal kernel $\mathcal{K}(\tau, \Delta v)$, describing the covariance of spectral perturbations across both time lag and velocity separation. Such an extension could help break degeneracies that are present in photometry alone: in particular, the inclination $I$ and spot latitude $\Phi$ enter differently through the line-of-sight velocity than through the projected area, and differential rotation $\kappa$ produces directly observable velocity-track separations for spots at different latitudes. Joint photometric–spectroscopic inference with a physically motivated kernel could therefore improve constraints on stellar surface properties, particularly in the context of extreme precision radial velocity (EPRV) surveys where disentangling stellar activity from planetary signals remains a central challenge.

### 11.2.3. *Further Development*

We make the kernel implementation publicly available as a Python package, `spotgp`, which provides a user-friendly interface for evaluating the kernel and performing GP regression with it. The package includes functions for computing the kernel given a set of hyperparameters, as well as utilities for

sampling the hyperparameters from data using MCMC. We also provide example Jupyter notebooks demonstrating how to use the kernel for modeling stellar lightcurves and inferring starspot properties.

Furthermore, we have implemented the package in a modular way such that users can easily adapt the kernel to their specific needs and assumptions (such as different spot latitude distributions, envelope functions, or spot parameter distributions). We hope that `spotgp` will be a valuable resource for the community and facilitate further research into stellar variability and starspot modeling.

### 11.3. *Limitations*

Our kernel model is based on several simplifying assumptions, such as circular spots with uniform contrast, a fixed spot latitude distribution, and independent spot evolution. In reality, starspots can have more complex shapes, non-uniform contrasts, and form in groups or active regions, which could introduce additional correlations in the lightcurve that are not captured by our model. Additionally, the single-population kernel derived here does not account for other forms of astrophysical variability, such as asteroseismic pulsations, flares, or granulation, which may be present in the lightcurve; additive kernel terms for granulation and white noise, and for several coexisting spot populations, are introduced in Paper II.

In the analysis of this paper, we have focused on dark starspots, however bright faculae can also contribute to photometric variability, particularly in Sun-like stars. In the next paper of this series, we will derive a kernel for multiband photometric data and show how dark spots and bright faculae can be disentangled using their different wavelength-dependent contrasts.

Our kernel is also derived under the assumption that all spots are drawn from the same distributions for $\Phi$, $\ell_{\rm spot}$, $\tau_{\rm spot}$, and $\alpha_{\rm max}$. In reality, there may be correlations between these parameters (e.g., larger spots may have longer lifetimes), which could affect the shape of the kernel. We also assume that these starspot properties are constant in time, such that the kernel is stationary. In practice, spot properties may evolve over time (e.g., due to activity cycles), leading to non-stationary behavior in the lightcurve. Future work could explore more complex models that allow for such correlations. However, the current model provides a useful starting point for understanding the key physical parameters that shape the photometric variability of spotted stars and can be extended in future work to relax some of these assumptions. We also provide several examples on the online documentation for how users can test non-stationary kernels and more complex spot models using the modular implementation of `spotgp`.

## 12. CONCLUSION

In conclusion, we have derived an analytic, physically motivated Gaussian process kernel for stellar photometric variability driven by rotating starspots. Starting from a forward model of circular spots on a differentially rotating star, we obtained a closed-form autocovariance function whose six hyperparameters, $P_{\rm eq}$, $\kappa$, $I$, $\ell_{\rm spot}$, $\tau_{\rm spot}$, and $\sigma_k$, map directly onto measurable stellar and spot properties. This stands in contrast to the phenomenological kernels widely used in the literature, whose parameters lack clear physical interpretation.

Using simulated lightcurves and MCMC/nested sampling, we demonstrated that the kernel can recover injected stellar parameters, including the stellar inclination and differential rotation coefficient $\kappa$, given sufficient observational baseline ($\gtrsim$10 spot lifetimes) and photometric precision ($\sigma_{\rm noise} \lesssim 10^{-2}$). Using Cramér–Rao analysis we further showed that $\kappa$ is best constrained for short-period,

high-inclination stars with long-lived spots. The application of the method to the *Kepler* lightcurve of Kepler-63 is presented in Paper II.

The compact support of the kernel enables a banded Cholesky GP solver that scales as $\mathcal{O}(Nb^2)$ rather than $\mathcal{O}(N^3)$, making inference tractable for the long-baseline, densely sampled lightcurves produced by *Kepler* and *TESS*. Combined with a JAX-based implementation that supports automatic differentiation and gradient-based sampling, this framework is well suited for population-level studies of stellar surface properties.

Looking ahead, systematic application of this kernel to large photometric surveys could yield differential rotation and spot lifetime measurements for thousands of stars across a range of spectral types and ages: quantities that are currently available for only a handful of targets. Such measurements would place new empirical constraints on stellar dynamo theory, the evolution of surface magnetic activity, and the impact of stellar variability on exoplanet detection and habitability assessments.

*Software:* `JAX` (Bradbury et al. 2018), `NumPy` (Harris et al. 2020), `SciPy` (Virtanen et al. 2020), `Matplotlib` (Hunter 2007), `Astropy` (Astropy Collaboration et al. 2013, 2018), `BlackJAX` (Cabezas et al. 2024), `SymPy` (Meurer et al. 2017)

## ACKNOWLEDGMENTS

EA was supported by National Aeronautics and Space Administration Exoplanet Research Program (NASA XRP) grant 80NSSC21K1111. This work was performed in part by the Virtual Planetary Laboratory Team, which is a member of the NASA Nexus for Exoplanet System Science, and funded via the NASA Astrobiology Program ICAR Grant 80NSSC23K1398.

## REFERENCES


Aigrain, S., Llama, J., Ceillier, T., et al. 2015, Mon Not R Astron Soc, 450, 3211, doi: 10.1093/mnras/stv853

Amard, L., & Matt, S. P. 2020, The Astrophysical Journal, 889, 108, doi: 10.3847/1538-4357/ab6173

Ambikasaran, S., Foreman-Mackey, D., Greengard, L., Hogg, D. W., & O'Neil, M. 2015, arXiv:1403.6015 [astro-ph, stat]. http://arxiv.org/abs/1403.6015

Angus, R., Morton, T., Aigrain, S., Foreman-Mackey, D., & Rajpaul, V. 2018, Mon Not R Astron Soc, 474, 2094, doi: 10.1093/mnras/stx2109

Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, \textbackslashaap, 558, A33, doi: 10.1051/0004-6361/201322068

Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, \textbackslashaj, 156, 123, doi: 10.3847/1538-3881/aabc4f

Augustson, K. C., Brown, B. P., Brun, A. S., Miesch, M. S., & Toomre, J. 2012, ApJ, 756, 169, doi: 10.1088/0004-637X/756/2/169

Barnes, S. A. 2003, The Astrophysical Journal, 586, 464, doi: 10.1086/367639

—. 2007, \textbackslashapj, 669, 1167, doi: 10.1086/519295

Basri, G., & Shah, R. 2020, The Information Content in Analytic Spot Models of Broadband Precision Light Curves. II. Spot Distributions and Lifetimes, Global and Differential Rotation, arXiv, doi: 10.48550/arXiv.2008.04969

Benomar, O., Bazot, M., Nielsen, M. B., et al. 2018, Science, 361, 1231, doi: 10.1126/science.aao6571

Birky, J., Agol, E., Barnes, R., Davenport, J. R. A., & Matt, S. 2026, Journal of Open Source Software, in preparation

Borucki, W. J., Koch, D., Basri, G., et al. 2010, Science, 327, 977, doi: 10.1126/science.1185402

Boyle, A. W., Bouma, L. G., & Mann, A. W. 2026, The TESS All-Sky Rotation Survey: Periods for 944,056 Stars Within 500 Pc, arXiv, doi: 10.48550/arXiv.2603.05586

Bradbury, J., Frostig, R., Hawkins, P., et al. 2018, JAX: composable transformations of Python+NumPy programs. http://github.com/jax-ml/jax

Brun, A. S., Strugarek, A., Varela, J., et al. 2017, ApJ, 836, 192, doi: 10.3847/1538-4357/aa5c40

Cabezas, A., Corenflos, A., Lao, J., & Louf, R. 2024, BlackJAX: A sampling library for JAX. http://github.com/blackjax-devs/blackjax

Charbonneau, P. 2020, Living Reviews in Solar Physics, 17, 4, doi: 10.1007/s41116-020-00025-6

Collier Cameron, A. 2007, Astronomische Nachrichten, 328, 1030, doi: 10.1002/asna.200710880

Foreman-Mackey, D. 2024, tinygp: The tiniest of Gaussian process libraries. https://github.com/dfm/tinygp

Foreman-Mackey, D., Agol, E., Ambikasaran, S., & Angus, R. 2017, The Astronomical Journal, 154, 220, doi: 10.3847/1538-3881/aa9332

Forgács-Dajka, E., Dobos, L., & Ballai, I. 2021, Astronomy & Astrophysics, 653, A50, doi: 10.1051/0004-6361/202140731

Gastine, T., Yadav, R. K., Morin, J., Reiners, A., & Wicht, J. 2013, Monthly Notices of the Royal Astronomical Society: Letters, 438, L76, doi: 10.1093/mnrasl/slt162

Giles, H. A. C., Cameron, A. C., & Haywood, R. D. 2017, Monthly Notices of the Royal Astronomical Society, 472, 1618, doi: 10.1093/mnras/stx1931

Gilliland, R. L., Jenkins, J. M., Borucki, W. J., et al. 2010, The Astrophysical Journal, 713, L160, doi: 10.1088/2041-8205/713/2/L160

Golub, G. H., & Van Loan, C. F. 2013, Matrix Computations, 4th edn. (Johns Hopkins University Press)

Gordon, T. A., Davenport, J. R. A., Angus, R., et al. 2021, The Astrophysical Journal, 913, 70, doi: 10.3847/1538-4357/abf63e

Guerrero, G., Smolarkiewicz, P. K., Kosovichev, A. G., & Mansour, N. N. 2013, The Astrophysical Journal, 779, 176, doi: 10.1088/0004-637X/779/2/176

Harris, C. R., Millman, K. J., van der Walt, S. J., et al. 2020, Nature, 585, 357, doi: 10.1038/s41586-020-2649-2

Hoffman, M. D., & Gelman, A. 2014, Journal of Machine Learning Research, 15, 1593. http://jmlr.org/papers/v15/hoffman14a.html

Howell, S. B., Sobeck, C., Haas, M., et al. 2014, Publications of the Astronomical Society of the Pacific, 126, 398, doi: 10.1086/676406
Hunter, J. D. 2007, Computing in Science and Engineering, 9, 90, doi: 10.1109/MCSE.2007.55
Ireland, L. G., Matt, S. P., Davey, C. R., et al. 2021, arXiv:2111.11082 [astro-ph]
Kawaler, S. D. 1988, The Astrophysical Journal, 333, 236, doi: 10.1086/166740
Kipping, D. M. 2012, Mon Not R Astron Soc, 427, 2487, doi: 10.1111/j.1365-2966.2012.22124.x
Käpylä, P. J., Browning, M. K., Brun, A. S., Guerrero, G., & Warnecke, J. 2023, Space Science Reviews, 219, 58, doi: 10.1007/s11214-023-01005-6
Luger, R., Agol, E., Foreman-Mackey, D., et al. 2019, AJ, 157, 64, doi: 10.3847/1538-3881/aae8e5
Luger, R., Bedell, M., Foreman-Mackey, D., et al. 2021a, Mapping stellar surfaces III: An Efficient, Scalable, and Open-Source Doppler Imaging Model, arXiv, doi: 10.48550/arXiv.2110.06271
Luger, R., Foreman-Mackey, D., Hedges, C., & Hogg, D. W. 2021b, The Astronomical Journal, 162, 123, doi: 10.3847/1538-3881/abfdb8
Matt, S. P., Brun, A. S., Baraffe, I., Bouvier, J., & Chabrier, G. 2015, The Astrophysical Journal Letters, 799, L23, doi: 10.1088/2041-8205/799/2/L23
McQuillan, A., Mazeh, T., & Aigrain, S. 2014, ApJS, 211, 24, doi: 10.1088/0067-0049/211/2/24
Meurer, A., Smith, C. P., Paprocki, M., et al. 2017, PeerJ Computer Science, 3, e103, doi: 10.7717/peerj-cs.103
Miesch, M. S., Brun, A. S., DeRosa, M. L., & Toomre, J. 2008, The Astrophysical Journal, 673, 557, doi: 10.1086/523838
Miesch, M. S., Elliott, J. R., Toomre, J., et al. 2000, The Astrophysical Journal, 532, 593, doi: 10.1086/308555
Morris, B. M., Davenport, J. R. A., Giles, H. A. C., et al. 2019, Monthly Notices of the Royal Astronomical Society, 484, 3244, doi: 10.1093/mnras/stz199
Morris, B. M., Hebb, L., Davenport, J. R. A., Rohn, G., & Hawley, S. L. 2017, ApJ, 846, 99, doi: 10.3847/1538-4357/aa8555
Nicholson, B. A., & Aigrain, S. 2022, Mon Not R Astron Soc, 515, 5251, doi: 10.1093/mnras/stac2097
Noraz, Q., Breton, S. N., Brun, A. S., et al. 2022a, Astronomy & Astrophysics, 667, A50, doi: 10.1051/0004-6361/202243890
Noraz, Q., Brun, A. S., Strugarek, A., & Depambour, G. 2022b, Astronomy & Astrophysics, 658, A144, doi: 10.1051/0004-6361/202141946
Rasmussen, C. E., & Williams, C. K. I. 2006, Gaussian Processes for Machine Learning (Cambridge, MA: MIT Press)
Reiners, A., & Mohanty, S. 2012, \textbackslashapj, 746, 43, doi: 10.1088/0004-637X/746/1/43
Ricker, G. R., Winn, J. N., Vanderspek, R., et al. 2014, in \textbackslashprocspie, Vol. 9143, Space Telescopes and Instrumentation 2014: Optical, Infrared, and Millimeter Wave, 914320, doi: 10.1117/12.2063489
Santos, A. R. G., Cunha, M. S., Avelino, P. P., García, R. A., & Mathur, S. 2017, Astronomy & Astrophysics, 599, A1, doi: 10.1051/0004-6361/201629923
Saunders, N., van Saders, J. L., Lyttle, A. J., et al. 2024, The Astrophysical Journal, 962, 138, doi: 10.3847/1538-4357/ad1516
Sills, A., Pinsonneault, M. H., & Terndrup, D. M. 2000, The Astrophysical Journal, 534, 335, doi: 10.1086/308739
Skumanich, A. 1972, \textbackslashapj, 171, 565, doi: 10.1086/151310
Speagle, J. S. 2020, Monthly Notices of the Royal Astronomical Society, 493, 3132, doi: 10.1093/mnras/staa278
van Saders, J. L., Ceillier, T., Metcalfe, T. S., et al. 2016, \textbackslashnat, 529, 181, doi: 10.1038/nature16168
van Saders, J. L., & Pinsonneault, M. H. 2013, The Astrophysical Journal, 776, 67, doi: 10.1088/0004-637X/776/2/67
Virtanen, P., Gommers, R., Oliphant, T. E., et al. 2020, Nature Methods, 17, 261, doi: 10.1038/s41592-019-0686-2
Weber, E. J., & Davis, Jr., L. 1967, The Astrophysical Journal, 148, 217, doi: 10.1086/149138

# APPENDIX

## A. FOURIER TRANSFORM PROPERTIES AND PAIRS

**Table 13.** Properties of the Fourier transform. We use the convention $\hat{f}(\omega) = \int_{-\infty}^{\infty} f(t)\, e^{-i\omega t}\, dt$. Here $a, b$ are constants, $t_0$ and $\omega_0$ are fixed shifts, and $f$, $g$ are functions with transforms $\hat{f}$, $\hat{g}$.

| No. | Property | Time domain $\longleftrightarrow$ Frequency domain |
|---|---|---|
| FT1 | Linearity | $a\, f(t) + b\, g(t) \;\longleftrightarrow\; a\, \hat{f}(\omega) + b\, \hat{g}(\omega)$ |
| FT2 | Time shifting | $f(t - t_0) \;\longleftrightarrow\; e^{-i\omega t_0}\, \hat{f}(\omega)$ |
| FT3 | Frequency shifting | $e^{i\omega_0 t}\, f(t) \;\longleftrightarrow\; \hat{f}(\omega - \omega_0)$ |
| FT4 | Convolution | $(f * g)(t) \;\longleftrightarrow\; \hat{f}(\omega) \cdot \hat{g}(\omega)$ |
| FT5 | Multiplication | $f(t) \cdot g(t) \;\longleftrightarrow\; \dfrac{1}{2\pi}\, \hat{f}(\omega) * \hat{g}(\omega)$ |
| FT6 | Even symmetry | $f(t)$ real and even $\implies$ $\hat{f}(\omega)$ purely real |
| FT7 | Odd symmetry | $f(t)$ real and odd $\implies$ $\hat{f}(\omega)$ purely imaginary |

**Table 14.** Fourier transform pairs used in this work. We use the convention $\hat{f}(\omega) = \int_{-\infty}^{\infty} f(t)\, e^{-i\omega t}\, dt$ and $\operatorname{sinc}(x) \equiv \sin(x)/x$.

| No. | Function | $f(t)$ | $\hat{f}(\omega)$ |
|---|---|---|---|
| P1 | Rectangular pulse | $\operatorname{rect}_W(t) = \begin{cases} 1 & \lvert t\rvert \le W/2 \\ 0 & \text{otherwise} \end{cases}$ | $W \operatorname{sinc}\left(\dfrac{\omega W}{2}\right)$ |
| P2 | Cosine | $\cos(\omega_0 t + \phi)$ | $\pi\left[e^{i\phi}\, \delta(\omega - \omega_0) + e^{-i\phi}\, \delta(\omega + \omega_0)\right]$ |
| P3 | Complex exponential | $e^{i\omega_0 t}$ | $2\pi\, \delta(\omega - \omega_0)$ |
| P4 | Square wave | $\operatorname{sgn}(\cos \omega_0 t)$ | $2\pi \displaystyle\sum_{n\,\text{odd}} \frac{(-1)^{(n-1)/2}}{n}\, \delta(\omega - n\omega_0)$ |
| P5 | Dirac delta | $\delta(t - t_0)$ | $e^{-i\omega t_0}$ |
| P6 | Dirac comb | $\displaystyle\sum_n \delta(t - nT)$ | $\dfrac{2\pi}{T} \displaystyle\sum_n \delta(\omega - n\omega_0), \quad \omega_0 = 2\pi/T$ |

## B. FOURIER COEFFICIENTS OF THE VISIBILITY FUNCTION

Here we derive the Fourier coefficients $c_n$ of the visibility function $\mathcal{V}(t) = \max\{\cos\beta(t), 0\}$ from first principles.[9]

### B.1. *Setup*

From Equation 6, we have:

$$\cos\beta(t) = \underbrace{\cos I \sin\Phi}_{b_0} + \underbrace{\sin I \cos\Phi}_{b_1} \cos(\Lambda_0 + \omega_0 t). \tag{B1}$$

This is a shifted cosine oscillating between $b_0 - b_1$ and $b_0 + b_1$. The spot is visible when $\cos\beta(t) > 0$, which (for $b_1 > 0$, $|b_0| < b_1$) occurs when:

$$|\omega_0 t + \Lambda_0| < \theta_{\rm vis}, \qquad \theta_{\rm vis} = \arccos\left(-\frac{b_0}{b_1}\right). \tag{B2}$$

This defines the *duty cycle* $\Delta = 2\theta_{\rm vis}/(2\pi) = \theta_{\rm vis}/\pi$ of the visibility window. When $b_0 + b_1 < 0$ the spot is never visible; when $b_0 - b_1 > 0$ the spot is always visible ($\theta_{\rm vis} = \pi$). Figure 30 illustrates the components of the visibility function $\mathcal{V}(t)$t.





**Figure 30.** Visibility function $\mathcal{V}(t)$ for a single spot. The spot is visible when $\cos\beta(t) > 0$, which occurs for a fraction $\Delta = 2\theta_{\rm vis}/(2\pi)$ of the rotation period. The visibility function is a truncated cosine, with amplitude $b_1$ and offset $b_0$.

Since $\mathcal{V}(t)$ is periodic with period $P = 2\pi/\omega_0$, we expand it in a Fourier series:

$$\mathcal{V}(t) = \sum_{n=-\infty}^{\infty} c_n\, e^{in\omega_0 t}, \tag{B3}$$

[9] Symbolic and numerical checks: `derive_ft_spot_rotation.ipynb`

where the Fourier coefficients are:

$$c_n = \frac{1}{T}\int_0^T \mathcal{V}(t)\, e^{-in\omega_0 t}\, dt = \frac{1}{2\pi}\int_{-\theta_{\text{vis}}}^{\theta_{\text{vis}}} \left(b_0 + b_1\cos\theta\right) e^{-in\theta}\, d\theta, \tag{B4}$$

with the substitution $\theta = \omega_0 t + \Lambda_0$ (and without loss of generality setting $\Lambda_0 = 0$, since a phase shift only affects the complex phase of $c_n$ but not $|c_n|^2$).

### B.2. *Evaluating the Integral*

We split the integrand into the constant and cosine parts:

$$c_n = \frac{1}{2\pi}\left[b_0\int_{-\theta_{\text{vis}}}^{\theta_{\text{vis}}} e^{-in\theta}\, d\theta + b_1\int_{-\theta_{\text{vis}}}^{\theta_{\text{vis}}} \cos\theta\; e^{-in\theta}\, d\theta\right]. \tag{B5}$$

*First integral (constant part).*—For $n \neq 0$:

$$\int_{-\theta_{\text{vis}}}^{\theta_{\text{vis}}} e^{-in\theta}\, d\theta = \left[\frac{e^{-in\theta}}{-in}\right]_{-\theta_{\text{vis}}}^{\theta_{\text{vis}}} = \frac{e^{-in\theta_{\text{vis}}} - e^{in\theta_{\text{vis}}}}{-in} = \frac{2\sin(n\,\theta_{\text{vis}})}{n}. \tag{B6}$$

For $n = 0$ this gives $2\theta_{\text{vis}}$.

*Second integral (cosine part).*—Using $\cos\theta = \frac{1}{2}(e^{i\theta} + e^{-i\theta})$:

$$\int_{-\theta_{\text{vis}}}^{\theta_{\text{vis}}} \cos\theta\; e^{-in\theta}\, d\theta = \frac{1}{2}\int_{-\theta_{\text{vis}}}^{\theta_{\text{vis}}} e^{i(1-n)\theta}\, d\theta + \frac{1}{2}\int_{-\theta_{\text{vis}}}^{\theta_{\text{vis}}} e^{-i(1+n)\theta}\, d\theta. \tag{B7}$$

Each integral is of the same form as the first. For $m \neq 0$:

$$\int_{-\theta_{\text{vis}}}^{\theta_{\text{vis}}} e^{im\theta}\, d\theta = \frac{2\sin(m\,\theta_{\text{vis}})}{m}. \tag{B8}$$

Therefore, for $n \neq \pm 1$:

$$\int_{-\theta_{\text{vis}}}^{\theta_{\text{vis}}} \cos\theta\; e^{-in\theta}\, d\theta = \frac{\sin\left((1-n)\theta_{\text{vis}}\right)}{1-n} + \frac{\sin\left((1+n)\theta_{\text{vis}}\right)}{1+n}. \tag{B9}$$

*Combining.*—Substituting back into Eq. (B5), for $n \neq 0, \pm 1$:

$$c_n = \frac{1}{\pi}\left[\frac{b_0\sin(n\,\theta_{\text{vis}})}{n} + \frac{b_1}{2}\left(\frac{\sin\left((n-1)\theta_{\text{vis}}\right)}{n-1} + \frac{\sin\left((n+1)\theta_{\text{vis}}\right)}{n+1}\right)\right], \tag{B10}$$

where we used $\sin\left((1-n)x\right) = -\sin\left((n-1)x\right)$ and $1/(1-n) = -1/(n-1)$.

## C. FOURIER TRANSFORM OF THE TRAPEZOIDAL ENVELOPE

Here we derive the Fourier transform and autocorrelation function of the trapezoidal envelope from Section 2.2; the transform of the unsquared trapezoid $\alpha(t)$ is derived in `derive_ft_trapezoid.ipynb`. These equations are derived and verified with `SymPy` in `derive_ft_squared_trapezoid.ipynb`.

### C.1. *Piecewise Definition of* $\Gamma(t)$

The trapezoidal spot evolution $\alpha(t)$, centered at $t = 0$, has a flat plateau of duration $\ell_{\rm spot}$ at amplitude $\alpha_{\rm max}$ and linear ramps of duration $\tau_{\rm spot}$ on each side:

$$\alpha(t) = \alpha_{\rm max} \begin{cases} 0 & t < -\frac{\ell_{\rm spot}}{2} - \tau_{\rm spot} \\ \dfrac{t + \frac{\ell_{\rm spot}}{2} + \tau_{\rm spot}}{\tau_{\rm spot}} & -\frac{\ell_{\rm spot}}{2} - \tau_{\rm spot} \le t < -\frac{\ell_{\rm spot}}{2} \\ 1 & -\frac{\ell_{\rm spot}}{2} \le t \le \frac{\ell_{\rm spot}}{2} \\ \dfrac{-t + \frac{\ell_{\rm spot}}{2} + \tau_{\rm spot}}{\tau_{\rm spot}} & \frac{\ell_{\rm spot}}{2} < t \le \frac{\ell_{\rm spot}}{2} + \tau_{\rm spot} \\ 0 & t > \frac{\ell_{\rm spot}}{2} + \tau_{\rm spot} \end{cases} \tag{C11}$$

Squaring and normalizing the trapezoid (C11):

$$\Gamma(t) = \frac{\alpha^2(t)}{\alpha_{\rm max}^2} = \begin{cases} 0 & |t| > \ell/2 + \tau \\ \left( \dfrac{|t| - \ell/2 - \tau}{-\tau} \right)^2 & \ell/2 < |t| \le \ell/2 + \tau \\ 1 & |t| \le \ell/2 \end{cases} \tag{C12}$$

where $\ell \equiv \ell_{\rm spot}$ and $\tau \equiv \tau_{\rm spot}$ for brevity. The ramps are now *parabolic* (concave) rather than linear.

### C.2. *Direct Integration*

Since $\Gamma(t)$ is real and even, $\hat{\Gamma}(\omega)$ is purely real:

$$\hat{\Gamma}(\omega) = 2\int_0^\infty \Gamma(t)\cos(\omega t)\,dt = 2\left[ \underbrace{\int_0^{\ell/2} \cos(\omega t)\,dt}_{I_1} + \underbrace{\int_{\ell/2}^{\ell/2+\tau} \left( \frac{\ell/2 + \tau - t}{\tau} \right)^2 \cos(\omega t)\,dt}_{I_2} \right]. \tag{C13}$$

*Integral $I_1$ (plateau):*—

$$I_1 = \frac{\sin(\omega\ell/2)}{\omega}. \tag{C14}$$

*Integral $I_2$ (parabolic ramp):*—With $a = \ell/2$ and $b = \ell/2 + \tau$, we integrate by parts twice. Setting $f(t) = \big((b-t)/\tau\big)^2$:

*First integration by parts* ($u = f(t)$, $dv = \cos(\omega t)\,dt$):

$$I_2 = \left[ \frac{f(t)\sin(\omega t)}{\omega} \right]_a^b + \frac{2}{\tau^2\omega}\int_a^b (b-t)\sin(\omega t)\,dt. \tag{C15}$$

The boundary term gives $0 - f(a)\sin(\omega a)/\omega = -\sin(\omega\ell/2)/\omega$ (since $f(a) = 1$).

*Second integration by parts* on $J = \int_a^b (b-t)\sin(\omega t)\,dt$ ($u = b - t$, $dv = \sin(\omega t)\,dt$):

$$J = \left[ -\frac{(b-t)\cos(\omega t)}{\omega} \right]_a^b - \frac{1}{\omega}\int_a^b \cos(\omega t)\,dt = \frac{\tau\cos(\omega\ell/2)}{\omega} - \frac{\sin(\omega b) - \sin(\omega\ell/2)}{\omega^2}. \tag{C16}$$

Combining:

$$I_2 = -\frac{\sin(\omega\ell/2)}{\omega} + \frac{2\cos(\omega\ell/2)}{\tau\omega^2} - \frac{2\left[\sin\left(\frac{\omega\ell}{2}+\omega\tau\right) - \sin\left(\frac{\omega\ell}{2}\right)\right]}{\tau^2\omega^3}. \tag{C17}$$

*Combining $I_1 + I_2$:*—The $\sin(\omega\ell/2)/\omega$ terms from $I_1$ and $I_2$ cancel:

$$I_1 + I_2 = \frac{2\cos(\omega\ell/2)}{\tau\omega^2} + \frac{2\sin(\omega\ell/2)}{\tau^2\omega^3} - \frac{2\sin\left(\frac{\omega\ell}{2}+\omega\tau\right)}{\tau^2\omega^3}. \tag{C18}$$

Therefore:

$$\hat{\Gamma}(\omega) = \frac{4}{\tau^2\omega^3}\left[\tau\omega\cos\left(\frac{\omega\ell}{2}\right) + \sin\left(\frac{\omega\ell}{2}\right) - \sin\left(\frac{\omega\ell}{2}+\omega\tau\right)\right]. \tag{C19}$$

## D. AUTOCORRELATION OF THE SQUARED ENVELOPE

Next we derive the closed-form autocorrelation $R_\Gamma(\tau) = \int_{-\infty}^{\infty} \Gamma(s)\,\Gamma(s+\tau)\,ds$ of the normalized squared trapezoidal envelope $\Gamma(t) = \alpha^2(t)/\alpha^2_{\max}$ defined in Eq. (C12). The `SymPy` derivation and its checks are in `derive_R_Gamma.ipynb`.

### D.1. *Setup*

The function $\Gamma(t)$ has three regions: a plateau of height 1 for $|t| \le \ell/2$, parabolic ramps for $\ell/2 < |t| \le S$, and zero otherwise. The autocorrelation integral

$$R_\Gamma(\tau) = \int_{-S}^{S-\tau} \Gamma(s)\,\Gamma(s+\tau)\,ds, \qquad \tau \ge 0, \tag{D20}$$

must be split into sub-intervals where both $\Gamma(s)$ and $\Gamma(s+\tau)$ have known polynomial forms. The breakpoints of $\Gamma(s)$ are at $s = \pm\ell/2$ and $s = \pm S$, while those of $\Gamma(s+\tau)$ are shifted by $-\tau$. The relevant sub-intervals and the character of the overlap change as $\tau$ increases through the four intervals $[0,\ \tau_s]$, $[\tau_s,\ \ell]$, $[\ell,\ \ell+\tau_s]$, and $[\ell+\tau_s,\ \ell+2\tau_s]$. The result below holds whenever $\ell \ge \tau_s$ (plateau at least as long as one ramp), which is typical for observed starspots; for $\ell < \tau_s$ the intervals reorder and the closed form no longer applies.

### D.2. *Result*

Evaluating the polynomial integrals in each sub-interval and summing (verified symbolically and numerically), the autocorrelation is:

$$R_\Gamma(\tau) = \begin{cases} \ell + \dfrac{2\tau_s}{5} - \dfrac{4\tau^2}{3\tau_s} + \dfrac{2\tau^3}{3\tau_s^2} - \dfrac{\tau^5}{15\tau_s^4} & 0 \le \tau \le \tau_s \\ \ell + \dfrac{2\tau_s}{3} - \tau & \tau_s \le \tau \le \ell \\ P_3(\tau) & \ell \le \tau \le \ell + \tau_s \\ \dfrac{(\ell + 2\tau_s - \tau)^5}{30\,\tau_s^4} & \ell + \tau_s \le \tau \le \ell + 2\tau_s \\ 0 & \tau > \ell + 2\tau_s \end{cases} \tag{D21}$$





**Figure 31.** Piecewise structure of the envelope autocorrelation. *Left:* Fourier transform $\hat{\Gamma}(\omega)$ (black) decomposed into the plateau sinc-like term $I_1$ (blue dashed) and the parabolic ramp term $I_2$ (orange dashed). *Right:* Autocorrelation $R_\Gamma(\tau)$ from Eq. (D21), with each of the four piecewise intervals shown in a different color. The thin gray curve shows the result of direct numerical convolution, confirming the closed-form expressions. Parameters: $\ell = 15\,\mathrm{d}$, $\tau_\mathrm{s} = 3\,\mathrm{d}$.

where the third piece is the degree-5 polynomial:

$$P_3(\tau) = \frac{\tau^5}{30\tau_\mathrm{s}^4} - \frac{(\ell + 2\tau_\mathrm{s})\,\tau^4}{6\tau_\mathrm{s}^4} + \frac{(\ell^2 + 4\ell\tau_\mathrm{s} + 2\tau_\mathrm{s}^2)\,\tau^3}{3\tau_\mathrm{s}^4} - \frac{\ell(\ell^2 + 6\ell\tau_\mathrm{s} + 6\tau_\mathrm{s}^2)\,\tau^2}{3\tau_\mathrm{s}^4} + \frac{(\ell^4 + 8\ell^3\tau_\mathrm{s} + 12\ell^2\tau_\mathrm{s}^2 - 6\tau_\mathrm{s}^4)\,\tau}{6\tau_\mathrm{s}^4} + \frac{-\ell^5 - 10\ell^4\tau_\mathrm{s} - 20\ell^3\tau_\mathrm{s}^2 + 30\ell\tau_\mathrm{s}^4 + 20\tau_\mathrm{s}^5}{30\tau_\mathrm{s}^4}. \tag{D22}$$

Figure 31 illustrates these results. The left panel shows the Fourier transform $\hat{\Gamma}(\omega)$ decomposed into the plateau contribution $I_1$ and the parabolic ramp contribution $I_2$. The right panel shows $R_\Gamma(\tau)$ with the four piecewise intervals color-coded, overlaid on a numerical convolution for verification.