Theoretical Limits in Constraining Tidal Quality Factors of Binary Stars

JESSICA BIRKY AND RORY BARNES

${\bf ABSTRACT}$

1. INTRODUCTION

2. METHODS

2.1. Observational Constraints


Model input parameters and prior constraints:

| Model Input | | Prior | Observational Constraint | Good Unc |
|-----------------------|---------------------------------|--------------------------|------------------------------------|----------|
| M_1 | primary mass $[M_{\odot}]$ | $\mathcal{N}(m,s)$ | kepler solution (lc eclipse + rvs) | 0.001 |
| M_2 | secondary mass $[M_{\odot}]$ | $\mathcal{N}(m,s)$ | kepler solution (lc eclipse + rvs) | 0.001 |
| $P_{\text{rot}1,i}$ | pri init rotation period [days] | $log \mathcal{N}(m,s)$ | dist in young open clusters | |
| $P_{\mathrm{rot}2,i}$ | sec init rotation period [days] | $log \mathcal{N}(m,s)$ | dist in young open clusters | |
| $P_{\mathrm{orb},i}$ | init orbital period [days] | $\mathcal{U}(4.0, 10.0)$ | uninformed | |
| e_i | init eccentricity | $\mathcal{U}(0,0.5)$ | uninformed | |
| age | system age [yr] | $\mathcal{N}(m,s)$ | open cluster age | 10% |
| $\varepsilon_{1,i}$ | pri init obliquity [deg] | $\mathcal{U}(0,30)$ | uninformed | |
| $\varepsilon_{2,i}$ | sec init obliquity [deg] | $\mathcal{U}(0,30)$ | uninformed | |
| \mathcal{Q}_1 | pri tidal phase lag | $\mathcal{U}(4,9)$ | uninformed | |
| \mathcal{Q}_2 | sec tidal phase lag | $\mathcal{U}(4,9)$ | uninformed | |
| $	au_1$ | pri tidal time lag [log(s)] | $\mathcal{U}(-4,2)$ | uninformed | |
| $	au_2$ | sec tidal time lag [log(s)] | $\mathcal{U}(-4,2)$ | uninformed | |

Model output parameters and likelihood constraints:

| | Model Output | Likelihood | Observational Constraint | Good Unc |
|-----------------------|------------------------------------|--------------------|-----------------------------|-----------|
| $P_{\text{rot}1,f}$ | pri final rotation period [days] | $\mathcal{N}(m,s)$ | lc autocorrelation function | 0.1 |
| $P_{\text{rot}2,f}$ | sec final rotation period [days] | $\mathcal{N}(m,s)$ | spectroscopic $v \sin i$ | 0.1 |
| $P_{\mathrm{orb},f}$ | final orbital period [days] | $\mathcal{N}(m,s)$ | lc lomb scargle | 10^{-5} |
| e_f | final eccentricity | $\mathcal{N}(m,s)$ | lc eclipse + rvs | 0.001 |
| $R_{1,f}$ | pri final radius $[R_{\odot}]$ | $\mathcal{N}(m,s)$ | stellar models + photometry | 0.01 |
| $R_{2,f}$ | sec final radius $[R_{\odot}]$ | $\mathcal{N}(m,s)$ | eclipse shape + pri radius | 0.01 |
| $L_{1,f}$ | pri final lumniosity $[L_{\odot}]$ | $\mathcal{N}(m,s)$ | stellar models + photometry | 0.1 |
| $L_{2,f}$ | sec final lumniosity $[L_{\odot}]$ | $\mathcal{N}(m,s)$ | stellar models + photometry | 0.1 |
| $T_{\mathrm{eff}1,f}$ | pri final temperature [K] | $\mathcal{N}(m,s)$ | stellar models + spectra | |
| $T_{{\rm eff}2,f}$ | sec final temperature [K] | $\mathcal{N}(m,s)$ | stellar models + spectra | |

BIRKY ET AL.








Figure 2. EQTIDE + STELLAR: CTL model



- 4. DISCUSSION
- 5. CONCLUSION



4 BIRKY ET AL.

APPENDIX

REFERENCES