

Regression

Part 1

Prof. Bisbee

Vanderbilt University

Slides Updated: 2024-01-09

Agenda

1. Modeling Conditional Variation
2. Adding Regression to the **Process**
3. Introducing the Data
4. Demonstrating Regressions

Regression & Conditional Analysis

- Recall our discussion of **conditional** analysis
 - Conditional → **depends on**
 - Analyze with **conditional means**

Reminder of the **Process**

1. Determine variable **type**

- I.e., categorical (ordered, unordered, binary) or continuous
- In R terms: `chr`, `fct` for categorical, `dbl` for continuous

2. Type informs **univariate analysis**


- I.e., histograms for continuous, barplots for categorical

3. Combination of types informs **conditional analysis**

- Categorical X Categorical: proportions by categories (`geom_bar`)
- Binary X Continuous: histograms by categories (`geom_histogram` / `geom_density`)
- Categorical X Continuous: distributions by categories (`geom_boxplot` / `geom_violin`)
- Continuous X Continuous: scatter plots (`geom_point`)


What is regression?

- Conditional means for continuous data




What is regression?

- People with hourly wages < \$20 spend ~\$50 on entertainment per week




What is regression?

- People with hourly wages $> \$40$ spend $\sim \$95$ on entertainment per week




What is regression?

- **Theory**: the more you earn, the more you spend




What is regression?

- But **conditional means** make a lot of mistakes. Can we do better?



What is regression?


- But **conditional means** make a lot of mistakes. Can we do better?




Regression

- Calculating a **line** that minimizes mistakes *for every observation*
 - NB: could be a curvey line! For now, just assume straight
- Recall from geometry how to graph a straight line
- $Y = a + bX$
 - a : the "intercept" (where the line intercepts the y-axis)
 - b : the "slope" (how much Y changes for each increase in X)
- (Data scientists use α and β instead of a and b b/c nerds)
- Regression analysis simply chooses the best line
 - "Best"?
 - The line that minimizes the mistakes (the **line of best fit**)


Linear Regression



Linear Regression




Linear Regression




- **Error/Residual:** mistake made by a line
 - In math: $u_i = y_i - \hat{y}_i$
 - In English: difference between true outcome value (y_i) and prediction (\hat{y}_i)

Linear Regression




- Use **errors** to find **line of best fit**
- **RMSE** (**R**oot **M**ean **S**quared **E**rror)
 - Square the errors
 - Take their average
 - Take the square root
- **RMSE** = 1.23

Linear Regression




- Use **errors** to find **line of best fit**
- **RMSE** (**R**oot **M**ean **S**quared **E**rror)
 - Square the errors
 - Take their average
 - Take the square root
- **RMSE** = 1.48

Linear Regression




- Use **errors** to find **line of best fit**
- **RMSE** (**R**oot **M**ean **S**quared **E**rror)
 - Square the errors
 - Take their average
 - Take the square root
- **RMSE** = 2.19

Linear Regression




- Use **errors** to find **line of best fit**
- **RMSE (Root Mean Squared Error)**
 - Square the errors
 - Take their average
 - Take the square root
- **RMSE = 1.03**

Visual Intuition



Visual Intuition



Regression

- The line is **substantively meaningful**
- Red line on scatter plot of spending and wages: $Y = 12 + 2 * X$
- α tells us the value of Y when X is zero
 - People who don't make any money spend \$12 per week on entertainment
- β tells us how much Y increases for each additional X
 - People spend an additional \$2 per week for each additional \$1 in hourly wages

Regression

- These are called "**linear models**"
 - **Not** because the line is straight (it might not be)
 - but because the components are additive ($\alpha + \beta X$)
- Can extend to multiple predictors (X 's)
 - $Y = \alpha + \beta_1 X_1 + \beta_2 X_2 + \dots + \varepsilon$
 - X_1 might be wages and X_2 might be age (for example)
 - The final term ε measures how bad our mistakes are

Regression

- Let's demonstrate with the `debt` data

```
require(tidyverse)

debt <-
read_rds('https://github.com/jbisbee1/DS1000_S2024/raw/main/data/sc_deb
glimpse(debt)
```

```
## Rows: 2,546
## Columns: 16
## $ unitid      <int> 100654, 100663, 100690, 100706, 100...
## $ instnm      <chr> "Alabama A & M University", "Univer...
## $ stabbr      <chr> "AL", "AL", "AL", "AL", "AL", "AL", ...
## $ grad_debt_mdn <int> 33375, 22500, 27334, 21607, 32000, ...
## $ control     <chr> "Public", "Public", "Private", "Pub...
## $ region      <chr> "Southeast", "Southeast", "Southeas...
## $ preddeg      <chr> "Bachelor's", "Bachelor's", "Associ...
## $ openadmp     <int> 2, 2, 1, 2, 2, 2, 1, NA, 2, 2, 2, 1...
## $ adm_rate     <dbl> 0.9175, 0.7366, NA, 0.8257, 0.9690, ...
## $ ccbasic      <int> 18, 15, 20, 16, 19, 15, 2, 22, 18, ...
## $ sat_avg      <int> 939, 1234, NA, 1319, 946, 1261, NA, ...
```

Research Camp

- **Research Question:** What is the relationship between SAT scores and median future earnings?
- **Theory:** Students with higher SAT scores work harder and have learned more. Employers reward these attributes with higher wages in the private market.
- **Hypothesis:** The relationship between SAT scores and future earnings should be positive.
 - **NB:** Important caveats to this simplistic theory!
 - Socioeconomic status: predicts both higher SAT scores and higher wages
 - **Correlation \neq Causation**

Set Up

- Linking Theory to Data
- Our SAT scores are theorized to explain future earnings
 - Thus the SAT scores are the independent / explanatory / predictor variable X
 - And earnings are the dependent / outcome variable Y

Regression

- There is a simple recipe to follow
- And it is exactly how the syllabus for the class is designed!
 1. Look at your data to identify missingness (**Wrangling: Lecture 5**)
 2. **Univariate** visualization of your variables (**Lecture 6**)
 3. **Multivariate** visualization of your variables (**Lectures 7-10**)
 4. **Regression** (today)
 5. Evaluation of **errors** (next lecture)

Step 1: Look

- Why worry about **missingness**?

1. **Substantive:** external validity

2. **Technical:** cross validation won't work! (Wednesday's lecture)

```
summary(debt %>% select(sat_avg,md_earn_wne_p6))
```

```
##      sat_avg      md_earn_wne_p6
##  Min.      : 737    Min.      : 10600
## 1st Qu.:1053    1st Qu.: 26100
## Median :1119    Median : 31500
## Mean   :1141    Mean   : 33028
## 3rd Qu.:1205    3rd Qu.: 37400
## Max.   :1557    Max.   :120400
## NA's   :1317    NA's   :240
```


Step 2: Univariate Viz

- Why visualize both Y and X ?
 1. **Substantive:** See which units you are talking about
 2. **Technical:** Adjust for *skew*

Step 2: Univariate Viz

- Why visualize both Y and X ?


```
debt %>%  
  ggplot(aes(x = sat_avg)) +  
  geom_histogram()
```



Step 2: Univariate Viz

- Why visualize both Y and X ?


```
debt %>%  
  ggplot(aes(x = md_earn_wne_p6)) +  
  geom_histogram()
```



Step 3: Multivariate

- Eyeball the relationship first!


```
debt %>%  
  ggplot(aes(x = sat_avg, y = md_earn_wne_p6)) +  
  geom_point()
```



Step 3: Multivariate Viz

- Adding regression line

```
debt %>%  
  ggplot(aes(x = sat_avg, y = md_earn_wne_p6)) +  
  geom_point() + geom_smooth(method = 'lm', se = F)
```



Step 3: Multivariate Viz


- Let's focus on two schools

```
toplot <- debt %>%  
  mutate(hl = ifelse(unitid %in% c(100654,179265), 'hl', 'none')) #  
Choosing two examples  
p2 <- toplot %>%  
  ggplot(aes(x = sat_avg, y = md_earn_wne_p6,color = hl,group =  
1,alpha = hl)) +  
  geom_point(data = toplot %>% filter(hl == 'none')) +  
  geom_point(data = toplot %>% filter(hl == 'hl'),size =3) +  
  scale_alpha_manual(values = c(1,.3)) +  
  scale_color_manual(values = c('red','black')) +  
  geom_smooth(method = 'lm',se = F) +  
  theme(legend.position = 'none') +  
  labs(title = "Graduate Earnings and SAT Scores",  
        subtitle = "By School",  
        x = "Average SAT Score",  
        y = "Median Earnings of Grads")
```

Step 3: Multivariate Viz

- Adding regression line

p2



Step 3: Multivariate Viz


- Defining ε

```
p3 <- toplot %>%
  ggplot(aes(x = sat_avg, y = md_earn_wne_p6,color = h1,group =
1,alpha = h1)) +
  geom_point(data = toplot %>% filter(h1 == 'none')) +
  geom_point(data = toplot %>% filter(h1 == 'h1'),size =3) +
  scale_alpha_manual(values = c(1,.3)) +
  scale_color_manual(values = c('red','black')) +
  geom_smooth(method = 'lm',se = F) +
  annotate(geom = 'segment',
    x = toplot %>% filter(h1 == 'h1') %>% .$sat_avg,
    y = toplot %>% filter(h1 == 'h1') %>% .$md_earn_wne_p6,
    xend = toplot %>% filter(h1 == 'h1') %>% .$sat_avg,
    yend = c(27500,41000),color = 'red',lwd = 1.2) +
  theme(legend.position = 'none') +
  labs(title = "Graduate Earnings and SAT Scores",
    subtitle = "By School",
    x = "Average SAT Score",
    y = "Median Earnings of Grads")
```

Step 3: Multivariate Viz

- Measuring errors

p3




The Data Scientist's Trade-off

- Those mistakes seem pretty big!
- Why not use a curvier line?


The Data Scientist's Trade-off

- Those mistakes seem pretty big!
- Why not use a curvier line?




The Data Scientist's Trade-off

- Those mistakes seem pretty big!
- Why not use a curvier line?




The Data Scientist's Trade-off

- Those mistakes seem pretty big!
- Why not use a curvier line?



The Data Scientist's Trade-off

- Those mistakes seem pretty big!
- Why not use a curvier line?



The Data Scientist's Trade-off

- Want to **reduce complexity**
- But also want to be **accurate**
- What is the right answer?
 - It depends on your **theory** and the **data**
 - It is context-dependent
- And this is still only using *linear regression models*!
 - This is a deep area of study, for those interested

Step 4: Regression

- Introducing the `lm(formula,data)` function
- Two inputs to care about:
 - `formula`: Code for $Y = \alpha + \beta X$
 - `data`: What is the data we are using?
- `formula` is written as $Y \sim X$
 - R will calculate α and β for us
 - Just need to tell it what is Y (`md_earn_wne_p6`) and X (`sat_avg`)
 - The tilde (`~`) is R's version of the equals sign
- Save the model to an object

```
model_earn_sat <- lm(formula = md_earn_wne_p6 ~ sat_avg, data = debt)
```

Step 4: Interpretation

- What is in this object?
- The regression results! Look at them with `summary()`

```
summary(model_earn_sat)
```

```
##
## Call:
## lm(formula = md_earn_wne_p6 ~ sat_avg, data = debt)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -23239  -4311   -852    2893   78695
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -12053.87   1939.80  -6.214 7.12e-10 ***
## sat_avg      42.60      1.69   25.203 < 2e-16 ***
## ---
## Signif. codes:
## 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
```

Step 4: Interpretation

- Starting with the first column called **Estimate**
- 1st Row (**Intercept**) is α : the predicted value of Y when X is zero
 - Schools with average SAT scores of 0 produce graduates who earn -\$12,053.87
 - Sensible?
- 2nd Row **sat_avg** is the β : the increase in Y when X increases by one
 - For each unit increase in the average SAT score, recent graduates earn \$42.60 more
 - Sensible?

Step 4: Interpretation

- Other 3 columns?
 - `Std. Error` is the "standard error"
 - `t value` is the "t-statistic"
 - `Pr(>|t|)` is the "p-value"
- $t\text{-statistic} = \text{Estimate} / \text{standard error}$
- $p\text{-value} = \text{function}(t\text{-statistic})$
 - Only really need to remember the p-value for this course
 - This is 1 minus confidence
 - The lower the p-value, the more confident we are that the `Estimate` is not zero

Step 4: Interpretation

```
summary(model_earn_sat)
```

```
##
## Call:
## lm(formula = md_earn_wne_p6 ~ sat_avg, data = debt)
##
## Residuals:
##      Min       1Q   Median       3Q      Max
## -23239  -4311   -852    2893   78695
##
## Coefficients:
##              Estimate Std. Error t value Pr(>|t|)
## (Intercept) -12053.87    1939.80  -6.214 7.12e-10 ***
## sat_avg      42.60       1.69   25.203 < 2e-16 ***
## ---
## Signif. codes:
## 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
## Residual standard error: 7594 on 1196 degrees of freedom
## (1348 observations deleted due to missingness)
## Multiple R-squared:  0.3469,    Adjusted R-squared:  0.3463
## F-statistic: 635.2 on 1 and 1196 DF,  p-value: < 2.2e-16
```

Another Example

- We will come back to the RMSE next class
- For now, let's try with a different research question!
- What is the relationship between admissions and future earnings?
 - Theory: More selective schools are more prestigious
 - Hypothesis: There should be a negative relationship between the admissions rate and future earnings

Do It Together!

1. Look at the data and acknowledge missingness
2. Univariate visualization of X and Y
3. Multivariate visualization of X and Y
4. Regression

Quiz & Homework

- Go to Brightspace and take the **11th** quiz
 - The password to take the quiz is ####
- **Homework:**
 1. Work through ds1000_hw_12.Rmd
 2. Problem set 7 (last before the midterm!)