

Supplemental Information for:
 “Mount Fuji’s stubby peak: the genotypic density of
 additive landscapes near maximal fitness”

Justin B. Kinney

May 6, 2026

Contents

S1 Cumulants of the fitness distribution for additive landscapes	1
S2 Supporting derivations for the near-peak scaling	2
S2.1 Evaluation of $\int_0^\infty \log(1 + e^{-u}) du$	2
S2.2 Generalized integral identity	3
S2.3 Derivation of the generalized scaling exponent	4
S3 Gap density for continuous distributions	5
S3.1 General formula from order statistics	5
S3.2 Evaluation for Gaussian effects	5
S4 Near-peak scaling for the Hamming-distance landscape	6
S5 Global epistasis coefficients	6

S1 Cumulants of the fitness distribution for additive landscapes

Using the expression for $\Phi(\beta)$ in Eq. 7 of the main text and the fact that Φ is a cumulant generating function, the mean and variance of $p_\beta(F)$ are found to be:

$$\mu_\beta = \Phi'(\beta) = \theta_0 + \sum_l \sum_c w_{lc} \theta_{lc}, \quad (\text{S1})$$

$$\sigma_\beta^2 = \Phi''(\beta) = \sum_l \left[\sum_c w_{lc} \theta_{lc}^2 - \left(\sum_c w_{lc} \theta_{lc} \right)^2 \right], \quad (\text{S2})$$

where the weights w_{lc} are defined to be

$$w_{lc} = \frac{e^{\beta \theta_{lc}}}{\sum_{c'} e^{\beta \theta_{lc'}}}. \quad (\text{S3})$$

More generally, the n th cumulant is given by $\kappa_n = \Phi^{(n)}(\beta)$. Φ scales linearly with L , and so do all these cumulants, including σ_β^2 . The standardized cumulants therefore scale as $\kappa_n / \sigma_\beta^n = O(L^{1-n/2})$,




Figure S1: Near-peak scaling for binary additive landscapes plotted against ϵ^α , where $\epsilon = (F_{\max} - F)/L$ and α is the analytic near-peak exponent for each panel. Curves are the same as in Fig. 3.

and thus vanish as $L \rightarrow \infty$ for all $n \geq 3$. This confirms that the tilted fitness distribution converges to a Gaussian in the large L limit (Eq. 8 of the main text).

S2 Supporting derivations for the near-peak scaling

This section provides mathematical details supporting the near-peak scaling results in Eqs. 17–18 of the main text.

S2.1 Evaluation of $\int_0^\infty \log(1 + e^{-u}) du$

The integral $\int_0^\infty \log(1 + e^{-u}) du$ appearing in Eq. 16 of the main text is evaluated by expanding the logarithm as a series:

$$\int_0^\infty \log(1 + e^{-u}) du = \sum_{k=1}^\infty \frac{(-1)^{k+1}}{k} \int_0^\infty e^{-ku} du = \sum_{k=1}^\infty \frac{(-1)^{k+1}}{k^2} = \eta(2), \quad (\text{S4})$$

where $\eta(s) = \sum_{k=1}^\infty (-1)^{k+1}/k^s$ is the Dirichlet eta function. To evaluate $\eta(s)$, we split the sum into odd and even terms:

$$\eta(s) = \sum_{k=1}^\infty \frac{1}{k^s} - 2 \sum_{k=1}^\infty \frac{1}{(2k)^s} = \zeta(s) - 2^{1-s} \zeta(s) = (1 - 2^{1-s}) \zeta(s), \quad (\text{S5})$$




Figure S2: Near-peak scaling for general alphabets plotted against ϵ^α , where α is fit separately for each panel. Curves are the same as in Fig. 4.

where $\zeta(s) = \sum_{k=1}^{\infty} 1/k^s$ is the Riemann zeta function. Using the Basel identity $\zeta(2) = \pi^2/6$, we obtain $\eta(2) = (1 - 2^{-1}) \cdot \pi^2/6 = \pi^2/12$, and thus

$$\int_0^\infty \log(1 + e^{-u}) du = \frac{\pi^2}{12}. \quad (\text{S6})$$

S2.2 Generalized integral identity

The derivation in Sec. S2.3 below relies on the identity

$$I(\gamma) \equiv \int_0^\infty \log(1 + e^{-u}) u^\gamma du = \Gamma(1 + \gamma) \eta(2 + \gamma), \quad \gamma > -1. \quad (\text{S7})$$

To derive this identity, expand $\log(1 + e^{-u}) = \sum_{k=1}^{\infty} (-1)^{k+1} e^{-ku}/k$ and integrate term by term:

$$I(\gamma) = \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} \int_0^\infty u^\gamma e^{-ku} du = \sum_{k=1}^{\infty} \frac{(-1)^{k+1} \Gamma(1 + \gamma)}{k^{2+\gamma}} = \Gamma(1 + \gamma) \eta(2 + \gamma). \quad (\text{S8})$$

As a consistency check, setting $\gamma = 0$ gives $I(0) = \Gamma(1) \eta(2) = \pi^2/12$, recovering Eq. S4.




Figure S3: Near-peak genotypic densities for four transcription factors (TFs). (a) The ABF1 motif reported by Mustonen et al. [1]. (b-d) The CTCF, MAX, and GATA1 motifs from HOCOMOCO v14 [2]. Left panels show additive effects shifted and scaled so that F values correspond to z-scores [$F_{\text{mean}} = 0, \text{var}(F) = 1$]. Middle panels compare the saddle-point approximation, Touzet density estimate [3], Gaussian bulk approximation, and fitted near-peak scaling behavior (i.e., $A + B\epsilon^\alpha$) as functions of F . Right panels show the same curves as functions of ϵ^α . The values of A , B , and α used in each row were empirically determined from the corresponding saddle-point approximation.

S2.3 Derivation of the generalized scaling exponent

When the gap distribution behaves as $p_{\text{gap}}(\Delta) \sim c \Delta^\gamma$ near $\Delta = 0$ with $\gamma > -1$, the substitution $u = \beta\Delta$ in the integral form of $E_{\text{bin}}(\beta)$ (Eq. 15 of the main text) gives

$$E_{\text{bin}}(\beta) \sim \frac{c}{\beta^{1+\gamma}} \int_0^\infty \log(1 + e^{-u}) u^\gamma du = \frac{b}{\beta^{1+\gamma}}, \quad (\text{S9})$$

where $b = c\Gamma(1+\gamma)\eta(2+\gamma)$ using the identity in Eq. S7. The saddle-point equation $\epsilon = -E'_{\text{bin}}(\beta)$ then gives

$$\epsilon = \frac{(1+\gamma)b}{\beta^{2+\gamma}}, \quad \text{so} \quad \beta = \left[\frac{(1+\gamma)b}{\epsilon} \right]^{1/(2+\gamma)}. \quad (\text{S10})$$

From Eq. 13 of the main text, $\frac{1}{L} \log \rho_{\text{saddle}} \approx E_{\text{bin}}(\beta) + \beta\epsilon$. Combining these two terms:

$$E_{\text{bin}}(\beta) + \beta\epsilon = \frac{b}{\beta^{1+\gamma}} + \beta\epsilon = \frac{\epsilon\beta}{1+\gamma} + \beta\epsilon = \frac{(2+\gamma)}{(1+\gamma)} \beta\epsilon. \quad (\text{S11})$$

Substituting the expression for β yields

$$\frac{1}{L} \log \rho \sim \underbrace{\frac{(2+\gamma)}{(1+\gamma)} [(1+\gamma)b]^{1/(2+\gamma)}}_B \overbrace{\epsilon^{(1+\gamma)/(2+\gamma)}}^\alpha. \quad (\text{S12})$$

This yields the scaling exponent $\alpha = (1+\gamma)/(2+\gamma)$ and the coefficient B stated in Eq. 18 of the main text.

S3 Gap density for continuous distributions

The main text states that the condition $0 < p_{\text{gap}}(0) < \infty$ is always satisfied when the fitness effects θ_{lc} are independently drawn from the same continuous distribution. Here we prove this claim and evaluate $p_{\text{gap}}(0)$ for the Gaussian distribution used in simulations.

S3.1 General formula from order statistics

Let $\theta_{(1)} \geq \theta_{(2)} \geq \dots \geq \theta_{(C)}$ denote the order statistics of C i.i.d. draws from a density p_θ with CDF P_θ . The gap $\Delta = \theta_{(1)} - \theta_{(2)}$ between the largest and second-largest values has density

$$p_{\text{gap}}(\delta) = C(C-1) \int_{-\infty}^{\infty} p_\theta(m+\delta) p_\theta(m) P_\theta(m)^{C-2} dm, \quad \delta \geq 0, \quad (\text{S13})$$

which follows from the joint density of the top two order statistics, $f_{\theta_{(1)}, \theta_{(2)}}(x_1, x_2) = C(C-1) p_\theta(x_1) p_\theta(x_2) P_\theta(x_2)^{C-2}$ for $x_1 \geq x_2$, by substituting $\delta = x_1 - x_2$ and integrating over $m = x_2$. Setting $\delta = 0$ gives

$$p_{\text{gap}}(0) = C(C-1) \int_{-\infty}^{\infty} p_\theta(m)^2 P_\theta(m)^{C-2} dm. \quad (\text{S14})$$

We now show that $0 < p_{\text{gap}}(0) < \infty$ for any continuous density p_θ . The integrand $C(C-1) p_\theta(m)^2 P_\theta(m)^{C-2}$ is non-negative everywhere and strictly positive on the interior of the support of p_θ (where $p_\theta(m) > 0$ and $0 < P_\theta(m) < 1$). It follows that $p_{\text{gap}}(0) > 0$. For the upper bound, $p_\theta(m)^2 \leq \|p_\theta\|_\infty p_\theta(m)$ and $P_\theta(m)^{C-2} \leq 1$, so $p_{\text{gap}}(0) \leq C(C-1) \|p_\theta\|_\infty < \infty$.

S3.2 Evaluation for Gaussian effects

For $\theta_{lc} \sim \mathcal{N}(0, s^2)$, substituting $p_\theta(m) = \varphi(m/s)/s$ and $P_\theta(m) = \Phi_N(m/s)$ into Eq. S14 and changing variables to $z = m/s$ gives

$$p_{\text{gap}}(0) = \frac{C(C-1)}{s} \int_{-\infty}^{\infty} \varphi(z)^2 \Phi_N(z)^{C-2} dz, \quad (\text{S15})$$

where φ and Φ_N denote the standard normal density and CDF, respectively. For $C = 2$ the integral evaluates to $1/(2\sqrt{\pi})$, giving $p_{\text{gap}}(0) = 1/(\sqrt{\pi}s)$. For general C the integral must be computed numerically; the resulting dimensionless product $s \cdot p_{\text{gap}}(0)$ is tabulated below for the alphabet sizes used in the main text.

C	$s \cdot p_{\text{gap}}(0)$
2	0.564
4	1.029
20	1.867

S4 Near-peak scaling for the Hamming-distance landscape

The Hamming-distance landscape [4] assumes that every mutation away from the optimal sequence x_{max} incurs the same fitness cost $\Delta_0 > 0$. A sequence at Hamming distance k from x_{max} therefore has fitness $F_{\text{max}} - k\Delta_0$. Because there are $\binom{L}{k}(C-1)^k$ such sequences, the exact genotypic density is

$$\rho(F_{\text{max}} - k\Delta_0) = \binom{L}{k}(C-1)^k, \quad k = 0, 1, \dots, L. \quad (\text{S16})$$

For small per-site fitness deficit $\epsilon = k\Delta_0/L$ with $k \ll L$, Stirling's approximation $\log \binom{L}{k} \approx k \log(eL/k)$ gives

$$\frac{1}{L} \log \rho(F_{\text{max}} - L\epsilon) \approx \frac{\epsilon}{\Delta_0} \log \frac{e(C-1)\Delta_0}{\epsilon}. \quad (\text{S17})$$

For $C = 2$ this reduces to $(\epsilon/\Delta_0) \log(e\Delta_0/\epsilon)$.

This scaling is qualitatively different from the power-law form $\log \rho \sim \epsilon^\alpha$ derived in the main text. The difference is traceable to the gap distribution: in the Hamming-distance landscape every position has the same gap $\Delta_0 > 0$, so $p_{\text{gap}}(\Delta) = \delta(\Delta - \Delta_0)$. This is not consistent with the assumption $p_{\text{gap}}(\Delta) \sim c\Delta^\gamma$ near $\Delta = 0$ for any c and γ , so the power-law scaling does not apply.

The same conclusion follows from the tilting framework. The cumulant generating function is $\Phi(\beta) = L \log[1 + (C-1)e^{-\beta\Delta_0}]$, and the saddle-point condition gives

$$\epsilon = (C-1)\Delta_0 \frac{e^{-\beta\Delta_0}}{1 + (C-1)e^{-\beta\Delta_0}} \approx (C-1)\Delta_0 e^{-\beta\Delta_0} \quad (\text{S18})$$

for large β . The fitness deficit decreases exponentially in β , rather than as a power law of β as in the generic case (Eq. S9), confirming the breakdown of the power-law scaling mechanism.

S5 Global epistasis coefficients

Here we derive the coefficients A_F and B_F appearing in Eq. 23 of the main text. The observed fitness is $F = g(\phi)$, where g is a nonlinear function of the additive trait ϕ . We assume that g attains its maximum at ϕ_{max} and that $g'_{\text{max}} \equiv g'(\phi_{\text{max}}) \neq 0$, so that g is locally invertible near the peak. In this region the genotypic density transforms by the standard change of variables:

$$\rho_F(F) = \frac{\rho_\phi(\phi)}{g'(\phi)}, \quad \phi = g^{-1}(F). \quad (\text{S19})$$

Define per-site trait and fitness deficits $\epsilon_\phi = (\phi_{\text{max}} - \phi)/L$ and $\epsilon_F = (F_{\text{max}} - F)/L$. Taylor-expanding $g(\phi_{\text{max}} - L\epsilon_\phi)$ around ϕ_{max} gives

$$\epsilon_F = g'_{\text{max}} \epsilon_\phi + O(\epsilon_\phi^2), \quad (\text{S20})$$

so that $\epsilon_\phi \approx \epsilon_F/g'_{\max}$ when $\epsilon_\phi \ll 2g'_{\max}/|g''_{\max}|$. Taking logarithms of Eq. S19 and dividing by L :

$$\frac{1}{L} \log \rho_F = \frac{1}{L} \log \rho_\phi - \frac{1}{L} \log g'(\phi). \quad (\text{S21})$$

Near the peak, $g'(\phi) \approx g'_{\max}$, so the second term contributes $-\frac{1}{L} \log g'_{\max}$. Using Eq. 22 of the main text, $L^{-1} \log \rho_\phi \approx A_\phi + B_\phi \epsilon_\phi^\alpha$, and substituting $\epsilon_\phi \approx \epsilon_F/g'_{\max}$:

$$\frac{1}{L} \log \rho_F \approx \underbrace{\left(A_\phi - \frac{\log g'_{\max}}{L} \right)}_{A_F} + \underbrace{\frac{B_\phi}{(g'_{\max})^\alpha}}_{B_F} \epsilon_F^\alpha. \quad (\text{S22})$$

The scaling exponent α is unchanged. This approximation is valid when $\epsilon_F \ll 2(g'_{\max})^2/|g''_{\max}|$, i.e., when the nonlinearity is well-approximated by its local linearization near the peak.

References

- [1] Ville Mustonen, Justin Kinney, Curtis G. Callan, and Michael Lässig. Energy-dependent fitness: A quantitative model for the evolution of yeast transcription factor binding sites. *Proceedings of the National Academy of Sciences*, 105(34):12376–12381, 08 2008. ISSN 0027-8424. doi: 10.1073/pnas.0805909105.
- [2] Ilya E Vorontsov, Irina A Eliseeva, Arsenii Zinkevich, Mikhail Nikonov, Sergey Abramov, Alexandr Boytsov, Vasily Kamenets, Alexandra Kasianova, Semyon Kolmykov, Ivan S Yevshin, Alexander Favorov, Yulia A Medvedeva, Arttu Jolma, Fedor Kolpakov, Vsevolod J Makeev, and Ivan V Kulakovskiy. HOCOMOCO in 2024: a rebuild of the curated collection of binding models for human and mouse transcription factors. *Nucleic Acids Research*, 52(D1):D154–D163, 2024. ISSN 0305-1048. doi: 10.1093/nar/gkad1077.
- [3] Hélène Touzet and Jean-Stéphane Varré. Efficient and accurate P-value computation for Position Weight Matrices. *Algorithms for Molecular Biology*, 2:15, 2007. doi: 10.1186/1748-7188-2-15.
- [4] Otto G. Berg and Peter H. von Hippel. Selection of DNA binding sites by regulatory proteins. Statistical-mechanical theory and application to operators and promoters. *Journal of Molecular Biology*, 193:723–750, 1987.