

Discussion 1 Questions

Quick Questions

One of these

- Why can't we say probability when interpreting an observed confidence interval?
- When we say we are '95% confident' a parameter is captured in an interval. What does this mean?

One of these

- If we're trying to do method of moments estimation and the first raw moment of our distribution is 0, what can we do?
- If we're trying to do maximum likelihood estimation and our support has a relationship with our parameter, what can we do?

More Involved Questions

One of these

- We've covered the idea of estimation using method of moments. Can you talk about what we are doing with estimation and the concepts behind method of moments?
 - What are some of the good properties of MOM estimators? What are some of the drawbacks?
- We've covered the idea of estimation using maximum likelihood estimation. Can you talk about what we are doing with estimation and the concepts behind maximum likelihood?
 - What are some of the good properties of MLEs? What are some of the drawbacks?

One of these

- We discussed the idea of using sufficient statistics. What do we mean by a sufficient statistic and why do we care about them?

- What is their relationship to our two types of estimation, MOM and ML?
- We discussed properties of estimators. One property was MSE. Why do we like this criteria?
 - What is its relationship to consistency and why?

One of these

- We looked at the concept of bootstrapping. What is the idea of bootstrapping and what can we use it for?
- We found confidence intervals using pivots. Can you describe what a pivot is and the process we use to create a confidence interval from a pivot?