(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
10 November 2022 (10.11.2022)

WIPO | PCT

(10) International Publication Number
WO 2022/234524 A1

(51) International Patent Classification:
A61K 31/437 (2006.01) *C07D 471/14* (2006.01)
A61K 31/4375 (2006.01) *A61P 25/00* (2006.01)

(21) International Application Number:
PCT/IB2022/054187

(22) International Filing Date:
06 May 2022 (06.05.2022)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
63/185,326 06 May 2021 (06.05.2021) US


(71) Applicant: **KURES, INC.** [US/US]; c/o ATAI Life Sciences US, Inc., 180 Varick St, Floor 6, New York, New York 10014 (US).

(72) Inventor: **KRUEGEL, Andrew**; 766 Humboldt Street, Fl 2, Secaucus, New Jersey 07094 (US).


(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IQ, IR, IS, IT, JM, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: SALTS AND POLYMORPHS OF MITRAGYNINE AND 3-DEUTEROMITRAGYNINE




(I)



(II)

Figure 30 - XRPD overlay of 3-DM Glycolate crystal forms



(57) **Abstract:** The present disclosure provides: one or more salts or crystalline forms of 3-deuteromitragynine of Formula I; and one or more salts or crystalline forms of mitragynine of Formula II. Pharmaceutical compositions comprising salts and crystalline forms of Formulae I and II and methods of treatment are also disclosed. Methods of preparing and purifying the salts of Formulae I and II are further disclosed.

[Continued on next page]

WO 2022/234524 A1

Published:

— *with international search report (Art. 21(3))*
— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
— *in black and white; the international application as filed contained color or greyscale and is available for download from PATENTSCOPE*


SALTS AND POLYMORPHS OF MITRAGYNINE AND 3-DEUTEROMITRAGYNINE

[0001] CROSS-REFERENCE TO RELATED APPLICATIONS

[0002] The present application claims the benefit of priority of U.S. Provisional Application Serial No. 63/185,326, filed on May 6, 2021, the disclosure of which is incorporated herein by reference in their entireties for all purposes.

[0003] Background of the Invention

[0004] Mitragynine is a psychoactive compound derived from the leaves of the Southeast Asian plant *Mitragyna speciosa*, of the Rubiaceae (coffee family). Mitragynine is known to have strong analgesic effects, so it has been suggested as a useful compound in the treatment of pain and opioid addiction (as a replacement therapy).




Mitragynine

[0005] The structure of mitragynine in free base form is depicted above. Mitragynine is insoluble in water but soluble in conventional organic solvents, including acetone, acetic acid, alcohols, chloroform and diethyl ether. Mitragynine distils at 230–240 °C at 5 mmHg. It forms white, amorphous crystals that melt at 102–106 °C. The melting point of mitragynine hydrochloride is 243°C; the picrate melts at 223–224 °C and the acetate at 142 °C. (https://www.emcdda.europa.eu/publications/drug-profiles/kratom_en)


[0006] The above mentioned mitragynine hydrochloride salt is known to form a gel in aqueous solution, which makes it difficult to formulate as a solution for injectable use as described by Ellen Field in *J. Chem. Soc., Trans.,* 1921, **119**, 887- 891; XCVIII Murrayanine and Mitraversine, Two New Alkaloids from Species of Mitragyne.

[0007] Mitragynine isolated from natural sources is frequently contaminated with the closely related alkaloid corynantheidine (CR, pictured below), which is very difficult to remove by known methods.




corynantheidine (CR)

[0008] Recently, 3-deuteromitragynine (3-DM), pictured below, has been found to be a useful derivative of mitragynine as described in PCT/US2020/015898, published as WO2020160280A1, which is hereby incorporated by reference in its entirety for all purposes.



3-deuteromitragyine (3-DM)

[0009] This compound has potential therapeutic properties similar to mitragynine and since it is a deuterated derivative, its physicochemical properties are also very similar. For example, its hydrochloride salt also forms a gel in aqueous solution. Further, 3-DM may be contaminated with 3-deuterocorynantheidine (3-DCR).




3-deuterocorynantheidine (3-DCR)

[0010] There is an ongoing need for alternative salts and crystalline forms of mitragynine and 3-DM that better exclude impurities, reduce gel formation, and offer different physicochemical properties, such as pKa, solubility, and melting point.

[0011] SUMMARY OF THE INVENTION

[0012] In one aspect, the present disclosure provides a salt of 3-deuteromitragynine of Formula I:



Formula I

[0013] wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

[0014] In some embodiments, the anion is glycolate. In some embodiments, the salt of 3-deuteromitragynine of Formula I is glycolate Type A, glycolate Type B, glycolate Type C, glycolate Type D, glycolate Type E, or glycolate Type F.

[0015] In some embodiments, the salt of 3-deuteromitragynine of Formula I is glycolate Type A. In some embodiments, the glycolate Type A is characterized by peaks in an X-ray diffraction (XRPD) pattern at 7.1±0.2, 10.1±0.2, and 11.2±0.2 °2θ.

[0016] In some embodiments, the salt of 3-deuteromitragynine of Formula I is glycolate Type B. In some embodiments, the glycolate Type B is characterized by peaks in an XRPD pattern at 5.3±0.2, 5.7±0.2, and 7.5±0.2 °2θ.

[0017] In some embodiments, the salt of 3-deuteromitragynine of Formula I is glycolate Type C. In some embodiments, the glycolate Type C is characterized by peaks in an XRPD pattern at 6.0±0.2, 7.4±0.2, and 24.2±0.2 °2θ.

[0018] In some embodiments, the salt of 3-deuteromitragynine of Formula I is glycolate Type D. In some embodiments, the glycolate Type D is characterized by peaks in an XRPD pattern at 5.3±0.2, 6.8±0.2, and 9.0±0.2 °2θ.

[0019] In some embodiments, the salt of 3-deuteromitragynine of Formula I is glycolate Type E. In some embodiments, the glycolate Type E is characterized by peaks in an XRPD pattern at 5.1±0.2, 7.8±0.2, and 8.8±0.2 °2θ.

[0020] In some embodiments, the salt of 3-deuteromitragynine of Formula I is glycolate Type F. In some embodiments, the glycolate Type F is characterized by peaks in an XRPD pattern at 5.9±0.2, 6.4±0.2, and 7.2±0.2 °2θ.

[0021] In some embodiments, the salt of 3-deuteromitragynine of Formula I is L-lactate. In some embodiments, the L-lactate salt is characterized by peaks in an XRPD pattern at 6.9±0.2, 10.0±0.2, and 11.0±0.2 °2θ.


[0022] In some embodiments, the salt of 3-deuteromitragynine of Formula I is succinate. In some embodiments, the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 17.6±0.2, and 19.3±0.2 °2θ.

[0023] In some embodiments, the salt of 3-deuteromitragynine of Formula I is fumarate. In some embodiments, the fumarate salt is characterized by peaks in an XRPD pattern at 8.4±0.2, 17.5±0.2, and 19.2±0.2 °2θ.

[0024] In some embodiments, the salt of 3-deuteromitragynine of Formula I is mesylate. In some embodiments, the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 16.7±0.2, and 17.3±0.2 °2θ.

[0025] In some embodiments, the present disclosure provides pharmaceutical compositions comprising a salt of 3-deuteromitragynine of Formula I as described herein.

[0026] In one aspect, the present disclosure provides a salt of mitragynine of Formula II



Formula II

[0027] wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

[0028] In some embodiments, the salt of mitragynine of Formula II is glycolate. In some embodiments, the glycolate salt is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.2±0.2, and 11.3±0.2 °2θ.

[0029] In some embodiments, the salt of mitragynine of Formula II is L-lactate. In some embodiments, the L-lactate salt is characterized by peaks in an XRPD pattern at 7.0±0.2, 10.1±0.2, and 11.2±0.2 °2θ.

[0030] In some embodiments, the salt of mitragynine of Formula II is succinate. In some embodiments, the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 17.6±0.2, and 19.3±0.2 °2θ.

[0031] In some embodiments, the salt of mitragynine of Formula II is fumarate. In some embodiments, the fumarate salt is characterized by peaks in an XRPD pattern at 8.3±0.2, 19.1±0.2, and 19.2±0.2 °2θ.

[0032] In some embodiments, the salt of mitragynine of Formula II is mesylate. In some embodiments, the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 16.7±0.2, and 17.4±0.2 °2θ.

[0033] In some embodiments, the present disclosure provides pharmaceutical compositions comprising a salt of mitragynine of Formula II as described herein.

[0034] In one aspect, the present disclosure provides a method of treating a subject afflicted with acute pain, chronic pain, a depressive disorder, a mood disorder, an anxiety disorder, borderline personality disorder, a substance use disorder, opioid use disorder, opioid withdrawal symptoms, alcohol use disorder, or alcohol withdrawal disorder, comprising administering an effective amount of a salt of 3-deuteromitragynine of Formula I or a salt of mitragynine of Formula II as described herein.

[0035] In some embodiments, present disclosure provides a method of treating opioid use disorder. In some embodiments, present disclosure provides a method of treating opioid withdrawal.

DEFINITIONS

[0036] In each instance herein, in descriptions, embodiments, and examples of the present disclosure, the terms "comprising", "including", etc., are to be read expansively, without limitation. Thus, unless the context clearly requires otherwise, throughout the description and the claims, the words "comprise", "comprising", and the like are to be construed in an inclusive sense as opposed to an exclusive sense, that is to say in the sense of "including but not limited to".

[0037] In the present description, the articles "a" and "an" are used to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, "an element" can be taken to mean one element or more than one element.

[0038] Throughout this description, the term "about" is used to indicate that a value includes the standard deviation of error for the method being employed to determine the value, for example, dosage levels, as described in detail herein. In particular, the term "about" encompasses a 10% to 15% deviation (positive and negative) in the stated value or range, particularly 10% deviation (positive and negative) in the stated value or range.

[0039] The present disclosure is also intended to include all isotopes of atoms occurring on the salt(s) disclosed herein. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium. Isotopes of carbon include C-13 and C-14.

[0040] It will be noted that any notation of a carbon in structures throughout this application, when used without further notation, are intended to represent all isotopes of carbon, such as ^{12}C, ^{13}C, or ^{14}C. Furthermore, any salt(s) containing ^{13}C or ^{14}C may specifically have the structure of any of the salt(s) disclosed herein.

[0041] It will also be noted that any notation of a hydrogen in structures throughout this application, when used without further notation, are intended to represent all isotopes of hydrogen, such as ^{1}H, ^{2}H, or ^{3}H. Furthermore, any salt(s) containing ^{2}H or ^{3}H may specifically have the structure of any of the salt(s) disclosed herein.

[0042] Isotopically-labeled salt(s) can generally be prepared by conventional techniques known to those skilled in the art using appropriate isotopically-labeled reagents in place of the non-labeled reagents employed.

[0043] All references, including patents and patent applications, cited in this specification are hereby incorporated by reference. No admission is made that any reference constitutes prior art. Nor does discussion of any reference constitute an admission that such reference forms part of the common general knowledge in the art, in any country.

Brief Description of the Figures

[0044] **FIG. 1** shows an XRPD pattern of amorphous 3-DM free base starting material (810397-01-A).

[0045] **FIG. 2** shows TGA/mDSC curves of amorphous 3-DM free base starting material (810397-01-A).

[0046] FIG. 3 shows an ^{1}H NMR spectrum of amorphous 3-DM free base starting material (810397-01-A).

[0047] FIG. 4 shows an XRPD pattern of Fumarate Type A 3-DM Salt.

FIG. 5 shows an XRPD pattern of L-Lactate Type A 3-DM Salt.

[0048] FIG. 6 shows an XRPD pattern of Glycolate Type A 3-DM Salt.

[0049] FIG. 7 shows an XRPD pattern of Succinate Type A 3-DM Salt.

[0050] FIG. 8 shows an XRPD pattern of Mesylate Type A 3-DM Salt.

[0051] FIG. 9 shows an XRPD pattern of HCl Type A 3-DM Salt (810397-04-B1).

[0052] FIG. 10 shows an XRPD pattern of HCl Type B 3-DM Salt (810397-04-D1).

[0053] FIG. 11 XRPD pattern of Phosphate Type A 3-DM Salt (810397-04-D3).

[0054] FIG. 12 shows an XRPD pattern of Maleate Type A 3-DM Salt (810397-04-C4).

[0055] FIG. 13 shows an XRPD pattern of L-Tartrate Type A 3-DM Salt (810397-04-B5).

[0056] FIG. 14 shows an XRPD pattern of Glycolate Type B 3-DM Salt (810397-04-C8).

[0057] FIG. 15 shows an XRPD pattern of Adipate Type A 3-DM Salt (810397-04-A12).

[0058] FIG. 16 shows an XRPD pattern of Adipate Type B 3-DM Salt (810397-04-C12).

[0059] FIG. 17 shows an XRPD pattern of Acetate Type A 3-DM Salt (810397-04-D13).

[0060] FIG. 18 shows an XRPD pattern of Malonate Type A 3-DM Salt (810397-04-B17).

[0061] FIG. 19 shows an XRPD pattern of Malonate Type B 3-DM Salt (810397-04-D17)

[0062] FIG. 20 shows an XRPD pattern of Gentisate Type A 3-DM Salt (810397-04-B18).

[0063] FIG. 21 shows an XRPD pattern of Gentisate Type B 3-DM Salt (810397-04-C18).

[0064] FIG. 22 shows an XRPD pattern of Edisylate Type A 3-DM Salt (810397-04-A19).

[0065] FIG. 23 shows an XRPD pattern of Edisylate Type B 3-DM Salt (810397-04-C19).

[0066] FIG. 24 shows an XRPD pattern of Benzoate Type A 3-DM Salt (810397-04-D20).

[0067] FIG. 25 shows an XRPD pattern of Esylate Type A 3-DM Salt (810397-04-C21).

[0068] FIG. 26 shows an XRPD pattern of Besylate Type A 3-DM Salt (810397-04-D23).

[0069] FIG. 27 shows an XRPD pattern of Tosylate Type A 3-DM Salt (810397-04-B24).

[0070] FIG. 28 shows an XRPD pattern of Oxalate Type A 3-DM Salt (810397-04-A25).

[0071] FIG. 29 shows an XRPD pattern of Oxalate Type B 3-DM Salt (810397-04-B25).

[0072] FIG. 30 shows an XRPD overlay of 3-DM Glycolate crystal forms.

[0073] FIG. 31 shows a polarized light microscopy (PLM) image of 3-DM Glycolate Type A starting material used in crystallization process development experiments (**Examples 18-21**).

[0074] FIG. 32 shows an XRPD pattern of 3-DM Free Base Type A (810081-04-C1_dry).

[0075] FIG. 33 shows a polarized light microscopy (PLM) image of 3-DM Glycolate Type A prepared at 5-g scale (810082-25-B).

[0076] FIG. 34 shows an XRPD pattern of amorphous mitragynine free base starting material (810080-01-A).

[0077] FIG. 35 shows TGA/mDSC curves of amorphous mitragynine free base starting material (810080-01-A).

[0078] FIG. 36 shows an ^{1}H NMR spectrum of amorphous mitragynine free base starting material (810080-01-A).

[0079] FIG. 37 shows an XRPD pattern of crude alkaloid extract starting material (810080-01-B).

[0080] FIG. 38 shows a TGA curve of crude alkaloid extract starting material (810080-01-B).

[0081] FIG. 39 shows an XRPD pattern of Fumarate Type A Mitragynine Salt.

[0082] FIG. 40 shows an XRPD pattern of L-Lactate Type A Mitragynine Salt.

[0083] FIG. 41 shows an XRPD pattern of Glycolate Type A Mitragynine Salt.

[0084] FIG. 42 shows an XRPD pattern of Succinate Type A Mitragynine Salt.


[0085] FIG. 43 shows an XRPD pattern of Mesylate Type A Mitragynine Salt.

DETAILED DESCRIPTION

[0086] The following description sets forth numerous exemplary configurations, parameters, and the like. It should be recognized, however, that such description is not intended as a limitation on the scope of the present disclosure but is instead provided as a description of exemplary embodiments.

Salts of 3-deuteromitragynine

[0087] In one aspect the present disclosure provides one or more salts of 3-deuteromitragynine of Formula I:



Formula I

[0088] wherein the anion is glycolate, L-lactate, succinate, fumarate or mesylate.

[0089] In some embodiments, the anion is glycolate (i.e., the salt is a glycolate salt).

[0090] In some embodiments, the glycolate salt of 3-deuteromitragynine exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately:

Pos. [°2θ]
7.06
10.12
11.25
15.97
17.99
18.11
19.49
19.69
20.34
20.88
22.58

Pos. [°2θ]
25.16
27.60

±0.2 degrees 2 theta.

[0091] In some embodiments, the salt of 3-deuteromitragynine of Formula I is glycolate Type A, glycolate Type B, glycolate Type C, glycolate Type D, glycolate Type E, glycolate Type F, or combinations thereof.

[0092] In some embodiments, the salt of 3-DM is glycolate Type A.

[0093] In some embodiments, the glycolate Type A is characterized by peaks in an X-ray diffraction (XRPD) pattern at 7.1±0.2, 10.1±0.2, and 11.2±0.2 °2θ. In some embodiments, the glycolate Type A is further characterized by at least one XRPD peak selected from 16.0±0.2, 18.0±0.2, 19.5±0.2, 20.9±0.2, 22.6±0.2, and 25.2±0.2 °2θ.

[0094] In some embodiments, the glycolate Type A is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.1±0.2, 13.2±0.2, 16.0±0.2, 18.0±0.2, 19.5±0.2, 20.9±0.2, 22.6±0.2, and 25.2±0.2 °2θ. In some embodiments, the glycolate Type A is further characterized by at least one XRPD peak selected from 13.2±0.2, 14.1±0.2, 15.0±0.2, 18.5±0.2, 19.2±0.2, 19.7±0.2, 20.3±0.2, 23.7±0.2, 240±0.2, 27.6±0.2, 29.5±0.2, 30.1±0.2, 31.6±0.2, and 34.1±0.2 °2θ.

[0095] In some embodiments, the glycolate Type A is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.1±0.2, and 11.2±0.2 °2θ and at least one XRPD peak selected from 16.0±0.2, 18.0±0.2, 19.5±0.2, 19.7±0.2, 20.3±0.2, 20.9±0.2, 22.6±0.2, 25.2±0.2 and 27.6±0.2 °2θ.

[0096] In some embodiments, the glycolate Type A is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.1±0.2, 11.2±0.2, 13.2±0.2, 14.1±0.2, 15.1±0.2, 16.0±0.2, 18.0±0.2, 18.5±0.2, 19.2±0.2, 19.5±0.2, 19.7±0.2, 20.3±0.2, 20.9±0.2, 22.6±0.2, 23.7±0.2, 24.0±0.2, 25.2±0.2, 27.6±0.2, 29.5±0.2, 30.2±0.2, 31.6±0.2, and 34.1±0.2 °2θ.

[0097] In some embodiments, the glycolate Type A exhibits a weight loss of about 1% up to a temperature of about 150 °C as measured by thermogravimetric (TGA) analysis.

[0098] In some embodiments, the glycolate Type A exhibits a Differential Scanning Calorimetry (DSC) thermogram comprising an endotherm peak at about 222±2.5 °C.

[0099]

[0100] In some embodiments, a glycolate Type A salt of 3-DM is provided exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at about:

Pos. [°2θ]
7.06
10.11
11.24
15.96
18.01
19.49
19.70
20.34
20.88
22.57
25.15
27.62

±0.2 degrees 2 theta.

[0101] In some embodiments, the salt of 3-DM is glycolate Type B.

[0102] In some embodiments, the glycolate Type B is characterized by peaks in an XRPD pattern at 5.3±0.2, 5.7±0.2, and 7.5±0.2 °2θ. In some embodiments, the glycolate Type B is further characterized by at least one XRPD peak selected from 6.8±0.2, 10.8±0.2, 13.7±0.2, 19.9±0.2, 22.7±0.2, and 27.4±0.2 °2θ.

[0103] In some embodiments, the glycolate Type B is characterized by peaks in an XRPD pattern at 5.3±0.2, 5.7±0.2, 6.8±0.2, 7.5±0.2, 10.8±0.2, 13.7±0.2, 19.9±0.2, 22.7±0.2, and 27.4±0.2 °2θ. In some embodiments, the glycolate Type B is further characterized by at least one XRPD peak selected from 9.0±0.2, 14.7±0.2, 17.4±0.2, 21.2±0.2, 24.1±0.2, and 25.4±0.2 °2θ.

[0104] In some embodiments, the glycolate Type B is characterized by peaks in an XRPD pattern at 5.3±0.2, 5.7±0.2, and 7.5±0.2 °2θ and at least one XRPD peak selected from 10.8±0.2, 13.7±0.2, 19.9±0.2, 21.2±0.2, 22.7±0.2, and 24.0±0.2 °2θ.

[0105] In some embodiments, the glycolate Type B is characterized by peaks in an XRPD pattern at 5.3±0.2, 5.7±0.2, 6.8±0.2, 7.5±0.2, 9.0±0.2, 10.8±0.2, 13.7±0.2, 14.7±0.2, 17.4±0.2, 19.9±0.2, 21.2±0.2, 22.7±0.2, 24.0±0.2, 25.4±0.2, and 27.4±0.2 °2θ.

[0106] In some embodiments, the glycolate Type B exhibits a weight loss of about 4% up to a temperature of about 120 °C as measured by TGA analysis. In some embodiments, the glycolate Type B further exhibits a weight loss of about 8% between a temperature ranging from about 120 °C to about 160 °C as measured by TGA analysis.

[0107] In some embodiments, the glycolate Type B exhibits a DSC thermogram comprising an endothermic peak at about 147±2.5 °C and 223±2.5 °C.

[0108] In some embodiments, a glycolate Type B salt of 3-DM is provided exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at about:

Pos. [°2θ]
5.29
5.68
7.45
10.78
13.65
19.86
21.22
22.72
24.05

±0.2 degrees 2 theta.

[0109] In some embodiments, the salt of 3-DM is glycolate Type C.

[0110] In some embodiments, the glycolate Type C is characterized by peaks in an XRPD pattern at 6.0±0.2, 7.4±0.2, and 24.2±0.2 °2θ. In some embodiments, the glycolate Type C is further characterized by at least one XRPD peak selected from 14.2±0.2, 16.3±0.2, 18.1±0.2, 20.1±0.2, 26.2±0.2, and 27.6±0.2 °2θ.

[0111] In some embodiments, the glycolate Type C is characterized by peaks in an XRPD pattern at 6.0±0.2, 7.4±0.2, 14.2±0.2, 16.3±0.2, 18.1±0.2, 20.1±0.2, 24.2±0.2, 26.2±0.2, and 27.6±0.2 °2θ. In some embodiments, the glycolate Type C is further characterized by at least one XRPD peak selected from 10.6±0.2, 11.0±0.2, 11.4±0.2, 12.8±0.2, 13.7±0.2, 15.7±0.2, 18.8±0.2, 21.2±0.2, 21.5±0.2, 22.6±0.2, 22.9±0.2, 25.1±0.2, 28.6±0.2, 29.4±0.2, 31.6±0.2, 33.7±0.2, 35.2±0.2, and 38.3±0.2 °2θ.

[0112] In some embodiments, the glycolate Type C is characterized by peaks in an XRPD pattern at 6.0±0.2, 7.4±0.2, and 24.2±0.2 °2θ and at least one XRPD peak at

11.0±0.2, 11.4±0.2, 13.7±0.2, 14.2±0.2, 16.3±0.2, 18.1±0.2, 18.8±0.2, 20.1±0.2, 21.5±0.2, 22.9±0.2, and 26.2±0.2 °2θ.

[0113] In some embodiments, the glycolate Type C is characterized by peaks in an XRPD pattern at 6.0±0.2, 7.4±0.2, 10.6±0.2, 11.0±0.2, 11.4±0.2, 12.8±0.2, 13.7±0.2, 14.2±0.2, 15.7±0.2, 16.3±0.2, 17.6±0.2, 18.1±0.2, 18.8±0.2, 20.1±0.2, 21.2±0.2, 21.5±0.2, 22.6±0.2, 22.9±0.2, 24.2±0.2, 25.2±0.2, 26.2±0.2, 27.6±0.2, 28.6±0.2, 29.4±0.2, 31.6±0.2, 33.7±0.2, 35.3±0.2, and 38.3±0.2 °2θ.

[0114] In some embodiments, the glycolate Type C exhibits a weight less of about 6% up to a temperature of about 150 °C as measured by TGA.

[0115] In some embodiments, the glycolate Type C exhibits a DSC thermogram comprising an endotherm peak at about 61±2.5 °C, 141±2.5 °C, and about 222±2.5 °C.

[0116] In some embodiments, a glycolate Type C salt of 3-DM is provided exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at about:

Pos. [°2θ]
6.03
7.43
10.99
11.39
13.65
14.22
16.30
18.07
18.84
20.08
21.54
22.87
24.16
26.22

±0.2 degrees 2 theta.

[0117] In some embodiments, the salt of 3-DM is glycolate Type D.

[0118] The salt of embodiment 32, wherein the glycolate Type D is characterized by peaks in an XRPD pattern at 5.3±0.2, 6.8±0.2, and 9.0±0.2 °2θ. In some

embodiments, the glycolate Type D is further characterized by at least one XRPD peak selected from 11.0±0.2, 13.5±0.2, 17.3±0.2, 19.5±0.2, 20.1±0.2, and 21.3±0.2 °2θ.

[0119] In some embodiments, the glycolate Type D is characterized by peaks in an XRPD pattern at 5.3±0.2, 6.8±0.2, 9.0±0.2, 11.0±0.2, 13.5±0.2, 17.3±0.2, 19.5±0.2, 20.1±0.2, and 21.3±0.2 °2θ. In some embodiments, the glycolate Type D is further characterized by at least one XRPD peak selected from 10.1±0.2, 11.3±0.2, 13.9±0.2, 16.5±0.2, 17.0±0.2, 18.0±0.2, 22.9±0.2, 23.7±0.2, 25.5±0.2, and 27.3±0.2 °2θ.

[0120] In some embodiments, the glycolate Type D is characterized by peaks in an XRPD pattern at 5.3±0.2, 6.8±0.2, and 9.0±0.2 °2θ and at least one XRPD peak selected from 11.0±0.2, 11.3±0.2, 13.5±0.2, 13.9±0.2, 17.3±0.2, 19.5±0.2, 20.1±0.2, 21.3±0.2, and 23.7±0.2 °2θ.

[0121] In some embodiments, the glycolate Type D is characterized peaks in an XRPD pattern at 5.3±0.2, 6.8±0.2, 9.0±0.2, 10.1±0.2, 11.0±0.2, 11.3±0.2, 13.5±0.2, 13.9±0.2, 16.5±0.2, 17.0±0.2, 17.3±0.2, 18.0±0.2, 19.5±0.2, 20.1±0.2, 21.3±0.2, 22.9±0.2, 23.7±0.2, 25.5±0.2, and 27.3±0.2 °2θ.

[0122] In some embodiments, the glycolate Type D exhibits a weight less of about 3% up to a temperature of about 100 °C as measured by TGA.

[0123] In some embodiments, the glycolate Type D exhibits a DSC thermogram comprising an endotherm peak at about 63±2.5 °C, about 210±2.5 °C, and about 123±2.5 °C.

[0124] In some embodiments, a glycolate Type D salt of 3-DM is provided exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at about:

Pos. [°2θ]
5.33
6.82
9.04
11.02
11.33
13.53
13.85
17.28
19.50
20.08
21.28

Pos. [°2θ]
23.71

±0.2 degrees 2 theta.

[0125] In some embodiments, the salt of 3-DM is glycolate Type E.

[0126] In some embodiments, the glycolate Type E is characterized by peaks in an XRPD pattern at 5.1±0.2, 7.8±0.2, and 8.8±0.2 °2θ. In some embodiments, the glycolate Type E is further characterized by at least one XRPD peak selected from 11.0±0.2, 12.0±0.2, 15.1±0.2, 16.8±0.2, 19.1±0.2, and 21.0±0.2 °2θ.

[0127] In some embodiments, the glycolate Type E is characterized by peaks in an XRPD pattern at 5.1±0.2, 7.8±0.2, 8.8±0.2, 11.0±0.2, 12.0±0.2, 15.1±0.2, 16.8±0.2, 19.1±0.2, and 21.0±0.2 °2θ. In some embodiments, the glycolate Type E is further characterized by at least one XRPD peak selected from 18.4±0.2 and 22.3±0.2 °2θ.

[0128] In some embodiments, the glycolate Type E is characterized by peaks in an XRPD pattern at 5.1±0.2, 7.8±0.2, and 8.8±0.2 °2θ and at least one peak selected from 11.0±0.2, 12.0±0.2, 16.8±0.2, 18.4±0.2, 19.1±0.2, 20.9±0.2, and 22.3±0.2 °2θ.

[0129] In some embodiments, the glycolate Type E is characterized by peaks in an XRPD pattern at 5.1±0.2, 7.8±0.2, 8.8±0.2, 11.0±0.2, 12.0±0.2, 15.1±0.2, 16.8±0.2, 18.4±0.2, 19.1±0.2, 20.9±0.2, and 22.3±0.2 °2θ.

[0130] In some embodiments, a glycolate Type E salt of 3-DM is provided exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at about:

Pos. [°2θ]
5.10
7.81
8.82
10.97
12.02
16.79
18.40
19.08
20.91
22.25

±0.2 degrees 2 theta.

[0131] In some embodiments, the salt of 3-DM is glycolate Type F.

[0132] In some embodiments, the glycolate Type F is characterized by peaks in an XRPD pattern at 5.9±0.2, 6.4±0.2, and 7.2±0.2 °2θ. In some embodiments, the glycolate Type F is further characterized by at least one XRPD peak selected from 13.4±0.2, 14.1±0.2, 18.4±0.2, 19.8±0.2, 24.7±0.2, and 23.9 ±0.2 °2θ.

[0133] In some embodiments, the glycolate Type F is characterized by peaks in an XRPD pattern at 5.9±0.2, 6.4±0.2, 7.2±0.2, 13.4±0.2, 14.1±0.2, 18.4±0.2, 19.8±0.2, 24.7±0.2, and 23.9±0.2 °2θ. In some embodiments, the glycolate Type F is further characterized by at least one XRPD peak selected from 10.3±0.2, 10.8±0.2, 11.2±0.2, 12.3±0.2, 16.1±0.2, 16.8±0.2, 17.2±0.2, 17.8±0.2, 20.6±0.2, 21.4±0.2, 22.1±0.2, 24.4±0.2, 26.0±0.2, 26.4±0.2, and 27.2±0.2 °2θ.

[0134] In some embodiments, the glycolate Type F is characterized by peaks in an XRPD pattern at 5.9±0.2, 6.4±0.2, and 7.2±0.2 °2θ and at least one XRPD peak selected from 10.8±0.2, 11.2±0.2, 13.4±0.2, 14.1±0.2, 16.11±0.2, 17.8±0.2, 18.4±0.2, 19.8±0.2, 21.4±0.2, 22.1±0.2, 23.7±0.2, 23.9±0.2 and 24.4±0.2 °2θ.

[0135] In some embodiments, the glycolate Type F is characterized by peaks in an XRPD pattern at 5.9±0.2, 6.4±0.2, 7.2±0.2, 10.3±0.2, 10.8±0.2, 11.2±0.2, 12.3±0.2, 13.4±0.2, 14.1±0.2, 16.1±0.2, 16.8±0.2, 17.2±0.2, 17.8±0.2, 18.4±0.2, 19.8±0.2, 20.6±0.2, 21.4±0.2, 22.1±0.2, 23.7±0.2, 23.9±0.2, 24.4±0.2, 26.0±0.2, 26.4±0.2, and 27.2±0.2 °2θ.

[0136] In some embodiments, a glycolate Type F salt of 3-DM is provided exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at about:

Pos. [°2θ]
5.86
6.39
7.21
10.83
11.22
13.41
14.05
16.13
17.76
18.42
19.81
21.36

Pos. [°2θ]
22.07
23.68
23.87
24.38

±0.2 degrees 2 theta.

[0137] In some embodiments, the anion is L-lactate (i.e., the salt is a L-lactate salt).

[0138] In some embodiments, the L-lactate salt is characterized by peaks in an XRPD pattern at 6.9±0.2, 10.0±0.2, and 11.0±0.2 °2θ. In some embodiments, the L-lactate salt is further characterized by at least one XRPD peak selected from 15.7±0.2, 20.6±0.2, 22.3±0.2, and 24.8±0.2 °2θ.

[0139] In some embodiments, the L-lactate salt is characterized by peaks in an XRPD pattern at 6.9±0.2, 10.0±0.2, 11.0±0.2, 15.7±0.2, 20.6±0.2, 22.3±0.2, and 24.8±0.2 °2θ. In some embodiments, the L-lactate salt is further characterized by at least one XRPD peak selected from 10.7±0.2, 13.0±0.2, 13.8±0.2, 17.7±0.2, 18.1±0.2, 18.8±0.2, 19.3±0.2, 19.8±0.2, 23.6±0.2, 24.4±0.2, 27.0±0.2, 28.0±0.2, 29.3±0.2, 31.2±0.2, 33.8±0.2, 35.6±0.2°2θ.

[0140] In some embodiments, the L-lactate salt is characterized by peaks in an XRPD pattern at 6.9±0.2, 10.0±0.2, 1.07±0.2, 11.0±0.2, 13.0±0.2, 13.8±0.2, 15.8±0.2, 17.7±0.2, 18.1±0.2, 18.8±0.2, 19.3±0.2, 19.8±0.2, 20.6±0.2, 22.3±0.2, 23.6±0.2, 24.4±0.2, 24.8±0.2, 27.0±0.2, 27.1±0.2, and 35.6±0.2 °2θ.

[0141] In some embodiments, the L-lactate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 5**.

[0142] In some embodiments, the L-lactate salt exhibits a weight less of about 1% up to a temperature of about 150 °C as measured by TGA.

[0143] In some embodiments, the L-lactate salt exhibits a DSC thermogram comprising an endotherm peak at about 218±2.5 °C.

[0144] In some embodiments, the anion is L-lactate and the 3-deuteromitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately:

Pos. [°2θ]
6.86
9.96
10.73

Pos. [°2θ]
11.05
15.74
19.77
20.63
22.26
24.75

±0.2 degrees 2 theta.

[0145] In some embodiments, the anion is succinate (i.e., the salt is a succinate salt).

[0146] In some embodiments, the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 17.6±0.2, and 19.3±0.2 °2θ. In some embodiments, the succinate salt is further characterized by at least one XRPD peak selected 9.6±0.2, 21.7±0.2, 23.1±0.2, 25.5±0.2, and 25.9±0.2 °2θ.

[0147] In some embodiments, the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 9.6±0.2, 17.6±0.2, 19.3±0.2, 21.7±0.2, 23.1±0.2, 25.5±0.2, and 25.9±0.2 °2θ. In some embodiments, the succinate salt is further characterized by at least one XRPD peak selected from 6.2±0.2, 10.1±0.2, 14.4±0.2, 15.7±0.2, 16.1±0.2, 16.9±0.2, 18.7±0.2, 21.2±0.2, 22.3±0.2, 23.7±0.2, 24.5±0.2, 25.0±0.2, 25.3±0.2, 27.0±0.2, 29.1±0.2, 30.5±0.2, 33.0±0.2, and 34.3±0.2 °2θ.

[0148] In some embodiments, the succinate salt is characterized by peaks in an XRPD pattern at 6.2±0.2, 8.5±0.2, 9.6±0.2, 10.0±0.2, 14.4±0.2, 15.7±0.2, 16.1±0.2, 16.9±0.2, 17.6±0.2, 18.7±0.2, 19.3±0.2, 21.2±0.2, 21.7±0.2, 22.3±0.2, 23.1±0.2, 23.7±0.2, 24.5±0.2, 25.0±0.2, 25.3±0.2, 25.5±0.2, 25.9±0.2, 27.0±0.2, 29.1±0.2, 30.5±0.2, 31.3±0.2, 33.0±0.2, and 34.0±0.2 °2θ.

[0149] In some embodiments, the succinate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 7**.

[0150] In some embodiments, the succinate salt exhibits a weight less of about 2 % up to a temperature of about 150 °C as measured by TGA.

[0151] In some embodiments, the succinate salt exhibits a DSC thermogram comprising an endothermic peak at about 198±2.5 °C and about 202±2.5 °C.

[0152] In some embodiments, the anion is succinate and the 3-deuteromitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately:

Pos. [°2θ]
8.46
10.06
17.56
19.27
21.71
23.13
25.53
25.94
31.27

±0.2 degrees 2 theta.

[0153] In some embodiments, the anion is fumarate (i.e., the salt is a fumarate salt).

[0154] The salt of embodiment 73, wherein the fumarate salt is characterized by peaks in an XRPD pattern at 8.4±0.2, 17.5±0.2, and 19.2±0.2 °2θ. In some embodiments, the fumarate salt is further characterized by at least one XRPD peak selected from 9.6±0.2, 21.6±0.2, 25.4±0.2, 25.8±0.2, and 31.1±0.2 °2θ.

[0155] In some embodiments, the fumarate salt is characterized by peaks in an XRPD pattern at 8.4±0.2, 9.6±0.2, 17.5±0.2, 19.2±0.2, 21.6±0.2, 25.4±0.2, 25.8±0.2, and 31.1±0.2 °2θ. In some embodiments, the fumarate salt is further characterized by at least one XRPD peak selected from 13.4±0.2, 14.4±0.2, 15.6±0.2, 16.2±0.2, 16.9±0.2, 18.7±0.2, 22.4±0.2, 23.0±0.2, 23.4±0.2, 23.8±0.2, 27.0±0.2, 28.9±0.2, 32.8±0.2, 34.0±0.2, and 38.0±0.2 °2θ.

[0156] In some embodiments, the fumarate salt is characterized by peaks in an XRPD pattern at 8.4±0.2, 9.6±0.2, 13.4±0.2, 14.4±0.2, 15.6±0.2, 16.2±0.2, 16.9±0.2, 17.5±0.2, 18.7±0.2, 19.2±0.2, 21.6±0.2, 23.4±0.2, 23.0±0.2, 23.4±0.2, 23.8±0.2, 25.4±0.2, 25.8±0.2, 27.0±0.2, 28.9±0.2, 31.1±0.2, 32.8±0.2, 34.0±0.2, and 38.0±0.2 °2θ.

[0157] In some embodiments, the fumarate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 4**.

[0158] In some embodiments, the fumarate salt exhibits a weight less of about 1% up to a temperature of about 150 °C as measured by TGA.

[0159] In some embodiments, the fumarate salt exhibits a DSC thermogram comprising an endothermic peak at about 255±2.5 °C.

[0160] In some embodiments, the anion is fumarate and the 3-deuteromitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately:

Pos. [°2θ]
8.41
9.58
17.46
19.20
23.03
25.36
25.77
31.11

±0.2 degrees 2 theta.

[0161] In some embodiments, the anion is mesylate (i.e., the salt is a mesylate salt).

[0162] In some embodiments, the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 16.7±0.2, and 17.3±0.2 °2θ. In some embodiments, the mesylate salt is further characterized by at least one XRPD peak selected from 11.6±0.2, 13.3±0.2, 18.6±0.2, 18.9±0.2, and 20.0±0.2 °2θ.

[0163] In some embodiments, the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 11.6±0.2, 13.3±0.2, 16.7±0.2, 17.3±0.2, 18.6±0.2, 18.9±0.2, and 20.0±0.2 °2θ. In some embodiments, the mesylate salt further characterized by at least one XRPD peak selected from 8.2±0.2, 10.0±0.2, 14.9±0.2, 15.3±0.2, 19.8±0.2, 21.1±0.2, 21.4±0.2, 22.2±0.2, 22.7±0.2, 23.7±0.2, 24.4±0.2, 25.1±0.2, 26.0±0.2, 26.9±0.2, 28.5±0.2, and 32.8±0.2 °2θ.

[0164] In some embodiments, the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 8.2±0.2, 10.0±0.2, 11.6±0.2, 13.3±0.2, 14.9±0.2, 15.3±0.2, 16.7±0.2, 17.3±0.2, 18.6±0.2, 18.9±0.2, 19.8±0.2, 20.0±0.2, 21.1±0.2, 21.4±0.2, 22.2±0.2, 22.7±0.2, 23.7±0.2, 24.4±0.2, 25.1±0.2, 26.0±0.2, 26.9±0.2, 28.5±0.2, and 32.8±0.2 °2θ.

[0165] In some embodiments, the mesylate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 8**.

[0166] In some embodiments, the mesylate salt exhibits a weight less of about 1% up to a temperature of about 150 °C as measured by TGA.

[0167] In some embodiments, the mesylate salt exhibits a DSC thermogram comprising an endothermic peak at about 266±2.5 °C.


[0168] In some embodiments, the anion is mesylate and the 3-deuteromitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately:

Pos. [°2θ]
6.67
11.56
13.34
14.93
16.69
17.34
18.60
18.89
20.03
22.21
22.72
25.97

±0.2 degrees 2 theta.

Salts of mitragynine

[0169] In one aspect the present disclosure provides one or more salts of mitragynine of Formula II



Formula II

[0170] wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

[0171] In some embodiments, the anion is glycolate (i.e., the salt is a glycolate salt).

[0172] The salt of embodiment 92, wherein the glycolate salt is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.2±0.2, and 11.3±0.2 °2θ. In some embodiments,

the glycolate salt is further characterized by at least one XRPD peak selected from 16.0±0.2, 18.0±0.2, 19.5±0.2, 20.9±0.2, 22.6±0.2, and 25.2±0.2 °2θ.

[0173] In some embodiments, the glycolate salt is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.2±0.2, 11.3±0.2, 16.0±0.2, 18.0±0.2, 19.5±0.2, 20.9±0.2, 22.6±0.2, and 25.2±0.2 °2θ. In some embodiments, the glycolate salt is further characterized by at least one XRPD peak selected from 13.2±0.2, 14.1±0.2, 15.1±0.2, 15.7±0.2, 18.5±0.2, 18.9±0.2, 19.2±0.2, 19.7±0.2, 20.4±0.2, 23.3±0.2, 23.5±0.2, 23.4±0.2, 24.0±0.2, 24.9±0.2, 25.9±0.2, 27.6±0.2, 28.3±0.2, 29.1±0.2, 29.6±0.2, 30.2±0.2, 30.6±0.2, 32.7±0.2, 32.2±0.2, 34.3±0.2, 35.2±0.2, 36.0±0.2, and 36.6±0.2 °2θ.

[0174] In some embodiments, the glycolate salt is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.2±0.2, 11.3±0.2, 13.2±0.2, 14.1±0.2, 15.1±0.2, 15.7±0.2, 16.0±0.2, 18.0±0.2, 18.5±0.2, 18.9±0.2 19.2±0.2, 19.7±0.2, 20.4±0.2, 23.3±0.2, 23.5±0.2, 23.8±0.2, 24.1±0.2, 24.9±0.2, 25.9±0.2, 27.6±0.2, 28.3±0.2, 29.1±0.2, 29.6±0.2, 30.2±0.2, 30.6±0.2, 31.7±0.2, 32.2±0.2, 34.2±0.2, 35.2±0.2, 36.0±0.2, and 36.6±0.2 °2θ.

[0175] In some embodiments, the glycolate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 41**.

[0176] In some embodiments, the glycolate salt exhibits a weight less of about 2% up to a temperature of about 150 °C as measured by TGA.

[0177] T In some embodiments, the glycolate salt exhibits a DSC thermogram comprising an endothermic peak at about 220±2.5 °C.

[0178] In some embodiments, the anion is glycolate and the mitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at about:

Pos. [°2θ]
7.09
10.16
11.29
13.23
14.14
15.74
16.01
18.03
19.54

Pos. [°2θ]
19.72
20.36
20.93
22.62
25.20
27.63

±0.2 degrees 2 theta.

[0179] In some embodiments, the anion is L-lactate (i.e., the salt is a L-lactate salt).

[0180] In some embodiments, the L-lactate salt is characterized by peaks in an XRPD pattern at 7.0±0.2, 10.1±0.2, and 11.2±0.2 °2θ. In some embodiments, the L-lactate salt is further characterized by at least one XRPD peak selected from 15.9±0.2, 17.9±0.2, 20.8±0.2, 22.4±0.2, and 24.9±0.2 °2θ.

[0181] In some embodiments, the L-lactate salt is characterized by peaks in an XRPD pattern at 7.0±0.2, 10.1±0.2, 11.2±0.2, 15.9±0.2, 17.9±0.2, 20.8±0.2, 22.4±0.2, and 24.9 ±0.2 °2θ. In some embodiments, the L-lactate salt is further characterized by at least one XRPD peak selected from 10.9±0.2, 13.2±0.2, 13.9±0.2, 15.1±0.2, 15.6±0.2, 18.3±0.2, 19.0±0.2, 19.9±0.2, 21.2±0.2, 21.8±0.2, 22.9±0.2, 23.4±0.2, 23.8±0.2, 24.5±0.2, 25.8±0.2, 27.1±0.2, 27.3±0.2, 28.2±0.2, 29.5±0.2, 30.7±0.2, 31.4±0.2, 34.0±0.2, 35.7±0.2, 37.4±0.2, and 38.1±0.2 °2θ.

[0182] In some embodiments, the L-lactate salt is characterized by peaks in an XRPD pattern at 7.0±0.2, 10.1±0.2, 10.9±0.2, 11.2±0.2, 13.2±0.2, 13.9±0.2, 15.1±0.2, 15.6±0.2, 15.9±0.2, 17.9±0.2, 18.3±0.2, 19.0±0.2, 19.9±0.2, 20.8±0.2, 21.2±0.2, 21.8±0.2, 22.4±0.2, 22.9±0.2, 23.4±0.2, 23.8±0.2, 24.5±0.2, 24.9±0.2, 25.8±0.2, 27.1±0.2, 27.3±0.2, 28.2±0.2, 29.5±0.2, 30.7±0.2, 31.4±0.2, 34.0±0.2, 35.7±0.2, 37.4±0.2, and 38.1±0.2 °2θ.

[0183] In some embodiments, the L-lactate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 40**.

[0184] In some embodiments, the L-lactate salt exhibits a weight less of about 3% up to a temperature of about 150 °C as measured by TGA.

[0185] In some embodiments, the L-lactate salt exhibits a DSC thermogram comprising an endothermic peak at about 226±2.5 °C.

[0186] In some embodiments, the anion is L-Lactate and the mitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at about:

Pos. [°2θ]
7.01
10.12
11.20
15.57
15.90
17.88
19.44
19.91
20.80
22.42
24.90

±0.2 degrees 2 theta.

[0187] In some embodiments, the anion is succinate (i.e., the salt is a succinate salt).

[0188] In some embodiments, the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 17.6±0.2, and 19.3±0.2 °2θ. In some embodiments, the succinate salt is further characterized by at least one XRPD peak selected from 9.6±0.2, 14.4±0.2, 21.7±0.2, 23.1±0.2, 25.5±0.2, and 25.9±0.2 °2θ.

[0189] In some embodiments, the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 9.6±0.2, 14.4±0.2, 17.6±0.2, 21.7±0.2, 23.1±0.2, 25.5±0.2, and 25.9±0.2 °2θ. In some embodiments, the succinate salt is further characterized by at least one XRPD peak selected from 6.2±0.2, 9.1±0.2, 10.1±0.2, 13.5±0.2, 15.7±0.2, 16.1±0.2, 17.0±0.2, 18.7±0.2, 21.2±0.2, 22.3±0.2, 23.7±0.2, 24.5±0.2, 25.0±0.2, 25.3±0.2, 27.0±0.2, 28.6±0.2, 29.1±0.2, 30.5±0.2, 31.6±0.2, 33.0±0.2, 34.3±0.2, 34.6±0.2, 37.0±0.2, and 39.1±0.2 °2θ.

[0190] In some embodiments, the succinate salt is characterized by peaks in an XRPD pattern at 6.2±0.2, 8.5±0.2, 9.1±0.2, 9.6±0.2, 10.1±0.2, 13.5±0.2, 14.4±0.2, 15.7±0.2, 16.1±0.2, 17.0±0.2, 17.6±0.2, 18.7±0.2, 21.2±0.2, 21.7±0.2, 22.3±0.2, 23.1±0.2, 23.7±0.2, 24.5±0.2, 25.0±0.2, 25.3±0.2, 25.5±0.2, 25.9±0.2, 27.0±0.2, 28.6±0.2, 29.1±0.2, 30.5±0.2, 31.6±0.2, 33.0±0.2, 34.3±0.2, 34.6±0.2, 37.0±0.2, and 39.1±0.2 °2θ.

[0191] In some embodiments, the succinate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 42**.

[0192] In some embodiments, the succinate salt exhibits a weight less of about 4% up to a temperature of about 150 °C as measured by TGA.

[0193] In some embodiments, the succinate salt exhibits a DSC thermogram comprising an endothermic peak at about 198±2.5 °C and about 202±2.5 °C.

[0194] In some embodiments, the anion is succinate and the mitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at about:

Pos. [°2θ]
8.47
9.62
14.36
19.27
21.71
23.13
25.52
25.93
31.27

±0.2 degrees 2 theta.

[0195] In some embodiments, the anion is fumarate (i.e., the salt is a fumarate salt).

[0196] In some embodiments, the fumarate salt is characterized by peaks in an XRPD pattern at 8.3±0.2, 19.1±0.2, and 19.2±0.2 °2θ.

In some embodiments, the fumarate salt is further characterized by at least one XRPD peak selected from 14.3±0.2, 17.3±0.2, 18.6±0.2, 25.2±0.2, and 25.6±0.2 °2θ.

[0197] In some embodiments, the fumarate salt is characterized by peaks in an XRPD pattern at 8.3±0.2, 14.3±0.2, 17.3±0.2, 18.6±0.2, 19.1±0.2, and 19.2±0.2, 25.2±0.2, and 25.6±0.2 °2θ. In some embodiments, the fumarate salt is further characterized by at least one XRPD peak selected from 9.5±0.2, 15.1±0.2, 15.5±0.2, 16.0±0.2, 16.7±0.2, 19.9±0.2, 21.2±0.2, 21.4±0.2, 22.2±0.2, 22.9±0.2, 23.2±0.2, 23.6±0.2, 24.5±0.2, 26.8±0.2, 28.8±0.2, 31.0±0.2, and 34.1±0.2 °2θ.

[0198] In some embodiments, the fumarate salt is characterized by peaks in an XRPD pattern at 8.3±0.2, 9.5±0.2, 14.3±0.2, 15.1±0.2, 15.5±0.2, 16.0±0.2, 16.7±0.2, 17.3±0.2, 18.6±0.2, 19.1±0.2, 19.2±0.2, 19.9±0.2, 21.2±0.2, 21.4±0.2, 22.2±0.2, 22.9±0.2, 23.2±0.2, 23.6±0.2, 24.5±0.2, 25.3±0.2, 25.6±0.2, 26.8±0.2, 28.8±0.2, 31.0±0.2, and 34.1±0.2 °2θ.

[0199] In some embodiments, the fumarate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 39**.

[0200] In some embodiments, the fumarate salt exhibits a weight less of about 3% up to a temperature of about 150 °C as measured by TGA.

[0201] In some embodiments, the fumarate salt exhibits a DSC thermogram comprising an endothermic peak at about 226±2.5 °C.

[0202] In some embodiments, the anion is fumarate and the mitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at about;

Pos. [°2θ]
8.30
14.29
16.03
16.73
17.33
18.55
19.06
19.22
21.19
21.44
22.89
23.61
25.20
25.62
28.80

±0.2 degrees 2 theta.

[0203] In some embodiments, the anion is mesylate (i.e., the salt is a mesylate salt).

[0204] In some embodiments, the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 16.7±0.2, and 17.4±0.2 °2θ. The salt of embodiment 129, wherein the mesylate salt is further characterized by at least one XRPD peak selected from 11.6±0.2, 18.6±0.2, 18.9±0.2, 20.1±0.2, and 26.0±0.2 °2θ.

[0205] In some embodiments, the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 11.6±0.2, 16.7±0.2, 17.4±0.2, 18.6±0.2, 18.9±0.2, 20.1±0.2, and 26.0±0.2 °2θ. In some embodiments, the mesylate salt is further characterized by at least one XRPD peak selected from 8.2±0.2, 10.0±0.2, 13.0±0.2, 13.4±0.2, 15.3±0.2,

16.4±0.2, 18.3±0.2, 19.8±0.2, 21.2±0.2, 21.4±0.2, 22.2±0.2, 22.7±0.2, 23.0±0.2, 23.7±0.2, 24.2±0.2, 24.4±0.2, 24.6±0.2, 25.1±0.2, 26.8±0.2, 27.1±0.2, 28.5±0.2, 30.1±0.2, 32.9±0.2, 33.7±0.2, and 37.1±0.2 °2θ.

[0206] In some embodiments, the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 8.2±0.2, 10.0±0.2, 11.6±0.2, 13.0±0.2, 13.4±0.2, 15.3±0.2, 16.4±0.2, 16.7±0.2, 17.4±0.2, 18.3±0.2, 18.6±0.2, 18.9±0.2, 19.8±0.2, 20.1±0.2, 21.2±0.2, 21.4±0.2, 22.2±0.2, 22.7±0.2, 23.0±0.2, 23.7±0.2, 24.2±0.2, 24.4±0.2, 24.6±0.2, 25.1±0.2, 26.0±0.2, 26.8±0.2, 27.1±0.2, 28.5±0.2, 30.1±0.2, 32.9±0.2, 33.7±0.2, and 37.1±0.2 °2θ.

[0207] In some embodiments, the mesylate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 43**.

[0208] In some embodiments, the mesylate salt exhibits a weight less of about 2% up to a temperature of about 150 °C as measured by TGA.

[0209] In some embodiments, the mesylate salt exhibits a DSC thermogram comprising an endothermic peak at about 275±2.5 °C.

[0210] In some embodiments, the anion is mesylate and the mitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at about:

Pos. [°2θ]
6.71
11.59
13.38
14.94
16.71
17.37
18.62
18.93
20.07
25.99

±0.2 degrees 2 theta.

[0211] In one aspect the present disclosure provides a crystalline glycolate salt of mitragynine.

[0212] In one aspect the present disclosure provides a glycolate Type A salt of mitragynine exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at about:

Pos. [°2θ]
7.09
10.16
11.29
13.23
14.14
15.74
16.01
18.03
19.54
19.72
20.36
20.93
22.62
25.20
27.63

±0.2 degrees 2 theta.

[0213] In some embodiments, a glycolate Type B salt of mitragynine is provided.

[0214] In some embodiments, a glycolate Type C salt of mitragynine is provided.

[0215] In some embodiments, a glycolate Type D salt of mitragynine is provided.

[0216] In some embodiments, a glycolate Type E salt of mitragynine is provided.

[0217] In some embodiments, a glycolate Type F salt of mitragynine is provided.

Methods of making the salts of the present disclosure

[0218] In one aspect there is provided a process for producing a glycolate salt of 3-deuteromitragynine, the process including the step of crystallizing a glycolate salt of 3-deuteromitragynine from a solution of isopropyl alcohol.

[0219] In some embodiments, the solution of isopropyl alcohol includes water.

[0220] In one aspect there is provided a process for producing a glycolate salt of mitragynine, the process including the step of crystallizing a glycolate salt of mitragynine from a solution of isopropyl alcohol.

[0221] In some embodiments, the solution of isopropyl alcohol includes water.

[0222] In some embodiments, the glycolate salt of mitragynine is derived from a crude alkaloid extract of *Mitragyna speciosa*.

[0223] In one aspect there is provided a process of purifying 3-deuteromitragynine or mitragynine, the purification process including a step of crystallizing any one of the 3-deuteromitragynine or mitragynine salts as defined above.

[0224] In some embodiments, the purified mitragynine is derived from a crude alkaloid extract of *Mitragyna speciosa*.

[0225] In some embodiments, the purification of 3-deuteromitragynine or mitragynine is comprises crystallizing the glycolate, L-lactate, succinate, fumarate, or mesylate salt.

[0226] In some embodiments, the purification of 3-deuteromitragynine or mitragynine comprises crystallizing the glycolate salt.

[0227] In some embodiments, the purification of 3-deuteromitragynine or mitragynine comprises crystallizing the L-lactate salt.

[0228] In some embodiments, the purification of 3-deuteromitragynine or mitragynine comprises crystallizing the succinate salt.

[0229] In some embodiments, the purification of 3-deuteromitragynine or mitragynine comprises crystallizing the fumarate salt.

[0230] In some embodiments, the purification of 3-deuteromitragynine or mitragynine comprises crystallizing the mesylate salt.

[0231] In some embodiments, the purified 3-deuteromitragynine or mitragynine salt is at least 90% free of other compounds or impurities.

[0232] In some embodiments, the purified 3-deuteromitragynine or mitragynine salt is at least 95% free of other compounds or impurities.

[0233] In some embodiments, the purified 3-deuteromitragynine or mitragynine salt is at least 98% free of other compounds or impurities.

[0234] In some embodiments, the purified 3-deuteromitragynine or mitragynine salt is at least 99% free of other compounds or impurities.

[0235] In some embodiments, the purified 3-deuteromitragynine salt has less than about 3% of the impurity 3-deuterocorynantheidine (3-DCR).

[0236] In some embodiments, the purified 3-deuteromitragynine salt has less than about 2% of the impurity 3-DCR.

[0237] In some embodiments, the purified 3-deuteromitragynine salt has less than about 1% of the impurity 3-DCR.

[0238] In some embodiments, the purified 3-deuteromitragynine salt has less than about 0.5% of the impurity 3-DCR.

[0239] In some embodiments, the purified mitragynine salt has less than about 3% of the impurity corynantheidine (CR).


[0240] In some embodiments, the purified mitragynine salt has less than about 2% of the impurity CR.

[0241] In some embodiments, the purified mitragynine salt has less than about 1% of the impurity CR.

[0242] In some embodiments, the purified mitragynine salt has less than about 0.5% of the impurity CR.

Pharmaceutical Compositions

[0243] The present disclosure further provides a pharmaceutical composition comprising an amount of one or more salts of 3-deuteromitragynine having the structure:



[0244] In some embodiments, the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

[0245] In some embodiments, the anion is glycolate.

[0246] In some embodiments, the anion is L-lactate.


[0247] In some embodiments, the anion is succinate.

[0248] In some embodiments, the anion is fumarate.

[0249] In some embodiments, the anion is mesylate.

[0250] In some embodiments, the composition further includes a pharmaceutically acceptable carrier.

[0251] The present disclosure further provides a pharmaceutical composition comprising an amount of one or more salts of mitragynine having the structure:



[0252] In some embodiments, the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

[0253] In some embodiments, the anion is glycolate.

[0254] In some embodiments, the anion is L-lactate.

[0255] In some embodiments, the anion is succinate.

[0256] In some embodiments, the anion is fumarate.

[0257] In some embodiments, the anion is mesylate.

[0258] In some embodiments, the composition further includes a pharmaceutically acceptable carrier.

[0259] As used herein, a "pharmaceutically acceptable carrier" is a pharmaceutically acceptable solvent, suspending agent or vehicle, for delivering the instant salt(s) to the animal or human. The carrier may be liquid or solid and is selected with the planned manner of administration in mind. Liposomes are also a pharmaceutically acceptable carrier, as are capsules, tablets, coatings, and various syringes.

[0260] A dosage unit of the salt(s) used in the method of the present disclosure may comprise a single salt or mixtures thereof. The salt(s) can be administered in oral dosage forms as tablets, capsules, pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. The salt(s) may also be administered in intravenous (bolus or infusion), intraperitoneal, subcutaneous, or intramuscular form, or introduced directly, e.g. by injection, topical application, or other methods, into or onto a site of disease, all using dosage forms well known to those of ordinary skill in the pharmaceutical arts.

[0261] The salts used in the present disclosure can be administered in admixture with suitable pharmaceutical diluents, extenders, excipients, or carriers (collectively referred to herein as a pharmaceutically acceptable carrier) suitably selected with respect to the intended form of administration and as consistent with conventional pharmaceutical practices. Extended-release formulations are specifically encompassed. The unit will be in a form suitable for oral, rectal, topical, intravenous or direct injection or parenteral administration. The salt(s) can be administered alone or mixed with a pharmaceutically acceptable carrier. This carrier can be a solid or liquid, and the type of carrier is generally chosen based on the type of administration being used. The active agent can be co-administered in the form of a tablet or capsule, liposome, as an agglomerated powder or in a liquid form. Examples of suitable solid

carriers include lactose, sucrose, gelatin and agar. Capsule or tablets can be easily formulated and can be made easy to swallow or chew; other solid forms include granules, and bulk powders. Tablets may contain suitable binders, lubricants, diluents, disintegrating agents, coloring agents, flavoring agents, flow-inducing agents, and melting agents. Examples of suitable liquid dosage forms include solutions or suspensions in water, pharmaceutically acceptable fats and oils, alcohols or other organic solvents, including esters, emulsions, syrups or elixirs, suspensions, solutions and/or suspensions reconstituted from non-effervescent granules and effervescent preparations reconstituted from effervescent granules. Such liquid dosage forms may contain, for example, suitable solvents, preservatives, emulsifying agents, suspending agents, diluents, sweeteners, thickeners, and melting agents. Oral dosage forms optionally contain flavorants and coloring agents. Parenteral and intravenous forms may also include minerals and other materials to make them compatible with the type of injection or delivery system chosen.

[0262] Techniques and compositions for making dosage forms useful in the present disclosure are described in the following references: 7 Modern Pharmaceutics, Chapters 9 and 10 (Banker & Rhodes, Editors, 1979); Pharmaceutical Dosage Forms: Tablets (Lieberman et al., 1981); Ansel, Introduction to Pharmaceutical Dosage Forms 2nd Edition (1976); Remington's Pharmaceutical Sciences, 17th ed. (Mack Publishing Company, Easton, Pa., 1985); Advances in Pharmaceutical Sciences (David Ganderton, Trevor Jones, Eds., 1992); Advances in Pharmaceutical Sciences Vol. 7. (David Ganderton, Trevor Jones, James McGinity, Eds., 1995); Aqueous Polymeric Coatings for Pharmaceutical Dosage Forms (Drugs and the Pharmaceutical Sciences, Series 36 (James McGinity, Ed., 1989); Pharmaceutical Particulate Carriers: Therapeutic Applications: Drugs and the Pharmaceutical Sciences, Vol 61 (Alain Rolland, Ed., 1993); Drug Delivery to the Gastrointestinal Tract (Ellis Horwood Books in the Biological Sciences. Series in Pharmaceutical Technology; J. G. Hardy, S. S. Davis, Clive G. Wilson, Eds.); Modem Pharmaceutics Drugs and the Pharmaceutical Sciences, Vol 40 (Gilbert S. Banker, Christopher T. Rhodes, Eds.). All of the aforementioned publications are incorporated by reference herein.

[0263] Tablets may contain suitable binders, lubricants, disintegrating agents, coloring agents, flavoring agents, flow-inducing agents, and melting agents. For instance, for oral administration in the dosage unit form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable,

inert carrier such as lactose, gelatin, agar, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth, or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

[0264] The salt(s) used in the method of the present disclosure may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamallar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

[0265] Gelatin capsules may contain the salt(s) and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as immediate release products or as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

[0266] For oral administration in liquid dosage form, the oral drug components are combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Examples of suitable liquid dosage forms include solutions or suspensions in water, pharmaceutically acceptable fats and oils, alcohols or other organic solvents, including esters, emulsions, syrups or elixirs, suspensions, solutions and/or suspensions reconstituted from non-effervescent granules and effervescent preparations reconstituted from effervescent granules. Such liquid dosage forms may contain, for example, suitable solvents, preservatives, emulsifying agents, suspending agents, diluents, sweeteners, thickeners, and melting agents.

[0267] Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance. In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol

or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propyl-paraben, and chlorobutanol. Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences, Mack Publishing Company, a standard reference text in this field.

[0268] The salt(s) used in the present disclosure may also be administered in intranasal form via use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will generally be continuous rather than intermittent throughout the dosage regimen.

[0269] Parenteral and intravenous forms may also include minerals and other materials to make them compatible with the type of injection or delivery system chosen.

Methods of Treatment

[0270] In one aspect there is provided a method of treating a subject afflicted with acute pain or chronic pain, a depressive disorder, a mood disorder, an anxiety disorder, borderline personality disorder, a substance use disorder, opioid use disorder, opioid withdrawal symptoms, alcohol use disorder, or alcohol withdrawal disorder, comprising administering an effective amount of a salt or a composition as defined above to the subject so as to thereby treat the subject afflicted with acute pain or chronic pain, the depressive disorder, mood disorder, anxiety disorder, borderline personality disorder, a substance use disorder, opioid use disorder, opioid withdrawal symptoms, alcohol use disorder, or alcohol withdrawal disorder.

[0271] In some embodiments, the present disclosure provides methods of treating a subject afflicted with acute pain, chronic pain, a depressive disorder, a mood disorder, an anxiety disorder, borderline personality disorder, a substance use disorder, opioid use disorder, opioid withdrawal symptoms, alcohol use disorder, or alcohol withdrawal

disorder, comprising administering an effective amount of a salt of any of the salts or pharmaceutical compositions described herein to the subject.

[0272] The method of embodiment 139, the subject is afflicted with an opioid use disorder.

[0273] The method of embodiment 140, wherein the subject is afflicted with opioid withdrawal.

[0274] Administration of one or more salt(s) and/or one or more compositions (e.g., pharmaceutical compositions) disclosed herein may be used for preventing, slowing, halting, or reversing the progression of acute pain or chronic pain, a depressive disorder, a mood disorder, an anxiety disorder, borderline personality disorder, a substance use disorder, opioid use disorder, opioid withdrawal symptoms, alcohol use disorder, or alcohol withdrawal disorder as set out herein. Administration may also improve one or more symptoms of acute pain or chronic pain, a depressive disorder, a mood disorder, an anxiety disorder, borderline personality disorder, a substance use disorder, opioid use disorder, opioid withdrawal symptoms, alcohol use disorder, or alcohol withdrawal disorder.

[0275] The salt(s) used in the method of the present disclosure may be administered in various forms, including those detailed herein. The treatment with the salt(s) may be a component of a combination therapy or an adjunct therapy, i.e. the subject or patient in need of the drug is treated or given another drug for the disease in conjunction with one or more of the instant salt(s). This combination therapy can be sequential therapy where the patient is treated first with one drug and then the other or the two drugs are given simultaneously. These can be administered independently by the same route or by two or more different routes of administration depending on the dosage forms employed.

[0276] The dosage of the salt(s) administered in treatment will vary depending upon factors such as the pharmacodynamic characteristics of a specific chemotherapeutic agent and its mode and route of administration; the age, sex, metabolic rate, absorptive efficiency, health and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment being administered; the frequency of treatment with; and the desired therapeutic effect.

[0277] In some embodiments, a disclosed salt may be administered at a dosage unit of about 0.1 mg to about 1000 mg, or about 1 mg to about 400 mg, or about 5 mg to about 300 mg, about 10 mg to about 200 mg, about 100 mg to about 200 mg, or at

least 400 mg, at least 300 mg, at least 200 mg, at least 150 mg, at least 120 mg, at least 100 mg, at least 50 mg, at least 40 mg, at least 30 mg, at least 20 mg, at least 10 mg, at least 9 mg, at least 9.5 mg, at least 8 mg, at least 7.5 mg, at least 7 mg, at least 6.5 mg, at least 6 mg, at least 5.5 mg, at least 5 mg, at least 4.5 mg, at least 4 mg, at least 3.5 mg, at least 3 mg, at least 2.5 mg, at least 2 mg, or at least 1 mg.

[0278] In some embodiments, about 5 mg to about 100 mg, including about 5 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 85 mg, about 90 mg, about 95 mg or about 100 mg, including all values and ranges therebetween, of a salt of deuterated mitragynine as described herein is administered to a patient in need thereof. In some embodiments, about 10 mg to about 90 mg of a salt of deuterated mitragynine as described herein is administered to a patient in need thereof.

[0279] Each embodiment disclosed herein is contemplated as being applicable to each of the other disclosed embodiments. Thus, all combinations of the various elements described herein are within the scope of this disclosure.

[0280] This invention will be better understood by reference to the Experimental Details which follow, but those skilled in the art will readily appreciate that the specific experiments detailed are only illustrative and that the invention is described in the claims which follow thereafter.

[0281] EXAMPLES

[0282] General Experimental Methods

[0283] The general analytical methods used throughout the following examples are described below.

[0284] X-Ray Powder Diffraction (XRPD)

[0285] For XRPD analysis, Empyrean and X'pert 3 X-ray powder diffractometers were used. Samples were spread on the middle of a zero-background Si holder. The XRPD parameters used are described in Table 1.

Table 1. Parameters for XRPD testing

Model	Empyrean	X'Pert3
X-Ray wavelength	Cu, kα, Kα1 (Å): 1.540598, Kα2 (Å): 1.544426 Kα2/Kα1 intensity ratio: 0.50	
X-Ray tube setting	45 kV, 40 mA	45 kV, 40 mA
Divergence slit	Automatic	1/8°
Scan mode	Continuous	Continuous
Scan range (°2TH)	3°-40°	3°-40°
Scan step time (s)	33.020	46.667
Step size (°2TH)	0.0167	0.0263
Test time	10 min 10 s	5 min 4 s

[0286] Thermogravimetric Analysis (TGA) and Differential Scanning Calorimetry (DSC)

[0287] TGA data were collected using a TA Q5000 and Discovery TGA 5500 TGA from TA Instruments. DSC and mDSC were performed using a TA Q2000 DSC and Discovery DSC 2500 from TA Instruments. The detailed parameters used are described in Table 2 and Table 3.

Table 2. Parameters for TGA and DSC testing

Parameters	TGA	DSC
Method	Ramp	Ramp
Sample pan	Aluminum, open	Aluminum, crimped

Temperature	RT to 350 °C	25 °C to target temperature
Heating rate	10 °C/min	10 °C/min
Purge gas	N_2	N_2

Table 3. Parameters for mDSC testing

Parameters	mDSC
Sample pan	Aluminum, crimped
Temperature	0-250 ºC
Heating rate	3 ºC/min
Purge gas	N_2
Amplitude	± 1 ºC
Modulation period	60 s

[0288] Dynamic Vapor Sorption (DVS)

[0289] DVS was measured via an SMS (Surface Measurement Systems) DVS Intrinsic. The relative humidity at 25 °C was calibrated against the deliquescence point of LiCl, $Mg(NO_3)_2$, and KCl. Parameters for DVS testing are described in Table 4.

Table 4. Parameters for DVS testing.

Parameters	DVS
Temperature	25°C
Sample size	10 ~ 20 mg
Gas and flow rate	N_2, 200 mL/min
dm/dt	0.002%/min
Min. Dm/dtstabilityduration	10 min
Max. equilibrium time	180 min
RH range	0%RH ~ 95%RH ~ 0%RH
RH step size	10%RH from 0%RH to 90%RH & 90%RH to 0%RH 5%RH from 90%RH to 95%RH & 95%RH to 90%RH

[0290] **High Pressure Liquid Chromatography (HPLC)**

[0291] An Agilent 1260 HPLC with DAD/VWD detector and Agilent 1100 with DAD detector were utilized. Detailed chromatographic conditions for purity and stoichiometric ratio analysis are described in Table 5-8.

Table 5. Chromatographic conditions for purity analysis, Method 1.

HPLC	**Agilent 1260 HPLC with DAD/VWD detector & Agilent 1100 HPLC with DAD detector**	
Column	Supelco Ascentis Express 2.7 µm 4.6 x 100mm	
Mobile phase	A: H_2O/MeOH (80:20, v/v) + 0.1% H_3PO_4	
	B: MeOH + 0.1% H_3PO_4	
Gradient table	Time (min)	%B
	0.0	0
	20.0	50
	25.0	50
	26.0	70
	27.0	0
	30.0	0
Run time	30.0	
Post time	0.0	
Flow rate	1.0 mL/min	
Injection volume	5 µL	
Detector wavelength	220 nm	
Column temperature	50 ºC	
Sampler temperature	RT	
Diluent	MeOH	

Table 6. Chromatographic conditions for purity analysis, Method 2.

HPLC	**Agilent 1260 HPLC with DAD/VWD detector**	
Column	ZORBAX SB-Phenyl C18, 4.6 x 150 mm, 3.5 µm	
Mobile phase	A: H_2O/MeOH (80:20, v/v) + 0.1% H_3PO_4	
	B: MeOH + 0.1% H_3PO_4	
Gradient table	Time (min)	%B

HPLC	Agilent 1260 HPLC with DAD/VWD detector	
	0.0	0
	20.0	50
	25.0	50
	26.0	70
	27.0	0
	30.0	0
Run time	30.0	
Post time	0.0	
Flow rate	1.0 mL/min	
Injection volume	5 µL	
Detector wavelength	220 nm	
Column temperature	50 ºC	
Sampler temperature	RT	
Diluent	MeOH	

Table 7. Chromatographic conditions for stoichiometric ratio and solubility analysis, Method 1.

HPLC	Agilent 1260 HPLC with DAD/VWD detector & Agilent 1100 HPLC with DAD detector	
Column	Supelco Ascentis Express 2.7 µm 4.6 x 100mm	
Mobile phase	A: H_2O/MeOH (80:20, v/v) + 0.1% H_3PO_4	
	B: MeOH + 0.1% H_3PO_4	
Gradient table	Time (min)	%B
	0.00	0
	8.00	60
	10.00	60
	10.01	0
	12.00	0
Run time	12.0 min	
Post time	0.0 min	
Flow rate	1.0 mL/min	

HPLC	Agilent 1260 HPLC with DAD/VWD detector & Agilent 1100 HPLC with DAD detector
Injection volume	5 µL
Detector wavelength	220 nm
Column temperature	50 ºC
Sampler temperature	RT
Diluent	MeOH

Table 8. Chromatographic conditions for stoichiometric ratio and solubility analysis, Method 2.

HPLC	Agilent 1260 HPLC with DAD/VWD detector	
Column	ZORBAX SB-Phenyl C18, 4.6 x 150 mm, 3.5 µm	
Mobile phase	A: H_2O/MeOH (80:20, v/v) + 0.1% H_3PO_4 B: MeOH + 0.1% H_3PO_4	
Gradient table	Time (min)	%B
	0.00	0
	8.00	60
	10.00	60
	10.01	0
	12.00	0
Run time	12.0 min	
Post time	0.0 min	
Flow rate	1.0 mL/min	
Injection volume	5 µL	
Detector wavelength	220 nm	
Column temperature	50 ºC	
Sampler temperature	RT	
Diluent	MeOH	

[0292] **Ion Chromatography (IC)**

[0293] A ThermoFisher ICS-1100 was utilized for ion chromatography (IC) and detailed conditions are described in Table 9.

Table 9. Ion chromatography conditions and parameters.

IC	ThermoFisher ICS-1100
Column	IonPac AS18 Analytical Column (4 × 250 mm)
Mobile phase	25 mM NaOH
Injection volume	25 µL
Flow rate	1.0 mL/min
Cell temperature	35 °C
Column temperature	35 °C
Current	80 mA
Run time	7.0 min for Cl^-, 16.0 min for $C_2O_4^{2-}$, 40.0 min for PO_4^{3-}

[0294] **Polarized Light Microscopy (PLM)**

[0295] PLM images were captured with a ZEISS Scope.A1 microscope.

[0296] **Karl Fischer (KF) Titration**

[0297] The instrument (Metrohm 870 KF Titrinoplus) was calibrated using purified water and the titration reagent was Hydranal® R-Composite 5 provided by Sigma-Aldrich. HPLC grade methanol was used to dissolve samples.

[0298] **^{1}H Solution Nuclear Magnetic Resonance (NMR)**


^{1}H solution NMR was collected on a Bruker 400 MHz NMR spectrometer using deuterated DMSO as solvent.

[0299] **Example 1. Preparation and Characterization of 3-Deuteromitragynine (3-DM) Free Base for Salt Screening**

[0300] 3-Deuteromitragynine (3-DM) free base for use in salt and polymorph screening experiments was prepared as previously described in WO2020160280 – entitled "Deuterated mitragynine analogs as safer opioid modulators in the mitragynine class, which reference is incorporated herein by reference." Reagents and solvents were obtained from commercial sources and were used without further purification unless otherwise stated. Reactions were monitored by TLC using solvent mixtures appropriate to each reaction. All column chromatography was performed on silica gel (40-63 µm). Preparative TLC was conducted on glass plates coated with a 1 mm silica

layer. Nuclear magnetic resonance spectra were recorded on Bruker 400 or 500 MHz instruments, as indicated. Chemical shifts are reported as δ values in ppm referenced to $CDCl_3$ (^1H NMR = 7.26 and ^{13}C NMR = 77.16) or methanol-d_4 (^1H NMR = 3.31 and ^{13}C NMR = 49.00). Multiplicity is indicated as follows: s (singlet); d (doublet); t (triplet); dd (doublet of doublets); td (triplet of doublets); dt (doublet of triplets); ddd (doublet of doublet of doublets); m (multiplet); br (broad). All carbon peaks are rounded to one decimal place unless such rounding would cause two close peaks to become identical; in these cases, two decimal places are retained. Low-resolution mass spectra were recorded on an Advion quadrupole instrument (ionization mode: APCI+). Percent deuteration was determined by mass spectrometry on a high-resolution quadrupole-time-of-flight instrument (ionization mode: ESI+) by quantitative comparison of the isotope pattern of deuterated compounds to controls having natural isotopic abundance. The following Scheme 1 also shows the process involved in preparing 3-DM.

Scheme 1. Preparation of 3-deuteromitragyine.



[0301] Mitragynine. Mitragynine free base was obtained by extraction from powdered Mitragyna speciosa leaves as previously described (Kruegel et al. 2016). Spectral and

physical properties were in agreement with those previously reported (Kruegel et al. 2016).

[0302] 7-Hydroxymitragynine (7-OH) Procedure 1). Mitragynine (1.99 g, 5.00 mmol) was dissolved in acetone (100 mL), saturated aqueous $NaHCO_3$ (10 mL) was added, and the mixture was cooled to 0 °C. A solution of Oxone monopersulfate ($2KHSO_5 \cdot KHSO_4 \cdot K_2SO_4$; 2.31 g, 3.75 mmol) in water (10 mL) was then added dropwise over 35 minutes and the mixture left to stir at 0 °C. After 45 minutes, additional Oxone monopersulfate (769 mg, 1.25 mmol) in water (3.3 mL) was added over ~2 minutes and stirring was continued at 0 °C for an additional 15 minutes. At this time, the reaction was diluted with water (150 mL) and extracted with EtOAc (3 x 50 mL). The combined organics were washed with brine (50 mL), dried over Na_2SO_4, and concentrated *in vacuo* to give the crude product as a tan foam (1.42 g). This material was purified by column chromatography (6:4 hexanes:EtOAc + 2% Et_3N) to provide pure 7-hydroxymitragynine as an amorphous, pale-yellow solid (882 mg, 43%). Spectral and physical properties were in agreement with those previously reported (Kruegel et al. 2016).

[0303] 7-Hydroxymitragynine (7-OH) (Procedure 2 - Larger Scale). Mitragynine (9.96 g, 25.00 mmol) was dissolved in acetone (750 mL), saturated aqueous $NaHCO_3$ (500 mL) was added, and the mixture was cooled to 0 °C. A solution of Oxone monopersulfate ($2KHSO_5 \cdot KHSO_4 \cdot K_2SO_4$; 15.39 g, 25.00 mmol) in water (250 mL) was then pre-cooled to 0 °C and added dropwise over 30 minutes (mixture was hard to stir at first but became less viscous over the course of the addition). TLC at the end of the Oxone addition showed no starting material so the reaction was worked up (at 15 minutes after the end of the addition). EtOAc (500 mL) and water (500 mL) were added to the reaction mixture while it was still stirring at 0 °C and the resulting mixture was then poured into a separatory funnel containing additional water (1,000 mL). The organic layer was separated and the aqueous phase extracted with additional EtOAc (2 x 500 mL). The combined organics were washed with brine (300 mL), dried over Na_2SO_4, and concentrated *in vacuo* to give the crude product as a yellow ochre foam (7.35 g). This material was purified by silica column chromatography (320 g silica; 600 mL column volume; 60 mL fractions; step gradient: 20% → 30% → 35% → 40% → 45% → 50% → 55% EtOAc in hexanes + 2% Et_3N, 1 column volume per step) to provide the following fractions: fractions 49-51 = very pale-yellow amorphous solid, 7-hydroxymitragynine + ~2% 7-hydroxycorynantheidine, 1.09 g (11%); fractions 52-64 =

pale-yellow amorphous solid, 7-hydroxymitragynine, 2.99 g (29%). Spectral properties were in agreement with those previously reported (Kruegel et al. 2016).

[0304] 3-Dehydromitragynine hydrochloride (DHM) (Procedure 1). To a solution of 7-hydroxymitragynine (746 mg, 1.80 mmol) in anhydrous CH_2Cl_2 (27 mL) under argon was added 2.0 M HC1 in Et_20 (9.0 mL) and the resulting mixture was stirred at room temperature for 45 minutes (all solids dissolved to give a transparent yellow solution after 2-3 minutes). The reaction mixture was then concentrated directly *in vacuo* to give pure 3-dehydromitragynine hydrochloride as a yellow solid (797 mg, quantitative). ^{1}H NMR (500 MHz, CDC13) d 13.56 (br s,1H), 7.49 (s, 1H) , 7.42 (d, J= 8.4 Hz, 1H), 7.25 (t, J = 8.0 Hz 1H), 6.38 (d, J = 7.7 Hz, 1H), 4.03 - 3.81 (m, 3H), 3.89 (s, 3H) , 3.80 - 3.65 (m, 1H), 3.76 (s, 3H) , 3.65 - 3.53 (m, 2H) , 3.61 (s, 3H) , 3.53 - 3.36 (m, 2H) , 3.29 (t, J= 12.6 Hz, 1H) , 2.10 (br s, 1H), 1.55 - 1.43 (m, 1H), 1.22 - 1.10 (m, 1H) , 0.98 (t, J = 7.4 Hz, 3H).

[0305] 3-Dehydromitragynine hydrochloride (DHM) (Procedure 2 - Larger Scale). To a solution of 7-hydroxymitragynine (14.09 g, 34.00 mmol) in anhydrous CH_2Cl_2 (510 mL) under argon was added 2.0 M HC1 in Et_20 (170 mL) (yellow suspension forms and slight warming occurs on HCl addition) and the resulting mixture was stirred at room temperature for 40 minutes (all solids dissolved to give a transparent yellow-orange solution after 2-3 minutes). The reaction mixture was then concentrated directly *in vacuo* to give pure 3-dehydromitragynine hydrochloride as a yellow solid (15.87 g, quantitative). The NMR spectra of this material were identical to those of material obtained via Procedure 1 above.

[0306] 3-Deuteromitragynine (3-DM) (Procedure 1). To a solution of 3-dehydromitragynine hydrochloride (606 mg, 1.40 mmol) in methanol-d_4 (28 mL) at 0 °C was added $NaBD_4$ (293 mg, 7.00 mmol) and the yellow solution was stirred at 0 °C for 20 minutes. The reaction was then diluted with water (100 mL) and extracted with CH_2Cl_2 (3 x 50 mL). The combined organics were washed with water (2 x 50 mL), dried over Na_2SO_4, and concentrated *in vacuo* to give the crude product as a very pale-yellow foam (0.52 g). This material was purified by column chromatography (8:2 hexanes:EtOAc + 2% Et_3N, 4 column volumes → 7:3 hexanes:EtOAc + 2% Et_3N, 3 column volumes) to provide pure 3-deuteromitragynine as an amorphous, off-white solid (480 mg, 86%) . 1H NMR (500 MHz, $CDCl_3$) d 7.70 (br s, 1H), 7.43 (s, 1H), 6.99 (t, J = 7.9 Hz, 1H), 6.90 (d, J= 7.7 Hz, 1H),· 6.46 (d, J = 7.7 Hz, 1H), 3.88 (s, 3H), 3.73 (s, 3H), 3.71 (s, 3H), 3.12 (ddd, J = 15.8, 11.6, 5.9 Hz, 1H), 3.07 - 2.89 (m, 4H), 2.58

- 2.42 (m, 3H) , 1.85 - 1.73 (m, 2H), 1.66 - 1.58 (m, 1H), 1.25 - 1.15 (m, 1H), 0.87 (t, J = 7.4 Hz, 3H);

[0307] ^{13}C NMR (126 MHz, $CDCl_3$) d 169.4, 160.7, 154.6, 137.4, 133.8, 121.9, 117.7, 111.6, 107.9, 104.4, 99.8, 61.6, 60.9 (t, JCD = 19.5 Hz), 57.9, 55.4, 53.9, 51.5, 40.8, 40.1, 29.9, 24.1, 19.2, 13.0; HR-MS calcd. for $C_{23}H_{30}DN_2O_4$ $[M+H]^+$ 400.2341, found 400.2332; Deuterium Enrichment = °97.5 - 97.7 atom % D (by HR-MS).

[0308] 3-Deuteromitragynine (3-DM) (Procedure 2). To a solution of 3-dehydromitragynine hydrochloride (54.1 mg, 0.125 mmol) in MeOH (2.5 mL) at 0 °C was added NaBD4 (26.2 mg, 0.625 mmol) and the yellow solution was allowed to warm to room temperature and stirred for 25 minutes. The reaction was then diluted with water (10 mL) and extracted with CH_2Cl_2 (3 x 5 mL). The combined organics were washed with water (2 x 5 mL), dried over Na2SC>4, and concentrated *in vacuo* to give the crude product as a foamy yellow glass (47.2 mg). This material was purified by column chromatography (7:3 hexanes:EtOAc + 2% Et_3N) to provide pure 3-deuteromitragynine as an amorphous, yellow solid (39.4 mg, 79%). The NMR spectra of this material were identical to those of material obtained via Procedure 1 above, with the exception of visible residual peaks for undeuterated mitragynine in both the proton and carbon spectra. Deuterium Enrichment = 93.5 - 93.8 atom % D (by HR-MS).

[0309] 3-Deuteromitragynine (3-DM) (Procedure 3). To a solution of 3-dehydromitragynine hydrochloride (14.72 g, 34.00 mmol = 15.85 g of crude containing CH_2Cl_2 from last step) in methanol-OD (CH_3OD; 170 mL) at 0 °C was added $NaBD_4$ (2.85 mg, 68.00 mmol) and the yellow solution (clouds immediately after NaBD4 addition) was stirred at 0 °C for 20 minutes (effervescence stops after 10 minutes). The reaction was then diluted with water (500 mL) and extracted with CH_2Cl_2 (3 x 250 mL). The combined organics were washed with water (2 x 250 mL), dried over Na_2SO_4, and concentrated *in vacuo* to give the crude product as a pale-yellow foam (14.28 g). This material was purified by silica column chromatography (320 g silica; 600 mL column volume; 60 mL fractions; step gradient: 10% (2 column volumes) → 20% (2 column volumes) → 30% (4 column volumes) EtOAc in hexanes + 2% Et_3N, first 2 column volumes discarded) to provide the following fractions: fractions 19-45 = cream-colored amorphous solid, 3-deuteromitragynine, 11.86 g (87%); fractions 17-18 + 46-55 = pale-yellow amorphous solid, impure 3-deuteromitragynine, 0.66 g (~5%). The NMR spectra of this material were identical to those of material obtained

via Procedures 1 and 2 above. Deuterium Enrichment = 98.2 - 98.4 atom % D (by HR-MS).

[0310] 3-DM free base prepared by the above procedures for use in the salt and polymorph screening described below was characterized by XRPD, TGA, and mDSC and ^{1}H NMR. As the XRPD pattern in Figure 1 shows, the starting material was amorphous. TGA/mDSC curves of starting material are shown in Figure 2. The TGA curve showed a weight loss of 1.8% up to 140 ºC. The mDSC curve showed a glass transition temperature (Tg) at 89.8 ºC (middle temperature). ^{1}H NMR data in Figure 3 was collected using DMSO-d_6 as solvent.

[0311] The approximate solubility of the free base starting material was determined in 27 solvent systems at RT. Approximate 2 mg of material was added into a 3-mL glass vial. The solvents in Table 10 were then added stepwise into the vials until the solids were dissolved visually or a total volume of 1 mL was reached. The solubility results summarized in Table 10 were used to guide the solvent selection in salt screening design.

Table 10. Approximate solubility of amorphous 3-DM free base starting material (810397-01-A) at RT.

Solvent	Solubility (mg/mL)	Solvent	Solubility (mg/mL)
MeOH	S>46.0	MTBE	S>46.0
EtOH	S>38.0	THF	S>40.0
IPA	S>36.0	2-MeTHF	S>48.0
Acetone	S>38.0	1,4-dioxane	S>42.0
MIBK	S>40.0	ACN	S>46.0
EtOAc	S>40.0	DCM	S>42.0
IPAc	S>42.0	$CHCl_3$	S>46.0
n-Heptane	S<2.3	IPA/H_2O (4:1)	7.0<S<21.0
Toluene	S>36.0	Acetone/n-Heptane (4:1)	S>42.0

Solvent	Solubility (mg/mL)	Solvent	Solubility (mg/mL)
DMSO	S>34.0	Acetone/n-Heptane (1:1)	S>38.0
DMAc	S>46.0	Acetone/n-Heptane (1:4)	S>46.0
NMP	S>44.0	Acetone/n-Heptane (1:9)	7.7<S<23.0
H_2O	S<2.3	Acetone/n-Heptane (1:15)	4.0<S<6.7
--	--	IPA/H_2O (9:1)	23.0<S<46.0

[0312] The purity and 3-deuterocorynantheidine (3-DCR) content of the 3-DM free base starting material were also determined by HPLC, with the results summarized in Table 11.

Table 11. Impurities summary of amorphous 3-DM free base starting material (810397-01-A).

#Peak	RRT	Area (%)
1	0.70	0.27
2	0.71	0.43
3	0.83	0.12
4	0.90 (3-DCR)	2.68
5	0.94	0.21
6	0.95	0.08
7	**1.00 (3-DM)**	**96.21**

[0313] **Example 2. Salt Screening of 3-Deuteromitragynine**

[0314] According to the approximate solubility of the free base starting material at room temperature (RT, 24 ± 2 ºC) and simulated pKa value (7.33, basic), a total of 100 salt screening experiments were performed using 25 acids in four solvent systems via solvent-assisted reaction crystallization. For each experiment, about 20 mg of starting material and equimolar acid were mixed into each HPLC vial. 0.5 mL of corresponding

solvent was then added to form a suspension, which was magnetically stirred (~1,000 rpm) at RT for about three days. Solids were isolated and dried at 50 ºC under vacuum for 4 hrs for XRPD analysis. As summarized in Table 12, a total of 26 crystalline salt hits and one crystalline form of free base were obtained during screening. Purity of all crystalline samples was tested, with results summarized in Table 13. All salt hits were characterized by XRPD, TGA, and DSC. The stoichiometric ratio was determined by ^{1}H NMR or HPLC/IC. Characterization results of salt hits and free base form are listed in Table 14.

Table 12. Summary of salt screening results for 3-DM.

#	Acid	Solvent (v/v)			
		A IPA/H_2O (4:1)	**B** EtOAc	**C** MTBE	**D** Acetone/n-Heptane (1:15)
0	Blank	Free Base Type A	Amorphous*	Amorphous*	Amorphous
1	HCl	Amorphous#	HCl salt Type A	HCl salt Type B	HCl salt Type B
2	H_2SO_4	Free Base Type A	Amorphous	Amorphous	Amorphous
3	H_3PO_4	Gel	Phosphate Type A	Phosphate Type A	Phosphate Type A
4	Maleic acid	Amorphous#	Amorphous	Maleate Type A	Maleic acid (one peak)
5	L-tartaric acid	Gel	L-tartrate Type A	Amorphous	L-tartaric acid
6	Fumaric acid	Fumarate Type A	Fumarate Type A	Fumarate Type A	Fumarate Type A + Fumaric acid
7	Citric acid	Gel	Amorphous	Amorphous + peaks	Citric acid
8	Glycolic acid	Glycolate Type A	Glycolate Type A+ B	Glycolate Type B	Glycolate Type A

#	Acid	Solvent (v/v)			
		A IPA/H_2O (4:1)	**B** EtOAc	**C** MTBE	**D** Acetone/n-Heptane (1:15)
9	L-Malic acid	Amorphous*	Amorphous	Amorphous	Amorphous
10	Hippuric acid	Amorphous*	Hippuric acid	Hippuric acid	Hippuric acid
11	DL-Lactic acid	Lactate Type A	Lactate Type A	Lactate Type A	Lactate Type A
12	Adipic acid	Adipate Type A	Adipate Type B	Adipate Type B	Adipate Type A+ Adipic acid
13	Acetic acid	Amorphous*	Acetate Type A	Acetate Type A	Acetate Type A
14	L-Ascorbic acid	Gel	L-Ascorbic acid	L-Ascorbic acid	L-Ascorbic acid
15	Succinic acid	Succinate Type A	Succinate Type A	Succinate Type A + Succinic acid	Succinate Type A + Succinic acid
16	L-Aspartic acid	Free Base Type A + peaks	L-Aspartic acid	L-Aspartic acid	L-Aspartic acid
17	Malonic acid	Amorphous*	Malonate Type A	Amorphous	Malonate Type B
18	Gentisic acid	Amorphous*	Gentisate Type A	Gentisate Type B	Weakly crystalline
19	Ethane-1.2-disulfonic acid	Edisylate Type A	Edisylate Type A	Edisylate Type B	Edisylate Type A
20	Benzoic acid	Amorphous*	Amorphous*	Amorphous*	Benzoate Type A
21	Ethanesulfonic acid	Esylate Type A	Esylate Type A	Esylate Type A	Esylate Type A
22	Methanesulfonic acid	Mesylate Type A#	Mesylate Type A	Mesylate Type A	Mesylate Type A
23	Benzenesulfonic acid	Amorphous#	Weakly crystalline	Amorphous	Besylate Type A

#	Acid	Solvent (v/v)			
		A IPA/H_2O (4:1)	**B** EtOAc	**C** MTBE	**D** Acetone/n-Heptane (1:15)
24	p-Toluenesulfonic acid	Tosylate Type A[#]	Tosylate Type A	Tosylate Type A	Tosylate Type A
25	Oxalic acid	Oxalate Type A	Oxalate Type B	Oxalate Type B	Weakly crystalline

[0315][#]: Clear solution was obtained after slurry at RT. Anti-solvent MTBE was added for precipitation.

[0316][*]: Clear solution was obtained after adding anti-solvent MTBE, which was transferred to evaporate at RT.

Table 13. Purity summary of crystalline hits for 3-DM (3-DCR = 3-deuterocorynantheidine).

#	Acid	Area (%)							
		A		B		C		D	
		3-DCR	3-DM	3-DCR	3-DM	3-DCR	3-DM	3-DCR	3-DM
0	Blank	1.01	98.63	--	--	--	--	--	--
1	HCl	--	--	2.62	96.22	2.02	94.59	2.46	92.60
2	H_2SO_4	1.41	98.14	--	--	--	--	--	--
3	H_3PO_4	--	--	2.26	97.29	2.22	97.33	2.55	96.06
4	Maleic acid	--	--	--	--	2.67	96.21	--	--
5	L-tartaric acid	--	--	2.52	96.89	--	--	2.23	96.50
6	Fumaric acid	2.53	96.81	2.64	96.48	2.67	96.25	--	--
7	Citric acid	--	--	--	--	--	--	2.28	95.75
8	Glycolic acid	0.84	98.90	2.29	97.45	2.39	97.13	3.13	95.49
9	L-Malic acid	--	--	--	--	--	--	--	--
10	Hippuric acid	--	--	--	--	--	--	--	--
11	DL-Lactic acid	1.16	98.55	2.47	96.55	2.59	96.20	2.69	96.14
12	Adipic acid	1.89	97.45	2.08	97.07	2.46	96.49	--	--
13	Acetic acid	--	--	0.76	99.05	1.37	98.21	1.75	97.42
14	L-Ascorbic acid	--	--	--	--	--	--	--	--
15	Succinic acid	1.96	97.66	2.47	96.90	--	--	--	--
16	L-Aspartic acid	--	--	2.84	96.69	2.66	96.14	2.22	96.44
17	Malonic acid	--	--	--	--	2.51	96.39	2.40	96.69
18	Gentisic acid	--	--	2.32	97.16	2.33	97.04	1.72	97.51
19	Ethane-1.2-disulfonic acid	0.89	98.75	2.05	95.15	2.47	87.63	2.44	91.05
20	Benzoic acid	--	--	2.50	96.62	--	--	2.25	96.98
21	Ethanesulfonic acid	1.61	97.44	2.22	97.10	1.89	97.39	1.85	96.97
22	Methanesulfonic acid	2.68	95.94	1.57	98.11	2.41	96.48	2.33	92.68

#	Acid	Area (%)							
		A		B		C		D	
		3-DCR	3-DM	3-DCR	3-DM	3-DCR	3-DM	3-DCR	3-DM
23	Benzenesulfonic acid	--	--	--	--	--	--	2.49	94.92
24	p-Toluenesulfonic acid	2.29	96.99	2.57	93.90	2.28	96.26	2.58	94.60
25	Oxalic acid	3.07	96.42	2.61	95.98	2.62	96.39	--	--

Table 14. Characterization results of salt hits for 3-DM.

Salt Hit (Sample ID)	TGA Weight loss (%, temp.)	DSC endotherm (peak, ºC)	Molar ratio (acid/free base)
HCl salt Type A (810397-04-B1)	2.6 (to 150 ºC)	202.2, 262.2	1.1:1
HCl salt Type B (810397-04-D1)	9.3 (to 175 ºC)	69.5, 198.3, 256.4	1.3:1
Phosphate Type A (810397-04-D3)	7.9 (to 200 ºC)	72.4, 178.1, 194.3, 232.2	1.2:1
Maleate Type A (810397-04-C4)	10.8 (to 150 ºC)	91.0, 168.9, 212.2	1.0:1
L-Tartrate Type A (810397-04-B5)	5.9 (to 150 ºC)	96.1, 146.4, 158.6	1.0:1
Fumarate Type A (810397-04-C6)	3.6 (to 150 ºC)	256.4	1.0:1
Glycolate Type A (810397-04-D8)	3.6 (to 150 ºC)	216.9	1.1:1
Glycolate Type B (810397-04-C8)	1.9 (to 150 ºC)	69.1, 126.0, 220.8	1.1:1
Lactate Type A (810397-04-B11)	1.5 (to 150 ºC)	221.5	1.1:1

Salt Hit (Sample ID)	TGA Weight loss (%, temp.)	DSC endotherm (peak, ºC)	Molar ratio (acid/free base)
Adipate Type A (810397-04-A12)	5.0 (to 170 ºC)	212.0	0.9:1
Adipate Type B (810397-04-C12)	4.9 (to 170 ºC)	211.1	1.0:1
Acetate Type A (810397-04-D13)	13.0 (to 180 ºC)	152.3	1.2:1
Succinate Type A (810397-04-B15)	1.4 (to 120 ºC)	201.2	1.1:1
Malonate Type A (810397-04-B17)	3.6 (to 110 ºC)	110.7, 154.4	1.2:1
Malonate Type B (810397-04-D17)	3.3 (to 100 ºC)	79.2, 109.7, 155.7	2.0:1
Gentisate Type A (810397-04-B18)	8.8 (to 200 ºC)	182.4	1.0:1
Gentisate Type B (810397-04-C18)	8.8 (to 200 ºC)	171.9, 179.5	1.1:1
Edisylate Type A (810397-04-A19)	16.7 (to 200 ºC)	69.2, 148.2, 259.1	0.5:1
Edisylate Type B (810397-04-C19)	16.0 (to 200 ºC)	70.7, 167.3, 191.5, 225.2	0.6:1
Benzoate Type A (810397-04-D20)	9.8 (to 130 ºC)	108.5, 212.6, 241.2	1:1
Esylate Type A (810397-04-C21)	2.5 (to 150 ºC)	230.0, 269.7	0.9:1
Mesylate Type A (810397-04-C22)	2.2 (to 150 ºC)	271.9	1.0:1
Besylate Type A (810397-04-D23)	7.0 (to 170 ºC)	144.4, 261.6	0.9:1
Tosylate Type A (810397-04-B24)	8.0 (to 170 ºC)	95.2, 234.0	1.0:1
Oxalate Type A	6.2 (to 150 ºC)	82.0, 240.8	0.8:1

Salt Hit (Sample ID)	TGA Weight loss (%, temp.)	DSC endotherm (peak, °C)	Molar ratio (acid/free base)
(810397-04-A25) Oxalate Type B (810397-04-B25)	6.0 (to 150 °C)	243.7	1.1:1
Free Base Type A (810397-04-A0)	11.7 (to 110 °C)	61.8, 80.6	--

[0317] XRPD results for 3-DM salt leads are provided below in Table 15 and the respective XRPD traces are shown in Figures 4 to 8.

Table 15. XRPD copper radiation results of salt leads for 3-DM.

3-DM Salt	Pos. [°2θ]	Height [cts]	FWHM Left [°2θ]	d-spacing [Å]	Rel. Int. [%]
Fumarate Type A (Fig 4)	6.23	466.62	0.1279	14.19	3.06
	8.41	15233.33	0.1279	10.51	100.00
	9.58	1168.41	0.0768	9.23	7.67
	13.43	337.14	0.0768	6.60	2.21
	14.40	327.55	0.1279	6.15	2.15
	15.62	223.48	0.1023	5.67	1.47
	16.16	300.94	0.1023	5.49	1.98
	16.85	508.24	0.1023	5.26	3.34
	17.46	2895.93	0.1279	5.08	19.01
	18.68	456.62	0.1279	4.75	3.00
	19.20	2583.93	0.1279	4.62	16.96
	21.59	1167.87	0.1279	4.12	7.67
	22.35	293.44	0.1023	3.98	1.93
	23.03	871.16	0.1023	3.86	5.72
	23.35	350.40	0.1023	3.81	2.30
	23.75	496.99	0.1279	3.75	3.26
	25.36	1302.70	0.2047	3.51	8.55
	25.77	2088.21	0.1535	3.46	13.71
	26.96	454.35	0.1279	3.31	2.98

	28.94	481.81	0.1279	3.09	3.16
	31.11	892.87	0.1279	2.87	5.86
	32.82	223.25	0.1791	2.73	1.47
	34.04	153.07	0.1535	2.63	1.00
	37.97	37.54	0.3070	2.37	0.25
L-Lactate Type A (Fig 5)	6.86	912.47	0.1023	12.88	41.23
	9.96	2213.14	0.1023	8.88	100.00
	10.73	383.13	0.0768	8.24	17.31
	11.05	1598.46	0.1023	8.01	72.23
	13.02	279.14	0.1023	6.80	12.61
	13.80	135.41	0.1023	6.42	6.12
	15.74	1336.33	0.1023	5.63	60.38
	17.74	684.95	0.1279	5.00	30.95
	18.11	126.02	0.1535	4.90	5.69
	18.80	239.00	0.0768	4.72	10.80
	19.29	732.83	0.1023	4.60	33.11
	19.77	501.70	0.1023	4.49	22.67
	20.63	1665.56	0.1279	4.31	75.26
	22.26	1777.09	0.1279	3.99	80.30
	23.62	162.88	0.2047	3.77	7.36
	24.38	294.07	0.1279	3.65	13.29
	24.75	1351.16	0.1023	3.60	61.05
	26.96	266.73	0.1023	3.31	12.05
	27.18	294.90	0.1023	3.28	13.33
	28.01	105.83	0.1535	3.19	4.78
	29.30	186.15	0.1023	3.05	8.41
	31.22	245.17	0.1023	2.86	11.08
	33.81	111.51	0.1791	2.65	5.04
	35.59	60.07	0.4093	2.52	2.71
Glycolate Type A (Fig 6)	7.06	2750.57	0.0768	12.53	48.83
	10.12	5633.51	0.1023	8.74	100.00
	11.25	3840.63	0.0768	7.87	68.17
	13.18	448.79	0.1023	6.72	7.97
	14.11	670.06	0.0768	6.28	11.89

	15.04	93.51	0.1535	5.89	1.66
	15.97	2524.11	0.1023	5.55	44.81
	17.99	1120.78	0.1023	4.93	19.89
	18.11	988.75	0.0768	4.90	17.55
	18.51	291.60	0.1023	4.79	5.18
	18.87	171.68	0.1023	4.70	3.05
	19.19	375.37	0.1023	4.62	6.66
	19.49	1396.21	0.1023	4.55	24.78
	19.69	863.13	0.0768	4.51	15.32
	20.34	1027.37	0.1023	4.37	18.24
	20.88	3514.53	0.1023	4.25	62.39
	22.58	2504.54	0.1279	3.94	44.46
	23.24	183.27	0.1023	3.83	3.25
	23.72	214.78	0.1279	3.75	3.81
	24.02	259.94	0.1023	3.71	4.61
	24.90	417.41	0.1279	3.58	7.41
	25.16	2177.96	0.1023	3.54	38.66
	27.60	777.81	0.1279	3.23	13.81
	28.27	106.24	0.1535	3.16	1.89
	29.55	140.41	0.1535	3.02	2.49
	30.15	281.33	0.1535	2.96	4.99
	30.59	189.47	0.1279	2.92	3.36
	31.61	447.05	0.1023	2.83	7.94
	32.24	128.20	0.1535	2.78	2.28
	34.18	121.06	0.1535	2.62	2.15
Succinate Type A (Fig 7)	6.23	495.96	0.0768	14.19	3.58
	8.46	13846.33	0.1023	10.45	100.00
	9.62	919.89	0.1023	9.20	6.64
	10.06	187.37	0.1023	8.79	1.35
	14.35	804.02	0.1023	6.17	5.81
	15.65	585.24	0.1023	5.66	4.23
	16.12	599.70	0.1023	5.50	4.33
	16.94	382.32	0.0768	5.23	2.76
	17.56	2944.00	0.1023	5.05	21.26

	18.72	326.94	0.0768	4.74	2.36
	19.27	2840.79	0.1279	4.61	20.52
	21.19	166.60	0.1535	4.19	1.20
	21.71	1024.55	0.1279	4.09	7.40
	22.33	378.95	0.1023	3.98	2.74
	23.13	984.83	0.1023	3.84	7.11
	23.73	395.60	0.1279	3.75	2.86
	24.48	113.21	0.1535	3.64	0.82
	25.03	318.43	0.1023	3.56	2.30
	25.34	647.53	0.0768	3.52	4.68
	25.53	1081.82	0.1023	3.49	7.81
	25.94	1289.20	0.1279	3.44	9.31
	26.98	297.64	0.1535	3.30	2.15
	29.08	270.84	0.1023	3.07	1.96
	30.54	135.43	0.1535	2.93	0.98
	31.27	845.96	0.1279	2.86	6.11
	32.99	168.07	0.1279	2.72	1.21
	34.25	105.34	0.1535	2.62	0.76
Mesylate Type A (Fig 8)	6.67	2264.45	0.1279	13.26	100.00
	8.18	238.23	0.0768	10.81	10.52
	10.00	169.74	0.1023	8.85	7.50
	11.56	728.33	0.1279	7.66	32.16
	13.34	524.16	0.1279	6.64	23.15
	14.93	504.38	0.1023	5.94	22.27
	15.29	241.58	0.1279	5.79	10.67
	16.69	1058.47	0.1279	5.31	46.74
	17.34	941.15	0.1535	5.12	41.56
	18.60	593.38	0.1279	4.77	26.20
	18.89	715.15	0.1535	4.70	31.58
	19.76	357.10	0.1023	4.49	15.77
	20.03	700.95	0.1279	4.43	30.95
	21.13	190.26	0.1535	4.20	8.40
	21.42	211.16	0.1535	4.15	9.33
	22.21	242.58	0.1279	4.00	10.71

	22.72	308.31	0.1023	3.91	13.62
	23.69	187.94	0.1023	3.76	8.30
	24.40	295.39	0.1535	3.65	13.04
	25.08	208.62	0.1791	3.55	9.21
	25.97	465.61	0.1791	3.43	20.56
	26.92	117.37	0.3070	3.31	5.18
	28.51	96.14	0.2047	3.13	4.25
	32.80	52.09	0.3070	2.73	2.30

[0318] XRPD data for the other 3-DM salt forms made and characterized are listed in Table 16 and their respective XRPD traces are shown in Figures 9 to 29.

Table 16. XRPD copper radiation results of other salts for 3-DM.

3-DM Salt	Pos. [°2θ]	Height [cts]	FWHM Left [°2θ]	d-spacing [Å]	Rel. Int. [%]
HCl Type A (810397-04-B1) (Fig 9)	6.01	5471.24	0.1023	14.69	100.00
	10.11	36.23	0.3070	8.75	0.66
	12.03	770.84	0.0768	7.36	14.09
	13.19	139.55	0.1023	6.71	2.55
	13.98	136.40	0.1023	6.34	2.49
	14.69	342.73	0.1279	6.03	6.26
	16.43	313.97	0.1279	5.39	5.74
	17.76	180.39	0.1279	5.00	3.30
	18.08	665.54	0.1023	4.91	12.16
	18.42	220.46	0.1023	4.82	4.03
	19.58	254.85	0.1535	4.53	4.66
	20.48	201.58	0.1535	4.34	3.68
	21.07	63.97	0.1535	4.22	1.17
	21.63	288.45	0.1279	4.11	5.27
	22.63	88.17	0.2047	3.93	1.61
	24.22	141.32	0.2047	3.67	2.58
	24.53	92.47	0.1279	3.63	1.69
	25.64	51.08	0.3070	3.47	0.93
	26.98	84.57	0.1535	3.30	1.55

	27.98	139.09	0.2047	3.19	2.54
	28.53	84.66	0.1535	3.13	1.55
	29.41	105.78	0.2047	3.04	1.93
	31.46	75.16	0.2047	2.84	1.37
HCl Type B (810397-04-D1) (Fig 10)	6.40	322.11	0.1023	13.82	100.00
	12.73	75.53	0.3070	6.95	23.45
	14.95	61.29	0.1535	5.93	19.03
	15.90	225.46	0.0768	5.57	69.99
	17.44	123.71	0.1535	5.09	38.40
	19.22	96.49	0.2047	4.62	29.95
	20.16	115.59	0.1535	4.40	35.89
	23.68	64.14	0.3070	3.76	19.91
Phosphate Type A (810397-04-D3) (Fig 11)	5.89	2174.99	0.1791	15.02	100.00
	8.88	92.40	0.1535	9.96	4.25
	13.14	93.55	0.3070	6.74	4.30
	13.98	214.05	0.2047	6.34	9.84
	15.24	53.48	0.3070	5.81	2.46
	19.57	146.86	0.4093	4.54	6.75
	22.52	113.58	0.3070	3.95	5.22
	25.07	102.83	0.4093	3.55	4.73
	26.71	42.05	0.6140	3.34	1.93
Maleate Type A (810397-04-C4 (Fig 12)	7.05	369.14	0.1023	12.54	100.00
	8.06	325.32	0.1535	10.97	88.13
	10.58	119.23	0.1535	8.36	32.30
	16.23	145.05	0.3070	5.46	39.30
	17.75	188.43	0.1535	5.00	51.04
	19.54	145.00	0.1791	4.54	39.28
	21.57	65.58	0.8187	4.12	17.77

	22.63	148.06	0.2558	3.93	40.11
	27.58	50.71	0.3070	3.23	13.74
L-Tartrate Type A (810397-04-B5) (Fig 13)	6.78	3012.28	0.1279	13.04	100.00
	11.89	233.63	0.1279	7.44	7.76
	14.46	282.46	0.1023	6.12	9.38
	15.51	138.47	0.2047	5.71	4.60
	16.60	537.37	0.1023	5.34	17.84
	18.62	302.16	0.1279	4.77	10.03
	19.35	141.27	0.1535	4.59	4.69
	20.36	131.67	0.1535	4.36	4.37
	21.11	101.91	0.2047	4.21	3.38
	23.82	202.50	0.3070	3.74	6.72
	24.65	234.43	0.1279	3.61	7.78
Glycolate Type B (810397-04-C8 (Fig 14)	5.31	559.43	0.2047	16.65	100.00
	6.81	207.66	0.2047	12.98	37.12
	8.98	63.50	0.3070	9.85	11.35
	10.99	184.46	0.1023	8.05	32.97
	11.32	127.38	0.1535	7.82	22.77
	13.53	211.79	0.2047	6.55	37.86
	17.32	66.45	0.4093	5.12	11.88
	20.05	115.12	0.2558	4.43	20.58
	23.88	60.26	0.5117	3.73	10.77
Adipate Type A (810397-04-A12) (Fig 15)	6.64	375.12	0.0768	13.31	40.46
	7.07	232.49	0.1279	12.51	25.08
	9.73	294.49	0.1023	9.09	31.76
	10.61	269.69	0.1023	8.34	29.09
	11.29	103.80	0.4093	7.83	11.20

	12.10	167.05	0.1535	7.31	18.02
	13.40	82.24	0.3070	6.61	8.87
	15.16	135.71	0.1535	5.84	14.64
	15.84	309.04	0.1023	5.59	33.33
	16.70	272.88	0.1791	5.31	29.43
	18.23	144.67	0.1535	4.87	15.60
	19.44	68.42	0.5117	4.57	7.38
	21.38	927.12	0.1279	4.16	100.00
	21.99	187.93	0.2047	4.04	20.27
	23.91	227.45	0.1535	3.72	24.53
	24.29	372.75	0.1535	3.66	40.21
	26.27	183.35	0.2047	3.39	19.78
Adipate Type B (810397-04-C12) (Fig 16)	6.64	1344.11	0.1023	13.32	100.00
	9.74	486.46	0.1023	9.09	36.19
	10.44	314.32	0.1023	8.47	23.39
	11.64	149.61	0.0768	7.60	11.13
	15.66	342.03	0.1023	5.66	25.45
	16.47	739.71	0.0768	5.38	55.03
	19.57	118.26	0.1279	4.54	8.80
	20.47	224.99	0.1535	4.34	16.74
	21.39	691.20	0.1279	4.15	51.42
	22.81	141.34	0.1023	3.90	10.52
	23.36	88.24	0.1535	3.81	6.56
	23.98	177.26	0.1279	3.71	13.19
	27.14	52.73	0.2047	3.29	3.92
Acetate Type A (810397-04-D13) (Fig 17)	7.10	841.05	0.1023	12.46	57.38
	10.14	1465.74	0.1535	8.72	100.00
	11.30	1153.88	0.1023	7.83	78.72
	13.16	873.36	0.1023	6.73	59.59
	14.20	294.00	0.1023	6.24	20.06

	15.72	802.23	0.0768	5.64	54.73
	16.04	708.13	0.1791	5.52	48.31
	18.03	991.88	0.1023	4.92	67.67
	18.22	503.56	0.0768	4.87	34.36
	19.55	862.57	0.1279	4.54	58.85
	20.87	380.61	0.1279	4.26	25.97
	21.02	396.45	0.0768	4.23	27.05
	21.35	158.28	0.1535	4.16	10.80
	22.72	540.35	0.2047	3.91	36.87
	23.39	274.39	0.1023	3.80	18.72
	25.33	632.04	0.1535	3.52	43.12
	27.74	231.17	0.1535	3.22	15.77
	31.77	130.77	0.2047	2.82	8.92
	34.12	26.88	0.6140	2.63	1.83
Malonate Type A (810397-04-B17) (Fig 18)	7.51	1325.29	0.1279	11.77	100.00
	8.74	350.43	0.1279	10.12	26.44
	11.56	191.83	0.1535	7.66	14.47
	13.70	991.43	0.1279	6.46	74.81
	15.13	156.86	0.2047	5.85	11.84
	15.78	522.99	0.2047	5.62	39.46
	17.39	286.23	0.1535	5.10	21.60
	18.01	223.95	0.1023	4.93	16.90
	18.70	558.37	0.1023	4.75	42.13
	19.32	229.70	0.1791	4.59	17.33
	20.10	131.16	0.1535	4.42	9.90
	20.51	145.23	0.1535	4.33	10.96
	21.15	145.63	0.1535	4.20	10.99
	22.82	507.87	0.1791	3.90	38.32
	23.28	176.20	0.1535	3.82	13.30
	24.28	303.51	0.2558	3.67	22.90
	24.95	223.65	0.2047	3.57	16.88
	25.54	210.44	0.2047	3.49	15.88

	26.81	197.95	0.2047	3.33	14.94
	27.48	153.28	0.2558	3.25	11.57
	28.44	40.35	0.3070	3.14	3.04
	29.85	53.29	0.3070	2.99	4.02
	37.80	24.41	0.6140	2.38	1.84
Malonate Type B (810397-04-D17) (Fig 19)	7.393905	911.630700	0.153504	11.95636	100.00
	8.705001	129.308500	0.307008	10.15829	14.18
	11.443480	133.246600	0.204672	7.73277	14.62
	13.679360	261.793200	0.153504	6.47348	28.72
	15.786770	298.816000	0.127920	5.61375	32.78
	17.205240	247.311000	0.281424	5.15399	27.13
	18.427380	260.767600	0.076752	4.81483	28.60
	18.675700	426.904700	0.076752	4.75137	46.83
	18.991190	238.203700	0.204672	4.67314	26.13
	20.427050	151.108200	0.153504	4.34779	16.58
	21.201710	85.153030	0.307008	4.19065	9.34
	22.711740	515.768900	0.153504	3.91533	56.58
	24.125140	246.694100	0.179088	3.68905	27.06
	24.744300	144.154700	0.204672	3.59813	15.81
	27.009250	84.520030	0.614016	3.30132	9.27
Gentisate Type A (810397-04-B18) (Fig 20)	8.06	1582.31	0.1023	10.97	100.00
	10.03	225.27	0.1023	8.82	14.24
	10.66	309.03	0.1023	8.30	19.53
	11.98	181.06	0.1023	7.39	11.44
	12.65	221.15	0.1023	7.00	13.98

	13.91	470.72	0.1023	6.37	29.75
	14.65	359.01	0.1023	6.05	22.69
	15.94	317.32	0.1279	5.56	20.05
	16.34	1170.18	0.0768	5.42	73.95
	17.20	203.02	0.0768	5.16	12.83
	17.45	274.73	0.0768	5.08	17.36
	17.73	474.84	0.1023	5.00	30.01
	18.88	881.85	0.1023	4.70	55.73
	19.71	210.08	0.1279	4.50	13.28
	21.15	398.05	0.0768	4.20	25.16
	21.37	513.08	0.0768	4.16	32.43
	22.60	130.17	0.1535	3.93	8.23
	22.94	91.47	0.1535	3.88	5.78
	23.81	126.46	0.1279	3.74	7.99
	24.38	527.21	0.1791	3.65	33.32
	24.79	510.89	0.1535	3.59	32.29
	26.03	108.51	0.1791	3.42	6.86
	26.93	229.84	0.1023	3.31	14.53
	27.55	70.69	0.2047	3.24	4.47
	29.51	51.86	0.3070	3.03	3.28
	38.37	30.75	0.3070	2.35	1.94
Gentisate Type B (810397-04-C18) (Fig 21)	8.26	2365.32	0.0768	10.71	100.00
	9.94	322.72	0.1023	8.90	13.64
	10.52	437.18	0.1023	8.41	18.48
	11.94	234.71	0.1023	7.41	9.92
	12.76	419.68	0.1023	6.94	17.74
	13.68	403.02	0.1023	6.47	17.04
	14.16	890.33	0.1279	6.26	37.64
	14.67	349.21	0.0768	6.04	14.76
	15.84	799.58	0.1279	5.59	33.80
	16.50	1311.53	0.1023	5.37	55.45
	16.86	157.48	0.1535	5.26	6.66

	17.30	218.60	0.1535	5.13	9.24
	17.88	510.62	0.1023	4.96	21.59
	18.67	1101.93	0.1023	4.75	46.59
	19.63	285.81	0.1279	4.52	12.08
	21.43	884.16	0.1279	4.15	37.38
	21.84	741.10	0.1279	4.07	31.33
	22.88	136.29	0.1535	3.89	5.76
	23.87	253.19	0.1279	3.73	10.70
	24.31	316.14	0.1791	3.66	13.37
	24.91	578.38	0.1279	3.57	24.45
	25.20	287.42	0.1279	3.53	12.15
	25.69	125.67	0.1535	3.47	5.31
	26.13	157.32	0.1535	3.41	6.65
	26.99	187.12	0.1279	3.30	7.91
	29.14	79.31	0.1535	3.06	3.35
	29.56	121.70	0.1279	3.02	5.15
	31.53	58.79	0.1535	2.84	2.49
	34.09	65.23	0.2047	2.63	2.76
	36.28	90.02	0.2558	2.48	3.81
Edisylate Type A (810397-04-A19) (Fig 22)	6.84	1772.74	0.1023	12.91	100.00
	9.73	239.23	0.1023	9.09	13.49
	10.89	62.52	0.3070	8.12	3.53
	13.15	358.17	0.1023	6.73	20.20
	13.53	553.52	0.0768	6.54	31.22
	15.14	295.09	0.1023	5.85	16.65
	15.89	108.43	0.1535	5.58	6.12
	16.52	192.88	0.2558	5.37	10.88
	17.50	146.03	0.4093	5.07	8.24
	19.72	254.81	0.1535	4.50	14.37
	20.13	335.61	0.0768	4.41	18.93
	21.34	387.88	0.0768	4.16	21.88
	21.57	453.85	0.1279	4.12	25.60

	23.26	306.91	0.1023	3.82	17.31
	23.46	366.60	0.0768	3.79	20.68
	23.89	332.69	0.1023	3.72	18.77
	24.40	258.49	0.1023	3.65	14.58
	25.43	229.38	0.1279	3.50	12.94
	27.17	125.77	0.1535	3.28	7.09
	31.33	63.59	0.2047	2.86	3.59
Edisylate Type B (810397-04-C19) (Fig 23)	6.47	190.15	0.1279	13.67	100.00
	12.89	39.88	0.6140	6.87	20.97
	14.75	72.81	0.3070	6.01	38.29
	17.08	67.53	0.3070	5.19	35.51
	20.92	30.55	0.4093	4.25	16.06
	24.18	56.49	0.3070	3.68	29.71
Benzoate Type A (810397-04-D20) (Fig 24)	6.88	137.63	0.1535	12.84	59.95
	7.38	112.52	0.2047	11.98	49.02
	9.02	50.52	0.3070	9.81	22.01
	10.55	90.55	0.1535	8.39	39.45
	11.59	40.77	0.4093	7.64	17.76
	13.23	229.56	0.1279	6.69	100.00
	14.60	113.18	0.2047	6.07	49.30
	15.74	118.47	0.1535	5.63	51.61
	16.28	105.42	0.3070	5.45	45.92
	17.66	169.37	0.1279	5.02	73.78
	18.44	50.63	0.3070	4.81	22.06
	20.45	148.57	0.2047	4.34	64.72
	21.30	85.90	0.4093	4.17	37.42
	23.38	86.26	0.4093	3.80	37.58
	23.93	81.57	0.1535	3.72	35.53
	25.07	48.36	0.3070	3.55	21.07

Esylate Type A (810397-04-C21) (Fig 25)	6.67	3667.35	0.1279	13.25	100.00
	8.13	218.87	0.1023	10.87	5.97
	9.97	111.78	0.1535	8.87	3.05
	11.16	521.74	0.1023	7.93	14.23
	12.98	556.17	0.1023	6.82	15.17
	13.37	183.35	0.1279	6.62	5.00
	14.85	292.74	0.1023	5.97	7.98
	15.21	183.06	0.1279	5.82	4.99
	16.42	1158.26	0.1279	5.40	31.58
	17.12	696.45	0.1279	5.18	18.99
	18.30	711.73	0.1023	4.85	19.41
	18.65	450.76	0.1279	4.76	12.29
	19.50	357.56	0.1279	4.55	9.75
	20.10	485.39	0.1535	4.42	13.24
	20.87	105.87	0.1535	4.26	2.89
	21.38	293.58	0.1279	4.16	8.01
	22.59	374.76	0.1279	3.94	10.22
	23.67	260.88	0.1279	3.76	7.11
	23.97	166.36	0.1535	3.71	4.54
	24.70	141.85	0.2558	3.60	3.87
	25.69	431.64	0.1535	3.47	11.77
	26.12	157.28	0.1023	3.41	4.29
	26.89	248.50	0.2558	3.32	6.78
	28.25	134.62	0.1535	3.16	3.67
	29.00	52.72	0.2047	3.08	1.44
	32.62	57.06	0.3070	2.75	1.56
	33.89	44.97	0.2047	2.65	1.23
	34.69	41.22	0.2047	2.59	1.12
	38.74	25.68	0.3070	2.32	0.70
Besylate Type A (810397-04-D23) (Fig 26)	5.49	254.56	0.0768	16.11	17.38

	6.98	1464.97	0.1023	12.67	100.00
	9.92	936.45	0.1023	8.92	63.92
	11.00	299.09	0.3070	8.05	20.42
	11.78	600.93	0.1023	7.51	41.02
	12.35	225.11	0.1023	7.17	15.37
	13.09	838.56	0.1023	6.77	57.24
	13.77	192.66	0.1279	6.43	13.15
	15.01	52.71	0.3070	5.90	3.60
	16.47	286.20	0.1023	5.38	19.54
	17.24	244.41	0.1023	5.14	16.68
	17.85	294.68	0.1023	4.97	20.12
	18.48	503.28	0.1023	4.80	34.35
	20.88	328.01	0.1535	4.25	22.39
	21.24	245.26	0.1023	4.18	16.74
	21.76	373.87	0.2558	4.08	25.52
	23.72	372.74	0.1279	3.75	25.44
	25.20	155.89	0.1535	3.53	10.64
	26.46	175.45	0.3070	3.37	11.98
	27.21	131.25	0.1535	3.28	8.96
	27.98	119.10	0.3070	3.19	8.13
Tosylate Type A (810397-04-B24) (Fig 27)	6.29	7162.69	0.1023	14.06	100.00
	9.42	102.41	0.1535	9.39	1.43
	12.09	134.52	0.1535	7.32	1.88
	13.63	67.80	0.1535	6.49	0.95
	15.27	160.36	0.1535	5.80	2.24
	15.92	161.50	0.1023	5.57	2.25
	16.42	76.74	0.1535	5.40	1.07
	17.48	105.68	0.1535	5.07	1.48
	18.93	524.42	0.1023	4.69	7.32
	19.88	109.82	0.2047	4.47	1.53
	20.37	119.82	0.2047	4.36	1.67
	20.91	135.14	0.1535	4.25	1.89

	23.89	119.27	0.2047	3.72	1.67
	25.30	93.55	0.1535	3.52	1.31
	26.82	26.25	0.6140	3.32	0.37
	28.53	118.19	0.1791	3.13	1.65
Oxalate Type A (810397-04-A25) (Fig 28)	6.49	6650.19	0.1535	13.62	100.00
	7.93	100.70	0.3070	11.15	1.51
	8.92	181.89	0.1535	9.91	2.74
	10.32	85.94	0.3070	8.57	1.29
	11.33	103.40	0.2047	7.81	1.55
	12.92	167.51	0.3070	6.85	2.52
	13.68	269.32	0.2047	6.47	4.05
	15.75	162.38	0.3070	5.63	2.44
	19.45	453.55	0.1535	4.56	6.82
Oxalate Type B (810397-04-B25) (Fig 29)	6.47	2862.81	0.1023	13.66	100.00
	9.90	151.67	0.1023	8.94	5.30
	10.63	183.15	0.1279	8.32	6.40
	12.96	139.44	0.1791	6.83	4.87
	13.95	84.95	0.4093	6.35	2.97
	16.00	90.27	0.1535	5.54	3.15
	16.72	80.26	0.3070	5.30	2.80
	19.50	206.28	0.1279	4.55	7.21
	19.93	157.42	0.2047	4.46	5.50
	21.32	75.05	0.3070	4.17	2.62
	22.28	140.88	0.1535	3.99	4.92
	24.25	104.26	0.3070	3.67	3.64
	26.75	62.82	0.3070	3.33	2.19
	27.65	59.74	0.3070	3.23	2.09

[0319] Example 3. Re-preparation and Characterization of 3-Deuteromitragynine Salt Leads

[0320] Based on the solid-state characterization of 3-DM salts in **Example 2**, Fumarate Type A, Glycolate Type A, L-Lactate Type A, Succinate Type A and Mesylate Type A were selected as salt leads of 3-DM for scale up and further evaluation. Detailed re-preparation procedures of the salt leads are listed in Table 17. XRPD comparison results confirmed all salt leads were generated on scale up. Characterization results of re-prepared salt leads are summarized in Table 18.

Table 17. Preparation procedures for 3-DM salt leads.

3-DM Salt (Sample ID)	Procedure
Fumarate Type A (810397-28-B1)	1. Weigh 199.4 mg of free base (810397-01-A) and 58.9 mg of fumaric acid into a 20-mL glass vial. 2. Add 5 mL IPA/H_2O (4:1, v/v) to form a suspension. Magnetically stir (~1000 rpm) at RT for 39 hrs. 3. Isolate suspension by centrifugation (10000 rpm, 2 min). Dry solids at 50 ºC under vacuum for ~7 hrs. 4. Collect solids, a total of 232.8 mg of material was obtained (Yield: 90.1%).
Fumarate Type A (810397-36-B1)	1. Weigh 200.6 mg of free base (810397-01-A) and 59.1 mg of fumaric acid into a 20-mL glass vial. 2. Add 5 mL IPA/H_2O (4:1, v/v) to form a suspension. Magnetically stir (~1000 rpm) at RT for 48 hrs. 3. Isolate solids by centrifugation (10000 rpm, 2 min). Dry solids at 50 ºC under vacuum for ~7 hrs. Collect solids.
Glycolate Type A (810397-28-B2)	1. Weigh 300.2 mg of free base (810397-01-A) and 58.2 mg of glycolic acid into a 20-mL glass vial. 2. Add 5 mL IPA/H_2O (4:1, v/v) to form a suspension. Magnetically stir (~1000 rpm) at RT for 39 hrs. 3. Isolate solids by centrifugation (10000 rpm, 2 min). Dry solids at 50 ºC under vacuum for ~7 hrs. 4. Collect solids, a total of 230.5 mg of material was obtained (Yield: 64.3%).

3-DM Salt (Sample ID)	Procedure
Glycolate Type A (810397-36-B2)	1. Weigh 299.8 mg of free base (810397-01-A) and 58.3 mg of glycolic acid into a 20-mL glass vial. 2. Add 5 mL IPA/H_2O (4:1, v/v) to form a suspension. Magnetically stir (~1000 rpm) at RT for 48 hrs. 3. Isolate solids by centrifugation (10000 rpm, 2 min). Dry solids at 50 ºC under vacuum for ~7 hrs. Collect solids.
L-Lactate Type A (810397-28-B3)	1. Weigh 55.0 mg of L-lactic acid into a 20-mL glass vial, add 5 mL IPA/H_2O (4:1, v/v) to dissolve acid. 2. Weigh 200.3 mg of free base (810397-01-A) into the acid solution. Magnetically stir (~1000 rpm) at RT for 39 hrs. 3. Isolate solids by centrifugation (10000 rpm, 2 min). Dry solids at 50 ºC under vacuum for ~7 hrs. 4. Collect solids, a total of 138.3 mg of material was obtained (Yield: 54.2%).
L-Lactate Type A (810397-36-B3)	1. Weigh 83.1 mg of L-Lactic acid into a 20-mL glass vial, add 5 mL IPA/H_2O (4:1, v/v) to dissolve acid. 2. Weigh 300.3 mg of free base (810397-01-A) into the acid solution. Magnetically stir (~1000 rpm) at RT for 48 hrs. 3. Isolate solids by centrifugation (10000 rpm, 2 min). Dry solids at 50 ºC under vacuum for ~7 hrs. Collect solids.
Succinate Type A (810397-28-B4)	1. Weigh 199.6 mg of free base (810397-01-A) and 60.3 mg of succinic acid into a 20-mL glass vial. 2. Add 5 mL IPA/H_2O (4:1, v/v) to form a suspension. Magnetically stir (~1000 rpm) at RT for 39 hrs. 3. Isolate solids by centrifugation (10000 rpm, 2 min). Dry solids at 50 ºC under vacuum for ~7 hrs. 4. Collect solids, a total of 179.2 mg of material was obtained (Yield: 68.9%).
Succinate Type A (810397-36-B4)	1. Weigh 200.5 mg of free base (810397-01-A) and 60.7 mg of succinic acid into a 20-mL glass vial.

3-DM Salt (Sample ID)	Procedure
	2. Add 5 mL IPA/H_2O (4:1, v/v) to form a suspension. Magnetically stir (~1000 rpm) at RT for 48 hrs. 3. Isolate solids by centrifugation (10000 rpm, 2 min). Dry solids at 50 ºC under vacuum for ~7 hrs. Collect solids.
Mesylate Type A (810397-28-B5)	1. Weigh 79.2 mg of Methanesulfonic acid into a 20-mL glass vial, add 5 mL EtOAc to dissolve acid. 2. Weigh 326.6 mg of free base (810397-01-A) into the acid solution. Magnetically stir (~1000 rpm) at RT for 39 hrs. 3. Isolate solids by centrifugation (10000 rpm, 2 min). Dry solids at 50 ºC under vacuum for ~7 hrs. 4. Collect solids, a total of 378.5 mg of material was obtained (Yield: 93.3%).

Table 18. Characterization results of re-prepared 3-DM salt leads.

3-DM Salt (Sample ID)	TGA weight loss (%)	DSC endotherm (ºC, onset)	Molar ratio (acid/base)	Water content by KF (%)	Purity (area%)	3-DCR (area%)
Fumarate Type A (810397-28-B1)	0.3 (to 150 ºC)	256.1	1:1	0.71	96.89	2.51
Fumarate Type A (810397-36-B1)	1.6 (to 150 ºC)	255.2	1:1	--	97.03	2.45
Glycolate Type A (810397-28-B2)	0.5 (to 150 ºC)	212.1	1:1	0.43	98.69	1.10
Glycolate Type A (810397-36-B2)	1.1 (to 150 ºC)	212.1	1:1	--	97.53	1.93
L-Lactate Type A (810397-28-B3)	0.9 (to 150 ºC)	211.6	1:1	0.31	98.18	1.50

3-DM Salt (Sample ID)	TGA weight loss (%)	DSC endotherm (ºC, onset)	Molar ratio (acid/base)	Water content by KF (%)	Purity (area%)	3-DCR (area%)
L-Lactate Type A (810397-36-B3)	1.9 (to 150 ºC)	224.2	0.9:1	--	98.82	0.94
Succinate Type A (810397-28-B4)	2.3 (to 150 ºC)	198.2*, 202.3*	1:1	0.41	97.92	1.87
Succinate Type A (810397-36-B4)	1.1 (to 150 ºC)	198.0*, 202.6*	1.1:1	--	97.89	1.81
Mesylate Type A (810397-28-B5)	1.3 (to 150 ºC)	266.4	1:1	0.50	96.48	2.65

[0321] *: Peak temperature.

[0322] **Example 4: Solubility of 3-Deuteromitragynine Salt Leads**

[0323] Solubility of 3-DM Fumarate Type A, Glycolate Type A, L-Lactate Type A, Succinate Type A, and Mesylate Type A was measured after 4 hrs at 37 ºC in water and the following bio-relevant media: simulated gastric fluid (SGF), fasting-state simulated intestinal fluid (FaSSIF), and fed-state simulated intestinal fluid (FeSSIF). Amorphous free base was also assessed as a comparison.

[0324] **Preparation of SGF.** Weighed 49.5 mg of NaCl and 25.4 mg of Triton X-100 into a 100-mL volumetric flask. Added appropriate volume of purified water and sonicated until all solids were completely dissolved. Added about 1.632 mL of aq. HCl solution (1 M) and sufficient purified water to achieve the target volume and adjust to pH 1.8. The pH value was checked with a pH meter and found to be 1.83.

[0325] **Preparation of FaSSIF Dissolving Buffer.** Weighed 340.8 mg of NaH_2PO_4, 43.0 mg of NaOH, and 619.6 mg of NaCl into a 100-mL volumetric flask. Added appropriate volume of purified water and sonicated until all solids were completely dissolved. Added sufficient purified water to achieve the target volume and adjust to pH 6.5. The pH value was checked with a pH meter and found to be 6.54.

[0326] Preparation of FaSSIF. Weighed 110.4 mg of simulated intestinal fluid (SIF) powder into a 50-mL volumetric flask. Added appropriate volume of FaSSIF dissolving buffer and sonicated until SIF powder was completely dissolved. Then diluted to volume with FaSSIF dissolving buffer and mixed well. The FaSSIF solution was equilibrated to RT for 2 hrs before use.

[0327] Preparation of FeSSIF Dissolving Buffer. Weighed 0.82 mL of glacial acetic acid, 404.9 mg of NaOH, and 1188.2 mg of NaCl into a 100-mL volumetric flask. Added appropriate volume of purified water and sonicated until all solids were completely dissolved. Added sufficient purified water to achieve the target volume and adjust to pH 5.0. The pH value was checked with a pH meter and found to be 4.96.

[0328] Preparation of FeSSIF. Weighed 559.6 mg of simulated intestinal fluid (SIF) powder into a 50-mL volumetric flask. Added appropriate volume of FeSSIF dissolving buffer and sonicated until SIF powder was completely dissolved. Then diluted to volume with FeSSIF dissolving buffer and mixed well. The FeSSIF solution was equilibrated to RT for 2 hrs before use.

[0329] Solubility Measurement. Solids were suspended in selected media at a loading of 10 mg/mL (calculated based on free base). The suspensions were agitated on a rolling incubator at 25 rpm and 37 ºC, prior to sampling at 4 hrs. At this time, suspensions were separated by centrifugation (~10,000 rpm, 37 ºC, 3 min) and the supernatants were filtered through 0.45 µm PTFE membranes. The pH of the filtrates was determined and they were analyzed by HPLC to determine solubility. Residual solids were analyzed by XRPD to determine their crystal form. Results are summarized in Table 19.

[0330] All salt leads showed higher solubility than amorphous free base in all media, except for Fumarate Type A in SGF. Glycolate Type A showed the highest solubility in SGF. XRPD overlay of the residual solids from solubility samples showed that no form change occurred for any of the tested salt forms or amorphous free base in any medium after 4 h. However, for Fumarate Type A, Glycolate Type A, and amorphous free base, an additional peak at 8.9º was observed in the XRPD spectra of the residual solids from FeSSIF experiments, which may result from sodium acetate.

Table 19. Summary of solubility results for 3-DM salts.

Salt lead	H_2O			SGF			FaSSIF			FeSSIF		
(810397-)	S	pH	FC	S	pH	FC	S	pH	FC	S	pH	FC
Fumarate Type A (36-B1)	0.30	3.4	No	1.07	1.9	No	2.35	5.6	No	1.49	4.9	No
Glycolate Type A (36-B2)	5.32	5.2	No	7.43	3.2	No	1.16	6.2	No	2.16	5.0	No
L-Lactate Type A (36-B3)	1.54	5.4	No	6.89	2.9	No	1.03	6.2	No	2.09	5.0	No
Succinate Type A (36-B4)	2.58	4.5	No	6.78	3.2	No	4.16	5.3	No	2.67	4.9	No
Mesylate Type A (28-B5)	13.1*	3.2	No	7.21	1.8	--	2.41	5.8	No	2.17	4.8	No
Amorphous free base (01-A)	0.013	5.9	No	5.32	5.5	No	0.65	6.6	No	0.008	5.1	No

[0331] S: Solubility, mg/mL. FC: Form change.

[0332] *: Additional 23.2 mg of salt was added because clear solution was observed at 10 mg/mL.

[0333] --: XRPD test was not performed because limited amount of solid was obtained after solubility test.

[0334] **Example 5: Hygroscopicity of 3-Deuteromitragynine Salt Leads.**

[0335] To investigate the solid form stability as a function of humidity, DVS isotherm plots of 3-DM Fumarate Type A, Glycolate Type A, L-Lactate Type A, Succinate Type A, and Mesylate Type A were collected at 25 ºC between 0 and 95% relative humidity (RH). After DVS testing, samples were characterized by XRPD to check for form conversion. The results are summarized in Table 20.

[0336] For Fumarate Type A, Glycolate Type A, and L-Lactate Type A, water uptake at 80% RH was 0.41%, 0.098%, and 0.31%, respectively. No form change was observed by XRPD after DVS test for any of these 3 salts.

[0337] For Succinate Type A and Mesylate Type A, water uptake at 80% RH was 0.16% and 0.49%, respectively. In these samples, some residual water in initial samples was removed during desorption cycle, which resulted in a negative mass change. No form change was observed by XRPD after DVS test for either of these salts.

Table 20. DVS results for 3-DM salts.

3-DM Salt(Sample ID)	Water uptake at 80% RH	Form change
Fumarate Type A (810397-28-B1)	0.41%	No
Glycolate Type A (810397-28-B2)	0.098%	No
L-Lactate Type A (810397-28-B3)	0.31%	No
Succinate Type A (810397-28-B4)	0.16%	No
Mesylate Type A (810397-28-B5)	0.49%	No

[0338] **Example 6: Physical and Chemical Stability of 3-Deuteromitragynine Salt Leads**

[0339] The physical and chemical stability of 3-DM Fumarate Type A, Glycolate Type A, L-Lactate Type A, Succinate Type A, and Mesylate Type A were evaluated under conditions of 25 ºC/60% RH and 40 ºC/75% RH for 1 week. Each sample was added into 3-mL glass vials, sealed by parafilm with several holes, and stored under one of the indicated test conditions. After one week, samples were taken for XRPD, KF, and HPLC purity tests. All the resulting characterization data are summarized in Table 21.

[0340] For all tested salt leads, no purity decrease (by HPLC) or form change (by XRPD) was observed after storage under either of the two conditions for one week. Impurity summaries for all salt leads as determined by HPLC are shown in Table 22 to Table 26. The peak at RRT = 0.90 corresponds to impurity 3-DCR. For Glycolate Type A and L-Lactate Type A, the area percentage of 3-DCR slightly decreased after storage under 25 ºC/60% RH for one week, while no significant change was observed for the other three salt leads.

Table 21. Summary of physicochemical stability results for 3-DM salts.

3-DM Salt (810397-)	**Initial**		**25 ºC/60% RH/1 week**			**40 ºC/75% RH/1 week**		
	Purity (area%)	KF (wt%)	Form change	Purity/Initial purity (%)	KF (wt%)	Form change	Purity/Initial purity (%)	KF (wt%)
Fumarate Type A (28-B1)	96.89	0.71	No	100.3	0.79	No	100.3	0.31
Glycolate Type A (28-B2)	98.69	0.43	No	100.4	0.28	No	100.4	0.26
L-Lactate Type A (28-B3)	98.18	0.31	No	100.7	0.64	No	100.1	0.67
Succinate Type A (28-B4)	97.92	0.41	No	100.0	0.39	No	99.9	0.35
Mesylate Type A (28-B5)	96.48	0.50	No	100.3	0.54	No	100.3	0.54

Table 22. Impurity summary of 3-DM Fumarate Type A.

#Peak	RRT	Area (%)		
		Initial	25 ºC/60% RH	40 ºC/75% RH
1	0.68	0.14	0.08	0.09
2	0.70	0.26	0.18	0.20
3	0.90	2.51	2.42	2.45
4	0.93	0.15	0.10	0.11
5	0.94	0.05	<0.05	<0.05
6	**1.00**	**96.89**	**97.21**	**97.16**

Table 23. Impurity summary of 3-DM Glycolate Type A.

#Peak	RRT	Area (%)		
		Initial	25 ºC/60% RH	40 ºC/75% RH
1	0.68	0.06	<0.04	<0.04
2	0.70	0.09	0.04	0.04
3	0.90	1.10	0.74	0.77
4	0.93	0.06	0.12	0.05
5	**1.00**	**98.69**	**99.10**	**99.13**

Table 24. Impurity summary of 3-DM L-Lactate Type A.

#Peak	RRT	Area (%)		
		Initial	25 ºC/60%RH	40 ºC/75%RH
1	0.68	0.10	<0.05	0.07
2	0.70	0.15	0.05	0.11
3	0.90	1.50	0.96	1.46
4	0.93	0.07	0.12	0.07
5	**1.00**	**98.18**	**98.87**	**98.28**

Table 25. Impurity summary of 3-DM Succinate Type A.

#Peak	RRT	Area (%)		
		Initial	25 ºC/60%RH	40 ºC/75%RH
1	0.68	0.05	0.04	0.05

#Peak	RRT	Area (%)		
		Initial	25 ºC/60%RH	40 ºC/75%RH
2	0.70	0.09	0.08	0.11
3	0.90	1.87	1.86	1.96
4	0.93	0.07	0.07	0.08
5	**1.00**	**97.92**	**97.96**	**97.80**

Table 26. Impurity summary of 3-DM Mesylate Type A.

#Peak	RRT	Area (%)		
		Initial	25 ºC/60%RH	40 ºC/75%RH
1	0.68	0.23	0.16	0.15
2	0.70	0.36	0.27	0.27
3	0.90	2.65	2.61	2.62
4	0.93	0.17	0.11	0.11
5	0.94	0.06	<0.05	<0.05
6	**1.00**	**96.48**	**96.79**	**96.79**
7	1.10	<0.05	0.06	0.07

[0341] **Example 7: Scale-up of 3-Deuteromitragynine Glycolate Type A**

[0342] Based on evaluation results of salt leads, 3-DM Glycolate Type A was selected as a favorable solid form. To select a suitable solvent for scale up of Glycolate Type A with higher yield, approximate solubility of 3-DM Glycolate Type A and 3-DM free base was measured, with the results summarized in Table 27. Based on these findings, isopropyl alcohol (IPA) was selected as the solvent of choice since 3-DM free base showed high solubility and 3-DM Glycolate Type A showed low solubility in IPA. A total of three batches of Glycolate Type A were prepared using solution crystallization methods. In the first batch (100 mg scale), pure IPA was used as solvent. The purity of the product was 97.56% (area percentage), which was lower than the sample obtained from screening. Since a sample obtained from IPA/H_2O during screening showed higher purity, a small amount of water was added to the crystallization solvent in the latter two bathes. Characterization results for the 3 batches are summarized in Table 28. Detailed preparation procedures for the 3 scaled-up batches can be found below.

Table 27. Approximate solubility of 3-DM Glycolate Type A and 3-DM free base.

3-DM Glycolate Type A (810397-36-B2)		3-DM Free Base (810397-01-A)	
Solvent (v/v)	**Solubility (mg/mL)**	**Solvent (v/v)**	**Solubility (mg/mL)**
MeOH	S>36.0	MeOH	S>46.0
EtOH	6.7<S<10.0	EtOH	S>38.0
IPA	1.1<S<2.1	IPA	S>36.0
Acetone	6.7<S<10.0	Acetone	S>38.0
EtOAc	1.1<S<2.1	EtOAc	S>40.0
Acetone/H_2O (2:1)	S>38.0	--	--
Acetone/n-heptane (1:1)	1.2<S<2.4	Acetone/n-heptane (1:1)	S>42.0
EtOH/n-heptane (1:1)	2.0<S<4.0	--	--

Table 28. Summary data for scaled up batches of 3-DM Glycolate Type A.

Sample ID		810397-56-B	810397-62-B	810397-64-B
Scale		100 mg	1 g	5 g
Final solvent system		IPA	IPA/H_2O (19:1, v/v)	IPA/H_2O (19:1, v/v)
Yield (%)		88.3 (105.4 mg)	78.8 (945.4 mg)	83.7 (4989.7 mg)
XRPD		Glycolate Type A	Glycolate Type A	Glycolate Type A
TGA weight loss (%)		2.2 (to 150 ºC)	1.7 (to 150 ºC)	1.1 (to 150 ºC)
DSC endotherm (ºC, onset)		220.2	223.3	222.1
Molar ratio (acid/base)		1:1	1:1	1:1
Morphology		Needle-like	Needle-like	Rod-like
Purity (area%)	3-DM	97.56	98.89	98.81
	3-DCR	1.96	0.95	1.02

[0343] Preparation of 3-DM Glycolate Type A at 100-mg Scale

1) Weighed 28.1 mg of glycolic acid into a 3-mL glass vial. Added 724 µL IPA to dissolve solids for acid solution.
2) Weighed 99.9 mg of 3-DM free base into a 5-mL glass vial. Added 1.5 mL IPA to dissolve solids for free base solution.
3) Added 0.5 mL acid solution dropwise into free base solution while stirring. Precipitate was observed.
4) Magnetically stirred the suspension at RT for ~5 min.
5) Isolated solids by centrifugation (10,000 rpm, 2 min). Dried solids at RT under vacuum for ~17 hrs.
6) A total of 105.4 mg of white solids (needle-like crystals by PLM) were obtained (Yield: 88.3%).

[0344] Preparation of 3-DM Glycolate Type A at 1-g Scale

1) Weighed 197.8 mg of glycolic acid into a 20-mL glass vial. Added 5 mL IPA/H_2O (4:1, v/v) to dissolve solids for acid solution.
2) Weighed 1002.0 mg of 3-DM free base (810397-01-A) into a 50-mL glass vial. Added 15 mL IPA to dissolve solids for free base solution.
3) Added all acid solution dropwise into free base solution while stirring. Thick suspension was observed.
4) Magnetically stirred the suspension at RT for ~17 hrs.
5) Isolated solids by vacuum filtration. Dried solids at 50ºC under vacuum for ~3 hrs.
6) A total of 945.4 mg of white solids (needle-like crystals by PLM) were obtained (Yield: 78.8%).

[0345] Preparation of 3-DM Glycolate Type A at 5-g Scale

1) Weighed 961.5 mg of glycolic acid into a 50-mL glass vial. Added 22.5 mL IPA/H_2O (4:1, v/v) to dissolve solids for acid solution.
2) Weighed 5002.5 mg of 3-DM free base (810397-01-A) into a 100-mL glass vial. Added 67.5 mL IPA to dissolve solids for free base solution.
3) Overhead stirred (250 rpm) the free base solution at RT. Added all acid solution into free base solution over 2 hrs.
4) Overhead stirred the obtained suspension at RT for ~13 hrs.
5) Isolated solids by vacuum filtration. Dried solids at 50 ºC under vacuum for ~5 hrs.
6) A total of 4989.7 mg of white solids (rod-like crystals by PLM) were obtained (Yield: 83.7%).

[0346] **Example 8: Solvent Solubility of 3-Deuteromitragynine Glycolate Type A**

[0347] The approximate solubility of 3-DM Glycolate Type A was measured in 28 solvent systems at RT. Approximately 2 mg samples were added into 3-mL glass vials. The solvents in Table 29 were then added stepwise (50/50/200/700 µL) into the vials until the solids were dissolved visually or a total volume of 1 mL was reached. The solubility results summarized in Table 29 were used to guide the solvent selection for polymorph screening experiments.

Table 29. Approximate solubility of 3-DM Glycolate Type A at RT.

Solvent	Solubility (mg/mL)	Solvent (v/v)	Solubility (mg/mL)
MeOH	S>36.0	1,4-Dioxane	10.0<S<20.0
EtOH	6.7<S<10.0	ACN	2.2<S<3.7
IPA	1.1<S<2.1	DCM	2.2<S<3.7
Acetone	6.7<S<10.0	$CHCl_3$	1.1<S<2.2
MIBK	2.1<S<3.5	n-Heptane	S<1.1
EtOAc	1.1<S<2.1	Toluene	S<1.0
IPAc	0.9<S<1.8	DMSO	S>44.0
MTBE	S<1.0	DMF	S>40.0
THF	19.0<S<38.0	NMP	S>42.0
2-MeTHF	6.7<S<10.0	H_2O	S=5.3*

[0348] *Solubility was measured in water at 37 ºC after rolling for 24 hrs.

[0349] **Example 9: Polymorph Screening of 3-DM Glycolate**

Polymorph screening experiments were performed under 100 conditions using different solid transition or solution crystallization methods. Methods and results are summarized in Table 30, and further described in detail in the subsections below. A total of six crystal forms of 3-DM Glycolate were obtained as different polymorphic forms.

Table 30. Summary of methods and results from polymorph screening of 3-DM Glycolate.

Method	No. of Experiments	Results
Anti-solvent addition	12	Glycolate Type A, B, C, A+B, amorphous
Solid vapor diffusion	12	Glycolate Type A, C
Slurry at RT	25	Glycolate Type A
Slurry at 50 ºC	16	Glycolate Type A, C
Slow evaporation	7	Glycolate Type A, C, E
Slow cooling	6	Glycolate Type A, B, C
Liquid-vapor diffusion	12	Glycolate Type A, B, D, A+C
Polymer induced crystallization	6	Glycolate Type A, C, amorphous
Grind	4	Glycolate Type A
Total	**100**	Glycolate Type A-E, A+B, A+C, amorphous

[0350] The XRPD spectra of all polymorphic forms are shown in Figure 30 and the 2-theta peaks are shown below in Table 31.

Table 31. XRPD copper radiation results for 3-DM Glycolate Types.

3-DM Glycolate Salt	Pos. [°2θ]	Height [cts]	FWHM Left [°2θ]	d-spacing [Å]	Rel. Int. [%]
Type A	7.06	2109.45	0.0768	12.53	61.58
	10.11	3425.57	0.1535	8.75	100.00

	11.24	2400.98	0.1023	7.87	70.09
	13.21	492.74	0.1023	6.70	14.38
	14.09	571.47	0.1023	6.29	16.68
	15.05	86.69	0.1535	5.89	2.53
	15.96	1576.20	0.1791	5.55	46.01
	18.01	1283.35	0.1535	4.93	37.46
	18.53	290.83	0.1023	4.79	8.49
	19.21	399.41	0.0768	4.62	11.66
	19.49	1013.08	0.1535	4.55	29.57
	19.70	605.75	0.1023	4.51	17.68
	20.34	782.93	0.1023	4.37	22.86
	20.88	1902.18	0.1791	4.25	55.53
	22.57	1536.47	0.2558	3.94	44.85
	23.66	149.95	0.2047	3.76	4.38
	24.00	227.34	0.1023	3.71	6.64
	25.15	1552.35	0.1535	3.54	45.32
	27.62	546.00	0.1535	3.23	15.94
	29.53	93.41	0.2047	3.02	2.73
	30.14	254.98	0.0768	2.96	7.44
	31.61	252.54	0.1535	2.83	7.37
	34.14	65.70	0.3070	2.63	1.92
Type B	5.29	1520.48	0.0768	16.70	16.07
	5.68	9462.35	0.1279	15.57	100.00
	6.78	377.43	0.1023	13.04	3.99
	7.45	639.06	0.1535	11.87	6.75
	9.01	67.39	0.4093	9.81	0.71
	10.78	224.66	0.1023	8.21	2.37
	13.65	303.10	0.1279	6.49	3.20
	14.71	99.83	0.3070	6.02	1.05
	17.35	171.94	0.1535	5.11	1.82
	19.86	353.67	0.2047	4.47	3.74
	21.22	161.91	0.5117	4.19	1.71
	22.72	405.05	0.1535	3.91	4.28
	24.05	135.60	0.3070	3.70	1.43
	25.35	73.51	0.6140	3.51	0.78

	27.43	198.64	0.2558	3.25	2.10
Type C	6.03	19774.56	0.1023	14.66	100.00
	7.43	2767.27	0.1279	11.90	13.99
	10.57	212.10	0.1023	8.37	1.07
	10.99	381.18	0.1023	8.05	1.93
	11.39	351.48	0.1023	7.77	1.78
	12.81	162.61	0.1023	6.91	0.82
	13.65	396.34	0.1023	6.49	2.00
	14.22	429.31	0.1279	6.23	2.17
	15.66	124.37	0.1535	5.66	0.63
	16.30	524.59	0.1279	5.44	2.65
	17.61	185.19	0.3582	5.04	0.94
	18.07	541.80	0.1791	4.91	2.74
	18.84	460.35	0.1023	4.71	2.33
	20.08	786.88	0.1279	4.42	3.98
	21.15	258.68	0.3070	4.20	1.31
	21.54	366.62	0.0768	4.13	1.85
	22.57	161.98	0.4093	3.94	0.82
	22.87	370.26	0.1279	3.89	1.87
	24.16	1612.07	0.1535	3.68	8.15
	25.14	157.20	0.3070	3.54	0.79
	26.22	441.23	0.1535	3.40	2.23
	27.55	465.02	0.1535	3.24	2.35
	28.60	111.96	0.2047	3.12	0.57
	29.38	79.84	0.2047	3.04	0.40
	31.63	79.96	0.3070	2.83	0.40
	33.74	84.29	0.1535	2.66	0.43
	35.22	56.07	0.3070	2.55	0.28
	38.31	44.82	0.3070	2.35	0.23
Type D	5.33	2619.84	0.1023	16.57	100.00
	6.82	1316.79	0.1023	12.97	50.26
	9.04	588.62	0.1791	9.78	22.47
	10.09	189.02	0.1023	8.77	7.21
	11.02	486.13	0.1023	8.03	18.56
	11.33	328.81	0.1023	7.81	12.55

	13.53	616.04	0.0768	6.54	23.51
	13.85	369.18	0.1023	6.40	14.09
	16.48	149.73	0.1535	5.38	5.72
	16.99	249.21	0.0768	5.22	9.51
	17.28	386.64	0.1279	5.13	14.76
	17.98	138.74	0.3070	4.93	5.30
	19.50	390.10	0.1023	4.55	14.89
	20.08	385.54	0.1023	4.42	14.72
	21.28	484.13	0.1279	4.18	18.48
	22.88	193.33	0.1535	3.89	7.38
	23.71	353.50	0.1535	3.75	13.49
	25.50	97.46	0.1535	3.49	3.72
	27.27	145.81	0.1535	3.27	5.57
Type E	5.10	166.04	0.1535	17.32	52.30
	7.81	317.45	0.1791	11.32	100.00
	8.82	179.65	0.2047	10.03	56.59
	10.97	151.26	0.2047	8.06	47.65
	12.02	138.74	0.1535	7.36	43.70
	15.14	70.19	0.5117	5.85	22.11
	16.79	111.00	0.1535	5.28	34.97
	18.40	66.56	0.4093	4.82	20.97
	19.08	97.22	0.3070	4.65	30.63
	20.91	76.12	0.4093	4.25	23.98
	22.25	69.11	0.6140	4.00	21.77
Type F	5.86	9925.04	0.1004	15.09	100.00
	6.39	3049.85	0.0836	13.84	30.73
	7.21	1668.11	0.0836	12.26	16.81
	10.26	342.75	0.1338	8.62	3.45
	10.83	1177.50	0.0669	8.17	11.86
	11.22	1052.61	0.0836	7.88	10.61
	12.30	344.98	0.2007	7.20	3.48
	13.41	1274.77	0.0836	6.60	12.84
	14.05	1246.30	0.0836	6.30	12.56
	16.13	1222.17	0.1004	5.49	12.31

16.84	803.92	0.2007	5.27	8.10
17.23	750.76	0.2007	5.15	7.56
17.76	1065.21	0.1338	4.99	10.73
18.42	1503.52	0.2007	4.82	15.15
19.81	1904.97	0.0669	4.48	19.19
20.58	859.81	0.1004	4.32	8.66
21.36	1393.30	0.0836	4.16	14.04
22.07	1013.11	0.1171	4.03	10.21
23.68	1580.69	0.0669	3.76	15.93
23.87	1542.75	0.1004	3.73	15.54
24.38	999.19	0.2342	3.65	10.07
25.99	634.61	0.2676	3.43	6.39
26.44	649.77	0.1338	3.37	6.55
27.19	596.23	0.2342	3.28	6.01

[0351] All crystal forms that could be obtained at ambient conditions were further characterized by TGA, DSC, and ^{1}H NMR.

[0352] 3-DM Glycolate Type A

[0353] 3-DM Glycolate Type A was prepared via solution crystallization in IPA/H_2O solvent system. Detailed preparation procedures and characterization results can be found in Examples 2-8 above. Glycolate Type A was speculated to be an anhydrate due to the small TGA weight loss and neat DSC signal.

[0354] 3-DM Glycolate Type B

[0355] 3-DM Glycolate Type B was obtained via vapor diffusion of a THF solution of 3-DM Glycolate Type A in cyclohexane atmosphere. The XRPD pattern is shown in Figure 30, which is consistent with that of Glycolate Type B obtained during salt screening. TGA showed a weight loss of 3.6% up to 120 ºC and a stepwise weight loss of 8.2% between 120 ºC and 160 ºC. DSC showed a weak endotherm at 146.9 ºC and a sharp endotherm at 222.7 ºC (peak). ^{1}H NMR indicated that the molar ratio of residual solvent (cyclohexane) to 3-DM was 0.4:1 (the corresponding TGA weight loss was 6.3%). The molar ratio of acid to free base was 1:1. XRPD showed that no form change was observed after heating 3-DM Glycolate Type B to 120 ºC and cooling

to RT. For the sample after heating to 120 ºC, TGA, DSC and ^{1}H NMR characterization were performed again. TGA showed a weight loss of 3.5% up to 120 ºC and a stepwise weight loss of 7.6% between 120 ºC and 170 ºC. DSC showed a weak endotherm at 146.3 ºC and a sharp endotherm at 214.7 ºC (peak). ^{1}H NMR showed that the molar ratio of residual solvent (cyclohexane) to 3-DM had decreased to 0.2:1 (the corresponding TGA weight loss was 3.9%). As no form change occurred after removal of much of the solvent, the residual solvent might be surface-adsorbed. Based on the results, 3-DM Glycolate Type B is likely to be a hydrate or anhydrate. After heating to 160 ºC and cooling to RT, 3-DM Glycolate Type A was obtained.

[0356] 3-DM Glycolate Type C

[0357] 3-DM Glycolate Type C was obtained via vapor diffusion of 3-DM Glycolate Type A in $CHCl_3$ atmosphere. The XRPD pattern is shown in Figure 30. TGA showed a weight loss of 5.9% up to 150 ºC. DSC showed two endotherms at 61.2 ºC and 222.0 ºC (peak), and an exotherm at 140.8 ºC (peak). ^{1}H NMR showed that the molar ratio of acid to free base was 1:1, while the solvent $CHCl_3$ was not detected. After heating 3-DM Glycolate Type C to 100 ºC and cooling to RT, no form change was observed. 3-DM Glycolate Type A was obtained after heating to 150 ºC and cooling to RT.

[0358] Another batch of 3-DM Glycolate Type C was obtained via slurry of 3-DM Glycolate Type A in $CHCl_3$ at 50 ºC. XRPD was consistent with the material obtained by vapor diffusion. TGA showed a weight loss of 2.9% up to 150 ºC. DSC showed two endotherms at 57.3 ºC and 219.7 ºC (peak), and an exotherm at 134.7 ºC (peak). ^{1}H NMR indicated that the molar ratio of acid to free base was 1:1, while the solvent $CHCl_3$ was not detected. After N_2 sweeping 3-DM Glycolate Type C for 20 min at 30 ºC, a new form was observed, which was classified as 3-DM Glycolate Type F. After exposure of Type F to ambient conditions for 1 hr, Glycolate Type C was re-obtained. Based on these results, glycolate Type C was speculated to be a hydrate as it could be re-obtained after absorbing water in air during exposure to ambient conditions.

[0359] 3-DM Glycolate Type D

[0360] 3-DM Glycolate Type D was obtained via vapor diffusion of 1,4-dioxane solution of 3-DM Glycolate Type A in an MTBE atmosphere. The XRPD pattern is shown in Figure 30. TGA showed a weight loss of 2.6% up to 100 ºC. DSC showed two

endotherms at 63.0 ºC and 210.3 ºC (peak), and an exotherm at 123.2 ºC (peak). ^{1}H NMR showed that the molar ratio of acid to free base was 1:1, while the solvents 1,4-dioxane and MTBE were not detected. Variable temperature XRPD was performed for further characterization of 3-DM Glycolate Type D. No form change was observed after heating 3-DM Glycolate Type D to 90 ºC with N_2 protection. As a result, 3-DM Glycolate Type D was speculated to be an anhydrate. However, after heating Glycolate Type D (810397-74-A8) to 150 ºC and cooling to RT, Glycolate Type A was observed by XRPD.

[0361] 3-DM Glycolate Type E

[0362] 3-DM Glycolate Type E was obtained via slow evaporation of an EtOH solution of 3-DM Glycolate Type A at RT. The XRPD pattern is shown in Figure 30. Form conversion to Glycolate Type A was observed after drying Glycolate Type E at RT.

[0363] 3-DM Glycolate Type F

[0364] 3-DM Glycolate Type F was obtained after N_2 sweeping 3-DM Glycolate Type C at 30 ºC for 20 min. The XRPD pattern is shown in Figure 30. After exposure of Glycolate Type F to ambient conditions for 30 min, Glycolate Type C was re-obtained.

[0365] Anti-solvent Addition

[0366] For each experiment, approximately 15 mg of 3-DM Glycolate Type A was added into a 20-mL glass vial and dissolved in 0.1-1.6 mL of corresponding solvent to obtain a clear solution. The solution was magnetically stirred with addition of anti-solvent until precipitates appeared or the total volume of anti-solvent reached 10 mL. The obtained solids were isolated for XRPD analysis. If solids were not obtained, slurry at 5 ºC and/or evaporation at RT was performed. The results (Table 32) showed that Glycolate Type A, B, C, A+B, and amorphous samples were generated.

Table 32. Summary of anti-solvent addition experiments for 3-DM Glycolate Type A.

Experiment ID	Solvent (v/v)	Anti-solvent	Result
810397-68-A1	MeOH	MTBE	Glycolate Type B
810397-68-A2		Toluene	Glycolate Type A
810397-68-A3		$CHCl_3$	Glycolate Type C

Experiment ID	Solvent (v/v)	Anti-solvent	Result
810397-68-A4		H_2O	Amorphous
810397-68-A5	THF	n-Heptane	Glycolate Type C
810397-68-A6		Toluene	Glycolate Type C
810397-68-A7		MTBE	Glycolate Type A+B
810397-68-A8		Cyclohexane	Glycolate Type C
810397-68-A9	DMSO	MTBE	Glycolate Type A
810397-68-A10		Toluene	Glycolate Type A
810397-68-A11		IPAc	Glycolate Type A+B
810397-68-A12		H_2O	Glycolate Type A

[0367] Solid-vapor Diffusion

[0368] Solid vapor diffusion experiments were conducted using 12 different solvents. For each experiment, approximately 15-mg of 3-DM Glycolate Type A was weighed into a 3-mL vial, which was placed into a 20-mL vial with 4 mL of volatile solvent. The 20-mL vial was sealed with a cap and kept at RT for 7 days allowing solvent vapor to interact with the sample. The solids were tested by XRPD and the results, summarized in Table 33, showed that Glycolate Type A and C were obtained.

Table 33. Summary of solid-vapor diffusion experiments for 3-DM Glycolate Type A.

Experiment ID	Solvent	Result
810397-69-A1	H_2O	Glycolate Type A
810397-69-A2	DCM	Glycolate Type A
810397-69-A3	EtOH	Glycolate Type A
810397-69-A4	MeOH	Glycolate Type A
810397-69-A5	ACN	Glycolate Type A
810397-69-A6	THF	Glycolate Type A
810397-69-A7	$CHCl_3$	Glycolate Type C
810397-69-A8	Acetone	Glycolate Type A
810397-69-A9	DMSO	Glycolate Type A
810397-69-A10	EtOAc	Glycolate Type A
810397-69-A11	1,4-Dioxane	Glycolate Type A

Experiment ID	Solvent	Result
810397-69-A12	IPA	Glycolate Type A

[0369] Slurry at RT

[0370] For each experiment, approximately 15 mg of 3-DM Glycolate Type A was suspended in 0.5 mL of corresponding solvent in an HPLC vial. After the suspension was stirred magnetically (~1000 rpm) for 4 days at RT, the remaining solids were isolated by centrifugation for XRPD analysis. The results, summarized in Table 34, indicated that only Glycolate Type A was generated.

Table 34. Summary of slurry conversion experiments at RT for 3-DM Glycolate Type A.

Sample ID	Solvent (v/v)	Crystal Form
810397-70-A1	EtOH	Glycolate Type A
810397-70-A2	IPA	Glycolate Type A
810397-70-A3	Acetone	Glycolate Type A
810397-70-A4	MIBK	Glycolate Type A
810397-70-A5	EtOAc	Glycolate Type A
810397-70-A6	IPAc	Glycolate Type A
810397-70-A7	MTBE	Glycolate Type A
810397-70-A8	2-MeTHF	Glycolate Type A
810397-70-A9	ACN	Glycolate Type A
810397-70-A10	DCM	Glycolate Type A
810397-70-A11	Toluene	Glycolate Type A
810397-70-A12	H_2O	Glycolate Type A
810397-70-A13	MeOH/Toluene (1:4)	Glycolate Type A
810397-70-A14	Acetone/EtOAc (1:1)	Glycolate Type A
810397-70-A15	THF/Toluene (1:1)	Glycolate Type A
810397-70-A16	EtOH/MEK (1:1)	Glycolate Type A
810397-70-A17	DMSO/H_2O (1:9)	Glycolate Type A
810397-70-A18	CPME/n-Heptane (1:1)	Glycolate Type A
810397-70-A19	ACN/IPAc (1:1)	Glycolate Type A
810397-70-A20	1,4-Dioxane/$CHCl_3$ (1:1)	Glycolate Type A

Sample ID	Solvent (v/v)	Crystal Form
810397-70-A21	MIBK/MTBE (1:1)	Glycolate Type A
810397-70-A22	IPA/H_2O (aw~0.2)	Glycolate Type A
810397-70-A23	IPA/H_2O (aw~0.4)	Glycolate Type A
810397-70-A24	IPA/H_2O (aw~0.6)	Glycolate Type A
810397-70-A25	IPA/H_2O (aw~0.8)	Glycolate Type A

[0371] Slurry at 50 °C

[0372] For each experiment, approximately 15 mg of 3-DM Glycolate Type A was suspended in 0.5 mL of corresponding solvent in an HPLC vial. After the suspension was magnetically stirred (~1000 rpm) for 3 days at 50 ºC, the remaining solids were isolated by centrifugation for XRPD analysis. The results, summarized in Table 35, indicated that Glycolate Type A and C were generated.

Table 35. Summary of slurry conversion experiments at 50 ºC for 3-DM Glycolate Type A.

Experiment ID	Solvent (v/v)	Result
810397-71-A1	IPA	Glycolate Type A
810397-71-A2	MIBK	Glycolate Type A
810397-71-A3	EtOAc	Glycolate Type A
810397-71-A4	MTBE	Glycolate Type A
810397-71-A5	ACN	Glycolate Type A
810397-71-A6	$CHCl_3$	Glycolate Type C
810397-71-A7	Toluene	Glycolate Type A
810397-71-A8	H_2O	Glycolate Type A
810397-71-A9	EtOH/n-Heptane (1:1)	Glycolate Type A
810397-71-A10	Acetone/Toluene (1:1)	Glycolate Type A
810397-71-A11	1,4-Dioxane/H_2O (1:1)	Glycolate Type A
810397-71-A12	IPA/IPAc (1:1)	Glycolate Type A
810397-71-A13	ACN/2-MeTHF (1:1)	Glycolate Type A
810397-71-A14	MIBK/MTBE (1:1)	Glycolate Type A
810397-71-A15	THF/n-Heptane (1:1)	Glycolate Type A

Experiment ID	Solvent (v/v)	Result
810397-71-A16	NMP/H_2O (1:4)	Glycolate Type A

[0373] Slow Evaporation

[0374] Slow evaporation experiments were performed under seven conditions. For each experiment, approximately 15 mg of 3-DM Glycolate Type A was dissolved in 0.4-2.0 mL of corresponding solvent in a 4-mL glass vial and filtered using a 0.45 µm PTFE membrane. The visually clear solution was subjected to evaporation at RT in a vial sealed by a transparent PE-Plug (poke 4 small holes). The solids were isolated for XRPD analysis and the results, summarized in Table 36, indicated that Glycolate Type A, C, and E were obtained.

Table 36. Summary of slow evaporation experiments for 3-DM Glycolate Type A.

Experiment ID	Solvent	Result
810397-72-A1	MeOH	Glycolate Type A
810397-72-A2	EtOH	Glycolate Type E
810397-72-A3	Acetone	Glycolate Type A
810397-72-A4	EtOAc	Glycolate Type A
810397-72-A5	THF	Glycolate Type C
810397-72-A6	DCM	Glycolate Type C
810397-72-A7	ACN	Glycolate Type A

[0375] Slow Cooling

[0376] For each experiment, approximately 15 mg of 3-DM Glycolate Type A was suspended in 1.0-2.0 mL of corresponding solvent in a 3-mL glass vial at RT. The suspension was then heated to 50 ºC, stirred for 1.5 hrs, and filtered into a new vial using a 0.45 µm PTFE membrane. Filtrates were slowly cooled down to 5 ºC at a rate of 0.1 ºC/min. The obtained solids were kept isothermal at 5 ºC before isolation for XRPD analysis. The results, summarized in Table 37, indicated that Glycolate Type A, B, and C were generated.

Table 37. Summary of slow cooling experiments for 3-DM Glycolate Type A.

Experiment ID	Solvent (v/v)	Result
810397-73-A1#	EtOH	Glycolate Type A
810397-73-A2*	Acetone	Glycolate Type C
810397-73-A3	ACN	Glycolate Type A
810397-73-A4*	EtOAc/MEK (1:1)	Glycolate Type C
810397-73-A5#	IPA/H_2O (9:1)	Glycolate Type B
810397-73-A6	THF/$CHCl_3$ (1:1)	Glycolate Type C

[0377] * Clear solution was obtained after cooling to 5 ºC, which was transferred to -20 ºC for precipitation.

[0378] #: Clear solution was obtained after cooling to 5 ºC and -20 ºC, so solution was transferred to evaporate at RT.

[0379] Liquid-Vapor Diffusion

[0380] For each experiment, approximately 15 mg of 3-DM Glycolate Type A was dissolved in 0.1-1.6 mL of corresponding solvent in a 3-mL glass vial to obtain a clear solution. The 3-mL vial was then placed into a 20-mL vial with 4 mL of anti-solvent. The 20-mL vial was sealed with a cap and kept at RT for 7 days, allowing organic vapor to interact with the solution. The solids were isolated for XRPD analysis. The results, summarized in Table 38, showed that Glycolate Type A, B, D, and A+C were generated.

Table 38. Summary of liquid-vapor diffusion experiments for 3-DM Glycolate Type A.

Experiment ID	Solvent	Anti-solvent	Result
810397-74-A1	EtOH	n-Hexane	Glycolate Type A
810397-74-A2*		MTBE	Glycolate Type A+C
810397-74-A3		Toluene	Glycolate Type A
810397-74-A4	THF	n-Hexane	Glycolate Type A
810397-74-A5		Cyclohexane	Glycolate Type B
810397-74-A6		MTBE	Glycolate Type A
810397-74-A7	1,4-Dioxane	n-Heptane	Glycolate Type A+C
810397-74-A8*		MTBE	Glycolate Type D
810397-74-A9*		H_2O	Amorphous

Experiment ID	Solvent	Anti-solvent	Result
810397-74-A10*	NMP	Toluene	Glycolate Type A
810397-74-A11*		MTBE	Glycolate Type A
810397-74-A12		H_2O	Glycolate Type A

[0381] *: No solids were obtained after vapor diffusion, so solution was transferred to evaporate at RT.

[0382] Polymer-Induced Crystallization

[0383] For each experiment, approximately 15 mg of 3-DM Glycolate Type A was dissolved in 1.0-2.0 mL of corresponding solvent in a 4-mL glass vial and filtered using a 0.45 μm PTFE membrane to a new vial containing ~2 mg of polymers. The visually clear solution was subjected to evaporation at RT in a vial sealed by a transparent PE-Plug (poke 4 small holes). The solids were isolated for XRPD analysis. The results, summarized in Table 39, indicated that Glycolate Type A, C, and amorphous samples were obtained.

Table 39. Summary of polymer-induced crystallization experiments for 3-DM Glycolate Type A.

Exp. ID	Solvent (v/v)	Polymer	Result
810397-75-A1	EtOH	Polymer mixture A	Glycolate Type A
810397-75-A2	MIBK		Glycolate Type A
810397-75-A3	ACN/H_2O (1:1)		Glycolate Type C + peaks
810397-75-A4	MTBE	Polymer mixture B	Glycolate Type A
810397-75-A5	IPAc		Glycolate Type A
810397-75-A6	MeOH/DCM (1:1)		Amorphous

[0384] Polymer mixture A: polyvinyl pyrrolidone (PVP), polyvinyl alcohol (PVA), polyvinylchloride (PVC), polyvinyl acetate (PVAC), hypromellose (HPMC), methyl cellulose (MC) (mass ratio of 1:1:1:1:1:1)

Polymer mixture B: polycaprolactone (PCL), polyethylene glycol (PEG), poly (methyl methacrylate) (PMMA) sodium alginate (SA), and hydroxyethyl cellulose (HEC) (mass ratio of 1:1:1:1:1).

[0385] Grinding

[0386] For each experiment, approximately 15 mg of 3-DM Glycolate Type A was added into an agate mortar and manually ground using a pestle for about 3 min after adding the corresponding solvent. The solids were checked by XRPD and the results, summarized in Table 40, showed that only Glycolate Type A was obtained.

Table 40. Summary of grinding experiments for 3-DM Glycolate Type A.

Exp. ID	Solvent	Result
810397-76-A1	None	Glycolate Type A
810397-76-A2	IPA	Glycolate Type A
810397-76-A3	MIBK	Glycolate Type A
810397-76-A4	EtOAc	Glycolate Type A

[0387] **Example 10: Re-Preparation of 3-DM Glycolate Forms**

[0388] 3-DM Glycolate Type B, Type C, and Type D were re-prepared for slurry competition experiments. The characterization results are summarized in Table 41.

Table 41. Characterization results of 3-DM Glycolate Type B/C/D.

Crystal form	TGA weight loss (%)	DSC endotherm (ºC, peak)	Molar ratio (acid/API)	Purity (area%)
Glycolate Type B (810397-81-A1)	3.1 (to 120 ºC) 5.0 (120 - 160 ºC)	139.1, 219.7	1:1	99.4#
Glycolate Type C (810397-69-A7)	8.2 (to 150 ºC)	135.0*, 224.0, 226.7	1:1	99.1
Glycolate Type D (810397-74-A8)	3.5 (to 100 ºC)	129.0, 135.7*, 210.9	1:1	93.8

[0389] *: Exotherm peak.

[0390] #: Glycolate Type B converted to Type A+B after storage at ambient condition before HPLC purity test.

[0391] Re-preparation of 3-DM Glycolate Type B

[0392] 3-DM Glycolate Type B was prepared via vapor diffusion of THF solution of 3-DM Glycolate Type A in cyclohexane atmosphere. XRPD was consistent with the material prepared during polymorph screening (**Example 9**). TGA showed a weight loss of 3.1% up to 120 ºC and a stepwise weight loss of 5.0% between 120 ºC and 160 ºC. DSC showed a broad endotherm at 139.1 ºC and a sharp endotherm at 219.7 ºC (peak). ^{1}H NMR showed that the molar ratio of residual solvent (cyclohexane) to 3-DM was 0.2:1 (the corresponding TGA weight loss was 3.8%) and that the acid/free base ratio was 1:1. HPLC purity was determined as 99.4% (area).

[0393] Re-preparation of 3-DM Glycolate Type C

[0394] 3-DM Glycolate Type C was prepared via a slurry of 3-DM Glycolate Type A in $CHCl_3$ at 50 ºC, followed by drying of the solids at RT. XRPD was consistent with the material prepared during polymorph screening (**Example 9**). TGA showed a weight loss of 8.2% up to 150 ºC. DSC showed an exotherm at 135.0 ºC and two endotherms at 224.0 ºC and 226.7 ºC (peak). ^{1}H NMR showed that the molar ratio of residual solvent ($CHCl_3$) to 3-DM was 0.4:1 (the corresponding TGA weight loss was 9.6%)

and that the acid/free base ratio was 1:1. HPLC purity was determined as 99.1% (area).

[0395] Re-preparation of 3-DM Glycolate Type D

[0396] 3-DM Glycolate Type D was prepared via adding the anti-solvent MTBE into a 1,4-Dioxane solution of 3-DM Glycolate Type A, followed by evaporation at RT. XRPD was consistent with the material prepared during polymorph screening (**Example 9**). TGA showed a weight loss of 3.5% up to 100 ºC. DSC showed two endotherms at 129.0 ºC and 210.9 ºC (peak) and an exotherm at 135.7 ºC (peak). ^{1}H NMR showed that the molar ratio of residual solvent (1,4-dioxane) to 3-DM was 0.3:1 (the corresponding TGA weight loss was 4.9%) and that the acid/free base ratio was 1:1. HPLC purity was determined as 93.8% (area).

[0397] **Example 11: Interconversion Relationships Between 3-DM Glycolate Forms**

[0398] Interconversion Relationship Between 3-DM Glycolate Types A/B/D

[0399] To investigate the thermodynamic stability relationship between 3-DM Glycolate anhydrates, slurry competition experiments between anhydrates Glycolate Type A and Type D and anhydrate/hydrate Type B were performed at RT and 50 ºC in EtOH and MIBK. Detailed procedures were as follows.

1) 3-DM Glycolate Type A was used to saturate the corresponding solvent systems at RT and 50 ºC before filtration to obtain a near-saturated solution.
2) Equal amounts (about 6 mg each) of 3-DM Glycolate Type A, Type B, and Type D were added into 1 mL of the saturated solution of Type A at RT or 50 ºC to form a suspension.
3) The suspension was magnetically stirred at RT or 50 ºC for 2 days and the solids were then isolated for XRPD analysis.

[0400] Glycolate Type A was obtained under all conditions, indicating that Glycolate Type A was more thermodynamically stable than Glycolate Type B and Type D in the temperature range of RT to 50 ºC.

[0401] Interconversion Relationship Between 3-DM Glycolate Types A/B/C

[0402] To investigate the interconversion relationship between 3-DM Glycolate Type A, Type B, and Type C, slurry competition experiments were performed in IPA/H_2O with water activity aw=0-1 at RT. Detailed procedures are as follows.

1) 3-DM Glycolate Type A was used to saturate the corresponding solvent systems at RT before filtration to obtain a near-saturated solution.
2) Equal amounts (about 7 mg each) of 3-DM Glycolate Type A, Type B, and Type C were added into 1 mL of the saturated solution of Type A at RT to form a suspension.
3) The suspension was magnetically stirred at RT for 2 days and the solids were then isolated for XRPD analysis.

[0403] The results are summarized in Table 42. Glycolate Type A was obtained under all conditions, indicating that Glycolate Type A was more thermodynamically stable than Glycolate Type B and Type C under the tested conditions.

Table 42. Results of slurry competition experiment between 3-DM Glycolate Types A/B/C.

Experiment ID	Starting form	Solvent (aw)	Temperature	Final form
810397-84-A1	Glycolate Type A, B, and C*	IPA	RT	Glycolate Type A
810397-84-A2		IPA/H_2O (aw~0.2)	RT	
810397-84-A3		IPA/H_2O (aw~0.4)	RT	
810397-84-A4		IPA/H_2O (aw~0.6)	RT	
810397-84-A5		IPA/H_2O (aw~0.8)	RT	
810397-84-A6		H_2O	RT	

[0404] *: Glycolate Type B and Type C converted to a mixture of Glycolate Type B+A and Glycolate Type C+A, respectively, before slurry competition.

[0405] **Example 12: Solid State Stability Evaluation of 3-DM Glycolate Type A**

[0406] According to the characterization and interconversion relationship study results, 3-DM Glycolate Type A was selected as the leading form for solid state stability evaluation. 3-DM Glycolate Type A samples were stored under 60 ºC/closed for 1 day, 25 ºC/60% RH/open for 2 weeks, and 40 ºC/75% RH/open for 2 weeks. All the stability samples were characterized by XRPD, HPLC, and Karl Fischer (KF) titration, with the results summarized in Table 43. No form change or significant HPLC purity decrease was observed for 3-DM Glycolate Type A under any condition, indicating good physical and chemical stability.

Table 43. Solid state stability evaluation results of 3-DM Glycolate Type A.

Condition	Initial Purity (area %)	Purity (area %)	Final/ Initial Purity (%)	Form Change	H_2O by KF (wt%)
60 ºC/closed		98.81	100.1	No	0.21
25 ºC/60% RH/open	98.76	98.80	100.0	No	0.40
40 ºC/75% RH/open		98.75	100.0	No	0.23

[0407] **Example 13: Summary of Polymorph Screening of 3-DM Glycolate**

[0408] Using compound 3-DM Glycolate as the starting material, 100 polymorph screening experiments were performed via anti-solvent addition, solid-vapor diffusion, slurry, slow evaporation, slow cooling, liquid-vapor diffusion, polymer induced crystallization and grinding. A total of six forms of 3-DM Glycolate were obtained during screening, as shown by XRPD: Glycolate Type A (the starting form) and Types B through F. All crystal forms except Glycolate Type E and Type F were further characterized by thermogravimetric analysis (TGA), differential scanning calorimetry (DSC) and proton nuclear magnetic resonance (^{1}H NMR). A characterization summary for all crystal forms is shown in Table 44. Based on characterization results, Glycolate Type A and D were anhydrates, Glycolate Type C was a hydrate, Glycolate Type B was a hydrate or anhydrate, and Glycolate Type E and F were metastable forms.

[0409] Interconversion among Glycolate Types A, B, and D was studied in EtOH and MIBK at room temperature (RT, 23±3 ºC) and 50 ºC. Glycolate Type A was obtained

under all conditions, indicating Glycolate Type A was thermodynamically more stable than Glycolate Types B and D from RT to 50 ºC. Slurry competition experiments among Glycolate Types A, B, and C were conducted in IPA/H_2O (aw=0~1) at RT. Glycolate Type A was obtained under all conditions. An interconversion diagram of the different forms is displayed below in Scheme 2.

[0410] 3-DM Glycolate Type A was further characterized in solid state stability studies. Samples were stored under 60 ºC for one day, 25 ºC/60% RH for 2 weeks, and 40 ºC/75% RH for two weeks. No form change or purity decrease was observed for Glycolate Type A under any condition tested.

Table 44. Characterization of 3-DM Glycolate crystal forms.


Crystal form	TGA weight loss (%)	DSC endotherm (ºC, peak)	Molar ratio (acid/API)	Speculated form
Glycolate Type A (810397-64-B)	1.1 (to 150 ºC)	222.1	1:1	Anhydrate
Glycolate Type B (810397-74-A5)	3.6 (to 120 ºC) 8.2 (120 ~160 ºC)	146.9, 222.7	1:1	Anhydrate/ hydrate
Glycolate Type C (810397-69-A7)	5.9 (to 150 ºC)	61.2, 140.8*, 222.0	1:1	Hydrate
Glycolate Type D (810397-74-A8)	2.6 (to 100 ºC)	63.0, 123.2*, 210.3	1:1	Anhydrate
Glycolate Type E (810397-72-A2)	NA	NA	NA	Metastable
Glycolate Type F (810397-80-A)	--	--	--	Metastable

[0411] *: Exotherm peak temperature.

[0412] NA: Form converted to Type A after drying, so TGA, DSC, and ^1H NMR characterization were not performed.

[0413] --: Type F was only observed under N_2 protection and converted to Type C after exposure to ambient conditions.

Scheme 2. Interconversion diagram of 3-DM Glycolate crystal forms.



[0414] A summary of each numbered process step in Scheme 2 above is provided below:

[0415] (1) Slurry in EtOH or MIBK at RT and 50 ºC.

[0416] (2) Slurry in IPA/H2O (aw=0~1) at RT.

[0417] (3) Heating to 160 ºC and cooling to 30 ºC.

[0418] (4) Heating to 150 ºC and cooling to 30 ºC.

[0419] (5) Solid vapor diffusion in CHCl3 atmosphere.

[0420] (6) Slurry in CHCl3 at 50 ºC.

[0421] (7) Drying at RT.

[0422] (8) N2 sweeping for 20 min 30 ºC.

[0423] (9) Exposure to ambient conditions for 30 min.

[0424] Example 14: Characterization of Starting Materials for 3-DM Glycolate Type A Crystallization Process Development

[0425] 3-DM Free Base

[0426] 3-Deuteromitragynine (3-DM) free base for use in 3-DM Glycolate Type A crystallization process development experiments (see **Examples 18-21**) was prepared as previously described above and characterized by XRPD, TGA, mDSC, ^{1}H NMR, and HPLC. XRPD showed that the material was amorphous. TGA/mDSC revealed a weight loss of 1.6% up to 200 ºC and a possible glass transition (Tg) signal

at 93.1 ºC (middle temperature). ^{1}H NMR confirmed the identity of the material. HPLC purity was determined as 99.57% (area), with detailed results listed in Table 45. The area percentage of impurity 3-deuterocorynantheidine (3-DCR) was 0.05%.

Table 45. HPLC data of 3-DM free base starting material.

#Peak	RRT	Area (%)
1	0.69	0.08
2	0.70	0.18
3	0.73	0.12
4	0.90 (3-DCR)	0.05
5	**1.00 (3-DM)**	**99.57**

[0427] 3-DM Glycolate Type A

[0428] 3-DM Glycolate Type A for use in crystallization process development experiments (see **Examples 18-21**) of this same material was characterized by XRPD, TGA, mDSC, ^{1}H NMR, PLM, PSD, and HPLC. XRPD confirmed that the material was Glycolate Type A. TGA/DSC revealed a weight loss of 1.2% up to 170 ºC and a sharp endotherm at 221.9 ºC (onset). ^{1}H NMR indicated that the molar ratio of acid to free base was 1.0:1 and that the residual solvent (IPA) to free base ratio was 0.08:1 (the corresponding TGA weight loss was 1.0 wt%). PLM as shown in Figure 31 showed that the material consisted of short rod-like crystals. Particle size distribution (PSD) showed that the particle size D90 was 10.03 µm and a unimodal distribution was observed after sonication for 30 s under 30-Watt power (Table 46). HPLC purity was determined as 99.38% (area) and the area percentage of impurity 3-DCR was 0.07%, with detailed results summarized in Table 47.

Table 46. PSD results of 3-DM Glycolate Type A starting material.

Sonication time (s)	MV (µm)	SD (µm)	D10 (µm)	D50 (µm)	D90 (µm)
0	106.7	168.4	1.92	5.37	425.1
30	5.17	4.60	1.55	3.88	10.03

Table 47. HPLC data of 3-DM Glycolate Type A starting material.

#Peak	RRT	Area (%)	#Peak	RRT	Area (%)
1	0.69	0.08	5	0.91 (3-DCR)	0.07
2	0.70	0.18	6	0.94	0.03
3	0.73	0.12	**7**	**1.00 (3-DM)**	**99.38**
4	0.82	0.10	8	1.24	0.05

[0429] **Example 15: Solubility Measurements of 3-DM Glycolate Type A**

[0430] In order to guide the optimization of the solution crystallization of 3-DM Glycolate Type A (see **Examples 18-21**), the solubility of 3-DM free base and 3-DM Glycolate Type A was measured in IPA and IPA/H_2O (19:1, v/v) at 5 ºC, 20 ºC, 40 ºC, and 60 ºC. Detailed experimental procedures were as follows.

1) Weighed about 50 mg of 3-DM free base or 25 mg of 3-DM Glycolate Type A into 3-mL glass vials. Added 1.0 mL of the corresponding solvent (IPA or IPA/H_2O (19:1, v/v)) into each glass vial to form a suspension.
2) Suspensions were mixed at 5 ºC, 20 ºC, 40 ºC, or 60 ºC for 24 hrs.
3) After 24 hrs, suspensions were extracted into centrifugation tubes, followed by centrifugation (10,000 rpm, RT, 0.5 min).
4) The supernatants were filtered through a 0.45 μm PTFE membrane into 10-mL volumetric flasks, diluted to the volume by IPA, and these solutions were used for HPLC solubility tests. The residual solids were analyzed by XRPD.

[0431] XRPD showed that Free Base Type A (a crystalline form of 3-DM free base, see **Example 16**) and Glycolate Type A were obtained after solubility tests of 3-DM free base (amorphous) and 3-DM Glycolate Type A, respectively. Solubility results are summarized in Tables 48 and 49. The solubility of free base was much higher than that of Glycolate Type A in both IPA and IPA/H_2O (19:1, v/v) at each temperature. Therefore, both IPA and IPA/H_2O (19:1, v/v) were considered suitable for crystallization. In addition, a significant solubility decrease for Glycolate Type A was observed when the temperature was decreased from 60 ºC to 5 ºC. Therefore, cooling was expected to further improve the yield. Furthermore, solubility of Free Base Type A in IPA/H_2O (19:1, v/v) was higher than IPA, so IPA/H_2O (19:1, v/v) was expected to result in a better volumetric efficiency. As there was not expected to be a significant difference in yield using either solvent, and IPA/H_2O (19:1, v/v) was used in previous

formation of Glycolate Type A (see **Example 7**), it was selected as the preferred solvent.

Table 48. Solubility of 3-DM Free Base and Glycolate Type A in IPA.

3-DM Material (Sample ID)	Experiment ID	Temp. (ºC)	Solubility (mg/mL)	Solid Form
Amorphous Free Base (810081-01-A)	810081-04-A1	5	11.08	Free Base Type A
	810081-04-B1	20	13.16	Free Base Type A
	810081-45-C1	40	24.74#	Free Base Type A&
	810081-05-D1*	60	56.70#	Free Base Type A
Glycolate Type A (810397-64-B)	810081-04-A3	5	2.29	Glycolate Type A
	810081-04-B3	20	2.93	Glycolate Type A
	810081-45-C3	40	3.33#	Glycolate Type A&
	810081-45-D3	60	9.09#	Glycolate Type A&

[0432]*: The initial loading concentration was 152 mg/mL.

[0433]#: Centrifugation (~8,000 rpm, 5 min) and filtration was performed at corresponding temperature, 40 or 60 ºC.

[0434]&: Solids from another experiment were collected for XRPD analysis.

Table 49. Solubility of 3-DM Free Base and Glycolate Type A in IPA/H_2O (19:1, v/v).

3-DM Material (Sample ID)	Experiment ID	Temp. (ºC)	Solubility (mg/mL)	Solid Form
Amorphous Free Base	810081-04-A2	5	13.63	Free Base Type A

3-DM Material (Sample ID)	Experiment ID	Temp. (ºC)	Solubility (mg/mL)	Solid Form
(810081-01-A)	810081-04-B2	20	18.38	Free Base Type A
	810081-45-C2	40	33.65#	Free Base Type A&
	810082-08-A1*	60	73.94#	Free Base Type A&
Glycolate Type A (810397-64-B)	810081-04-A4	5	5.30	Glycolate Type A
	810081-04-B4	20	5.88	Glycolate Type A
	810081-45-C4	40	9.72#	Glycolate Type A&
	810081-45-D4	60	17.56#	Glycolate Type A&

[0435] *: The initial loading concentration was ~75 mg/mL.

[0436] #: Centrifugation (~8,000 rpm, 5 min) and filtration was performed at corresponding temperature, 40 or 60 ºC.

[0437] &: Solids from another experiment were collected for XRPD analysis.

[0438] **Example 16: Characterization of 3-DM Free Base Type A**

[0439] 3-DM Free Base Type A was obtained via stirring of a slurry of amorphous 3-DM free base in IPA at 40 ºC for one day and drying the resulting solid at RT under vacuum for 3 hrs. The XRPD pattern of the resulting material is displayed in Figure 32. TGA revealed a weight loss of 2.2% up to 80 ºC and a stepwise weight loss of 12.3% between 80 ºC and 110 ºC. DSC showed a sharp endotherm at 94.6 ºC (onset). ^{1}H NMR showed that the molar ratio of residual solvent (IPA) to free base was 0.9:1 (corresponding TGA weight loss was 12.2 wt%, close to the second step weight loss). Based on data above, Free Base Type A was identified as an IPA solvate.

[0440] **Example 17: Solution Stability of 3-DM Free Base and 3-DM Glycolate Type A**

[0441] To ensure that instability would not confound crystallization process development experiments (see **Examples 18-21**), solution stability of 3-DM free base and 3-DM Glycolate Type A in IPA and IPA/H_2O (19:1, v/v) at 20 ºC, 40 ºC, 60 ºC, and 65 ºC was assessed. Detailed experimental procedures were as follows.

1) For each sample, weighed approximately 1 mg of material (3-DM free base or 3-DM Glycolate Type A) into a 3-mL glass vial. Added 1 mL of solvent to dissolve solid completely.
2) Stored solution samples at different temperatures (20 ºC, 40 ºC, 60 ºC, and 65 ºC) for 38 hrs.
3) Tested HPLC purity for all solution samples.

[0442] Solution stability results are summarized in Table 50. No decrease in HPLC purity or increase in 3-DCR area percentage was observed under any condition, indicating good solution stability for amorphous 3-DM free base and 3-DM Glycolate Type A in IPA and IPA/H_2O (19:1, v/v) in the range of 20 ºC - 65 ºC.

Table 50. Solution stability results of 3-DM Free Base and 3-DM Glycolate Type A.

3-DM Material (Sample ID)	Solvent (v/v)	HPLC purity of initial sample (area%)	HPLC Final Purity / Initial Purity (%)			
			20 ºC	40 ºC	60 ºC	65 ºC
Amorphous free base (810081-01-A)	IPA	99.57	100.3	100.4	100.3	100.3
	IPA/H_2O (19:1)		100.3	100.3	100.3	100.3
Glycolate Type A (810397-64-B)	IPA	98.81	100.2	100.2	100.2	100.2
	IPA/H_2O (19:1)		100.2	100.2	100.2	100.2

[0443] **Example 18: Preliminary Crystallization Process Development for 3-DM Glycolate Type A (300-mg Scale)**

[0444] Preliminary crystallization experiments were performed in IPA/H_2O (19:1, v/v) using slow cooling method. Detailed procedures are described in Table 51. The process parameters and characterization results are summarized in Table 52. XRPD indicated that 3-DM Glycolate Type A was obtained in both batches prepared.

[0445] *For Batch 810081-18-A2_dry*: A weight loss of 1.7% up to 150 ºC, a weak endotherm at 202.8 ºC (peak), and a sharp endotherm at 220.8 ºC (onset) were observed in TGA/DSC. ^{1}H NMR showed that the molar ratio of acid/free base was 1.0:1, and residual IPA/free base was 0.04:1 (0.5 wt%). PLM showed needle-like or short rod-like crystals. HPLC purity was determined as 99.86% (area).

[0446] *For Batch 810081-28-B*: A weight loss of 0.8% up to 150 ºC and a sharp endotherm at 223.0 ºC (onset) were observed in TGA/DSC. ^{1}H NMR showed that the molar ratio of acid/free base was 1.0:1, and negligible solvent (IPA) was detected. PLM showed rod-like crystals. Results of PSD characterization are listed in Table 53. The particle size (D90) was 47.05 µm (after sonication for 30 s / 30 Watt), and a

unimodal distribution was observed before and after sonication. HPLC purity was determined as 99.89% (area).

[0447] The main differences between these two batches were initial temperature and seed loading. As the initial concentration was very close to the equilibrium solubility of Free Base Type A in IPA/H_2O (19:1, v/v) at 60 ºC (73.94 mg/mL), for Batch 810081-28-B, the temperature was increased to 65 ºC to prevent possible precipitation. For Batch 810081-18-A2_dry, no seeds were added and particles with small size were obtained. For batch 810081-28-B, ~2% seeds were added for generation of crystals with larger particle size.

[0448] However, the yield of both batches was relatively low. For batches at only 300-mg scale, it was suspected that material loss during filtration and collection of solids might be very high. To confirm if a higher yield could be achieved on larger scale, additional experiments were conducted at 900-mg scale (see **Example 19**).

Table 51. Procedures for preliminary crystallization development experiments for 3-DM Glycolate Type A at 300-mg scale.

Batch No.	Experimental Procedure
810081-18-A2_dry	1. Transfer 301.1 mg of free base (810081-01-A) into a 20-mL glass vial. Add 4.0 mL IPA/H_2O (19:1, v/v). Magnetically stir at 60 ºC for 30 mins and filter the solution through 0.45 µm PTFE membrane. 2. Transfer 56.4 mg of glycolic acid into another 20-mL glass vial. Add 4.0 mL IPA/H_2O (19:1, v/v) to dissolve solids. 3. Inject acid solution into the API solution. A suspension was observed after ~1.4 mL of acid solution was added. 4. Slurry the suspension at 60 ºC for 2 h, and cool to 5 ºC at 0.1 ºC/min. Keep isothermal at 5 ºC. 5. Isolate solids by centrifugation (10,000 rpm, 2 mins). Dry solids under vacuum at 50 ºC and collect solids after drying. 6. A total of 257.0 mg of solids were obtained (Yield: 71.9%).
810081-28-B	1. Transfer 300.1 mg of free base (810081-01-A) into a 20-mL

Batch No.	Experimental Procedure
	glass vial. Add 4 mL IPA/H_2O (19:1, v/v). Transfer the suspension to slurry at 65 ºC for 30 min before filtered with a 0.45 µm PTFE membrane. 2. Transfer 59.3 mg of glycolic acid into another 20-mL glass vial. Add 3.0 mL IPA/H_2O (19:1, v/v) to dissolve solids. 3. Transfer the API solution into a 50-mL reactor and stir at 65 ºC. 4. Add acid solution into API solution over 30 min. 5. Add 6.9 mg of Glycolate Type A (810081-18-A2_dry) into the solution as seeds. Stir at 65 ºC for 2 hrs. 6. Cool to 25 ºC at 0.1 ºC/min and keep slurry at 25 ºC for 15.5 hrs. Then cool to 5 ºC at 0.1 ºC/min and keep isothermal at 5 ºC overnight. 7. Collect solids by vacuum filtration. Dry solids at 50 ºC under vacuum for 4 h. A total of 214.0 mg of solids were obtained (Yield: 59.6%).

Table 52. Parameters and characterization results for preliminary crystallization development experiments for 3-DM Glycolate Type A at 300-mg scale.

Sample ID	810081-18-A2_dry	810081-28-B
Scale (free base)	300 mg	300 mg
Initial temperature	60 ºC	65 ºC
Initial concentration (free base)	~75 mg/mL	~75 mg/mL
Seeds	NA	~2%
XRPD	Glycolate Type A	Glycolate Type A
TGA weight loss (%)	1.7 (to 150 ºC)	0.8 (to 150 ºC)
DSC endotherm (ºC, onset)	202.8*, 220.8	223.0
Molar ratio (acid/free base)	1.0:1	1.0:1
Residual IPA (wt%)	0.5	Not detected
Morphology	Needle-like or short rod-like	Rod-like
HPLC purity (area%)	99.86	99.89

Sample ID	810081-18-A2_dry	810081-28-B
Yield (%)	71.9	59.6

[0449] *Peak temperature.

Table 53. PSD results of 3-DM Glycolate Type A (Batch 810081-28-B).

Sonication Time (30 Watt)	MV (µm)	SD (µm)	D10 (µm)	D50 (µm)	D90 (µm)
0 s	28.68	18.45	7.51	19.96	58.66
30 s	22.77	14.81	5.92	15.93	47.05

[0450] **Example 19: Additional Crystallization Process Development for 3-DM Glycolate Type A (900-mg Scale)**

[0451] In order to improve the yield of 3-DM Glycolate Type A crystallization, two additional experiments were performed at 900-mg scale. Detailed procedures are described in Table 54. The process parameters and characterization results are summarized in Table 55. Glycolate Type A was obtained in both batches with ~80% yield, which was higher than observed at 300-mg scale.

[0452] *For Batch 810081-50-A*: A weight loss of 0.7% up to 150 ºC, a weak endotherm at 204.5 ºC (peak), and a sharp endotherm at 223.2 ºC (onset) were observed in TGA/DSC. ^{1}H NMR showed that the molar ratio of acid/free base was 1.0:1, and residual IPA/free base was 0.01:1 (0.19 wt%). PLM showed that the sample was needle-like or rod-like crystals. HPLC purity was determined as 99.94% (area).

[0453] *For Batch 810082-13-B*: A weight loss of 1.0% up to 150 ºC, a weak endotherm at 205.2 ºC (peak), and a sharp endotherm at 222.4 ºC (onset) were observed in TGA/DSC. ^{1}H NMR showed that the molar ratio of acid/free base was 1.0:1, and residual IPA/free base was 0.01:1 (0.19 wt%). PLM showed that the sample was rod-like or short rod-like crystals. HPLC purity was determined as 99.91% (area).

[0454] The main difference between these two batches was seed loading. For Batch 810082-13-B, ~2% seeds were added and the particle size was larger than that of

Batch 810081-50-A, which was crystallized without seed addition. On the basis of these results, it was decided that seeding was likely to improve reproducibility at different scales and therefore, that seeds would be added at larger scale (see **Example 20**).

Table 54. Procedures for crystallization development experiments for 3-DM Glycolate Type A at 900-mg scale.

Batch No.	Experimental Procedure
810081-50-A	1. Transfer 900.8 mg of free base (810081-01-A) and 12 mL IPA/H_2O (19:1, v/v) into a 20-mL glass vial. Magnetically stir the suspension at 65 ºC for 2 h. Clear API solution was obtained. 2. Transfer 180.4 mg of glycolic acid into another 20-mL glass vial. Add 9 mL IPA/H_2O (19:1, v/v) to dissolve solid for acid solution. 3. Transfer the API solution into a 50-mL reactor and stir at 65 ºC. 4. Add acid solution into API solution over 1 h, and magnetically stir (300 rpm) at 65 ºC for 2 h. 5. Cool to 5 ºC at 0.1 ºC/min and keep slurry at 5 ºC for 10 h. 6. Sample 1 mL of suspension, filter it by 0.45 µm PTFE membrane to obtain clear solution for concentration test (3.76 mg/mL). 7. After slurrying at 5 ºC for 3.5 h, isolate suspension by vacuum filtration. Wash the wet cake with 4 mL cold IPA. 8. Dry solids under vacuum at RT for 13.5 h and then at 50 ºC for 2 h. 9. A total of 815.0 mg of solids were obtained (Yield: 79.8%).
810082-13-B	1. Transfer 899.7 mg of free base (810081-01-A) and 12 mL IPA/H_2O (19:1, v/v) into a 20-mL glass vial. Magnetically stir the suspension at 65 ºC for 2 h. Clear API solution was obtained.

Batch No.	Experimental Procedure
	2. Transfer 180.2 mg of glycolic acid into another 20-mL glass vial. Add 9 mL IPA/H_2O (19:1, v/v) to dissolve solid for acid solution. 3. Transfer the API solution into a 50-mL reactor and stir at 65 ºC. 4. Add 3 mL acid solution into API solution over 20 min. Then add 18.2 mg of seeds (810081-18-A2_dry). A turbid suspension was observed. 5. Add 6 mL acid solution into API solution over 40 min. Magnetically stir (300 rpm) at 65 ºC for 2 h. 6. Cool to 5 ºC at 0.1 ºC/min and keep slurry at 5 ºC for ~13.5 h. 7. Isolate solids by vacuum filtration. Wash the wet cake by 4 mL of cold IPA. 8. Dry solids under vacuum at 50 ºC for 2 h and then at RT for 13 h. 9. A total of 861.7 mg of solids were obtained (Yield: 78.7%).

Table 55. Parameters and characterization results for crystallization development experiments for 3-DM Glycolate Type A at 900-mg scale.

Sample ID	810081-50-A	810082-13-B
Scale (free base)	900 mg	900 mg
Initial temperature	65 ºC	65 ºC
Initial concentration (free base)	~75 mg/mL	~75 mg/mL
Seeds	NA	~2%
XRPD	Glycolate Type A	Glycolate Type A
TGA weight loss (%)	0.7 (to 150 ºC)	1.0 (to 150 ºC)
DSC endotherm (ºC, onset)	204.5*, 223.2	205.2*, 222.4
Molar ratio (acid/free base)	1.0:1	1.0:1
Residual IPA (wt%)	0.19	0.19

Sample ID	810081-50-A	810082-13-B
Morphology	Needle-like or rod-like	Rod-like or short rod-like
HPLC purity (area%)	99.94	99.91
Yield (%)	79.8	78.7

[0455] *: Peak temperature.

[0456] **Example 20: Crystallization of 3-DM Glycolate Type A at 5-g Scale**

[0457] To further evaluate the robustness of the optimized crystallization method (see **Example 19**), an experiment at 5-g scale was conducted using the optimized parameters for solution crystallization of 3-DM Glycolate Type A, as follows:

Initial free base concentration: ~75 mg/mL in IPA/H_2O (19:1, v/v)
Initial acid concentration: ~20 mg/mL in IPA/H_2O (19:1, v/v)
Volume ratio (free base/acid): 4:3
Initial temperature: 65 ºC
Seed loading: ~2%
Ending temperature: 5 ºC
Cooling rate: 0.1 ºC/min

[0458] The detailed procedure is described in Table 56. XRPD showed that 3-DM Glycolate Type A was successfully obtained with a yield of 85.1%. TGA/DSC showed a weight loss of 1.0% up to 150 ºC, a weak endotherm at 206.9 ºC (peak), and a sharp endotherm at 222.2 ºC (onset). ^1H NMR showed that the molar ratio of acid/free base was 1.0:1, and residual IPA/free base was 0.02:1 (0.32 wt%). PLM showed that the product was rod-like crystals (Figure 33). HPLC revealed a purity of 99.89% (area) and impurity 3-DCR was not detected (area percentage was ~0.01%), as detailed in Table 57.

Table 56. Procedure for crystallization of 3-DM Glycolate Type A at 5-g scale.

Procedure
1. Weigh 1.001 g of glycolic acid into a 100-mL glass vial. Add 50 mL IPA/H_2O (19:1, v/v) to dissolve solid for acid solution.
2. Add 5.010 g of free base (810081-01-A) and 66.7 mL IPA/H_2O (19:1, v/v) to a 250-mL reactor. Overhead stir the suspension at 65 ºC (250 rpm) for 2 h for

Procedure
API solution. 3. Add 16 mL acid solution into API solution over 20 min. Then add 100.7 mg of seeds (810081-28-B). A turbid suspension was observed. 4. Add 34 mL acid solution into API solution over 40 min. Overhead stir (300 rpm) at 65 ºC for 2 h. 5. Cool to 5 ºC at 0.1 ºC/min and keep slurry at 5 ºC for ~13.5 h. 6. Isolate solids by vacuum filtration. Wash the wet cake with ~10 mL cold IPA. 7. Dry solids under vacuum at RT for ~13.5 h and then at 50 ºC for 3 h. 8. A total of 5.175 g of solids were obtained (Yield: 85.1%).

[0459] A polarized light microscopy image of the 3-DM Glycolate Type A crystals prepared at 5-g scale (810082-25-B) is shown in Figure 33.

Table 57. HPLC data of 3-DM Glycolate Type A prepared at 5-g scale (810082-25-B).

#Peak	RRT	Area (%)
1	0.70	0.05
2	0.73	0.06
3 (3-DM)	**1.00**	**99.89**

[0460] **Example 21: Crystallization of 3-DM Glycolate Type A Using IPA as Solvent (300-mg Scale)**

[0461] Two crystallization experiments were conducted with 3-DM Glycolate using IPA as solvent via a slow cooling method. Detailed procedures are described in Table 58. XRPD revealed that Glycolate Type A was obtained in both batches, which were further characterized with the results summarized in Table 59.

[0462] *For Batch 810082-18-B*: A weight loss of 2.5% up to 150 ºC, a weak endotherm at 203.7 ºC (peak), and a sharp endotherm at 221.8 ºC (onset) were observed in TGA/DSC. ^{1}H NMR showed that the molar ratio of acid/free base was 1.0:1, and residual IPA/free base was 0.03:1 (0.6 wt%). PLM showed that the product was rod-like or short rod-like crystals.

[0463] *For Batch 810082-20-A*: A weight loss of 1.3% up to 150 ºC, a weak endotherm at 203.9 ºC (peak), and a sharp endotherm at 222.2 ºC (onset) were observed in TGA/DS. ^{1}H NMR showed that the molar ratio of acid/free base was 1.0:1, and residual IPA/free base was 0.04:1 (0.7 wt%). PLM showed that the product was rod-like or short rod-like crystals.

[0464] When switching to IPA, the residual solvent increased compared to batches prepared using IPA/H_2O (19:1, v/v). In addition, the IPA/H_2O (19:1, v/v) procedure could be run at the higher concentration of 75 mg/mL with full dissolution, whereas a reduction to 50 mg/mL was required in the IPA procedure.

Table 58. Procedures for crystallization of 3-DM Glycolate Type A from IPA.

Batch No.	Experimental Procedure
810082-18-B	1. Add 300.6 mg of free base (810081-01-A) and 4 mL IPA into a 20-mL glass vial. Magnetically stir the suspension at 65 ºC for 1 h. 2. Additional 2 mL of IPA was added as some solids were not dissolved. Magnetically stir at 65 ºC for additional 2 h for API solution. 3. Transfer 68.0 mg of glycolic acid into another 20-mL glass vial. Add 3.4 mL IPA to dissolve solid for acid solution. 4. Transfer the API solution into a 50-mL reactor and stir at 65 ºC. 5. Add 1 mL acid solution into API solution over 7 min. Then add 6.0 mg of seeds (810081-18-A2_dry). A turbid suspension was observed. 6. Add 2 mL acid solution into API solution over 13 min. Magnetically stir (300 rpm) at 65 ºC for 2 h. 7. Cool to 5 ºC at 0.1 ºC/min and keep slurry at 5 ºC for ~13.5 h. 8. Isolate solids by vacuum filtration. Wash the wet cake with ~4 mL cold IPA. 9. Dry solids under vacuum at 50 ºC for 2 h and then at RT for 13

Batch No.	Experimental Procedure
	h. 10. A total of 270.2 mg of solids were obtained (Yield: 73.8%).
810082-20-A	1. Add 300.4 mg of free base (810081-01-A) and 4 mL IPA into a 50-mL reactor. Magnetically stir at 65 ºC for ~1.5 h. Solids were not dissolved completely. 2. Transfer 67.9 mg of glycolic acid into another 20-mL glass vial. Add 3.4 mL IPA to dissolve solid for acid solution. 3. Add 1 mL acid solution into API suspension over 7 min. Then add 6.1 mg of seeds (810081-18-A2_dry). A turbid suspension was observed. 4. Add 2 mL acid solution into API solution over 13 min. Magnetically stir (300 rpm) at 65 ºC for 2 h. 5. Cool to 5 ºC at 0.1 ºC/min and keep slurry at 5 ºC for ~13.5 h. 6. Isolate solids by vacuum filtration. Wash the wet cake with ~4 mL cold IPA. 7. Dry solids under vacuum at 50 ºC for 2 h and then at RT for 13 h. 8. A total of 300.2 mg of solids were obtained (Yield: 82.2%).

Table 59. Parameters and characterization results of 3-DM Glycolate Type A crystallized from IPA.

Sample ID	810082-18-B	810082-20-A
Scale (free base)	300 mg	300 mg
Initial temperature	65 ºC	65 ºC
Initial concentration (free base)	~50 mg/mL	~75 mg/mL
Seeds	~2%	~2%
XRPD	Glycolate Type A	Glycolate Type A
TGA weight loss (%)	2.5 (to 150 ºC)	1.3 (to 150 ºC)
DSC endotherm (ºC, onset)	203.7*, 221.8	203.9*, 222.2

Sample ID	810082-18-B	810082-20-A
Molar ratio (acid/free base)	1.0:1	1.0:1
Residual IPA (wt%)	0.6	0.7
Morphology	Rod-like or short rod-like	Rod-like or short rod-like
HPLC purity (area%)	99.91	99.91
Yield (%)	73.8	82.2

[0465] *: Peak temperature.

[0466] **Example 22. Characterization of Mitragynine Free Base and Crude Alkaloid Extract for Salt Preparation**

[0467] Pure mitragynine free base (810080-01-A) for use in salt preparation experiments was commercially obtained. The material was characterized by XRPD, TGA, mDSC, ^{1}H NMR, and HPLC. XRPD showed that the material was amorphous, with several weak peaks (Figure 34). The TGA curve showed a weight loss of 1.3% up to 200 ºC and mDSC curve revealed a possible glass transition (Tg) signal at 94.2 ºC (middle temperature) (Figure 35). ^{1}H NMR confirmed the structure of the material (Figure 36). As shown in Table 60, HPLC purity was determined as 98.98% (area) and the content of corynantheidine (CR) impurity was determined as 0.48% (area).

[0468] Crude alkaloid extract of *Mitragyna speciosa* (810080-01-B) was commercially obtained and characterized by XRPD, TGA, and HPLC. XRPD showed that the material was amorphous (See Figure 37). TGA showed a weight loss of 2.3% up to 200 ºC (See Figure 38). The weight content of mitragynine was found to be 61.0% by HPLC (75.94 area%) using pure mitragynine (810080-01-A) as a reference standard. The CR content was found to be 1.19% (area).

Table 60. Individual impurities of amorphous mitragynine free base starting material (810080-01-A) as determined by HPLC.

#Peak	RRT	Area(%)
1	0.69	0.11
2	0.86	0.06

#Peak	RRT	Area(%)
3	0.90 (CR)	0.48
4	**1.00 (mitragynine)**	**98.98**
5	1.06	0.05
6	1.17	0.06
7	1.20	0.11
8	1.23	0.05
9	1.29	0.10

[0469] **Example 23: Preparation and Characterization of Mitragynine Salts**

[0470] Using mitragynine free base as the starting material, mitragynine Fumarate Type A, L-Lactate Type A, Glycolate Type A, Succinate Type A, and Mesylate Type A were prepared at 500-mg or 750-mg scale using procedures analogous to those used to prepare the corresponding 3-DM salts. The detailed preparation procedures are described in Table 61.

Table 61. Preparation procedures for mitragynine salts.

Mitragynine Salt Lead (Sample ID)	Procedure
Fumarate Type A (810080-06-B1)	1. Transfer 500.9 mg of free base (810080-01-A) and 149.9 mg of fumaric acid into a 20-mL glass vial. 2. Add 12 mL isopropanol IPA/H_2O (4:1, v/v) to form a suspension. Magnetically stir (~1000 rpm) at RT for 2 days. 3. Isolate solids by vacuum filtration. Dry solids at RT under vacuum for ~15 h. 4. A total of 439.7 mg of material was obtained (Yield: 67.6%).
L-Lactate Type A (810080-06-B2)	1. Transfer 126.4 mg of L-lactic acid into a 20-mL glass vial, add 8 mL IPA/H_2O (4:1, v/v) to dissolve acid. 2. Transfer 499.5 mg of free base (810080-01-A) into the acid solution. Magnetically stir (~1000 rpm) at RT for 2 days. 3. Isolate solids by vacuum filtration. Dry solids at RT under

Mitragynine Salt Lead (Sample ID)	Procedure
	vacuum for ~15 h. 4. A total of 266.8 mg of material was obtained (Yield: 42.6%).
Glycolate Type A (810080-07-B3)	1. Transfer 499.8 mg of free base (810080-01-A) and 98.4 mg of glycolic acid into a 20-mL glass vial. 2. Add 12 mL IPA/H_2O (4:1, v/v) to form a suspension. Magnetically stir (~1000 rpm) at RT for 2 days. 3. Isolate solids by vacuum filtration. Dry solids at RT under vacuum for ~15 h. 4. A total of 203.4 mg of material was obtained (Yield: 34.0%, with ID: 810080-06-B3). 5. Weigh 250.3 mg of free base (810080-01-A) and 49.6 mg of glycolic acid into the filtrate. Magnetically stir (~1000 rpm) at RT for 2 days. 6. Add the dried sample (810080-06-B3) into the suspension, magnetically stir (~1000 rpm) at RT for 4 h. 7. Isolate solids by vacuum filtration. Dry solids at 50 ºC under vacuum for ~2 h. 8. A total of 427.6 mg of material was obtained (Yield: 47.6%).
Succinate Type A (810080-06-B4)	1. Transfer 500.8 mg of free base (810080-01-A) and 151.4 mg of succinic acid into a 20-mL glass vial. 2. Add 12 mL IPA/H_2O (4:1, v/v) to form a suspension. Magnetically stir (~1000 rpm) at RT for 2 days. 3. Isolate solids by vacuum filtration. Dry solids at RT under vacuum for ~15 h. 4. A total of 362.7 mg of material was obtained (Yield: 55.6 %).
Mesylate Type A (810080-06-B5)	1. Transfer 126.7 mg of methanesulfonic acid into a 20-mL glass vial, add 10 mL ethyl acetate (EtOAc) to dissolve acid. 2. Transfer 501.0 mg of free base (810080-01-A) into the acid

Mitragynine Salt Lead (Sample ID)	Procedure
	solution. Magnetically stir (~1000 rpm) at RT for 2 days. 3. Isolate solids by vacuum filtration. Dry solids at RT under vacuum for ~15 h. 4. A total of 468.0 mg of material was obtained (Yield: 74.6 %).

[0471] XRPD revealed that the mitragynine salts obtained had similar crystalline properties to those obtained from the deuterated material, 3-DM (Table 62). The full XRPD traces are also shown in Figures 39-43.

Table 62. XRPD copper radiation results for mitragynine salts.

Mitragynine Salt	Pos. [°2θ]	Height [cts]	FWHM Left [°2θ]	d-spacing [Å]	Rel. Int. [%]
Fumarate Type A	8.30	877.10	0.1023	10.65	100.00
	9.47	149.70	0.1023	9.34	17.07
	14.29	311.81	0.1535	6.20	35.55
	15.06	83.30	0.1535	5.88	9.50
	15.47	64.83	0.1535	5.73	7.39
	16.03	211.03	0.1279	5.53	24.06
	16.73	234.50	0.1023	5.30	26.74
	17.33	513.93	0.1279	5.12	58.59
	18.55	338.24	0.1023	4.78	38.56
	19.06	765.72	0.1023	4.66	87.30
	19.22	788.22	0.1023	4.62	89.87
	19.89	95.70	0.1535	4.46	10.91
	21.19	221.58	0.1023	4.19	25.26
	21.44	238.21	0.1023	4.14	27.16
	22.22	175.89	0.1279	4.00	20.05
	22.89	204.35	0.1279	3.89	23.30
	23.22	299.25	0.1023	3.83	34.12
	23.61	265.88	0.1023	3.77	30.31

	24.51	126.44	0.1279	3.63	14.42
	25.20	373.37	0.1791	3.53	42.57
	25.62	564.11	0.1279	3.48	64.32
	26.81	199.64	0.1279	3.33	22.76
	28.80	227.71	0.1023	3.10	25.96
	30.95	198.61	0.1535	2.89	22.64
	34.11	20.76	0.6140	2.63	2.37
L-Lactate Type A	7.01	1984.77	0.1023	12.61	46.69
	10.12	4250.55	0.1023	8.74	100.00
	10.87	836.75	0.1023	8.14	19.69
	11.20	3069.37	0.1279	7.90	72.21
	13.17	972.58	0.1023	6.72	22.88
	13.94	298.46	0.1023	6.35	7.02
	15.06	347.19	0.1023	5.88	8.17
	15.57	873.48	0.1023	5.69	20.55
	15.90	2338.73	0.1791	5.57	55.02
	17.88	1729.79	0.1535	4.96	40.70
	18.25	456.60	0.1023	4.86	10.74
	18.95	727.87	0.0768	4.68	17.12
	19.44	1859.59	0.1023	4.57	43.75
	19.91	1020.38	0.1023	4.46	24.01
	20.80	2959.52	0.1279	4.27	69.63
	21.18	199.50	0.1023	4.20	4.69
	21.76	142.00	0.1535	4.08	3.34
	22.42	2886.58	0.1279	3.96	67.91
	22.86	373.87	0.1535	3.89	8.80
	23.44	231.96	0.0768	3.79	5.46
	23.77	382.95	0.1023	3.74	9.01
	24.53	543.69	0.1279	3.63	12.79
	24.90	2134.64	0.1279	3.58	50.22
	25.80	80.85	0.2047	3.45	1.90
	27.09	590.27	0.1023	3.29	13.89
	27.34	631.24	0.1023	3.26	14.85
	28.16	202.54	0.1279	3.17	4.77

	29.45	367.99	0.1279	3.03	8.66
	30.73	88.24	0.3070	2.91	2.08
	31.36	461.46	0.1279	2.85	10.86
	33.97	197.90	0.1023	2.64	4.66
	35.72	148.09	0.2047	2.51	3.48
	37.41	66.97	0.2558	2.40	1.58
	38.12	42.44	0.3070	2.36	1.00
Glycolate Type A	7.09	2227.09	0.1023	12.46	59.02
	10.16	3773.50	0.1023	8.71	100.00
	11.29	2780.95	0.1023	7.84	73.70
	13.23	1002.77	0.1023	6.69	26.57
	14.14	569.61	0.1023	6.26	15.09
	15.10	226.01	0.1023	5.87	5.99
	15.74	1100.63	0.0768	5.63	29.17
	16.01	2026.54	0.0768	5.53	53.70
	18.03	1749.18	0.1279	4.92	46.35
	18.54	640.19	0.1023	4.79	16.97
	18.90	243.68	0.1023	4.69	6.46
	19.22	710.87	0.0768	4.62	18.84
	19.54	1864.83	0.0768	4.54	49.42
	19.72	1408.53	0.0768	4.50	37.33
	20.36	1575.92	0.1023	4.36	41.76
	20.93	3005.06	0.1023	4.24	79.64
	22.62	2357.70	0.1279	3.93	62.48
	23.26	297.77	0.1023	3.82	7.89
	23.52	246.96	0.0768	3.78	6.54
	23.76	284.00	0.1023	3.74	7.53
	24.05	457.06	0.1023	3.70	12.11
	24.90	563.37	0.1279	3.58	14.93
	25.20	2346.48	0.1279	3.53	62.18
	25.94	100.05	0.1535	3.43	2.65
	27.63	1346.88	0.1023	3.23	35.69
	28.30	161.35	0.1023	3.15	4.28
	29.12	93.42	0.1535	3.07	2.48

	29.64	149.00	0.2047	3.01	3.95
	30.15	670.34	0.1023	2.96	17.76
	30.64	193.78	0.1023	2.92	5.14
	31.66	576.95	0.1023	2.83	15.29
	32.24	183.42	0.1279	2.78	4.86
	34.20	147.65	0.1535	2.62	3.91
	35.16	131.39	0.1023	2.55	3.48
	36.03	132.20	0.1023	2.49	3.50
	36.63	124.79	0.1023	2.45	3.31
Succinate Type A	6.24	979.78	0.1023	14.17	4.93
	8.47	19866.64	0.1023	10.44	100.00
	9.14	570.92	0.1279	9.67	2.87
	9.62	1525.98	0.1023	9.20	7.68
	10.07	468.63	0.1023	8.78	2.36
	13.48	87.24	0.1535	6.57	0.44
	14.36	2064.94	0.1023	6.17	10.39
	15.66	1012.50	0.1023	5.66	5.10
	16.13	1233.90	0.1023	5.50	6.21
	16.95	573.33	0.1023	5.23	2.89
	17.56	4829.41	0.1023	5.05	24.31
	18.72	618.40	0.1023	4.74	3.11
	19.27	4490.83	0.1023	4.61	22.60
	21.21	187.16	0.1023	4.19	0.94
	21.71	1704.95	0.1023	4.09	8.58
	22.33	888.53	0.1023	3.98	4.47
	23.13	1576.63	0.1023	3.84	7.94
	23.72	346.57	0.1023	3.75	1.74
	24.49	183.17	0.1023	3.63	0.92
	25.03	676.61	0.1023	3.56	3.41
	25.33	1045.48	0.0768	3.52	5.26
	25.52	1637.15	0.1023	3.49	8.24
	25.93	1819.07	0.1279	3.44	9.16
	26.97	339.78	0.0768	3.31	1.71
	28.63	200.76	0.1023	3.12	1.01

	29.08	491.71	0.1023	3.07	2.48
	30.52	139.21	0.1279	2.93	0.70
	31.27	1486.70	0.1279	2.86	7.48
	31.56	322.01	0.1023	2.83	1.62
	32.97	249.55	0.1279	2.72	1.26
	34.22	179.42	0.1023	2.62	0.90
	34.55	163.67	0.1023	2.60	0.82
	37.03	94.16	0.3070	2.43	0.47
	39.05	158.57	0.1023	2.31	0.80
Mesylate Type A	6.71	2834.93	0.1023	13.18	100.00
	8.21	375.62	0.1023	10.77	13.25
	10.03	218.11	0.1023	8.82	7.69
	11.59	1208.66	0.1023	7.64	42.63
	12.95	161.80	0.1279	6.84	5.71
	13.38	809.76	0.1023	6.62	28.56
	14.94	866.79	0.1023	5.93	30.58
	15.31	438.33	0.1023	5.79	15.46
	16.36	220.09	0.1023	5.42	7.76
	16.71	1770.92	0.1023	5.30	62.47
	17.37	1374.36	0.1023	5.10	48.48
	18.31	256.80	0.0768	4.84	9.06
	18.62	932.18	0.1023	4.76	32.88
	18.93	1154.31	0.1023	4.69	40.72
	19.79	564.35	0.1023	4.49	19.91
	20.07	909.38	0.1023	4.42	32.08
	21.17	308.16	0.1023	4.20	10.87
	21.44	348.36	0.1023	4.14	12.29
	22.24	475.02	0.1023	4.00	16.76
	22.71	576.59	0.1023	3.92	20.34
	22.96	319.53	0.1023	3.87	11.27
	23.71	276.64	0.0768	3.75	9.76
	24.17	378.94	0.0768	3.68	13.37
	24.42	503.38	0.1023	3.65	17.76
	24.64	227.58	0.1023	3.61	8.03

25.08	366.85	0.1023	3.55	12.94
25.99	920.51	0.1279	3.43	32.47
26.83	212.37	0.1023	3.32	7.49
27.08	260.83	0.1023	3.29	9.20
28.51	206.55	0.1279	3.13	7.29
30.08	88.12	0.1535	2.97	3.11
32.85	141.62	0.1023	2.73	5.00
33.73	73.68	0.1535	2.66	2.60
37.05	44.01	0.3070	2.43	1.55

[0472] All mitragynine salts obtained were further characterized by TGA, DSC, ^{1}H NMR, PLM, and HPLC, with the results summarized in Table 63.

Table 63. Characterization results of mitragynine salts.

Mitragynine Salt (Sample ID)	TGA weight loss (%)	DSC endotherm (ºC, onset)	Molar ratio (acid/base)	PLM Appearance	Purity (area %)	CR (area %)	Water Content by KF (%)
Fumarate Type A (810080-06-B1)	3.3 (to 150 ºC)	253.3	1:1	fine particles	99.41	0.41	0.47
Glycolate Type A (810080-07-B3)	2.3 (to 150 ºC)	220.3	1.1:1	rod-like crystals	99.89	0.06	0.60
L-Lactate Type A (810080-06-B2)	2.6 (to 150 ºC)	226.2	1:1	short rod-like crystals	99.81	0.11	0.31
Succinate Type A (810080-06-B4)	3.9 (to 150 ºC)	197.7*, 201.9*	1.1:1	needle-like crystals	99.63	0.26	0.42
Mesylate Type A (810080-06-B5)	1.6 (to 150 ºC)	274.9	1:1	fine particles	99.09	0.48	0.47

[0473] *: Peak temperature.

[0474] **Example 24: Solubility of Mitragynine Salts**

[0475] Solubility of mitragynine Fumarate Type A, L-Lactate Type A, Glycolate Type A, Succinate Type A, and Mesylate Type A in water and bio-relevant media (SGF, FaSSIF, and FeSSIF) was measured after 4 h at 37 ºC. Bio-relevant media were prepared in the same manner as **Example 4**. Solids were suspended in media at a solid loading of 10 mg/mL (calculated based on free base). The suspensions were agitated on a rolling incubator at 25 rpm at 37 ºC, prior to sampling at 4 h. At sampling time point, suspensions were separated by centrifugation (~10,000 rpm, 37 ºC, 3 min) prior to filtering the supernatants through 0.45 µm PTFE membranes. The filtrates were analyzed for HPLC solubility and pH, and the residual solids were used for XRPD analysis. Results are summarized in Table 64. No form change was observed by XRPD for any of the salts in any of the media after 4 h.

Table 64. Summary of solubility results of mitragynine salts.

Salt lead (810080-)	H_2O			SGF			FaSSIF			FeSSIF		
	S	pH	FC	S	pH	FC	S	pH	FC	S	pH	FC
Fumarate Type A (06-B1)	0.40	4.3	No	1.28	1.9	No	2.31	5.6	No	1.91	4.9	No
L-Lactate Type A (06-B2)	1.87	5.6	No	7.04	3.0	No	0.97	6.1	No	2.28	4.9	No
Glycolate Type A (07-B3)	7.88	5.3	No	7.76	3.4	--	1.03	6.1	No	2.22	4.9	No
Succinate Type A (06-B4)	3.14	4.9	No	7.63	3.4	No	4.63	5.3	No	2.81	4.9	No
Mesylate Type A (06-B5)	13.83*	4.6	No	8.11	1.8	--	1.72	5.9	No	2.36	4.9	No

[0476] S: Solubility, mg/mL, calculated based on free base.

[0477] FC: Form change.

[0478] *: The initial loading concentration was ~20 mg/mL (calculated based on free base).

--: Only limited amount of gel-like sample was obtained after solubility experiment. Therefore, XRPD was not conducted.

[0479] **Example 25: Hygroscopicity of Mitragynine Salts**

[0480] To investigate the solid form stability and hygroscopicity of mitragynine salts as a function of humidity, DVS isotherm plots of mitragynine Fumarate Type A, L-Lactate Type A, Glycolate Type A, Succinate Type A, and Mesylate Type A were collected at 25 ºC between 0 and 95% RH. After DVS test, samples were characterized by XRPD. The results are summarized in Table 65. No form change was observed by XRPD for any of the salts after DVS test.

Table 65. Summary of DVS results for mitragynine salts.

Mitragynine Salt (Sample ID)	Water uptake at 80% RH	Form change
Fumarate Type A (810080-06-B1)	0.18%	No
L-Lactate Type A (810080-46-A1_dry)*	0.03%	No
Glycolate Type A (810080-07-B3)	0.25%	No
Succinate Type A (810080-06-B4)	0.09%	No
Mesylate Type A (810080-06-B5)	0.40%	No

[0481] *: The remaining L-Lactate Type A (810080-06-B2) was not sufficient for DVS test, so another batch of L-Lactate Type A (810080-46-A1_dry) was used.

[0482] **Example 26: Chemical and Solid Form Stability of Mitragynine Salts**

[0483] The solid state stability of mitragynine Fumarate Type A, L-Lactate Type A, Glycolate Type A, Succinate Type A, and Mesylate Type A was evaluated under conditions of 25 ºC/60% RH and 40 ºC/75% RH for 1 week. Each sample was placed in a glass vial, sealed by parafilm with several holes, and stored under the designated tested condition. After one week, samples were taken for XRPD, KF, and HPLC purity test. The results are summarized in Table 66. No purity decrease, form change, or water content difference was observed for any of the salts after storage under either condition for one week.

Table 66. Summary of physicochemical stability evaluation of mitragynine salts.

Mitragynine Salt (810080-)	Initial		25 ºC/60%RH/1 week			40 ºC/75%RH/1 week		
	Purity (area%)	KF (wt%)	Purity/Initial purity (%)	KF (wt%)	Form change	Purity/Initial purity (%)	KF (wt%)	Form change
Fumarate Type A (06-B1)	99.41	0.47	99.9	0.24	No	100.2	0.34	No
L-Lactate Type A (06-B2)	99.81	0.31	100.1	0.43	No	100.1	0.68	No
Glycolate Type A (07-B3)	99.89	0.60	100.1	0.54	No	100.1	0.43	No
Succinate Type A (06-B4)	99.63	0.42	99.9	0.46	No	100.1	0.24	No
Mesylate Type A (06-B5)	99.09	0.47	100.1	0.43	No	100.1	0.44	No

[0484] **Example 27: Preparation of Mitragynine Salts from Crude Alkaloid Extract**

[0485] Crude *Mitragynina speciosa* alkaloid extract (810080-01-B), with weight content of mitragynine of 61.0% (75.94 area%), was used as the starting material for preparation of mitragynine salts, in order to test the ability of the different salts to separate mitragynine from a crude mixture. Two sets of experiments were set up with different charge ratios by loading different amounts of crude material and the same amount of counter ion, as follows:

(1) Treating the crude starting material as pure mitragynine free base with molar charge ratio 1:1 (Experiment ID has the prefix 810080-20-A).

(2) According to the weight purity, calculating the actual amount of mitragynine free base in crude material with molar charge ratio 1:1 (Experiment ID has the prefix 810080-20-B).

[0486] XRPD confirmed that Fumarate Type A and Mesylate Type A were obtained from charge ratio (1) and (2), L-Lactate Type A and Glycolate Type A were obtained from charge ratio (2). No L-Lactate Type A or Glycolate Type A was obtained with charge ratio (1) and no Succinate Type A was obtained with either charge ratio after slurry at RT for 3 days and then at 5 ºC for 1 day. The detailed crystallization procedures are described in Table 67. All crystalline salts obtained were further characterized by ^{1}H NMR and HPLC, with the results summarized in Table 68.

[0487] All salts obtained were more enriched in mitragynine than the starting material, with Glycolate Type A and L-Lactate Type A having the highest mitragynine content. In contrast, only Glycolate Type A and L-Lactate Type A afforded significant purging of the corynantheidine (CR) impurity relative to the starting material.

Table 67. Crystallization procedures for mitragynine salts using crude alkaloid extract (61.0 wt%, 75.94 area% mitragynine) as the starting material.

Mitragynine Salt (Sample ID)	Procedure
Fumarate Type A (810080-20-A1_dry)	1. Weigh 14.9 mg of fumaric acid and 50.1 mg of crude free base (810080-01-B) into a 3-mL glass vial. 2. Add 1.2 mL of IPA/H_2O (4:1, v/v) to form a suspension. 3. Magnetically stir (~1,000 rpm) at RT for 3 days. 4. Isolate solids by centrifugation (10,000 rpm, 2 min). Dry solids at RT under vacuum for 15.5 h.
Fumarate Type A (810080-20-B1_dry)	1. Weigh 14.9 mg of fumaric acid and 82.1 mg of crude free base (810080-01-B) into a 3-mL glass vial. 2. Add 1.2 mL of IPA/H_2O (4:1, v/v) to form a suspension. 3. Magnetically stir (~1,000 rpm) at RT for 3 days. 4. Isolate solids by centrifugation (10,000 rpm, 2 min). Dry solids at RT under vacuum for 15.5 h.
L-Lactate Type A (810080-20-A2)	1. Weigh 12.5 mg of L-lactic acid into a 3-mL glass vial. 2. Add 0.8 mL of IPA/H_2O (4:1, v/v) to dissolve acid. 3. Weigh 50.0 mg of crude free base (810080-01-B) into the acid solution. 4. Magnetically stir (~1,000 rpm) at RT for 3 days. Clear solution was obtained. 5. Transferred to slurry at 5 ºC for 1 day. Clear solution was obtained.
L-Lactate Type A (810080-20-B2_dry)	1. Weigh 12.6 mg of L-lactic acid into a 3-mL glass vial. 2. Add 0.8 mL of IPA/H_2O (4:1, v/v) to dissolve acid. 3. Weigh 82.0 mg of crude free base (810080-01-B) into the acid solution. 4. Magnetically stir (~1,000 rpm) at RT for 3 days. 5. Isolate solids by centrifugation (10,000 rpm, 2 min). Dry solids at RT under vacuum for 15.5 h.
Glycolate Type A (810080-20-A3)	1. Weigh 9.7 mg of glycolic acid and 50.2 mg of crude free base (810080-01-B) into a 3-mL glass vial. 2. Add 0.8 mL of IPA/H_2O (4:1, v/v) to form a suspension. 3. Magnetically stir (~1,000 rpm) at RT for 3 days. Clear solution was obtained. 4. Transferred to slurry at 5 ºC for 1 day. Clear solution was obtained.
Glycolate Type A (810080-20-	1. Weigh 9.8 mg of glycolic acid and 82.1 mg of crude free base (810080-01-B) into a 3-mL glass vial.

Mitragynine Salt (Sample ID)	Procedure
B3_dry)	2. Add 0.8 mL of IPA/H_2O (4:1, v/v) to form a suspension. 3. Magnetically stir (~1,000 rpm) at RT for 3 days. 4. Isolate solids by centrifugation (10,000 rpm, 2 min). Dry solids at RT under vacuum for 15.5 h.
Succinate Type A (810080-20-A4)	1. Weigh 15.0 mg of succinic acid and 50.1 mg of crude free base (810080-01-B) into a 3-mL glass vial. 2. Add 1.2 mL of IPA/H_2O (4:1, v/v) to form a suspension. 3. Magnetically stir (~1,000 rpm) at RT for 3 days. Clear solution was obtained. 4. Transferred to slurry at 5 ºC for 1 day. Clear solution was obtained.
Succinate Type A (810080-20-B4)	1. Weigh 15.1 mg of succinic acid and 81.9 mg of crude free base (810080-01-B) into a 3-mL glass vial. 2. Add 1.2 mL of IPA/H_2O (4:1, v/v) to form a suspension. 3. Magnetically stir (~1,000 rpm) at RT for 3 days. Clear solution was obtained. 4. Transferred to slurry at 5 ºC for 1 day. Clear solution was obtained.
Mesylate Type A (810080-20-A5_dry)	1. Weigh 12.9 mg of methanesulfonic acid into a 3-mL glass vial. 2. Add 1.0 mL of EtOAc to dissolve acid. 3. Weigh 49.9 mg of crude free base (810080-01-B) into the acid solution. 4. Magnetically stir (~1,000 rpm) at RT for 3 days. 5. Isolate solids by centrifugation (10,000 rpm, 2 min). Dry solids at 50 ºC under vacuum for 2 h.
Mesylate Type A (810080-20-B5_dry)	1. Weigh 12.9 mg of methanesulfonic acid into a 3-mL glass vial. 2. Add 1.0 mL of EtOAc to dissolve acid. 3. Weigh 82.2 mg of crude free base (810080-01-B) into the acid solution. 4. Magnetically stir (~1,000 rpm) at RT for 3 days. 5. Isolate solids by centrifugation (10,000 rpm, 2 min). Dry solids at RT under vacuum for 15.5 h.

Table 68. Characterization results of mitragynine salts obtained starting from crude alkaloid extract (61.0 wt%, 75.94 area% mitragynine).

Mitragynine Salt (Sample ID: 810080-20)	Target form	Charge ratio	Obtained form	HPLC purity (area%)	Weight purity (%)	CR Content (area%)
810080-01-B	Crude Alkaloid Extract	-	-	75.94	61.0	1.19
A1_dry	Fumarate Type A	(1)	Fumarate Type A	87.66	87.25	1.31
B1_dry	Fumarate Type A	(2)	Fumarate Type A	89.33	86.39	1.18
A2	L-Lactate Type A	(1)	NA	NA	NA	NA
B2_dry	L-Lactate Type A	(2)	L-Lactate Type A	93.97	91.45	0.41
A3	Gycolate Type A	(1)	NA	NA	NA	NA
B3_dry	Gycolate Type A	(2)	Gycolate Type A	94.45	88.84	0.36
A4	Succinate Type A	(1)	NA	NA	NA	NA
B4	Succinate Type A	(2)	NA	NA	NA	NA
A5_dry	Mesylate Type A	(1)	Mesylate Type A	86.98	75.44	1.47
B5_dry	Mesylate Type A	(2)	Mesylate Type A	84.51	75.01	1.39


[0488] NA: No solid was obtained after slurry at RT for 3 days and then at 5 ºC for 1 day.

[0489] (1): Treating the crude starting material as pure mitragynine free base with charge ratio 1:1.

[0490] (2): Based on weight purity, calculating the actual amount of mitragynine free base in crude material with charge ratio 1:1.

Numbered Embodiments:

1 A salt of 3-deuteromitragynine of Formula I



Formula I

wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

2 The salt of embodiment 1, wherein the anion is glycolate.

3 The salt of embodiment 1 or embodiment 2, wherein the anion is glycolate and the 3-deuteromitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately 7.06. 10.12, 11.25. 15.97, 17.99, 18.11, 19.49, 19.69, 20.34, 20.88, 22.58, 25.16 and 27.60 ±0.2 degrees 2 theta.

4 The salt of embodiment 1, wherein the anion is L-lactate.

5 The salt of embodiment 1 or embodiment 4, wherein the anion is L-lactate and the 3-deuteromitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately 6.86, 9.96, 10.73, 11.05, 15.74, 19.77, 20.63, 22.26 and 24.75 ±0.2 degrees 2 theta.

6 The salt of embodiment 1, wherein the anion is succinate.

7 The salt of embodiment 1 or embodiment 6, wherein the anion is succinate and the 3-deuteromitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately 8.46, 10.06, 17.56, 19.27, 21.71, 23.13, 25.53, 25.94 and 31.27 ±0.2 degrees 2 theta.


8 The salt of embodiment 1 wherein the anion is fumarate.

9 The salt of embodiment 1 or embodiment 8 wherein the anion is fumarate and the 3-deuteromitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately 8.41, 9.58, 17.46, 19.20, 23.03, 25.36, 25.77 and 31.11 ±0.2 degrees 2 theta.

10 The salt of embodiment 1 wherein the anion is mesylate.

11 The salt of embodiment 1 or embodiment 10, wherein the anion is mesylate and the 3-deuteromitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately 6.67; 11.56, 13.34, 14.93, 16.69, 17.34, 18.60, 18.89, 20.03, 22.21, 22.71 and 25.97 ±0.2 degrees 2 theta.

12 A salt of mitragynine of Formula II



Formula II

wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

13 The salt of embodiment 12, wherein the anion is glycolate.

14 The salt of embodiment 12 or embodiment 13, wherein the anion is glycolate and the mitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately 7.09, 10.16, 11.29, 13.23, 14.14, 15.74, 16.01, 18.03, 19.54, 19.72, 20.36, 20.93, 22.62, 25.20 and 27.63 ±0.2 degrees 2 theta.

15 The salt of embodiment 12, wherein the anion is L-lactate.

16 The salt of embodiment 12 or embodiment 15 wherein the anion is L-Lactate and the mitragynine salt exhibits an XRPD spectrum with copper radiation having 2

theta peaks at approximately 7.01, 10.12, 11.20, 15.57, 15.90, 17.88, 19.44, 19.91, 20.80, 22.42 and 24.90 ±0.2 degrees 2 theta.

17 The salt of embodiment 12, wherein the anion is succinate.

18 The salt of embodiment 12 or embodiment 17, wherein the anion is succinate and the mitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately 8.47, 9.62, 14.36, 19.27, 21.71, 23.13, 25.52, 25.93 and 31.27 ±0.2 degrees 2 theta.

19 The salt of embodiment 12, wherein the anion is fumarate.

20 The salt of embodiment 12 or embodiment 19, wherein the anion is fumarate and the mitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately 8.30, 14.29. 16.03, 16.73, 17.33, 18.55, 19.06, 19.22, 21.19. 21.44, 22.89, 23.61, 25.20, 25.62 and 28.80 ±0.2 degrees 2 theta.

21 The salt of embodiment 12, wherein the anion is mesylate.

22 The salt of embodiment 12 or embodiment 21, wherein the anion is mesylate and the mitragynine salt exhibits an XRPD spectrum with copper radiation having 2 theta peaks at approximately 6.71, 11.59, 13.38, 14.94, 16.71, 17.37, 18.62, 18.93, 20.07 and 25.99 ±0.2 degrees 2 theta.

23 A crystalline glycolate salt of 3-deuteromitragynine (3-DM), wherein the salt is glycolate Type A, glycolate Type B, glycolate Type C, glycolate Type D, glycolate Type E, glycolate Type F, or combinations thereof.

24 The Type A of 3-DM glycolate salt of embodiment 23 exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at approximately 7.06, 10.11, 11.24, 15.96, 18.01, 19.49, 19.70, 20.34, 20.88, 22.57, 25.15 and 27.62 ±0.2 degrees 2 theta.

25 The Type B of 3-DM glycolate salt of embodiment 23 exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at approximately 5.29, 5.68, 7.45, 10.78, 13.65, 19.86, 21.22, 22.72 and 24.05 ±0.2 degrees 2 theta.

26 The Type C of 3-DM glycolate salt of embodiment 23 exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at approximately 6.03, 7.43, 10.99, 11.39, 13.65, 14.22, 16.30, 18.07, 18.84, 20.08, 21.54, 22.87, 24.16 and 26.22 ±0.2 degrees 2 theta.

27 The Type D of 3-DM glycolate salt of embodiment 23 exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at approximately 5.33, 6.82, 9.04, 11.02, 11.33, 13.53, 13.85, 17.28, 19.50, 20.08, 21.28 and 23.71 ±0.2 degrees 2 theta.


28 The Type E of 3-DM glycolate salt of embodiment 23 exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at approximately 5.10, 7.81, 8.82, 10.97, 12.02, 16.79, 18.40, 19.08, 20.91 and 22.25 ±0.2 degrees 2 theta.

29 The Type F of 3-DM glycolate salt of embodiment 23 exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at approximately 5.86, 6.39, 7.21, 10.83, 11.22, 13.41, 14.05, 16.13, 17.7618.42, 19.81, 21.36, 22.07, 23.68, 23.87 and 24.38 ±0.2 degrees 2 theta.

30 A crystalline glycolate salt of mitragynine.

31 The crystalline mitragynine glycolate salt of embodiment 30 exhibiting an XRPD spectrum with copper radiation having 2 theta peaks at approximately 7.09, 10.16, 11.29, 13.23, 14.14, 15.74, 16.01, 18.03, 19.54, 19.72, 20.36, 20.93, 22.62, 25.20 and 27.63 ±0.2 degrees 2 theta and wherein the crystalline mitragynine glycolate salt is a Type A.

32 A pharmaceutical composition comprising an amount of one or more salts of 3-deuteromitragynine of Formula 1



Formula I.

33 The pharmaceutical composition of embodiment 32, wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

34 The pharmaceutical composition of embodiment 32 or embodiment 33, wherein the anion is glycolate.

35 The pharmaceutical composition of embodiment 32 or embodiment 33, wherein the anion is L-lactate.


36 The pharmaceutical composition of embodiment 32 or embodiment 33, wherein the anion is succinate.

37 The pharmaceutical composition of embodiment 32 or embodiment 33, wherein the anion is fumarate.

38 The pharmaceutical composition of embodiment 32 or embodiment 33, wherein the anion is mesylate.

39 The pharmaceutical composition of any one of embodiments 32 to 38, wherein the composition further includes a pharmaceutically acceptable carrier.

40 A pharmaceutical composition comprising an amount of one or more salts of mitragynine of Formula II



Formula II.

41 The pharmaceutical composition of embodiment 39-40, wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

42 The pharmaceutical composition of embodiment 40 or embodiment 41, wherein the anion is glycolate.

43 The pharmaceutical composition of embodiment 40 or embodiment 41, wherein the anion is L-lactate.

44 The pharmaceutical composition of embodiment 40 or embodiment 41, wherein the anion is succinate.

45 The pharmaceutical composition of embodiment 40 or embodiment 41, wherein the anion is fumarate.

46 The pharmaceutical composition of embodiment 40 or embodiment 41, wherein the anion is mesylate.

47 The pharmaceutical composition of any one of embodiments 40 to 46, wherein the composition further includes a pharmaceutically acceptable carrier.

48 A method of treating a subject afflicted with acute pain, chronic pain, a depressive disorder, a mood disorder, an anxiety disorder, borderline personality disorder, a substance use disorder, opioid use disorder, opioid withdrawal symptoms, alcohol use disorder, or alcohol withdrawal disorder, comprising administering an effective amount of a salt or a composition as defined in any one of embodiments 1 to 47 to the subject so as to thereby treat the subject afflicted with acute pain, chronic pain, a depressive disorder, a mood disorder, an anxiety disorder, borderline personality disorder, a substance use disorder, opioid use disorder, opioid withdrawal symptoms, alcohol use disorder, or alcohol withdrawal disorder.

49 A process for producing a glycolate salt of 3-deuteromitragynine, the process including the step of crystallizing a glycolate salt of 3-deuteromitragynine from a solution of isopropyl alcohol.

50 The process of embodiment 49, wherein the solution of isopropyl alcohol includes water.

51 A process for producing a glycolate salt of mitragynine, the process including the step of crystallizing a glycolate salt of mitragynine from a solution of isopropyl alcohol.

52 The process of embodiment 51, wherein the solution of isopropyl alcohol includes water.

53 The process of embodiment 51 or embodiment 52, wherein the glycolate salt of mitragynine is derived from a crude alkaloid extract of *Mitragyna speciosa*.

54 A process of purifying 3-deuteromitragynine or mitragynine, the purification process including a step of crystallizing any one of the 3-deuteromitragynine or mitragynine salts as defined above in any one of embodiments 1 to 22.

55 The process of embodiment 54 wherein the resulting purified 3-deuteromitragynine or mitragynine salt is at least 90% free of other compounds or impurities.

56 The process of embodiment 55 wherein the purified 3-deuteromitragynine or mitragynine salt is at least 95% free of other compounds or impurities.

57 The process of embodiment 55 or embodiment 56 wherein the purified 3-deuteromitragynine or mitragynine salt is at least 98% free of other compounds or impurities.

58 The process of any one of embodiments 55 to 57 wherein the purified 3-deuteromitragynine or mitragynine salt is at least 99% free of other compounds or impurities.

59 The process of embodiment 54 wherein the purified 3-deuteromitragynine salt has less than about 3% of the impurity 3-deuterocorynantheidine (3-DCR).

60 The process of embodiment 59 wherein the purified 3-deuteromitragynine salt has less than about 2% of the impurity 3-DCR.

61 The process of embodiment 59 or embodiment 60 wherein the purified 3-deuteromitragynine salt has less than about 1% of the impurity 3-DCR.

62 The process of any one of embodiments 59 to 61 wherein the purified 3-deuteromitragynine salt has less than about 0.5% of the impurity 3-DCR.

63 The process of embodiment 54 wherein the purified mitragynine salt has less than about 3% of the impurity corynantheidine (CR).

64 The process of embodiment 63 wherein the purified mitragynine salt has less than about 2% of the impurity CR.

65 The process of embodiment 63 or embodiment 64 wherein the purified mitragynine salt has less than about 1% of the impurity CR.

66 The process of any one of embodiments 63 to 65 wherein the purified mitragynine salt has less than about 0.5% of the impurity CR.


67 The process of any one of embodiments 54 to 58 or embodiments 63 to 66, wherein the purified mitragynine is derived from a crude alkaloid extract of *Mitragyna speciosa*.

68 The process of any one of embodiments 54 to 67 wherein the purification comprises crystallizing glycolate, L-lactate, succinate, fumarate, or mesylate salt.

69 The process of any one of embodiments 54 to 68 wherein the purification comprises crystallizing the glycolate salt.

Numbered Embodiments:

1. A salt of 3-deuteromitragynine of Formula I:



Formula I

wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

2. The salt of embodiment 1, wherein the anion is glycolate.

3. The salt of embodiment 2, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type A, glycolate Type B, glycolate Type C, glycolate Type D, glycolate Type E, glycolate Type F, or combinations thereof.

4. The salt of embodiment 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type A.

5. The salt of embodiment 4, wherein the glycolate Type A is characterized by peaks in an X-ray diffraction (XRPD) pattern at 7.1±0.2, 10.1±0.2, and 11.2±0.2 °2θ.

6. The salt of embodiment 5, wherein the glycolate Type A is further characterized by at least one XRPD peak selected from 16.0±0.2, 18.0±0.2, 19.5±0.2, 20.9±0.2, 22.6±0.2, and 25.2±0.2 °2θ.

7. The salt of any of embodiments 4-6, wherein the glycolate Type A is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.1±0.2, 13.2±0.2, 16.0±0.2, 18.0±0.2, 19.5±0.2, 20.9±0.2, 22.6±0.2, and 25.2±0.2 °2θ.

8. The salt of any of embodiments 5-7, wherein the glycolate Type A is further characterized by at least one XRPD peak selected from 13.2±0.2, 14.1±0.2, 15.0±0.2, 18.5±0.2, 19.2±0.2, 19.7±0.2, 20.3±0.2, 23.7±0.2, 240±0.2, 27.6±0.2, 29.5±0.2, 30.1±0.2, 31.6±0.2, and 34.1±0.2 °2θ.

9. The salt of embodiment 4, wherein the glycolate Type A is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.1±0.2, and 11.2±0.2 °2θ and at least one XRPD peak selected from 16.0±0.2, 18.0±0.2, 19.5±0.2, 19.7±0.2, 20.3±0.2, 20.9±0.2, 22.6±0.2, 25.2±0.2 and 27.6±0.2 °2θ.

10. The salt of any of embodiments 4-9, wherein the glycolate Type A is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.1±0.2, 11.2±0.2, 13.2±0.2, 14.1±0.2, 15.1±0.2, 16.0±0.2, 18.0±0.2, 18.5±0.2, 19.2±0.2, 19.5±0.2, 19.7±0.2, 20.3±0.2, 20.9±0.2, 22.6±0.2, 23.7±0.2, 24.0±0.2, 25.2±0.2, 27.6±0.2, 29.5±0.2, 30.2±0.2, 31.6±0.2, and 34.1±0.2 °2θ.

11. The salt of any of embodiments 4-10, wherein the glycolate Type A exhibits a weight loss of about 1% up to a temperature of about 150 °C as measured by thermogravimetric (TGA) analysis.

12. The salt of any of embodiments 4-11, wherein the glycolate Type A exhibits a Differential Scanning Calorimetry (DSC) thermogram comprising an endotherm peak at about 222±2.5 °C.

13. The salt of embodiment 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type B.

14. The salt of embodiment 13, wherein the glycolate Type B is characterized by peaks in an XRPD pattern at 5.3±0.2, 5.7±0.2, and 7.5±0.2 °2θ.

15. The salt of embodiment 14, wherein the glycolate Type B is further characterized by at least one XRPD peak selected from 6.8±0.2, 10.8±0.2, 13.7±0.2, 19.9±0.2, 22.7±0.2, and 27.4±0.2 °2θ.

16. The salt of any of embodiments 13-15, wherein the glycolate Type B is characterized by peaks in an XRPD pattern at 5.3±0.2, 5.7±0.2, 6.8±0.2, 7.5±0.2, 10.8±0.2, 13.7±0.2, 19.9±0.2, 22.7±0.2, and 27.4±0.2 °2θ.

17. The salt of any of embodiments 14-16, wherein the glycolate Type B is further characterized by at least one XRPD peak selected from 9.0±0.2, 14.7±0.2, 17.4±0.2, 21.2±0.2, 24.1±0.2, and 25.4±0.2 °2θ.

18. The salt of embodiment 13, wherein the glycolate Type B is characterized by peaks in an XRPD pattern at 5.3±0.2, 5.7±0.2, and 7.5±0.2 °2θ and at least one XRPD peak selected from 10.8±0.2, 13.7±0.2, 19.9±0.2, 21.2±0.2, 22.7±0.2, and 24.0±0.2 °2θ.

19. The salt of any of embodiments 13-18, wherein the glycolate Type B is characterized by peaks in an XRPD pattern at 5.3±0.2, 5.7±0.2, 6.8±0.2, 7.5±0.2, 9.0±0.2, 10.8±0.2, 13.7±0.2, 14.7±0.2, 17.4±0.2, 19.9±0.2, 21.2±0.2, 22.7±0.2, 24.0±0.2, 25.4±0.2, and 27.4±0.2 °2θ.

20. The salt of any of embodiments 13-19, wherein the glycolate Type B exhibits a weight loss of about 4% up to a temperature of about 120 °C as measured by TGA analysis.

21. The salt of embodiment 20, wherein the glycolate Type B further exhibits a weight loss of about 8% between a temperature ranging from about 120 °C to about 160 °C as measured by TGA analysis.

22. The salt of any of embodiments 12-21, wherein the glycolate Type B exhibits a DSC thermogram comprising an endothermic peak at about 147±2.5 °C and 223±2.5 °C.

23. The salt of embodiment 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type C.

24. The salt of embodiment 23, wherein the glycolate Type C is characterized by peaks in an XRPD pattern at 6.0±0.2, 7.4±0.2, and 24.2±0.2 °2θ.

25. The salt of embodiment 24, wherein the glycolate Type C is further characterized by at least one XRPD peak selected from 14.2±0.2, 16.3±0.2, 18.1±0.2, 20.1±0.2, 26.2±0.2, and 27.6±0.2 °2θ.

26. The salt of any of embodiments 23-25, wherein the glycolate Type C is characterized by peaks in an XRPD pattern at 6.0±0.2, 7.4±0.2, 14.2±0.2, 16.3±0.2, 18.1±0.2, 20.1±0.2, 24.2±0.2, 26.2±0.2, and 27.6±0.2 °2θ.

27. The salt of any of embodiments 24-26, wherein the glycolate Type C is further characterized by at least one XRPD peak selected from 10.6±0.2, 11.0±0.2, 11.4±0.2, 12.8±0.2, 13.7±0.2, 15.7±0.2, 18.8±0.2, 21.2±0.2, 21.5±0.2, 22.6±0.2, 22.9±0.2, 25.1±0.2, 28.6±0.2, 29.4±0.2, 31.6±0.2, 33.7±0.2, 35.2±0.2, and 38.3±0.2 °2θ.

28. The salt of embodiment 23, wherein the glycolate Type C is characterized by peaks in an XRPD pattern at 6.0±0.2, 7.4±0.2, and 24.2±0.2 °2θ and at least one XRPD peak at 11.0±0.2, 11.4±0.2, 13.7±0.2, 14.2±0.2, 16.3±0.2, 18.1±0.2, 18.8±0.2, 20.1±0.2, 21.5±0.2, 22.9±0.2, and 26.2±0.2 °2θ.

29. The salt of any of embodiments 23-28, wherein the glycolate Type C is characterized by peaks in an XRPD pattern at 6.0±0.2, 7.4±0.2, 10.6±0.2, 11.0±0.2, 11.4±0.2, 12.8±0.2, 13.7±0.2, 14.2±0.2, 15.7±0.2, 16.3±0.2, 17.6±0.2, 18.1±0.2, 18.8±0.2, 20.1±0.2, 21.2±0.2, 21.5±0.2, 22.6±0.2, 22.9±0.2, 24.2±0.2, 25.2±0.2, 26.2±0.2, 27.6±0.2, 28.6±0.2, 29.4±0.2, 31.6±0.2, 33.7±0.2, 35.3±0.2, and 38.3±0.2 °2θ.

30. The salt of any of embodiments 23-29, wherein the glycolate Type C exhibits a weight less of about 6% up to a temperature of about 150 °C as measured by TGA.

31. The salt of any of embodiments 23-30, wherein the glycolate Type C exhibits a DSC thermogram comprising an endotherm peak at about 61±2.5 °C, 141±2.5 °C, and about 222±2.5 °C.

32. The salt of embodiment 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type D.

33. The salt of embodiment 32, wherein the glycolate Type D is characterized by peaks in an XRPD pattern at 5.3±0.2, 6.8±0.2, and 9.0±0.2 °2θ.

34. The salt of embodiment 33, wherein the glycolate Type D is further characterized by at least one XRPD peak selected from 11.0±0.2, 13.5±0.2, 17.3±0.2, 19.5±0.2, 20.1±0.2, and 21.3±0.2 °2θ.

35. The salt of any of embodiments 32-34, wherein the glycolate Type D is characterized by peaks in an XRPD pattern at 5.3±0.2, 6.8±0.2, 9.0±0.2, 11.0±0.2, 13.5±0.2, 17.3±0.2, 19.5±0.2, 20.1±0.2, and 21.3±0.2 °2θ.

36. The salt of any of embodiments 33-35, wherein the glycolate Type D is further characterized by at least one XRPD peak selected from 10.1±0.2, 11.3±0.2, 13.9±0.2, 16.5±0.2, 17.0±0.2, 18.0±0.2, 22.9±0.2, 23.7±0.2, 25.5±0.2, and 27.3±0.2 °2θ.

37. The salt of embodiment 32, wherein the glycolate Type D is characterized by peaks in an XRPD pattern at 5.3±0.2, 6.8±0.2, and 9.0±0.2 °2θ and at least one XRPD peak selected from 11.0±0.2, 11.3±0.2, 13.5±0.2, 13.9±0.2, 17.3±0.2, 19.5±0.2, 20.1±0.2, 21.3±0.2, and 23.7±0.2 °2θ.

38. The salt of any of embodiments 32-37, wherein the glycolate Type D is characterized peaks in an XRPD pattern at 5.3±0.2, 6.8±0.2, 9.0±0.2, 10.1±0.2, 11.0±0.2, 11.3±0.2, 13.5±0.2, 13.9±0.2, 16.5±0.2, 17.0±0.2, 17.3±0.2, 18.0±0.2, 19.5±0.2, 20.1±0.2, 21.3±0.2, 22.9±0.2, 23.7±0.2, 25.5±0.2, and 27.3±0.2 °2θ.

39. The salt of any of embodiments 32-38, wherein the glycolate Type D exhibits a weight less of about 3% up to a temperature of about 100 °C as measured by TGA.

40. The salt of any of embodiments 32-39, wherein the glycolate Type D exhibits a DSC thermogram comprising an endotherm peak at about 63±2.5 °C, about 210±2.5 °C, and about 123±2.5 °C.

41. The salt of embodiment 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type E.

42. The salt of embodiment 41, wherein the glycolate Type E is characterized by peaks in an XRPD pattern at 5.1±0.2, 7.8±0.2, and 8.8±0.2 °2θ.

43. The salt of embodiment 42, wherein the glycolate Type E is further characterized by at least one XRPD peak selected from 11.0±0.2, 12.0±0.2, 15.1±0.2, 16.8±0.2, 19.1±0.2, and 21.0±0.2 °2θ.

44. The salt of any of embodiments 41-43, wherein the glycolate Type E is characterized by peaks in an XRPD pattern at 5.1±0.2, 7.8±0.2, 8.8±0.2, 11.0±0.2, 12.0±0.2, 15.1±0.2, 16.8±0.2, 19.1±0.2, and 21.0±0.2 °2θ.

45. The salt of any of embodiments 42-44, wherein the glycolate Type E is further characterized by at least one XRPD peak selected from 18.4±0.2 and 22.3±0.2 °2θ.

46. The salt of embodiment 41, wherein the glycolate Type E is characterized by peaks in an XRPD pattern at 5.1±0.2, 7.8±0.2, and 8.8±0.2 °2θ and at least one peak selected from 11.0±0.2, 12.0±0.2, 16.8±0.2, 18.4±0.2, 19.1±0.2, 20.9±0.2, and 22.3±0.2 °2θ.

47. The salt of any of embodiments 41-46, wherein the glycolate Type E is characterized by peaks in an XRPD pattern at 5.1±0.2, 7.8±0.2, 8.8±0.2, 11.0±0.2, 12.0±0.2, 15.1±0.2, 16.8±0.2, 18.4±0.2, 19.1±0.2, 20.9±0.2, and 22.3±0.2 °2θ.

48. The salt of embodiment 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type F.

49. The salt of embodiment 48, wherein the glycolate Type F is characterized by peaks in an XRPD pattern at 5.9±0.2, 6.4±0.2, and 7.2±0.2 °2θ.

50. The salt of embodiment 49, wherein the glycolate Type F is further characterized by at least one XRPD peak selected from 13.4±0.2, 14.1±0.2, 18.4±0.2, 19.8±0.2, 24.7±0.2, and 23.9 ±0.2 °2θ.

51. The salt of any of embodiments 48-50, wherein the glycolate Type F is characterized by peaks in an XRPD pattern at 5.9±0.2, 6.4±0.2, 7.2±0.2, 13.4±0.2, 14.1±0.2, 18.4±0.2, 19.8±0.2, 24.7±0.2, and 23.9±0.2 °2θ.

52. The salt of any of embodiments 49-51, wherein the glycolate Type F is further characterized by at least one XRPD peak selected from 10.3±0.2, 10.8±0.2, 11.2±0.2, 12.3±0.2, 16.1±0.2, 16.8±0.2, 17.2±0.2, 17.8±0.2, 20.6±0.2, 21.4±0.2, 22.1±0.2, 24.4±0.2, 26.0±0.2, 26.4±0.2, and 27.2±0.2 °2θ.

53. The salt of embodiment 48, wherein the glycolate Type F is characterized by peaks in an XRPD pattern at 5.9±0.2, 6.4±0.2, and 7.2±0.2 °2θ and at least one XRPD

peak selected from 10.8±0.2, 11.2±0.2, 13.4±0.2, 14.1±0.2, 16.11±0.2, 17.8±0.2, 18.4±0.2, 19.8±0.2, 21.4±0.2, 22.1±0.2, 23.7±0.2, 23.9±0.2 and 24.4±0.2 °2θ.

54. The salt of any of embodiments 48-53, wherein the glycolate Type F is characterized by peaks in an XRPD pattern at 5.9±0.2, 6.4±0.2, 7.2±0.2, 10.3±0.2, 10.8±0.2, 11.2±0.2, 12.3±0.2, 13.4±0.2, 14.1±0.2, 16.1±0.2, 16.8±0.2, 17.2±0.2, 17.8±0.2, 18.4±0.2, 19.8±0.2, 20.6±0.2, 21.4±0.2, 22.1±0.2, 23.7±0.2, 23.9±0.2, 24.4±0.2, 26.0±0.2, 26.4±0.2, and 27.2±0.2 °2θ.

55. The salt of embodiment 1, wherein the anion is L-lactate.

56. The salt of embodiment 55, wherein the L-lactate salt is characterized by peaks in an XRPD pattern at 6.9±0.2, 10.0±0.2, and 11.0±0.2 °2θ.

57. The salt of embodiment 56, wherein the L-lactate salt is further characterized by at least one XRPD peak selected from 15.7±0.2, 20.6±0.2, 22.3±0.2, and 24.8±0.2 °2θ.

58. The salt of any of embodiments 55-57, wherein the L-lactate salt is characterized by peaks in an XRPD pattern at 6.9±0.2, 10.0±0.2, 11.0±0.2, 15.7±0.2, 20.6±0.2, 22.3±0.2, and 24.8±0.2 °2θ.

59. The salt of any of embodiments 56-58, wherein the L-lactate salt is further characterized by at least one XRPD peak selected from 10.7±0.2, 13.0±0.2, 13.8±0.2, 17.7±0.2, 18.1±0.2, 18.8±0.2, 19.3±0.2, 19.8±0.2, 23.6±0.2, 24.4±0.2, 27.0±0.2, 28.0±0.2, 29.3±0.2, 31.2±0.2, 33.8±0.2, 35.6±0.2°2θ.

60. The salt of any of embodiments 55-59, wherein the L-lactate salt is characterized by peaks in an XRPD pattern at 6.9±0.2, 10.0±0.2, 1.07±0.2, 11.0±0.2, 13.0±0.2, 13.8±0.2, 15.8±0.2, 17.7±0.2, 18.1±0.2, 18.8±0.2, 19.3±0.2, 19.8±0.2, 20.6±0.2, 22.3±0.2, 23.6±0.2, 24.4±0.2, 24.8±0.2, 27.0±0.2, 27.1±0.2, and 35.6±0.2 °2θ.

61. The salt of any of embodiments 55-60, wherein the L-lactate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 5**.

62. The salt of any of embodiments 55-61, wherein the L-lactate salt exhibits a weight less of about 1% up to a temperature of about 150 °C as measured by TGA.

63. The salt of any of embodiments 55-62, wherein the L-lactate salt exhibits a DSC thermogram comprising an endotherm peak at about 218±2.5 °C.

64. The salt of embodiment 1, wherein the anion is succinate.

65. The salt of embodiment 64, wherein the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 17.6±0.2, and 19.3±0.2 °2θ.

66. The salt of embodiment 65, wherein the succinate salt is further characterized by at least one XRPD peak selected 9.6±0.2, 21.7±0.2, 23.1±0.2, 25.5±0.2, and 25.9±0.2 °2θ.

67. The salt of any of embodiments 64-66, wherein the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 9.6±0.2, 17.6±0.2, 19.3±0.2, 21.7±0.2, 23.1±0.2, 25.5±0.2, and 25.9±0.2 °2θ.

68. The salt of any of embodiments 65-67, wherein the succinate salt is further characterized by at least one XRPD peak selected from 6.2±0.2, 10.1±0.2, 14.4±0.2, 15.7±0.2, 16.1±0.2, 16.9±0.2, 18.7±0.2, 21.2±0.2, 22.3±0.2, 23.7±0.2, 24.5±0.2, 25.0±0.2, 25.3±0.2, 27.0±0.2, 29.1±0.2, 30.5±0.2, 33.0±0.2, and 34.3±0.2 °2θ.

69. The salt of any of embodiments 64-68, wherein the succinate salt is characterized by peaks in an XRPD pattern at 6.2±0.2, 8.5±0.2, 9.6±0.2, 10.0±0.2, 14.4±0.2, 15.7±0.2, 16.1±0.2, 16.9±0.2, 17.6±0.2, 18.7±0.2, 19.3±0.2, 21.2±0.2, 21.7±0.2, 22.3±0.2, 23.1±0.2, 23.7±0.2, 24.5±0.2, 25.0±0.2, 25.3±0.2, 25.5±0.2, 25.9±0.2, 27.0±0.2, 29.1±0.2, 30.5±0.2, 31.3±0.2, 33.0±0.2, and 34.0±0.2 °2θ.

70. The salt of any of embodiments 64-70, wherein the succinate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 7**.

71. The salt of any of embodiments 64-69, wherein the succinate salt exhibits a weight less of about 2 % up to a temperature of about 150 °C as measured by TGA.

72. The salt of any of embodiments 64-70, wherein the succinate salt exhibits a DSC thermogram comprising an endothermic peak at about 198±2.5 °C and about 202±2.5 °C.

73. The salt of embodiment 1, wherein the anion is fumarate.

74. The salt of embodiment 73, wherein the fumarate salt is characterized by peaks in an XRPD pattern at 8.4±0.2, 17.5±0.2, and 19.2±0.2 °2θ.

75. The salt of embodiment 74, wherein the fumarate salt is further characterized by at least one XRPD peak selected from 9.6±0.2, 21.6±0.2, 25.4±0.2, 25.8±0.2, and 31.1±0.2 °2θ.

76. The salt of any of embodiments 73-75, wherein the fumarate salt is characterized by peaks in an XRPD pattern at 8.4±0.2, 9.6±0.2, 17.5±0.2, 19.2±0.2, 21.6±0.2, 25.4±0.2, 25.8±0.2, and 31.1±0.2 °2θ.

77. The salt of any of embodiments 74-76, wherein the fumarate salt is further characterized by at least one XRPD peak selected from 13.4±0.2, 14.4±0.2, 15.6±0.2, 16.2±0.2, 16.9±0.2, 18.7±0.2, 22.4±0.2, 23.0±0.2, 23.4±0.2, 23.8±0.2, 27.0±0.2, 28.9±0.2, 32.8±0.2, 34.0±0.2, and 38.0±0.2 °2θ.

78. The salt of any of embodiments 73-77, wherein the fumarate salt is characterized by peaks in an XRPD pattern at 8.4±0.2, 9.6±0.2, 13.4±0.2, 14.4±0.2, 15.6±0.2, 16.2±0.2, 16.9±0.2, 17.5±0.2, 18.7±0.2, 19.2±0.2, 21.6±0.2, 23.4±0.2, 23.0±0.2, 23.4±0.2, 23.8±0.2, 25.4±0.2, 25.8±0.2, 27.0±0.2, 28.9±0.2, 31.1±0.2, 32.8±0.2, 34.0±0.2, and 38.0±0.2 °2θ.

79. The salt of any of embodiments 73-78, wherein the fumarate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 4**.

80. The salt of any of embodiments 73-79, wherein the fumarate salt exhibits a weight less of about 1% up to a temperature of about 150 °C as measured by TGA.

81. The salt of any of embodiments 73-80, wherein the fumarate salt exhibits a DSC thermogram comprising an endothermic peak at about 255±2.5 °C.

82. The salt of embodiment 1, wherein the anion is mesylate.

83. The salt of embodiment 82, wherein the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 16.7±0.2, and 17.3±0.2 °2θ.

84. The salt of embodiment 83, wherein the mesylate salt is further characterized by at least one XRPD peak selected from 11.6±0.2, 13.3±0.2, 18.6±0.2, 18.9±0.2, and 20.0±0.2 °2θ.

85. The salt of any of embodiments 82-84, wherein the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 11.6±0.2, 13.3±0.2, 16.7±0.2, 17.3±0.2, 18.6±0.2, 18.9±0.2, and 20.0±0.2 °2θ.

86. The salt of any of embodiments 82-85, wherein the mesylate salt is further characterized by at least one XRPD peak selected from 8.2±0.2, 10.0±0.2, 14.9±0.2, 15.3±0.2, 19.8±0.2, 21.1±0.2, 21.4±0.2, 22.2±0.2, 22.7±0.2, 23.7±0.2, 24.4±0.2, 25.1±0.2, 26.0±0.2, 26.9±0.2, 28.5±0.2, and 32.8±0.2 °2θ.


87. The salt of any of embodiments 82-86, wherein the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 8.2±0.2, 10.0±0.2, 11.6±0.2, 13.3±0.2, 14.9±0.2, 15.3±0.2, 16.7±0.2, 17.3±0.2, 18.6±0.2, 18.9±0.2, 19.8±0.2, 20.0±0.2, 21.1±0.2, 21.4±0.2, 22.2±0.2, 22.7±0.2, 23.7±0.2, 24.4±0.2, 25.1±0.2, 26.0±0.2, 26.9±0.2, 28.5±0.2, and 32.8±0.2 °2θ.

88. The salt of any of embodiments 82-87, wherein the mesylate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 8**.

89. The salt of any of embodiments 82-88, wherein the mesylate salt exhibits a weight less of about 1% up to a temperature of about 150 °C as measured by TGA.

90. The salt of any of embodiments 82-89, wherein the mesylate salt exhibits a DSC thermogram comprising an endothermic peak at about 266±2.5 °C.

91. A salt of mitragynine of Formula II:



Formula II

wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

92. The salt of embodiment 91, wherein the anion is glycolate.

93. The salt of embodiment 92, wherein the glycolate salt is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.2±0.2, and 11.3±0.2 °2θ.

94. The salt of embodiment 93, wherein the glycolate salt is further characterized by at least one XRPD peak selected from 16.0±0.2, 18.0±0.2, 19.5±0.2, 20.9±0.2, 22.6±0.2, and 25.2±0.2 °2θ.

95. The salt of any of embodiments 92-94, wherein the glycolate salt is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.2±0.2, 11.3±0.2, 16.0±0.2, 18.0±0.2, 19.5±0.2, 20.9±0.2, 22.6±0.2, and 25.2±0.2 °2θ.

96. The salt of any of embodiments 93-95, wherein the glycolate salt is further characterized by at least one XRPD peak selected from 13.2±0.2, 14.1±0.2, 15.1±0.2, 15.7±0.2, 18.5±0.2, 18.9±0.2, 19.2±0.2, 19.7±0.2, 20.4±0.2, 23.3±0.2, 23.5±0.2, 23.4±0.2, 24.0±0.2, 24.9±0.2, 25.9±0.2, 27.6±0.2, 28.3±0.2, 29.1±0.2, 29.6±0.2, 30.2±0.2, 30.6±0.2, 32.7±0.2, 32.2±0.2, 34.3±0.2, 35.2±0.2, 36.0±0.2, and 36.6±0.2 °2θ.

97. The salt of any of embodiments 93-96, wherein the glycolate salt is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.2±0.2, 11.3±0.2, 13.2±0.2, 14.1±0.2, 15.1±0.2, 15.7±0.2, 16.0±0.2, 18.0±0.2, 18.5±0.2, 18.9±0.2 19.2±0.2, 19.7±0.2, 20.4±0.2, 23.3±0.2, 23.5±0.2, 23.8±0.2, 24.1±0.2, 24.9±0.2, 25.9±0.2, 27.6±0.2, 28.3±0.2, 29.1±0.2, 29.6±0.2, 30.2±0.2, 30.6±0.2, 31.7±0.2, 32.2±0.2, 34.2±0.2, 35.2±0.2, 36.0±0.2, and 36.6±0.2 °2θ.

98. The salt of any of embodiments 92-97, wherein the glycolate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 41**.

99. The salt of any of embodiments 92-98, wherein the glycolate salt exhibits a weight less of about 2% up to a temperature of about 150 °C as measured by TGA.

100. The salt of any of embodiments 91-99, wherein the glycolate salt exhibits a DSC thermogram comprising an endothermic peak at about 220±2.5 °C.

101. The salt of embodiment 91, wherein the anion is L-lactate.

102. The salt of embodiment 101, wherein the L-lactate salt is characterized by peaks in an XRPD pattern at 7.0±0.2, 10.1±0.2, and 11.2±0.2 °2θ.

103. The salt of embodiment 102, wherein the L-lactate salt is further characterized by at least one XRPD peak selected from 15.9±0.2, 17.9±0.2, 20.8±0.2, 22.4±0.2, and 24.9±0.2 °2θ.

104. The salt of any of embodiments 101-103, wherein the L-lactate salt is characterized by peaks in an XRPD pattern at 7.0±0.2, 10.1±0.2, 11.2±0.2, 15.9±0.2, 17.9±0.2, 20.8±0.2, 22.4±0.2, and 24.9 ±0.2 °2θ.

105. The salt of any of embodiments 102-104, wherein the L-lactate salt is further characterized by at least one XRPD peak selected from 10.9±0.2, 13.2±0.2, 13.9±0.2, 15.1±0.2, 15.6±0.2, 18.3±0.2, 19.0±0.2, 19.9±0.2, 21.2±0.2, 21.8±0.2, 22.9±0.2, 23.4±0.2, 23.8±0.2, 24.5±0.2, 25.8±0.2, 27.1±0.2, 27.3±0.2, 28.2±0.2, 29.5±0.2, 30.7±0.2, 31.4±0.2, 34.0±0.2, 35.7±0.2, 37.4±0.2, and 38.1±0.2 °2θ.

106. The salt of any of embodiments 101-105, wherein the L-lactate salt is characterized by peaks in an XRPD pattern at 7.0±0.2, 10.1±0.2, 10.9±0.2, 11.2±0.2, 13.2±0.2, 13.9±0.2, 15.1±0.2, 15.6±0.2, 15.9±0.2, 17.9±0.2, 18.3±0.2, 19.0±0.2, 19.9±0.2, 20.8±0.2, 21.2±0.2, 21.8±0.2, 22.4±0.2, 22.9±0.2, 23.4±0.2, 23.8±0.2, 24.5±0.2, 24.9±0.2, 25.8±0.2, 27.1±0.2, 27.3±0.2, 28.2±0.2, 29.5±0.2, 30.7±0.2, 31.4±0.2, 34.0±0.2, 35.7±0.2, 37.4±0.2, and 38.1±0.2 °2θ.

107. The salt of any of embodiments 101-106, wherein the L-lactate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 40**.

108. The salt of any of embodiments 101-107, wherein the L-lactate salt exhibits a weight less of about 3% up to a temperature of about 150 °C as measured by TGA.

109. The salt of any of embodiments 101-108, wherein the L-lactate salt exhibits a DSC thermogram comprising an endothermic peak at about 226±2.5 °C.

110. The salt of embodiment 91, wherein the anion is succinate.

111. The salt of embodiment 110, wherein the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 17.6±0.2, and 19.3±0.2 °2θ.

112. The salt of embodiment 111, wherein the succinate salt is further characterized by at least one XRPD peak selected from 9.6±0.2, 14.4±0.2, 21.7±0.2, 23.1±0.2, 25.5±0.2, and 25.9±0.2 °2θ.

113. The salt of any of embodiments 110-112, wherein the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 9.6±0.2, 14.4±0.2, 17.6±0.2, 21.7±0.2, 23.1±0.2, 25.5±0.2, and 25.9±0.2 °2θ.

114. The salt of any of embodiments 110-113, wherein the succinate salt is further characterized by at least one XRPD peak selected from 6.2±0.2, 9.1±0.2, 10.1±0.2, 13.5±0.2, 15.7±0.2, 16.1±0.2, 17.0±0.2, 18.7±0.2, 21.2±0.2, 22.3±0.2, 23.7±0.2, 24.5±0.2, 25.0±0.2, 25.3±0.2, 27.0±0.2, 28.6±0.2, 29.1±0.2, 30.5±0.2, 31.6±0.2, 33.0±0.2, 34.3±0.2, 34.6±0.2, 37.0±0.2, and 39.1±0.2 °2θ.

115. The salt of any of embodiments 110-114, wherein the succinate salt is characterized by peaks in an XRPD pattern at 6.2±0.2, 8.5±0.2, 9.1±0.2, 9.6±0.2, 10.1±0.2, 13.5±0.2, 14.4±0.2, 15.7±0.2, 16.1±0.2, 17.0±0.2, 17.6±0.2, 18.7±0.2, 21.2±0.2, 21.7±0.2, 22.3±0.2, 23.1±0.2, 23.7±0.2, 24.5±0.2, 25.0±0.2, 25.3±0.2, 25.5±0.2, 25.9±0.2, 27.0±0.2, 28.6±0.2, 29.1±0.2, 30.5±0.2, 31.6±0.2, 33.0±0.2, 34.3±0.2, 34.6±0.2, 37.0±0.2, and 39.1±0.2 °2θ.

116. The salt of any of embodiments 110-115, wherein the succinate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 42**.

117. The salt of any of embodiments 110-116, wherein the succinate salt exhibits a weight less of about 4% up to a temperature of about 150 °C as measured by TGA.

118. The salt of any of embodiments 110-117, wherein the succinate salt exhibits a DSC thermogram comprising an endothermic peak at about 198±2.5 °C and about 202±2.5 °C.

119. The salt of embodiment 91, wherein the anion is fumarate.

120. The salt of embodiment 119, wherein the fumarate salt is characterized by peaks in an XRPD pattern at 8.3±0.2, 19.1±0.2, and 19.2±0.2 °2θ.

121. The salt of embodiment 120, wherein the fumarate salt is further characterized by at least one XRPD peak selected from 14.3±0.2, 17.3±0.2, 18.6±0.2, 25.2±0.2, and 25.6±0.2 °2θ.

122. The salt of any of embodiments 119-121, wherein the fumarate salt is characterized by peaks in an XRPD pattern at 8.3±0.2, 14.3±0.2, 17.3±0.2, 18.6±0.2, 19.1±0.2, and 19.2±0.2, 25.2±0.2, and 25.6±0.2 °2θ.

123. The salt of any of embodiments 120-122, wherein the fumarate salt is further characterized by at least one XRPD peak selected from 9.5±0.2, 15.1±0.2, 15.5±0.2, 16.0±0.2, 16.7±0.2, 19.9±0.2, 21.2±0.2, 21.4±0.2, 22.2±0.2, 22.9±0.2, 23.2±0.2, 23.6±0.2, 24.5±0.2, 26.8±0.2, 28.8±0.2, 31.0±0.2, and 34.1±0.2 °2θ.

124. The salt of any of embodiments 119-123, wherein the fumarate salt is characterized by peaks in an XRPD pattern at 8.3±0.2, 9.5±0.2, 14.3±0.2, 15.1±0.2, 15.5±0.2, 16.0±0.2, 16.7±0.2, 17.3±0.2, 18.6±0.2, 19.1±0.2, 19.2±0.2, 19.9±0.2, 21.2±0.2, 21.4±0.2, 22.2±0.2, 22.9±0.2, 23.2±0.2, 23.6±0.2, 24.5±0.2, 25.3±0.2, 25.6±0.2, 26.8±0.2, 28.8±0.2, 31.0±0.2, and 34.1±0.2 °2θ.

125. The salt of any of embodiments 119-124, wherein the fumarate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 39**.

126. The salt of any of embodiments 119-125, wherein the fumarate salt exhibits a weight less of about 3% up to a temperature of about 150 °C as measured by TGA.

127. The salt of any of embodiments 119-126, wherein the fumarate salt exhibits a DSC thermogram comprising an endothermic peak at about 226±2.5 °C.

128. The salt of embodiment 91, wherein the anion is mesylate.

129. The salt of embodiment 128, wherein the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 16.7±0.2, and 17.4±0.2 °2θ.

130. The salt of embodiment 129, wherein the mesylate salt is further characterized by at least one XRPD peak selected from 11.6±0.2, 18.6±0.2, 18.9±0.2, 20.1±0.2, and 26.0±0.2 °2θ.

131. The salt of any of embodiments 128-130, wherein the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 11.6±0.2, 16.7±0.2, 17.4±0.2, 18.6±0.2, 18.9±0.2, 20.1±0.2, and 26.0±0.2 °2θ.

132. The salt of any of embodiments 129-131, wherein the mesylate salt is further characterized by at least one XRPD peak selected from 8.2±0.2, 10.0±0.2, 13.0±0.2, 13.4±0.2, 15.3±0.2, 16.4±0.2, 18.3±0.2, 19.8±0.2, 21.2±0.2, 21.4±0.2, 22.2±0.2, 22.7±0.2, 23.0±0.2, 23.7±0.2, 24.2±0.2, 24.4±0.2, 24.6±0.2, 25.1±0.2, 26.8±0.2, 27.1±0.2, 28.5±0.2, 30.1±0.2, 32.9±0.2, 33.7±0.2, and 37.1±0.2 °2θ.

133. The salt of any of embodiments 128-132, wherein the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 8.2±0.2, 10.0±0.2, 11.6±0.2, 13.0±0.2, 13.4±0.2, 15.3±0.2, 16.4±0.2, 16.7±0.2, 17.4±0.2, 18.3±0.2, 18.6±0.2, 18.9±0.2, 19.8±0.2, 20.1±0.2, 21.2±0.2, 21.4±0.2, 22.2±0.2, 22.7±0.2, 23.0±0.2, 23.7±0.2, 24.2±0.2, 24.4±0.2, 24.6±0.2, 25.1±0.2, 26.0±0.2, 26.8±0.2, 27.1±0.2, 28.5±0.2, 30.1±0.2, 32.9±0.2, 33.7±0.2, and 37.1±0.2 °2θ.

134. The salt of any of embodiments 128-133, wherein the mesylate salt is characterized by an XRPD pattern substantially similar to that shown in **Figure 43**.

135. The salt of any of embodiments 128-134, wherein the mesylate salt exhibits a weight less of about 2% up to a temperature of about 150 °C as measured by TGA.

136. The salt of any of embodiments 128-135, wherein the mesylate salt exhibits a DSC thermogram comprising an endothermic peak at about 275±2.5 °C.

137. A pharmaceutical composition comprising a salt of any of embodiments 1-136.

138. The pharmaceutical composition of embodiment 137 further comprising a pharmaceutically acceptable excipient.

139. A method of treating a subject afflicted with acute pain, chronic pain, a depressive disorder, a mood disorder, an anxiety disorder, borderline personality

disorder, a substance use disorder, opioid use disorder, opioid withdrawal symptoms, alcohol use disorder, or alcohol withdrawal disorder, comprising administering an effective amount of a salt of any of embodiments 1-136 or a pharmaceutical composition of embodiments 137-138 to the subject.

140. The method of embodiment 139, wherein the subject is afflicted with an opioid use disorder.


141. The method of embodiment 139, wherein the subject is afflicted with opioid withdrawal.

142. The method of any one of embodiments 139-142, wherein the subject is administered about 10 mg to about 90 mg of a salt of any of embodiments 1-136.

143. The method of embodiment 142, wherein the subject is administered a glycolate salt of 3-deuteromitragynine of any one of embodiments 2-54.

What is claimed is:

1. A salt of 3-deuteromitragynine of Formula I:



Formula I

wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

2. The salt of claim 1, wherein the anion is glycolate.

3. The salt of claim 2, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type A, glycolate Type B, glycolate Type C, glycolate Type D, glycolate Type E, glycolate Type F, or combinations thereof.

4. The salt of claim 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type A.

5. The salt of claim 4, wherein the glycolate Type A is characterized by peaks in an X-ray diffraction (XRPD) pattern at 7.1±0.2, 10.1±0.2, and 11.2±0.2 °2θ.

6. The salt of claim 5, wherein the glycolate Type A is further characterized by at least one XRPD peak selected from 16.0±0.2, 18.0±0.2, 19.5±0.2, 20.9±0.2, 22.6±0.2, and 25.2±0.2 °2θ.

7. The salt of claim 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type B.

8. The salt of claim 7, wherein the glycolate Type B is characterized by peaks in an XRPD pattern at 5.3±0.2, 5.7±0.2, and 7.5±0.2 °2θ.

9. The salt of claim 8, wherein the glycolate Type B is further characterized by at least one XRPD peak selected from 6.8±0.2, 10.8±0.2, 13.7±0.2, 19.9±0.2, 22.7±0.2, and 27.4±0.2 °2θ.

10. The salt of claim 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type C.

11. The salt of claim 10, wherein the glycolate Type C is characterized by peaks in an XRPD pattern at 6.0±0.2, 7.4±0.2, and 24.2±0.2 °2θ.

12. The salt of claim 11, wherein the glycolate Type C is further characterized by at least one XRPD peak selected from 14.2±0.2, 16.3±0.2, 18.1±0.2, 20.1±0.2, 26.2±0.2, and 27.6±0.2 °2θ.

13. The salt of claim 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type D.

14. The salt of claim 13, wherein the glycolate Type D is characterized by peaks in an XRPD pattern at 5.3±0.2, 6.8±0.2, and 9.0±0.2 °2θ.

15. The salt of claim 14, wherein the glycolate Type D is further characterized by at least one XRPD peak selected from 11.0±0.2, 13.5±0.2, 17.3±0.2, 19.5±0.2, 20.1±0.2, and 21.3±0.2 °2θ.

16. The salt of claim 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type E.

17. The salt of claim 16, wherein the glycolate Type E is characterized by peaks in an XRPD pattern at 5.1±0.2, 7.8±0.2, and 8.8±0.2 °2θ.

18. The salt of claim 17, wherein the glycolate Type E is further characterized by at least one XRPD peak selected from 11.0±0.2, 12.0±0.2, 15.1±0.2, 16.8±0.2, 19.1±0.2, and 21.0±0.2 °2θ.

19. The salt of claim 3, wherein the salt of 3-deuteromitragynine of Formula I is glycolate Type F.

20. The salt of claim 19, wherein the glycolate Type F is characterized by peaks in an XRPD pattern at 5.9±0.2, 6.4±0.2, and 7.2±0.2 °2θ.

21. The salt of claim 20, wherein the glycolate Type F is further characterized by at least one XRPD peak selected from 13.4±0.2, 14.1±0.2, 18.4±0.2, 19.8±0.2, 24.7±0.2, and 23.9 ±0.2 °2θ.

22. The salt of claim 1, wherein the anion is L-lactate.

23. The salt of claim 22, wherein the L-lactate salt is characterized by peaks in an XRPD pattern at 6.9±0.2, 10.0±0.2, and 11.0±0.2 °2θ.

24. The salt of claim 23, wherein the L-lactate salt is further characterized by at least one XRPD peak selected from 15.7±0.2, 20.6±0.2, 22.3±0.2, and 24.8±0.2 °2θ.

25. The salt of claim 1, wherein the anion is succinate.

26. The salt of claim 25, wherein the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 17.6±0.2, and 19.3±0.2 °2θ.

27. The salt of claim 26, wherein the succinate salt is further characterized by at least one XRPD peak selected 9.6±0.2, 21.7±0.2, 23.1±0.2, 25.5±0.2, and 25.9±0.2 °2θ.

28. The salt of claim 1, wherein the anion is fumarate.

29. The salt of claim 28, wherein the fumarate salt is characterized by peaks in an XRPD pattern at 8.4±0.2, 17.5±0.2, and 19.2±0.2 °2θ.


30. The salt of claim 29, wherein the fumarate salt is further characterized by at least one XRPD peak selected from 9.6±0.2, 21.6±0.2, 25.4±0.2, 25.8±0.2, and 31.1±0.2 °2θ.

31. The salt of claim 1, wherein the anion is mesylate.

32. The salt of claim 31, wherein the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 16.7±0.2, and 17.3±0.2 °2θ.

33. The salt of claim 32, wherein the mesylate salt is further characterized by at least one XRPD peak selected from 11.6±0.2, 13.3±0.2, 18.6±0.2, 18.9±0.2, and 20.0±0.2 °2θ.

34. A salt of mitragynine of Formula II



Formula II

wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate.

35. The salt of claim 34, wherein the anion is glycolate.

36. The salt of claim 35, wherein the glycolate salt is characterized by peaks in an XRPD pattern at 7.1±0.2, 10.2±0.2, and 11.3±0.2 °2θ.

37. The salt of claim 36, wherein the glycolate salt is further characterized by at least one XRPD peak selected from 16.0±0.2, 18.0±0.2, 19.5±0.2, 20.9±0.2, 22.6±0.2, and 25.2±0.2 °2θ.

38. The salt of claim 34, wherein the anion is L-lactate.

39. The salt of claim 38, wherein the L-lactate salt is characterized by peaks in an XRPD pattern at 7.0±0.2, 10.1±0.2, and 11.2±0.2 °2θ.

40. The salt of claim 39, wherein the L-lactate salt is further characterized by at least one XRPD peak selected from 15.9±0.2, 17.9±0.2, 20.8±0.2, 22.4±0.2, and 24.9±0.2 °2θ.

41. The salt of claim 34, wherein the anion is succinate.

42. The salt of claim 41, wherein the succinate salt is characterized by peaks in an XRPD pattern at 8.5±0.2, 17.6±0.2, and 19.3±0.2 °2θ.

43. The salt of claim 42, wherein the succinate salt is further characterized by at least one XRPD peak selected from 9.6±0.2, 14.4±0.2, 21.7±0.2, 23.1±0.2, 25.5±0.2, and 25.9±0.2 °2θ.

44. The salt of claim 34, wherein the anion is fumarate.

45. The salt of claim 44, wherein the fumarate salt is characterized by peaks in an XRPD pattern at 8.3±0.2, 19.1±0.2, and 19.2±0.2 °2θ.

46. The salt of claim 45, wherein the fumarate salt is further characterized by at least one XRPD peak selected from 14.3±0.2, 17.3±0.2, 18.6±0.2, 25.2±0.2, and 25.6±0.2 °2θ.

47. The salt of claim 34, wherein the anion is mesylate.

48. The salt of claim 47, wherein the mesylate salt is characterized by peaks in an XRPD pattern at 6.7±0.2, 16.7±0.2, and 17.4±0.2 °2θ.

49. The salt of claim 48, wherein the mesylate salt is further characterized by at least one XRPD peak selected from 11.6±0.2, 18.6±0.2, 18.9±0.2, 20.1±0.2, and 26.0±0.2 °2θ.

50. A pharmaceutical composition comprising a salt of any of claims 1-49.

51. The pharmaceutical composition of claim 50, further comprising a pharmaceutically acceptable excipient.

52. A method of treating a subject afflicted with acute pain, chronic pain, a depressive disorder, a mood disorder, an anxiety disorder, borderline personality disorder, a substance use disorder, opioid use disorder, opioid withdrawal symptoms, alcohol use disorder, or alcohol withdrawal disorder, comprising administering an effective amount of a salt of any of claims 1-49 or a pharmaceutical composition of claims 50-51 to the subject.

53. The method of claim 52, wherein the subject is afflicted with an opioid use disorder.

54. The method of claim 52, wherein the subject is afflicted with opioid withdrawal.

Figure 1 – XRPD pattern of amorphous 3-DM free base starting material




Figure 2 – TGA/mDSC curves of amorphous 3-DM free base starting material (810397-01-A)




Figure 3 - ^{1}H NMR spectrum of amorphous 3-DM free base starting material (810397-01-A)




Figure 4 – XRPD pattern of Fumarate Type A 3-DM Salt




Figure 5 – XRPD pattern of L-Lactate Type A 3-DM Salt




Figure 6 – XRPD pattern of Glycolate Type A 3-DM Salt




Figure 7 – XRPD pattern of Succinate Type A 3-DM Salt




Figure 8– XRPD pattern of Mesylate Type A 3-DM Salt




Figure 9 – XRPD pattern of HCl Type A 3-DM Salt (810397-04-B1)




Figure 10 - XRPD pattern of HCl Type B 3-DM Salt (810397-04-D1)




Figure 11 - XRPD pattern of Phosphate Type A 3-DM Salt (810397-04-D3)




Figure 12 - XRPD pattern of Maleate Type A 3-DM Salt (810397-04-C4)




Figure 13 - XRPD pattern of L-Tartrate Type A 3-DM Salt (810397-04-B5)




Figure 14 - XRPD pattern of Glycolate Type B 3-DM Salt (810397-04-C8)




Figure 15 - XRPD pattern of Adipate Type A 3-DM Salt (810397-04-A12)




Figure 16 - XRPD pattern of Adipate Type B 3-DM Salt (810397-04-C12)




Figure 17 - XRPD pattern of Acetate Type A 3-DM Salt (810397-04-D13)




Figure 18 - XRPD pattern of Malonate Type A 3-DM Salt (810397-04-B17)




Figure 19 - XRPD pattern of Malonate Type B 3-DM Salt (810397-04-D17)




Figure 20 - XRPD pattern of Gentisate Type A 3-DM Salt (810397-04-B18)




Figure 21 - XRPD pattern of Gentisate Type B 3-DM Salt (810397-04-C18)




Figure 22 - XRPD pattern of Edisylate Type A 3-DM Salt (810397-04-A19)




Figure 23 - XRPD pattern of Edisylate Type B 3-DM Salt (810397-04-C19)




Figure 24 - XRPD pattern of Benzoate Type A 3-DM Salt (810397-04-D20)




Figure 25 - XRPD pattern of Esylate Type A 3-DM Salt (810397-04-C21)




Figure 26 - XRPD pattern of Besylate Type A 3-DM Salt (810397-04-D23)




Figure 27 - XRPD pattern of Tosylate Type A 3-DM Salt (810397-04-B24)




Figure 28 - XRPD pattern of Oxalate Type A 3-DM Salt (810397-04-A25)




Figure 29 - XRPD pattern of Oxalate Type B 3-DM Salt (810397-04-B25)




Figure 30 - XRPD overlay of 3-DM Glycolate crystal forms




Figure 31 – Polarized light microscopy (PLM) image of 3-DM Glycolate Type A starting material used in crystallization process development experiments (**Examples 18-21**)




Figure 32 – XRPD pattern of 3-DM Free Base Type A (810081-04-C1_dry)




Figure 33 - Polarized light microscopy (PLM) image of 3-DM Glycolate Type A prepared at 5-g scale (810082-25-B)




Figure 34 – XRPD pattern of amorphous mitragynine free base starting material (810080-01-A)




Figure 35 - TGA/mDSC curves of amorphous mitragynine free base starting material (810080-01-A)




Figure 36 - ^{1}H NMR spectrum of amorphous mitragynine free base starting material (810080-01-A)




Figure 37 – XRPD pattern of crude alkaloid extract starting material (810080-01-B)




Figure 38 - TGA curve of crude alkaloid extract starting material (810080-01-B)




Figure 39 – XRPD pattern of Fumarate Type A Mitragynine Salt




Figure 40 – XRPD pattern of L-Lactate Type A Mitragynine Salt




Figure 41 – XRPD pattern of Glycolate Type A Mitragynine Salt




Figure 42 – XRPD pattern of Succinate Type A Mitragynine Salt





Figure 43 – XRPD pattern of Mesylate Type A Mitragynine Salt



INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB 22/54187

A. CLASSIFICATION OF SUBJECT MATTER

IPC - INV. A61K 31/437, A61K 31/4375, C07D 471/14; ADD. A61P 25/00 (2022.01)

CPC - INV. A61K 31/437, A61K 31/4375, C07D 471/14; ADD. A61P 25/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
See Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y --- A	WO 2020/160280 A1 (The Trustees of Columbia University in the City of New York, et al.) 06 August 2020 (06.08.2020) pg. 48, ln 5-10;pg. 57, ln 12-14;pg. 66, ln 22 to pg. 67, ln 10;pg. 68, ln 20-30	1-2; (50-51)/(1-2) ---------- 3-21; (50-51)/(3-21)
Y --- A	WO 2016/176657 A1 (Memorial Sloan-Kettering Cancer Center) 03 November 2016 (03.11.2016) para [0070];[0110]	1-2; (50-51)/(1-2) ---------- 3-21; (50-51)/(3-21)
A	ZACHARIAS, et al. 'The Structure of Mitragynine Hydrochloride', Acta Cryst., 1965, Vol.18, pp. 1039-1043. Abstract; pg 1039, Col 2, para 23, Table 1	3-21; (50-51)/(3-21)
A	US 2019/0084983 A1 (The Trustees of Columbia University in the City of New York) 21 March 2019 (21.03.2019) para [0370];[0457]	3-21; (50-51)/(3-21)
A	LIU et al. 'Computational Study on the Conformations of Mitragynine and Mitragynaline', Theochem., 2010, Vol. 945(1-3), pp. 57-63. ENTIRE DOCUMENT	1-21; 50;51

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:
"A" document defining the general state of the art which is not considered to be of particular relevance
"D" document cited by the applicant in the international application
"E" earlier application or patent but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed
"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
17 JULY 2022	OCT 05 2022
Name and mailing address of the ISA/US Mail Stop PCT, Attn: ISA/US, Commissioner for Patents P.O. Box 1450, Alexandria, Virginia 22313-1450 Facsimile No. 571-273-8300	Authorized officer Kari Rodriquez Telephone No. PCT Helpdesk: 571-272-4300

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB 22/54187

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 52-54
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
--Please see attached sheet---

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-21, 50 and 51

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.

☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.

☐ No protest accompanied the payment of additional search fees.

Attachment to Box.No.III:

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I+: Claims 1-51, directed to a salt of 3-deuteromitragynine of Formula (I), wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate; a salt of mitragynine of Formula II, wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate; and to a pharmaceutical composition comprising said salt. The salt and pharmaceutical composition will be searched to the extent that the salt encompasses a glycolate salt of 3-deuteromitragynine of Formula (I). It is believed that claims 1-21, 50 and 51 read on this first named invention, and thus these claims will be searched without fee to the extent that they encompass a glycolate salt of 3-deuteromitragynine of Formula (I). This first named invention has been selected based on the guidance set forth in section 10.54 of the PCT International Search and Preliminary Examination Guidelines.Applicant is invited to elect additional salt(s) wherein each additional salt elected will require one additional invention fee. Applicants must specify the claims that encompass any additionally elected salt. Applicants must further indicate, if applicable, the claims which encompass the first named invention, if different than what was indicated above for this group. Failure to clearly identify how any paid additional invention fees are to be applied to the '+' group(s) will result in only the first claimed invention to be searched. Additionally, an exemplary election wherein different actual variables are selected is suggested. An exemplary election would be a L-lactate salt of 3-deuteromitragynine of Formula (I) and a pharmaceutical composition comprising said salt (i.e., claims 1, 22-24, 50 and 51).

The group of inventions listed above do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Special Technical Features:

Group I+ includes the technical feature of a unique salt of 3-deuteromitragynine or mitragynine, which is not required by any other invention of Group I+.

Common technical features:

The inventions of Group I+ share the technical feature of a salt of 3-deuteromitragynine of Formula (I), or a salt of mitragynine of Formula II, wherein the anion is glycolate, L-lactate, succinate, fumarate, or mesylate; and a pharmaceutical composition comprising said salt.

These shared technical features, however, do not provide a contribution over the prior art, as being obvious over WO 2020/160280 A1 to The Trustees of Columbia University in the City of New York, et al. (hereinafter "Columbia"), which discloses a pharmaceutical composition comprising a salt of 3-deuteromitragynine of formula I (pg. 19, ln 5 to pg. 20, ln 7; pg. 20, ln 24, first structure; pg. 57, ln 6-7), wherein the salt can be a lactate, succinate or mesylate (pg. 66, ln 22 to pg. 67, ln 10). Columbia does not disclose a specific example of the salt of formula I. However, it would have been obvious to one of ordinary skill in the art to combine the teachings of Columbia with routine experimentation to make a lactate, succinate or mesylate salt of formula I for use in the pharmaceutical composition, in order to enhance the solubility of said compound and the efficacy of the resulting composition toward treating a neurological disorder (pg 37, ln 4-9), because Columbia discloses that these anions can be used to make the salt of the 3-deuteromitragynine of formula I.

As said salt and pharmaceutical composition were known in the art at the time of the invention, these cannot be considered special technical features, that would otherwise unify the inventions of Group I+.

The inventions of Group I+, thus, lack unity under PCT Rule 13.

Note reg. item 4: Claims 52-54 are unsearchable because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).