

STAA552 Homework 1

Due Friday, October 28, 2016.

1. Read Sections 1.1, 1.2, 1.3, 1.4 (you can skip 1.4.4 if you like), 1.5.1–1.5.5 and 2.1 of Agresti.
2. This is a HUGE EDIT.
3. Yet another edit.
4. Complete exercises 1.2, 2.1, 2.2a–b, and 2.3 of Agresti.
5. One version of the *prosecutor's fallacy* occurs when the probability of some evidence given that a defendant is not guilty is presented as if it were the probability that the defendant is not guilty given the evidence. For example, let M be the event that DNA known to have been left by the perpetrator at a crime scene matches the DNA of the defendant. Let G denote the event that the defendant is guilty and \bar{G} the event that the defendant is innocent. Suppose that $P(M \mid \bar{G}) = 1 \times 10^{-6}$ (that is, the DNA could match by chance even if the defendant is not guilty). Because the defendant's DNA matches the crime scene DNA, the prosecutor claims that there is a one in a million chance that the defendant is not guilty. Suppose that the weight of all other evidence leads to $P(G) = 1 \times 10^{-4}$ (for example, other evidence narrows the set of suspects to 10,000 individuals, one of whom is the defendant). Use the provided information to compute $P(\bar{G} \mid M)$. Comment on your result and the prosecutor's claim.
6. Use the data of Example 1.5.4 in Agresti. Assume that the count of yellow seeds among the $n = 8023$ hybrid seeds in Mendel's experiment follows a $\text{Binomial}(n, \pi)$ distribution.
 - (a) Plot the log-likelihood function of the unknown parameter π given the data, for a fine grid of values in the interval $(0, 1)$.
 - (b) Zoom in by plotting the log-likelihood function of the unknown parameter π given the data, for a fine grid of values in the interval $(0.65, 0.85)$. Large-sample normal approximations rely on the approximate quadratic shape of the log-likelihood in a neighborhood of the true value. Does this quadratic approximation appear plausible here?

- (c) Add a vertical line to your plot to indicate the location of $\hat{\pi}$, the maximum likelihood estimator of π .
 - (d) Use equation (1.11) of Agresti to compute the score statistic for testing the null hypothesis $H_0 : \pi = 0.75$ versus the alternative $H_a : \pi \neq 0.75$. Compute the p -value of your test statistic using the normal approximation, and compute the p -value of your *squared* test statistic using the χ_1^2 approximation. Interpret your results.
 - (e) Use equation (1.16) of Agresti to compute Pearson's chi-squared statistic X^2 for Mendel's data. Compute the p -value of the test statistic using the χ_1^2 approximation, and compare to the results of the previous problem (5d).
7. Complete exercise 1.10 of Agresti. Use five categories: 0 deaths, 1 death, 2 deaths, 3 deaths, or ≥ 4 deaths, and note that you must estimate one parameter (see Agresti §1.5.5).