


## Review of Lecture 5


- Dichotomies



- Growth function

$$m_{\mathcal{H}}(N) = \max_{\mathbf{x}_1, \dots, \mathbf{x}_N \in \mathcal{X}} |\mathcal{H}(\mathbf{x}_1, \dots, \mathbf{x}_N)|$$

- Break point



- Maximum # of dichotomies

$\mathbf{x}_1$	$\mathbf{x}_2$	$\mathbf{x}_3$
○	○	○
○	○	●
○	●	○
●	○	○


# Learning From Data

Yaser S. Abu-Mostafa  
*California Institute of Technology*

## Lecture 6: Theory of Generalization



Sponsored by Caltech's Provost Office, E&AS Division, and IST • Thursday, April 19, 2012



# Outline

- Proof that  $m_{\mathcal{H}}(N)$  is polynomial
- Proof that  $m_{\mathcal{H}}(N)$  can replace  $M$

# Bounding $m_{\mathcal{H}}(N)$

To show:  $m_{\mathcal{H}}(N)$  is polynomial

We show:  $m_{\mathcal{H}}(N) \leq \dots \leq \dots \leq$  a polynomial

Key quantity:

$B(N, k)$ : Maximum number of dichotomies on  $N$  points, with break point  $k$

# Recursive bound on $B(N, k)$

Consider the following table:

$$B(N, k) = \alpha + 2\beta$$

	# of rows	$\mathbf{x}_1$	$\mathbf{x}_2$	$\dots$	$\mathbf{x}_{N-1}$	$\mathbf{x}_N$
$S_1$	$\alpha$	+1	+1	$\dots$	+1	+1
		-1	+1	$\dots$	+1	-1
		$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
		+1	-1	$\dots$	-1	-1
		-1	+1	$\dots$	-1	+1
$S_2$	$\beta$	+1	-1	$\dots$	+1	+1
		-1	-1	$\dots$	+1	+1
		$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
		+1	-1	$\dots$	+1	+1
		-1	-1	$\dots$	-1	+1
$S_2^-$	$\beta$	+1	-1	$\dots$	+1	-1
		-1	-1	$\dots$	+1	-1
		$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
		+1	-1	$\dots$	+1	-1
		-1	-1	$\dots$	-1	-1

# Estimating $\alpha$ and $\beta$

Focus on  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{N-1}$  columns:

$$\alpha + \beta \leq B(N-1, k)$$

	# of rows	$\mathbf{x}_1$	$\mathbf{x}_2$	$\dots$	$\mathbf{x}_{N-1}$	$\mathbf{x}_N$
$S_1$	$\alpha$	+1	+1	$\dots$	+1	+1
		-1	+1	$\dots$	+1	-1
		$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
		+1	-1	$\dots$	-1	-1
		-1	+1	$\dots$	-1	+1
$S_2$	$\beta$	+1	-1	$\dots$	+1	+1
		-1	-1	$\dots$	+1	+1
		$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
		+1	-1	$\dots$	+1	+1
		-1	-1	$\dots$	-1	+1
$S_2^-$	$\beta$	+1	-1	$\dots$	+1	-1
		-1	-1	$\dots$	+1	-1
		$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
		+1	-1	$\dots$	+1	-1
		-1	-1	$\dots$	-1	-1

# Estimating $\beta$ by itself

Now, focus on the  $S_2 = S_2^+ \cup S_2^-$  rows:

$$\beta \leq B(N-1, k-1)$$

	# of rows	$\mathbf{x}_1$	$\mathbf{x}_2$	$\dots$	$\mathbf{x}_{N-1}$	$\mathbf{x}_N$
$S_1$	$\alpha$	+1	+1	$\dots$	+1	+1
		-1	+1	$\dots$	+1	-1
		$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
		+1	-1	$\dots$	-1	-1
		-1	+1	$\dots$	-1	+1
$S_2$	$S_2^+$ $\beta$	+1	-1	$\dots$	+1	+1
		-1	-1	$\dots$	+1	+1
		$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
		+1	-1	$\dots$	+1	+1
		-1	-1	$\dots$	-1	+1
$S_2^-$ $\beta$	$\beta$	+1	-1	$\dots$	+1	-1
		-1	-1	$\dots$	+1	-1
		$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
		+1	-1	$\dots$	+1	-1
		-1	-1	$\dots$	-1	-1

# Putting it together

$$B(N, k) = \alpha + 2\beta$$

$$\alpha + \beta \leq B(N - 1, k)$$

$$\beta \leq B(N - 1, k - 1)$$

$$B(N, k) \leq$$


$$B(N - 1, k) + B(N - 1, k - 1)$$

		# of rows	$\mathbf{x}_1$	$\mathbf{x}_2$	$\dots$	$\mathbf{x}_{N-1}$	$\mathbf{x}_N$
$S_1$	$\alpha$		+1	+1	$\dots$	+1	+1
			-1	+1	$\dots$	+1	-1
			$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
			+1	-1	$\dots$	-1	-1
			-1	+1	$\dots$	-1	+1
$S_2$	$S_2^+$	$\beta$	+1	-1	$\dots$	+1	+1
			-1	-1	$\dots$	+1	+1
			$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
			+1	-1	$\dots$	+1	+1
			-1	-1	$\dots$	-1	+1
	$S_2^-$	$\beta$	+1	-1	$\dots$	+1	-1
			-1	-1	$\dots$	+1	-1
			$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
			+1	-1	$\dots$	+1	-1
			-1	-1	$\dots$	-1	-1



# Numerical computation of $B(N, k)$ bound

$$B(N, k) \leq B(N - 1, k) + B(N - 1, k - 1)$$



		$k$						
		1	2	3	4	5	6	..
$N$	1	1	2	2	2	2	2	..
	2	1	3	4	4	4	4	..
	3	1	4	7	8	8	8	..
	4	1	5	11	..	..	..	..
	5	1	6	:	.			
	6	1	7	:		.		
	:	:	:	:			.	

# Analytic solution for $B(N, k)$ bound


$$B(N, k) \leq B(N - 1, k) + B(N - 1, k - 1)$$

Theorem:

$$B(N, k) \leq \sum_{i=0}^{k-1} \binom{N}{i}$$


1. Boundary conditions: easy

		$k$						
		1	2	3	4	5	6	..
$N$	1	1	2	2	2	2	2	..
	2	1						
	3	1						
	4	1						
	5	1						
	6	1						
	:	:						



## 2. The induction step

$$\begin{aligned}\sum_{i=0}^{k-1} \binom{N}{i} &= \sum_{i=0}^{k-1} \binom{N-1}{i} + \sum_{i=0}^{k-2} \binom{N-1}{i} \text{ ?} \\ &= 1 + \sum_{i=1}^{k-1} \binom{N-1}{i} + \sum_{i=1}^{k-1} \binom{N-1}{i-1} \\ &= 1 + \sum_{i=1}^{k-1} \left[ \binom{N-1}{i} + \binom{N-1}{i-1} \right] \\ &= 1 + \sum_{i=1}^{k-1} \binom{N}{i} = \sum_{i=0}^{k-1} \binom{N}{i} \checkmark\end{aligned}$$



# It is polynomial!

For a given  $\mathcal{H}$ , the break point  $k$  is fixed

$$m_{\mathcal{H}}(N) \leq \underbrace{\sum_{i=0}^{k-1} \binom{N}{i}}_{\text{maximum power is } N^{k-1}}$$

## Three examples

$$\sum_{i=0}^{k-1} \binom{N}{i}$$

- $\mathcal{H}$  is positive rays: (break point  $k = 2$ )

$$m_{\mathcal{H}}(N) = N + 1 \leq N + 1$$

- $\mathcal{H}$  is positive intervals: (break point  $k = 3$ )

$$m_{\mathcal{H}}(N) = \frac{1}{2}N^2 + \frac{1}{2}N + 1 \leq \frac{1}{2}N^2 + \frac{1}{2}N + 1$$

- $\mathcal{H}$  is 2D perceptrons: (break point  $k = 4$ )

$$m_{\mathcal{H}}(N) = ? \leq \frac{1}{6}N^3 + \frac{5}{6}N + 1$$

# Outline

- Proof that  $m_{\mathcal{H}}(N)$  is polynomial
- Proof that  $m_{\mathcal{H}}(N)$  can replace  $M$

## What we want

Instead of:

$$\mathbb{P} \left[ |E_{\text{in}}(g) - E_{\text{out}}(g)| > \epsilon \right] \leq 2 \quad \textcolor{red}{M} \quad e^{-2\epsilon^2 N}$$

We want:


$$\mathbb{P} \left[ |E_{\text{in}}(g) - E_{\text{out}}(g)| > \epsilon \right] \leq 2 \quad \textcolor{red}{m}_{\mathcal{H}}(N) \quad e^{-2\epsilon^2 N}$$

## Pictorial proof ☺

- How does  $m_{\mathcal{H}}(N)$  relate to overlaps?
- What to do about  $E_{\text{out}}$ ?
- Putting it together




## Hoeffding Inequality




(a)

## Union Bound




(b)

## VC Bound



(c)

# What to do about $E_{\text{out}}$



## Putting it together

Not quite:

$$\mathbb{P} \left[ |E_{\text{in}}(g) - E_{\text{out}}(g)| > \epsilon \right] \leq 2 m_{\mathcal{H}}(N) e^{-2 \epsilon^2 N}$$

but rather:

$$\mathbb{P} \left[ |E_{\text{in}}(g) - E_{\text{out}}(g)| > \epsilon \right] \leq 4 m_{\mathcal{H}}(2N) e^{-\frac{1}{8} \epsilon^2 N}$$

## The Vapnik-Chervonenkis Inequality