ESAIM: Mathematical Modelling and Numerical Analysis

Mixed-dimensional models of poroelasticity

--Manuscript Draft--

Manuscript Number:	
Article Type:	Research article
Full Title:	Mixed-dimensional models of poroelasticity
Short Title:	
Corresponding Author:	Jan Stebel, Ph.D. Technicka Univerzita v Liberci Fakulta Mechatroniky Informatiky a Mezioborovych Studii Liberec, CZECH REPUBLIC
Corresponding Author E-Mail:	jan.stebel@tul.cz
Order of Authors:	Jan Stebel, Ph.D. Jan Březina
Manuscript Region of Origin:	CZECH REPUBLIC
Keywords:	Biot's poroelasticity; dimension reduction; fixed-stress splitting.
Manuscript Classifications:	10.10: Fluid mechanics; 10.20: Solid mechanics; 10.100: Geophysics; 20.90: Model reduction; 20.170: Variational methods
Abstract:	The paper deals with derivation and analysis of a poroelasticity model in a domain with lower-dimensional fracture. The resulting system of equations, which is obtained by dimension reduction, consists of saturated flow and linear elasticity both in the domain and in the fracture, and of appropriate interface conditions. Existence and uniqueness of a weak solution is proved with help of a fixed-point argument.
Suggested Reviewers:	Jakub Wiktor Both jakub.both@uib.no Katja Hanowski katja.hanowski@tu-dresden.de Andro Mikelic mikelic@univ-lyon1.fr
Opposed Reviewers:	
Author Comments:	
Funding Information:	European Social Fund (CZ.02.1.01/0.0/0.0/16 013/0001792) — Not applicable

Powered by Editorial Manager® and ProduXion Manager® from Aries Systems Corporation

Manuscript Click here to access/download;Manuscript;brst-m2an.pdf

Mathematical Modelling and Numerical Analysis Will be set by the publisher
Modélisation Mathématique et Analyse Numérique

MIXED-DIMENSIONAL MODELS OF POROELASTICITY *

Jan Březina [1, 2] and Jan Stebel [2]

Abstract. The paper deals with derivation and analysis of a poroelasticity model in a domain with lower-dimensional fracture. The resulting system of equations, which is obtained by dimension reduction, consists of saturated flow and linear elasticity both in the domain and in the fracture, and of appropriate interface conditions. Existence and uniqueness of a weak solution is proved with help of a fixed-point argument.

2010 Mathematics Subject Classification. 35M33, 35Q86, 74S05, 74F10.

April 1, 2020.

1. Introduction

The interaction of fluid flow and rock mechanics plays a significant role in many important applications such as geothermal power utilization, nuclear waste deposition or CO_2 storage [17]. Appropriate mathematical models must take into account heterogeneities and fault zones of various scales and shapes. An important role is played by models describing fractures as codimension-one manifolds, greatly simplifying the treatment of extremely different geometrical scales. The challenge in mathematical and numerical methods for these problems is very often related to opposite character of flow and mechanics: intact rock (matrix) is stiff but almost impermeable, while fractures are permeable and with small or negligible elastic modulus. It turns out that simply neglecting the less dominant of these two processess often leads to inaccurate description. For example, the widely-used *discrete fracture network* (DFN) models are not able to capture the mass transfer between fractures that are not topologically connected. The communication between fractures and matrix is a part of so-called *discrete fracture-matrix* (DFM) models, also called *mixed-dimensional*. In modelling hydro-geomechanical coupling, the mechanical response of fractures is often neglected. However, phenomena like shear-induced opening, friction or healing may play essential role, raising the need for appropriate mathematical description of fracture mechanics. Our aim is to develop a mixed-dimensional model taking into accout both flow and mechanics in matrix and fractures.

The original idea of dimension reduction comes from [12], where it was applied to Darcian flow and one separating fracture. Later on the model has been extended e.g. by [1] for immersed, polygonal fracture with variable aperture. DFN models with intersections and junctions were analyzed in e.g. [9, 13, 16], for results on

Keywords and phrases: Biot's poroelasticity, dimension reduction, fixed-stress splitting

* *The paper was created with support of the project No CZ.02.1.01/0.0/0.0/16_013/0001792 "RINGEN - research infrastructure upgrade", co-funded by the EU Operational Programme "Research, Development and Education".*

[1] Institute of New Technologies and Applied Informatics, Faculty of Mechatronics, Informatics and Interdisciplinary Studies, Technical University of Liberec, Studentská 1402/2, 461 17 Liberec, Czech Republic; e-mail: {jan.brezina,jan.stebel}@tul.cz

[2] Department of Nanotechnology and Informatics, Institute for Nanomaterials, Advanced Technologies and Innovation, Technical University of Liberec, Studentská 1402/2, 461 17 Liberec, Czech Republic

© EDP Sciences, SMAI 1999

DFM we refer to [18]. Mixed-dimensional poroelasticity models were studied e.g. in [10] where Biot's system in matrix was coupled with fracture flow based on the cubic law. In [11] the Biot system in matrix is coupled with Darcian flow, but in addition fracture width and length can vary according to rock deformation. Concerning mechanics in fracture, there are not so many papers on dimension reduction. A general framework for mixed-dimensional elliptic equations was developed in [4, 5] and applied to linear elasticity systems. We also mention the work [14] where a shell model is derived for poroelastic membrane.

In this paper we describe a dimension reduction process for the Biot system in a domain separated by a fracture. To cope with the fact that the mechanical part of the problem is a system of equations, we develop a special calculus for tangential and normal parts of functions and operators. In contrast to other works, we also consider general conductivity and elasticity tensors not necessarily aligned with the fracture orientation. The obtained equations are expressed in terms of mean pressure and mean displacement in the fracture. The resulting problem consists of equations of flow and mechanics in the fracture and in the surrounding domain, accompanied by appropriate interface conditions.

The second part of the paper deals with the analysis of well-posedness of the mixed-dimensional poroelasticity. There are two main strategies of solving poromechanics systems: monolithic methods (see e.g. [19, 21]) and iterative splittings. The latter approach has a great advantage in the design of numerical methods where the splitting acts as a preconditioner [20] that allows sequential use of different solvers for flow and mechanics. Here we follow the optimized fixed-stress iterative splitting proposed in [15] and show the contractivity of the appropriate mapping with optimal estimates of splitting parameters.

The structure of the paper is as follows. In section 2 we describe the full and reduced geometry and the Biot's poroelasticity model. Then in section 3 we present the tangential and normal calculus which will be used in the dimension reduction process in section 4. The result of dimension reduction is the system (4.8), which describes flow and mechanics in matrix and fracture with appropriate interface conditions. We further formulate a modified problem (4.10), which is valid under the assumption of permeable fracture. In section 5 we analyze the existence and uniqueness of solution to the latter problem, first studying separately the mechanics and flow and later on combining them in the iterative scheme.

2. Biot's poroelasticity in domain with fracture

Let Ω be a bounded, simply connected domain in the Euclidean space $\mathbf{R}^d$, $d \in \{2, 3\}$, with Lipschitz boundary $\partial\Omega$. We assume that Ω is composed of two subdomains: the fracture Ω_f and the matrix $\Omega_m := \Omega \setminus \overline{\Omega}_f$, and that the fracture divides the matrix into two parts (see Figure 1). For simplicity we consider a straight fracture, i.e. there is a subset γ of a hyperplane in $\mathbf{R}^d$ with unit normal vector ν and a number $\delta > 0$ such that

$$\Omega_f = \{\mathbf{x} + s\nu;\ \mathbf{x} \in \gamma,\ s \in (-\tfrac{\delta}{2}, \tfrac{\delta}{2})\}.$$

The parameter δ is usually called the aperture or the width of the fracture. Then the two parts of Ω_m are interacting with Ω_f via the interfaces

$$\gamma^+ := \{\mathbf{x} + \tfrac{\delta}{2}\nu;\ \mathbf{x} \in \gamma\}, \quad \gamma^- := \{\mathbf{x} - \tfrac{\delta}{2}\nu;\ \mathbf{x} \in \gamma\}.$$

The symbol $\partial\gamma$ shall denote the relative boundary of γ.

The basic hydro-mechanical interaction in a porous medium is described by the Biot system [3]. Let $I = (0, T)$ be a finite time interval, $\mathbf{f}$ the body force, g the fluid source and p_0 the initial pressure distribution. Then the balance of mass and forces in matrix and fracture is given by the equations

$$-\operatorname{div} \sigma + \nabla(\alpha p) = \mathbf{f} \qquad \text{in } I \times (\Omega_m \cup \Omega_f), \tag{2.1a}$$

$$\partial_t \left(Sp + \operatorname{div}(\alpha \mathbf{u})\right) + \operatorname{div} \mathbf{q} = g \qquad \text{in } I \times (\Omega_m \cup \Omega_f). \tag{2.1b}$$

Here, the displacement $\mathbf{u}$ and the pressure p are the principal unknowns; further α is the Biot effective stress parameter, S the storativity. The stress tensor σ and the flux $\mathbf{q}$ are determined by the Hooke and Darcy law,




FIGURE 1. The domain of the full model (left) and the reduced geometry (right).

respectively:

$$\mathfrak{o} = \mathbb{C}\nabla\mathbf{u}, \quad \mathbf{q} = -\mathbb{K}\nabla p,$$

via the 4th-order elasticity tensor $\mathbb{C}$ and the hydraulic conductivity tensor $\mathbb{K}$. We consider the following initial and boundary conditions:

$$p(0,\cdot) = p_0 \qquad \text{in } \Omega_m \cup \Omega_f, \tag{2.1c}$$

$$p = 0, \quad \mathbf{u} = \mathbf{0} \qquad \text{on } I \times \partial\Omega. \tag{2.1d}$$

To complete the system of equations, we impose the continuity of pressure, displacement, flux and normal poroelastic stress on the interfaces between matrix and fracture:

$$p, \mathbf{u}, \mathbf{q}\cdot\nu, \mathfrak{o}\nu - \alpha p\nu \text{ are continuous on } I \times \gamma^{\pm}. \tag{2.1e}$$

In what follows, we shall assume that the physical parameters $\alpha, S, \mathbb{C}, \mathbb{K}$ are constant in Ω_m, Ω_f, respectively. To distinguish values in Ω_m and Ω_f, we shall use the subscripts "m" and "f", i.e. $\alpha_m := \alpha_{|\Omega_m}$, $\alpha_f := \alpha_{|\Omega_f}$ etc. We also impose standard requirements for the data:

- $\alpha_* \geq 0$, $S_* > 0$ for $* \in \{m, f\}$;
- $\mathbb{C}_*$ and $\mathbb{K}_*$, $* \in \{m, f\}$, have the usual symmetries:

$$\forall \mathbb{A}, \mathbb{B} \in \mathbf{R}^{d\times d}: \ \mathbb{C}_*\mathbb{A} : \mathbb{B} = \mathbb{C}_*\mathbb{A}^\top : \mathbb{B} = \mathbb{C}_*\mathbb{A} : \mathbb{B}^\top = \mathbb{C}_*\mathbb{B} : \mathbb{A},$$

$$\mathbb{K}_* = \mathbb{K}_*^\top;$$

- $\mathbb{C}_*$ and $\mathbb{K}_*$, $* \in \{m, f\}$, are positive definite: There exist positive constants μ_m, μ_f, λ_m, λ_f, κ_m, κ_f, such that

$$\forall \mathbb{A} \in \mathbf{R}^{d\times d}: \ \mathbb{C}_*\mathbb{A} : \mathbb{A} \geq \mu_* \left|\mathbb{A} + \mathbb{A}^\top\right|^2 + \lambda_* |\mathbb{I} : \mathbb{A}|^2, \tag{2.2}$$

$$\forall \mathbf{v} \in \mathbf{R}^d: \ \mathbb{K}_*\mathbf{v}\cdot\mathbf{v} \geq \kappa_* |\mathbf{v}|^2, \quad * \in \{m, f\}. \tag{2.3}$$

Here ":" stands for the scalar product in $\mathbf{R}^{d\times d}$ and $|\mathbb{A}|$ denotes the Frobenius norm, i.e. $|\mathbb{A}|^2 = \mathbb{A} : \mathbb{A}$.

3. Tangential and normal calculus in fracture

In this section we develop tools that will be used later in the derivation of the mixed-dimensional Biot system. Let $\mathbb{P} := \nu \otimes \nu = [\nu_i \nu_j]_{i,j=1}^d$ be the orthogonal projector to γ. For any $\mathbf{v} \in \mathbf{R}^d$ and $\mathbb{A} \in \mathbf{R}^{d\times d}$ we introduce the orthogonal decomposition into normal and tangential direction to γ:

$$\mathbf{v} = \mathbb{P}\mathbf{v} + (\mathbb{I} - \mathbb{P})\mathbf{v} =: \mathbf{v}_\nu + \mathbf{v}_\tau, \qquad \mathbb{A} = \mathbb{A}\mathbb{P} + \mathbb{A}(\mathbb{I} - \mathbb{P}) =: \mathbb{A}_\nu + \mathbb{A}_\tau.$$

In the rest of this section, f, $\mathbf{v}$, $\mathbb{A}$ will be scalar-, vector- and tensor-valued functions defined in $\overline{\Omega}_f$. The gradient and divergence operators are decomposed into the normal and tangential part as follows:

$$\nabla f = (\nabla f)_\nu + (\nabla f)_\tau =: \nabla_\nu f + \nabla_\tau f,$$

$$\nabla \mathbf{v} = (\nabla \mathbf{v})_\nu + (\nabla \mathbf{v})_\tau =: \nabla_\nu \mathbf{v} + \nabla_\tau \mathbf{v},$$
$$\operatorname{div} \mathbf{v} = \operatorname{div} \mathbf{v}_\nu + \operatorname{div} \mathbf{v}_\tau =: \operatorname{div}_\nu \mathbf{v} + \operatorname{div}_\tau \mathbf{v},$$
$$\operatorname{div} \mathbb{A} = \operatorname{div}(\mathbb{P}\mathbb{A}) + \operatorname{div}((\mathbb{I} - \mathbb{P})\mathbb{A}) =: \operatorname{div}_\nu \mathbb{A} + \operatorname{div}_\tau \mathbb{A}.$$

We recall that the components of the Jacobian matrix and of the tensor divergence are given by $\nabla \mathbf{v} = \left[\frac{\partial v_i}{\partial x_j}\right]_{i,j=1}^d$ and $\operatorname{div} \mathbb{A} = \left[\sum_{j=1}^d \frac{\partial a_{ij}}{\partial x_j}\right]_{i=1}^d$, respectively. The symbols $f^\oplus$ and $f^\ominus$ will denote the trace of f on γ^+, γ^-, respectively, i.e.

$$f^{\oplus}(\mathbf{x}) := f(\mathbf{x} \pm \tfrac{\delta}{2}\nu), \ \mathbf{x} \in \gamma.$$

Using these symbols we can introduce the average and jump operators:

$$\{\!\{f\}\!\} := \frac{f^\oplus + f^\ominus}{2}, \quad [\![f]\!] := f^\oplus - f^\ominus.$$

The same notation will be used for vector- and tensor-valued functions. By $\overline{f}$ we shall denote the integral mean of f across the fracture aperture, i.e.

$$\overline{f}(\mathbf{x}) := \frac{1}{\delta} \int_{-\frac{\delta}{2}}^{\frac{\delta}{2}} f(\mathbf{x} + s\nu) \, \mathrm{d}s, \ \mathbf{x} \in \gamma.$$

The approximate gradient and divergence operators in the positive and negative half of fracture are defined as follows:

$$\widetilde{\nabla}^\oplus[f, \overline{f}] := \nabla_\tau \overline{f} \pm \frac{2}{\delta}(f^\oplus - \overline{f})\nu, \qquad \widetilde{\nabla}^\oplus[\mathbf{v}, \overline{\mathbf{v}}] := \nabla_\tau \overline{\mathbf{v}} \pm \frac{2}{\delta}(\mathbf{v}^\oplus - \overline{\mathbf{v}}) \otimes \nu,$$
$$\widetilde{\operatorname{div}}^\oplus[\mathbf{v}, \overline{\mathbf{v}}] := \operatorname{div}_\tau \overline{\mathbf{v}} \pm \frac{2}{\delta}(\mathbf{v}^\oplus - \overline{\mathbf{v}}) \cdot \nu, \qquad \widetilde{\operatorname{div}}^\oplus[\mathbb{A}, \overline{\mathbb{A}}] := \operatorname{div}_\tau \overline{\mathbb{A}} \pm \frac{2}{\delta}(\mathbb{A}^\oplus - \overline{\mathbb{A}})^\top \nu.$$

Remark 3.1. In the previous definitions we have used two-point differences to approximate derivatives in the normal direction. We note that it is possible to use more general relations using all three values $f^\oplus$, $\overline{f}$, $f^\ominus$ (see e.g. [12]). The two-point formulas are suitable for possible generalization at fracture intersections and junctions.

If $f \in \mathrm{C}^2(\overline{\Omega}_f)$ then from Taylor's expansion we infer that

$$f^\oplus = \overline{f} + O(\delta), \quad \nabla f^\oplus = \widetilde{\nabla}[f, \overline{f}] + O(\delta), \quad \operatorname{div} f^\oplus = \widetilde{\operatorname{div}}[f, \overline{f}] + O(\delta). \tag{3.1}$$

Here the remainder $O(\delta)$ represents an expression that can be estimated from above by $\delta C \, \|f\|_{\mathrm{C}^2(\overline{\Omega}_f)}$ and $C > 0$ is a constant independent of f and δ. Analogous properties hold for vector- and tensor-valued functions. Despite the indicated approximations, the following relations between the integral means and averaged approximates still hold exactly:

Lemma 3.2. *For all admissible functions f, $\mathbf{v}$ and $\mathbb{A}$ it holds:*

$$\overline{\nabla f} = \nabla_\tau \overline{f} + \frac{1}{\delta}\,[\![f]\!]\,\nu = \left\{\!\!\left\{ \widetilde{\nabla}[f, \overline{f}] \right\}\!\!\right\}, \tag{3.2}$$

$$\overline{\nabla \mathbf{v}} = \nabla_\tau \overline{\mathbf{v}} + \frac{1}{\delta}\,[\![\mathbf{v}]\!] \otimes \nu = \left\{\!\!\left\{ \widetilde{\nabla}[\mathbf{v}, \overline{\mathbf{v}}] \right\}\!\!\right\}, \tag{3.3}$$

$$\overline{\operatorname{div} \mathbf{v}} = \operatorname{div}_\tau \overline{\mathbf{v}} + \frac{1}{\delta}\,[\![\mathbf{v}]\!] \cdot \nu = \left\{\!\!\left\{ \widetilde{\operatorname{div}}[\mathbf{v}, \overline{\mathbf{v}}] \right\}\!\!\right\}, \tag{3.4}$$

$$\overline{\operatorname{div} \mathbb{A}} = \operatorname{div}_\tau \overline{\mathbb{A}} + \frac{1}{\delta}\,[\![\mathbb{A}^\top]\!]\,\nu = \left\{\!\!\left\{ \widetilde{\operatorname{div}}[\mathbb{A}, \overline{\mathbb{A}}] \right\}\!\!\right\}. \tag{3.5}$$

Proof. It is readily seen that $\left\{\!\!\left\{ \widetilde{\nabla} f, \overline{f} \right\}\!\!\right\} = \nabla_\tau \overline{f} + \frac{1}{\delta}\,[\![f]\!]$. Further we have:

$$\overline{\nabla_\tau f} = \frac{1}{\delta} \int_{-\frac{\delta}{2}}^{\frac{\delta}{2}} \nabla_\tau f(\cdot + s\nu)\, \mathrm{d}s = \nabla_\tau \left(\frac{1}{\delta} \int_{-\frac{\delta}{2}}^{\frac{\delta}{2}} f(\cdot + s\nu)\, \mathrm{d}s \right) = \nabla_\tau \overline{f},$$

$$\overline{\nabla_\nu f} = \frac{1}{\delta} \int_{-\frac{\delta}{2}}^{\frac{\delta}{2}} \nu \frac{\mathrm{d}}{\mathrm{d}s} \left(f(\cdot + s\nu) \right)\, \mathrm{d}s = \frac{1}{\delta}\,[\![f]\!]\,\nu,$$

so that $\overline{\nabla f} = \overline{\nabla_\tau f} + \overline{\nabla_\nu f}$ satisfies (3.2). The identities (3.3)–(3.5) can be proved analogously. □

We also have the Green identity in γ:

$$\int_\gamma (\operatorname{div}_\tau \mathbf{v}) f = \int_{\partial\gamma} (\mathbf{v} \cdot \mathbf{n}) f - \int_\gamma \mathbf{v} \cdot \nabla_\tau f,$$

where $\mathbf{n}$ denotes the unit outward normal to $\partial\gamma$.

4. Dimension reduction

Let us assume that $(\mathbf{u}, p)$ is a smooth solution of (2.1). We shall use the calculus from section 3 to derive appropriate equations that are satisfied by the averages $(\overline{\mathbf{u}}, \overline{p})$ in $I \times \gamma$.

4.1. Fracture model for elasticity

Integrating (2.1a) over the fracture aperture yields:

$$-\delta \overline{\operatorname{div} \sigma} + \delta \alpha_f \overline{\nabla p} = \delta \overline{\mathbf{f}} \text{ in } I \times \gamma.$$

To express the averaged terms on the left hand side we use (3.5), (3.3) and (3.2). In particular, we obtain:

$$\overline{\operatorname{div} \sigma} = \operatorname{div}_\tau \left(\mathbb{C}_f \left\{\!\!\left\{ \widetilde{\nabla}[\mathbf{u}, \overline{\mathbf{u}}] \right\}\!\!\right\} \right) + \frac{1}{\delta} \left[\!\!\left[\mathbb{C}_f \nabla \mathbf{u}_{|\Omega_f} \right]\!\!\right] \nu.$$

The normal stress $(\mathbb{C}_f \nabla \mathbf{u}_{|\Omega_f})^{\oplus} \nu$ will be approximated by the quantity

$$\mathbf{t}^{\oplus}[\mathbf{u}, \overline{\mathbf{u}}] := \left(\mathbb{C}_f \widetilde{\nabla}^{\oplus}[\mathbf{u}, \overline{\mathbf{u}}] \right) \nu. \tag{4.1}$$

Then, taking into account (3.1), we obtain the reduced counterpart of (2.1a):

$$-\delta \operatorname{div}_\tau \left(\mathbb{C}_f \left\{\!\!\left\{ \widetilde{\nabla}[\mathbf{u}, \overline{\mathbf{u}}] \right\}\!\!\right\} \right) + \delta \alpha_f \left\{\!\!\left\{ \widetilde{\nabla}[p, \overline{p}] \right\}\!\!\right\} - [\![\mathbf{t}[\mathbf{u}, \overline{\mathbf{u}}]]\!] = \delta \overline{\mathbf{f}} + O(\delta) \text{ in } I \times \gamma. \tag{4.2}$$

The continuity of normal poroelastic stress $\varpi\nu - \alpha p\nu$ on $\gamma^{\pm}$ becomes

$$(\mathbb{C}_m(\nabla \mathbf{u}_{|\Omega_m})^{\oplus} - \alpha_m p^{\oplus}\mathbb{I})\nu = \mathbf{t}^{\oplus}[\mathbf{u}, \overline{\mathbf{u}}] - \alpha_f p^{\oplus}\nu + O(\delta) \text{ in } I \times \gamma. \tag{4.3}$$

4.2. **Fracture model for flow**

Let us integrate (2.1b) over the fracture aperture. We get:

$$\partial_t \left(\delta S_f \overline{p} + \delta \alpha_f \overline{\operatorname{div} \mathbf{u}}\right) + \delta \overline{\operatorname{div} \mathbf{q}} = \delta \overline{g} \text{ in } I \times \gamma. \tag{4.4}$$

From (3.4), (3.2) and the definition of $\mathbf{q}$ we obtain:

$$\overline{\operatorname{div} \mathbf{q}} = \operatorname{div}_\tau \overline{\mathbf{q}} + \frac{1}{\delta}[\![\mathbf{q}]\!] \cdot \nu = -\operatorname{div}_\tau \left(\mathbb{K}_f \left\{\!\!\left\{ \widetilde{\nabla}[p, \overline{p}] \right\}\!\!\right\}\right) + \frac{1}{\delta}[\![\mathbf{q}]\!] \cdot \nu.$$

To express the approximation of the normal flux $-\mathbf{q}^{\oplus} \cdot \nu$ we use the quantity

$$\varphi^{\oplus}[p, \overline{p}] := \mathbb{K}_f \widetilde{\nabla}^{\oplus}[p, \overline{p}] \cdot \nu. \tag{4.5}$$

Hence, we obtain from (4.4)–(4.5) and (3.4) the reduced flow equation:

$$\delta \partial_t \left(S_f \overline{p} + \alpha_f \left\{\!\!\left\{ \widetilde{\operatorname{div}}[\mathbf{u}, \overline{\mathbf{u}}] \right\}\!\!\right\}\right) - \delta \operatorname{div}_\tau \left(\mathbb{K}_f \left\{\!\!\left\{ \widetilde{\nabla}[p, \overline{p}] \right\}\!\!\right\}\right) - [\![\varphi[p, \overline{p}]]\!] = \delta \overline{g} + O(\delta) \text{ in } I \times \gamma. \tag{4.6}$$

Continuity of flux through $\gamma^{\pm}$ yields:

$$\mathbb{K}_m(\nabla p_{|\Omega_m})^{\oplus} \cdot \nu = \varphi^{\oplus}[p, \overline{p}] + O(\delta) \text{ in } I \times \gamma. \tag{4.7}$$

4.3. **Mixed-dimensional poroelasticity**

The mixed-dimensional coupled problem is obtained after neglecting the remainders $O(\delta)$ in (4.2), (4.3), (4.6), (4.7). Analysis of the truncation error can be done similarly as in [7], however in this paper we omit it. Let $\Gamma := \partial\Omega_m \cap \partial\Omega$, $P_0 := \overline{p}_0$, $\mathbf{F} := \delta\overline{\mathbf{f}}$ and $G := \delta\overline{g}$. We want to find functions $\mathbf{u}, p$ defined in $I \times \Omega_m$ and $\mathbf{U}, P$ defined in $I \times \gamma$, which satisfy:

(i) Biot's equations for the matrix:

$$-\operatorname{div}(\mathbb{C}_m \nabla \mathbf{u}) + \alpha_m \nabla p = \mathbf{f} \qquad \text{in } I \times \Omega_m, \tag{4.8a}$$

$$\partial_t \left(S_m p + \alpha_m \operatorname{div} \mathbf{u}\right) - \operatorname{div}(\mathbb{K}_m \nabla p) = g \qquad \text{in } I \times \Omega_m, \tag{4.8b}$$

(ii) averaged Biot's equations for the fracture:

$$-\delta \operatorname{div}_\tau \left(\mathbb{C}_f \left\{\!\!\left\{ \widetilde{\nabla}[\mathbf{u}, \mathbf{U}] \right\}\!\!\right\}\right) + \delta \alpha_f \left\{\!\!\left\{ \widetilde{\nabla}[p, P] \right\}\!\!\right\} - [\![\mathbf{t}[\mathbf{u}, \mathbf{U}]]\!] = \mathbf{F} \text{ in } I \times \gamma, \tag{4.8c}$$

$$\delta \partial_t \left(S_f P + \alpha_f \left\{\!\!\left\{ \widetilde{\operatorname{div}}[\mathbf{u}, \mathbf{U}] \right\}\!\!\right\}\right) - \delta \operatorname{div}_\tau \left(\mathbb{K}_f \left\{\!\!\left\{ \widetilde{\nabla}[p, P] \right\}\!\!\right\}\right) - [\![\varphi[p, P]]\!] = G \text{ in } I \times \gamma, \tag{4.8d}$$

(iii) interface conditions:

$$(\mathbb{C}_m(\nabla \mathbf{u})^{\oplus} - \alpha_m p^{\oplus}\mathbb{I})\nu = \mathbf{t}^{\oplus}[\mathbf{u}, \mathbf{U}] - \alpha_f p^{\oplus}\nu \qquad \text{in } I \times \gamma, \tag{4.8e}$$

$$\mathbb{K}_m(\nabla p)^{\oplus} \cdot \nu = \varphi^{\oplus}[p, P] \qquad \text{in } I \times \gamma, \tag{4.8f}$$

(iv) boundary conditions:

$$\mathbf{u} = \mathbf{0}, \quad p = 0 \qquad \text{on } I \times \Gamma, \tag{4.8g}$$

$$\mathbf{U} = \mathbf{0}, \quad P = 0 \qquad \text{on } I \times \partial\gamma, \tag{4.8h}$$

(v) initial conditions:

$$p(0,\cdot) = p_0 \qquad \text{in } \Omega_m, \tag{4.8i}$$

$$P(0,\cdot) = P_0 \qquad \text{in } \gamma. \tag{4.8j}$$

We point out that $\mathbf{U}$ and P play the role of approximations of the averages $\overline{\mathbf{u}}$ and $\overline{p}$ in Ω_f.

At this moment we note that the problem (4.8) is not suitable for further analysis. Namely, formal calculations (see appendix B) lead to the following "energy equality":

$$\begin{aligned}\int_{\Omega_m} \mathbb{C}_m \nabla\partial_t\mathbf{u} : \nabla\partial_t\mathbf{u} + \delta \int_\gamma \left\{\!\!\left\{ \mathbb{C}_f \widetilde{\nabla}[\partial_t\mathbf{u}, \partial_t\mathbf{U}] : \widetilde{\nabla}[\partial_t\mathbf{u}, \partial_t\mathbf{U}] \right\}\!\!\right\} + S_m \int_{\Omega_m} |\partial_t p|^2 + \delta S_f \int_\gamma |\partial_t P|^2 \\ + \frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\int_{\Omega_m} \mathbb{K}_m \nabla p \cdot \nabla p + \frac{\delta}{2}\frac{\mathrm{d}}{\mathrm{d}t}\int_\gamma \left\{\!\!\left\{ \mathbb{K}_f \widetilde{\nabla}[p,P] \cdot \widetilde{\nabla}[p,P] \right\}\!\!\right\} + \int_\gamma \alpha_f \left[\!\left[\partial_t p \partial_t(\mathbf{U} - \mathbf{u}) + \partial_t P \partial_t \mathbf{u} \right]\!\right] \cdot \nu \\ = \int_{\Omega_m} (\partial_t \mathbf{f} \cdot \partial_t \mathbf{u} + \partial_t g \partial_t p) + \int_\gamma (\partial_t \mathbf{F} \cdot \partial_t \mathbf{U} + \partial_t G \partial_t P). \end{aligned} \tag{4.9}$$

If the the last term on the left hand side was not present, the identity would yield a priori bounds of the solution. However, this last term cannot be estimated by the other terms and may possibly be a source of uncontrolled "energy". Due to this fact we are unable to prove the global in time existence of weak solution without any further restriction of the model data.

4.4. **Permeable fracture**

In what follows we shall consider the case when the fracture is much more permeable compared to the matrix, in particular that the largest eigenvalue of $\mathbb{K}_m$ is much smaller than the smallest eigenvalue of $\mathbb{K}_f$:

$$\lambda_{max}(\mathbb{K}_m) \ll \lambda_{min}(\mathbb{K}_f).$$

Under such conditions the variation of pressure across the fracture width is usually very small compared to the tangential direction. In the mixed-dimensional setting it means that

$$\frac{1}{\delta} |\left[\!\left[p \right]\!\right]| \ll |\nabla_\tau P| \text{ and } P \approx p^{\oplus}.$$

Then, the last term on the left of (4.9) becomes negligible:

$$\int_\gamma \alpha_f \left[\!\left[\partial_t p \partial_t(\mathbf{U} - \mathbf{u}) + \partial_t P \partial_t \mathbf{u} \right]\!\right] \cdot \nu = \delta \int_\gamma \alpha_f \partial_t \left(\frac{1}{\delta} \left[\!\left[p \right]\!\right] \right) \partial_t \mathbf{U} \cdot \nu + \int_\gamma \alpha_f \left[\!\left[\partial_t (P - p) \partial_t \mathbf{u} \right]\!\right] \cdot \nu \approx 0.$$

Motivated by the above considerations, we replace the elasticity equation (4.8c) and the interface condition (4.8e) by the following equations:

$$-\delta \operatorname{div}_\tau \left(\mathbb{C}_f \left\{\!\!\left\{ \widetilde{\nabla}[\mathbf{u}, \mathbf{U}] \right\}\!\!\right\} \right) + \delta \alpha_f \nabla_\tau P - \left[\!\left[\mathbf{t}[\mathbf{u}, \mathbf{U}] \right]\!\right] = \mathbf{F} \text{ in } I \times \gamma, \tag{4.8c'}$$

$$(\mathbb{C}_m (\nabla \mathbf{u})^{\oplus} - \alpha_m p^{\oplus} \mathbb{I})\nu = \mathbf{t}^{\oplus}[\mathbf{u}, \mathbf{U}] - \alpha_f P \nu \text{ in } I \times \gamma. \tag{4.8e'}$$

The mixed-dimensional Biot system for permeable fracture, which will be analyzed in the next section, reads as follows:

Find a quadruplet $(\mathbf{u}, \mathbf{U}, p, P)$ which satisfies (4.8a)–(4.8b), (4.8c'), (4.8d), (4.8e'), (4.8f)–(4.8j). (4.10)

5. Well-posedness of mixed-dimensional poroelasticity

In this section we first introduce the weak formulation of (4.10). Then we solve independently the mechanical and the flow part of the problem. Finally we combine both subproblems in the fixed-stress iterative process and prove its convergence to the solution of the coupled problem.

In what follows we shall use the symbols $\mathrm{L}^2(D;Y)$, $\mathrm{H}^1(D;Y)$ for the Lebesgue and the Sobolev space of functions defined in D and taking values in Y. If $Y = \mathbf{R}$ then we shall write simply $\mathrm{L}^2(D)$, $\mathrm{H}^1(D)$. Further, $\mathrm{H}^1_0(D;Y)$ and $\mathrm{H}^1_B(D;Y)$ will be the subspaces of $\mathrm{H}^1(D;Y)$ for functions vanishing on the boundary and on a subset B of the boundary, respectively. The symbol $\mathrm{L}^2(I;X)$ will denote the Bochner space and $\mathrm{H}^1(I;X)$ its subspace of functions with time derivative in $\mathrm{L}^2(I;X)$. By $\langle y, x\rangle_X$ we shall denote the duality pairing between a Banach space X and its dual X^*, with $x \in X$ and $y \in X^*$.

For the weak formulation we introduce the spaces

$$\mathcal{V} := \mathrm{H}^1_\Gamma(\Omega_m;\mathbf{R}^d) \times \mathrm{H}^1_0(\gamma;\mathbf{R}^d), \quad \mathscr{L} := \mathrm{L}^2(\Omega_m) \times \mathrm{L}^2(\gamma), \quad \mathcal{V} := \mathrm{H}^1_\Gamma(\Omega_m) \times \mathrm{H}^1_0(\gamma),$$

equipped by the norms

$$\begin{aligned}
\|(\mathbf{v},\mathbf{V})\|_{\mathcal{V}} &:= (\|\nabla \mathbf{v}\|^2_{\mathrm{L}^2(\Omega)} + \|\nabla_\tau \mathbf{V}\|^2_{\mathrm{L}^2(\gamma)})^{1/2},\\
\|(q,Q)\|_{\mathscr{L}} &:= (\|q\|^2_{\mathrm{L}^2(\gamma)} + \|Q\|^2_{\mathrm{L}^2(\gamma)})^{1/2},\\
\|(q,Q)\|_{\mathcal{V}} &:= (\|\nabla q\|^2_{\mathrm{L}^2(\gamma)} + \|\nabla_\tau Q\|^2_{\mathrm{L}^2(\gamma)})^{1/2}.
\end{aligned}$$

Then the weak formulation of (4.10) reads (see appendix C for the derivation):

Find $(\mathbf{u},\mathbf{U}) \in \mathrm{H}^1(I;\mathcal{V})$ *and* $(p,P) \in L^\infty(I;\mathcal{V}) \cap \mathrm{H}^1(I;\mathscr{L})$ *such that*

$$(p,P)(0,\cdot) = (p_0,P_0); \tag{5.1a}$$

for all $(\mathbf{v},\mathbf{V}) \in \mathcal{V}$ *and a.a.* $t \in I$*:*

$$\int_{\Omega_m} (\mathbb{C}_m \nabla \mathbf{u} : \nabla \mathbf{v} - \alpha_m p \operatorname{div} \mathbf{v}) + \delta \int_\gamma \left\{\!\!\left\{ \mathbb{C}_f \widetilde{\nabla}[\mathbf{u},\mathbf{U}] : \widetilde{\nabla}[\mathbf{v},\mathbf{V}] - \alpha_f P \widetilde{\operatorname{div}}[\mathbf{v},\mathbf{V}] \right\}\!\!\right\} = \langle (\mathbf{f},\mathbf{F}), (\mathbf{v},\mathbf{V})\rangle_{\mathcal{V}}\,; \tag{5.1b}$$

for all $(q,Q) \in \mathcal{V}$ *and a.a.* $t \in I$*:*

$$\begin{aligned}
\int_{\Omega_m} (q\partial_t(S_m p + \alpha_m \operatorname{div} \mathbf{u}) + \mathbb{K}_m \nabla p \cdot \nabla q) + \delta \int_\gamma Q \partial_t \left(S_f P + \alpha_f \left\{\!\!\left\{ \widetilde{\operatorname{div}}[\mathbf{u},\mathbf{U}] \right\}\!\!\right\} \right) \\
+ \delta \int_\gamma \left\{\!\!\left\{ \mathbb{K}_f \widetilde{\nabla}[p,P] \cdot \widetilde{\nabla}[q,Q] \right\}\!\!\right\} = \langle (g,G), (q,Q)\rangle_{\mathscr{L}}\,.
\end{aligned} \tag{5.1c}$$

We state the main result of the paper.

Theorem 5.1. *Let* $(\mathbf{f},\mathbf{F}) \in \mathrm{H}^1(I;\mathcal{V}^*)$*,* $(g,G) \in \mathrm{L}^2(I;\mathscr{L})$ *and* $(p_0,P_0) \in \mathcal{V}$*. Then* (5.1) *has a unique solution.*

The proof will be done in the following subsections.

5.1. **Mixed-dimensional linear elasticity**

In this section we consider only the mechanical part of the problem (5.1). Assuming that p and P are given pressure fields at some time $t \in I$, we define the following forms in $\mathcal{V}$:

$$a((\mathbf{u},\mathbf{U}),(\mathbf{v},\mathbf{V})) := \int_{\Omega_m} \mathbb{C}_m \nabla\mathbf{u} : \nabla\mathbf{v} + \delta \int_\gamma \left\{\!\!\left\{ \mathbb{C}_f \widetilde{\nabla}[\mathbf{u},\mathbf{U}] : \widetilde{\nabla}[\mathbf{v},\mathbf{V}] \right\}\!\!\right\},$$

$$l((\mathbf{v},\mathbf{V})) := \langle (\mathbf{f},\mathbf{F}),(\mathbf{v},\mathbf{V})\rangle_{\mathcal{V}} + \int_{\Omega_m} \alpha_m p \operatorname{div}\mathbf{v} + \delta \int_\gamma \alpha_f P \left\{\!\!\left\{ \widetilde{\operatorname{div}}[\mathbf{v},\mathbf{V}] \right\}\!\!\right\}.$$

Then we introduce the problem:

$$\text{Find } (\mathbf{u},\mathbf{U}) \in \mathcal{V} \text{ such that for all } (\mathbf{v},\mathbf{V}) \in \mathcal{V}\text{: } a((\mathbf{u},\mathbf{U}),(\mathbf{v},\mathbf{V})) = l((\mathbf{v},\mathbf{V})). \tag{5.2}$$

The existence and uniqueness of the solution to (5.2) will be proved using the Lax-Milgram theorem. To ensure ellipticity of the bilinear form a, we first prove a suitable inequality for functions in $\mathcal{V}$.

5.1.1. *Korn type inequalities*

Let us recall the classical Korn inequality: For any bounded domain $D \subset \mathbf{R}^d$ with Lipschitz boundary and a non-trivial part $B \subset \partial D$ there is a constant $C_K := C_K(D,B) > 0$ such that

$$\forall \mathbf{v} \in \mathrm{H}^1_B(D;\mathbf{R}^d):\ C_K \left\|\nabla\mathbf{v}\right\|_{\mathrm{L}^2(D)} \le \left\|\varepsilon(\mathbf{v})\right\|_{\mathrm{L}^2(D)}, \tag{5.3}$$

where $\varepsilon(\mathbf{v}) := \frac{1}{2}(\nabla\mathbf{v} + (\nabla\mathbf{v})^\top)$. For any function $\mathbf{V} \in \mathrm{H}^1(\gamma;\mathbf{R}^d)$ we denote its symmetrized tangential gradient by

$$\varepsilon_\tau(\mathbf{V}) := \frac{1}{2}(\nabla_\tau\mathbf{V} + (\nabla_\tau\mathbf{V})^\top).$$

Similarly for a pair of functions $(\mathbf{v},\mathbf{V}) \in \mathcal{V}$ we define its symmetrized approximate gradient by

$$\widetilde{\varepsilon}^{\oplus}[\mathbf{v},\mathbf{V}] := \frac{1}{2}(\widetilde{\nabla}^{\oplus}[\mathbf{v},\mathbf{V}] + (\widetilde{\nabla}^{\oplus}[\mathbf{v},\mathbf{V}])^\top).$$

We have the following Korn inequality in $\mathrm{H}^1_0(\gamma)$.

Lemma 5.2. *There exists a constant $C_1 > 0$ such that for all $\mathbf{V} \in \mathrm{H}^1_0(\gamma;\mathbf{R}^d)$ it holds:*

$$C_1 \left\|\nabla_\tau\mathbf{V}\right\|^2_{\mathrm{L}^2(\gamma)} \le \left\|\varepsilon_\tau(\mathbf{V})\right\|^2_{\mathrm{L}^2(\gamma)}. \tag{5.4}$$

Proof. Let $\mathbf{V} \in \mathrm{H}^1_0(\gamma;\mathbf{R}^d)$. From the definition of the tangential operator it follows that

$$\left\|\varepsilon_\tau(\mathbf{V}_\nu)\right\|^2_{\mathrm{L}^2(\gamma)} = \tfrac{1}{2}\left\|\nabla_\tau\mathbf{V}_\nu\right\|^2_{\mathrm{L}^2(\gamma)},$$

which implies

$$\left\|\varepsilon_\tau(\mathbf{V})\right\|^2_{\mathrm{L}^2(\gamma)} = \left\|\varepsilon_\tau(\mathbf{V}_\tau)\right\|^2_{\mathrm{L}^2(\gamma)} + \tfrac{1}{2}\left\|\nabla_\tau\mathbf{V}_\nu\right\|^2_{\mathrm{L}^2(\gamma)}.$$

Further, by mapping $\mathbf{V}$ isometrically from γ to a subset of $\mathbf{R}^{d-1}$, we can apply (5.3) to the tangential part $\mathbf{V}_\tau$, which implies:

$$\left\|\varepsilon_\tau(\mathbf{V}_\tau)\right\|^2_{\mathrm{L}^2(\gamma)} \ge C_K \left\|\nabla_\tau\mathbf{V}_\tau\right\|^2_{\mathrm{L}^2(\gamma)}.$$

Altogether we have:

$$\left\|\varepsilon_\tau(\mathbf{V})\right\|^2_{\mathrm{L}^2(\gamma)} \ge \min\{C_K, \tfrac{1}{2}\}\left\|\nabla_\tau\mathbf{V}\right\|^2_{\mathrm{L}^2(\gamma)},$$

which proves (5.4). □

Now we can prove a variant of the Korn inequality in $\mathcal{V}$, which takes into account the approximate gradient.

Lemma 5.3. *There exists a constant $C_2 > 0$ such that for all $(\mathbf{v}, \mathbf{V}) \in \mathcal{V}$ it holds:*

$$C_2 \|(\mathbf{v}, \mathbf{V})\|^2_{\mathcal{V}} \leq \|\varepsilon(\mathbf{v})\|^2_{\mathrm{L}^2(\Omega_m)} + \left\{\!\!\left\{ \|\tilde{\varepsilon}[\mathbf{v}, \mathbf{V}]\|^2_{\mathrm{L}^2(\gamma)} \right\}\!\!\right\}. \tag{5.5}$$

Proof. Let us assume for contradiction that there is a sequence $\{(\mathbf{v}_n, \mathbf{V}_n)\}_{n=1}^{\infty}$ in $\mathcal{V}$ such that

$$\|\varepsilon(\mathbf{v}_n)\|^2_{\mathrm{L}^2(\Omega_m)} + \left\{\!\!\left\{ \|\tilde{\varepsilon}[\mathbf{v}_n, \mathbf{V}_n]\|^2_{\mathrm{L}^2(\gamma)} \right\}\!\!\right\} < \frac{1}{n} \|(\mathbf{v}_n, \mathbf{V}_n)\|^2_{\mathcal{V}}. \tag{5.6}$$

Without loss of generality we may assume that $\|(\mathbf{v}_n, \mathbf{V}_n)\|_{\mathcal{V}} = 1$, so that

$$(\mathbf{v}_n, \mathbf{V}_n) \rightharpoonup (\mathbf{v}, \mathbf{V}) \text{ weakly in } \mathcal{V},\ n \to \infty, \tag{5.7}$$

where $(\mathbf{v}, \mathbf{V}) \in \mathcal{V}$, passing eventually to a subsequence. From (5.6) and the Korn inequality in $\mathrm{H}^1_\Gamma(\Omega_m; \mathbf{R}^d)$ it follows that

$$\mathbf{v}_n \to \mathbf{0} \text{ strongly in } \mathrm{H}^1_\Gamma(\Omega_m; \mathbf{R}^d),\ n \to \infty, \tag{5.8}$$

implying that $\mathbf{v} = \mathbf{0}$. From (5.6) it also follows that

$$\tilde{\varepsilon}^{\oplus}[\mathbf{v}_n, \mathbf{V}_n] \to 0 \text{ strongly in } \mathrm{L}^2(\gamma; \mathbf{R}^{d\times d}),\ n \to \infty. \tag{5.9}$$

Using this together with the definition of $\tilde{\varepsilon}^{\oplus}$, (5.7), (5.8) and the embedding $\mathrm{H}^1(\Omega_m) \hookrightarrow \mathrm{L}^2(\gamma)$, we obtain:

$$\varepsilon_\tau(\mathbf{V}) = \pm\frac{1}{\delta}(\mathbf{V} \otimes \nu + \nu \otimes \mathbf{V}),$$

from which it follows that $\mathbf{V} \otimes \nu + \nu \otimes \mathbf{V} = \mathbb{0}$ and thus $\mathbf{V} = \mathbf{0}$.

With help of the compact embedding $\mathrm{H}^1(\gamma) \hookrightarrow \mathrm{L}^2(\gamma)$ we deduce from (5.9) that $\varepsilon_\tau(\mathbf{V}_n) \to 0$ strongly in $\mathrm{L}^2(\gamma; \mathbf{R}^{d\times d})$, $n \to \infty$. Lemma 5.2 then implies that also $\mathbf{V}_n \to \mathbf{0}$ strongly in $\mathrm{H}^1_0(\gamma; \mathbf{R}^d)$, $n \to \infty$. Finally, having shown the strong convergence of $\{(\mathbf{v}_n, \mathbf{V}_n)\}_{n=1}^{\infty}$, we observe that

$$1 = \lim_{n\to\infty} \|(\mathbf{v}_n, \mathbf{V}_n)\|_{\mathcal{V}} = \|(\mathbf{v}, \mathbf{V})\|_{\mathcal{V}} = 0,$$

which is a contradiction. Therefore the proof is finished. □

5.1.2. *Existence and uniqueness of solutions*

In the following lemma we prove essential properties of a and l.

Lemma 5.4. *The bilinear form a is bounded and elliptic in $\mathcal{V}$ and the linear form l is bounded in $\mathcal{V}$. In particular, there exist constants $C_3 = C_3(C_2, \delta, \mu_m, \mu_f) > 0$ and $C_4 = C_4(\alpha_m, \alpha_f, \delta, \gamma^+, \gamma^-, \Omega_m) > 0$ such that*

$$a((\mathbf{v}, \mathbf{V}), (\mathbf{v}, \mathbf{V})) \geq C_3 \|(\mathbf{v}, \mathbf{V})\|^2_{\mathcal{V}},$$

$$l((\mathbf{v}, \mathbf{V})) \leq (\|(\mathbf{f}, \mathbf{F})\|_{\mathcal{V}^*} + C_4 \|(p, P)\|_{\mathscr{L}}) \|(\mathbf{v}, \mathbf{V})\|_{\mathcal{V}}$$

for all $(\mathbf{v}, \mathbf{V}) \in \mathcal{V}$.

Proof. The boundedness of a and l is a consequence of the embeddings $\mathrm{H}^1(\Omega_m) \hookrightarrow \mathrm{L}^2(\gamma^\pm)$ and standard inequalities; we prove only the ellipticity. For any $(\mathbf{v}, \mathbf{V}) \in \mathcal{V}$ we have:

$$a((\mathbf{v}, \mathbf{V}), (\mathbf{v}, \mathbf{V})) = \int_{\Omega_m} \mathbb{C}_m \nabla\mathbf{v} : \nabla\mathbf{v} + \delta \int_\gamma \left\{\!\!\left\{ \mathbb{C}_f \widetilde{\nabla}[\mathbf{v}, \mathbf{V}] : \widetilde{\nabla}[\mathbf{v}, \mathbf{V}] \right\}\!\!\right\}.$$

Now we can apply (2.2) and (5.5), which yields:

$$a((\mathbf{v},\mathbf{V}),(\mathbf{v},\mathbf{V})) \geq 2\mu_m \left\|\varepsilon(\mathbf{v})\right\|^2_{\mathrm{L}^2(\Omega_m)} + 2\mu_f\delta \left\{\!\!\left\{ \left\|\tilde{\varepsilon}[\mathbf{v},\mathbf{V}]\right\|^2_{\mathrm{L}^2(\gamma)} \right\}\!\!\right\} \geq 2C_2 \min\{\mu_m,\mu_f\delta\} \left\|(\mathbf{v},\mathbf{V})\right\|^2_{\mathcal{V}}.$$

From this we see that a is elliptic with the constant $C_3 = 2C_2\min\{\mu_m,\mu_f\delta\}$. □

Having shown the above properties of the forms a and l, the following result is a direct consequence of the Lax-Milgram theorem.

Theorem 5.5. *Let $(\mathbf{f},\mathbf{F}) \in \mathcal{V}^*$ and $(p,P) \in \mathscr{L}$. Then problem (5.2) has a unique solution $(\mathbf{u},\mathbf{U}) \in \mathcal{V}$ which satisfies:*

$$\left\|(\mathbf{u},\mathbf{U})\right\|_{\mathcal{V}} \leq \frac{1}{C_3}\Big(\left\|(\mathbf{f},\mathbf{F})\right\|_{\mathcal{V}^*} + C_4 \left\|(p,P)\right\|_{\mathscr{L}} \Big).$$

Now we consider the time-dependent case. If $(\mathbf{f},\mathbf{F},p,P)$ are defined in the time interval I, then by differentiating the identity in (5.2) with respect to time we observe that the pair $(\partial_t\mathbf{u},\partial_t\mathbf{U})$ satisfies for all $(\mathbf{v},\mathbf{V}) \in \mathcal{V}$:

$$a((\partial_t\mathbf{u},\partial_t\mathbf{U}),(\mathbf{v},\mathbf{V})) = \partial_t l((\mathbf{v},\mathbf{V})), \tag{5.10}$$

where $\partial_t l$ stands for the linear form

$$\partial_t l((\mathbf{v},\mathbf{V})) := \langle(\partial_t\mathbf{f},\partial_t\mathbf{F}),(\mathbf{v},\mathbf{V})\rangle_{\mathcal{V}} + \int_{\Omega_m} \alpha_m \partial_t p \operatorname{div}\mathbf{v} + \delta\int_\gamma \alpha_f \partial_t P \left\{\!\!\left\{ \widetilde{\operatorname{div}}[\mathbf{v},\mathbf{V}] \right\}\!\!\right\}.$$

Similarly as in Lemma 5.4, one can show that $\partial_t l$ is bounded and estimate it in terms of the time derivatives of the data:

$$\partial_t l((\mathbf{v},\mathbf{V})) \leq \left(\left\|(\partial_t\mathbf{f},\partial_t\mathbf{F})\right\|_{\mathcal{V}^*} + C_4\left\|(\partial_t p,\partial_t P)\right\|_{\mathscr{L}}\right)\left\|(\mathbf{v},\mathbf{V})\right\|_{\mathcal{V}}.$$

Since the solution mappings $(\mathbf{f},\mathbf{F},p,P) \mapsto (\mathbf{u},\mathbf{U})$ and $(\partial_t\mathbf{f},\partial_t\mathbf{F},\partial_t p,\partial_t P) \mapsto (\partial_t\mathbf{u},\partial_t\mathbf{U})$ are linear and bounded, we obtain:

Corollary 5.6. *Let $(\mathbf{f},\mathbf{F}) \in \mathrm{L}^2(I;\mathcal{V}^*)$ and $(p,P) \in \mathrm{L}^2(I;\mathscr{L})$. Denoting by $(\mathbf{u}[t],\mathbf{U}[t])$ the solution to (5.2) with the data $\mathbf{f}[t],\mathbf{F}[t],p[t],P[t]$, then $(\mathbf{u},\mathbf{U}) \in \mathrm{L}^2(I;\mathcal{V})$. If in addition $\partial_t(\mathbf{f},\mathbf{F}) \in \mathrm{L}^2(I;\mathcal{V}^*)$ and $\partial_t(p,P) \in \mathrm{L}^2(I;\mathscr{L})$, then also $\partial_t(\mathbf{u},\mathbf{U}) \in \mathrm{L}^2(I;\mathcal{V})$.*

Based on the above result, we define the mapping $\mathcal{E} : \mathrm{H}^1(I;\mathscr{L}) \to \mathrm{H}^1(I;\mathcal{V})$ by the relation

$$\mathcal{E}(p,P)[t] := (\mathbf{u}[t],\mathbf{U}[t]).$$

We shall use it in the definition of the fixed-stress splitting in section 5.3.

5.2. **Mixed-dimensional flow**

We now turn to the flow part of the mixed-dimensional problem (5.1), assuming zero displacements. We want to solve the following problem:

Find $(p,P) \in L^\infty(I;\mathcal{V}) \cap \mathrm{H}^1(I;\mathscr{L})$ such that

$$(p,P)(0,\cdot) = (p_0,P_0); \tag{5.11a}$$

and for all $(q,Q) \in \mathcal{V}$ and a.a. $t \in I$:

$$\int_{\Omega_m} (S_m q\partial_t p + \mathbb{K}_m\nabla p\cdot\nabla q) + \delta S_f\int_\gamma Q\partial_t P + \delta\int_\gamma \left\{\!\!\left\{ \mathbb{K}_f\widetilde{\nabla}[p,P]\cdot\widetilde{\nabla}[q,Q] \right\}\!\!\right\} = \Big\langle(\tilde{g},\tilde{G}),(q,Q)\Big\rangle_{\mathscr{L}}. \tag{5.11b}$$

Here $(\tilde{g},\tilde{G})$ are modified fluid sources to be specified in section 5.3. The well-posedness of (5.11) will be established on the base of standard theory of parabolic problems (see e.g. [8], chap. 7.1). We formulate the result in the following theorem.

Theorem 5.7. *Let $(p_0, P_0) \in \mathcal{V}$ and $(\tilde{g}, \tilde{G}) \in \mathrm{L}^2(I; \mathscr{L})$. Then there exists a unique solution to* (5.11).

Proof. Existence: Let $\{w_j\}_{j=1}^{\infty}$ and $\{W_j\}_{j=1}^{\infty}$ be bases in $\mathrm{H}^1_\Gamma(\Omega_m)$, $\mathrm{H}^1_0(\gamma)$, respectively, orthonormal with respect to the L^2-scalar products. For any $n = 1, 2, ...$ we define the Galerkin approximation (p_n, P_n) as a pair of functions in the form $p_n(t, \mathbf{x}) := \sum_{j=1}^n \psi_j^n(t) w_j(\mathbf{x})$, $P_n(t, \mathbf{x}) := \sum_{j=1}^n \Psi_j^n(t) W_j(\mathbf{x})$, satisfying for a.a. $t \in I$ and for $i = 1, ..., n$ the integral identities

$$S_m \int_{\Omega_m} w_i \partial_t p_n + \int_{\Omega_m} \mathbb{K}_m \nabla p_n \cdot \nabla w_i + \delta \int_\gamma \left\{\!\!\left\{ \mathbb{K}_f \widetilde{\nabla}[p_n, P_n] \cdot \widetilde{\nabla}[w_i, 0] \right\}\!\!\right\} + \int_\gamma [\![\varphi[p_n, P_n] w_i]\!] = \int_{\Omega_m} \tilde{g} w_i, \quad (5.12)$$

$$\delta S_f \int_\gamma W_i \partial_t P_n + \delta \int_\gamma \mathbb{K}_f \left\{\!\!\left\{ \widetilde{\nabla}[p_n, P_n] \right\}\!\!\right\} \cdot \nabla_\tau W_i - \int_\gamma [\![\varphi[p_n, P_n] W_i]\!] = \int_\gamma \tilde{G} W_i, \quad (5.13)$$

and the initial conditions

$$\psi_i(0) = \int_{\Omega_m} p_0 w_i, \quad \Psi_i(0) = \int_\gamma P_0 W_i.$$

The existence and uniqueness of (p_n, P_n) follows from the Carathéodory theory of ODEs. Multiplying (5.12) by ψ_i' and (5.13) by Ψ_i', summing over $i = 1, ..., n$ and integrating in time, we arrive at the identity

$$\left\| \partial_t (\sqrt{S_m} p_n, \sqrt{\delta S_f} P_n) \right\|_{\mathscr{L}}^2 + \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \left(\int_{\Omega_m} \mathbb{K}_m \nabla p_n \cdot \nabla p_n + \delta \int_\gamma \left\{\!\!\left\{ \mathbb{K}_f \widetilde{\nabla}[p_n, P_n] \cdot \widetilde{\nabla}[p_n, P_n] \right\}\!\!\right\} \right)$$
$$= \left\langle (\tilde{g}, \tilde{G}), \partial_t (p_n, P_n) \right\rangle_{\mathcal{V}}. \quad (5.14)$$

From the positive definiteness of $\mathbb{K}_f$ and the definition of $\widetilde{\nabla}^{\oplus}$ we infer that

$$\left\{\!\!\left\{ \mathbb{K}_f \widetilde{\nabla}[p_n, P_n] \cdot \widetilde{\nabla}[p_n, P_n] \right\}\!\!\right\} \geq \kappa_f \left(|\nabla_\tau P_n|^2 + \frac{2}{\delta} |p_n^{\oplus} - P_n|^2 \right). \quad (5.15)$$

Hence, after integrating (5.14) in time, using the positive definiteness of $\mathbb{K}_m$, (5.15), the initial conditions and a suitable estimate of the right hand side, we obtain:

$$\int_0^T \|\partial_t (p_n, P_n)\|_{\mathscr{L}}^2 + \sup_{t \in I} \|(p_n, P_n)\|_{\mathcal{V}}^2 (t) \leq C \left(\|(p_0, P_0)\|_{\mathcal{V}}^2 + \int_0^T \left\| (\tilde{g}, \tilde{G}) \right\|_{\mathscr{L}}^2 \right), \quad (5.16)$$

where $C > 0$ may depend on $S_m, S_f, \delta, \kappa_m, \kappa_f, |\mathbb{K}_m|, |\mathbb{K}_f|$.

Hence the sequence $\{(p_n, P_n)\}_{n=1}^{\infty}$ is bounded in $L^\infty(I; \mathcal{V}) \cap \mathrm{H}^1(I; \mathscr{L})$. Choosing a subsequence weakly convergent in the above spaces to a pair (p, P), we can pass to the limit in (5.12)–(5.13), which proves that (p, P) is a solution of (5.11).

Uniqueness: Let (p^1, P^1) and (p^2, P^2) be two solutions of (5.11) and denote their difference by $(\Delta p, \Delta P)$. Subtracting the integral identities in (5.11b) for the two solutions and testing by $(\Delta p, \Delta P)$ we obtain:

$$\frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \left\| (\sqrt{S_m} \Delta p, \sqrt{\delta S_f} \Delta P) \right\|_{\mathscr{L}}^2 + \int_{\Omega_m} \mathbb{K}_m \nabla \Delta p \cdot \nabla \Delta p + \delta \int_\gamma \left\{\!\!\left\{ \mathbb{K}_f \widetilde{\nabla}[\Delta p, \Delta P] \cdot \widetilde{\nabla}[\Delta p, \Delta P] \right\}\!\!\right\} = 0. \quad (5.17)$$

Since all but the first term on the left hand side of (5.17) are nonnegative, we obtain the inequality

$$\frac{\mathrm{d}}{\mathrm{d}t} \|(\Delta p, \Delta P)\|_{\mathscr{L}}^2 \leq 0,$$

which together with the fact that $(\Delta p, \Delta P)(0, \cdot) = (0, 0)$ implies $(p^1, P^1) = (p^2, P^2)$. □

Having proved the above result, we introduce the solution map $\mathcal{F} : \mathrm{L}^2(I;\mathscr{L}) \to L^\infty(I;\mathcal{V}) \cap \mathrm{H}^1(I;\mathscr{L})$ by the relation

$$\mathcal{F}(\tilde{g},\tilde{G}) := (p,P),$$

where (p,P) is the solution of (5.11). Let $\boldsymbol{\beta} := (\beta_m,\beta_f)$ be a pair of positive parameters. We define a similar map $\mathcal{F}_{\boldsymbol{\beta}}$ as the solution operator for the flow problem with increased storativity $S_m+\beta_m$, $S_f+\beta_f$, respectively. In particular, $(p,P) := \mathcal{F}_{\boldsymbol{\beta}}(\tilde{g},\tilde{G})$ satisfies the initial conditions (5.11a) and the integral identity

$$(S_m+\beta_m)\int_{\Omega_m} q\partial_t p + \int_{\Omega_m} \mathbb{K}_m\nabla p\cdot\nabla q + \delta(S_f+\beta_f)\int_\gamma Q\partial_t P + \delta\int_\gamma \left\{\!\!\left\{ \mathbb{K}_f\widetilde{\nabla}[p,P]\cdot\widetilde{\nabla}[q,Q]\right\}\!\!\right\} = \Big\langle (\tilde{g},\tilde{G}),(q,Q)\Big\rangle_{\mathscr{L}} \quad (5.18)$$

for all $(q,Q)\in\mathcal{V}$ and a.a. $t\in I$.

5.3. **Fixed-stress iterations**

We follow the idea of optimized fixed-stress splitting from [15]. Let $\mathcal{X} := L^\infty(I;\mathcal{V})\cap \mathrm{H}^1(I;\mathscr{L})$. For any $n\in\{0,1,2,...\}$ and $(p^n,P^n)\in\mathcal{X}$ we define

$$(\mathbf{u}^n,\mathbf{U}^n) := \mathcal{E}(p^n,P^n).$$

Next we choose a pair of positive parameters $\boldsymbol{\beta} := (\beta_m,\beta_f)$ and define

$$(p^{n+1},P^{n+1}) := \mathcal{F}_{\boldsymbol{\beta}}(\tilde{g}^n,\tilde{G}^n),$$

where

$$\tilde{g}^n := g - \alpha_m \operatorname{div}\partial_t\mathbf{u}^n + \beta_m\partial_t p^n,$$
$$\tilde{G}^n := G - \delta\alpha_f\left\{\!\!\left\{\widetilde{\operatorname{div}}[\partial_t\mathbf{u}^n,\partial_t\mathbf{U}^n]\right\}\!\!\right\} + \delta\beta_f\partial_t P^n.$$

From the results of sections 5.1 and 5.2 we infer that whenever $(g,G)\in\mathrm{L}^2(I;\mathscr{L})$ then also $(\tilde{g}^n,\tilde{G}^n)\in\mathrm{L}^2(I;\mathscr{L})$. The above scheme defines the mapping

$$\mathcal{M}_{\boldsymbol{\beta}} : (p^n,P^n)\mapsto(p^{n+1},P^{n+1})$$

from $\mathcal{X}$ into itself. If $(\hat{p},\hat{P})$ is a fixed point of this map and $(\hat{\mathbf{u}},\hat{\mathbf{U}}) := \mathcal{E}(\hat{p},\hat{P})$, then $(\hat{\mathbf{u}},\hat{\mathbf{U}},\hat{p},\hat{P})$ is a solution to (5.1). Let

$$\omega := \min\left\{\frac{S_m+\beta_m}{\beta_m},\frac{S_f+\beta_f}{\beta_f}\right\}.$$

Then the following expression $\rho_{\boldsymbol{\beta}}$ is a distance in $\mathcal{X}$:

$$\begin{aligned}\rho_{\boldsymbol{\beta}}((p_1,P_1),(p_2,P_2))^2 := {}& \frac{\omega}{\beta_m}\int_0^T \left\|\partial_t\left(\alpha_m\operatorname{div}(\mathbf{u}_1-\mathbf{u}_2)-\beta_m(p_1-p_2)\right)\right\|^2_{\mathrm{L}^2(\Omega_m)}\\ &+\frac{\delta\omega}{\beta_f}\int_0^T\left\{\!\!\left\{\left\|\partial_t\left(\alpha_f\widetilde{\operatorname{div}}[\mathbf{u}_1-\mathbf{u}_2,\mathbf{U}_1-\mathbf{U}_2]-\beta_f(P_1-P_2)\right)\right\|^2_{\mathrm{L}^2(\gamma)}\right\}\!\!\right\}\\ &+\sup_{t\in I}\left(\kappa_m\left\|\nabla(p_1-p_2)\right\|^2_{\mathrm{L}^2(\Omega_m)}+\delta\kappa_f\left\{\!\!\left\{\left\|\widetilde{\nabla}[p_1-p_2,P_1-P_2]\right\|^2_{\mathrm{L}^2(\gamma)}\right\}\!\!\right\}\right),\end{aligned}$$

where $(\mathbf{u}_i,\mathbf{U}_i) := \mathcal{E}(p_i,P_i)$, $i=1,2$. Further, $(\mathcal{X},\rho_{\boldsymbol{\beta}})$ is a complete metric space. We shall prove that $\mathcal{M}_{\boldsymbol{\beta}}$ is a contraction with respect to $\rho_{\boldsymbol{\beta}}$. Then $\mathcal{M}_{\boldsymbol{\beta}}$ has a unique fixed point and (5.1) has a unique solution, thus Theorem 5.1 will be proved.

Theorem 5.8. *Let* $(\mathbf{f},\mathbf{F})\in \mathrm{H}^1(I;\mathcal{V}^*)$, $(g,G)\in \mathrm{L}^2(I;\mathscr{L})$ *and* $(p_0,P_0)\in\mathcal{V}$. *If the pair* $\boldsymbol{\beta}=(\beta_m,\beta_f)$ *satisfies*

$$\beta_m \geq \frac{\alpha_m^2}{2\left(\frac{2\mu_m}{d}+\lambda_m\right)} \quad \textit{and} \quad \beta_f \geq \frac{\alpha_f^2}{2\left(\frac{2\mu_f}{d}+\lambda_f\right)}, \tag{5.19}$$

then $\mathcal{M}_{\boldsymbol{\beta}}$ *is a contraction in* $\mathcal{X}$ *with respect to the distance* $\rho_{\boldsymbol{\beta}}$. *Hence the sequence* $\{(p^n,P^n)\}_{n=0}^{\infty}$ *with an arbitrary initial guess* $(p^0,P^0)\in\mathcal{X}$, *has a unique fixed point* $(\hat{p},\hat{P})$. *Denoting* $(\hat{\mathbf{u}},\hat{\mathbf{U}}):=\mathcal{E}(\hat{p},\hat{P})$, *then the quadruplet* $(\hat{\mathbf{u}},\hat{\mathbf{U}},\hat{p},\hat{P})$ *is the unique solution of the coupled problem* (5.1). *The contraction constant takes the value* $1/\omega$, *which becomes smallest when equalities hold in* (5.19).

Proof. For $n=1,2,...$ we denote:

$$\begin{aligned}&\Delta_p^n:=p^n-p^{n-1},\quad e_p^n:=\partial_t\Delta_p^n,\quad \Delta_P^n:=P^n-P^{n-1},\quad e_P^n:=\partial_t\Delta_P^n,\\ &\mathbf{e}_{\mathbf{u}}^n:=\partial_t(\mathbf{u}^n-\mathbf{u}^{n-1}),\quad \mathbf{e}_{\mathbf{U}}^n:=\partial_t(\mathbf{U}^n-\mathbf{U}^{n-1}).\end{aligned}$$

We subtract the equations (5.10) for $(\partial_t\mathbf{u}^{n+1},\partial_t\mathbf{U}^{n+1})$ and $(\partial_t\mathbf{u}^n,\partial_t\mathbf{U}^n)$ and test by $(\mathbf{e}_{\mathbf{u}}^{n+1},\mathbf{e}_{\mathbf{U}}^{n+1})$. We obtain:

$$\begin{aligned}\int_{\Omega_m}\mathbb{C}_m\nabla\mathbf{e}_{\mathbf{u}}^{n+1}:\nabla\mathbf{e}_{\mathbf{u}}^{n+1}+\delta\int_\gamma\left\{\!\!\left\{\mathbb{C}_f\widetilde{\nabla}[\mathbf{e}_{\mathbf{u}}^{n+1},\mathbf{e}_{\mathbf{U}}^{n+1}]:\widetilde{\nabla}[\mathbf{e}_{\mathbf{u}}^{n+1},\mathbf{e}_{\mathbf{U}}^{n+1}]\right\}\!\!\right\}\\ =\int_{\Omega_m}e_p^{n+1}\alpha_m\operatorname{div}\mathbf{e}_{\mathbf{u}}^{n+1}+\delta\int_\gamma e_P^{n+1}\left\{\!\!\left\{\alpha_f\widetilde{\operatorname{div}}[\mathbf{e}_{\mathbf{u}}^{n+1},\mathbf{e}_{\mathbf{U}}^{n+1}]\right\}\!\!\right\}.\end{aligned}$$

Using (2.2) and the inequalities (A.1)–(A.2) from appendix A we obtain:

$$\begin{aligned}\left(\frac{2\mu_m}{d}+\lambda_m\right)\left\|\operatorname{div}\mathbf{e}_{\mathbf{u}}^{n+1}\right\|^2_{\mathrm{L}^2(\Omega_m)}+\delta\left(\frac{2\mu_f}{d}+\lambda_f\right)\left\{\!\!\left\{\left\|\widetilde{\operatorname{div}}[\mathbf{e}_{\mathbf{u}}^{n+1},\mathbf{e}_{\mathbf{U}}^{n+1}]\right\|^2_{\mathrm{L}^2(\gamma)}\right\}\!\!\right\}\\ \leq\int_{\Omega_m}e_p^{n+1}\alpha_m\operatorname{div}\mathbf{e}_{\mathbf{u}}^{n+1}+\delta\int_\gamma e_P^{n+1}\left\{\!\!\left\{\alpha_f\widetilde{\operatorname{div}}[\mathbf{e}_{\mathbf{u}}^{n+1},\mathbf{e}_{\mathbf{U}}^{n+1}]\right\}\!\!\right\},\end{aligned}$$

which together with (5.19) yields:

$$\begin{aligned}\frac{1}{2\beta_m}\left\|\alpha_m\operatorname{div}\mathbf{e}_{\mathbf{u}}^{n+1}\right\|^2_{\mathrm{L}^2(\Omega_m)}+\frac{\delta}{2\beta_f}\left\{\!\!\left\{\left\|\alpha_f\widetilde{\operatorname{div}}[\mathbf{e}_{\mathbf{u}}^{n+1},\mathbf{e}_{\mathbf{U}}^{n+1}]\right\|^2_{\mathrm{L}^2(\gamma)}\right\}\!\!\right\}\\ -\frac{1}{\beta_m}\int_{\Omega_m}\beta_m e_p^{n+1}\alpha_m\operatorname{div}\mathbf{e}_{\mathbf{u}}^{n+1}-\frac{\delta}{\beta_f}\int_\gamma\beta_f e_P^{n+1}\alpha_f\left\{\!\!\left\{\widetilde{\operatorname{div}}[\mathbf{e}_{\mathbf{u}}^{n+1},\mathbf{e}_{\mathbf{U}}^{n+1}]\right\}\!\!\right\}\leq 0.\end{aligned}\tag{5.20}$$

Next we subtract the equations (5.18) for (p^{n+1},P^{n+1}) and (p^n,P^n) with appropriate right hand sides $(\tilde{g}^n,\tilde{G}^n)$ and $(\tilde{g}^{n-1},\tilde{G}^{n-1})$, respectively, test by (e_p^{n+1},e_P^{n+1}), integrate in time and use (2.3), which yields for a.a. $t\in I$:

$$\begin{aligned}(S_m+\beta_m)\int_0^t\left\|e_p^{n+1}\right\|^2_{\mathrm{L}^2(\Omega_m)}+\delta(S_f+\beta_f)\int_0^t\left\|e_P^{n+1}\right\|^2_{\mathrm{L}^2(\gamma)}\\ +\frac{\kappa_m}{2}\left\|\nabla\Delta_p^{n+1}\right\|^2_{\mathrm{L}^2(\Omega_m)}(t)+\frac{\delta\kappa_f}{2}\left\{\!\!\left\{\left\|\widetilde{\nabla}[\Delta_p^{n+1},\Delta_P^{n+1}]\right\|^2_{\mathrm{L}^2(\gamma)}(t)\right\}\!\!\right\}\\ \leq-\int_0^t\int_{\Omega_m}\underbrace{\left(\alpha_m\operatorname{div}\mathbf{e}_{\mathbf{u}}^n-\beta_m e_p^n\right)}_{=:e_\sigma^n}e_p^{n+1}-\delta\int_0^t\int_\gamma\left\{\!\!\left\{\underbrace{\alpha_f\widetilde{\operatorname{div}}[\mathbf{e}_{\mathbf{u}}^n,\mathbf{e}_{\mathbf{U}}^n]-\beta_f e_P^n}_{=:e_\Sigma^n}\right\}\!\!\right\}e_P^{n+1}.\end{aligned}\tag{5.21}$$

We apply Hölder's and Young's inequality to the right hand side of (5.21) and obtain:

$$\begin{aligned}\frac{S_m+\beta_m}{\beta_m^2}\int_0^t \left\|\beta_m e_p^{n+1}\right\|^2_{\mathrm{L}^2(\Omega_m)} + \delta\frac{S_f+\beta_f}{\beta_f^2}\int_0^t \left\|\beta_f e_P^{n+1}\right\|^2_{\mathrm{L}^2(\gamma)} \\ + \frac{\kappa_m}{2}\left\|\nabla\Delta_p^{n+1}\right\|^2_{\mathrm{L}^2(\Omega_m)}(t) + \frac{\delta\kappa_f}{2}\left\{\!\!\left\{\left\|\widetilde{\nabla}[\Delta_p^{n+1},\Delta_P^{n+1}]\right\|^2_{\mathrm{L}^2(\gamma)}(t)\right\}\!\!\right\} \\ \leq \frac{S_m+\beta_m}{2\beta_m^2}\int_0^t \left\|\beta_m e_p^{n+1}\right\|^2_{\mathrm{L}^2(\Omega_m)} + \frac{1}{2(S_m+\beta_m)}\int_0^t \|e_\sigma^n\|^2_{\mathrm{L}^2(\Omega_m)} \\ + \frac{\delta(S_f+\beta_f)}{2\beta_f^2}\int_0^t \|\beta_f e_P^n\|^2_{\mathrm{L}^2(\gamma)} + \frac{\delta}{2(S_f+\beta_f)}\int_0^t \left\{\!\!\left\{\|e_\Sigma^n\|^2_{\mathrm{L}^2(\gamma)}\right\}\!\!\right\}. \quad (5.22)\end{aligned}$$

Subtracting the first and the third term on the right hand side of (5.22) and estimating the coefficients $(S_*+\beta_*)/\beta_*^2$ from below by ω/β_*, $*\in\{m,f\}$, the inequality simplifies as follows:

$$\begin{aligned}\frac{\omega}{2\beta_m}\int_0^t \left\|\beta_m e_p^{n+1}\right\|^2_{\mathrm{L}^2(\Omega_m)} + \frac{\delta\omega}{2\beta_f}\int_0^t \left\|\beta_f e_P^{n+1}\right\|^2_{\mathrm{L}^2(\gamma)} \\ + \frac{\kappa_m}{2}\left\|\nabla\Delta_p^{n+1}\right\|^2_{\mathrm{L}^2(\Omega_m)}(t) + \frac{\delta\kappa_f}{2}\left\{\!\!\left\{\left\|\widetilde{\nabla}[\Delta_p^{n+1},\Delta_P^{n+1}]\right\|^2_{\mathrm{L}^2(\gamma)}(t)\right\}\!\!\right\} \\ \leq \frac{1}{2(S_m+\beta_m)}\int_0^t \|e_\sigma^n\|^2_{\mathrm{L}^2(\Omega_m)} + \frac{\delta}{2(S_f+\beta_f)}\int_0^t \left\{\!\!\left\{\|e_\Sigma^n\|^2_{\mathrm{L}^2(\gamma)}\right\}\!\!\right\}. \quad (5.23)\end{aligned}$$

Now we multiply (5.20) by ω, integrate in time and add to (5.23). The square of the difference formula then yields:

$$\begin{aligned}\frac{\omega}{2\beta_m}\int_0^t \left\|e_\sigma^{n+1}\right\|^2_{\mathrm{L}^2(\Omega_m)} + \frac{\delta\omega}{2\beta_f}\int_0^t \left\{\!\!\left\{\left\|e_\Sigma^{n+1}\right\|^2_{\mathrm{L}^2(\gamma)}\right\}\!\!\right\} \\ + \frac{\kappa_m}{2}\left\|\nabla\Delta_p^{n+1}\right\|^2_{\mathrm{L}^2(\Omega_m)}(t) + \frac{\delta\kappa_f}{2}\left\{\!\!\left\{\left\|\widetilde{\nabla}[\Delta_p^{n+1},\Delta_P^{n+1}]\right\|^2_{\mathrm{L}^2(\gamma)}(t)\right\}\!\!\right\} \\ \leq \frac{1}{2(S_m+\beta_m)}\int_0^t \|e_\sigma^n\|^2_{\mathrm{L}^2(\Omega_m)} + \frac{\delta}{2(S_f+\beta_f)}\int_0^t \left\{\!\!\left\{\|e_\Sigma^n\|^2_{\mathrm{L}^2(\gamma)}\right\}\!\!\right\}.\end{aligned}$$

Using the fact that $\frac{1}{S_*+\beta_*} \leq \frac{1}{\omega^2}\frac{\omega}{\beta_*}$, $*\in\{m,f\}$, the right hand side of the above inequality can be estimated from above and we obtain:

$$\begin{aligned}\frac{1}{2}\left(\rho_{\boldsymbol{\beta}}((p^{n+1},P^{n+1}),(p^n,P^n))\right)^2 \leq \frac{1}{\omega^2}\left(\frac{\omega}{2\beta_m}\int_0^t \|e_\sigma^n\|^2_{\mathrm{L}^2(\Omega_m)} + \frac{\delta\omega}{2\beta_f}\int_0^t \left\{\!\!\left\{\|e_\Sigma^n\|^2_{\mathrm{L}^2(\gamma)}\right\}\!\!\right\}\right) \\ \leq \frac{1}{2\omega^2}\left(\rho_{\boldsymbol{\beta}}((p^n,P^n),(p^{n-1},P^{n-1}))\right)^2.\end{aligned}$$

Hence

$$\rho_{\boldsymbol{\beta}}((p^{n+1},P^{n+1}),(p^n,P^n)) \leq \frac{1}{\omega}\rho_{\boldsymbol{\beta}}((p^n,P^n),(p^{n-1},P^{n-1})).$$

It is readily seen that $\omega > 1$. Since ω is non-increasing with respect to β_m and β_f, the smallest constant of contraction is obtained for the smallest possible values of β_m, β_f. □

As we noted above, Theorem 5.8 implies the existence of a unique solution to (5.1). Hence Theorem 5.1 is proved.

Conclusion

We have derived a mixed-dimensional model of flow and linear elasticity in fractured porous media. Under the assumption of permeable fracture, the existence and uniqueness of weak solution was proved using the fixed-stress splitting scheme. The convergence of this scheme has been analyzed and optimal values of iteration parameters were found in agreement e.g. with [6]. The numerical solution based on the proposed splitting will be the subject of a following paper. The problem can be further extended to the case of embedded fracture with variable aperture and intersecting fractures, further one should consider contact and friction conditions (such as in [2]), to name a few challenging tasks.

References

[1] P. Angot, F. Boyer, and F. Hubert. Asymptotic and numerical modelling of flows in fractured porous media. *ESAIM: Mathematical Modelling and Numerical Analysis*, 43(2):239–275, 2009.

[2] R. L. Berge, I. Berre, E. Keilegavlen, J. M. Nordbotten, and B. Wohlmuth. Finite volume discretization for poroelastic media with fractures modeled by contact mechanics. *International Journal for Numerical Methods in Engineering*, 2019.

[3] M. A. Biot. General theory of three-dimensional consolidation. *Journal of Applied Physics*, 12(2):155–164, 1941.

[4] W. M. Boon and J. M. Nordbotten. Stable mixed finite elements for linear elasticity with thin inclusions. *arXiv preprint arXiv:1903.01757*, 2019.

[5] W. M. Boon, J. M. Nordbotten, and J. E. Vatne. Functional analysis and exterior calculus on mixed-dimensional geometries. *arXiv preprint arXiv:1710.00556*, 2017.

[6] J. W. Both, M. Borregales, J. M. Nordbotten, K. Kumar, and F. A. Radu. Robust fixed stress splitting for Biot's equations in heterogeneous media. *Applied Mathematics Letters*, 68:101–108, 2017.

[7] J. Březina and J. Stebel. Analysis of model error for a continuum-fracture model of porous media flow. In *High Performance Computing in Science and Engineering, Lecture Notes in Computer Science*, pages 152–160. Springer, 2016.

[8] L. C. Evans. *Partial differential equations*, volume 19. American Mathematical Society, 2010.

[9] L. Formaggia, A. Fumagalli, A. Scotti, and P. Ruffo. A reduced model for Darcy's problem in networks of fractures. *ESAIM: Mathematical Modelling and Numerical Analysis*, 48(4):1089–1116, 2014.

[10] B. Ganis, V. Girault, M. Mear, G. Singh, and M. Wheeler. Modeling fractures in a poro-elastic medium. *Oil & Gas Science and Technology–Revue d'IFP Energies nouvelles*, 69(4):515–528, 2014.

[11] K. K. Hanowski and O. Sander. The hydromechanical equilibrium state of poroelastic media with a static fracture: A dimension-reduced model with existence results in weighted Sobolev spaces and simulations with an XFEM discretization. *Mathematical Models and Methods in Applied Sciences*, 28(13):2511–2556, 2018.

[12] V. Martin, J. Jaffré, and J. E. Roberts. Modeling fractures and barriers as interfaces for flow in porous media. *SIAM Journal on Scientific Computing*, 26(5):1667, 2005.

[13] J. Maryška, O. Severýn, and M. Vohralík. Numerical simulation of fracture flow with a mixed-hybrid FEM stochastic discrete fracture network model. *Computational Geosciences*, 8(3):217–234, 2005.

[14] A. Mikelić and J. Tambača. Derivation of a poroelastic elliptic membrane shell model. *Applicable Analysis*, 98(1-2):136–161, 2019.

[15] A. Mikelić and M. F. Wheeler. Convergence of iterative coupling for coupled flow and geomechanics. *Computational Geosciences*, 17(3):455–461, 2013.

[16] G. Pichot, J. Erhel, and J.-R. De Dreuzy. A generalized mixed hybrid mortar method for solving flow in stochastic discrete fracture networks. *SIAM Journal on Scientific Computing*, 34(1):B86–B105, 2012.

[17] J. Rutqvist and O. Stephansson. The role of hydromechanical coupling in fractured rock engineering. *Hydrogeology Journal*, 11(1):7–40, 2003.

[18] N. Schwenck, B. Flemisch, R. Helmig, and B. I. Wohlmuth. Dimensionally reduced flow models in fractured porous media: crossings and boundaries. *Computational Geosciences*, 19(6):1219–1230, 2015.
[19] R. E. Showalter. Diffusion in poro-elastic media. *Journal of Mathematical Analysis and Applications*, 251 (1):310–340, 2000.
[20] J. A. White, N. Castelletto, and H. A. Tchelepi. Block-partitioned solvers for coupled poromechanics: A unified framework. *Computer Methods in Applied Mechanics and Engineering*, 303:55–74, 2016.
[21] A. Ženíšek. The existence and uniqueness theorem in Biot's consolidation theory. *Aplikace Matematiky*, 29 (3):194–211, 1984.

Appendix A. Auxiliary inequalities and identities

In what follows we prove some technical relations that are used in the paper.

Lemma A.1. *For all functions* $(\mathbf{v}, \mathbf{V}) \in \mathcal{V}$ *it holds:*

$$\|\operatorname{div} \mathbf{v}\|^2_{\mathrm{L}^2(\Omega_m)} \leq d \, \|\varepsilon(\mathbf{v})\|^2_{\mathrm{L}^2(\Omega_m)}\,, \tag{A.1}$$

$$\left\|\widetilde{\operatorname{div}}^{\oplus}[\mathbf{v}, \mathbf{V}]\right\|^2_{\mathrm{L}^2(\gamma)} \leq d \left\|\tilde{\varepsilon}^{\oplus}[\mathbf{v}, \mathbf{V}]\right\|^2_{\mathrm{L}^2(\gamma)}. \tag{A.2}$$

Proof. For any tensor $\mathbb{A} \in \mathbf{R}^{d\times d}$, the AM-QM inequality implies

$$|\mathbb{I} : \mathbb{A}|^2 \leq d|\mathbb{A}|^2.$$

Choosing $\mathbb{A} := \varepsilon(\mathbf{v})$ and integrating the inequality over Ω_m we obtain (A.1). Similarly, (A.2) is obtained for $\mathbb{A} := \tilde{\varepsilon}^{\oplus}[\mathbf{v}, \mathbf{V}]$ and integration over γ. □

Lemma A.2. *The functions* $\mathbf{t}^{\oplus}$ *and* $\varphi^{\oplus}$*, defined in* (4.1) *and* (4.5)*, satisfy:*

$$[\![\mathbf{t}[\mathbf{v}, \mathbf{V}] \cdot (\mathbf{w} - \mathbf{W})]\!] = \delta \left\{\!\!\left\{ \mathbb{C}_f \widetilde{\nabla}[\mathbf{v}, \mathbf{V}] : \widetilde{\nabla}_\nu[\mathbf{w}, \mathbf{W}] \right\}\!\!\right\}, \tag{A.3}$$

$$[\![\varphi[q, Q](r - R)]\!] = \delta \left\{\!\!\left\{ \mathbb{K}_f \widetilde{\nabla}[q, Q] \cdot \widetilde{\nabla}_\nu[r, R] \right\}\!\!\right\} \tag{A.4}$$

for all admissible functions $\mathbf{v}, \mathbf{V}, \mathbf{w}, \mathbf{W}, q, Q, r, R$.

Proof. From the definition of $\mathbf{t}^{\oplus}$ and $\widetilde{\nabla}^{\oplus}$ it follows that

$$\mathbf{t}^{\oplus}[\mathbf{v}, \mathbf{V}] \cdot (\mathbf{w}^{\oplus} - \mathbf{W}) = \pm\frac{\delta}{2}\mathbb{C}_f \widetilde{\nabla}^{\oplus}[\mathbf{v}, \mathbf{V}] : \left(\pm\frac{2}{\delta}\right)(\mathbf{w}^{\oplus} - \mathbf{W}) \otimes \nu = \pm\frac{\delta}{2}\mathbb{C}_f \widetilde{\nabla}^{\oplus}[\mathbf{v}, \mathbf{V}] : \widetilde{\nabla}^{\oplus}_\nu[\mathbf{w}, \mathbf{W}].$$

Taking the jump of the above expression then yields (A.3). The second identity can be proved in analogous way. □

Appendix B. Energy equality for (4.8)

In this section we derive the identity (4.9). We take the time derivative of (4.8a), multiply by $\partial_t \mathbf{u}$ and integrate over Ω_m. Then we apply the Green identity, interface and boundary conditions (4.8e), (4.8g), which yields:

$$\int_{\Omega_m} (\mathbb{C}_m \nabla \partial_t \mathbf{u} : \nabla \partial_t \mathbf{u} - \alpha_m \partial_t p \operatorname{div} \partial_t \mathbf{u}) + \int_\gamma [\![\mathbf{t}[\partial_t \mathbf{u}, \partial_t \mathbf{U}] \cdot \partial_t \mathbf{u}]\!] - \int_\gamma \alpha_f \, [\![\partial_t p \partial_t \mathbf{u}]\!] \cdot \nu = \int_{\Omega_m} \partial_t \mathbf{f} \cdot \partial_t \mathbf{u}. \tag{B.1}$$

Similarly we differentiate (4.8c) with respect to time, multiply by $\partial_t \mathbf{U}$, integrate over γ, use Green's formula and apply the boundary conditions (4.8h). We arrive at the identity

$$\delta \int_\gamma \mathbb{C}_f \left\{\!\!\left\{ \widetilde{\nabla}[\partial_t \mathbf{u}, \partial_t \mathbf{U}] \right\}\!\!\right\} : \nabla_\tau \partial_t \mathbf{U} - \delta \int_\gamma \alpha_f \partial_t P \operatorname{div}_\tau \partial_t \mathbf{U} + \int_\gamma \alpha_f \left[\!\left[\partial_t p \right]\!\right] \partial_t \mathbf{U} \cdot \nu - \int_\gamma \left[\!\left[\mathbf{t}[\partial_t \mathbf{u}, \partial_t \mathbf{U}] \right]\!\right] \cdot \partial_t \mathbf{U} = \int_\gamma \partial_t \mathbf{F} \cdot \partial_t \mathbf{U}. \quad \text{(B.2)}$$

Next we multiply (4.8b) by $\partial_t p$ and (4.8d) by $\partial_t P$, integrate over Ω_m, γ, respectively, use Green's formula, interface and boundary conditions (4.8f), (4.8g), (4.8h). We get the following equalities:

$$S_m \int_{\Omega_m} |\partial_t p|^2 + \int_{\Omega_m} \alpha_m \partial_t p \operatorname{div} \partial_t \mathbf{u} + \int_\gamma \left[\!\left[\varphi[p, P] \partial_t p \right]\!\right] + \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \int_{\Omega_m} \mathbb{K}_m \nabla p \cdot \nabla p = \int_{\Omega_m} \partial_t g \partial_t p, \quad \text{(B.3)}$$

$$\delta S_f \int_\gamma |\partial_t P|^2 + \delta \int_\gamma \alpha_f \partial_t P \operatorname{div}_\tau \partial_t \mathbf{U} + \int_\gamma \alpha_f \partial_t P \left[\!\left[\partial_t \mathbf{u} \right]\!\right] \cdot \nu + \delta \int_\gamma \mathbb{K}_f \left\{\!\!\left\{ \widetilde{\nabla}[p, P] \right\}\!\!\right\} \cdot \nabla_\tau \partial_t P - \int_\gamma \left[\!\left[\varphi[p, P] \right]\!\right] \partial_t P = \int_\gamma \partial_t G \partial_t P. \quad \text{(B.4)}$$

Summing up (B.1)–(B.4) some terms cancel out and we obtain:

$$\begin{aligned} &\int_{\Omega_m} (\mathbb{C}_m \nabla \partial_t \mathbf{u} : \nabla \partial_t \mathbf{u}) + \int_\gamma \left[\!\left[\mathbf{t}[\partial_t \mathbf{u}, \partial_t \mathbf{U}] \cdot \partial_t (\mathbf{u} - \mathbf{U}) \right]\!\right] - \int_\gamma \alpha_f \left[\!\left[\partial_t p \partial_t \mathbf{u} \right]\!\right] \cdot \nu \\ &\quad + \delta \int_\gamma \mathbb{C}_f \left\{\!\!\left\{ \widetilde{\nabla}[\partial_t \mathbf{u}, \partial_t \mathbf{U}] \right\}\!\!\right\} : \nabla_\tau \partial_t \mathbf{U} + \int_\gamma \alpha_f \left[\!\left[\partial_t p \right]\!\right] \partial_t \mathbf{U} \cdot \nu + S_m \int_{\Omega_m} |\partial_t p|^2 + \int_\gamma \left[\!\left[\varphi[p, P] \partial_t (p - P) \right]\!\right] \\ &\quad + \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \int_{\Omega_m} \mathbb{K}_m \nabla p \cdot \nabla p + \delta S_f \int_\gamma |\partial_t P|^2 + \int_\gamma \alpha_f \partial_t P \left[\!\left[\partial_t \mathbf{u} \right]\!\right] \cdot \nu + \delta \int_\gamma \mathbb{K}_f \left\{\!\!\left\{ \widetilde{\nabla}[p, P] \right\}\!\!\right\} \cdot \nabla_\tau \partial_t P \\ &\quad = \int_{\Omega_m} (\partial_t \mathbf{f} \cdot \partial_t \mathbf{u} + \partial_t g \partial_t p) + \int_\gamma (\partial_t \mathbf{F} \cdot \partial_t \mathbf{U} + \partial_t G \partial_t P). \end{aligned} \quad \text{(B.5)}$$

Using (A.3)–(A.4) in (B.5) leads to the identity

$$\begin{aligned} &\int_{\Omega_m} (\mathbb{C}_m \nabla \partial_t \mathbf{u} : \nabla \partial_t \mathbf{u}) + \delta \int_\gamma \left\{\!\!\left\{ \mathbb{C}_f \widetilde{\nabla}[\partial_t \mathbf{u}, \partial_t \mathbf{U}] : \widetilde{\nabla}[\partial_t \mathbf{u}, \partial_t \mathbf{U}] \right\}\!\!\right\} + S_m \int_{\Omega_m} |\partial_t p|^2 + \delta S_f \int_\gamma |\partial_t P|^2 \\ &\quad + \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \int_{\Omega_m} \mathbb{K}_m \nabla p \cdot \nabla p + \delta \int_\gamma \left\{\!\!\left\{ \mathbb{K}_f \widetilde{\nabla}[p, P] \cdot \widetilde{\nabla}[\partial_t p, \partial_t P] \right\}\!\!\right\} - \int_\gamma \alpha_f \left[\!\left[\partial_t p \partial_t \mathbf{u} \right]\!\right] \cdot \nu + \int_\gamma \alpha_f \left[\!\left[\partial_t p \right]\!\right] \partial_t \mathbf{U} \cdot \nu \\ &\quad + \int_\gamma \alpha_f \partial_t P \left[\!\left[\partial_t \mathbf{u} \right]\!\right] \cdot \nu = \int_{\Omega_m} (\partial_t \mathbf{f} \cdot \partial_t \mathbf{u} + \partial_t g \partial_t p) + \int_\gamma (\partial_t \mathbf{F} \cdot \partial_t \mathbf{U} + \partial_t G \partial_t P), \end{aligned}$$

which is in fact (4.9).

Appendix C. Weak formulation of (4.10)

In what follows we derive the integral identities (5.1b) – (5.1c). Multiplying (4.8a) and (4.8c') by test functions $(\mathbf{v}, \mathbf{V}) \in \mathcal{V}$, integrating them by parts, summing and applying the boundary conditions (4.8g)–(4.8h),

we arrive at the identity

$$\int_{\Omega_m} (\mathbb{C}_m \nabla \mathbf{u} : \nabla \mathbf{v} - \alpha_m p \operatorname{div} \mathbf{v}) + \int_\gamma [\![(\mathbb{C}_m \nabla \mathbf{u} - \alpha_m p \mathbb{I}) \nu \cdot \mathbf{v}]\!] + \delta \int_\gamma \mathbb{C}_f \left\{\!\!\left\{ \widetilde{\nabla}[\mathbf{u}, \mathbf{U}] \right\}\!\!\right\} : \nabla_\tau \mathbf{V} \\ - \int_\gamma \alpha_f \delta P \operatorname{div}_\tau \mathbf{V} - \int_\gamma [\![\mathbf{t}[\mathbf{u}, \mathbf{U}] \cdot \mathbf{V}]\!] = \langle (\mathbf{f}, \mathbf{F}), (\mathbf{v}, \mathbf{V}) \rangle_{\mathcal{V}} \,. \quad \text{(C.1)}$$

The second term in (C.1) can be rewritten using the interface conditions (4.8e'), which yields:

$$\int_{\Omega_m} (\mathbb{C}_m \nabla \mathbf{u} : \nabla \mathbf{v} - \alpha_m p \operatorname{div} \mathbf{v}) + \delta \int_\gamma \mathbb{C}_f \left\{\!\!\left\{ \widetilde{\nabla}[\mathbf{u}, \mathbf{U}] \right\}\!\!\right\} : \nabla_\tau \mathbf{V} + \int_\gamma [\![\mathbf{t}[\mathbf{u}, \mathbf{U}] \cdot (\mathbf{v} - \mathbf{V})]\!] \\ - \delta \int_\gamma \alpha_f P \left\{\!\!\left\{ \widetilde{\operatorname{div}}[\mathbf{v}, \mathbf{V}] \right\}\!\!\right\} = \langle (\mathbf{f}, \mathbf{F}), (\mathbf{v}, \mathbf{V}) \rangle_{\mathcal{V}} \,. \quad \text{(C.2)}$$

Using (A.3), we can rewrite (C.2) as follows:

$$\int_{\Omega_m} (\mathbb{C}_m \nabla \mathbf{u} : \nabla \mathbf{v} - \alpha_m p \operatorname{div} \mathbf{v}) + \delta \int_\gamma \left\{\!\!\left\{ \mathbb{C}_f \widetilde{\nabla}[\mathbf{u}, \mathbf{U}] : \widetilde{\nabla}[\mathbf{v}, \mathbf{V}] - \alpha_f P \widetilde{\operatorname{div}}[\mathbf{v}, \mathbf{V}] \right\}\!\!\right\} = \langle (\mathbf{f}, \mathbf{F}), (\mathbf{v}, \mathbf{V}) \rangle_{\mathcal{V}} \,,$$

which is (5.1b). Similarly, multiplying the equations (4.8b), (4.8d) by test functions $(q, Q) \in \mathcal{V}$, integrating them by parts, and applying the boundary conditions (4.8g)–(4.8h), we obtain:

$$\int_{\Omega_m} (q \partial_t (S_m p + \alpha_m \operatorname{div} \mathbf{u}) + \mathbb{K}_m \nabla p \cdot \nabla q) + \int_\gamma [\![\mathbb{K}_m \nabla p \cdot \nu q]\!] + \delta \int_\gamma Q \partial_t \left(S_f P + \alpha_f \left\{\!\!\left\{ \widetilde{\operatorname{div}}[\mathbf{u}, \mathbf{U}] \right\}\!\!\right\} \right) \\ + \delta \int_\gamma \mathbb{K}_f \left\{\!\!\left\{ \widetilde{\nabla}[p, P] \right\}\!\!\right\} \cdot \nabla_\tau Q - \int_\gamma [\![\varphi[p, P]]\!] Q = \langle (g, G), (q, Q) \rangle_{\mathscr{L}} \,.$$

Application of the interface conditions (4.8f) then gives rise to the identity

$$\int_{\Omega_m} (q \partial_t (S_m p + \alpha_m \operatorname{div} \mathbf{u}) + \mathbb{K}_m \nabla p \cdot \nabla q) + \delta \int_\gamma Q \partial_t \left(S_f P + \alpha_f \left\{\!\!\left\{ \widetilde{\operatorname{div}}[\mathbf{u}, \mathbf{U}] \right\}\!\!\right\} \right) \\ + \delta \int_\gamma \mathbb{K}_f \left\{\!\!\left\{ \widetilde{\nabla}[p, P] \right\}\!\!\right\} \cdot \nabla_\tau Q + \int_\gamma [\![\varphi[p, P](q - Q)]\!] = \langle (g, G), (q, Q) \rangle_{\mathscr{L}} \,. \quad \text{(C.3)}$$

Due to (A.4), (C.3) becomes

$$\int_{\Omega_m} (q \partial_t (S_m p + \alpha_m \operatorname{div} \mathbf{u}) + \mathbb{K}_m \nabla p \cdot \nabla q) + \delta \int_\gamma Q \partial_t \left(S_f P + \alpha_f \left\{\!\!\left\{ \widetilde{\operatorname{div}}[\mathbf{u}, \mathbf{U}] \right\}\!\!\right\} \right) \\ + \delta \int_\gamma \left\{\!\!\left\{ \mathbb{K}_f \widetilde{\nabla}[p, P] \cdot \widetilde{\nabla}[q, Q] \right\}\!\!\right\} = \langle (g, G), (q, Q) \rangle_{\mathscr{L}} \,,$$

which is (5.1c).

Dear editors,

we are submitting our manuscript „Mixed-dimensional models of poroelasticity“, dealing with derivation and analysis of a coupled problem of porous media flow and mechanics in a domain with fracture. We believe you find it useful for publication in ESAIM:M2AN.

Sincerely,

Jan Březina and Jan Stebel