

# TRACE IDENTITIES FOR THE TOPOLOGICAL VERTEX

JIM BRYAN, MARTIJN KOOL, BEN YOUNG

ABSTRACT. The topological vertex is a universal series which can be regarded as an object in combinatorics, representation theory, geometry, or physics. It encodes the combinatorics of 3D partitions, the action of vertex operators on Fock space, the Donaldson-Thomas theory of toric Calabi-Yau threefolds, or the open string partition function of  $\mathbb{C}^3$ .

We prove several identities in which a sum over terms involving the topological vertex is expressed as a closed formula, often a product of simple terms, closely related to Fourier expansions of Jacobi forms. We use purely combinatorial and representation theoretic methods to prove our formulas, but we discuss applications to the Donaldson-Thomas invariants of elliptically fibered Calabi-Yau threefolds at the end of the paper.

## 1. INTRODUCTION

The topological vertex  $V_{\lambda\mu\nu} = V_{\lambda\mu\nu}(p)$  is a universal formal Laurent series in  $p$  depending on a triple of partitions  $(\lambda, \mu, \nu)$ . It can be considered as an object in combinatorics, representation theory, geometry, or physics. In combinatorics,  $V_{\lambda\mu\nu}$  is the generating function for the number of 3D partitions with asymptotic legs of type  $(\lambda, \mu, \nu)$  (see Definition 2). In representation theory,  $V_{\lambda\mu\nu}$  is given as the matrix coefficients of a certain vertex operator on Fock space [11]. In geometry,  $V_{\lambda\mu\nu}$  is the basic building block for computing the Donaldson-Thomas/Gromov-Witten invariants of toric Calabi-Yau threefolds [8]; in this incarnation, it can be realized as the generating function for the Euler characteristics of certain Hilbert schemes of curves in  $\mathbb{C}^3$  (see § 6). The topological vertex was first discovered in physics as an open string partition function in type IIA string theory on  $\mathbb{C}^3$  [1]. An explicit expression for  $V_{\lambda\mu\nu}$  in terms of Schur functions was given by [11] (see section 3).

In this paper we prove several “trace identities” in which a sum over certain combinations of the vertex is expressed as a closed formula, often a product of simple terms. The products are closely related to the Fourier expansions of Jacobi forms. Applications of these identities are used to

compute the Donaldson-Thomas partition functions of certain Calabi-Yau threefolds in terms of Jacobi forms [5, 4, 6].

## 2. DEFINITIONS AND THE MAIN RESULT.

In this section we give the combinatorial definition of the vertex and we state our main identities.

**Definition 1.** Let  $(\lambda, \mu, \nu)$  be a triple of ordinary partitions. A 3D partition  $\pi$  asymptotic to  $(\lambda, \mu, \nu)$  is a subset

$$\pi \subset (\mathbb{Z}_{\geq 0})^3$$

satisfying

- (1) if any of  $(i+1, j, k)$ ,  $(i, j+1, k)$ , and  $(i, j, k+1)$  is in  $\pi$ , then  $(i, j, k)$  is also in  $\pi$ , and
- (2) (a)  $(j, k) \in \lambda$  if and only if  $(i, j, k) \in \pi$  for all  $i \gg 0$ ,  
 (b)  $(k, i) \in \mu$  if and only if  $(i, j, k) \in \pi$  for all  $j \gg 0$ ,  
 (c)  $(i, j) \in \nu$  if and only if  $(i, j, k) \in \pi$  for all  $k \gg 0$ .

where we regard ordinary partitions as finite subsets of  $(\mathbb{Z}_{\geq 0})^2$  via their diagram.

Intuitively,  $\pi$  is a pile of boxes in the positive octant of 3-space. Condition (1) means that the boxes are stacked stably with gravity pulling them in the  $(-1, -1, -1)$  direction; condition (2) means that the pile of boxes is infinite along the coordinate axes with cross-sections asymptotically given by  $\lambda$ ,  $\mu$ , and  $\nu$ .

The subset  $\{(i, j, k) : (j, k) \in \lambda\} \subset \pi$  will be called the *leg* of  $\pi$  in the  $i$  direction, and the legs in the  $j$  and  $k$  directions are defined analogously. Let

$$\xi_\pi(i, j, k) = 1 - \# \text{ of legs of } \pi \text{ containing } (i, j, k).$$

We define the renormalized volume of  $\pi$  by

$$|\pi| = \sum_{(i,j,k) \in \pi} \xi_\pi(i, j, k).$$

Note that  $|\pi|$  can be negative.

**Definition 2.** The topological vertex  $V_{\lambda\mu\nu}$  is defined to be

$$V_{\lambda\mu\nu} = \sum_{\pi} p^{|\pi|}$$

where the sum is taken over all 3D partitions  $\pi$  asymptotic to  $(\lambda, \mu, \nu)$ . We regard  $V_{\lambda\mu\nu}$  as a formal Laurent series in  $p$ . Note that  $V_{\lambda\mu\nu}$  is clearly cyclically symmetric in the indices, and reflection about the  $i = j$  plane yields

$$V_{\lambda\mu\nu} = V_{\mu'\lambda'\nu'}$$

where  $'$  denotes conjugate partition:

$$\lambda' = \{(i, j) : (j, i) \in \lambda\}.$$

This definition of topological vertex differs from the vertex  $C(\lambda, \mu, \nu)$  of the physics literature by a normalization factor (and we use the variable  $p$  instead of  $q$ ). Our  $V_{\lambda\mu\nu}$  is equal to  $P(\lambda, \mu, \nu)$  defined by Okounkov, Reshetikhin, and Vafa [11, eqn 3.16]. They derive an explicit formula for  $V_{\lambda\mu\nu} = P(\lambda, \mu, \nu)$  in terms of Schur functions [11, eqns 3.20 and 3.21].

The *rows* or *parts* of  $\lambda$  are the integers  $\lambda_j = \min\{i \mid (i, j) \notin \lambda\}$ , for  $j \geq 0$ . We use the notation

$$|\lambda| = \sum_j \lambda_j, \quad ||\lambda||^2 = \sum_j \lambda_j^2.$$

Let  $\square$  denote the partition with a single part of size 1.

We also use the notation

$$M(p, q) = \prod_{m=1}^{\infty} (1 - p^m q)^{-m}$$

and the shorthand  $M(p) = M(p, 1)$ .

We can now state our main result.

**Theorem 3.** *The following identities hold:*

(1)

$$\sum_{\lambda} q^{|\lambda|} = \prod_{m=1}^{\infty} (1 - q^m)^{-1}$$

(2)

$$\sum_{\lambda} q^{|\lambda|} p^{||\lambda||^2} v_{\lambda' \lambda \emptyset} = M(p) \prod_{d=1}^{\infty} (1 - q^d)^{-1} M(p, q^d)$$

(3)

$$\sum_{\lambda} q^{|\lambda|} \frac{v_{\lambda \square \emptyset}}{v_{\lambda \emptyset \emptyset}} = (1 - p)^{-1} \prod_{d=1}^{\infty} \frac{(1 - q^d)}{(1 - pq^d)(1 - p^{-1}q^d)}$$

(4)

$$\sum_{\lambda} q^{|\lambda|} p \frac{v_{\square \square \lambda}}{v_{\emptyset \emptyset \lambda}} = \prod_{m=1}^{\infty} (1 - q^m)^{-1} \cdot \left\{ 1 + \frac{p}{(1 - p)^2} + \sum_{d=1}^{\infty} \sum_{k|d} k(p^k + p^{-k})q^d \right\}$$

(5)

$$\sum_{\lambda} q^{|\lambda|} p^{||\lambda||^2} \frac{v_{\lambda \square \emptyset}}{v_{\lambda \emptyset \emptyset}} v_{\lambda' \lambda \emptyset} = (1 - p)^{-1} M(p) \prod_{d=1}^{\infty} \frac{M(p, q^d)}{(1 - pq^d)(1 - p^{-1}q^d)}$$

We call these formulas “trace formulas” since the left hand side can be expressed as the traces of certain operators on Fock space. This will be made explicit in section 5.

We note that formula (1) is elementary and well known. We prove Formula (2) in section 3 using the orthogonality properties of skew Schur functions. Formulas (3) and (4) are proved in section 4 using a theorem of Bloch-Okounkov [2]. The most difficult identity to prove is equation (5) which we do in section 5. There we prove that the left hand side of equation (5) is given as the trace of a certain product of operators on Fock space. To compute the trace, we introduce a new trick which involves an “infinite number” of permutations of the operators.

### 3. THE TOPOLOGICAL VERTEX AND SCHUR FUNCTIONS

Okounkov-Reshetikhin-Vafa derived a formula for the topological vertex in terms of skew Schur functions. Translating their formulas [11, 3.20&

3.21] into our notation, we get:

(6)

$$V_{\lambda\mu\nu}(p) = M(p)p^{-\frac{1}{2}(\|\lambda\|^2 + \|\mu\|^2 + \|\nu\|^2)} s_{\nu'}(p^{-\rho}) \sum_{\eta} s_{\lambda'/\eta}(p^{-\nu-\rho}) s_{\mu/\eta}(p^{\nu'-\rho}).$$

Here,  $s_{\alpha/\beta}(x_1, x_2, \dots)$  is the skew Schur function (see for example [7, § 5]) and

$$\rho = \left(-\frac{1}{2}, -\frac{3}{2}, -\frac{5}{2}, \dots\right)$$

so that  $p^{-\nu-\rho}$  is notation for the variable list

$$p^{-\nu-\rho} = \left(p^{-\nu_1+\frac{1}{2}}, p^{-\nu_2+\frac{3}{2}}, \dots\right).$$

Using equation (6) and orthogonality of skew Schur functions [7, 28(a) pg 94] we prove equation (2) as follows:

$$V_{\lambda'\lambda\emptyset} = M(p)p^{-\|\lambda\|^2} \sum_{\eta} s_{\lambda/\eta}(p^{-\rho})^2$$

so

$$\begin{aligned} \sum_{\lambda} q^{|\lambda|} p^{\|\lambda\|^2} V_{\lambda'\lambda\emptyset} &= M(p) \sum_{\lambda, \eta} q^{|\lambda|} (s_{\lambda/\eta}(p^{\frac{1}{2}}, p^{\frac{3}{2}}, \dots))^2 \\ &= M(p) \prod_{d=1}^{\infty} \left( (1 - q^d)^{-1} \prod_{j,k=1}^{\infty} (1 - q^d p^{i-\frac{1}{2}+j-\frac{1}{2}})^{-1} \right) \\ &= M(p) \prod_{d=1}^{\infty} (1 - q^d)^{-1} \prod_{m=1}^{\infty} (1 - q^d p^m)^{-m} \\ &= M(p) \prod_{d=1}^{\infty} (1 - q^d)^{-1} M(p, q^d). \end{aligned}$$

We also use equation 6 to derive the following key formulas:

**Lemma 4.** *The following hold:*

$$\begin{aligned} p^{\frac{1}{2}} \frac{V_{\lambda\Box\emptyset}}{V_{\lambda\emptyset\emptyset}} &= \sum_{i=1}^{\infty} p^{-\lambda_i+i-\frac{1}{2}} \\ p \frac{V_{\lambda\Box\Box}}{V_{\lambda\emptyset\emptyset}} &= 1 - \left( \sum_{i=1}^{\infty} p^{-\lambda_i+i-\frac{1}{2}} \right) \left( \sum_{j=1}^{\infty} p^{\lambda_j-j+\frac{1}{2}} \right) \end{aligned}$$

*Proof.* Applying equation (6) to  $V_{\lambda\Box\emptyset}/V_{\lambda\emptyset\emptyset} = V_{\Box\emptyset\lambda}/V_{\emptyset\emptyset\lambda}$  we see that the left hand side of the first equation is given by

$$s_{\Box}(p^{-\lambda-\rho}) = s_{\Box}(p^{-\lambda_1+\frac{1}{2}}, p^{-\lambda_2+\frac{3}{2}}, \dots) = \sum_{i=1}^{\infty} p^{-\lambda_i+i-\frac{1}{2}}.$$

Similarly,

$$\begin{aligned} p \frac{V_{\lambda \square \square}}{V_{\lambda \emptyset \emptyset}} &= p \frac{V_{\square \square \lambda}}{V_{\emptyset \emptyset \lambda}} = \sum_{\eta} s_{\square/\eta}(p^{-\lambda-\rho}) s_{\square/\eta}(p^{-\lambda'-\rho}) \\ &= 1 + s_{\square}(p^{-\lambda-\rho}) s_{\square}(p^{-\lambda'-\rho}). \end{aligned}$$

In general we have the following relation (see [11, Eqn (3.10)])<sup>1</sup>

$$s_{\lambda/\mu}(p^{\nu+\rho}) = (-1)^{|\lambda|-|\mu|} s_{\lambda'/\mu'}(p^{-\nu'-\rho})$$

so in particular  $s_{\square}(p^{\nu+\rho}) = -s_{\square}(p^{-\nu'-\rho})$  and thus

$$\begin{aligned} p \frac{V_{\lambda \square \square}}{V_{\lambda \emptyset \emptyset}} &= 1 - s_{\square}(p^{-\lambda-\rho}) s_{\square}(p^{\lambda+\rho}) \\ &= 1 - \left( \sum_{i=1}^{\infty} p^{-\lambda_i + i - \frac{1}{2}} \right) \left( \sum_{j=1}^{\infty} p^{\lambda_j - j + \frac{1}{2}} \right) \end{aligned}$$

which proves the lemma.

#### 4. APPLICATIONS OF A THEOREM OF BLOCH-OKOUNKOV

We summarize a result of Bloch-Okounkov [2] and use it apply it to proving more of our trace formulas.

We define the following theta function

$$\Theta(p, q) = \eta(q)^{-3} \sum_{n \in \mathbb{Z}} (-1)^n q^{\frac{1}{2}(n+\frac{1}{2})^2} p^{n+\frac{1}{2}}$$

which, by the Jacobi triple product formula is given by

$$\Theta(p, q) = (p^{\frac{1}{2}} - p^{-\frac{1}{2}}) \prod_{m=1}^{\infty} \frac{(1 - pq^m)(1 - p^{-1}q^m)}{(1 - q^m)^2}.$$

We suppress the  $q$  from the notation:  $\Theta(p) = \Theta(p, q)$ , and we note that

$$\Theta(p) = -\Theta(p^{-1}).$$

**Theorem 5** (Bloch-Okounkov [2]). *Define the  $n$  point correlation function by the formula*

$$F(p_1, \dots, p_n) = \prod_{m=1}^{\infty} (1 - q^m) \sum_{\lambda} q^{\lambda} \prod_{k=1}^n \left( \sum_{i=1}^{\infty} p_k^{\lambda_i - i + \frac{1}{2}} \right).$$

Then

$$F(p) = \frac{1}{\Theta(p)}$$

---

<sup>1</sup>There is a typo in equation 3.10 in [11] — the exponent on the right hand side should be  $\nu' - \rho$ .

and

$$F(p_1, p_2) = \frac{1}{\Theta(p_1 p_2)} \left( p_1 \frac{d}{dp_1} \log(\Theta(p_1)) + p_2 \frac{d}{dp_2} \log(\Theta(p_2)) \right).$$

In [2], formulas for the general  $n$  variable function is given, but we will only need the cases of  $n = 1$  and  $n = 2$ .

Using this theorem, we will prove equations (3) and (4) of the main theorem.

#### 4.1. Proofs of equations (3) and (4).

We apply Lemma 4 and Theorem 5:

$$\begin{aligned} \sum_{\lambda} (1-p) q^{|\lambda|} \frac{V_{\lambda \square \emptyset}}{V_{\lambda \emptyset \emptyset}} &= (p^{-\frac{1}{2}} - p^{\frac{1}{2}}) \sum_{\lambda} q^{|\lambda|} \sum_{i=1}^{\infty} p^{-\lambda_i + i - \frac{1}{2}} \\ &= (p^{-\frac{1}{2}} - p^{\frac{1}{2}}) \prod_{m=1}^{\infty} (1 - q^m)^{-1} F(p^{-1}) \\ &= (p^{-\frac{1}{2}} - p^{\frac{1}{2}}) \prod_{m=1}^{\infty} (1 - q^m)^{-1} \frac{1}{-\Theta(p)} \\ &= \prod_{m=1}^{\infty} \frac{(1 - q^m)}{(1 - pq^m)(1 - p^{-1}q^m)} \end{aligned}$$

which proves equation (3).

Again we apply Lemma 4 and Theorem 5:

$$\begin{aligned} \sum_{\lambda} q^{|\lambda|} p \frac{V_{\lambda \square \square}}{V_{\lambda \emptyset \emptyset}} &= \sum_{\lambda} q^{|\lambda|} \left\{ 1 - \left( \sum_{i=1}^{\infty} p^{-\lambda_i + i - \frac{1}{2}} \right) \left( \sum_{j=1}^{\infty} p^{-\lambda_j - j + \frac{1}{2}} \right) \right\} \\ &= \prod_{m=1}^{\infty} (1 - q^m)^{-1} (1 - F(p, p^{-1})). \end{aligned}$$

From Theorem 5, we see that

$$F(p, p^{-1}) = \lim_{(p_1, p_2) \rightarrow (p, p^{-1})} \frac{1}{\Theta(p_1 p_2)} \left( p_1 \frac{d}{dp_1} \log(\Theta(p_1)) + p_2 \frac{d}{dp_2} \log(\Theta(p_2)) \right).$$

To evaluate this limit, we simplify the above expression. A short computation shows that

$$p \frac{d}{dp} \log(\Theta(p)) = \frac{1}{2} \frac{p+1}{p-1} + \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} (-p^k + p^{-k}) q^{mk}.$$

Thus

$$\begin{aligned}
F(p, p^{-1}) &= \lim_{\substack{(p_1, p_2) \rightarrow \\ (p, p^{-1})}} \left( (p_1 p_2)^{\frac{1}{2}} - (p_1 p_2)^{-\frac{1}{2}} \right)^{-1} \prod_{m=1}^{\infty} \frac{(1 - q^m)^2}{(1 - (p_1 p_2) q^m)(1 - (p_1 p_2)^{-1} q^m)} \\
&\quad \cdot \left\{ \frac{1}{2} \cdot \frac{p_1 + 1}{p_1 - 1} + \frac{1}{2} \cdot \frac{p_2 + 1}{p_2 - 1} + \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} (-p_1^k - p_2^k + p_1^{-k} + p_2^{-k}) q^{mk} \right\} \\
&= \lim_{\substack{(p_1, p_2) \rightarrow \\ (p, p^{-1})}} \frac{-(p_1 p_2)^{\frac{1}{2}}}{1 - p_1 p_2} \cdot \left\{ \frac{p_1 p_2 - 1}{(p_1 - 1)(p_2 - 1)} + \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} (1 - p_1^k p_2^k)(p_1^{-k} + p_2^{-k}) q^{mk} \right\} \\
&= \lim_{\substack{(p_1, p_2) \rightarrow \\ (p, p^{-1})}} (p_1 p_2)^{\frac{1}{2}} \left\{ \frac{1}{(1 - p_1)(1 - p_2)} - \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} \frac{1 - (p_1 p_2)^k}{1 - p_1 p_2} (p_1^{-k} + p_2^{-k}) q^{mk} \right\} \\
&= \frac{1}{(1 - p)(1 - p^{-1})} - \sum_{m=1}^{\infty} \sum_{k=1}^{\infty} k(p^k + p^{-k}) q^{mk}.
\end{aligned}$$

Therefore

$$1 - F(p, p^{-1}) = 1 + \frac{p}{(1 - p)^2} + \sum_{d=1}^{\infty} \sum_{k|d} k(p^k + p^{-k}) q^d$$

which finishes the proof of equation (4).

## 5. VERTEX OPERATORS AND THE PROOF OF EQUATION (5)

There are several sources for vertex operators and the infinite wedge formalism. For consistency, we will follow the notation and conventions of [10, Appendix A].

Let  $V$  be the vector space with basis  $\{\underline{k}\}$ ,  $k \in \mathbb{Z} + \frac{1}{2}$ . We define *Fock space*  $\Lambda^{\frac{\infty}{2}} V$  to be the vector space spanned by vectors

$$v_S = \underline{s_1} \wedge \underline{s_2} \wedge \dots$$

where  $S = \{s_1 > s_2 > \dots\} \subset \mathbb{Z} + \frac{1}{2}$  is any subset such that the sets

$$S_+ = S \cap \left( \mathbb{Z} + \frac{1}{2} \right)_{>0} \quad \text{and} \quad S_- = S^c \cap \left( \mathbb{Z} + \frac{1}{2} \right)_{<0}$$

are both finite. Let  $(\cdot, \cdot)$  be the inner product on  $\Lambda^{\frac{\infty}{2}} V$  such that the basis  $\{v_S\}$  is orthonormal.

For any  $k \in \mathbb{Z} + \frac{1}{2}$  let  $\psi_k$  be the operator

$$\psi_k(f) = \underline{k} \wedge f$$

and let  $\psi_k^*$  be its adjoint.



For any partition  $\lambda = \{\lambda_1 \geq \lambda_2 \geq \dots\}$ , we define the vector

$$v_\lambda = \underbrace{(\lambda_1 - \frac{1}{2})}_{\text{}} \wedge \underbrace{(\lambda_2 - \frac{3}{2})}_{\text{}} \wedge \dots$$

Let  $\Lambda_0^{\frac{\infty}{2}} V \subset \Lambda^{\frac{\infty}{2}} V$  be the subspace spanned by the vectors  $\{v_\lambda\}$  where  $\lambda$  runs over all partitions. We call this *charge zero Fock space*.

The *energy operator*

$$H = \sum_{k>0} k (\psi_k \psi_k^* + \psi_{-k}^* \psi_{-k})$$

acts on the basis  $v_\lambda$  by

$$H v_\lambda = |\lambda| v_\lambda$$

and so the operator  $q^H$  acts by

$$q^H v_\lambda = q^{|\lambda|} v_\lambda$$

where  $q$  is a formal parameter.

For  $n \in \mathbb{Z}$ ,  $n \neq 0$  define

$$\alpha_n = \sum_k \psi_{k-n} \psi_k^*$$

and observe that  $\alpha_n^* = \alpha_{-n}$ .

Following [10], we define the *vertex operators*  $\Gamma_\pm(\mathbf{x})$  which are operators on  $\Lambda_0^{\frac{\infty}{2}} V$  over the coefficient ring given by symmetric functions in an infinite set of variables  $\mathbf{x} = (x_1, x_2, x_3, \dots)$ . Let  $\mathbf{s} = (s_1, s_2, \dots)$

$$s_k = \frac{1}{k} \sum_{i=1}^{\infty} x_i^k$$

be the power sum basis for the ring of symmetric functions and let <sup>2</sup>

$$\Gamma_\pm(\mathbf{x}) = \exp \left( \sum_{n=1}^{\infty} s_n \alpha_\pm \right).$$

Observe that  $\Gamma_\pm^* = \Gamma_\mp$ .

The matrix coefficients of the vertex operators in the  $\{v_\lambda\}$  basis are given by skew Schur functions:

$$(7) \quad (\Gamma_-(\mathbf{x}) v_\mu, v_\lambda) = (v_\mu, \Gamma_+(\mathbf{x}) v_\lambda) = s_{\lambda/\mu}(\mathbf{x}).$$

---

<sup>2</sup>In [10], the argument of  $\Gamma_\pm$  is  $\mathbf{s}$ , and the dependence on the underlying set of variables  $\mathbf{x}$  is left implicit. We prefer to make  $\mathbf{x}$  the explicit argument.

Orthogonality of the skew Schur functions then gives rise to the following commutation equation:

$$\Gamma_+(\mathbf{x})\Gamma_-(\mathbf{y}) = \prod_{i,j} (1 - x_j y_j)^{-1} \Gamma_-(\mathbf{y})\Gamma_+(\mathbf{x}),$$

in particular

$$(8) \quad \Gamma_+(up^{-\rho})\Gamma_-(vp^{-\rho}) = M(p, uv)\Gamma_-(vp^{-\rho})\Gamma_+(up^{-\rho})$$

where recall that  $up^{-\rho} = (up^{\frac{1}{2}}, up^{\frac{3}{2}}, up^{\frac{5}{2}}, \dots)$ .

We let

$$\psi(z) = \sum_i z^i \psi_i \quad \text{and} \quad \psi^*(w) = \sum_j w^{-j} \psi_j^*.$$

The commutation relations of these operators with the vertex operators is given by

$$(9) \quad \begin{aligned} \Gamma_{\pm}(\mathbf{x})\psi(z) &= \prod_{i=1}^{\infty} (1 - x_i z^{\pm 1})^{-1} \psi(z)\Gamma_{\pm}(\mathbf{x}) \\ \Gamma_{\pm}(\mathbf{x})\psi^*(w) &= \prod_{i=1}^{\infty} (1 - x_i w^{\pm 1}) \psi^*(w)\Gamma_{\pm}(\mathbf{x}). \end{aligned}$$

We use operators  $\mathcal{E}_r$  introduced by Okounkov-Pandharipande in [9, § 2.2.4]. For  $r \in \mathbb{Z}$ , let<sup>3</sup>

$$\mathcal{E}_r(p) = \sum_{k \in \mathbb{Z} + \frac{1}{2}} p^{-k + \frac{r}{2}} \psi_{k-r} \psi_k^*.$$

Our variable  $p$  is related to the variable  $z$  in [9] by  $p = e^{-z}$ .

From [9, Eqns 2.9 and 0.18], we see that  $\mathcal{E}_0$  is the diagonal operator given by

$$(10) \quad \mathcal{E}_0(p)v_{\lambda} = \left( \sum_{i=1}^{\infty} p^{-\lambda_i + i - \frac{1}{2}} \right) v_{\lambda}.$$

We define

$$\mathcal{E}(a, p) = \sum_{r \in \mathbb{Z}} a^{-r} \mathcal{E}_r(p)$$

where  $a$  is a formal parameter. A short computation shows that

$$\mathcal{E}(a, p) = \psi(ap^{\frac{1}{2}})\psi^*(ap^{-\frac{1}{2}}).$$

From equation (9) we get

$$\Gamma_{\pm}(\mathbf{x})\mathcal{E}(a, p) = \prod_{i=1}^{\infty} \frac{(1 - a^{\mp 1} p^{\mp \frac{1}{2}} x_i)}{(1 - a^{\mp 1} p^{\pm \frac{1}{2}} x_i)} \mathcal{E}(a, p)\Gamma_{\pm}(\mathbf{x}).$$

---

<sup>3</sup>We avoid the use of normal ordering by allowing coefficients which are Laurent series in  $p^{\frac{1}{2}}$ .

For  $\mathbf{x} = up^{-\rho} = (up^{\frac{1}{2}}, up^{\frac{3}{2}}, up^{\frac{5}{2}}, \dots)$  the above simplifies to

$$(11) \quad \begin{aligned} \Gamma_+(up^{-\rho})\mathcal{E}(a, p) &= (1 - au)\mathcal{E}(a, p)\Gamma_+(up^{-\rho}) \\ \mathcal{E}(a, p)\Gamma_-(up^{-\rho}) &= (1 - a^{-1}u)\Gamma_-(up^{-\rho})\mathcal{E}(a, p). \end{aligned}$$

Finally, it follows from equation (7) that

$$(12) \quad \Gamma_{\pm}(\mathbf{x})q^H = q^H\Gamma_{\pm}(q^{\pm 1}\mathbf{x}).$$

We now write the left hand side of equation (5) in the main theorem as a trace of operators on charge zero Fock space.

**Lemma 6.**

$$\begin{aligned} \sum_{\lambda} q^{|\lambda|} p^{||\lambda||^2} \frac{V_{\lambda \square \emptyset}}{V_{\lambda \emptyset \emptyset}} V_{\lambda' \lambda \emptyset} &= p^{-\frac{1}{2}} \text{Tr} \left( \mathcal{E}_0(p) \Gamma_+(p^{-\rho}) \Gamma_-(p^{-\rho}) q^H \right) \\ &= p^{-\frac{1}{2}} \text{Coeff}_{a^0} \left\{ \text{Tr} \left( \mathcal{E}(a, p) \Gamma_+(p^{-\rho}) \Gamma_-(p^{-\rho}) q^H \right) \right\}. \end{aligned}$$

*Proof.* The following computation uses, in order, the commutation relation for  $\Gamma_+$  and  $\Gamma_-$  (equation (8)), the definition of trace, the formula for  $\mathcal{E}_0(p)$  (equation (10)), Lemma 4, equation (7), and finally Lemma 4 again:

$$\begin{aligned} & p^{-\frac{1}{2}} \text{tr} \left( \mathcal{E}_0(p) \Gamma_+(p^{-\rho}) \Gamma_-(p^{-\rho}) q^H \right) \\ &= p^{-\frac{1}{2}} M(p) \text{tr} \left( \mathcal{E}_0(p) \Gamma_-(p^{-\rho}) \Gamma_+(p^{-\rho}) q^H \right) \\ &= p^{-\frac{1}{2}} M(p) \sum_{\lambda} (v_{\lambda}, \mathcal{E}_0(p) \Gamma_-(p^{-\rho}) \Gamma_+(p^{-\rho}) q^H v_{\lambda}) \\ &= p^{-\frac{1}{2}} \sum_{\lambda} q^{|\lambda|} \left( \sum_{i=1}^{\infty} p^{-\lambda_i + i - \frac{1}{2}} \right) (v_{\lambda}, \Gamma_-(p^{-\rho}) \Gamma_+(p^{-\rho}) v_{\lambda}) \\ &= M(p) \sum_{\lambda} q^{|\lambda|} \frac{V_{\lambda \square \emptyset}}{V_{\lambda \emptyset \emptyset}} (\Gamma_+(p^{-\rho}) V_{\lambda}, \Gamma_+(p^{-\rho}) V_{\lambda}) \\ &= \sum_{\lambda} q^{|\lambda|} \frac{V_{\lambda \square \emptyset}}{V_{\lambda \emptyset \emptyset}} M(p) \sum_{\eta} (s_{\lambda/\eta}(p^{-\rho}))^2 \\ &= \sum_{\lambda} q^{|\lambda|} \frac{V_{\lambda \square \emptyset}}{V_{\lambda \emptyset \emptyset}} p^{||\lambda||^2} V_{\lambda' \lambda \emptyset}. \end{aligned}$$

□

While the operator  $\mathcal{E}_0(p)$  does not have good commutation relations with the vertex operators, the operator  $\mathcal{E}(a, p)$  does. Hence we first replace  $\mathcal{E}_0(p)$  with the more general  $\mathcal{E}(a, p)$ , compute the trace, and then specialize to the  $a^0$  coefficient.

**Lemma 7.**

$$\begin{aligned} \text{tr}(\mathcal{E}(a, p)\Gamma_+(p^{-\rho})\Gamma_-(p^{-\rho})q^H) = \\ \frac{1}{p^{-\frac{1}{2}} - p^{\frac{1}{2}}} M(p) \prod_{m=1}^{\infty} \frac{(1 - q^m a)(1 - q^{m-1} a^{-1})(1 - q^m) M(p, q^m)}{(1 - pq^m)(1 - p^{-1} q^m)}. \end{aligned}$$

*Proof.* Our strategy is the following. We use the cyclic invariance of trace along with the commutation relations for  $\Gamma_+$  to move the operator  $\Gamma_+$  past the other operators cyclically to the right until the operators are back to their original positions, but with new arguments. We perform this operation a countable number of times, eventually making the  $\Gamma_+$  operator disappear<sup>4</sup>. We then employ the same strategy moving  $\Gamma_-$  cyclically to the left a countable number of times until it disappears and we are left with a term which we can evaluate with the Okounkov-Bloch theorem (theorem 5).

We first cyclically commute the operator  $\Gamma_+$  to the right using equations (8), (11), and (12):

$$\begin{aligned} & \text{tr}(\mathcal{E}(a, p)\Gamma_+(p^{-\rho})\Gamma_-(p^{-\rho})q^H) \\ &= M(p) \text{tr}(\mathcal{E}(a, p)\Gamma_-(p^{-\rho})\Gamma_+(p^{-\rho})q^H) \\ &= M(p) \text{tr}(\mathcal{E}(a, p)\Gamma_-(p^{-\rho})q^H\Gamma_+(qp^{-\rho})) \\ &= M(p) \text{tr}(\Gamma_+(qp^{-\rho})\mathcal{E}(a, p)\Gamma_-(p^{-\rho})q^H) \\ &= M(p)(1 - qa) \text{tr}(\mathcal{E}(a, p)\Gamma_+(qp^{-\rho})\Gamma_-(p^{-\rho})q^H). \end{aligned}$$

Cyclically commuting  $\Gamma_+$  to the right a second time we get:

$$\begin{aligned} & \text{tr}(\mathcal{E}(a, p)\Gamma_+(p^{-\rho})\Gamma_-(p^{-\rho})q^H) = \\ & M(p)(1 - qa)M(p, q)(1 - q^2 a) \text{tr}(\mathcal{E}(a, p)\Gamma_+(q^2 p^{-\rho})\Gamma_-(p^{-\rho})q^H). \end{aligned}$$

After performing  $N$  iterations of this strategy, we arrive at

$$\begin{aligned} & \text{tr}(\mathcal{E}(a, p)\Gamma_+(p^{-\rho})\Gamma_-(p^{-\rho})q^H) = \\ & \prod_{d=1}^N M(p, q^{d-1})(1 - q^d a) \text{tr}(\mathcal{E}(a, p)\Gamma_+(q^N p^{-\rho})\Gamma_-(p^{-\rho})q^H). \end{aligned}$$

It follows from Equation (7) that

$$\Gamma_{\pm}(q^N p^{-\rho}) \equiv \text{Id} \pmod{q^N}.$$

---

<sup>4</sup>We learned this trick from [3]

So the above two equations imply that the equation

$$\begin{aligned} \text{tr}(\mathcal{E}(a, p)\Gamma_+(p^{-\rho})\Gamma_-(p^{-\rho})q^H) = \\ \prod_{d=1}^{\infty} M(p, q^{d-1})(1 - q^d a) \text{tr}(\mathcal{E}(a, p)\Gamma_-(p^{-\rho})q^H) \end{aligned}$$

holds to all orders in  $q$  and is hence true as a formal power series in  $q$ .

We now apply the same strategy commuting  $\Gamma_-$  to the left:

$$\begin{aligned} \text{tr}(\mathcal{E}(a, p)\Gamma_-(p^{-\rho})q^H) &= (1 - a^{-1}) \text{tr}(\mathcal{E}(a, p)\Gamma_-(qp^{-\rho})q^H) \\ &= (1 - a^{-1})(1 - a^{-1}q) \text{tr}(\mathcal{E}(a, p)\Gamma_-(q^2p^{-\rho})q^H) \\ &= \dots \\ &= \prod_{d=1}^{\infty} (1 - a^{-1}q^{d-1}) \text{tr}(\mathcal{E}(a, p)q^H) \end{aligned}$$

and so we have proved

$$\begin{aligned} \text{tr}(\mathcal{E}(a, p)\Gamma_+(p^{-\rho})\Gamma_-(p^{-\rho})q^H) = \\ \prod_{d=1}^{\infty} M(p, q^{d-1})(1 - q^d a)(1 - a^{-1}q^{d-1}) \text{tr}(\mathcal{E}(a, p)q^H). \end{aligned}$$

From the definition of  $\mathcal{E}_r(p)$  we see that its matrix entries are all off-diagonal if  $r \neq 0$ . Therefore

$$\begin{aligned} \text{tr}(\mathcal{E}(a, p)q^H) &= \text{tr}(\mathcal{E}_0(p)q^H) \\ &= \sum_{\lambda} q^{|\lambda|} \sum_{i=1}^{\infty} p^{-\lambda+i-\frac{1}{2}} \\ &= (p^{-\frac{1}{2}} - p^{\frac{1}{2}})^{-1} \prod_{m=1}^{\infty} \frac{(1 - q^m)}{(1 - pq^m)(1 - p^{-1}q^m)} \end{aligned}$$

where the last equality follows from the computation in the proof of equation (3) in § 4.1. Combining this with the previous computations finishes the proof of the lemma.  $\square$

Combining Lemmas 6 and 7, we get

$$\begin{aligned} \sum_{\lambda} q^{|\lambda|} p^{||\lambda||^2} \frac{V_{\lambda \square \emptyset}}{V_{\lambda \emptyset \emptyset}} V_{\lambda' \lambda \emptyset} &= \frac{1}{1-p} M(p) \prod_{m=1}^{\infty} \frac{M(p, q^m)}{(1 - pq^m)(1 - p^{-1}q^m)} \\ &\quad \cdot \text{Coeff}_{a^0} \left\{ \prod_{m=1}^{\infty} (1 - q^m a)(1 - q^m a^{-1})(1 - q^m) \right\}. \end{aligned}$$

By the Jacobi triple product identity, we have

$$\prod_{m=1}^{\infty} (1 - q^m a)(1 - q^m a^{-1})(1 - q^m) = \sum_{n=-\infty}^{\infty} q^{\binom{n}{2}} (-a)^n$$

whose  $a^0$  coefficient is 1. Plugging into the previous equation we finish the proof of equation (5).

## 6. GEOMETRY AND APPLICATIONS

$V_{\lambda\mu\nu}(p)$  as generating function for Euler characteristics of Hilbert schemes on  $\mathbb{C}^3$ .

short description of local contributions to DT invariants in threefolds with elliptic fibrations.

- (1) multiple of smooth fiber
- (2) multiple of nodal fiber
- (3) multiple of smooth fiber attached to section
- (4) multiple of nodal fiber attached to section
- (5) multiple of smooth fiber attached to node

connection of formulas with Jacobi forms.

## REFERENCES

- [1] Mina Aganagic, Albrecht Klemm, Marcos Mariño, and Cumrun Vafa. The topological vertex. *Comm. Math. Phys.*, 254(2):425–478, 2005. arXiv:hep-th/0305132.
- [2] Spencer Bloch and Andrei Okounkov. The character of the infinite wedge representation. *Adv. Math.*, 149(1):1–60, 2000. arXiv:alg-geom/9712009.
- [3] Jrmie Bouttier, Guillaume Chapuy, and Sylvie Corteel. From Aztec diamonds to pyramids: steep tilings. arXiv:math/1407.0665.
- [4] Jim Bryan. The Donaldson-Thomas theory of  $K3 \times E$  via the topological vertex. arXiv:math/1504.02920.
- [5] Jim Bryan and Martijn Kool. Donaldson-Thomas invariants of local elliptic surfaces via the topological vertex. In preparation.
- [6] Jim Bryan, George Oberdieck, Rahul Pandharipande, and Qizheng Yin. Curve counting on abelian surfaces and threefolds. arXiv:math/1506.00841.
- [7] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.
- [8] D. Maulik, N. Nekrasov, A. Okounkov, and R. Pandharipande. Gromov-Witten theory and Donaldson-Thomas theory. I. *Compos. Math.*, 142(5):1263–1285, 2006. arXiv:math.AG/0312059.
- [9] A. Okounkov and R. Pandharipande. Gromov-Witten theory, Hurwitz theory, and completed cycles. *Ann. of Math. (2)*, 163(2):517–560, 2006. arXiv:math.AG/0204305.
- [10] Andrei Okounkov. Infinite wedge and random partitions. *Selecta Math. (N.S.)*, 7(1):57–81, 2001. arXiv:9907127.

- [11] Andrei Okounkov, Nikolai Reshetikhin, and Cumrun Vafa. Quantum Calabi-Yau and classical crystals. In *The unity of mathematics*, volume 244 of *Progr. Math.*, pages 597–618. Birkhäuser Boston, Boston, MA, 2006. arXiv:hep-th/0309208.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, ROOM  
121, 1984 MATHEMATICS ROAD, VANCOUVER, B.C., CANADA V6T 1Z2