
Table of Contents

.....	1
Part 1: The finite difference approximation	1
Part 2: The finite difference solution to the diffusion equation	1
Part 3: Implementing the numerical solution	2
Part 4: Discussion	4

John Shuler

GEOS597 Homework #3: Hillslope evolution and plotting

Due: 9/19/2016

```
close all;  
clear all;  
clc;
```

Part 1: The finite difference approximation

Step 1: Taylor Series expansion

$$f(x, t + k) = f(x, t) + k * (\partial^2 f(x, t)) / (\partial t) + (k^2 / 2) * ((\partial^2 f(x, t)) / \partial t^2) + (k^3 / 6) \dots$$

Step 2: The forward difference operator for first derivatives

a.) $O(h)$ represents all terms of order h and higher. For the purposes of finite difference approximation here, these terms are dropped.

b.)
$$\partial f(x, t) / \partial t = (f(x, t + k) - f(x, t)) / k + O(k)$$

Step 3: The centered difference operator for second derivatives

$$\partial^2 f(x, t + k) / \partial t^2 \approx (f(x, t + k) - 2f(x, t) + f(x, t - k)) / k^2$$

Step 4: Approximate the derivatives

$$\partial f(x, t) / \partial t \approx (f(x, t + k) - f(x, t)) / k$$

Part 2: The finite difference solution to the diffusion equation

Step 1: Approximate the partial differential equation

$$(z(x, t + k) - z(x, t))/k \approx \kappa * ((z(x + h, t) - 2(z(x, t) + z(x - h, t))/h^2)$$

Step 2: Solve for the value of the function at time $t+k$

$$z(x, t + k) \approx \kappa * k * ((z(x + h, t) - 2(z(x, t) + z(x - h, t))/h^2) + z(x, t)$$


Part 3: Implementing the numerical solution


Step 1: Define parameters and constants

```
kappa = 2*10^-3;           % topographic diffusivity [m^2/year]
dt     = 1;                % time step in [years]
dx     = 1;                % space interval [m]
%
%
% *Step 2: Make the initial model*
%
z = [0 0 0 0 0 1 2 3 4 5 6 7 8 9 10 9 8 7 6 5 4 3 2 1 0 0 0 0 0]; %
    [m]
nNode = numel( z ); % [No] number of elements in the x-direction
xArray = ( 0 : nNode - 1 ) .* dx; % [m] make the x-position vector
%
% *At x = 14 m, z = 10 m, the maximum elevation.*
%
figure (1);
plot (xArray,z,'o');
title ('Initial Hillslope Topography')
xlabel ('Position (m)')
ylabel ('Elevation (m)')
legend ('Slope surface')
axis ([0 30 0 12])
set(gcf,'PaperUnits','inches','PaperPosition',[0 0 6 6])
print -dpng InitialCondition -r100
grid on
%
% *Step 3: Loop through time to compute the topography at* _t+dt_
%
tMax = 100;                % max time steps [years]
t0   = dt;
zNew = zeros (nNode,tMax);
zNew(:,1) = z;
for it=t0+dt:dt:tMax;
    for ix=2:nNode-1;
        zNew(ix,it)=dt*kappa* ((zNew(ix+1,it-1) - 2*(zNew(ix,it-1)) +
            zNew(ix-1,it-1))/(dx^2)) + zNew(ix,it-1);
    end
end
%
% *Step 4: Plot results*
%
figure (2);
plot (xArray,z)
```

```

hold on;
plot (xArray,zNew(:,tMax))
legend ('Initial Slope Profile','Profile After 100 years')
title ('Moraine Erosion Model')
xlabel ('Position (m)')
ylabel ('Elevation (m)')
axis ([0 30 0 12])
grid on
%
```





Part 4: Discussion

Increasing t_{Max} from 100 years to 1 million years significantly increases the rounding of the top of the moraine. More material is removed and deposited at the foot if the duration of exposure is increased, all other variables being constant. This is consistent with our current understanding of moraine erosion and deposition (Putkonen et al, 2008). In this example, our final moraine height varied from ~9.5 m after 100 years to < 0.5 m after 100,000 years.

Values of K_s most often vary within one order of magnitude of the initial value used here, according to our lecture notes. Decreasing K_s by an order of magnitude results in decreased sediment movement. Increasing K_s by an order of magnitude has the opposite effect, illustrating the necessity of accurate estimates of K_s to increase model performance. In this example, for $t_{\text{Max}} = 1000$, the final height of the moraine varied by ~ 5 meters from $\kappa = 2e^{-4}$ to $\kappa = 2e^{-2}$.

Published with MATLAB® R2016a