

SWAT

Generated by Doxygen 1.8.16

1 SWAT	1
2 Modules Index	7
2.1 Modules List	7
3 Data Type Index	9
3.1 Data Types List	9
4 File Index	11
4.1 File List	11
5 Module Documentation	15
5.1 parm Module Reference	15
5.1.1 Detailed Description	99
5.1.2 Variable Documentation	99
5.1.2.1 igropt	99
6 Data Type Documentation	101
6.1 parm::fcgd Interface Reference	101
6.2 parm::ndenit Interface Reference	101
6.3 parm::regres Interface Reference	101
6.4 parm::tair Interface Reference	102
6.5 parm::theta Interface Reference	102
7 File Documentation	103
7.1 albedo.f90 File Reference	103
7.1.1 Detailed Description	103
7.2 allocate_parms.f90 File Reference	103
7.2.1 Detailed Description	103
7.3 alph.f90 File Reference	104
7.3.1 Detailed Description	104
7.3.2 Function/Subroutine Documentation	104
7.3.2.1 alph()	104
7.4 apply.f90 File Reference	104
7.4.1 Detailed Description	105
7.4.2 Function/Subroutine Documentation	105
7.4.2.1 apply()	105
7.5 ascrv.f90 File Reference	105
7.5.1 Detailed Description	105
7.5.2 Function/Subroutine Documentation	105
7.5.2.1 ascrv()	106
7.6 atri.f90 File Reference	106
7.6.1 Detailed Description	106
7.6.2 Function/Subroutine Documentation	106

7.6.2.1 atri()	107
7.7 aunif.f90 File Reference	107
7.7.1 Detailed Description	107
7.7.2 Function/Subroutine Documentation	107
7.7.2.1 aunif()	107
7.8 bmpinit.f90 File Reference	108
7.8.1 Detailed Description	108
7.8.2 Function/Subroutine Documentation	108
7.8.2.1 bmpinit()	108
7.9 burnop.f90 File Reference	108
7.9.1 Detailed Description	109
7.9.2 Function/Subroutine Documentation	109
7.9.2.1 burnop()	109
7.10 canopyint.f90 File Reference	109
7.10.1 Detailed Description	109
7.11 caps.f90 File Reference	109
7.11.1 Detailed Description	110
7.11.2 Function/Subroutine Documentation	110
7.11.2.1 caps()	110
7.12 cfactor.f90 File Reference	110
7.12.1 Detailed Description	110
7.13 clgen.f90 File Reference	110
7.13.1 Detailed Description	111
7.13.2 Function/Subroutine Documentation	111
7.13.2.1 clgen()	111
7.14 clicon.f90 File Reference	111
7.14.1 Detailed Description	111
7.14.2 Function/Subroutine Documentation	111
7.14.2.1 clicon()	112
7.15 command.f90 File Reference	112
7.15.1 Detailed Description	112
7.15.2 Function/Subroutine Documentation	112
7.15.2.1 command()	112
7.16 crackflow.f90 File Reference	113
7.16.1 Detailed Description	113
7.17 crackvol.f90 File Reference	113
7.17.1 Detailed Description	113
7.18 curno.f90 File Reference	113
7.18.1 Detailed Description	113
7.18.2 Function/Subroutine Documentation	114
7.18.2.1 curno()	114
7.19 dailycn.f90 File Reference	114

7.19.1 Detailed Description	114
7.20 dstn1.f90 File Reference	114
7.20.1 Detailed Description	115
7.20.2 Function/Subroutine Documentation	115
7.20.2.1 dstn1()	115
7.21 ee.f90 File Reference	115
7.21.1 Detailed Description	115
7.21.2 Function/Subroutine Documentation	116
7.21.2.1 ee()	116
7.22 eiusle.f90 File Reference	116
7.22.1 Detailed Description	116
7.23 estimate_ksat.f90 File Reference	116
7.23.1 Detailed Description	116
7.23.2 Function/Subroutine Documentation	117
7.23.2.1 estimate_ksat()	117
7.24 etpot.f90 File Reference	117
7.24.1 Detailed Description	117
7.24.2 Function/Subroutine Documentation	117
7.24.2.1 etpot()	117
7.25 expo.f90 File Reference	118
7.25.1 Detailed Description	118
7.25.2 Function/Subroutine Documentation	118
7.25.2.1 expo()	118
7.26 fert.f90 File Reference	118
7.26.1 Detailed Description	119
7.26.2 Function/Subroutine Documentation	119
7.26.2.1 fert()	119
7.27 finalbal.f90 File Reference	119
7.27.1 Detailed Description	119
7.28 gcycl.f90 File Reference	119
7.28.1 Detailed Description	120
7.29 getallo.f90 File Reference	120
7.29.1 Detailed Description	120
7.30 graze.f90 File Reference	120
7.30.1 Detailed Description	120
7.30.2 Function/Subroutine Documentation	120
7.30.2.1 graze()	120
7.31 h2omgt_init.f90 File Reference	121
7.31.1 Detailed Description	121
7.32 harvestop.f90 File Reference	121
7.32.1 Detailed Description	121
7.32.2 Function/Subroutine Documentation	121

7.32.2.1 harvestop()	121
7.33 harvkillop.f90 File Reference	122
7.33.1 Detailed Description	122
7.33.2 Function/Subroutine Documentation	122
7.33.2.1 harvkillop()	122
7.34 headout.f90 File Reference	122
7.34.1 Detailed Description	123
7.35 hmeas.f90 File Reference	123
7.35.1 Detailed Description	123
7.36 hruaa.f90 File Reference	123
7.36.1 Detailed Description	123
7.36.2 Function/Subroutine Documentation	123
7.36.2.1 hruaa()	123
7.37 hruallo.f90 File Reference	124
7.37.1 Detailed Description	124
7.38 hrumon.f90 File Reference	124
7.38.1 Detailed Description	124
7.39 hruyr.f90 File Reference	124
7.39.1 Detailed Description	125
7.40 hydroinit.f90 File Reference	125
7.40.1 Detailed Description	125
7.41 icl.f90 File Reference	125
7.41.1 Detailed Description	125
7.41.2 Function/Subroutine Documentation	125
7.41.2.1 icl()	125
7.42 impnd_init.f90 File Reference	126
7.42.1 Detailed Description	126
7.43 impndmon.f90 File Reference	126
7.43.1 Detailed Description	126
7.44 impndyr.f90 File Reference	126
7.44.1 Detailed Description	127
7.45 irrigate.f90 File Reference	127
7.45.1 Detailed Description	127
7.45.2 Function/Subroutine Documentation	127
7.45.2.1 irrigate()	127
7.46 irrsb.f90 File Reference	127
7.46.1 Detailed Description	128
7.46.2 Function/Subroutine Documentation	128
7.46.2.1 irrsb()	128
7.47 jdt.f90 File Reference	128
7.47.1 Detailed Description	128
7.47.2 Function/Subroutine Documentation	128

7.47.2.1 jdt()	129
7.48 killop.f90 File Reference	129
7.48.1 Detailed Description	129
7.48.2 Function/Subroutine Documentation	129
7.48.2.1 killop()	129
7.49 lid_cistern.f90 File Reference	130
7.49.1 Detailed Description	130
7.49.2 Function/Subroutine Documentation	130
7.49.2.1 lid_cistern()	130
7.50 lid_greenroof.f90 File Reference	130
7.50.1 Detailed Description	131
7.50.2 Function/Subroutine Documentation	131
7.50.2.1 lid_greenroof()	131
7.51 lid_porpavement.f90 File Reference	131
7.51.1 Detailed Description	131
7.51.2 Function/Subroutine Documentation	132
7.51.2.1 lid_porpavement()	132
7.52 lid_raingarden.f90 File Reference	132
7.52.1 Detailed Description	132
7.52.2 Function/Subroutine Documentation	132
7.52.2.1 lid_raingarden()	132
7.53 lidinit.f90 File Reference	133
7.53.1 Detailed Description	133
7.53.2 Function/Subroutine Documentation	133
7.53.2.1 lidinit()	133
7.54 lids.f90 File Reference	133
7.54.1 Detailed Description	134
7.54.2 Function/Subroutine Documentation	134
7.54.2.1 lids()	134
7.55 lwqdef.f90 File Reference	134
7.55.1 Detailed Description	134
7.55.2 Function/Subroutine Documentation	135
7.55.2.1 lwqdef()	135
7.56 main.f90 File Reference	135
7.56.1 Detailed Description	135
7.57 modparm.f90 File Reference	135
7.57.1 Detailed Description	219
7.58 newtillmix.f90 File Reference	219
7.58.1 Detailed Description	219
7.58.2 Function/Subroutine Documentation	219
7.58.2.1 newtillmix()	219
7.59 openwth.f90 File Reference	220

7.59.1 Detailed Description	220
7.60 ovr_sed.f90 File Reference	220
7.60.1 Detailed Description	220
7.61 pgen.f90 File Reference	220
7.61.1 Detailed Description	221
7.61.2 Function/Subroutine Documentation	221
7.61.2.1 pgen()	221
7.62 pgenhr.f90 File Reference	221
7.62.1 Detailed Description	221
7.63 pkq.f90 File Reference	221
7.63.1 Detailed Description	222
7.63.2 Function/Subroutine Documentation	222
7.63.2.1 pkq()	222
7.64 plantop.f90 File Reference	222
7.64.1 Detailed Description	222
7.64.2 Function/Subroutine Documentation	222
7.64.2.1 plantop()	223
7.65 pmeas.f90 File Reference	223
7.65.1 Detailed Description	223
7.65.2 Function/Subroutine Documentation	223
7.65.2.1 pmeas()	223
7.66 qman.f90 File Reference	224
7.66.1 Detailed Description	224
7.66.2 Function/Subroutine Documentation	224
7.66.2.1 qman()	224
7.67 rchaa.f90 File Reference	224
7.67.1 Detailed Description	225
7.67.2 Function/Subroutine Documentation	225
7.67.2.1 rchaa()	225
7.68 rchday.f90 File Reference	225
7.68.1 Detailed Description	225
7.69 rchmon.f90 File Reference	225
7.69.1 Detailed Description	226
7.69.2 Function/Subroutine Documentation	226
7.69.2.1 rchmon()	226
7.70 rchyr.f90 File Reference	226
7.70.1 Detailed Description	226
7.70.2 Function/Subroutine Documentation	226
7.70.2.1 rchyr()	226
7.71 readatmodep.f90 File Reference	227
7.71.1 Detailed Description	227
7.72 readbsn.f90 File Reference	227

7.72.1 Detailed Description	227
7.73 readchm.f90 File Reference	227
7.73.1 Detailed Description	228
7.73.2 Function/Subroutine Documentation	228
7.73.2.1 readchm()	228
7.74 readcnst.f90 File Reference	228
7.74.1 Detailed Description	228
7.74.2 Function/Subroutine Documentation	228
7.74.2.1 readcnst()	228
7.75 readfcst.f90 File Reference	229
7.75.1 Detailed Description	229
7.76 readfert.f90 File Reference	229
7.76.1 Detailed Description	229
7.77 readfig.f90 File Reference	229
7.77.1 Detailed Description	230
7.78 readfile.f90 File Reference	230
7.78.1 Detailed Description	230
7.79 readgw.f90 File Reference	230
7.79.1 Detailed Description	230
7.79.2 Function/Subroutine Documentation	230
7.79.2.1 readgw()	230
7.80 readhru.f90 File Reference	231
7.80.1 Detailed Description	231
7.80.2 Function/Subroutine Documentation	231
7.80.2.1 readhru()	231
7.81 readinpt.f90 File Reference	231
7.81.1 Detailed Description	232
7.82 readlup.f90 File Reference	232
7.82.1 Detailed Description	232
7.83 readlwq.f90 File Reference	232
7.83.1 Detailed Description	232
7.83.2 Function/Subroutine Documentation	232
7.83.2.1 readlwq()	232
7.84 readmgt.f90 File Reference	233
7.84.1 Detailed Description	233
7.84.2 Function/Subroutine Documentation	233
7.84.2.1 readmgt()	233
7.85 readmon.f90 File Reference	233
7.85.1 Detailed Description	234
7.86 readops.f90 File Reference	234
7.86.1 Detailed Description	234
7.86.2 Function/Subroutine Documentation	234

7.86.2.1 readops()	234
7.87 readpest.f90 File Reference	234
7.87.1 Detailed Description	235
7.88 readplant.f90 File Reference	235
7.88.1 Detailed Description	235
7.89 readpnd.f90 File Reference	235
7.89.1 Detailed Description	235
7.89.2 Function/Subroutine Documentation	235
7.89.2.1 readpnd()	235
7.90 readres.f90 File Reference	236
7.90.1 Detailed Description	236
7.90.2 Function/Subroutine Documentation	236
7.90.2.1 readres()	236
7.91 readrte.f90 File Reference	236
7.91.1 Detailed Description	237
7.92 readru.f90 File Reference	237
7.92.1 Detailed Description	237
7.92.2 Function/Subroutine Documentation	237
7.92.2.1 readru()	237
7.93 readsdr.f90 File Reference	237
7.93.1 Detailed Description	238
7.93.2 Function/Subroutine Documentation	238
7.93.2.1 readsdr()	238
7.94 readsepticbz.f90 File Reference	238
7.94.1 Detailed Description	238
7.94.2 Function/Subroutine Documentation	238
7.94.2.1 readsepticbz()	239
7.95 readseptwq.f90 File Reference	240
7.95.1 Detailed Description	240
7.95.2 Function/Subroutine Documentation	240
7.95.2.1 readseptwq()	240
7.96 readsno.f90 File Reference	240
7.96.1 Detailed Description	241
7.96.2 Function/Subroutine Documentation	241
7.96.2.1 readsno()	241
7.97 readsol.f90 File Reference	241
7.97.1 Detailed Description	241
7.97.2 Function/Subroutine Documentation	241
7.97.2.1 readsol()	241
7.98 readsub.f90 File Reference	242
7.98.1 Detailed Description	242
7.98.2 Function/Subroutine Documentation	242

7.98.2.1 readsub()	242
7.99 readswq.f90 File Reference	242
7.99.1 Detailed Description	243
7.100 readtill.f90 File Reference	243
7.100.1 Detailed Description	243
7.101 readurban.f90 File Reference	243
7.101.1 Detailed Description	243
7.102 readwgn.f90 File Reference	243
7.102.1 Detailed Description	244
7.102.2 Function/Subroutine Documentation	244
7.102.2.1 readwgn()	244
7.103 readwus.f90 File Reference	244
7.103.1 Detailed Description	244
7.103.2 Function/Subroutine Documentation	244
7.103.2.1 readwus()	245
7.104 readwwq.f90 File Reference	245
7.104.1 Detailed Description	245
7.105 readyr.f90 File Reference	245
7.105.1 Detailed Description	245
7.105.2 Function/Subroutine Documentation	245
7.105.2.1 readyr()	245
7.106 resetlu.f90 File Reference	246
7.106.1 Detailed Description	246
7.107 rhgen.f90 File Reference	246
7.107.1 Detailed Description	246
7.108 rootfr.f90 File Reference	246
7.108.1 Detailed Description	247
7.108.2 Function/Subroutine Documentation	247
7.108.2.1 rootfr()	247
7.109 rsedaa.f90 File Reference	247
7.109.1 Detailed Description	247
7.109.2 Function/Subroutine Documentation	247
7.109.2.1 rsedaa()	247
7.110 rseday.f90 File Reference	248
7.110.1 Detailed Description	248
7.111 rsedmon.f90 File Reference	248
7.111.1 Detailed Description	248
7.111.2 Function/Subroutine Documentation	248
7.111.2.1 rsedmon()	248
7.112 rsedyr.f90 File Reference	249
7.112.1 Detailed Description	249
7.113 rteinit.f90 File Reference	249

7.113.1 Detailed Description	249
7.114 sched_mgt.f90 File Reference	249
7.114.1 Detailed Description	250
7.114.2 Function/Subroutine Documentation	250
7.114.2.1 sched_mgt()	250
7.115 schedule_ops.f90 File Reference	250
7.115.1 Detailed Description	250
7.115.2 Function/Subroutine Documentation	250
7.115.2.1 schedule_ops()	250
7.116 sim_inityr.f90 File Reference	251
7.116.1 Detailed Description	251
7.117 simulate.f90 File Reference	251
7.117.1 Detailed Description	251
7.118 slrgen.f90 File Reference	251
7.118.1 Detailed Description	252
7.118.2 Function/Subroutine Documentation	252
7.118.2.1 slrgen()	252
7.119 smeas.f90 File Reference	252
7.119.1 Detailed Description	252
7.120 snom.f90 File Reference	252
7.120.1 Detailed Description	253
7.120.2 Function/Subroutine Documentation	253
7.120.2.1 snom()	253
7.121 soil_chem.f90 File Reference	253
7.121.1 Detailed Description	253
7.121.2 Function/Subroutine Documentation	253
7.121.2.1 soil_chem()	253
7.122 soil_phys.f90 File Reference	254
7.122.1 Detailed Description	254
7.122.2 Function/Subroutine Documentation	254
7.122.2.1 soil_phys()	254
7.123 soil_write.f90 File Reference	254
7.123.1 Detailed Description	255
7.123.2 Function/Subroutine Documentation	255
7.123.2.1 soil_write()	255
7.124 solt.f90 File Reference	255
7.124.1 Detailed Description	255
7.125 std1.f90 File Reference	255
7.125.1 Detailed Description	256
7.126 std2.f90 File Reference	256
7.126.1 Detailed Description	256
7.127 std3.f90 File Reference	256

7.127.1 Detailed Description	256
7.128 storeinitial.f90 File Reference	256
7.128.1 Detailed Description	257
7.129 subbasin.f90 File Reference	257
7.129.1 Detailed Description	257
7.129.2 Function/Subroutine Documentation	257
7.129.2.1 subbasin()	257
7.130 submon.f90 File Reference	257
7.130.1 Detailed Description	258
7.131 subyr.f90 File Reference	258
7.131.1 Detailed Description	258
7.132 surface.f90 File Reference	258
7.132.1 Detailed Description	258
7.132.2 Function/Subroutine Documentation	258
7.132.2.1 surface()	258
7.133 surfst_h2o.f90 File Reference	259
7.133.1 Detailed Description	259
7.133.2 Function/Subroutine Documentation	259
7.133.2.1 surfst_h2o()	259
7.134 surq_daycn.f90 File Reference	259
7.134.1 Detailed Description	260
7.134.2 Function/Subroutine Documentation	260
7.134.2.1 surq_daycn()	260
7.135 surq_greenampt.f90 File Reference	260
7.135.1 Detailed Description	260
7.135.2 Function/Subroutine Documentation	260
7.135.2.1 surq_greenampt()	261
7.136 swbl.f90 File Reference	261
7.136.1 Detailed Description	261
7.136.2 Function/Subroutine Documentation	261
7.136.2.1 swbl()	261
7.137 tgen.f90 File Reference	262
7.137.1 Detailed Description	262
7.137.2 Function/Subroutine Documentation	262
7.137.2.1 tgen()	262
7.138 tillfactor.f90 File Reference	262
7.138.1 Detailed Description	262
7.138.2 Function/Subroutine Documentation	263
7.138.2.1 tillfactor()	263
7.139 tmeas.f90 File Reference	263
7.139.1 Detailed Description	263
7.140 tran.f90 File Reference	264

7.140.1 Detailed Description	264
7.140.2 Function/Subroutine Documentation	264
7.140.2.1 tran()	264
7.141 ttcoef.f90 File Reference	264
7.141.1 Detailed Description	264
7.141.2 Function/Subroutine Documentation	265
7.141.2.1 ttcoef()	265
7.142 ttcoef_wway.f90 File Reference	265
7.142.1 Detailed Description	265
7.143 varinit.f90 File Reference	265
7.143.1 Detailed Description	266
7.143.2 Function/Subroutine Documentation	266
7.143.2.1 varinit()	266
7.144 vbl.f90 File Reference	266
7.144.1 Detailed Description	266
7.144.2 Function/Subroutine Documentation	266
7.144.2.1 vbl()	267
7.145 volq.f90 File Reference	267
7.145.1 Detailed Description	267
7.145.2 Function/Subroutine Documentation	268
7.145.2.1 volq()	268
7.146 water_hru.f90 File Reference	268
7.146.1 Detailed Description	268
7.147 wattable.f90 File Reference	268
7.147.1 Detailed Description	268
7.148 weatgn.f90 File Reference	269
7.148.1 Detailed Description	269
7.148.2 Function/Subroutine Documentation	269
7.148.2.1 weatgn()	269
7.149 wmeas.f90 File Reference	269
7.149.1 Detailed Description	269
7.150 wndgen.f90 File Reference	270
7.150.1 Detailed Description	270
7.150.2 Function/Subroutine Documentation	270
7.150.2.1 wndgen()	270
7.151 writea.f90 File Reference	270
7.151.1 Detailed Description	270
7.151.2 Function/Subroutine Documentation	271
7.151.2.1 writea()	271
7.152 writed.f90 File Reference	271
7.152.1 Detailed Description	271
7.153 writem.f90 File Reference	271

7.153.1 Detailed Description	271
7.153.2 Function/Subroutine Documentation	272
7.153.2.1 writem()	272
7.154 xmon.f90 File Reference	272
7.154.1 Detailed Description	272
7.155 ysed.f90 File Reference	272
7.155.1 Detailed Description	272
7.155.2 Function/Subroutine Documentation	273
7.155.2.1 ysed()	273
7.156 zero0.f90 File Reference	273
7.156.1 Detailed Description	273
7.157 zero1.f90 File Reference	273
7.157.1 Detailed Description	273
7.158 zero2.f90 File Reference	274
7.158.1 Detailed Description	274
7.159 zero_urban.f90 File Reference	274
7.159.1 Detailed Description	274
7.160 zeroini.f90 File Reference	274
7.160.1 Detailed Description	274
Bibliography	275
Index	277

Chapter 1

SWAT

An upgraded SWAT 2012 revision 670 code

Objectives

- Standard indentation and translation to Fortran 90 by using `findent`. See the `translate-fortran90.pl` perl script file (:heavy_check_mark:)
- Exhaustive use of the "implicit none" directive to detect bad variable usage (:heavy_check_mark:)
- Generate a GNU `Make` makefile and compile with GNU `GFortran`. See the `generate-makefile.pl` perl script file (:heavy_check_mark:)
- Remove non-used variables and format labels (:heavy_check_mark:)
- Detect and solve all uninitialized variables (:heavy_check_mark: :construction:, some proposed solutions could be incorrect)
- Remove unneeded variable initializations (:heavy_check_mark:) as:

```
j=0 ! this line is not necessary  
j=ihru
```
- Remove redundant code (:heavy_check_mark:)
- Exhaustive use of the "parameter" directive on constants (:heavy_check_mark:)
- Remove global counters (as `i`, `ihru`, `iihru` or `idum` in module `parm`). Using local counters or passing values as argument are preferred (:construction:)
- Generate a detailed list of issues detected in the original code (:heavy_check_mark:, see at the end of this README)
- Remove obsolete commented code (:x:)
- Update variable descriptions in comments (:construction:, a lot of work)
- Standardize comments by using Doxygen style in order to generate documentation. See at `latex/refman.pdf` (:construction:, a lot of work)

Required tools

- `GFortran` (to compile the source code)
- `Make` (to build the executable file)
- `Perl` (optional: to execute the perl scripts to update the makefile or to translate original files to Fortran 90)
- `Findent` (optional: to translate original files to Fortran 90 with a standard indentation)
- `Doxygen` (optional: to generate a reference programming manual from source code)
- `TeX Live` or `MiKTeX` (optional: to generate a reference programming manual from source code)
- On Microsoft Windows systems you have to install `MSYS2` and the required utilities (`GFortran` and `Make`). You can follow detailed instructions in `install-unix`

Instructions to generate Fortran 90 style code from original code

In order to generate Fortran 90 style code with standard indentation from original code you have to type on a UNIX type terminal (you need `Perl` and `Findent`):

```
$ perl translate-fortran90.pl
```

Instructions to generate an initial GNU make Makefile

Type on the UNIX type terminal, when translated the original code to Fortran 90 style (you need `Perl`):

```
$ perl generate-makefile.pl
```

Instructions to generate an executable to test

Type on the UNIX type terminal (you need `GFortran` and `Make`)

- In UNIX type operative systems:

```
$ make
```

- In a `MSYS2` terminal in Microsoft Windows:

```
$ EXE=".exe" LDFLAGS="-static" make
```

- Cross-compiling a 32 bits Microsoft Windows executable in a UNIX type operative system:

```
$ prefix="i686-w64-mingw32-" EXE=".exe" LDFLAGS="-static" make
```

- Cross-compiling a 64 bits Microsoft Windows executable in a UNIX type operative system:

```
$ prefix="x86_64-w64-mingw32-" EXE=".exe" LDFLAGS="-static" make
```

Instructions to generate an optimized executable file

Type on the UNIX type terminal (you need **GFortran** and **Make**)

- In UNIX type operative systems:

```
$ CFLAGS="-march=native -flto" LDFLAGS="-flto" make strip
```

- In a **MSYS2** terminal in Microsoft Windows:

```
$ EXE=".exe" CFLAGS="-flto" LDFLAGS="-flto -static" make strip
```

- Cross-compiling a 32 bits Microsoft Windows executable in a UNIX type operative system:

```
$ prefix="i686-w64-mingw32-" EXE=".exe" CFLAGS="-flto" LDFLAGS="-flto -static" make strip
```

- Cross-compiling a 64 bits Microsoft Windows executable in a UNIX type operative system:

```
$ prefix="x86_64-w64-mingw32-" EXE=".exe" CFLAGS="-flto" LDFLAGS="-flto -static" make strip
```

Instructions to generate a reference programming manual from source code

Type on the UNIX type terminal (you need **Doxygen** and **TeX Live** or **MiKTeX**):

```
$ make latex/refman.pdf
```

The reference programming manual file latex/refman.pdf is generated from source code in PDF format

Issues in the original source code

This is a list of possible issues detected in the original source code. These issues have been mostly detected by the **GFortran** compiler warnings. Some of them could not arise because the logic of the variables is not possible.

- In biofilm.f:

- `dcoef` is used but not initialized. `dcoef=3` as in `watqual.f`? Then, I propose at beginning: `real*8, parameter :: dcoef = 3.`

- In `bmp_ri_pond.f`:

- `qseep` and `qet` could be used not initialized at lines 133 and 134. However the problem only arises for `nstep<1`

- In `bmp_sand_filter.f`:

- `sed_removed` at line 342 could be used not initialized if `sfsedstdev<=0`

- In `bpm_sed_pond.f`:

- `bmp_sed_pond` seems to be `bmp_sed_pond` at line 186

- In `bmp_wet_pond.f`:

- `hvol` could be used not initialized in `ext_dpth` subroutine at line 267 in first bucle iteration

- In `clicon.f`:
 - `tmxbsb`, `tmnbsb`, `rbsb`, `rstopbsb`, `rhdbbsb`, `rabsb`, `rmxbsb`, `daylbsb`, `fradbsb` and `u10bsb` could be used not initialized at 186-207 lines
- In `conapply.f`:
 - `k` and `kk` could be used not initialized at 121-122 lines if `iday_pest(j) /= ipst_freq(j)` and `curyr > nyskip`
- In `confert.f`:
 - `ifrt` seems to be it at line 214
- In `curno.f`:
 - `smxold` could be used not initialized if `cn1(h) <= 1.e-6` and `curyr /= 0` at line 96
- In `drains.f`:
 - `nlayer` could be used not initialized at line 23. However, the problem only arises if it is not set in the previous bucle (`mlyr <= 1` or `sol_z(j1, j) <= 0`)
- In `etact.f`:
 - `sev` could be used not initialized at line 286 if `dep >= esd` and `ly == 2`
- In `filter.f`:
 - `remove21` seems to be `remove2` at line 316
- In `grass_wway.f`:
 - `sf_depth` and `sf_sed` could be used not initialized at lines 133 and 137 if `sf_area > 0` and `sf_area <= 1.e-6`
- In `headout.f`:
 - `hedr` array of column titles is written out of defined bounds at lines 118, 119, 121 and 133. It is written to `mrcho` (set to 62 in `allocate_parms.f` line 59) but in `modparm.f` the bound of `hedr` array is set to 46 (line 663)
- In `hhnoqual.f`:
 - `algon` seems to be `algcon` at line 190
- In `hhwatqual.f`
 - `orgnpin` seems to be `orgpin` at line 278
 - `thour=1.0` at line 377 overwrites previous `thour` calculation. It is wrong
- In `hmeas.f`:
 - `rhdbbsb` could be used not initialized at line 84
- In `killop.f`:
 - `ff1` and `ff2` are used but not initialized at lines 167 and 267. They are set in `harvkillop.f` file (lines 257-258). They have to be included in `modparm.f` to share `harvkillop.f` values? or they have to be redefined as in `harvkillop.f`?
- In `NCsed_leach.f90`:
 - `perc_clyr` could be used not initialized at line 221 if `sol_nly(j) < 2`
- In `nrain.f`:
 - `no2pcp` seems to be `no3pcp` at line 72

- In pmeas.f:
 - rbsb could be used not initialized at line 143
 - flag could be used not initialized if 'a==' 'at line 210 -rainsb could be used not initialized, however only if nstep<=0'
- In pminrl2.f:
 - at line 95 a comma is necessary between base and vara
 - ssp could be used not initialized at line 196 if xx<=1.e-6
- In pothole.f:
 - solp_tileo could be used not initialized at line 593 if pot_vol(j)<=1.e-6 or potvol_↵ tile<=1.e-6
- In potholehr.f:
 - potflow seems to be potflwo at line 447
- In readatmodep.f:
 - momax=12*nbyr is defined at line 65 but not used. It has to be mo_max? but then, it overwrites the file read
- In readops.f:
 - year = 0. seems to be iyear = 0 at line 98
 - mg13 seems to be mgt13 at line 206
- In readpnd.f:
 - vsetsetlpnd seems to be velsetlpnd at line 279
- In readru.f:
 - tck is used but not initialized at line 79
- In readsepticbz.f:
 - at line 135 4. e-8 seems to be 4.e-8
- In rewind_init.f:
 - orig_tnylda is used but not initialized at line 174
- In routels.f:
 - dstor is used but not initialized at line 134. It has to be calculated as in watbal.f? or as in the commented line 109?
 - latqout and gwqout could be used not initialized at lines 142-143
- In rtbact.f:
 - netwtr could be used not initialized at line 124, however only if nstep<1
- In rthpest.f:
 - thour=1.0 at line 183 overwrites previous thour calculation. It is wrong
 - frsol and frsrb could be used not initialized at lines 289-290 if hrtwtr(ii)>0.001 and hrtwtr(ii)/(idt*60)<=0.01
- In rtpest.f:
 - tday=1.0 at line 180 overwrites previous tday calculation. It is wrong
- In sched_mgt.f:

- `< =` seems to be `<=` at 202 line
 - `husc` and `igrow` at lines 264-265 are used but not initialized. `husc` has to be `phu_op(iop, ihru)` has in `readmgt.f?` `igrow` has to be `igro(ihru)` has in `readmgt.f?`
- In `smeas.f`:
 - `rabsb` could be used not initialized at line 86
- In `sweep.f`:
 - `fr_curb` is used but not initialized at line 56. It has to be added to `modparm.f` to share result with `sched_mgt.f?` or it has to be `mgt5op(nop(ihru), ihru)` as in `sched_mgt.f?`
- In `tmeas.f`:
 - `tmxbsb` and `tmnbsb` could be used not initialized at lines 109-110
- In `transfer.f`:
 - `ratio, xx` and `ratio1` could be used not initialized at lines 236, 239 and 241 if `ihout==2`
- In `wmeas.f`:
 - `u10bsb` could be used not initialized at line 85
- In `zero0.f`:
 - `sol_sumn03` seems to be `sol_sumno3` at line 508
- In `zero_urban.f`:
 - `stp_stagdis` seems to be `stp_stagdis` at line 84
 - `subdr_kg` seems to be `subdr_km` at line 149
 - `spl_eros` is not defined at line 21, it could be `eros_spl?`

Chapter 2

Modules Index

2.1 Modules List

Here is a list of all documented modules with brief descriptions:

parm	Main module containing the global variables	15
----------------------	---	--------------------

Chapter 3

Data Type Index

3.1 Data Types List

Here are the data types with brief descriptions:

parm::fcgd	101
parm::ndenit	101
parm::regres	101
parm::tair	102
parm::theta	102

Chapter 4

File Index

4.1 File List

Here is a list of all documented files with brief descriptions:

albedo.f90	103
allocate_parms.f90	103
alph.f90	104
apply.f90	104
ascrv.f90	105
atri.f90	106
aunif.f90	107
bmpinit.f90	108
burnop.f90	108
canopyint.f90	109
caps.f90	109
cfactor.f90	110
clgen.f90	110
clicon.f90	111
command.f90	112
crackflow.f90	113
crackvol.f90	113
curno.f90	113
dailycn.f90	114
dstn1.f90	114
ee.f90	115
eiusle.f90	116
estimate_ksat.f90	116
etpot.f90	117
expo.f90	118
fert.f90	118
finalbal.f90	119
gcycl.f90	119
getallo.f90	120
graze.f90	120
h2omgt_init.f90	121
harvestop.f90	121
harvkillop.f90	122
headout.f90	122
hmeas.f90	123

hruaa.f90	123
hruallo.f90	124
hrumon.f90	124
hruyr.f90	124
hydroinit.f90	125
icl.f90	125
impnd_init.f90	126
impndmon.f90	126
impndyr.f90	126
irrigate.f90	127
irrsb.f90	127
jdt.f90	128
killop.f90	129
lid_cistern.f90	130
lid_greenroof.f90	130
lid_por pavement.f90	131
lid_raingarden.f90	132
lidinit.f90	133
lids.f90	133
lwqdef.f90	134
main.f90	135
modparm.f90	135
newtillmix.f90	219
openwth.f90	220
ovr_sed.f90	220
pgen.f90	220
pgenhr.f90	221
pkq.f90	221
plantop.f90	222
pmeas.f90	223
qman.f90	224
rchaa.f90	224
rchday.f90	225
rchmon.f90	225
rchyr.f90	226
readatmodep.f90	227
readbsn.f90	227
readchm.f90	227
readcnst.f90	228
readfcst.f90	229
readfert.f90	229
readfig.f90	229
readfile.f90	230
readgw.f90	230
readhru.f90	231
readinpt.f90	231
readlup.f90	232
readlwq.f90	232
readmgt.f90	233
readmon.f90	233
readops.f90	234
readpest.f90	234
readplant.f90	235
readpnd.f90	235
readres.f90	236
readrte.f90	236
readru.f90	237
readsdr.f90	237

readsepticbz.f90	238
readseptwq.f90	240
readsno.f90	240
readsol.f90	241
readsub.f90	242
readswq.f90	242
readtill.f90	243
readurban.f90	243
readwgn.f90	243
readwus.f90	244
readwwq.f90	245
readyr.f90	245
resetlu.f90	246
rhgen.f90	246
rootfr.f90	246
rsedaa.f90	247
rseday.f90	248
rsedmon.f90	248
rsedyr.f90	249
rteinit.f90	249
sched_mgt.f90	249
schedule_ops.f90	250
sim_inityr.f90	251
simulate.f90	251
slrgen.f90	251
smeas.f90	252
snom.f90	252
soil_chem.f90	253
soil_phys.f90	254
soil_write.f90	254
solt.f90	255
std1.f90	255
std2.f90	256
std3.f90	256
storeinitial.f90	256
subbasin.f90	257
submon.f90	257
subyr.f90	258
surface.f90	258
surfst_h2o.f90	259
surq_daycn.f90	259
surq_greenampt.f90	260
swbl.f90	261
tgen.f90	262
tillfactor.f90	262
tmeas.f90	263
tran.f90	264
ttcoef.f90	264
ttcoef_wway.f90	265
varinit.f90	265
vbl.f90	266
volq.f90	267
water_hru.f90	268
wattable.f90	268
weatgn.f90	269
wmeas.f90	269
wndgen.f90	270
writea.f90	270

writed.f90	271
writem.f90	271
xmon.f90	272
ysed.f90	272
zero0.f90	273
zero1.f90	273
zero2.f90	274
zero_urban.f90	274
zeroini.f90	274

Chapter 5

Module Documentation

5.1 parm Module Reference

main module containing the global variables

Data Types

- interface [fcgd](#)
- interface [ndenit](#)
- interface [regres](#)
- interface [tair](#)
- interface [theta](#)

Variables

- integer, parameter [mvaro](#) = 33
max number of variables routed through the reach
- integer, parameter [mhruo](#) = 79
maximum number of variables written to HRU output file (output.hru) (none)
- integer, parameter [mrcho](#) = 62
maximum number of variables written to reach output file (.rch) (none)
- integer, parameter [msubo](#) = 24
maximum number of variables written to subbasin output file (output.sub) (none)
- integer, parameter [mstdo](#) = 113
max number of variables summarized in output.std
- integer, parameter **motot** = 600
- character(len=80), parameter [prog](#) = "SWAT Sep 7 VER 2018/Rev 670"
SWAT program header string (name and version)

- integer, dimension(13), parameter **ndays_leap** = (/0,31,60,91,121,152,182,213,244,274,305,335,366/)
- integer, dimension(13), parameter **ndays_noleap** = (/0,31,59,90,120,151,181,212,243,273,304,334,365/)
- integer **icalen**
- real *8 **prf_bsn**
Basinwide peak rate adjustment factor for sediment routing in the channel. Allows impact of peak flow rate on sediment routing and channel reshaping to be taken into account.
- real *8 **co2_x2**
- real *8 **co2_x**
- real *8, dimension(:), allocatable **alph_e**
- real *8, dimension(:), allocatable **cdn**
denitrification exponential rate coefficient
- real *8, dimension(:), allocatable **nperco**
nitrate percolation coefficient (0-1)
0:concentration of nitrate in surface runoff is zero
1:percolate has same concentration of nitrate as surface runoff
- real *8, dimension(:), allocatable **surlag**
Surface runoff lag time. This parameter is needed in subbasins where the time of concentration is greater than 1 day. SURLAG is used to create a "storage" for surface runoff to allow the runoff to take longer than 1 day to reach the subbasin outlet (days)
- real *8, dimension(:), allocatable **co_p**
- real *8, dimension(:), allocatable **cmn**
rate factor for humus mineralization on active organic N
- real *8, dimension(:), allocatable **phoskd**
Phosphorus soil partitioning coefficient. Ratio of soluble phosphorus in surface layer to soluble phosphorus in runoff.
- real *8, dimension(:), allocatable **psp**
Phosphorus availability index. The fraction of fertilizer P remaining in labile pool after initial rapid phase of P sorption (none)
- real *8, dimension(:), allocatable **sdnco**
denitrification threshold: fraction of field capacity triggering denitrification
- real *8 **r2adj_bsn**
basinwide retention parameter adjustment factor (greater than 1)
- real *8 **pst_kg**
amount of pesticide applied to HRU (kg/ha)
- real *8 **yield**
yield (dry weight) (kg)
- real *8 **burn_frlb**
fraction of biomass and residue that burn(input in management file) range (0 - 1.0) (none)
- real *8 **yieldgrn**
- real *8 **yieldbms**
- real *8 **yieldtbr**
- real *8 **yieldn**
- real *8 **yieldp**
- real *8 **hi_bms**
- real *8 **hi_rsd**
- real *8 **yieldrsd**
- real *8, dimension(:), allocatable **I_k1**
- real *8, dimension(:), allocatable **I_k2**
- real *8, dimension(:), allocatable **I_lambda**
- real *8, dimension(:), allocatable **I_beta**
- real *8, dimension(:), allocatable **I_gama**
- real *8, dimension(:), allocatable **I_harea**
- real *8, dimension(:), allocatable **I_vleng**
- real *8, dimension(:), allocatable **I_vslope**

- real *8, dimension(:), allocatable **l_ktc**
- real *8, dimension(:), allocatable **biofilm_mumax**
- real *8, dimension(:), allocatable **biofilm_kinv**
- real *8, dimension(:), allocatable **biofilm_klw**
- real *8, dimension(:), allocatable **biofilm_kla**
- real *8, dimension(:), allocatable **biofilm_cdet**
- real *8, dimension(:), allocatable **biofilm_bm**
- real *8, dimension(:, :), allocatable **hru_rufr**
- real *8, dimension(:, :), allocatable **daru_km**
- real *8, dimension(:, :), allocatable **ru_k**
- real *8, dimension(:, :), allocatable **ru_c**
- real *8, dimension(:, :), allocatable **ru_eiq**
- real *8, dimension(:, :), allocatable **ru_ovsl**
- real *8, dimension(:, :), allocatable **ru_a**
- real *8, dimension(:, :), allocatable **ru_ovs**
- real *8, dimension(:, :), allocatable **ru_ktc**
- real *8, dimension(:), allocatable **gwq_ru**
- real *8, dimension(:), allocatable **qdayout**
- integer, dimension(:), allocatable **ils2**
- integer, dimension(:), allocatable **ils2flag**
- integer **ipest**
pesticide identification number from pest.dat (none)
- integer **iru**
- integer **mrui**
- integer **irch**
- integer **isub**
- integer **mhyd_bsn**
- integer **ils_nofig**
- integer **mhru1**
- integer, dimension(:), allocatable **mhyd1**
- integer, dimension(:), allocatable **irtun**
- real *8 **wshd_sepno3**
- real *8 **wshd_sepnh3**
- real *8 **wshd_seporgn**
- real *8 **wshd_sepfon**
- real *8 **wshd_seporgp**
- real *8 **wshd_sepfop**
- real *8 **wshd_sepsolp**
- real *8 **wshd_sepbod**
- real *8 **wshd_sepmm**
- integer, dimension(:), allocatable **isep_hru**
- real *8 **fixco**
nitrogen fixation coefficient
- real *8 **nfixmx**
maximum daily n-fixation (kg/ha)
- real *8 **res_stlr_co**
reservoir sediment settling coefficient
- real *8 **rsd_covco**
residue cover factor for computing fraction of cover
- real *8 **vcrit**
critical velocity
- real *8 **wshd_snob**
average amount of water stored in snow at the beginning of the simulation for the entire watershed (mm H2O)

- real *8 **wshd_sw**
water in soil at beginning of simulation, or\ average amount of water stored in soil for the entire watershed, or\ difference between mass balance calculated from watershed averages and actual value for water in soil at end of simulation (goal is to have wshd_sw = 0.) (mm H2O)
- real *8 **wshd_pndfr**
fraction of watershed area which drains into ponds (none)
- real *8 **wshd_pndsed**
total amount of suspended sediment in ponds in the watershed (metric tons), or mass balance discrepancy for pond sediment expressed as loading per unit hectare of drainage area (metric tons/ha)
- real *8 **wshd_pndv**
total volume of water in ponds in the watershed (m^3), or mass balance discrepancy for pond water volume expressed as depth over drainage area (mm H2O)
- real *8 **percop**
*pesticide percolation coefficient (0-1)
0: concentration of pesticide in surface runoff is zero
1: percolate has same concentration of pesticide as surface runoff*
- real *8 **wshd_resfr**
fraction of watershed area that drains into reservoirs (none)
- real *8 **wshd_pndha**
watershed area in hectares which drains into ponds (ha)
- real *8 **wshd_resha**
watershed area in hectares which drains into reservoirs (ha)
- real *8 **wshd_wetfr**
fraction of watershed area which drains into wetlands (none)
- real *8 **wshd_fminp**
average annual amount of mineral P applied in watershed (kg P/ha)
- real *8 **wshd_fnh3**
average annual amount of NH3-N applied in watershed (kg N/ha)
- real *8 **wshd_fno3**
average annual amount of NO3-N applied in watershed (kg N/ha)
- real *8 **wshd_forgn**
average annual amount of organic N applied in watershed (kg N/ha)
- real *8 **wshd_ftotn**
average annual amount of N (mineral & organic) applied in watershed (kg N/ha)
- real *8 **wshd_forgp**
average annual amount of organic P applied in watershed (kg P/ha)
- real *8 **wshd_ftotp**
average annual amount of P (mineral & organic) applied in watershed (kg P/ha)
- real *8 **wshd_yldn**
amount of nitrogen removed from soil in watershed in the yield (kg N/ha)
- real *8 **wshd_yldp**
amount of phosphorus removed from soil in watershed in the yield (kg P/ha)
- real *8 **wshd_fixn**
- real *8 **wshd_pup**
- real *8 **wshd_wstrs**
- real *8 **wshd_nstrs**
- real *8 **wshd_pstrs**
- real *8 **wshd_tstrs**
- real *8 **wshd_astrs**
- real *8 **ffcb**
initial soil water content expressed as a fraction of field capacity
- real *8 **wshd_hmn**

- real *8 **wshd_rwn**
- real *8 **wshd_hmp**
- real *8 **wshd_rmn**
- real *8 **wshd_dnit**
- real *8 **wdpq**
die-off factor for persistent bacteria in soil solution (1/day)
- real *8 **wshd_rmp**
- real *8 **wshd_voln**
- real *8 **wshd_nitn**
- real *8 **wshd_pas**
- real *8 **wshd_pal**
- real *8 **wof_p**
wash off fraction for persistent bacteria on foliage during a rainfall event
- real *8 **wshd_plch**
- real *8 **wshd_raino3**
- real *8 **ressedc**
- real *8 **basno3f**
- real *8 **basorgnf**
- real *8 **wshd_pinlet**
- real *8 **wshd_ptile**
- real *8 **sftmp**
Snowfall temperature (deg C)
- real *8 **smfmn**
Minimum melt rate for snow during year (Dec. 21) where deg C refers to the air temperature. (mm/deg C/day)
- real *8 **smfmx**
Maximum melt rate for snow during year (June 21) where deg C refers to the air temperature. SMFMX and SMFMN allow the rate of snow melt to vary through the year. These parameters are accounting for the impact of soil temperature on snow melt. (mm/deg C/day)
- real *8 **smtmp**
Snow melt base temperature. Mean air temperature at which snow melt will occur. (deg C)
- real *8 **wgppq**
growth factor for persistent bacteria in soil solution (1/day)
- real *8 **basminpf**
- real *8 **basorgpf**
- real *8 **wdlpq**
die-off factor for less persistent bacteria in soil solution (1/day)
- real *8 **wshd_ressed**
*total amount of suspended sediment in reservoirs in the watershed (metric tons),
or mass balance discrepancy for reservoir sediment expressed as loading per unit hectare of drainage area (metric tons/ha)*
- real *8 **wshd_resv**
*total volume of water in all reservoirs in the watershed (m^3),
or mass balance discrepancy for reservoir water volume expressed as depth over drainage area (mm H₂O)*
- real *8 **basminpi**
average amount of phosphorus initially in the mineral P pool in watershed soil (kg P/ha)
- real *8 **basno3i**
average amount of nitrogen initially in the nitrate pool in watershed soil (kg N/ha)
- real *8 **basorgni**
average amount of nitrogen initially in the organic N pool in watershed soil (kg N/ha)
- real *8 **wdps**
die-off factor for persistent bacteria adsorbed to soil particles (1/day)
- real *8 **wglpq**
growth factor for less persistent bacteria in soil solution (1/day)

- real *8 **basorgpi**
average amount of phosphorus initially in the organic P pool in watershed soil (kg P/ha)
- real *8 **peakr**
peak runoff rate for the day in HRU (m^3/s)
- real *8 **albday**
albedo of ground for the day in HRU, the fraction of the solar radiation reflected at the soil surface back into space (none)
- real *8 **pndsedin**
sediment inflow to the pond from HRU (metric tons)
- real *8 **sw_excess**
amount of water stored in soil layer on the current day that exceeds field capacity (gravity drained water) (mm H2O)
- real *8 **timp**
*Snow pack temperature lag factor (0-1)
1 = no lag (snow pack temp=current day air temp) as the lag factor goes to zero, the snow pack's temperature will be less influenced by the current day's air temperature.*
- real *8 **wtabelo**
- real *8 **tilep**
- real *8 **wt_shall**
- real *8 **sq_rto**
- real *8 **qtile**
drainage tile flow in HRU soil layer for the day (mm H2O)
- real *8 **infilpcp**
amount of precipitation that infiltrates into soil (enters soil) (mm H2O)
- real *8 **crk**
percolation due to crack flow (mm H2O)
- real *8 **fixn**
amount of nitrogen added to plant biomass via fixation on the day in HRU (kg N/ha)
- real *8 **latlyr**
amount of water in lateral flow in layer in HRU for the day (mm H2O)
- real *8 **snofall**
amount of precipitation falling as freezing rain/snow on day in HRU (mm H2O)
- real *8 **snomlt**
amount of water in snow melt for the day in HRU (mm H2O)
- real *8 **tloss**
amount of water removed from surface runoff via transmission losses on day in HRU (mm H2O)
- real *8 **pndloss**
- real *8 **wetloss**
- real *8 **potloss**
- real *8 **lpndloss**
- real *8 **lwetloss**
- real *8 **bioday**
biomass generated on current day in HRU (kg)
- real *8 **cfertn**
amount of nitrogen added to soil in continuous fertilizer operation on day (kg N/ha)
- real *8 **cfertp**
amount of phosphorus added to soil in continuous fertilizer operation on day (kg P/ha)
- real *8 **fertn**
total amount of nitrogen applied to soil in HRU on day (kg N/ha)
- real *8 **sepday**
micropore percolation from bottom of the soil layer on day in HRU (mm H2O)
- real *8 **sol_rd**
current rooting depth (mm)

- real *8 **sedrch**
- real *8 **sepcrk**
- real *8 **sepcrktot**
- real *8 **fertno3**
- real *8 **fertnh3**
- real *8 **fertorgn**
- real *8 **fertsolp**
- real *8 **fertorgp**
- real *8 **wgps**
growth factor for persistent bacteria adsorbed to soil particles (1/day)
- real *8 **qdfr**
fraction of water yield that is surface runoff (none)
- real *8 **fertp**
total amount of phosphorus applied to soil in HRU on day (kg P/ha)
- real *8 **grazn**
amount of nitrogen added to soil in grazing on the day in HRU (kg N/ha)
- real *8 **grazp**
amount of phosphorus added to soil in grazing on the day in HRU (kg P/ha)
- real *8 **soxy**
saturation dissolved oxygen concentration (mg/L)
- real *8 **sdti**
- real *8 **rtwtr**
- real *8 **ressa**
- real *8 **wdlps**
die-off factor for less persistent bacteria absorbed to soil particles (1/day)
- real *8 **wglps**
growth factor for less persistent bacteria adsorbed to soil particles (1/day)
- real *8 **da_km**
area of the watershed in square kilometers (km²)
- real *8 **rttime**
- real *8 **rchdep**
- real *8 **rtevp**
- real *8 **rttlc**
- real *8 **resflwi**
- real *8 **wdprch**
die-off factor for persistent bacteria in streams (1/day)
- real *8 **resflwo**
- real *8 **respcp**
- real *8 **resev**
- real *8 **ressep**
- real *8 **ressedi**
- real *8 **ressedo**
- real *8 **dtot**
- real *8 **pperco_bsn**
phosphorus percolation coefficient. Ratio of soluble phosphorus in surface to soluble phosphorus in percolate
- real *8 **nperco_bsn**
basin nitrate percolation coefficient (0-1)
0:concentration of nitrate in surface runoff is zero
1:percolate has same concentration of nitrate as surface runoff
- real *8 **rsdco**
residue decomposition coefficient. The fraction of residue which will decompose in a day assuming optimal moisture, temperature, C:N ratio, and C:P ratio
- real *8 **voltot**

- total volume of cracks expressed as depth per unit area (mm)*
- real *8 **phoskd_bsn**
- real *8 **msk_x**
 - weighting factor controlling relative importance of inflow rate and outflow rate in determining storage on reach*
- real *8 **volcrmin**
 - minimum crack volume allowed in any soil layer (mm)*
- real *8 **bactkdq**
 - bacteria soil partitioning coefficient. Ratio of solution bacteria in surface layer to solution bacteria in runoff soluble and sorbed phase in surface runoff.*
- real *8 **wdpf**
 - die-off factor for persistent bacteria on foliage (1/day)*
- real *8 **canev**
 - amount of water evaporated from canopy storage (mm H₂O)*
- real *8 **precipday**
 - precipitation, or effective precipitation reaching soil surface, for the current day in HRU (mm H₂O)*
- real *8 **uno3d**
 - plant nitrogen deficiency for day in HRU (kg N/ha)*
- real *8 **usle**
 - daily soil loss predicted with USLE equation (metric tons/ha)*
- real *8 **rcn**
- real *8 **surlag_bsn**
- real *8 **thbact**
 - temperature adjustment factor for bacteria die-off/growth*
- real *8 **wlpq20**
 - overall rate change for less persistent bacteria in soil solution (1/day)*
- real *8 **wlps20**
 - overall rate change for less persistent bacteria adsorbed to soil particles (1/day)*
- real *8 **wpq20**
 - overall rate change for persistent bacteria in soil solution (1/day)*
- real *8 **wps20**
 - overall rate change for persistent bacteria adsorbed to soil particles (1/day)*
- real *8 **bactrop**
 - persistent bacteria transported to main channel with surface runoff (# colonies/ha)*
- real *8 **bactsedp**
 - persistent bacteria transported with sediment in surface runoff (# colonies/ha)*
- real *8 **wgpf**
 - growth factor for persistent bacteria on foliage (1/day)*
- real *8 **bactlchlp**
 - less persistent bacteria removed from soil surface layer by percolation (# colonies/ha)*
- real *8 **bactlchp**
 - persistent bacteria removed from soil surface layer by percolation (# colonies/ha)*
- real *8 **enratio**
 - enrichment ratio calculated for day in HRU (none)*
- real *8 **pndpcp**
 - precipitation on pond during day (m³ H₂O)*
- real *8 **wetpcp**
- real *8 **wetsep**
 - seepage from wetland bottom for day (m³ H₂O)*
- real *8 **pndev**
 - evaporation from pond on day (m³ H₂O)*
- real *8 **pndflwi**

- volume of water flowing into pond on day ($m^3 H_2O$)*

 - real *8 **pndsedo**
sediment leaving pond during day (metric tons)
 - real *8 **pndsep**
seepage from pond on day ($m^3 H_2O$)
 - real *8 **wetev**
evaporation from wetland for day ($m^3 H_2O$)
 - real *8 **wetflwi**
volume of water flowing in wetland on day ($m^3 H_2O$)
 - real *8 **wetsedo**
sediment loading from wetland for day (metric tons)
 - real *8 **da_ha**
drainage area of watershed in hectares (ha)
 - real *8 **pndflwo**
volume of water flowing out of pond on day ($m^3 H_2O$)
 - real *8 **vpd**
vapor pressure deficit (kPa)
 - real *8 **wetflwo**
volume of water flowing out wetland on day ($m^3 H_2O$)
 - real *8 **wetsedi**
sediment loading to wetland for day (metric tons)
 - real *8 **evlai**
leaf area index at which no evaporation occurs. This variable is used in ponded HRUs where evaporation from the water surface is restricted by the plant canopy cover. Evaporation from the water surface equals potential ET when LAI = 0 and decreased linearly to 0 when LAI = EVLAI
 - real *8 **evrch**
Reach evaporation adjustment factor. Evaporation from the reach is multiplied by EVRCH. This variable was created to limit the evaporation predicted in arid regions.
 - real *8 **wdlpf**
die-off factor for less persistent bacteria on foliage (1/day)
 - real *8 **ep_day**
actual amount of transpiration that occurs on day in HRU (mm H₂O)
 - real *8 **pet_day**
potential evapotranspiration on current day in HRU (mm H₂O)
 - real *8 **bactrolp**
less persistent bacteria transported to main channel with surface runoff (# colonies/ha)
 - real *8 **bactsedlp**
less persistent bacteria transported with sediment in surface runoff (# colonies/ha)
 - real *8 **adj_pkr**
peak rate adjustment factor in the subbasin. Used in the MUSLE equation to account for impact of peak flow on erosion (none)
 - real *8 **n_updis**
nitrogen uptake distribution parameter. This parameter controls the amount of nitrogen removed from the different soil layer layers by the plant. In particular, this parameter allows the amount of nitrogen removed from the surface layer via plant uptake to be controlled. While the relationship between UBN and N removed from the surface layer is affected by the depth of the soil profile, in general, as UBN increases the amount of N removed from the surface layer relative to the amount removed from the entire profile increases
 - real *8 **nactfr**
nitrogen active pool fraction. The fraction of organic nitrogen in the active pool (none)
 - real *8 **p_updis**

phosphorus uptake distribution parameter This parameter controls the amount of phosphorus removed from the different soil layers by the plant. In particular, this parameter allows the amount of phosphorus removed from the surface layer via plant uptake to be controlled. While the relationship between UBP and P uptake from the surface layer is affected by the depth of the soil profile, in general, as UBP increases the amount of P removed from the surface layer relative to the amount removed from the entire profile increases

- real *8 **snoev**
amount of water in snow lost through sublimation on current day in HRU (mm H2O)
- real *8 **sno3up**
- real *8 **reactw**
- real *8 **es_day**
actual amount of evaporation (soil et) that occurs on day in HRU (mm H2O)
- real *8 **sdiégropq**
- real *8 **sdiégrolpq**
- real *8 **sdiégrops**
- real *8 **sdiégrolps**
- real *8 **wof_lp**
wash off fraction for less persistent bacteria on foliage during a rainfall event
- real *8 **ep_max**
maximum amount of transpiration (plant et) that can occur on day in HRU (mm H2O)
- real *8 **sbactrop**
- real *8 **sbactrolp**
- real *8 **sbactsedp**
- real *8 **sbactsedlp**
- real *8 **sbactlchp**
- real *8 **sbactlchlp**
- real *8 **psp_bsn**
- real *8 **rchwtr**
- real *8 **resuspst**
- real *8 **setlpst**
- real *8 **bsprev**
surface runoff lagged from prior day of simulation (mm H2O)
- real *8 **bssprev**
lateral flow lagged from prior day of simulation (mm H2O)
- real *8 **spadyo**
- real *8 **spadyev**
- real *8 **spadysp**
- real *8 **spadyrfv**
- real *8 **spadyosp**
- real *8 **qday**
amount of surface runoff loading to main channel from HRU on current day (mm H2O)
- real *8 **al5**
fraction of total rainfall that occurs during 0.5h of highest intensity rain (none)
- real *8 **no3pcp**
nitrate added to the soil in rainfall (kg N/ha)
- real *8 **pndsedc**
net change in sediment in pond during day (metric tons)
- real *8 **usle_ei**
USLE rainfall erosion index on day for HRU (100(ft·tn in)/(acre·hr))
- real *8 **rcharea**
- real *8 **volatpst**
- real *8 **ubw**

water uptake distribution parameter. This parameter controls the amount of water removed from the different soil layers by the plant. In particular, this parameter allows the amount of water removed from the surface layer via plant uptake to be controlled. While the relationship between UBW and H2O removed from the surface layer is affected by the depth of the soil profile, in general, as UBW increases the amount of water removed from the surface layer relative to the amount removed from the entire profile increases

- real *8 **uobn**
nitrogen uptake normalization parameter. This variable normalizes the nitrogen uptake so that the model can easily verify that uptake from the different soil layers sums to 1.0
- real *8 **uobp**
phosphorus uptake normalization parameter. This variable normalizes the phosphorus uptake so that the model can easily verify that uptake from the different soil layers sums to 1.0
- real *8 **uobw**
water uptake normalization parameter. This variable normalizes the water uptake so that the model can easily verify that uptake from the different soil layers sums to 1.0
- real *8 **wglpf**
growth factor for less persistent bacteria on foliage (1/day)
- real *8 **wetsedc**
net change in sediment in wetland during day (metric tons)
- real *8 **respesti**
- real *8 **rcor**
correction coefficient for generated rainfall to ensure that the annual means for generated and observed values are comparable (needed only if IDIST=1)
- real *8 **rexp**
value of exponent for mixed exponential rainfall distribution (needed only if IDIST=1)
- real *8 **snocov1**
1st shape parameter for snow cover equation. This parameter is determined by solving the equation for 50% snow cover
- real *8 **snocov2**
2nd shape parameter for snow cover equation. This parameter is determined by solving the equation for 95% snow cover
- real *8 **snocovmx**
Minimum snow water content that corresponds to 100% snow cover. If the snow water content is less than SNOC \leftrightarrow OV \leftrightarrow MX, then a certain percentage of the ground will be bare (mm H2O)
- real *8 **lyrtile**
drainage tile flow in soil layer for day (mm H2O)
- real *8 **lyrtilex**
- real *8 **sno50cov**
Fraction of SNOCOVMX that corresponds to 50% snow cover. SWAT assumes a nonlinear relationship between snow water and snow cover.
- real *8 **ai0**
ratio of chlorophyll-a to algal biomass (ug chla/mg alg)
- real *8 **ai1**
fraction of algal biomass that is nitrogen (mg N/mg alg)
- real *8 **ai2**
fraction of algal biomass that is phosphorus (mg P/mg alg)
- real *8 **ai3**
the rate of oxygen production per unit of algal photosynthesis (mg O2/mg alg)
- real *8 **ai4**
the rate of oxygen uptake per unit of algae respiration (mg O2/mg alg)
- real *8 **ai5**
the rate of oxygen uptake per unit of NH3 nitrogen oxidation (mg O2/mg N)
- real *8 **ai6**
the rate of oxygen uptake per unit of NO2 nitrogen oxidation (mg O2/mg N)

- real *8 rhoq
algal respiration rate (1/day or 1/hr)
- real *8 tfact
fraction of solar radiation computed in the temperature heat balance that is photosynthetically active
- real *8 k_l
half-saturation coefficient for light (MJ/(m²*hr))
- real *8 k_n
michaelis-menton half-saturation constant for nitrogen (mg N/L)
- real *8 k_p
michaelis-menton half saturation constant for phosphorus (mg P/L)
- real *8 lambda0
non-algal portion of the light extinction coefficient (1/m)
- real *8 lambda1
linear algal self-shading coefficient (1/(m*ug chla/L))
- real *8 lambda2
nonlinear algal self-shading coefficient ((1/m)(ug chla/L)**(-2/3))
- real *8 mumax
maximum specific algal growth rate (1/day or 1/hr)
- real *8 p_n
algal preference factor for ammonia
- real *8 rnum1
variable to hold value for rnum1s(:) (none)
- real *8 etday
actual evapotranspiration occurring on day in HRU (mm H2O)
- real *8 auton
amount of nitrogen applied in auto-fert application (kg N/ha)
- real *8 autop
amount of phosphorus applied in auto-fert application (kg P/ha)
- real *8 hmntl
amount of nitrogen moving from active organic to nitrate pool in soil profile on current day in HRU (kg N/ha)
- real *8 hmptl
amount of phosphorus moving from active organic to nitrate pool in soil profile on current day in HRU (kg P/ha)
- real *8 rmn2tl
amount of nitrogen moving from the fresh organic (residue) to the nitrate(80%) and active organic(20%) pools in soil profile on current day in HRU (kg N/ha)
- real *8 rwnl
amount of nitrogen moving from active organic to stable organic pool in soil profile on current day in HRU (kg N/ha)
- real *8 gwseep
amount of water recharging deep aquifer on current day (mm H2O)
- real *8 revapday
amount of water moving from the shallow aquifer into the soil profile or being taken up by plant roots in the shallow aquifer (mm H2O)
- real *8 rmp1tl
amount of phosphorus moving from the labile mineral pool to the active mineral pool in the soil profile on the current day in the HRU (kg P/ha)
- real *8 rmptl
amount of phosphorus moving from the fresh organic (residue) to the labile(80%) and organic(20%) pools in soil profile on current day in HRU (kg P/ha)
- real *8 roctl
amount of phosphorus moving from the active mineral pool to the stable mineral pool in the soil profile on the current day in the HRU (kg P/ha)
- real *8 wdntl

- amount of nitrogen lost from nitrate pool by denitrification in soil profile on current day in HRU (kg N/ha)*

 - real *8 **cmn_bsn**
 - real *8 **reswtr**
 - real *8 **wdlprch**
- die-off factor for less persistent bacteria in streams (1/day)*

 - real *8 **wdpres**
- die-off factor for persistent bacteria in reservoirs (1/day)*

 - real *8 **petmeas**
- potential ET value read in for day (mm H2O)*

 - real *8 **bury**
 - real *8 **difus**
 - real *8 **reactb**
 - real *8 **solpesto**
 - real *8 **wdlpres**
- die-off factor for less persistent bacteria in reservoirs (1/day)*

 - real *8 **sorpesto**
 - real *8 **spcon_bsn**
 - real *8 **spexp_bsn**
 - real *8 **solpesti**
 - real *8 **sorpesti**
 - real *8 **msk_co1**
- calibration coefficient to control impact of the storage time constant for the reach at bankfull depth (phi(10,:)) upon the storage time constant for the reach used in the Muskingum flow method*

 - real *8 **msk_co2**
- calibration coefficient to control impact of the storage time constant for the reach at 0.1 bankfull depth (phi(13,:)) upon the storage time constant for the reach used in the Muskingum flow method*

 - real *8 **deepstp**
- depth of water in deep aquifer in HRU (mm H2O)*

 - real *8 **shallstp**
- depth of water in shallow aquifer in HRU on previous day (mm H2O)*

 - real *8 **snoprev**
- amount of water stored as snow on previous day (mm H2O)*

 - real *8 **swprev**
- amount of water stored in soil profile in the HRU on the previous day (mm H2O)*

 - real *8 **ressolpo**
 - real *8 **resorgno**
 - real *8 **resorgpo**
 - real *8 **resno3o**
 - real *8 **reschlao**
 - real *8 **resno2o**
 - real *8 **potevmm**
- volume of water evaporated from pothole expressed as depth over HRU (mm H2O)*

 - real *8 **potflwo**
- volume of water released to main channel from pothole expressed as depth over HRU (mm H2O)*

 - real *8 **potpcpmm**
- precipitation falling on pothole water body expressed as depth over HRU (mm H2O)*

 - real *8 **potsepmm**
- seepage from pothole expressed as depth over HRU (mm H2O)*

 - real *8 **resnh3o**
 - real *8 **qdbank**
 - real *8 **bactminlp**
- Threshold detection level for less persistent bacteria. When bacteria levels drop to this amount the model considers bacteria in the soil to be insignificant and sets the levels to zero (cfu/m²)*

- real *8 **bactminp**
Threshold detection level for persistent bacteria. When bacteria levels drop to this amount the model considers bacteria in the soil to be insignificant and sets the levels to zero (cfu/m²)
- real *8 **trnsrch**
fraction of transmission losses from main channel that enter deep aquifer
- real *8 **wp20p_plt**
overall rate change for persistent bacteria on foliage (1/day)
- real *8 **potsedo**
sediment released to main channel from HRU (metric tons/ha)
- real *8 **pest_sol**
- real *8 **bact_swf**
fraction of manure containing active colony forming units (cfu)
- real *8 **bactmx**
bacteria percolation coefficient. Ratio of solution bacteria in surface layer to solution bacteria in percolate
- real *8 **cncoef**
plant ET curve number coefficient
- real *8 **wp20lp_plt**
overall rate change for less persistent bacteria on foliage (1/day)
- real *8 **cdn_bsn**
- real *8 **sdnco_bsn**
- real *8 **bactmin**
- real *8 **cn_froz**
drainage coefficient (mm day⁻¹)
- real *8 **dorm_hr**
time threshold used to define dormant (hours)
- real *8 **smxco**
adjustment factor for max curve number s factor (0-1)
- real *8 **tb_adj**
adjustment factor for subdaily unit hydrograph basetime
- real *8 **chla_subco**
regional adjustment on sub chla_a loading (fraction)
- real *8 **depimp_bsn**
depth to impervious layer. Used to model perched water tables in all HRUs in watershed (mm)
- real *8 **ddrain_bsn**
depth to the sub-surface drain (mm)
- real *8 **tdrain_bsn**
time to drain soil to field capacity (hours)
- real *8 **gdrain_bsn**
- real *8 **rch_san**
- real *8 **rch_sil**
- real *8 **rch_cla**
- real *8 **rch_sag**
- real *8 **rch_lag**
- real *8 **rch_gra**
- real *8 **hlife_ngw_bsn**
Half-life of nitrogen in groundwater? (days)
- real *8 **ch_opco_bsn**
- real *8 **ch_onco_bsn**
- real *8 **decr_min**
Minimum daily residue decay.
- real *8 **rcn_sub_bsn**
Concentration of nitrogen in the rainfall (mg/kg)

- real *8 **bc1_bsn**
- real *8 **bc2_bsn**
- real *8 **bc3_bsn**
- real *8 **bc4_bsn**
- real *8 **anion_excl_bsn**
- real *8, dimension(:), allocatable **wat_tbl**
- real *8, dimension(:), allocatable **sol_swpwt**
- real *8, dimension(:,:), allocatable **vwt**
- real *8 **re_bsn**
Effective radius of drains (range 3.0 - 40.0) (mm)
- real *8 **sdrain_bsn**
Distance bewtween two drain or tile tubes (range 7600.0 - 30000.0) (mm)
- real *8 **sstmaxd_bsn**
- real *8 **drain_co_bsn**
Drainage coefficient (range 10.0 - 51.0) (mm-day-1)
- real *8 **latksatf_bsn**
Multiplication factor to determine lateral ksat from SWAT ksat input value for HRU (range 0.01 - 4.0)
- real *8 **pc_bsn**
Pump capacity (def val = 1.042 mm h-1 or 25 mm day-1) (mm h-1)
- integer **idlast**
number of days simulated in month (none)
- integer **i_subhw**
- integer **imgt**
- integer **iwtr**
- integer **ifrttyp**
- integer **mo_atmo**
- integer **mo_atmo1**
- integer **ifirstatmo**
- integer **iy_r_atmo**
- integer **iy_r_atmo1**
- integer **matmo**
- integer **mch**
maximum number of channels
- integer **mcr**
maximum number of crops grown per year
- integer **mcrdb**
maximum number of crops/landcover in database file (crop.dat)
- integer **mfcst**
maximum number of forecast stations
- integer **mfdb**
maximum number of fertilizers in fert.dat
- integer **mhru**
maximum number of HRUs in watershed
- integer **mhyd**
maximum number of hydrograph nodes
- integer **mpdb**
maximum number of pesticides in pest.dat
- integer **mrq**
maximum number of rainfall/temp gages (none)
- integer **mcut**
maximum number of cuttings per year
- integer **mqr**

- maximum number of grazings per year*
- integer **mnr**
 - maximum number of years of rotation*
- integer **myr**
 - maximum number of years of simulation*
- integer **isubwq**
 - subbasin water quality code*
 - 0 do not calculate algae/CBOD 1 calculate algae/CBOD drainmod tile equations*
- integer **ffcst**
- integer **isproj**
 - special project code (none):*
 - 1 test rewind (run simulation twice)*
- integer **nbyr**
 - number of calendar years simulated (none)*
- integer **irte**
 - water routing method (none):*
 - 0 variable storage method*
 - 1 Muskingum method*
- integer **nrch**
 - number of reaches in watershed (none)*
- integer **nres**
 - total number of reservoirs in watershed (none)*
- integer **nhru**
 - number of last HRU in previous subbasin or*
 - number of HRUs in watershed (none)*
- integer **i_mo**
 - current month being simulated or month of next day of simulation (none)*
- integer **immo**
 - current cumulative month of simulation (none)*
- integer **mo**
- integer **wndsim**
 - wind speed input code (noen)*
 - 1 measured data read for each subbasin*
 - 2 data simulated for each subbasin*
- integer **ihru**
 - HRU number (none)*
- integer **icode**
 - variable to hold value for icode(:) (none)*
- integer **ihout**
 - variable to hold value for ihouts(:) (none)*
- integer **inum1**
 - variable to hold value for inum1s(:) (subbasin number) (none)*
- integer **inum2**
 - variable to hold value for inum2s(:) (none)*
- integer **inum3**
 - variable to hold value for inum3s(:) (none)*
- integer **inum4**
 - variable to hold value for inum4s(:) (none)*
- integer **icfac**
 - icfac = 0 for C-factor calculation using Cmin (as described in manual)*
 - = 1 for new C-factor calculation from RUSLE (no minimum needed)*
- integer **inum5**

- integer **inum6**
- integer **inum7**
- integer **inum8**
- integer **mrech**
maximum number of rehour files
- integer **nrgage**
number of raingage files (none)
- integer **nrgfil**
number of rain gages per file (none)
- integer **nrtot**
total number of rain gages (none)
- integer **ntgage**
number of temperature gage files (none)
- integer **ntgfil**
number of temperature gages per file (none)
- integer **nttot**
total number of temperature gages (none)
- integer **tmpsim**
temperature input code (none)
1 measured data read for each subbasin
2 data simulated for each subbasin
- integer **icrk**
crack flow code
1: simulate crack flow in watershed
- integer **irtpest**
number of pesticide to be routed through the watershed. Redefined to the sequence number of pesticide in NPNO(:) which is to be routed through the watershed (none)
- integer **igropt**
Qual2E option for calculating the local specific growth rate of algae
1: multiplicative.
- integer **lao**
Qual2E light averaging option. Qual2E defines four light averaging options. The only option currently available in SWAT is #2.
- integer **npmx**
number of different pesticides used in the simulation (none)
- integer **curyr**
current year in simulation (sequence) (none)
- integer **itdrn**
tile drainage equations flag/code
1 simulate tile flow using subroutine drains(wt_shall)
0 simulate tile flow using subroutine origtile(wt_shall,d)
- integer **iwt dn**
water table depth algorithms flag/code
1 simulate wt_shall using subroutine new water table depth routine
0 simulate wt_shall using subroutine original water table depth routine
- integer **ismax**
maximum depressional storage selection flag/code
0 = static depressional storage
1 = dynamic storage based on tillage and cumulative rainfall
- integer **iroutunit**
not being implemented in this version drainmod tile equations
- integer **ires_nut**
- integer **iclb**

- auto-calibration flag*
- integer **mrecc**
 - maximum number of reccnst files*
- integer **mrecd**
 - maximum number of recday files*
- integer **mrecm**
 - maximum number of recmon files*
- integer **mtil**
 - max number of tillage types in till.dat*
- integer **mudb**
 - maximum number of urban land types in urban.dat*
- integer **idist**
 - rainfall distribution code*
 - 0 for skewed normal dist*
 - 1 for mixed exponential distribution*
- integer **mrecy**
 - maximum number of recyear files*
- integer **nyskip**
 - number of years to skip output summarization and printing (none)*
- integer **slrsim**
 - solar radiation input code (none)*
 - 1 measured data read for each subbasin*
 - 2 data simulated for each subbasin*
- integer **ideg**
 - channel degradation code*
 - 0: do not compute channel degradation*
 - 1: compute channel degradation (downcutting and widening)*
- integer **ievent**
 - rainfall/runoff code (none)*
 - 0 daily rainfall/curve number technique 1 sub-daily rainfall/Green&Ampt/hourly routing 3 sub-daily rainfall/↵ Green&Ampt/hourly routing*
- integer **ipet**
 - code for potential ET method (none)*
 - 0 Priestley-Taylor method*
 - 1 Penman/Monteith method*
 - 2 Hargreaves method*
 - 3 read in daily potential ET data*
- integer **iopera**
- integer **idaf**
 - beginning day of simulation (julian date)*
- integer **idal**
 - ending day of simulation (julian date)*
- integer **rhsim**
 - relative humidity input code (none)*
 - 1 measured data read for each subbasin*
 - 2 data simulated for each subbasin*
- integer **leapyr**
 - leap year flag (none)*
 - 0 leap year*
 - 1 regular year*
- integer **id1**
 - first day of simulation in current year (julian date)*
- integer **mo_chk**
 - current month of simulation (none)*

- integer **nhtot**
total number of relative humidity records in file
- integer **nstot**
total number of solar radiation records in file (none)
- integer **nwtot**
total number of wind speed records in file
- integer **ifirsts**
solar radiation data search code (none)
0 first day of solar radiation data located in file
1 first day of solar radiation data not located in file
- integer **ifirsth**
relative humidity data search code (none)
0 first day of relative humidity data located in file
1 first day of relative humidity data not located in file
- integer **firstw**
wind speed data search code (none)
0 first day of wind speed data located in file
1 first day of wind speed data not located in file
- integer **icst**
- integer **ilog**
streamflow print code (none)
0 print streamflow in reach
1 print Log10 streamflow in reach
- integer **itotr**
number of output variables printed (output.rch)
- integer **iy**
current year of simulation (year)
- integer **iwq**
stream water quality code
0 do not model stream water quality
1 model stream water quality (QUAL2E & pesticide transformations)
- integer **iskip**
flag for calculations performed only for the first year of simulation (none)
- integer **firstpet**
potential ET data search code (none)
0 first day of potential ET data located in file
1 first day of potential ET data not located in file
- integer **iprp**
print code for output.pst file
0 do not print pesticide output
1 print pesticide output
- integer **itotb**
number of output variables printed (output.sub)
- integer **itots**
number of output variables printed (output.hru)
- integer **itoh**
number of HRUs printed (output.hru/output.wtr)
- integer **pcpsim**
rainfall input code (none)
1 measured data read for each subbasin
2 data simulated for each subbasin
- integer **nd_30**
- integer **iops**
- integer **iphr**

- integer **isto**
- integer **isol**
- integer **fcstcycles**
number of times forecast period is simulated (using different weather generator seeds each time)
- integer **fcstday**
beginning date of forecast period (julian date)
- integer **fcstyr**
beginning year of forecast period
- integer **iscen**
scenarios counter
- integer **subtot**
number of subbasins in watershed (none)
- integer **ogen**
- integer **mapp**
maximum number of applications
- integer **mlyr**
maximum number of soil layers
- integer **mpst**
max number of pesticides used in wshed
- integer **mres**
maximum number of reservoirs
- integer **msub**
maximum number of subbasins
- integer **igen**
random number generator seed code (none):
0: use default numbers
1: generate new numbers in every simulation
- integer **iprint**
print code (none): 0=monthly, 1=daily, 2=annually
- integer **iida**
day being simulated (current julian date) (julian date)
- integer **icn**
CN method flag (for testing alternative method):
0 use traditional SWAT method which bases CN on soil moisture
1 use alternative method which bases CN on plant ET
2 use traditional SWAT method which bases CN on soil moisture but rention is adjusted for mildly-sloped tiled-drained watersheds.
- integer **ised_det**
max half-hour rainfall fraction calc option:
0 generate max half-hour rainfall fraction from triangular distribution
1 use monthly mean max half-hour rainfall fraction
- integer **fcstcnt**
- integer **mtran**
- integer **idtil**
- integer, dimension(100) **ida_lup**
- integer, dimension(100) **iy_r_lup**
- integer **no_lup**
- integer **no_up**
- integer **nostep**
- character(len=8) **date**
date simulation is performed where leftmost eight characters are set to a value of yyyyymmdd, where yyyy is the year, mm is the month and dd is the day
- character(len=10) **time**

- time simulation is performed where leftmost ten characters are set to a value of hhmmss.sss, where hh is the hour, mm is the minutes and ss.sss is the seconds and milliseconds*
- character(len=5) **zone**
 - time difference with respect to Coordinated Universal Time (ie Greenwich Mean Time)*
- character(len=13) **calfile**
 - name of file containing calibration parameters*
- character(len=13) **rhfile**
 - relative humidity file name (.hmd)*
- character(len=13) **slrfile**
 - solar radiation file name (.slr)*
- character(len=13) **wndfile**
 - wind speed file name (.wnd)*
- character(len=13) **petfile**
 - potential ET file name (.pet)*
- character(len=13) **atmofile**
- character(len=13) **lucfile**
- character(len=13) **septdb**
 - name of septic tank database file (septwq1.dat)*
- character(len=13) **dpd_file**
- character(len=13) **wpd_file**
- character(len=13) **rib_file**
- character(len=13) **sfb_file**
- character(len=13) **lid_file**
- integer, dimension(9) **idg**
 - array location of random number seed used for a given process*
- integer, dimension(:), allocatable **ifistr**
- integer, dimension(:), allocatable **ifisthr**
- integer, dimension(8) **values**
 - values(1): year simulation is performed*
 - values(2): month simulation is performed*
 - values(3): day in month simulation is performed*
 - values(4): time difference with respect to Coordinated Universal Time (ie Greenwich Mean Time)*
 - values(5): hour simulation is performed*
 - values(6): minute simulation is performed*
 - values(7): second simulation is performed*
 - values(8): millisecond simulation is performed*
- integer, dimension(13) **ndays**
 - julian date for last day of preceding month (where the array location is the number of the month). The dates are for leap years (julian date)*
- integer **mapex**
- real *8, dimension(:), allocatable **flodaya**
- real *8, dimension(:), allocatable **seddaya**
- real *8, dimension(:), allocatable **orgndaya**
- real *8, dimension(:), allocatable **orgpdaya**
- real *8, dimension(:), allocatable **no3daya**
- real *8, dimension(:), allocatable **minpdaya**
- real *8, dimension(:), allocatable **hi_targ**
 - harvest index target of cover defined at planting ((kg/ha)/(kg/ha))*
- real *8, dimension(:), allocatable **bio_targ**
 - biomass target (kg/ha)*
- real *8, dimension(:), allocatable **tnyld**
 - modifier for autofertilization target nitrogen content for plant (kg N/kg yield)*
- integer, dimension(:), allocatable **idapa**
- integer, dimension(:), allocatable **iypa**

- integer, dimension(:), allocatable **ifirsta**
- integer, dimension(100) **mo_transb**
- integer, dimension(100) **mo_transe**
- integer, dimension(100) **ih_tran**
- integer **msdb**
maximum number of sept wq data database (none)
- integer **iseptic**
- real *8, dimension(:), allocatable **sptqs**
flow rate of the septic tank effluent per capita (m3/d)
- real *8, dimension(:), allocatable **percp**
- real *8, dimension(:), allocatable **sptbodconcs**
Biological Oxygen Demand of the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **sptssconcs**
concentration of total suspended solid in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **spttnconcs**
concentration of total nitrogen in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **sptnh4concs**
concentration of total phosphorus of the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **sptno3concs**
concentration of nitrate in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **sptno2concs**
concentration of nitrite in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **sptorgnconcs**
concentration of organic nitrogen in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **spttpconcs**
concentration of total phosphorus in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **sptminps**
concentration of mineral phosphorus in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **sptorgps**
concentration of organic phosphorus in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **sptfcolis**
concentration of the facel caliform in the septic tank effluent (cfu/100ml)
- real *8, dimension(:), allocatable **failyr**
- real *8, dimension(:), allocatable **qstemm**
- real *8, dimension(:), allocatable **bio_amn**
- real *8, dimension(:), allocatable **bio_bod**
- real *8, dimension(:), allocatable **biom**
- real *8, dimension(:), allocatable **rbiom**
- real *8, dimension(:), allocatable **fcoli**
- real *8, dimension(:), allocatable **bio_ntr**
- real *8, dimension(:), allocatable **bz_perc**
- real *8, dimension(:), allocatable **sep_cap**
number of permanent residents in the hourse (none)
- real *8, dimension(:), allocatable **plqm**
- real *8, dimension(:), allocatable **bz_area**
- real *8, dimension(:), allocatable **bz_z**
Depth of biozone layer(mm)
- real *8, dimension(:), allocatable **bz_thk**
thickness of biozone (mm)
- real *8, dimension(:), allocatable **bio_bd**
density of biomass (kg/m³) carbon outputs for .hru file
- real *8, dimension(:), allocatable **cmup_kgh**

- real *8, dimension(:), allocatable **cmtot_kgh**
- real *8, dimension(:), allocatable **coeff_denitr**
denitrification rate coefficient (none)
- real *8, dimension(:), allocatable **coeff_bod_dc**
BOD decay rate coefficient (m^3/day)
- real *8, dimension(:), allocatable **coeff_bod_conv**
BOD to live bacteria biomass conversion factor (none)
- real *8, dimension(:), allocatable **coeff_fc1**
field capacity calibration parameter 1 (none)
- real *8, dimension(:), allocatable **coeff_fc2**
field capacity calibration parameter 2 (none)
- real *8, dimension(:), allocatable **coeff_fecal**
fecal coliform bacteria decay rate coefficient (m^3/day)
- real *8, dimension(:), allocatable **coeff_mrt**
mortality rate coefficient (none)
- real *8, dimension(:), allocatable **coeff_nitr**
nitrification rate coefficient (none)
- real *8, dimension(:), allocatable **coeff_plq**
conversion factor for plaque from TDS (none)
- real *8, dimension(:), allocatable **coeff_rsp**
respiration rate coefficient (none)
- real *8, dimension(:), allocatable **coeff_slg1**
slough-off calibration parameter (none)
- real *8, dimension(:), allocatable **coeff_slg2**
slough-off calibration parameter (none)
- real *8, dimension(:), allocatable **coeff_pdistrib**
- real *8, dimension(:), allocatable **coeff_solpslp**
- real *8, dimension(:), allocatable **coeff_solpintc**
- real *8, dimension(:), allocatable **coeff_psortpmax**
- integer, dimension(:), allocatable **isep_typ**
septic system type (none)
- integer, dimension(:), allocatable **i_sep**
- integer, dimension(:), allocatable **isep_opt**
septic system operation flag (1=active, 2=failing, 3=not operated) (none)
- integer, dimension(:), allocatable **sep_tsincefail**
- integer, dimension(:), allocatable **isep_tfail**
- integer, dimension(:), allocatable **isep_iyr**
- integer, dimension(:), allocatable **sep_strm_dist**
- integer, dimension(:), allocatable **sep_den**
- real *8, dimension(:), allocatable **sol_sumno3**
- real *8, dimension(:), allocatable **sol_sumsolp**
- real *8, dimension(:), allocatable **strsw_sum**
- real *8, dimension(:), allocatable **strstmp_sum**
- real *8, dimension(:), allocatable **strsn_sum**
- real *8, dimension(:), allocatable **strsp_sum**
- real *8, dimension(:), allocatable **strsa_sum**
- real *8, dimension(:), allocatable **spill_hru**
- real *8, dimension(:), allocatable **tile_out**
- real *8, dimension(:), allocatable **hru_in**
- real *8, dimension(:), allocatable **spill_precip**
- real *8, dimension(:), allocatable **pot_seep**
- real *8, dimension(:), allocatable **pot_evap**

- real *8, dimension(:), allocatable **pot_sedin**
- real *8, dimension(:), allocatable **pot_solp**
soluble P loss rate in the pothole (.01 - 0.5) (1/d)
- real *8, dimension(:), allocatable **pot_solpi**
- real *8, dimension(:), allocatable **pot_orgp**
- real *8, dimension(:), allocatable **pot_orgpi**
- real *8, dimension(:), allocatable **pot_orgn**
- real *8, dimension(:), allocatable **pot_orgni**
- real *8, dimension(:), allocatable **pot_mps**
- real *8, dimension(:), allocatable **pot_mpsi**
- real *8, dimension(:), allocatable **pot_mpa**
- real *8, dimension(:), allocatable **pot_mpai**
- real *8, dimension(:), allocatable **pot_no3i**
- real *8, dimension(:), allocatable **precip_in**
- real *8, dimension(:), allocatable **tile_sedo**
- real *8, dimension(:), allocatable **tile_no3o**
- real *8, dimension(:), allocatable **tile_solpo**
- real *8, dimension(:), allocatable **tile_orgno**
- real *8, dimension(:), allocatable **tile_orgpo**
- real *8, dimension(:), allocatable **tile_minpso**
- real *8, dimension(:), allocatable **tile_minpao**
- integer **ia_b**
- integer **ihumus**
- integer **itemp**
- integer **isnow**
- integer, dimension(46) **ipdvar**
output variable codes for output.rch file (none)
- integer, dimension(**mhruo**) **ipdvas**
output variable codes for output.hru file (none)
- integer, dimension(**msubo**) **ipdvab**
output variable codes for output.sub file (none)
- integer, dimension(:), allocatable **ipdhru**
HRUs whose output information will be printed to the output.hru and output.wtr files.
- real *8, dimension(**mstdo**) **wshddayo**
wshddayo(1) average amount of precipitation in watershed for the day (mm H2O)
wshddayo(3) surface runoff in watershed for day (mm H2O)
wshddayo(4) lateral flow contribution to streamflow in watershed for day (mm H2O)
wshddayo(5) water percolation past bottom of soil profile in watershed for day (mm H2O)
wshddayo(6) water yield to streamflow from HRUs in watershed for day (mm H2O)
wshddayo(7) actual evapotranspiration in watershed for day (mm H2O)
wshddayo(12) sediment yield from HRUs in watershed for day (metric tons or metric tons/ha)
wshddayo(35) amount of water stored in soil profile in watershed at end of day (mm H2O)
wshddayo(41) organic P loading to stream in watershed for day (kg P/ha)
wshddayo(42) nitrate loading to stream in surface runoff in watershed for day (kg N/ha)
wshddayo(43) soluble P loading to stream in watershed for day (kg P/ha)
wshddayo(44) plant uptake of N in watershed for day (kg N/ha)
wshddayo(45) nitrate loading to stream in lateral flow in watershed for day (kg N/ha)
wshddayo(46) nitrate percolation past bottom of soil profile in watershed for day (kg N/ha)
wshddayo(104) groundwater contribution to stream in watershed on day (mm H2O)
wshddayo(108) potential evapotranspiration in watershed on day (mm H2O)
wshddayo(109) drainage tile flow contribution to stream in watershed on day (mm H2O)
- real *8, dimension(**mstdo**) **wshdmono**

watershed monthly output array (see definitions for `wshddayo` array elements) (varies)
`wshdmono(1)` average amount of precipitation in watershed for the month (mm H₂O)
`wshdmono(3)` surface runoff in watershed for month (mm H₂O)
`wshdmono(4)` lateral flow contribution to streamflow in watershed for month (mm H₂O)
`wshdmono(5)` water percolation past bottom of soil profile in watershed for month (mm H₂O)
`wshdmono(6)` water yield to streamflow from HRUs in watershed for month (mm H₂O)
`wshdmono(7)` actual evapotranspiration in watershed for month (mm H₂O)
`wshdmono(8)` average maximum temperature in watershed for the month (deg C)
`wshdmono(9)` average minimum temperature in watershed for the month (deg C)
`wshdmono(12)` sediment yield from HRUs in watershed for the month (metric tons)
`wshdmono(39)` freezing rain/snow fall in watershed for the month (mm H₂O)
`wshdmono(40)` organic N loading to stream in watershed for the month (kg N/ha)
`wshdmono(41)` organic P loading to stream in watershed for the month (kg P/ha)
`wshdmono(42)` nitrate loading to stream in surface runoff in watershed for the month (kg N/ha)
`wshdmono(43)` soluble P loading to stream in watershed for the month (kg P/ha)
`wshdmono(44)` plant uptake of N in watershed for the month (kg N/ha)
`wshdmono(45)` nitrate loading to stream in lateral flow in watershed for the month (kg N/ha)
`wshdmono(46)` nitrate percolation past bottom of soil profile in watershed for the month (kg N/ha)
`wshdmono(104)` groundwater contribution to stream in watershed for the month (mm H₂O)
`wshdmono(108)` potential evapotranspiration in watershed for the month (mm H₂O)
`wshdmono(109)` drainage tile flow contribution to stream in watershed for the month (mm H₂O)

- real *8, dimension(`mstdo`) `wshdyro`

watershed annual output array (varies)
`wshdyro(1)` average amount of precipitation in watershed for the year (mm H₂O)
`wshdyro(3)` surface runoff in watershed for year (mm H₂O)
`wshdyro(4)` lateral flow contribution to streamflow in watershed for year (mm H₂O)
`wshdyro(5)` water percolation past bottom of soil profile in watershed for year (mm H₂O)
`wshdyro(6)` water yield to streamflow from HRUs in watershed for year (mm H₂O)
`wshdyro(7)` actual evapotranspiration in watershed for year (mm H₂O)
`wshdyro(8)` average maximum temperature in watershed for the year (deg C)
`wshdyro(9)` average minimum temperature in watershed for the year (deg C)
`wshdyro(12)` sediment yield from HRUs in watershed for the year (metric tons)
`wshdyro(40)` organic N loading to stream in watershed for the year (kg N/ha)
`wshdyro(41)` organic P loading to stream in watershed for the year (kg P/ha)
`wshdyro(42)` nitrate loading to stream in surface runoff in watershed for the year (kg N/ha)
`wshdyro(43)` soluble P loading to stream in watershed for the year (kg P/ha)
`wshdyro(44)` plant uptake of N in watershed for the year
`wshdyro(45)` nitrate loading to stream in lateral flow in watershed for the year (kg N/ha)
`wshdyro(46)` nitrate percolation past bottom of soil profile in watershed for the year (kg N/ha)
`wshdyro(104)` groundwater contribution to stream in watershed for the year (mm H₂O)
`wshdyro(108)` potential evapotranspiration in watershed for the year (mm H₂O)
`wshdyro(109)` drainage tile flow contribution to stream in watershed for the year (mm H₂O)

- real *8, dimension(16) `fcstaao`

- real *8, dimension(`mstdo`) `wshdaao`

watershed average annual output array (varies)
`wshdaao(1)` precipitation in watershed (mm H₂O)
`wshdaao(3)` surface runoff loading to main channel in watershed (mm H₂O)
`wshdaao(4)` lateral flow loading to main channel in watershed (mm H₂O)
`wshdaao(5)` percolation of water out of root zone in watershed (mm H₂O)
`wshdaao(7)` actual evapotranspiration in watershed (mm H₂O)
`wshdaao(13)` sediment loading to ponds in watershed (metric tons)
`wshdaao(14)` sediment loading from ponds in watershed (metric tons)
`wshdaao(15)` net change in sediment level in ponds in watershed (metric tons)
`wshdaao(19)` evaporation from ponds in watershed (m³ H₂O)
`wshdaao(20)` seepage from ponds in watershed (m³ H₂O)
`wshdaao(21)` precipitation on ponds in watershed (m³ H₂O)
`wshdaao(22)` volume of water entering ponds in watershed (m³ H₂O)
`wshdaao(23)` volume of water leaving ponds in watershed (m³ H₂O)
`wshdaao(38)` transmission losses in watershed (mm H₂O)

- real *8, dimension(:,.), allocatable `wpstdayo`
- real *8, dimension(:,.), allocatable `wpstmono`
- real *8, dimension(:,.), allocatable `wpstyro`

- real *8, dimension(:,:), allocatable [bio_hv](#)
harvested biomass (dry weight) (kg/ha)
- real *8, dimension(:,:), allocatable [yldkg](#)
yield (dry weight) by crop type in the HRU (kg/ha)
- real *8, dimension(:,:), allocatable [rchmono](#)
reach monthly output array (varies)
rchmono(1,:) flow into reach during month (m^3/s)
rchmono(2,:) flow out of reach during month (m^3/s)
rchmono(3,:) sediment transported into reach during month (metric tons)
rchmono(4,:) sediment transported out of reach during month (metric tons)
rchmono(5,:) sediment concentration in outflow during month (mg/L)
rchmono(6,:) organic N transported into reach during month (kg N)
rchmono(7,:) organic N transported out of reach during month (kg N)
rchmono(8,:) organic P transported into reach during month (kg P)
rchmono(9,:) organic P transported out of reach during month (kg P)
rchmono(10,:) evaporation from reach during month (m^3/s)
rchmono(11,:) transmission losses from reach during month (m^3/s)
rchmono(12,:) conservative metal #1 transported out of reach during month (kg)
rchmono(13,:) conservative metal #2 transported out of reach during month (kg)
rchmono(14,:) conservative metal #3 transported out of reach during month (kg)
rchmono(15,:) nitrate transported into reach during month (kg N)
rchmono(16,:) nitrate transported out of reach during month (kg N)
rchmono(17,:) soluble P transported into reach during month (kg P)
rchmono(18,:) soluble P transported out of reach during month (kg P)
rchmono(19,:) soluble pesticide transported into reach during month (mg pst)
rchmono(20,:) soluble pesticide transported out of reach during month (mg pst)
rchmono(21,:) sorbed pesticide transported into reach during month (mg pst)
rchmono(22,:) sorbed pesticide transported out of reach during month (mg pst)
rchmono(23,:) amount of pesticide lost through reactions in reach during month (mg pst)
rchmono(24,:) amount of pesticide lost through volatilization from reach during month (mg pst)
rchmono(25,:) amount of pesticide settling out of reach to bed sediment during month (mg pst)
rchmono(26,:) amount of pesticide resuspended from bed sediment to reach during month (mg pst)
rchmono(27,:) amount of pesticide diffusing from reach to bed sediment during month (mg pst)
rchmono(28,:) amount of pesticide in sediment layer lost through reactions during month (mg pst)
rchmono(29,:) amount of pesticide in sediment layer lost through burial during month (mg pst)
rchmono(30,:) chlorophyll-a transported into reach during month (kg chla)
rchmono(31,:) chlorophyll-a transported out of reach during month (kg chla)
rchmono(32,:) ammonia transported into reach during month (kg N)
rchmono(33,:) ammonia transported out of reach during month (kg N)
rchmono(34,:) nitrite transported into reach during month (kg N)
rchmono(35,:) nitrite transported out of reach during month (kg N)
rchmono(36,:) CBOD transported into reach during month (kg O₂)
rchmono(37,:) CBOD transported out of reach during month (kg O₂)
rchmono(38,:) dissolved oxygen transported into reach during month (kg O₂)
rchmono(39,:) dissolved oxygen transported out of reach during month (kg O₂)
rchmono(40,:) persistent bacteria transported out of reach during month (kg bact)
rchmono(41,:) less persistent bacteria transported out of reach during month (kg bact)
rchmono(43,:) total N (org N + no3 + no2 + nh4 outs) (kg)
rchmono(44,:) total P (org P + sol p outs) (kg)
- real *8, dimension(:,:), allocatable [rchyro](#)
reach annual output array (varies)
rchyro(1,:) flow into reach during year (m^3/s)
rchyro(2,:) flow out of reach during year (m^3/s)
rchyro(3,:) sediment transported into reach during year (metric tons)
rchyro(4,:) sediment transported out of reach during year (metric tons)
rchyro(5,:) sediment concentration in outflow during year (mg/L)
rchyro(6,:) organic N transported into reach during year (kg N)
rchyro(7,:) organic N transported out of reach during year (kg N)
rchyro(8,:) organic P transported into reach during year (kg P)
rchyro(9,:) organic P transported out of reach during year (kg P)
rchyro(10,:) evaporation from reach during year (m^3/s)
rchyro(11,:) transmission losses from reach during year (m^3/s)

rchyro(12,:) conservative metal #1 transported out of reach during year (kg)
rchyro(13,:) conservative metal #2 transported out of reach during year (kg)
rchyro(14,:) conservative metal #3 transported out of reach during year (kg)
rchyro(15,:) nitrate transported into reach during year (kg N)
rchyro(16,:) nitrate transported out of reach during year (kg N)
rchyro(17,:) soluble P transported into reach during year (kg P)
rchyro(18,:) soluble P transported out of reach during year (kg P)
rchyro(19,:) soluble pesticide transported into reach during year (mg pst)
rchyro(20,:) soluble pesticide transported out of reach during year (mg pst)
rchyro(21,:) sorbed pesticide transported into reach during year (mg pst)
rchyro(22,:) sorbed pesticide transported out of reach during year (mg pst)
rchyro(23,:) amount of pesticide lost through reactions in reach during year!> (mg pst)
rchyro(24,:) amount of pesticide lost through volatilization from reach during year (mg pst)
rchyro(25,:) amount of pesticide settling out of reach to bed sediment during year (mg pst)
rchyro(26,:) amount of pesticide resuspended from bed sediment to reach during year (mg pst)
rchyro(27,:) amount of pesticide diffusing from reach to bed sediment during year (mg pst)
rchyro(28,:) amount of pesticide in sediment layer lost through reactions during year (mg pst)
rchyro(29,:) amount of pesticide in sediment layer lost through burial during year (mg pst)
rchyro(30,:) chlorophyll-a transported into reach during year (kg chla)
rchyro(31,:) chlorophyll-a transported out of reach during year (kg chla)
rchyro(32,:) ammonia transported into reach during year (kg N)
rchyro(33,:) ammonia transported out of reach during year (kg N)
rchyro(34,:) nitrite transported into reach during year (kg N)
rchyro(35,:) nitrite transported out of reach during year (kg N)
rchyro(36,:) CBOD transported into reach during year (kg O2)
rchyro(37,:) CBOD transported out of reach during year (kg O2)
rchyro(38,:) dissolved oxygen transported into reach during year (kg O2)
rchyro(39,:) dissolved oxygen transported out of reach during year (kg O2)
rchyro(40,:) persistent bacteria transported out of reach during year (kg bact)
rchyro(41,:) less persistent bacteria transported out of reach during year (kg bact)

- real *8, dimension(:,:), allocatable **wpstaao**
- real *8, dimension(:,:), allocatable **hrumono**

HRU monthly output data array (varies)

hrumono(1,:) precipitation in HRU during month (mm H2O)
hrumono(2,:) amount of precipitation falling as freezing rain/snow in HRU during month (mm H2O)
hrumono(3,:) amount of snow melt in HRU during month (mm H2O)
hrumono(4,:) amount of surface runoff to main channel from HRU during month (ignores impact of transmission losses) (mm H2O)
hrumono(5,:) amount of lateral flow contribution to main channel from HRU during month (mm H2O)
hrumono(6,:) amount of groundwater flow contribution to main channel from HRU during month (mm H2O)
hrumono(7,:) amount of water moving from shallow aquifer to plants or soil profile in HRU during month (mm H2O)
hrumono(8,:) amount of water recharging deep aquifer in HRU during month (mm H2O)
hrumono(9,:) total amount of water entering both aquifers from HRU during month (mm H2O)
hrumono(10,:) water yield (total amount of water entering main channel) from HRU during month (mm H2O)
hrumono(11,:) amount of water percolating out of the soil profile and into the vadose zone in HRU during month (mm H2O)
hrumono(12,:) actual evapotranspiration in HRU during month (mm H2O)
hrumono(13,:) amount of transmission losses from tributary channels in HRU for month (mm H2O)
hrumono(14,:) sediment yield from HRU for month (metric tons/ha)
hrumono(15,:) actual amount of transpiration that occurs during month in HRU (mm H2O)
hrumono(16,:) actual amount of evaporation (from soil) that occurs during month in HRU (mm H2O)
hrumono(17,:) amount of nitrogen applied in continuous fertilizer operation during month in HRU (kg N/ha)
hrumono(18,:) amount of phosphorus applied in continuous fertilizer operation during month in HRU (kg P/ha)
hrumono(22,:) amount of irrigation water applied to HRU during month (mm H2O)
hrumono(23,:) amount of water removed from shallow aquifer in HRU for irrigation during month (mm H2O)
hrumono(24,:) amount of water removed from deep aquifer in HRU for irrigation during month (mm H2O)
hrumono(25,:) potential evapotranspiration in HRU during month (mm H2O)
hrumono(26,:) monthly amount of N (organic & mineral) applied in HRU during grazing (kg N/ha)
hrumono(27,:) monthly amount of P (organic & mineral) applied in HRU during grazing (kg P/ha)
hrumono(28,:) monthly amount of N (organic & mineral) auto-applied in HRU (kg N/ha)
hrumono(29,:) monthly amount of P (organic & mineral) auto-applied in HRU (kg P/ha)
hrumono(31,:) water stress days in HRU during month (stress days)
hrumono(32,:) temperature stress days in HRU during month (stress days)

- humono(33,): nitrogen stress days in HRU during month (stress days)*
 - humono(34,): phosphorus stress days in HRU during month (stress days)*
 - humono(35,): organic nitrogen in surface runoff in HRU during month (kg N/ha)*
 - humono(36,): organic phosphorus in surface runoff in HRU during month (kg P/ha)*
 - humono(37,): nitrate in surface runoff in HRU during month (kg N/ha)*
 - humono(38,): nitrate in lateral flow in HRU during month (kg N/ha)*
 - humono(39,): soluble phosphorus in surface runoff in HRU during month (kg P/ha)*
 - humono(40,): amount of nitrogen removed from soil by plant uptake in HRU during month (kg N/ha)*
 - humono(41,): nitrate percolating past bottom of soil profile in HRU during month (kg N/ha)*
 - humono(42,): amount of phosphorus removed from soil by plant uptake in HRU during month (kg P/ha)*
 - humono(43,): amount of phosphorus moving from labile mineral to active mineral pool in HRU during month (kg P/ha)*
 - humono(44,): amount of phosphorus moving from active mineral to stable mineral pool in HRU during month (kg P/ha)*
 - humono(45,): amount of nitrogen applied to HRU in fertilizer and grazing operations during month (kg N/ha)*
 - humono(46,): amount of phosphorus applied to HRU in fertilizer and grazing operations during month (kg P/ha)*
 - humono(47,): amount of nitrogen added to soil by fixation in HRU during month (kg N/ha)*
 - humono(48,): amount of nitrogen lost by denitrification in HRU during month (kg N/ha)*
 - humono(49,): amount of nitrogen moving from active organic to nitrate pool in HRU during month (kg N/ha)*
 - humono(50,): amount of nitrogen moving from active organic to stable organic pool in HRU during month (kg N/ha)*
 - humono(51,): amount of phosphorus moving from organic to labile mineral pool in HRU during month (kg P/ha)*
 - humono(52,): amount of nitrogen moving from fresh organic to nitrate and active organic pools in HRU during month (kg N/ha)*
 - humono(53,): amount of phosphorus moving from fresh organic to the labile mineral and organic pools in HRU during month (kg P/ha)*
 - humono(54,): amount of nitrogen added to soil in rain (kg N/ha)*
 - humono(61,): daily soil loss predicted with USLE equation (metric tons/ha)*
 - humono(63,): less persistent bacteria transported to main channel from HRU during month (#bacteria/ha)*
 - humono(64,): persistent bacteria transported to main channel from HRU during month (#bacteria/ha)*
 - humono(65,): nitrate loading from groundwater in HRU to main channel during month (kg N/ha)*
 - humono(66,): soluble P loading from groundwater in HRU to main channel during month (kg P/ha)*
 - humono(67,): loading of mineral P attached to sediment in HRU to main channel during month (kg P/ha)*
- **real *8, dimension(:,), allocatable [rchdy](#)**
 - rchdy(1,): flow into reach on day (m^3/s)*
 - rchdy(2,): flow out of reach on day (m^3/s)*
 - rchdy(3,): evaporation from reach on day (m^3/s)*
 - rchdy(4,): transmission losses from reach on day (m^3/s)*
 - rchdy(5,): sediment transported into reach on day (metric tons)*
 - rchdy(6,): sediment transported out of reach on day (metric tons)*
 - rchdy(7,): sediment concentration in outflow (mg/L)*
 - rchdy(8,): organic N transported into reach on day (kg N)*
 - rchdy(9,): organic N transported out of reach on day (kg N)*
 - rchdy(10,): organic P transported into reach on day (kg P)*
 - rchdy(11,): organic P transported out of reach on day (kg P)*
 - rchdy(12,): nitrate transported into reach on day (kg N)*
 - rchdy(13,): nitrate transported out of reach on day (kg N)*
 - rchdy(14,): ammonia transported into reach on day (kg N)*
 - rchdy(15,): ammonia transported out of reach on day (kg N)*
 - rchdy(16,): nitrite transported into reach on day (kg N)*
 - rchdy(17,): nitrite transported out of reach on day (kg N)*
 - rchdy(18,): soluble P transported into reach on day (kg P)*
 - rchdy(19,): soluble P transported out of reach on day (kg P)*
 - rchdy(20,): chlorophyll-a transported into reach on day (kg chl a)*
 - rchdy(21,): chlorophyll-a transported out of reach on day (kg chl a)*
 - rchdy(22,): CBOD transported into reach on day (kg O₂)*
 - rchdy(23,): CBOD transported out of reach on day (kg O₂)*
 - rchdy(24,): dissolved oxygen transported into reach on day (kg O₂)*
 - rchdy(25,): dissolved oxygen transported out of reach on day (kg O₂)*
 - rchdy(26,): soluble pesticide transported into reach on day (mg pst)*
 - rchdy(27,): soluble pesticide transported out of reach on day (mg pst)*
 - rchdy(28,): sorbed pesticide transported into reach on day (mg pst)*
 - rchdy(29,): sorbed pesticide transported out of reach on day (mg pst)*
 - rchdy(30,): amount of pesticide lost through reactions in reach on day (mg pst)*

- rchdy(31,:) amount of pesticide lost through volatilization from reach on day (mg pst)*
- rchdy(32,:) amount of pesticide settling out of reach to bed sediment on day (mg pst)*
- rchdy(33,:) amount of pesticide resuspended from bed sediment to reach on day (mg pst)*
- rchdy(34,:) amount of pesticide diffusing from reach to bed sediment on day (mg pst)*
- rchdy(35,:) amount of pesticide in sediment layer lost through reactions on day (mg pst)*
- rchdy(36,:) amount of pesticide in sediment layer lost through burial on day (mg pst)*
- rchdy(37,:) amount of pesticide stored in river bed sediments (mg pst)*
- rchdy(38,:) persistent bacteria transported out of reach on day (kg bact)*
- rchdy(39,:) less persistent bacteria transported out of reach on day (kg bact)*
- rchdy(40,:) amount of conservative metal #1 transported out of reach on day (kg)*
- rchdy(41,:) amount of conservative metal #2 transported out of reach on day (kg)*
- rchdy(42,:) amount of conservative metal #3 transported out of reach on day (kg)*
- rchdy(43,:) total N (org N + no3 + no2 + nh4 outs) (kg)*
- rchdy(44,:) total P (org P + sol p outs) (kg)*
- real *8, dimension(:,:), allocatable **hruyro**
 - HRU annual output array (varies) hruyro(1,:) precipitation in HRU during year (mm H2O)*
 - hruyro(2,:) amount of precipitation falling as freezing rain/snow in HRU during year (mm H2O)*
 - hruyro(3,:) amount of snow melt in HRU during year (mm H2O)*
 - hruyro(4,:) amount of surface runoff to main channel from HRU during year (ignores impact of transmission losses) (mm H2O)*
 - hruyro(5,:) amount of lateral flow contribution to main channel from HRU during year (mm H2O)*
 - hruyro(6,:) amount of groundwater flow contribution to main channel from HRU during year (mm H2O)*
 - hruyro(7,:) amount of water moving from shallow aquifer to plants or soil profile in HRU during year (mm H2O)*
 - hruyro(8,:) amount of water recharging deep aquifer in HRU during year (mm H2O)*
 - hruyro(9,:) total amount of water entering both aquifers from HRU during year (mm H2O)*
 - hruyro(10,:) water yield (total amount of water entering main channel) from HRU during year (mm H2O)*
 - hruyro(11,:) amount of water percolating out of the soil profile and into the vadose zone in HRU during year (mm H2O)*
 - hruyro(12,:) actual evapotranspiration in HRU during year (mm H2O)*
 - hruyro(13,:) amount of transmission losses from tributary channels in HRU for year (mm H2O)*
 - hruyro(14,:) sediment yield from HRU for year (metric tons/ha)*
 - hruyro(15,:) actual amount of transpiration that occurs during year in HRU (mm H2O)*
 - hruyro(16,:) actual amount of evaporation (from soil) that occurs during year in HRU (mm H2O)*
 - hruyro(17,:) amount of nitrogen applied in continuous fertilizer operation during year in HRU (kg N/ha)*
 - hruyro(18,:) amount of phosphorus applied in continuous fertilizer operation during year in HRU (kg P/ha)*
 - hruyro(23,:) amount of water removed from shallow aquifer in HRU for irrigation during year (mm H2O)*
 - hruyro(24,:) amount of water removed from deep aquifer in HRU for irrigation during year (mm H2O)*
 - hruyro(25,:) potential evapotranspiration in HRU during year (mm H2O)*
 - hruyro(26,:) annual amount of N (organic & mineral) applied in HRU during grazing (kg N/ha)*
 - hruyro(27,:) annual amount of P (organic & mineral) applied in HRU during grazing (kg P/ha)*
 - hruyro(28,:) annual amount of N (organic & mineral) auto-applied in HRU (kg N/ha)*
 - hruyro(29,:) annual amount of P (organic & mineral) auto-applied in HRU (kg P/ha)*
 - hruyro(31,:) water stress days in HRU during year (stress days)*
 - hruyro(32,:) temperature stress days in HRU during year (stress days)*
 - hruyro(33,:) nitrogen stress days in HRU during year (stress days)*
 - hruyro(34,:) phosphorus stress days in HRU during year (stress days)*
 - hruyro(35,:) organic nitrogen in surface runoff in HRU during year (kg N/ha)*
 - hruyro(36,:) organic phosphorus in surface runoff in HRU during year (kg P/ha)*
 - hruyro(37,:) nitrate in surface runoff in HRU during year (kg N/ha)*
 - hruyro(38,:) nitrate in lateral flow in HRU during year (kg N/ha)*
 - hruyro(39,:) soluble phosphorus in surface runoff in HRU during year (kg P/ha)*
 - hruyro(40,:) amount of nitrogen removed from soil by plant uptake in HRU during year (kg N/ha)*
 - hruyro(41,:) nitrate percolating past bottom of soil profile in HRU during year (kg N/ha)*
 - hruyro(42,:) amount of phosphorus removed from soil by plant uptake in HRU during year (kg P/ha)*
 - hruyro(43,:) amount of phosphorus moving from labile mineral to active mineral pool in HRU during year (kg P/ha)*
 - hruyro(44,:) amount of phosphorus moving from active mineral to stable mineral pool in HRU during year (kg P/ha)*
 - hruyro(45,:) amount of nitrogen applied to HRU in fertilizer and grazing operations during year (kg N/ha)*
 - hruyro(46,:) amount of phosphorus applied to HRU in fertilizer and grazing operations during year (kg P/ha)*
 - hruyro(47,:) amount of nitrogen added to soil by fixation in HRU during year (kg N/ha)*
 - hruyro(48,:) amount of nitrogen lost by denitrification in HRU during year (kg N/ha)*
 - hruyro(49,:) amount of nitrogen moving from active organic to nitrate pool in HRU during year (kg N/ha)*
 - hruyro(50,:) amount of nitrogen moving from active organic to stable organic pool in HRU during year (kg N/ha)*
 - hruyro(51,:) amount of phosphorus moving from organic to labile mineral pool in HRU during year (kg P/ha)*

- hruyro(52,:)* amount of nitrogen moving from fresh organic to nitrate and active organic pools in HRU during year (kg N/ha)
- hruyro(53,:)* amount of phosphorus moving from fresh organic to the labile mineral and organic pools in HRU during year (kg P/ha)
- hruyro(54,:)* amount of nitrogen added to soil in rain during year (kg N/ha)
- hruyro(61,:)* daily soil loss predicted with USLE equation (metric tons/ha)
- hruyro(63,:)* less persistent bacteria transported to main channel from HRU during year (# bacteria/ha)
- hruyro(64,:)* persistent bacteria transported to main channel from HRU during year (# bacteria/ha)
- hruyro(65,:)* nitrate loading from groundwater in HRU to main channel during year (kg N/ha)
- hruyro(66,:)* soluble P loading from groundwater in HRU to main channel during year (kg P/ha)
- hruyro(67,:)* loading of mineral P attached to sediment in HRU to main channel during year (kg P/ha)
- real *8, dimension(:,), allocatable **rchaao**
 - reach average annual output array (varies)
 - rchaao(1,:)* flow into reach during simulation (m^3/s)
 - rchaao(2,:)* flow out of reach during simulation (m^3/s)
 - rchaao(3,:)* sediment transported into reach during simulation (metric tons)
 - rchaao(4,:)* sediment transported out of reach during simulation (metric tons)
 - rchaao(5,:)* sediment concentration in outflow during simulation (mg/L)
 - rchaao(6,:)* organic N transported into reach during simulation (kg N)
 - rchaao(7,:)* organic N transported out of reach during simulation (kg N)
 - rchaao(8,:)* organic P transported into reach during simulation (kg P)
 - rchaao(9,:)* organic P transported out of reach during simulation (kg P)
 - rchaao(10,:)* evaporation from reach during simulation (m^3/s)
 - rchaao(11,:)* transmission losses from reach during simulation (m^3/s)
 - rchaao(12,:)* conservative metal #1 transported out of reach during simulation (kg)
 - rchaao(13,:)* conservative metal #2 transported out of reach during simulation (kg)
 - rchaao(14,:)* conservative metal #3 transported out of reach during simulation (kg)
 - rchaao(15,:)* nitrate transported into reach during simulation (kg N)
 - rchaao(16,:)* nitrate transported out of reach during simulation (kg N)
 - rchaao(17,:)* soluble P transported into reach during simulation (kg P)
 - rchaao(18,:)* soluble P transported out of reach during simulation (kg P)
 - rchaao(19,:)* soluble pesticide transported into reach during simulation
 - rchaao(20,:)* soluble pesticide transported out of reach during simulation
 - rchaao(21,:)* sorbed pesticide transported into reach during simulation
 - rchaao(22,:)* sorbed pesticide transported out of reach during simulation
 - rchaao(23,:)* amount of pesticide lost through reactions in reach during simulation
 - rchaao(24,:)* amount of pesticide lost through volatilization from reach during simulation
 - rchaao(25,:)* amount of pesticide settling out of reach to bed sediment during simulation
 - rchaao(26,:)* amount of pesticide resuspended from bed sediment to reach during simulation
 - rchaao(27,:)* amount of pesticide diffusing from reach to bed sediment during simulation
 - rchaao(28,:)* amount of pesticide in sediment layer lost through reactions during simulation
 - rchaao(29,:)* amount of pesticide in sediment layer lost through burial during simulation
 - rchaao(30,:)* chlorophyll-a transported into reach during simulation (kg chl-a)
 - rchaao(31,:)* chlorophyll-a transported out of reach during simulation (kg chl-a)
 - rchaao(32,:)* ammonia transported into reach during simulation (kg N)
 - rchaao(33,:)* ammonia transported out of reach during simulation (kg N)
 - rchaao(34,:)* nitrite transported into reach during simulation (kg N)
 - rchaao(35,:)* nitrite transported out of reach during simulation (kg N)
 - rchaao(36,:)* CBOD transported into reach during simulation (kg O₂)
 - rchaao(37,:)* CBOD transported out of reach during simulation (kg O₂)
 - rchaao(38,:)* dissolved oxygen transported into reach during simulation (kg O₂)
 - rchaao(39,:)* dissolved oxygen transported out of reach during simulation (kg O₂)
 - rchaao(40,:)* persistent bacteria transported out of reach during simulation (kg bact)
 - rchaao(41,:)* less persistent bacteria transported out of reach during simulation (kg bact)
 - rchaao(43,:)* Total N (org N + no₃ + no₂ + nh₄ outs) (kg)
 - rchaao(44,:)* Total P (org P + sol p outs) (kg)
- real *8, dimension(:,), allocatable **submono**
 - subbasin monthly output array (varies)
 - submono(1,:)* precipitation in subbasin for month (mm H₂O)
 - submono(2,:)* snow melt in subbasin for month (mm H₂O)
 - submono(3,:)* surface runoff loading in subbasin for month (mm H₂O)
 - submono(4,:)* water yield from subbasin for month (mm H₂O)
 - submono(5,:)* potential evapotranspiration in subbasin for month (mm H₂O)

- submono(6,:)* actual evapotranspiration in subbasin for month (mm H2O)
 - submono(7,:)* sediment yield from subbasin for month (metric tons/ha)
 - submono(8,:)* organic N loading from subbasin for month (kg N/ha)
 - submono(9,:)* organic P loading from subbasin for month (kg P/ha)
 - submono(10,:)* NO3 loading from surface runoff in subbasin for month (kg N/ha)
 - submono(11,:)* soluble P loading from subbasin for month (kg P/ha)
 - submono(12,:)* groundwater loading from subbasin for month (mm H2O)
 - submono(13,:)* percolation out of soil profile in subbasin for month (mm H2O)
 - submono(14,:)* loading to reach of mineral P attached to sediment from subbasin for month (kg P/ha)
- real *8, dimension(:,:), allocatable **subyro**
 - subbasin* annual output array (varies)
 - subyro(1,:)* precipitation in subbasin for year (mm H2O)
 - subyro(2,:)* snow melt in subbasin for year (mm H2O)
 - subyro(3,:)* surface runoff loading in subbasin for year (mm H2O)
 - subyro(4,:)* water yield from subbasin for year (mm H2O)
 - subyro(5,:)* potential evapotranspiration in subbasin for year (mm H2O)
 - subyro(6,:)* actual evapotranspiration in subbasin for year (mm H2O)
 - subyro(7,:)* sediment yield from subbasin for year (metric tons/ha)
 - subyro(8,:)* organic N loading from subbasin for year (kg N/ha)
 - subyro(9,:)* organic P loading from subbasin for year (kg P/ha)
 - subyro(10,:)* NO3 loading from surface runoff in subbasin for year (kg N/ha)
 - subyro(11,:)* soluble P loading from subbasin for year (kg P/ha)
 - subyro(12,:)* groundwater loading from subbasin for year (mm H2O)
 - subyro(13,:)* percolation out of soil profile in subbasin for year (mm H2O)
 - subyro(14,:)* loading to reach of mineral P attached to sediment from subbasin for year (kg P/ha)
- real *8, dimension(:,:), allocatable **hruaao**
 - HRU* average annual output array (varies)
 - hruaao(1,:)* precipitation in HRU during simulation (mm H2O)
 - hruaao(2,:)* amount of precipitation falling as freezing rain/snow in HRU during simulation (mm H2O)
 - hruaao(3,:)* amount of snow melt in HRU during simulation (mm H2O)
 - hruaao(4,:)* amount of surface runoff to main channel from HRU during simulation (ignores impact of transmission losses) (mm H2O)
 - hruaao(5,:)* amount of lateral flow contribution to main channel from HRU during simulation (mm H2O)
 - hruaao(6,:)* amount of groundwater flow contribution to main channel from HRU during simulation (mm H2O)
 - hruaao(7,:)* amount of water moving from shallow aquifer to plants or soil profile in HRU during simulation (mm H2O)
 - hruaao(8,:)* amount of water recharging deep aquifer in HRU during simulation (mm H2O)
 - hruaao(9,:)* total amount of water entering both aquifers from HRU during simulation (mm H2O)
 - hruaao(10,:)* water yield (total amount of water entering main channel) from HRU during simulation (mm H2O)
 - hruaao(11,:)* amount of water percolating out of the soil profile and into the vadose zone in HRU during simulation (mm H2O)
 - hruaao(12,:)* actual evapotranspiration in HRU during simulation
 - hruaao(13,:)* amount of transmission losses from tributary channels in HRU for simulation (mm H2O)
 - hruaao(14,:)* sediment yield from HRU for simulation (metric tons/ha)
 - hruaao(17,:)* amount of nitrogen applied in continuous fertilizer operation in HRU for simulation (kg N/ha)
 - hruaao(18,:)* amount of phosphorus applied in continuous fertilizer operation in HRU for simulation (kg P/ha)
 - hruaao(23,:)* amount of water removed from shallow aquifer in HRU for irrigation during simulation (mm H2O)
 - hruaao(24,:)* amount of water removed from deep aquifer in HRU for irrigation during simulation (mm H2O)
 - hruaao(25,:)* potential evapotranspiration in HRU during simulation (mm H2O)
 - hruaao(26,:)* annual amount of N (organic & mineral) applied in HRU during grazing (kg N/ha)
 - hruaao(27,:)* annual amount of P (organic & mineral) applied in HRU during grazing (kg P/ha)
 - hruaao(28,:)* average annual amount of N (organic & mineral) auto-applied in HRU (kg N/ha)
 - hruaao(29,:)* average annual amount of P (organic & mineral) auto-applied in HRU (kg P/ha)
 - hruaao(31,:)* water stress days in HRU during simulation (stress days)
 - hruaao(32,:)* temperature stress days in HRU during simulation (stress days)
 - hruaao(33,:)* nitrogen stress days in HRU during simulation (stress days)
 - hruaao(34,:)* phosphorus stress days in HRU during simulation (stress days)
 - hruaao(35,:)* organic nitrogen in surface runoff in HRU during simulation (kg N/ha)
 - hruaao(36,:)* organic phosphorus in surface runoff in HRU during simulation (kg P/ha)
 - hruaao(37,:)* nitrate in surface runoff in HRU during simulation (kg N/ha)
 - hruaao(38,:)* nitrate in lateral flow in HRU during simulation (kg N/ha)
 - hruaao(39,:)* soluble phosphorus in surface runoff in HRU during simulation (kg P/ha)
 - hruaao(40,:)* amount of nitrogen removed from soil by plant uptake in HRU during simulation (kg N/ha)
 - hruaao(41,:)* nitrate percolating past bottom of soil profile in HRU during simulation (kg N/ha)

- hruaao(42,:)* amount of phosphorus removed from soil by plant uptake in HRU during simulation (kg P/ha)
hruaao(43,:) amount of phosphorus moving from labile mineral to active mineral pool in HRU during simulation (kg P/ha)
hruaao(44,:) amount of phosphorus moving from active mineral to stable mineral pool in HRU during simulation (kg P/ha)
hruaao(45,:) amount of nitrogen applied to HRU in fertilizer and grazing operations during simulation (kg N/ha)
hruaao(46,:) amount of phosphorus applied to HRU in fertilizer and grazing operations during simulation (kg P/ha)
hruaao(47,:) amount of nitrogen added to soil by fixation in HRU during simulation (kg N/ha)
hruaao(48,:) amount of nitrogen lost by denitrification in HRU during simulation (kg N/ha)
hruaao(49,:) amount of nitrogen moving from active organic to nitrate pool in HRU during simulation (kg N/ha)
hruaao(50,:) amount of nitrogen moving from active organic to stable organic pool in HRU during simulation (kg N/ha)
hruaao(51,:) amount of phosphorus moving from organic to labile mineral pool in HRU during simulation (kg P/ha)
hruaao(52,:) amount of nitrogen moving from fresh organic to nitrate and active organic pools in HRU during simulation (kg N/ha)
hruaao(53,:) amount of phosphorus moving from fresh organic to the labile mineral and organic pools in HRU during simulation (kg P/ha)
hruaao(54,:) amount of nitrogen added to soil in rain during simulation (kg N/ha)
hruaao(61,:) daily soil loss predicted with USLE equation (metric tons/ha)
hruaao(63,:) less persistent bacteria transported to main channel from HRU during simulation (# bacteria/ha)
hruaao(64,:) persistent bacteria transported to main channel from HRU during simulation (# bacteria/ha)
hruaao(65,:) nitrate loading from groundwater in HRU to main channel during simulation (kg N/ha)
hruaao(66,:) soluble P loading from groundwater in HRU to main channel during simulation (kg P/ha)
hruaao(67,:) loading of mineral P attached to sediment in HRU to main channel during simulation (kg P/ha)
- real *8, dimension(:,:), allocatable [subaao](#)
 subbasin average annual output array (varies)
 - real *8, dimension(:,:), allocatable [resoutm](#)
 reservoir monthly output array (varies)
resoutm(1,:) flow into reservoir during month (m^3/s)
resoutm(2,:) flow out of reservoir during month (m^3/s)
resoutm(3,:) sediment entering reservoir during month (metric tons)
resoutm(4,:) sediment leaving reservoir during month (metric tons)
resoutm(5,:) sediment concentration in reservoir during month (mg/L)
resoutm(6,:) pesticide entering reservoir during month (mg pst)
resoutm(7,:) pesticide lost from reservoir through reactions during month (mg pst)
resoutm(8,:) pesticide lost from reservoir through volatilization during month (mg pst)
resoutm(9,:) pesticide moving from water to sediment through settling during month (mg pst)
resoutm(10,:) pesticide moving from sediment to water through resuspension during month (mg pst)
resoutm(11,:) pesticide moving from water to sediment through diffusion during month (mg pst)
resoutm(12,:) pesticide lost from reservoir sediment layer through reactions during month (mg pst)
resoutm(13,:) pesticide lost from reservoir sediment layer through burial during month (mg pst)
resoutm(14,:) pesticide transported out of reservoir during month (mg pst)
resoutm(15,:) pesticide concentration in reservoir water during month ($\text{mg pst}/\text{m}^3$)
resoutm(16,:) pesticide concentration in reservoir sediment layer during month ($\text{mg pst}/\text{m}^3$)
resoutm(17,:) evaporation from reservoir during month ($\text{m}^3 \text{H}_2\text{O}$)
resoutm(18,:) seepage from reservoir during month ($\text{m}^3 \text{H}_2\text{O}$)
resoutm(19,:) precipitation on reservoir during month ($\text{m}^3 \text{H}_2\text{O}$)
resoutm(22,:) organic N entering reservoir during month (kg N)
resoutm(23,:) organic N leaving reservoir during month (kg N)
resoutm(24,:) organic P entering reservoir during month (kg P)
resoutm(25,:) organic P leaving reservoir during month (kg P)
resoutm(26,:) nitrate entering reservoir during month (kg N)
resoutm(27,:) nitrate leaving reservoir during month (kg N)
resoutm(28,:) nitrite entering reservoir during month (kg N)
resoutm(29,:) nitrite leaving reservoir during month (kg N)
resoutm(30,:) ammonia entering reservoir during month (kg N)
resoutm(31,:) ammonia leaving reservoir during month (kg N)
resoutm(32,:) mineral P entering reservoir during month (kg P)
resoutm(33,:) mineral P leaving reservoir during month (kg P)
resoutm(34,:) chlorophyll-a entering reservoir during month (kg chl a)
resoutm(35,:) chlorophyll-a leaving reservoir during month (kg chl a)
resoutm(36,:) organic P concentration in reservoir water during month (mg P/L)
resoutm(37,:) mineral P concentration in reservoir water during month (mg P/L)
resoutm(38,:) organic N concentration in reservoir water during month (mg N/L)

resoutm(39,:) nitrate concentration in reservoir water during month (mg N/L)
resoutm(40,:) nitrite concentration in reservoir water during month (mg N/L)
resoutm(41,:) ammonia concentration in reservoir water during month (mg N/L)

- real *8, dimension(:,:), allocatable **resouty**

reservoir annual output array (varies)
resouty(1,:) flow into reservoir during year (m^3/s)
resouty(2,:) flow out of reservoir during year (m^3/s)
resouty(3,:) sediment entering reservoir during year (metric tons)
resouty(4,:) sediment leaving reservoir during year (metric tons)
resouty(5,:) sediment concentration in reservoir during year (mg/L)
resouty(6,:) pesticide entering reservoir during year (mg pst)
resouty(7,:) pesticide lost from reservoir through reactions during year (mg pst)
resouty(8,:) pesticide lost from reservoir through volatilization during year (mg pst)
resouty(9,:) pesticide moving from water to sediment through settling during year (mg pst)
resouty(10,:) pesticide moving from sediment to water through resuspension during year (mg pst)
resouty(11,:) pesticide moving from water to sediment through diffusion during year (mg pst)
resouty(12,:) pesticide lost from reservoir sediment layer through reactions during year (mg pst)
resouty(13,:) pesticide lost from reservoir sediment layer through burial during year (mg pst)
resouty(14,:) pesticide transported out of reservoir during year (mg pst)
resouty(15,:) pesticide concentration in reservoir water during year (mg pst/ m^3)
resouty(16,:) pesticide concentration in reservoir sediment layer during year (mg pst/ m^3)
resouty(17,:) evaporation from reservoir during year ($m^3 H_2O$)
resouty(18,:) seepage from reservoir during year ($m^3 H_2O$)
resouty(19,:) precipitation on reservoir during year ($m^3 H_2O$)
resouty(22,:) organic N entering reservoir during year (kg N)
resouty(23,:) organic N leaving reservoir during year (kg N)
resouty(24,:) organic P entering reservoir during year (kg P)
resouty(25,:) organic P leaving reservoir during year (kg P)
resouty(26,:) nitrate entering reservoir during year (kg N)
resouty(27,:) nitrate leaving reservoir during year (kg N)
resouty(28,:) nitrite entering reservoir during year (kg N)
resouty(29,:) nitrite leaving reservoir during year (kg N)
resouty(30,:) ammonia entering reservoir during year (kg N)
resouty(31,:) ammonia leaving reservoir during year (kg N)
resouty(32,:) mineral P entering reservoir during year (kg P)
resouty(33,:) mineral P leaving reservoir during year (kg P)
resouty(34,:) chlorophyll-a entering reservoir during year (kg chl a)
resouty(35,:) chlorophyll-a leaving reservoir during year (kg chl a)
resouty(36,:) organic P concentration in reservoir water during year (mg P/L)
resouty(37,:) mineral P concentration in reservoir water during year (mg P/L)
resouty(38,:) organic N concentration in reservoir water during year (mg N/L)
resouty(39,:) nitrate concentration in reservoir water during year (mg N/L)
resouty(40,:) nitrite concentration in reservoir water during year (mg N/L)
resouty(41,:) ammonia concentration in reservoir water during year (mg N/L)

- real *8, dimension(:,:), allocatable **resouta**

reservoir average annual output array (varies)
resouta(3,:) sediment entering reservoir during simulation (metric tons)
resouta(4,:) sediment leaving reservoir during simulation (metric tons)
resouta(17,:) evaporation from reservoir during simulation ($m^3 H_2O$)
resouta(18,:) seepage from reservoir during simulation ($m^3 H_2O$)
resouta(19,:) precipitation on reservoir during simulation ($m^3 H_2O$)
resouta(20,:) water entering reservoir during simulation ($m^3 H_2O$)
resouta(21,:) water leaving reservoir during simulation ($m^3 H_2O$)

- real *8, dimension(12, 8) **wshd_aamon**

wshd_aamon(:,1) average annual precipitation in watershed falling during month (mm H₂O)
wshd_aamon(:,2) average annual freezing rain in watershed falling during month (mm H₂O)
wshd_aamon(:,3) average annual surface runoff in watershed during month (mm H₂O)
wshd_aamon(:,4) average annual lateral flow in watershed during month (mm H₂O)
wshd_aamon(:,5) average annual water yield in watershed during month (mm H₂O)
wshd_aamon(:,6) average annual actual evapotranspiration in watershed during month (mm H₂O)
wshd_aamon(:,7) average annual sediment yield in watershed during month (metric tons)
wshd_aamon(:,8) average annual potential evapotranspiration in watershed during month (mm H₂O)

- real *8, dimension(:,:), allocatable [wtrmon](#)

HRU monthly output data array for impoundments (varies)

wtrmon(1,:) evaporation from ponds in HRU for month (mm H2O)
wtrmon(2,:) seepage from ponds in HRU for month (mm H2O)
wtrmon(3,:) precipitation on ponds in HRU for month (mm H2O)
wtrmon(4,:) amount of water entering ponds in HRU for month (mm H2O)
wtrmon(5,:) sediment entering ponds in HRU for month (metric tons/ha)
wtrmon(6,:) amount of water leaving ponds in HRU for month (mm H2O)
wtrmon(7,:) sediment leaving ponds in HRU for month (metric tons/ha)
wtrmon(8,:) precipitation on wetlands in HRU for month (mm H2O)
wtrmon(9,:) volume of water entering wetlands from HRU for month (mm H2O)
wtrmon(10,:) sediment loading to wetlands for month from HRU (metric tons/ha)
wtrmon(11,:) evaporation from wetlands in HRU for month (mm H2O)
wtrmon(12,:) seepage from wetlands in HRU for month (mm H2O)
wtrmon(13,:) volume of water leaving wetlands in HRU for month (mm H2O)
wtrmon(14,:) sediment loading from wetlands in HRU to main channel during month (metric tons/ha)
wtrmon(15,:) precipitation on potholes in HRU for month (mm H2O)
wtrmon(16,:) evaporation from potholes in HRU for month (mm H2O)
wtrmon(17,:) seepage from potholes in HRU for month (mm H2O)
wtrmon(18,:) water leaving potholes in HRU for month (mm H2O)
wtrmon(19,:) water entering potholes in HRU for month (mm H2O)
wtrmon(20,:) sediment entering potholes in HRU for month (metric tons/ha)
wtrmon(21,:) sediment leaving potholes in HRU for month (metric tons/ha)

- real *8, dimension(:,:), allocatable [wtryr](#)

HRU impoundment annual output array (varies)

wtryr(1,:) evaporation from ponds in HRU for year (mm H2O)
wtryr(2,:) seepage from ponds in HRU for year (mm H2O)
wtryr(3,:) precipitation on ponds in HRU for year (mm H2O)
wtryr(4,:) amount of water entering ponds in HRU for year (mm H2O)
wtryr(5,:) sediment entering ponds in HRU for year (metric tons/ha)
wtryr(6,:) amount of water leaving ponds in HRU for year (mm H2O)
wtryr(7,:) sediment leaving ponds in HRU for year (metric tons/ha)
wtryr(8,:) precipitation on wetlands in HRU for year (mm H2O)
wtryr(9,:) volume of water entering wetlands from HRU for year (mm H2O)
wtryr(10,:) sediment loading to wetlands for year from HRU (metric tons/ha)
wtryr(11,:) evaporation from wetlands in HRU for year (mm H2O)
wtryr(12,:) seepage from wetlands in HRU for year (mm H2O)
wtryr(13,:) volume of water leaving wetlands in HRU for year (mm H2O)
wtryr(14,:) sediment loading from wetlands in HRU to main channel during year (metric tons/ha)
wtryr(15,:) precipitation on potholes in HRU during year (mm H2O)
wtryr(16,:) evaporation from potholes in HRU during year (mm H2O)
wtryr(17,:) seepage from potholes in HRU during year (mm H2O)
wtryr(18,:) water leaving potholes in HRU during year (mm H2O)
wtryr(19,:) water entering potholes in HRU during year (mm H2O)
wtryr(20,:) sediment entering potholes in HRU during year (metric tons/ha)
wtryr(21,:) sediment leaving potholes in HRU during year (metric tons/ha)

- real *8, dimension(:,:), allocatable [wtraa](#)

HRU impoundment average annual output array (varies)

- real *8, dimension(:,:), allocatable [sub_smfmn](#)

max melt rate for snow during year (June 21) for subbasin(:) where deg C refers to the air temperature. SUB_SMFMX and SMFMN allow the rate of snow melt to vary through the year. These parameters are accounting for the impact of soil temperature on snow melt (range: -5.0/5.0) (mm/deg C/day)

- real *8, dimension(:,:), allocatable [sub_smfmn](#)

min melt rate for snow during year (Dec 21) for subbasin(:) (range: -5.0/5.0) where deg C refers to the air temperature (mm/deg C/day)

- real *8, dimension(:,:,:), allocatable [hrupstd](#)

hrupstd(:,1,:) amount of pesticide type in surface runoff contribution to stream from HRU on day (in solution) (mg pst)
hrupstd(:,2,:) amount of pesticide type in surface runoff contribution to stream from HRU on day (sorbed to sediment) (mg pst)

- real *8, dimension(:,:,:), allocatable [hrupstm](#)

- hrupstm(:, :, :)* HRU monthly pesticide output array (varies)
- hrupstm(:, 1, :)* amount of pesticide type in surface runoff contribution to stream from HRU during month (in solution) (mg pst)
- hrupstm(:, 2, :)* amount of pesticide type in surface runoff contribution to stream from HRU during month (sorbed to sediment) (mg pst)
- hrupstm(:, 3, :)* total pesticide loading to stream in surface runoff from HRU during month (mg pst)
- real *8, dimension(:, :, :), allocatable [hrupsta](#)
 - HRU average annual pesticide output array (varies)*
- real *8, dimension(:, :, :), allocatable [hrupsty](#)
 - hrupsty(:, :, :)* HRU annual pesticide output array (varies)
 - hrupsty(:, 1, :)* amount of pesticide type in surface runoff contribution to stream from HRU during year (in solution) (mg pst)
 - hrupsty(:, 2, :)* amount of pesticide type in surface runoff contribution to stream from HRU during year (sorbed to sediment) (mg pst)
- integer, dimension(:), allocatable [ifirstt](#)
 - temperature data search code (none)*
 - 0 first day of temperature data located in file*
 - 1 first day of temperature data not located in file*
- integer, dimension(:), allocatable **ifirstpcp**
- integer, dimension(:), allocatable [elevp](#)
 - elevation of precipitation gage station (m)*
- integer, dimension(:), allocatable [elevt](#)
 - elevation of temperature gage station (m)*
- real *8, dimension(:, :, :), allocatable [ftmpmn](#)
 - avg monthly minimum air temperature (deg C)*
- real *8, dimension(:, :, :), allocatable [ftmpmx](#)
 - avg monthly maximum air temperature (deg C)*
- real *8, dimension(:, :, :), allocatable [ftmpstdmn](#)
 - standard deviation for avg monthly minimum air temperature (deg C)*
- real *8, dimension(:, :, :), allocatable [ftmpstdmx](#)
 - standard deviation for avg monthly maximum air temperature (deg C)*
- real *8, dimension(:, :, :), allocatable [fpcp_stat](#)
 - fpcp_stat(:, 1, :)* average amount of precipitation falling in one day for the month (mm/day)
 - fpcp_stat(:, 2, :)* standard deviation for the average daily precipitation (mm/day)
 - fpcp_stat(:, 3, :)* skew coefficient for the average daily precipitation (none)
- real *8, dimension(:, :, :), allocatable [fpr_w1](#)
 - probability of wet day after dry day in month (none)*
- real *8, dimension(:, :, :), allocatable [fpr_w2](#)
 - probability of wet day after wet day in month (none)*
- real *8, dimension(:, :, :), allocatable [fpr_w3](#)
 - proportion of wet days in the month (none)*
- real *8, dimension(:), allocatable [ch_d](#)
 - average depth of main channel (m)*
- real *8, dimension(:), allocatable **flwin**
- real *8, dimension(:), allocatable **flwout**
- real *8, dimension(:), allocatable **bankst**
- real *8, dimension(:), allocatable **ch_wi**
- real *8, dimension(:), allocatable [ch_onco](#)
 - channel organic n concentration (ppm)*
- real *8, dimension(:), allocatable [ch_opco](#)
 - channel organic p concentration (ppm)*
- real *8, dimension(:), allocatable **ch_orgn**
- real *8, dimension(:), allocatable **ch_orgp**
- real *8, dimension(:), allocatable [drift](#)

- amount of pesticide drifting onto main channel in subbasin (kg)*
 - real *8, dimension(:), allocatable **rch_dox**
 - real *8, dimension(:), allocatable **rch_bactp**
 - real *8, dimension(:), allocatable **alpha_bnk**
- alpha factor for bank storage recession curve (days)*
 - real *8, dimension(:), allocatable **alpha_bnke**
- $\exp(-\alpha_{bnk})$ (none)
 - real *8, dimension(:), allocatable **disolvp**
 - real *8, dimension(:), allocatable **algae**
 - real *8, dimension(:), allocatable **sedst**
 - real *8, dimension(:), allocatable **rchstor**
 - real *8, dimension(:), allocatable **organicn**
 - real *8, dimension(:), allocatable **organicp**
 - real *8, dimension(:), allocatable **chlora**
 - real *8, dimension(:), allocatable **ch_li**
- initial length of main channel (km)*
 - real *8, dimension(:), allocatable **ch_si**
- initial slope of main channel (m/m)*
 - real *8, dimension(:), allocatable **nitraten**
 - real *8, dimension(:), allocatable **nitriten**
 - real *8, dimension(:), allocatable **ch_bnk_san**
 - real *8, dimension(:), allocatable **ch_bnk_sil**
 - real *8, dimension(:), allocatable **ch_bnk_cla**
 - real *8, dimension(:), allocatable **ch_bnk_gra**
 - real *8, dimension(:), allocatable **ch_bed_san**
 - real *8, dimension(:), allocatable **ch_bed_sil**
 - real *8, dimension(:), allocatable **ch_bed_cla**
 - real *8, dimension(:), allocatable **ch_bed_gra**
 - real *8, dimension(:), allocatable **depfp**
 - real *8, dimension(:), allocatable **depsanfp**
 - real *8, dimension(:), allocatable **depsilfp**
 - real *8, dimension(:), allocatable **depclafp**
 - real *8, dimension(:), allocatable **depsagfp**
 - real *8, dimension(:), allocatable **deplagfp**
 - real *8, dimension(:), allocatable **depch**
 - real *8, dimension(:), allocatable **depsanch**
 - real *8, dimension(:), allocatable **depsilch**
 - real *8, dimension(:), allocatable **depclach**
 - real *8, dimension(:), allocatable **depsagch**
 - real *8, dimension(:), allocatable **deplagch**
 - real *8, dimension(:), allocatable **depgrach**
 - real *8, dimension(:), allocatable **depgrafp**
 - real *8, dimension(:), allocatable **grast**
 - real *8, dimension(:), allocatable **r2adj**
- curve number retention parameter adjustment factor to adjust surface runoff for flat slopes (0.5 - 3.0) (dimensionless)*
 - real *8, dimension(:), allocatable **prf**
- Reach peak rate adjustment factor for sediment routing in the channel. Allows impact of peak flow rate on sediment routing and channel reshaping to be taken into account (none)*
 - real *8, dimension(:), allocatable **depprch**
 - real *8, dimension(:), allocatable **depprfp**
 - real *8, dimension(:), allocatable **spcon**
- linear parameter for calculating sediment reentrained in channel sediment routing*
 - real *8, dimension(:), allocatable **spexp**

exponent parameter for calculating sediment reentrained in channel sediment routing

- real *8, dimension(:), allocatable **sanst**
- real *8, dimension(:), allocatable **silst**
- real *8, dimension(:), allocatable **clast**
- real *8, dimension(:), allocatable **sagst**
- real *8, dimension(:), allocatable **lagst**
- real *8, dimension(:), allocatable **pot_san**
- real *8, dimension(:), allocatable **pot_sil**
- real *8, dimension(:), allocatable **pot_cla**
- real *8, dimension(:), allocatable **pot_sag**
- real *8, dimension(:), allocatable **pot_lag**
- real *8, dimension(:), allocatable **potsani**
- real *8, dimension(:), allocatable **potsili**
- real *8, dimension(:), allocatable **potclai**
- real *8, dimension(:), allocatable **potsagi**
- real *8, dimension(:), allocatable **potlagi**
- real *8, dimension(:), allocatable **sanyld**
- real *8, dimension(:), allocatable **silyld**
- real *8, dimension(:), allocatable **clayld**
- real *8, dimension(:), allocatable **sagyld**
- real *8, dimension(:), allocatable **lagyld**
- real *8, dimension(:), allocatable **grayld**
- real *8, dimension(:), allocatable **res_san**
- real *8, dimension(:), allocatable **res_sil**
- real *8, dimension(:), allocatable **res_cla**
- real *8, dimension(:), allocatable **res_sag**
- real *8, dimension(:), allocatable **res_lag**
- real *8, dimension(:), allocatable **res_gra**
- real *8, dimension(:), allocatable **pnd_san**
- real *8, dimension(:), allocatable **pnd_sil**
- real *8, dimension(:), allocatable **pnd_cla**
- real *8, dimension(:), allocatable **pnd_sag**
- real *8, dimension(:), allocatable **pnd_lag**
- real *8, dimension(:), allocatable **wet_san**
- real *8, dimension(:), allocatable **wet_sil**
- real *8, dimension(:), allocatable **wet_cla**
- real *8, dimension(:), allocatable **wet_lag**
- real *8, dimension(:), allocatable **wet_sag**
- real *8 **ressano**
- real *8 **ressilo**
- real *8 **resclao**
- real *8 **ressago**
- real *8 **reslago**
- real *8 **resgrao**
- real *8 **ressani**
- real *8 **ressili**
- real *8 **resclai**
- real *8 **ressagi**
- real *8 **reslagi**
- real *8 **resgrai**
- real *8 **potsano**
- real *8 **potsilo**
- real *8 **potclao**
- real *8 **potsago**
- real *8 **potlago**

- real *8 **pndsarin**
- real *8 **pndsilin**
- real *8 **pndclain**
- real *8 **pndsagin**
- real *8 **pndlagin**
- real *8 **pndsano**
- real *8 **pndsilo**
- real *8 **pndclao**
- real *8 **pndsago**
- real *8 **pndlago**
- real *8, dimension(:), allocatable **ch_di**
initial depth of main channel (m)
- real *8, dimension(:), allocatable **ch_erod**
channel erodibility factor (0.0-1.0) (none)
0 non-erosive channel
1 no resistance to erosion
- real *8, dimension(:), allocatable **ch_l2**
length of main channel (km)
- real *8, dimension(:), allocatable **ch_cov**
- real *8, dimension(:), allocatable **ch_bnk_bd**
bulk density of channel bank sediment (1.1-1.9) (g/cc)
- real *8, dimension(:), allocatable **ch_bed_bd**
bulk density of channel bed sediment (1.1-1.9) (g/cc)
- real *8, dimension(:), allocatable **ch_bnk_kd**
erodibility of channel bank sediment by jet test (Peter Allen needs to give more info on this)
- real *8, dimension(:), allocatable **ch_bed_kd**
erodibility of channel bed sediment by jet test (Peter Allen needs to give more info on this)
- real *8, dimension(:), allocatable **ch_bnk_d50**
D50(median) particle size diameter of channel bank sediment (0.001 - 20)
- real *8, dimension(:), allocatable **ch_bed_d50**
D50(median) particle size diameter of channel bed sediment (micrometers) (0.001 - 20)
- real *8, dimension(:), allocatable **ch_cov1**
channel erodibility factor (0.0-1.0) (none)
0 non-erosive channel
1 no resistance to erosion
- real *8, dimension(:), allocatable **ch_cov2**
channel cover factor (0.0-1.0) (none)
0 channel is completely protected from erosion by cover
1 no vegetative cover on channel
- real *8, dimension(:), allocatable **tc_bed**
critical shear stress of channel bed (N/m2)
- real *8, dimension(:), allocatable **tc_bnk**
critical shear stress of channel bank (N/m2)
- integer, dimension(:), allocatable **ch_eqn**
sediment routine methods (DAILY):
0 = original SWAT method
1 = Bagnold's
2 = Kodatie
3 = Molinas WU
4 = Yang
- real *8, dimension(:), allocatable **chpst_rea**
pesticide reaction coefficient in reach (1/day)
- real *8, dimension(:), allocatable **chpst_vol**
pesticide volatilization coefficient in reach (m/day)

- real *8, dimension(:), allocatable **chpst_conc**
- real *8, dimension(:), allocatable **chpst_koc**
pesticide partition coefficient between water and sediment in reach (m^3/g)
- real *8, dimension(:), allocatable **chpst_rsp**
resuspension velocity in reach for pesticide sorbed to sediment (m/day)
- real *8, dimension(:), allocatable **chpst_stl**
settling velocity in reach for pesticide sorbed to sediment (m/day)
- real *8, dimension(:), allocatable **ch_wdr**
channel width to depth ratio (m/m)
- real *8, dimension(:), allocatable **chpst_mix**
mixing velocity (diffusion/dispersion) for pesticide in reach (m/day)
- real *8, dimension(:), allocatable **sedpst_conc**
initil pesticide concentration in river bed sediment (mg/m^3)
- real *8, dimension(:), allocatable **sedpst_bry**
pesticide burial velocity in river bed sediment (m/day)
- real *8, dimension(:), allocatable **sedpst_rea**
pesticide reaction coefficient in river bed sediment (1/day)
- real *8, dimension(:), allocatable **sedpst_act**
depth of active sediment layer in reach for pesticide (m)
- real *8, dimension(:), allocatable **rch_cbod**
- real *8, dimension(:), allocatable **rch_bactlp**
- real *8, dimension(:), allocatable **chside**
change in horizontal distance per unit vertical distance (0.0 - 5)
0 = for vertical channel bank
5 = for channel bank with gentl side slope
- real *8, dimension(:), allocatable **rs1**
local algal settling rate in reach at 20 deg C (m/day or m/hour)
- real *8, dimension(:), allocatable **rs2**
*benthos source rate for dissolved phosphorus in reach at 20 deg C ($(mg\ disP-P)/(m^2*day)$ or $(mg\ disP-P)/(m^2*hour)$)*
- real *8, dimension(:), allocatable **rs3**
*benthos source rate for ammonia nitrogen in reach at 20 deg C ($(mg\ NH4-N)/(m^2*day)$ or $(mg\ NH4-N)/(m^2*hour)$)*
- real *8, dimension(:), allocatable **rs4**
rate coefficient for organic nitrogen settling in reach at 20 deg C (1/day or 1/hour)
- real *8, dimension(:), allocatable **rs5**
organic phosphorus settling rate in reach at 20 deg C (1/day or 1/hour)
- real *8, dimension(:), allocatable **rk1**
CBOD deoxygenation rate coefficient in reach at 20 deg C (1/day or 1/hour)
- real *8, dimension(:), allocatable **rk2**
reaeration rate in accordance with Fickian diffusion in reach at 20 deg C (1/day or 1/hour)
- real *8, dimension(:), allocatable **rk3**
rate of loss of CBOD due to settling in reach at 20 deg C (1/day or 1/hour)
- real *8, dimension(:), allocatable **rk4**
*sediment oxygen demand rate in reach at 20 deg C ($mg\ O_2/(m^2*day)$ or $mg\ O_2/(m^2*hour)$)*
- real *8, dimension(:), allocatable **rk5**
coliform die-off rate in reach (1/day)
- real *8, dimension(:), allocatable **rs6**
rate coefficient for settling of arbitrary non-conservative constituent in reach (1/day)
- real *8, dimension(:), allocatable **rs7**
*benthos source rate for arbitrary non-conservative constituent in reach ($(mg\ ANC)/(m^2*day)$)*
- real *8, dimension(:), allocatable **bc1**

- rate constant for biological oxidation of NH3 to NO2 in reach at 20 deg C (1/day or 1/hour)*
 - real *8, dimension(:), allocatable [bc2](#)
- rate constant for biological oxidation of NO2 to NO3 in reach at 20 deg C (1/day or 1/hour)*
 - real *8, dimension(:), allocatable [bc3](#)
- rate constant for hydrolysis of organic N to ammonia in reach at 20 deg C (1/day or 1/hour)*
 - real *8, dimension(:), allocatable [bc4](#)
- rate constant for the decay of organic P to dissolved P in reach at 20 deg C (1/day or 1/hour)*
 - real *8, dimension(:), allocatable [rk6](#)
- decay rate for arbitrary non-conservative constituent in reach (1/day)*
 - real *8, dimension(:), allocatable **ammonian**
 - real *8, dimension(:), allocatable **orig_sedpstconc**
 - real *8, dimension(:,:), allocatable [wurch](#)
- average daily water removal from the reach for the month ($10^4 \text{ m}^3/\text{day}$)*
 - integer, dimension(:), allocatable **icanal**
 - integer, dimension(:), allocatable **itb**
 - real *8, dimension(:), allocatable [ch_revap](#)
- revap coeff: this variable controls the amount of water moving from bank storage to the root zone as a result of soil moisture depletion(none)*
 - real *8, dimension(:), allocatable **dep_chan**
 - real *8, dimension(:), allocatable [harg_petco](#)
- coefficient related to radiation used in hargreaves eq (range: 0.0019 - 0.0032)*
 - real *8, dimension(:), allocatable **subfr_nowtr**
 - real *8, dimension(:), allocatable [cncoef_sub](#)
- soil water depletion coefficient used in the new (modified curve number method) same as soil index coeff used in APEX range: 0.5 - 2.0*
 - real *8, dimension(:), allocatable **dr_sub**
 - real *8, dimension(:), allocatable [sub_fr](#)
- fraction of total watershed area contained in subbasin (km2/km2)*
 - real *8, dimension(:), allocatable [sub_sw](#)
- water in soil profile in subbasin (mm H2O)*
 - real *8, dimension(:), allocatable **wcklsp**
 - real *8, dimension(:), allocatable **sub_minp**
 - real *8, dimension(:), allocatable **sub_sumfc**
 - real *8, dimension(:), allocatable **sub_gwno3**
 - real *8, dimension(:), allocatable **sub_gwsolp**
 - real *8, dimension(:), allocatable [co2](#)
- CO2 concentration (ppmv)*
 - real *8, dimension(:), allocatable [sub_km](#)
- area of subbasin in square kilometers (km^2)*
 - real *8, dimension(:), allocatable [wlat](#)
- latitude of weather station used to compile data (degrees)*
 - real *8, dimension(:), allocatable [sub_tc](#)
- time of concentration for subbasin (hour)*
 - real *8, dimension(:), allocatable **sub_pet**
 - real *8, dimension(:), allocatable [welev](#)
- elevation of weather station used to compile weather generator data (m)*
 - real *8, dimension(:), allocatable **sub_orgn**
 - real *8, dimension(:), allocatable **sub_orgp**
 - real *8, dimension(:), allocatable **sub_bd**
 - real *8, dimension(:), allocatable **sub_wtmp**
 - real *8, dimension(:), allocatable **sub_sedpa**
 - real *8, dimension(:), allocatable **sub_sedps**

- real *8, dimension(:), allocatable **daylmn**
shortest daylength occurring during the year (hour)
- real *8, dimension(:), allocatable **sub_minpa**
- real *8, dimension(:), allocatable **sub_minps**
- real *8, dimension(:), allocatable **latcos**
cos(latitude) (none)
- real *8, dimension(:), allocatable **latsin**
sin(latitude) (none)
- real *8, dimension(:), allocatable **phutot**
total potential heat units for year (used when no crop is growing) (heat unit)
- real *8, dimension(:), allocatable **plaps**
precipitation lapse rate: precipitation change due to change in elevation (mm H2O/km)
- real *8, dimension(:), allocatable **tlaps**
temperature lapse rate: temperature change due to change in elevation (deg C/km)
- real *8, dimension(:), allocatable **tmp_an**
average annual air temperature (deg C)
- real *8, dimension(:), allocatable **sub_precip**
amount of water reaching soil surface in subbasin (mm H2O)
- real *8, dimension(:), allocatable **rammo_sub**
atmospheric deposition of ammonium values for entire watershed (mg/l)
- real *8, dimension(:), allocatable **rcn_sub**
atmospheric deposition of nitrate for entire watershed (mg/l)
- real *8, dimension(:), allocatable **pcpdays**
- real *8, dimension(:), allocatable **atmo_day**
- real *8, dimension(:), allocatable **sub_snom**
amount of snow melt in subbasin on day (mm H2O)
- real *8, dimension(:), allocatable **sub_qd**
surface runoff that reaches main channel during day in subbasin (mm H2O)
- real *8, dimension(:), allocatable **sub_sedy**
- real *8, dimension(:), allocatable **sub_tran**
transmission losses on day in subbasin (mm H2O)
- real *8, dimension(:), allocatable **sub_no3**
- real *8, dimension(:), allocatable **sub_latno3**
- real *8, dimension(:,:), allocatable **sub_sftmp**
snowfall temperature for subbasin(:). Mean air temperature at which precip is equally likely to be rain as snow/freezing rain (range: -5.0/5.0) (deg C)
- real *8, dimension(:,:), allocatable **sub_smtmp**
snow melt base temperature for subbasin(:) mean air temperature at which snow melt will occur (range: -5.0/5.0) (deg C)
- real *8, dimension(:,:), allocatable **sub_timp**
snow pack temperature lag factor (0-1) (none)
1 = no lag (snow pack temp=current day air temp) as the lag factor goes to zero, the snow pack's temperature will be less influenced by the current day's air temperature
- real *8, dimension(:), allocatable **sub_tileno3**
- real *8, dimension(:), allocatable **sub_solp**
- real *8, dimension(:), allocatable **sub_subp**
- real *8, dimension(:), allocatable **sub_etday**
- real *8, dimension(:), allocatable **sub_elev**
average elevation of HRU (m)
- real *8, dimension(:), allocatable **sub_wyld**
- real *8, dimension(:), allocatable **sub_surfq**
- real *8, dimension(:), allocatable **qird**

- real *8, dimension(:), allocatable **sub_gwq**
- real *8, dimension(:), allocatable **sub_sep**
- real *8, dimension(:), allocatable **sub_chl**
- real *8, dimension(:), allocatable **sub_cbod**
- real *8, dimension(:), allocatable **sub_dox**
- real *8, dimension(:), allocatable **sub_solpst**
- real *8, dimension(:), allocatable **sub_sorpst**
- real *8, dimension(:), allocatable **sub_yorgn**
- real *8, dimension(:), allocatable **sub_yorgp**
- real *8, dimension(:), allocatable **sub_lat**
latitude of HRU/subbasin (degrees)
- real *8, dimension(:), allocatable **sub_bactp**
- real *8, dimension(:), allocatable **sub_bactlp**
- real *8, dimension(:), allocatable **sub_latq**
- real *8, dimension(:), allocatable **sub_gwq_d**
- real *8, dimension(:), allocatable **sub_tileq**
- real *8, dimension(:), allocatable **sub_vaptile**
- real *8, dimension(:), allocatable **sub_dsan**
- real *8, dimension(:), allocatable **sub_dsil**
- real *8, dimension(:), allocatable **sub_dcla**
- real *8, dimension(:), allocatable **sub_dsag**
- real *8, dimension(:), allocatable **sub_dlag**
- real *8 **vap_tile**
- real *8, dimension(:), allocatable **wnan**
- real *8, dimension(:, :), allocatable **sol_stpwt**
- real *8, dimension(:, :), allocatable **sub_pst**
- real *8, dimension(:, :), allocatable **sub_hhq**
- real *8, dimension(:, :), allocatable **sub_hhwtmp**
- real *8, dimension(:, :), allocatable **huminc**
monthly humidity adjustment. Daily values for relative humidity within the month are raised or lowered by the specified amount (used in climate change studies) (none)
- real *8, dimension(:, :), allocatable **radinc**
monthly solar radiation adjustment. Daily radiation within the month is raised or lowered by the specified amount (used in climate change studies) (MJ/m²)
- real *8, dimension(:, :), allocatable **rfinc**
monthly rainfall adjustment. Daily rainfall within the month is adjusted to the specified percentage of the original value (used in climate change studies)(%)
- real *8, dimension(:, :), allocatable **tmpinc**
monthly temperature adjustment. Daily maximum and minimum temperatures within the month are raised or lowered by the specified amount (used in climate change studies) (deg C)
- real *8, dimension(:), allocatable **ch_k1**
effective hydraulic conductivity of tributary channel alluvium (mm/hr)
- real *8, dimension(:), allocatable **ch_k2**
effective hydraulic conductivity of main channel alluvium (mm/hr)
- real *8, dimension(:, :), allocatable **elevb**
elevation at the center of the band (m)
- real *8, dimension(:, :), allocatable **elevb_fr**
fraction of subbasin area within elevation band (the same fractions should be listed for all HRUs within the subbasin) (none)
- real *8, dimension(:, :), allocatable **wndav**
average wind speed for the month (m/s)
- real *8, dimension(:), allocatable **ch_n1**
Manning's "n" value for the tributary channels (none)
- real *8, dimension(:), allocatable **ch_n2**

- Manning's "n" value for the main channel (none)*
- real *8, dimension(:), allocatable [ch_s1](#)
average slope of tributary channels (m/m)
- real *8, dimension(:), allocatable [ch_s2](#)
average slope of main channel (m/m)
- real *8, dimension(:), allocatable [ch_w1](#)
average width of tributary channels (m)
- real *8, dimension(:), allocatable [ch_w2](#)
average width of main channel (m)
- real *8, dimension(:,:), allocatable [dewpt](#)
average dew point temperature for the month (deg C)
- real *8, dimension(:,:), allocatable [amp_r](#)
average fraction of total daily rainfall occurring in maximum half-hour period for month (none)
- real *8, dimension(:,:), allocatable [solarav](#)
average daily solar radiation for the month (MJ/m²/day)
- real *8, dimension(:,:), allocatable [tmpstdmx](#)
standard deviation for avg monthly maximum air temperature (deg C)
- real *8, dimension(:,:), allocatable [pcf](#)
normalization coefficient for precipitation generated from skewed distribution (none)
- real *8, dimension(:,:), allocatable [tmpmn](#)
avg monthly minimum air temperature (deg C)
- real *8, dimension(:,:), allocatable [tmpmx](#)
avg monthly maximum air temperature (deg C)
- real *8, dimension(:,:), allocatable [tmpstdmn](#)
standard deviation for avg monthly minimum air temperature (deg C)
- real *8, dimension(:,:), allocatable [otmpstdmn](#)
- real *8, dimension(:,:), allocatable [otmpmn](#)
- real *8, dimension(:,:), allocatable [otmpmx](#)
- real *8, dimension(:,:), allocatable [otmpstdmx](#)
- real *8, dimension(:,:), allocatable [ch_erodmo](#)
- real *8, dimension(:,:), allocatable [uh](#)
- real *8, dimension(:,:), allocatable [hqdsave](#)
- real *8, dimension(:,:), allocatable [hsdsave](#)
- real *8, dimension(:,:), allocatable [pr_w1](#)
probability of wet day after dry day in month (none)
- real *8, dimension(:,:), allocatable [pr_w2](#)
probability of wet day after wet day in month (none)
- real *8, dimension(:,:), allocatable [pr_w3](#)
proportion of wet days in the month (none)
- real *8, dimension(:,:), allocatable [pcp_stat](#)
- real *8, dimension(:,:), allocatable [opr_w1](#)
- real *8, dimension(:,:), allocatable [opr_w2](#)
- real *8, dimension(:,:), allocatable [opr_w3](#)
- real *8, dimension(:,:), allocatable [opcp_stat](#)
- integer, dimension(:), allocatable [ireg](#)
precipitation category (none):
1 precipitation ≤ 508 mm/yr
2 precipitation > 508 and ≤ 1016 mm/yr
3 precipitation > 1016 mm/yr
- integer, dimension(:), allocatable [hrutot](#)
number of HRUs in subbasin (none)
- integer, dimension(:), allocatable [hru1](#)

- integer, dimension(:), allocatable [ihgage](#)
HRU relative humidity data code (gage # for relative humidity data used in as HRU) (none)
- integer, dimension(:), allocatable [isgage](#)
HRU solar radiation data code (record # for solar radiation used in HRU) (none)
- integer, dimension(:), allocatable [iwgage](#)
HRU wind speed gage data code (gage # for wind speed data used in HRU) (none)
- integer, dimension(:), allocatable [subgis](#)
GIS code printed to output files (output.sub, .rch) (none)
- integer, dimension(:), allocatable [irgage](#)
subbasin rain gage data code (gage # for rainfall data used in HRU) (none)
- integer, dimension(:), allocatable [itgage](#)
subbasin temp gage data code (gage # for temperature data used in HRU) (none)
- integer, dimension(:), allocatable [irelh](#)
(none) irelh = 0 (dewpoint)
irelh = 1 (relative humidity)
note: inputs > 1.0 (dewpoint)
inputs < 1.0 (relative hum)
- integer, dimension(:), allocatable **fcst_reg**
- real *8, dimension(:,:), allocatable [sol_aorgn](#)
amount of nitrogen stored in the active organic (humic) nitrogen pool (kg N/ha)
- real *8, dimension(:,:), allocatable [sol_fon](#)
amount of nitrogen stored in the fresh organic (residue) pool (kg N/ha)
- real *8, dimension(:,:), allocatable [sol_tmp](#)
average temperature of soil layer on previous day or
daily average temperature of soil layer (deg C)
- real *8, dimension(:,:), allocatable [sol_awc](#)
available water capacity of soil layer (mm H2O/mm soil)
- real *8, dimension(:,:), allocatable [volcr](#)
crack volume for soil layer (mm)
- real *8, dimension(:,:), allocatable [sol_prk](#)
percolation storage array (mm H2O)
- real *8, dimension(:,:), allocatable [pperco_sub](#)
subbasin phosphorus percolation coefficient. Ratio of soluble phosphorus in surface to soluble phosphorus in percolate
- real *8, dimension(:,:), allocatable [sol_stap](#)
amount of phosphorus in the soil layer stored in the stable mineral phosphorus pool (kg P/ha)
- real *8, dimension(:,:), allocatable [conv_wt](#)
factor which converts kg/kg soil to kg/ha (none)
- real *8, dimension(:,:), allocatable [sol_actp](#)
amount of phosphorus stored in the active mineral phosphorus pool (kg P/ha)
- real *8, dimension(:,:), allocatable [sol_solp](#)
soluble P concentration in top soil layer (mg P/kg soil) or
amount of inorganic phosphorus stored in solution. NOTE UNIT CHANGE! (kg P/ha)
- real *8, dimension(:,:), allocatable [crdep](#)
maximum or potential crack volume (mm)
- real *8, dimension(:,:), allocatable [sol_fc](#)
amount of water available to plants in soil layer at field capacity (fc - wp) (mm H2O)
- real *8, dimension(:,:), allocatable [sol_ul](#)
amount of water held in the soil layer at saturation (sat - wp water) (mm H2O)
- real *8, dimension(:,:), allocatable [sol_bd](#)
bulk density of the soil layer in HRU (Mg/m³)
- real *8, dimension(:,:), allocatable [sol_z](#)

- depth to bottom of soil layer (mm)*
- real *8, dimension(:,:), allocatable **sol_st**
amount of water stored in the soil layer on any given day (less wp water) (mm H2O)
- real *8, dimension(:,:), allocatable **sol_up**
water content of soil at -0.033 MPa (field capacity) (mm H2O/mm soil)
- real *8, dimension(:,:), allocatable **sol_clay**
percent clay content in soil material (UNIT CHANGE!) (% or none)
- real *8, dimension(:,:), allocatable **sol_hk**
beta coefficient to calculate hydraulic conductivity (none)
- real *8, dimension(:,:), allocatable **flat**
lateral flow storage array (mm H2O)
- real *8, dimension(:,:), allocatable **sol_nh3**
amount of nitrogen stored in the ammonium pool in soil layer (kg N/ha)
- real *8, dimension(:,:), allocatable **sol_ec**
electrical conductivity of soil layer (dS/m)
- real *8, dimension(:,:), allocatable **sol_orgn**
amount of nitrogen stored in the stable organic N pool. NOTE UNIT CHANGE! (mg N/kg soil or kg N/ha)
- real *8, dimension(:,:), allocatable **sol_por**
total porosity of soil layer expressed as a fraction of the total volume (none)
- real *8, dimension(:,:), allocatable **sol_wp**
water content of soil at -1.5 MPa (wilting point) (mm H2O/mm soil)
- real *8, dimension(:,:), allocatable **sol_orgp**
amount of phosphorus stored in the organic P pool. NOTE UNIT CHANGE! (mg P/kg soil or kg P/ha)
- real *8, dimension(:,:), allocatable **sol_hum**
amount of organic matter in the soil layer classified as humic substances (kg humus/ha)
- real *8, dimension(:,:), allocatable **sol_wpmmm**
water content of soil at -1.5 MPa (wilting point) (mm H2O)
- real *8, dimension(:,:), allocatable **sol_no3**
amount of nitrogen stored in the nitrate pool. This variable is read in as a concentration and converted to kg/ha (this value is read from the .sol file in units of mg/kg) (kg N/ha)
- real *8, dimension(:,:), allocatable **sol_cbn**
percent organic carbon in soil layer (%)
- real *8, dimension(:,:), allocatable **sol_k**
saturated hydraulic conductivity of soil layer (mm/hour)
- real *8, dimension(:,:), allocatable **sol_rsd**
amount of organic matter in the soil layer classified as residue (kg/ha)
- real *8, dimension(:,:), allocatable **sol_fop**
amount of phosphorus stored in the fresh organic (residue) pool (kg P/ha)
- real *8, dimension(:,:), allocatable **sol_rock**
percent of rock fragments in soil layer (%)
- real *8, dimension(:,:), allocatable **sol_silt**
percent silt content in soil material (UNIT CHANGE!) (% or none)
- real *8, dimension(:,:), allocatable **sol_sand**
percent sand content of soil material (%)
- real *8, dimension(:,:), allocatable **orig_solno3**
- real *8, dimension(:,:), allocatable **orig_solorgn**
- real *8, dimension(:,:), allocatable **orig_solsoip**
- real *8, dimension(:,:), allocatable **orig_solorgp**
- real *8, dimension(:,:), allocatable **orig_soltmp**
- real *8, dimension(:,:), allocatable **orig_solrsd**
- real *8, dimension(:,:), allocatable **orig_solfop**

- real *8, dimension(:,:), allocatable **orig_solfon**
- real *8, dimension(:,:), allocatable **orig_solaorgn**
- real *8, dimension(:,:), allocatable **orig_solst**
- real *8, dimension(:,:), allocatable **orig_solactp**
- real *8, dimension(:,:), allocatable **orig_solstap**
- real *8, dimension(:,:), allocatable **orig_volcr**
- real *8, dimension(:,:), allocatable **conk**
- real *8, dimension(:,:), allocatable **sol_pst**
sol_pst(:,1) initial amount of pesticide in first layer read in from .chm file (mg/kg)
sol_pst(:,,:) amount of pesticide in layer. NOTE UNIT CHANGE! (kg/ha)
- real *8, dimension(:,:), allocatable **sol_kp**
pesticide sorption coefficient, Kp; the ratio of the concentration in the solid phase to the concentration in solution ((mg/kg)/(mg/L))
- real *8, dimension(:,:), allocatable **orig_solpst**
- real *8, dimension(:), allocatable **velsetlr**
- real *8, dimension(:), allocatable **velsetlp**
- real *8, dimension(:), allocatable **br1**
1st shape parameter for reservoir surface area equation (none)
- real *8, dimension(:), allocatable **evrsv**
lake evaporation coefficient (none)
- real *8, dimension(:), allocatable **res_k**
hydraulic conductivity of the reservoir bottom (mm/hr)
- real *8, dimension(:), allocatable **lkpst_conc**
pesticide concentration in lake water (mg/m³)
- real *8, dimension(:), allocatable **res_evol**
volume of water needed to fill the reservoir to the emergency spillway (read in as 10⁴ m³ and converted to m³) (m³)
- real *8, dimension(:), allocatable **res_pvol**
volume of water needed to fill the reservoir to the principal spillway (read in as 10⁴ m³ and converted to m³) (m³)
- real *8, dimension(:), allocatable **res_vol**
reservoir volume (read in as 10⁴ m³ and converted to m³) (m³)
- real *8, dimension(:), allocatable **res_psa**
reservoir surface area when reservoir is filled to principal spillway (ha)
- real *8, dimension(:), allocatable **lkpst_rea**
pesticide reaction coefficient in lake water (1/day)
- real *8, dimension(:), allocatable **lkpst_vol**
pesticide volatilization coefficient in lake water (m/day)
- real *8, dimension(:), allocatable **br2**
2nd shape parameter for reservoir surface area equation (none)
- real *8, dimension(:), allocatable **res_rr**
average daily principal spillway release volume (read in as a release rate in m³/s and converted to m³/day) (m³/day)
- real *8, dimension(:), allocatable **res_sed**
amount of sediment in reservoir (read in as mg/L and converted to kg/L) (kg/L)
- real *8, dimension(:), allocatable **lkpst_koc**
pesticide partition coefficient between water and sediment in lake water (m³/g)
- real *8, dimension(:), allocatable **lkpst_mix**
mixing velocity (diffusion/dispersion) in lake water for pesticide (m/day)
- real *8, dimension(:), allocatable **lkpst_rsp**
resuspension velocity in lake water for pesticide sorbed to sediment (m/day)
- real *8, dimension(:), allocatable **lkpst_stl**

- settling velocity in lake water for pesticide sorbed to sediment (m/day)*

 - real *8, dimension(:), allocatable [lkspst_conc](#)
- pesticide concentration in lake bed sediment (mg/m³)*

 - real *8, dimension(:), allocatable [lkspst_rea](#)
- pesticide reaction coefficient in lake bed sediment (1/day)*

 - real *8, dimension(:), allocatable **theta_n**
 - real *8, dimension(:), allocatable **theta_p**
 - real *8, dimension(:), allocatable **con_nirr**
 - real *8, dimension(:), allocatable **con_pirr**
 - real *8, dimension(:), allocatable [lkspst_act](#)
- depth of active sediment layer in lake for for pesticide (m)*

 - real *8, dimension(:), allocatable [lkspst_bry](#)
- pesticide burial velocity in lake bed sediment (m/day)*

 - real *8, dimension(:), allocatable **sed_stlr**
 - real *8, dimension(7) **resdata**
 - real *8, dimension(:), allocatable [res_nsed](#)
- normal amount of sediment in reservoir (read in as mg/L and convert to kg/L) (kg/L)*

 - real *8, dimension(:), allocatable [wurtmf](#)
- fraction of water removed from the reservoir via WURESN which is returned and becomes flow from the reservoir outlet (none)*

 - real *8, dimension(:), allocatable [chlar](#)
- chlorophyll-a production coefficient for reservoir (none)*

 - real *8, dimension(:), allocatable [res_no3](#)
- amount of nitrate in reservoir (kg N)*

 - real *8, dimension(:), allocatable [res_orgn](#)
- amount of organic N in reservoir (kg N)*

 - real *8, dimension(:), allocatable [res_orgp](#)
- amount of organic P in reservoir (kg P)*

 - real *8, dimension(:), allocatable [res_solp](#)
- amount of soluble P in reservoir (kg P)*

 - real *8, dimension(:), allocatable [res_seci](#)
- secchi-disk depth (m)*

 - real *8, dimension(:), allocatable **res_chla**
 - real *8, dimension(:), allocatable [res_esa](#)
- reservoir surface area when reservoir is filled to emergency spillway (ha)*

 - real *8, dimension(:), allocatable [res_nh3](#)
- amount of ammonia in reservoir (kg N)*

 - real *8, dimension(:), allocatable [res_no2](#)
- amount of nitrite in reservoir (kg N)*

 - real *8, dimension(:), allocatable [seccir](#)
- water clarity coefficient for reservoir (none)*

 - real *8, dimension(:), allocatable **res_bactp**
 - real *8, dimension(:), allocatable **res_bactlp**
 - real *8, dimension(:), allocatable [oflowmn_fps](#)
- minimum reservoir outflow as a fraction of the principal spillway volume (0-1) (fraction)*

 - real *8, dimension(:), allocatable [starg_fps](#)
- target volume as a fraction of the principal spillway volume (.1-5) (fraction)*

 - real *8, dimension(:), allocatable **weirc**
 - real *8, dimension(:), allocatable **weirk**
 - real *8, dimension(:), allocatable **weirw**
 - real *8, dimension(:), allocatable **acoef**
 - real *8, dimension(:), allocatable **bcoef**

- real *8, dimension(:), allocatable **ccoef**
- real *8, dimension(:), allocatable **orig_resvol**
- real *8, dimension(:), allocatable **orig_ressed**
- real *8, dimension(:), allocatable **orig_lkpstconc**
- real *8, dimension(:), allocatable **orig_lkspstconc**
- real *8, dimension(:), allocatable **orig_ressolp**
- real *8, dimension(:), allocatable **orig_resorgp**
- real *8, dimension(:), allocatable **orig_resno3**
- real *8, dimension(:), allocatable **orig_resno2**
- real *8, dimension(:), allocatable **orig_resnh3**
- real *8, dimension(:), allocatable **orig_resorgn**
- real *8, dimension(:,:), allocatable **oflowmn**
minimum daily outflow for the month (read in as m^3/s and converted to m^3/day) (m^3/day)
- real *8, dimension(:,:), allocatable **oflowmx**
maximum daily outflow for the month (read in as m^3/s and converted to m^3/day) (m^3/day)
- real *8, dimension(:,:), allocatable **starg**
monthly target reservoir storage (needed if IRESKO=2) (read in as $10^4 m^3$ and converted to m^3) (m^3)
- real *8, dimension(:), allocatable **psetlr1**
phosphorus settling rate for mid-year period (read in as $m/year$ and converted to m/day) (m/day)
- real *8, dimension(:), allocatable **psetlr2**
phosphorus settling rate for remainder of year (read in as $m/year$ and converted to m/day) (m/day)
- real *8, dimension(:), allocatable **nsetlr1**
nitrogen settling rate for mid-year period (read in as $m/year$ and converted to m/day) (m/day)
- real *8, dimension(:), allocatable **nsetlr2**
nitrogen settling rate for remainder of year (read in as $m/year$ and converted to m/day) (m/day)
- real *8, dimension(:,:), allocatable **wuresn**
average amount of water withdrawn from reservoir each month for consumptive water use (read in as $10^4 m^3$ and converted to m^3) (m^3)
- real *8, dimension(:,:), allocatable **res_out**
measured average daily outflow from the reservoir for the month (needed if IRESKO=1) (read in as m^3/s and converted to m^3/day) (m^3/day)
- integer, dimension(:), allocatable **res_sub**
number of subbasin reservoir is in (weather for the subbasin is used for the reservoir) (none)
- integer, dimension(:), allocatable **ires1**
beginning of mid-year nutrient settling "season" (none)
- integer, dimension(:), allocatable **ires2**
end of mid-year nutrient settling "season" (none)
- integer, dimension(:), allocatable **iresco**
outflow simulation code (none):
0 compute outflow for uncontrolled reservoir with average annual release rate
1 measured monthly outflow
2 simulated controlled outflow-target release
3 measured daily outflow
4 stage/volume/outflow relationship
- integer, dimension(:), allocatable **iyres**
year of the simulation that the reservoir becomes operational (none)
- integer, dimension(:), allocatable **mores**
month the reservoir becomes operational (none)
- integer, dimension(:), allocatable **iflod1r**
beginning month of non-flood season (needed if IRESKO=2) (none)
- integer, dimension(:), allocatable **iflod2r**
ending month of non-flood season (needed if IRESKO=2) (none)
- integer, dimension(:), allocatable **ndtargr**

- number of days to reach target storage from current reservoir storage (needed if IRESCO=2) (days)*

 - real *8, dimension(:), allocatable [ap_ef](#)
application efficiency (0-1) (none)
 - real *8, dimension(:), allocatable [decay_f](#)
exponential of the rate constant for degradation of the pesticide on foliage (none)
 - real *8, dimension(:), allocatable [skoc](#)
soil adsorption coefficient normalized for soil organic carbon content ((mg/kg)/(mg/L))
 - real *8, dimension(:), allocatable [decay_s](#)
exponential of the rate constant for degradation of the pesticide in soil (none)
 - real *8, dimension(:), allocatable [hlife_f](#)
half-life of pesticide on foliage (days)
 - real *8, dimension(:), allocatable [hlife_s](#)
half-life of pesticide in soil (days)
 - real *8, dimension(:), allocatable [pst_wof](#)
fraction of pesticide on foliage which is washed-off by a rainfall event (none)
 - real *8, dimension(:), allocatable [pst_wsol](#)
solubility of chemical in water (mg/L (ppm))
 - real *8, dimension(:), allocatable [irramt](#)
depth of irrigation water applied to HRU (mm H2O)
 - real *8, dimension(:), allocatable **phusw**
 - real *8, dimension(:), allocatable **phusw_nocrop**
 - integer, dimension(:), allocatable [pstflg](#)
flag for types of pesticide used in watershed. Array location is pesticide ID number
0: pesticide not used
1: pesticide used
 - integer, dimension(:), allocatable [nope](#)
sequence number of pesticide in NPNO(:) (none)
 - integer, dimension(:), allocatable **nop**
 - integer, dimension(:), allocatable **yr_skip**
 - integer, dimension(:), allocatable **isweep**
 - integer, dimension(:), allocatable **icrmx**
 - integer, dimension(:), allocatable **nopmx**
 - integer, dimension(:,:), allocatable **mgtop**
 - integer, dimension(:,:), allocatable **idop**
 - integer, dimension(:,:), allocatable **mgt1iop**
 - integer, dimension(:,:), allocatable **mgt2iop**
 - integer, dimension(:,:), allocatable **mgt3iop**
 - real *8, dimension(:,:), allocatable **mgt4op**
 - real *8, dimension(:,:), allocatable **mgt5op**
 - real *8, dimension(:,:), allocatable **mgt6op**
 - real *8, dimension(:,:), allocatable **mgt7op**
 - real *8, dimension(:,:), allocatable **mgt8op**
 - real *8, dimension(:,:), allocatable **mgt9op**
 - real *8, dimension(:,:), allocatable **mgt10iop**
 - real *8, dimension(:,:), allocatable **phu_op**
 - real *8, dimension(:), allocatable [cnlyd](#)
fraction of nitrogen in yield (kg N/kg yield)
 - real *8, dimension(:), allocatable [rsdco_pl](#)
plant residue decomposition coefficient. The fraction of residue which will decompose in a day assuming optimal moisture, temperature, C:N ratio, and C:P ratio (none)
 - real *8, dimension(:), allocatable [wac21](#)
1st shape parameter for radiation use efficiency equation (none)
 - real *8, dimension(:), allocatable [wac22](#)

- 2nd shape parameter for radiation use efficiency equation (none)*

 - real *8, dimension(:), allocatable [alai_min](#)
minimum LAI during winter dormant period (m^2/m^2)
- 1st shape parameter for leaf area development equation (none)*

 - real *8, dimension(:), allocatable [leaf1](#)
- 2nd shape parameter for leaf area development equation (none)*

 - real *8, dimension(:), allocatable [leaf2](#)
- Value of harvest index between 0 and HVSTI which represents the lowest value expected due to water stress ((kg/ha)/(kg/ha))*

 - real *8, dimension(:), allocatable [bio_e](#)
*biomass-energy ratio. The potential (unstressed) growth rate per unit of intercepted photosynthetically active radiation. ((kg/ha)/(MJ/m**2))*
- harvest index: crop yield/aboveground biomass ((kg/ha)/(kg/ha))*

 - real *8, dimension(:), allocatable [hvsti](#)
- minimum temperature for plant growth (deg C)*

 - real *8, dimension(:), allocatable [t_base](#)
- optimal temperature for plant growth (deg C)*

 - real *8, dimension(:), allocatable [t_opt](#)
- maximum canopy height (m)*

 - real *8, dimension(:), allocatable [chtmx](#)
- natural log of USLE_C (the minimum value of the USLE C factor for the land cover) (none)*

 - real *8, dimension(:), allocatable [cvm](#)
- maximum stomatal conductance (m/s)*

 - real *8, dimension(:), allocatable [gsi](#)
- rate of decline in stomatal conductance per unit increase in vapor pressure deficit ((m/s)*(1/kPa))*

 - real *8, dimension(:), allocatable [vpd2](#)
- rate of decline in radiation use efficiency as a function of vapor pressure deficit (none)*

 - real *8, dimension(:), allocatable [wavp](#)
- fraction of leaf/needle biomass that drops during dormancy (for trees only) (none)*

 - real *8, dimension(:), allocatable [bio_leaf](#)
- maximum (potential) leaf area index (none)*

 - real *8, dimension(:), allocatable [blai](#)
- fraction of phosphorus in yield (kg P/kg yield)*

 - real *8, dimension(:), allocatable [cpyld](#)
- fraction of growing season when leaf area declines (none)*

 - real *8, dimension(:), allocatable [dlai](#)
- maximum root depth of plant (m)*

 - real *8, dimension(:), allocatable [rdmx](#)
- 1st shape parameter for plant N uptake equation (none)*

 - real *8, dimension(:), allocatable [bio_n1](#)
- 2nd shape parameter for plant N uptake equation (none)*

 - real *8, dimension(:), allocatable [bio_n2](#)
- 1st shape parameter for plant P uptake equation (none)*

 - real *8, dimension(:), allocatable [bio_p1](#)
- 2st shape parameter for plant P uptake equation (none)*

 - real *8, dimension(:), allocatable [bio_p2](#)
- fraction above ground biomass that dies off at dormancy (fraction)*

 - real *8, dimension(:), allocatable [bm_dieoff](#)
- fraction above ground biomass that dies off at dormancy (fraction)*

 - real *8, dimension(:), allocatable [bm_trees](#)
- fraction above ground biomass that dies off at dormancy (fraction)*

 - real *8, dimension(:), allocatable [ext_coef](#)

- real *8, dimension(:), allocatable [rsr1](#)
initial root to shoot ratio at the beg of growing season
- real *8, dimension(:), allocatable [rsr2](#)
root to shoot ratio at the end of the growing season
- real *8, dimension(:), allocatable [pltnfr1](#)
nitrogen uptake parameter #1: normal fraction of N in crop biomass at emergence (kg N/kg biomass)
- real *8, dimension(:), allocatable [pltnfr2](#)
nitrogen uptake parameter #2: normal fraction of N in crop biomass at 0.5 maturity (kg N/kg biomass)
- real *8, dimension(:), allocatable [pltnfr3](#)
nitrogen uptake parameter #3: normal fraction of N in crop biomass at maturity (kg N/kg biomass)
- real *8, dimension(:), allocatable [pltpfr1](#)
phosphorus uptake parameter #1: normal fraction of P in crop biomass at emergence (kg P/kg biomass)
- real *8, dimension(:), allocatable [pltpfr2](#)
phosphorus uptake parameter #2: normal fraction of P in crop biomass at 0.5 maturity (kg P/kg biomass)
- real *8, dimension(:), allocatable [pltpfr3](#)
phosphorus uptake parameter #3: normal fraction of P in crop biomass at maturity (kg P/kg biomass)
- integer, dimension(:), allocatable [idc](#)
crop/landcover category:
1 warm season annual legume
2 cold season annual legume
3 perennial legume
4 warm season annual
5 cold season annual
6 perennial
7 trees
- integer, dimension(:), allocatable [mat_yrs](#)
- real *8, dimension(:), allocatable [bactpdb](#)
concentration of persistent bacteria in manure (fertilizer) (cfu/g manure)
- real *8, dimension(:), allocatable [fminn](#)
fraction of fertilize/manure that is mineral N (NO3 + NH3) (kg minN/kg fert)
- real *8, dimension(:), allocatable [forgn](#)
fraction of organic N in fertilizer/manure (kg orgN/kg fert)
- real *8, dimension(:), allocatable [forgp](#)
fraction of fertilizer/manure that is organic P (kg orgP/kg fert)
- real *8, dimension(:), allocatable [bactkddb](#)
fraction of bacteria in solution (the remaining fraction is sorbed to soil particles) (none):
1: all bacteria in solution
0: all bacteria sorbed to soil particles
- real *8, dimension(:), allocatable [bactlpdb](#)
concentration of less persistent bacteria in manure (fertilizer) (cfu/g manure)
- real *8, dimension(:), allocatable [fminp](#)
fraction of fertilizer that is mineral P in fertilizer/manure (kg minP/kg fert)
- real *8, dimension(:), allocatable [fnh3n](#)
fraction of mineral N in fertilizer that is NH3-N in fertilizer/manure (kgNH3-N/kgminN)
- character(len=8), dimension(200) [fertnm](#)
name of fertilizer
- real *8, dimension(:), allocatable [curbden](#)
curb length density in HRU (km/ha)
- real *8, dimension(:), allocatable [dirtmx](#)
maximum amount of solids allowed to build up on impervious surfaces (kg/curb km)
- real *8, dimension(:), allocatable [fimp](#)
fraction of HRU area that is impervious (both directly and indirectly connected)(fraction)

- real *8, dimension(:), allocatable **urbcoef**
wash-off coefficient for removal of constituents from an impervious surface (1/mm)
- real *8, dimension(:), allocatable **thalf**
time for the amount of solids on impervious areas to build up to 1/2 the maximum level (days)
- real *8, dimension(:), allocatable **tnconc**
concentration of total nitrogen in suspended solid load from impervious areas (mg N/kg sed)
- real *8, dimension(:), allocatable **tno3conc**
concentration of NO3-N in suspended solid load from impervious areas (mg NO3-N/kg sed)
- real *8, dimension(:), allocatable **tpconc**
concentration of total phosphorus in suspended solid load from impervious areas (mg P/kg sed)
- real *8, dimension(:), allocatable **fcimp**
fraction of HRU area that is classified as directly connected impervious (fraction)
- real *8, dimension(:), allocatable **urbcn2**
SCS curve number for moisture condition II in impervious areas (none)
- real *8 **fr_curb**
availability factor, the fraction of the curb length that is sweepable (none)
- real *8 **frt_kg**
amount of fertilizer applied to HRU (kg/ha)
- real *8 **pst_dep**
depth of pesticide in the soil (mm)
- real *8 **swepeff**
- real *8, dimension(:), allocatable **ranrns_hru**
- integer, dimension(:), allocatable **itill**
- real *8, dimension(:), allocatable **deptil**
depth of mixing caused by tillage operation (mm)
- real *8, dimension(:), allocatable **effmix**
mixing efficiency of tillage operation (none)
- real *8, dimension(:), allocatable **ranrns**
random roughness of a given tillage operation (mm)
- character(len=8), dimension(550) **tillnm**
8-character name for the tillage operation
- real *8, dimension(:), allocatable **rnum1s**
For ICODES equal to (none)
0,1,3,5,9: not used
2: Fraction of flow in channel
4: amount of water transferred (as defined by INUM4S)
7,8,10,11: drainage area in square kilometers associated with the record file
12: reparation coefficient.
- real *8, dimension(:), allocatable **hyd_dakm**
total drainage area of hydrograph in square kilometers (km²)
- real *8, dimension(:,:), allocatable **varoute**
- real *8, dimension(:,:), allocatable **shyd**
- real *8, dimension(:,:), allocatable **vartran**
- real *8, dimension(:,:), allocatable **hhvaroute**
- integer, dimension(:), allocatable **icodes**
routing command code (none):
0 = finish
1 = subbasin
2 = route
3 = routres
4 = transfer
5 = add
6 = rechour
7 = recmon

- 8 = recyear
 - 9 = save
 - 10 = recday
 - 11 = reccnst
 - 12 = structure
 - 13 = apex
 - 14 = saveconc
 - 15 =
- integer, dimension(:), allocatable [ihouts](#)
 - For ICODES equal to (none)*
 - 0: not used
 - 1,2,3,5,7,8,10,11: hydrograph storage location number
 - 4: departure type (1=reach, 2=reservoir)
 - 9: hydrograph storage location of data to be printed to event file
 - 14: hydrograph storage location of data to be printed to saveconc file.
- integer, dimension(:), allocatable [inum1s](#)
 - For ICODES equal to (none)*
 - 0: not used
 - 1: subbasin number
 - 2: reach number
 - 3: reservoir number
 - 4: reach or res # flow is diverted from
 - 5: hydrograph storage location of 1st dataset to be added
 - 7,8,9,10,11,14: file number.
- integer, dimension(:), allocatable [inum2s](#)
 - For ICODES equal to (none)*
 - 0,1,7,8,10,11: not used
 - 2,3: inflow hydrograph storage location
 - 4: destination type (1=reach, 2=reservoir)
 - 5: hydrograph storage location of 2nd dataset to be added
 - 9,14: print frequency (0=daily, 1=hourly)
- integer, dimension(:), allocatable [inum3s](#)
 - For ICODES equal to (none)*
 - 0,1,5,7,8,10,11: not used
 - 2,3: subbasin number 4: destination number. Reach or reservoir receiving water
 - 9: print format (0=normal, fixed format; 1=txt format for AV interface, recday)
- integer, dimension(:), allocatable [inum4s](#)
 - For ICODES equal to (none)*
 - 0,2,3,5,7,8,9,10,11: not used
 - 1: GIS code printed to output file (optional)
 - 4: rule code governing transfer of water (1=fraction transferred out, 2=min volume or flow left, 3=exact amount transferred)
- integer, dimension(:), allocatable **inum5s**
- integer, dimension(:), allocatable **inum6s**
- integer, dimension(:), allocatable **inum7s**
- integer, dimension(:), allocatable **inum8s**
- integer, dimension(:), allocatable **subed**
- character(len=10), dimension(:), allocatable **recmonps**
- character(len=10), dimension(:), allocatable **reccnstps**
- character(len=5), dimension(:), allocatable **subnum**
- character(len=4), dimension(:), allocatable **hruno**
- real *8, dimension(:), allocatable [grwat_n](#)
 - Mannings's n for grassed waterway (none)*
- integer, dimension(:), allocatable [grwat_i](#)
 - flag for the simulation of grass waterways (none)*
 - = 0 inactive
 - = 1 active
- real *8, dimension(:), allocatable [grwat_l](#)

- length of grass waterway (km)*
- real *8, dimension(:), allocatable [grwat_w](#)
- average width of grassed waterway (m)*
- real *8, dimension(:), allocatable [grwat_d](#)
- depth of grassed waterway from top of bank to bottom (m)*
- real *8, dimension(:), allocatable [grwat_s](#)
- average slope of grassed waterway channel (m)*
- real *8, dimension(:), allocatable [grwat_spcon](#)
- linear parameter for calculating sediment in grassed waterways (none)*
- real *8, dimension(:), allocatable **tc_gwat**
- real *8, dimension(:), allocatable **pot_volmm**
- real *8, dimension(:), allocatable **pot_tilemm**
- real *8, dimension(:), allocatable **pot_volxmm**
- real *8, dimension(:), allocatable [pot_fr](#)
- fraction of HRU area that drains into pothole (km^2/km^2)*
- real *8, dimension(:), allocatable [pot_tile](#)
- average daily outflow to main channel from tile flow if drainage tiles are installed in pothole (needed only if current HRU is IPOT) (m^3/s)*
- real *8, dimension(:), allocatable [pot_vol](#)
- initial or current volume of water stored in the depression/impounded area (read in as mm and converted to m^3) (needed only if current HRU is IPOT) (mm or $\text{m}^3 \text{H2O}$)*
- real *8, dimension(:), allocatable [potsa](#)
- surface area of impounded water body (ha)*
- real *8, dimension(:), allocatable [pot_volx](#)
- maximum volume of water stored in the depression/impounded area (read in as mm and converted to m^3) (needed only if current HRU is IPOT) (mm)*
- real *8, dimension(:), allocatable [wfs](#)
- wetting front matric potential (average capillary suction at wetting front) (mm)*
- real *8, dimension(:), allocatable **potfliwi**
- real *8, dimension(:), allocatable **potsedi**
- real *8, dimension(:), allocatable [pot_no3l](#)
- nitrate decay rate in impounded area (1/day)*
- real *8, dimension(:), allocatable [pot_nsed](#)
- normal sediment concentration in impounded water (needed only if current HRU is IPOT)(mg/L)*
- real *8, dimension(:), allocatable [gwno3](#)
- nitrate-N concentration in groundwater loading to reach (mg N/L)*
- real *8, dimension(:), allocatable [newrti](#)
- infiltration rate for last time step from the previous day (mm/hr)*
- real *8, dimension(:), allocatable [fsred](#)
- reduction in bacteria loading from filter strip (none)*
- real *8, dimension(:), allocatable **pot_sed**
- real *8, dimension(:), allocatable **pot_no3**
- real *8, dimension(:), allocatable **tmpavp**
- real *8, dimension(:), allocatable [dis_stream](#)
- average distance to stream (m)*
- real *8, dimension(:), allocatable [evpot](#)
- pothole evaporation coefficient (none)*
- real *8, dimension(:), allocatable **pot_solpl**
- real *8, dimension(:), allocatable **sed_con**
- real *8, dimension(:), allocatable **orgn_con**
- real *8, dimension(:), allocatable **orgp_con**
- real *8, dimension(:), allocatable [pot_k](#)

- hydraulic conductivity of soil surface of pothole defaults to conductivity of upper soil (0.↔ 01-10.) layer*
- real *8, dimension(:), allocatable **soln_con**
 - real *8, dimension(:), allocatable **solp_con**
 - real *8, dimension(:), allocatable **n_reduc**
nitrogen uptake reduction factor (not currently used; defaulted 300.)
 - real *8, dimension(:), allocatable **n_lag**
lag coefficient for calculating nitrate concentration in subsurface drains (0.001 - 1.0) (dimensionless)
 - real *8, dimension(:), allocatable **n_ln**
power function exponent for calculating nitrate concentration in subsurface drains (1.0 - 3.0) (dimensionless)
 - real *8, dimension(:), allocatable **n_lnc**
coefficient for power function for calculating nitrate concentration in subsurface drains (0.5 - 4.0) (dimensionless)
 - integer, dimension(:), allocatable **ioper**
 - integer, dimension(:), allocatable **ngrwat**
 - real *8, dimension(:), allocatable **usle_ls**
USLE equation length slope (LS) factor (none)
 - real *8, dimension(:), allocatable **filterw**
filter strip width for bacteria transport (m)
 - real *8, dimension(:), allocatable **phuacc**
fraction of plant heat units accumulated (none)
 - real *8, dimension(:), allocatable **sumix**
sum of all tillage mixing efficiencies for HRU operation (none)
 - real *8, dimension(:), allocatable **epco**
plant water uptake compensation factor (0-1) (none)
 - real *8, dimension(:), allocatable **esco**
soil evaporation compensation factor (0-1) (none)
 - real *8, dimension(:), allocatable **hru_slp**
average slope steepness (m/m)
 - real *8, dimension(:), allocatable **slsubbsn**
average slope length for subbasin (m)
 - real *8, dimension(:), allocatable **erorgn**
organic N enrichment ratio, if left blank the model will calculate for every event (none)
 - real *8, dimension(:), allocatable **erorgp**
organic P enrichment ratio, if left blank the model will calculate for every event (none)
 - real *8, dimension(:), allocatable **biomix**
biological mixing efficiency. Mixing of soil due to activity of earthworms and other soil biota. Mixing is performed at the end of every calendar year (none)
 - real *8, dimension(:), allocatable **pnd_seci**
secchi-disk depth of pond (m)
 - real *8, dimension(:), allocatable **canmx**
maximum canopy storage (mm H2O)
 - real *8, dimension(:), allocatable **divmax**
maximum daily irrigation diversion from the reach (when IRRSC=1 or IRR=3): when value is positive the units are mm H2O; when the value is negative, the units are (10⁴ m³ H2O) (mm H2O or 10⁴ m³ H2O)
 - real *8, dimension(:), allocatable **flowmin**
minimum instream flow for irrigation diversions when IRRSC=1, irrigation water will be diverted only when streamflow is at or above FLOWMIN (m³/s)
 - real *8, dimension(:), allocatable **usle_p**
USLE equation support practice (P) factor (none)
 - real *8, dimension(:), allocatable **lat_sed**
sediment concentration in lateral flow (g/L)
 - real *8, dimension(:), allocatable **rch_dakm**

- total drainage area contributing to flow at the outlet (pour point) of the reach in square kilometers (km^2)*

 - real *8, dimension(:), allocatable [cn1](#)
- SCS runoff curve number for moisture condition I (none)*

 - real *8, dimension(:), allocatable [pnd_no3s](#)
- amount of nitrate originating from lateral flow in pond at end of day (kg N)*

 - real *8, dimension(:), allocatable [lat_ttime](#)
- lateral flow travel time or exponential of the lateral flow travel time (days or none)*

 - real *8, dimension(:), allocatable [cn2](#)
- SCS runoff curve number for moisture condition II (none)*

 - real *8, dimension(:), allocatable [flowfr](#)
- fraction of available flow in reach that is allowed to be applied to the HRU (none)*

 - real *8, dimension(:), allocatable [sol_zmx](#)
- maximum rooting depth (mm)*

 - real *8, dimension(:), allocatable [tile_ttime](#)
- exponential of the tile flow travel time (none)*

 - real *8, dimension(:), allocatable [slsoil](#)
- slope length for lateral subsurface flow (m)*

 - real *8, dimension(:), allocatable [gwminp](#)
- soluble P concentration in groundwater loading to reach (mg P/L)*

 - real *8, dimension(:), allocatable [sol_cov](#)
- amount of residue on soil surface (kg/ha)*

 - real *8, dimension(:), allocatable [sed_stl](#)
- fraction of sediment remaining suspended in impoundment after settling for one day (kg/kg)*

 - real *8, dimension(:), allocatable [ov_n](#)
- Manning's "n" value for overland flow (none)*

 - real *8, dimension(:), allocatable [pnd_no3](#)
- amount of nitrate originating from surface runoff in pond at end of day (kg N)*

 - real *8, dimension(:), allocatable [pnd_solp](#)
- amount of soluble P originating from surface runoff in pond at end of day (kg P)*

 - real *8, dimension(:), allocatable [yldanu](#)
- annual yield (dry weight) in the HRU (metric tons/ha)*

 - real *8, dimension(:), allocatable [driftco](#)
- coefficient for pesticide drift directly onto stream (none)*

 - real *8, dimension(:), allocatable [pnd_orgn](#)
- amount of organic N originating from surface runoff in pond at end of day (kg N)*

 - real *8, dimension(:), allocatable [pnd_orgp](#)
- amount of organic P originating from surface runoff in pond at end of day (kg P)*

 - real *8, dimension(:), allocatable [cn3](#)
- SCS runoff curve number for moisture condition III (none)*

 - real *8, dimension(:), allocatable [twlpnd](#)
- water lost through seepage from ponds on day in HRU (mm H₂O)*

 - real *8, dimension(:), allocatable [twlwet](#)
- water lost through seepage from wetlands on day in HRU (mm H₂O)*

 - real *8, dimension(:), allocatable [hru_fr](#)
- fraction of subbasin area contained in HRU (km^2/km^2)*

 - real *8, dimension(:), allocatable [sol_sumul](#)
- amount of water held in soil profile at saturation (mm H₂O)*

 - real *8, dimension(:), allocatable [pnd_chla](#)
- amount of chlorophyll-a in pond at end of day (kg chl_a)*

 - real *8, dimension(:), allocatable [hru_km](#)
- area of HRU in square kilometers (km^2)*

- real *8, dimension(:), allocatable [bio_ms](#)
land cover/crop biomass (dry weight) (kg/ha)
- real *8, dimension(:), allocatable [sol_alb](#)
albedo when soil is moist (none)
- real *8, dimension(:), allocatable [strsw](#)
fraction of potential plant growth achieved on the day where the reduction is caused by water stress (none)
- real *8, dimension(:), allocatable [pnd_fr](#)
fraction of HRU/subbasin area that drains into ponds (none)
- real *8, dimension(:), allocatable [pnd_k](#)
hydraulic conductivity through bottom of ponds (mm/hr)
- real *8, dimension(:), allocatable [pnd_psa](#)
surface area of ponds when filled to principal spillway (ha)
- real *8, dimension(:), allocatable [pnd_pvol](#)
runoff volume from catchment area needed to fill the ponds to the principal spillway (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)
- real *8, dimension(:), allocatable [pnd_esa](#)
surface area of ponds when filled to emergency spillway (ha)
- real *8, dimension(:), allocatable [pnd_evol](#)
runoff volume from catchment area needed to fill the ponds to the emergency spillway (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)
- real *8, dimension(:), allocatable [pnd_vol](#)
volume of water in ponds (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)
- real *8, dimension(:), allocatable [yldaa](#)
average annual yield (dry weight) in the HRU (metric tons)
- real *8, dimension(:), allocatable [pnd_nsed](#)
normal sediment concentration in pond water (UNIT CHANGE!) (mg/kg or kg/kg)
- real *8, dimension(:), allocatable [pnd_sed](#)
sediment concentration in pond water (UNIT CHANGE!) (mg/kg or kg/kg)
- real *8, dimension(:), allocatable [strsa](#)
- real *8, dimension(:), allocatable [dep_imp](#)
- real *8, dimension(:), allocatable [evpnd](#)
- real *8, dimension(:), allocatable [evwet](#)
- real *8, dimension(:), allocatable [wet_fr](#)
fraction of HRU/subbasin area that drains into wetlands (none)
- real *8, dimension(:), allocatable [wet_k](#)
hydraulic conductivity of bottom of wetlands (mm/hr)
- real *8, dimension(:), allocatable [wet_nsa](#)
surface area of wetlands in subbasin at normal water level (ha)
- real *8, dimension(:), allocatable [wet_nvol](#)
runoff volume from catchment area needed to fill wetlands to normal water level (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)
- integer, dimension(:), allocatable [iwetgw](#)
- integer, dimension(:), allocatable [iwetile](#)
- real *8, dimension(:), allocatable [wet_mxsa](#)
surface area of wetlands at maximum water level (ha)
- real *8, dimension(:), allocatable [wet_mxvol](#)
runoff volume from catchment area needed to fill wetlands to maximum water level (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)
- real *8, dimension(:), allocatable [wet_vol](#)
volume of water in wetlands (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)
- real *8, dimension(:), allocatable [wet_nsed](#)
normal sediment concentration in wetland water (UNIT CHANGE!) (mg/kg or kg/kg)

- real *8, dimension(:), allocatable [wet_sed](#)
sediment concentration in wetland water (UNIT CHANGE!) (mg/L or kg/L)
- real *8, dimension(:), allocatable [bp1](#)
1st shape parameter for pond surface area equation (none)
- real *8, dimension(:), allocatable [bp2](#)
2nd shape parameter for the pond surface area equation (none)
- real *8, dimension(:), allocatable [sci](#)
retention coefficient for CN method based on plant ET (none)
- real *8, dimension(:), allocatable [smx](#)
retention coefficient for CN method based on soil moisture (none)
- real *8, dimension(:), allocatable [bw1](#)
1st shape parameter for the wetland surface area equation (none)
- real *8, dimension(:), allocatable [bw2](#)
2nd shape parameter for the wetland surface area equation (none)
- real *8, dimension(:), allocatable [bactpq](#)
persistent bacteria in soil solution (# cfu/m²)
- real *8, dimension(:), allocatable [cnday](#)
curve number for current day, HRU and at current soil moisture (none)
- real *8, dimension(:), allocatable [bactlp_plt](#)
less persistent bacteria on foliage (# cfu/m²)
- real *8, dimension(:), allocatable [bactp_plt](#)
persistent bacteria on foliage (# cfu/m²)
- real *8, dimension(:), allocatable [auto_eff](#)
fertilizer application efficiency calculated as the amount of N applied divided by the amount of N removed at harvest (none)
- real *8, dimension(:), allocatable [secciw](#)
water clarity coefficient for wetland (none)
- real *8, dimension(:), allocatable [sol_sw](#)
amount of water stored in soil profile on current day (mm H₂O)
- real *8, dimension(:), allocatable [bactlpq](#)
less persistent bacteria in soil solution (# cfu/m²)
- real *8, dimension(:), allocatable [chlaw](#)
chlorophyll-a production coefficient for wetland (none)
- real *8, dimension(:), allocatable [tmpav](#)
average air temperature on current day in HRU (deg C)
- real *8, dimension(:), allocatable [bactlps](#)
less persistent bacteria attached to soil particles (# cfu/m²)
- real *8, dimension(:), allocatable [bactps](#)
persistent bacteria attached to soil particles (# cfu/m²)
- real *8, dimension(:), allocatable [sno_hru](#)
amount of water stored as snow in HRU on current day (mm H₂O)
- real *8, dimension(:), allocatable [wet_orn](#)
amount of organic N originating from surface runoff in wetland at end of day (kg N)
- real *8, dimension(:), allocatable [hru_ra](#)
solar radiation for the day in HRU (MJ/m²)
- real *8, dimension(:), allocatable [subp](#)
precipitation for the day in HRU (mm H₂O)
- real *8, dimension(:), allocatable [rsdin](#)
initial residue cover (kg/ha)
- real *8, dimension(:), allocatable [tmn](#)
minimum air temperature on current day in HRU (deg C)

- real *8, dimension(:), allocatable [tmx](#)
maximum air temperature on current day in HRU (deg C)
- real *8, dimension(:), allocatable [tmp_hi](#)
- real *8, dimension(:), allocatable [tmp_lo](#)
- real *8, dimension(:), allocatable [usle_k](#)
USLE equation soil erodibility (K) factor (none)
- real *8, dimension(:), allocatable [tconc](#)
time of concentration for HRU (hour)
- real *8, dimension(:), allocatable [hru_rmx](#)
maximum possible solar radiation for the day in HRU (MJ/m²)
- real *8, dimension(:), allocatable [rwt](#)
fraction of total plant biomass that is in roots (none)
- real *8, dimension(:), allocatable [olai](#)
- real *8, dimension(:), allocatable [usle_cf](#)
- real *8, dimension(:), allocatable [usle_eif](#)
- real *8, dimension(:), allocatable [sol_sumfc](#)
amount of water held in soil profile at field capacity (mm H₂O)
- real *8, dimension(:), allocatable [t_ov](#)
time for flow from farthest point in subbasin to enter a channel (hour)
- real *8, dimension(:), allocatable [anano3](#)
total amount of NO₃ applied during the year in auto-fertilization (kg N/ha)
- real *8, dimension(:), allocatable [aird](#)
amount of water applied to HRU on current day (mm H₂O)
- real *8, dimension(:), allocatable [wet_orgp](#)
amount of organic P originating from surface runoff in wetland at end of day (kg P)
- real *8, dimension(:), allocatable [sol_avpor](#)
average porosity for entire soil profile (none)
- real *8, dimension(:), allocatable [usle_mult](#)
product of USLE K,P,LS,exp(rock) (none)
- real *8, dimension(:), allocatable [rhd](#)
relative humidity for the day in HRU (none)
- real *8, dimension(:), allocatable [u10](#)
wind speed (measured at 10 meters above surface) for the day in HRU (m/s)
- real *8, dimension(:), allocatable [cht](#)
canopy height (m)
- real *8, dimension(:), allocatable [aairr](#)
average annual amount of irrigation water applied to HRU (mm H₂O)
- real *8, dimension(:), allocatable [lai_aamx](#)
maximum leaf area index for the entire period of simulation in the HRU (none)
- real *8, dimension(:), allocatable [deepirr](#)
amount of water removed from deep aquifer for irrigation (mm H₂O)
- real *8, dimension(:), allocatable [shallirr](#)
amount of water removed from shallow aquifer for irrigation (mm H₂O)
- real *8, dimension(:), allocatable [ch_l1](#)
longest tributary channel length in subbasin (km)
- real *8, dimension(:), allocatable [wet_no3](#)
amount of nitrate originating from surface runoff in wetland at end of day (kg N)
- real *8, dimension(:), allocatable [ovrlnd](#)
overland flow onto HRU from upstream routing unit (mm H₂O)
- real *8, dimension(:), allocatable [canstor](#)
amount of water held in canopy storage (mm H₂O)

- real *8, dimension(:), allocatable **irr_mx**
maximum irrigation amount per auto application (mm)
- real *8, dimension(:), allocatable **auto_wstr**
water stress factor which triggers auto irrigation (none or mm)
- real *8, dimension(:), allocatable **cfrt_id**
fertilizer/manure id number from database (none)
- real *8, dimension(:), allocatable **cfrt_kg**
amount of fertilzier applied to HRU on a given day (kg/ha)
- real *8, dimension(:), allocatable **cpst_id**
- real *8, dimension(:), allocatable **cpst_kg**
- real *8, dimension(:), allocatable **irr_asq**
surface runoff ratio
- real *8, dimension(:), allocatable **irr_eff**
- real *8, dimension(:), allocatable **irrsq**
surface runoff ratio (0-1) .1 is 10% surface runoff (frac)
- real *8, dimension(:), allocatable **irrsalt**
concentration of salt in irrigation water (mg/kg)
- real *8, dimension(:), allocatable **irrefm**
- real *8, dimension(:), allocatable **bio_eat**
dry weight of biomass removed by grazing daily ((kg/ha)/day)
- real *8, dimension(:), allocatable **bio_trmp**
dry weight of biomass removed by trampling daily ((kg/ha)/day)
- integer, dimension(:), allocatable **ipst_freq**
number of days between applications (days)
- integer, dimension(:), allocatable **ifrt_freq**
- integer, dimension(:), allocatable **irr_noa**
- integer, dimension(:), allocatable **irr_sc**
- integer, dimension(:), allocatable **irr_no**
- integer, dimension(:), allocatable **imp_trig**
release/impound action code (none):
0 begin impounding water
1 release impounded water
- integer, dimension(:), allocatable **fert_days**
- integer, dimension(:), allocatable **irr_sca**
- integer, dimension(:), allocatable **idplt**
land cover/crop identification code for first crop grown in HRU (the only crop if there is no rotation) (from crop.dat)
(none)
- integer, dimension(:), allocatable **pest_days**
- integer, dimension(:), allocatable **wstrs_id**
- real *8, dimension(:,:), allocatable **bio_aahv**
- real *8, dimension(:), allocatable **cumei**
- real *8, dimension(:), allocatable **cumeira**
- real *8, dimension(:), allocatable **cumrt**
- real *8, dimension(:), allocatable **cumrai**
- real *8, dimension(:), allocatable **wet_solp**
amount of soluble P originating from surface runoff in wetland at end of day (kg P)
- real *8, dimension(:), allocatable **wet_chla**
amount of chlorophyll-a in wetland at end of day (kg chla)
- real *8, dimension(:), allocatable **wet_no3s**
amount of nitrate originating from lateral flow in wetland at end of day (kg N)
- real *8, dimension(:), allocatable **pstsol**
soluble pesticide leached from bottom of soil profile (kg pst/ha)

- real *8, dimension(:), allocatable [pnd_no3g](#)
amount of nitrate originating from groundwater in pond at end of day (kg N)
- real *8, dimension(:), allocatable [wet_seci](#)
secchi-disk depth in wetland at end of day (m)
- real *8, dimension(:), allocatable [delay](#)
groundwater delay: time required for water leaving the bottom of the root zone to reach the shallow aquifer (days)
- real *8, dimension(:), allocatable [gwht](#)
groundwater height (m)
- real *8, dimension(:), allocatable [gw_q](#)
groundwater contribution to streamflow from HRU on current day (mm H2O)
- real *8, dimension(:), allocatable [pnd_solpg](#)
amount of soluble P originating from groundwater in pond at end of day (kg P)
- real *8, dimension(:), allocatable [alpha_bf](#)
alpha factor for groundwater recession curve (1/days)
- real *8, dimension(:), allocatable [alpha_bfe](#)
 $\exp(-\alpha_{bf})$ (none)
- real *8, dimension(:), allocatable [gw_spyld](#)
specific yield for shallow aquifer (m^3/m^3)
- real *8, dimension(:), allocatable [alpha_bf_d](#)
alpha factor for groundwater recession curve of the deep aquifer (1/days)
- real *8, dimension(:), allocatable [alpha_bfe_d](#)
 $\exp(-\alpha_{bf_d})$ for deep aquifer (none)
- real *8, dimension(:), allocatable [gw_qdeep](#)
- real *8, dimension(:), allocatable [gw_delaye](#)
 $\exp(-1/delay)$ (none)
- real *8, dimension(:), allocatable [gw_revap](#)
revap coeff: this variable controls the amount of water moving from the shallow aquifer to the root zone as a result of soil moisture depletion (none)
- real *8, dimension(:), allocatable [rchrg_dp](#)
recharge to deep aquifer: the fraction of root zone percolation that reaches the deep aquifer (none)
- real *8, dimension(:), allocatable [anion_excl](#)
fraction of porosity from which anions are excluded
- real *8, dimension(:), allocatable [revapmn](#)
threshold depth of water in shallow aquifer required to allow revap to occur (mm H2O)
- real *8, dimension(:), allocatable [rchrg](#)
- real *8, dimension(:), allocatable [bio_min](#)
minimum plant biomass for grazing (kg/ha)
- real *8, dimension(:), allocatable [ffc](#)
initial HRU soil water content expressed as fraction of field capacity (none)
- real *8, dimension(:), allocatable [surqsolp](#)
- real *8, dimension(:), allocatable [deepst](#)
depth of water in deep aquifer (mm H2O)
- real *8, dimension(:), allocatable [shallst](#)
depth of water in shallow aquifer in HRU (mm H2O)
- real *8, dimension(:), allocatable [wet_solpg](#)
amount of soluble P originating from groundwater in wetland at end of day (kg P)
- real *8, dimension(:), allocatable [cklsp](#)
- real *8, dimension(:), allocatable [rchrg_src](#)
- real *8, dimension(:), allocatable [trapeff](#)
filter strip trapping efficiency (used for everything but bacteria) (none)
- real *8, dimension(:), allocatable [sol_avbd](#)

- *average bulk density for soil profile (Mg/m^3)*
- real *8, dimension(:), allocatable [wet_no3g](#)
- *amount of nitrate originating from groundwater in wetland at end of day (kg N)*
- real *8, dimension(:), allocatable [tdrain](#)
- *time to drain soil to field capacity yield used in autofertilization (hours)*
- real *8, dimension(:), allocatable [gwqmn](#)
- *threshold depth of water in shallow aquifer required before groundwater flow will occur (mm H₂O)*
- real *8, dimension(:), allocatable [snotmp](#)
- *temperature of snow pack in HRU (deg C)*
- real *8, dimension(:), allocatable **ppInt**
- real *8, dimension(:), allocatable [gdrain](#)
- *drain tile lag time: the amount of time between the transfer of water from the soil to the drain tile and the release of the water from the drain tile to the reach (hours)*
- real *8, dimension(:), allocatable [ddrain](#)
- *depth to the sub-surface drain (mm)*
- real *8, dimension(:), allocatable [sol_crk](#)
- *crack volume potential of soil (none)*
- real *8, dimension(:), allocatable [brt](#)
- *fraction of surface runoff within the subbasin which takes 1 day or less to reach the subbasin outlet (none)*
- real *8, dimension(:), allocatable [dayl](#)
- *day length (hours)*
- real *8, dimension(:), allocatable [sstmaxd](#)
- *static maximum depressional storage; read from .sdr (mm)*
- real *8, dimension(:), allocatable [re](#)
- *effective radius of drains (mm)*
- real *8, dimension(:), allocatable [sdrain](#)
- *distance between two drain tubes or tiles (mm)*
- real *8, dimension(:), allocatable **ddrain_hru**
- real *8, dimension(:), allocatable [drain_co](#)
- *drainage coefficient (mm/day)*
- real *8, dimension(:), allocatable [latksatf](#)
- *multiplication factor to determine $\text{conk}(j1,j)$ from $\text{sol_k}(j1,j)$ for HRU (none)*
- real *8, dimension(:), allocatable [pc](#)
- *pump capacity (default pump capacity = 1.042mm/hr or 25mm/day) (mm/hr)*
- real *8, dimension(:), allocatable **stmaxd**
- real *8, dimension(:), allocatable [rnd3](#)
- *random number between 0.0 and 1.0 (none)*
- real *8, dimension(:), allocatable [rnd2](#)
- *random number between 0.0 and 1.0 (none)*
- real *8, dimension(:), allocatable **twash**
- real *8, dimension(:), allocatable [sol_cnsu](#)
- real *8, dimension(:), allocatable **doxq**
- real *8, dimension(:), allocatable [rnd8](#)
- *random number between 0.0 and 1.0 (none)*
- real *8, dimension(:), allocatable [rnd9](#)
- *random number between 0.0 and 1.0 (none)*
- real *8, dimension(:), allocatable **percn**
- real *8, dimension(:), allocatable [sol_sumwp](#)
- real *8, dimension(:), allocatable [qdr](#)
- *total amount of water entering main channel for day from HRU (mm H₂O)*
- real *8, dimension(:), allocatable **tauton**
- real *8, dimension(:), allocatable **tautop**

- real *8, dimension(:), allocatable **cbodu**
- real *8, dimension(:), allocatable **chl_a**
- real *8, dimension(:), allocatable **tfertn**
- real *8, dimension(:), allocatable **tfertp**
- real *8, dimension(:), allocatable **tgrazn**
- real *8, dimension(:), allocatable **tgrazp**
- real *8, dimension(:), allocatable **latq**
- total lateral flow in soil profile for the day in HRU (mm H2O)*
- real *8, dimension(:), allocatable **latno3**
- real *8, dimension(:), allocatable **minpgw**
- real *8, dimension(:), allocatable **no3gw**
- real *8, dimension(:), allocatable **npInt**
- real *8, dimension(:), allocatable **tileq**
- real *8, dimension(:), allocatable **tileno3**
- real *8, dimension(:), allocatable **sedminpa**
- real *8, dimension(:), allocatable **sedminps**
- real *8, dimension(:), allocatable **sedorgn**
- real *8, dimension(:), allocatable **sedyld**
- soil loss caused by water erosion for day in HRU (metric tons)*
- real *8, dimension(:), allocatable **sepbtm**
- percolation from bottom of soil profile for the day in HRU (mm H2O)*
- real *8, dimension(:), allocatable **sedorgp**
- real *8, dimension(:), allocatable **strsn**
- real *8, dimension(:), allocatable **surfq**
- surface runoff generated in HRU on the current day (mm H2O)*
- real *8, dimension(:), allocatable **strsp**
- real *8, dimension(:), allocatable **strstmp**
- real *8, dimension(:), allocatable **surqno3**
- real *8, dimension(:), allocatable **hru_ha**
- area of HRU in hectares (ha)*
- real *8, dimension(:), allocatable **hru_dafr**
- fraction of total watershed area contained in HRU (km2/km2)*
- real *8, dimension(:), allocatable **tcfrtn**
- real *8, dimension(:), allocatable **tcfrtp**
- real *8, dimension(:), allocatable **drydep_no3**
- atmospheric dry deposition of nitrates (kg/ha/yr)*
- real *8, dimension(:), allocatable **drydep_nh4**
- atmospheric dry deposition of ammonia (kg/ha/yr)*
- real *8, dimension(:), allocatable **bio_yrms**
- annual biomass (dry weight) in the HRU (metric tons/ha)*
- real *8, dimension(:), allocatable **phubase**
- base zero total heat units (used when no land cover is growing) (heat units)*
- real *8, dimension(:), allocatable **hvstiadj**
- optimal harvest index for current time during growing season ((kg/ha)/(kg/ha))*
- real *8, dimension(:), allocatable **laiday**
- leaf area index for HRU (m^2/m^2)*
- real *8, dimension(:), allocatable **chlap**
- chlorophyll-a production coefficient for pond (none)*
- real *8, dimension(:), allocatable **pnd_psed**
- amount of mineral P attached to sediment originating from surface runoff in pond at end of day (kg P)*
- real *8, dimension(:), allocatable **laimxfr**
- real *8, dimension(:), allocatable **seccip**

- water clarity coefficient for pond (none)*
- real *8, dimension(:), allocatable [plantn](#)
- amount of nitrogen in plant biomass (kg N/ha)*
- real *8, dimension(:), allocatable [plt_et](#)
- actual ET simulated during life of plant (mm H2O)*
- real *8, dimension(:), allocatable [wet_psed](#)
- amount of mineral P attached to sediment originating from surface runoff in wetland at end of day (kg P)*
- real *8, dimension(:), allocatable [bio_aams](#)
- average annual biomass (dry weight) in the HRU (metric tons)*
- real *8, dimension(:), allocatable [plantp](#)
- amount of phosphorus in plant biomass (kg P/ha)*
- real *8, dimension(:), allocatable [plt_pet](#)
- potential ET simulated during life of plant (mm H2O)*
- real *8, dimension(:), allocatable [dormhr](#)
- time threshold used to define dormant period for plant (when daylength is within the time specified by dl from the minimum daylength for the area, the plant will go dormant) (hour)*
- real *8, dimension(:), allocatable [lai_yrmx](#)
- maximum leaf area index for the year in the HRU (none)*
- real *8, dimension(:), allocatable [bio_aamx](#)
- real *8, dimension(:), allocatable [lat_pst](#)
- amount of pesticide in lateral flow in HRU for the day (kg pst/ha)*
- real *8, dimension(:), allocatable [fld_fr](#)
- fraction of HRU area that drains into floodplain (km^2/km^2)*
- real *8, dimension(:), allocatable [orig_snohru](#)
- real *8, dimension(:), allocatable [orig_potvol](#)
- real *8, dimension(:), allocatable [pltfr_n](#)
- fraction of plant biomass that is nitrogen (none)*
- real *8, dimension(:), allocatable [orig_alai](#)
- real *8, dimension(:), allocatable [orig_bioms](#)
- real *8, dimension(:), allocatable [pltfr_p](#)
- fraction of plant biomass that is phosphorus (none)*
- real *8, dimension(:), allocatable [orig_phuacc](#)
- real *8, dimension(:), allocatable [orig_sumix](#)
- real *8, dimension(:), allocatable [phu_plt](#)
- total number of heat units to bring plant to maturity (heat units)*
- real *8, dimension(:), allocatable [orig_phu](#)
- real *8, dimension(:), allocatable [orig_shallst](#)
- real *8, dimension(:), allocatable [orig_deepst](#)
- real *8, dimension(:), allocatable [rip_fr](#)
- fraction of HRU area that drains into riparian zone (km^2/km^2)*
- real *8, dimension(:), allocatable [orig_pndvol](#)
- real *8, dimension(:), allocatable [orig_pndsds](#)
- real *8, dimension(:), allocatable [orig_pndno3](#)
- real *8, dimension(:), allocatable [orig_pndsols](#)
- real *8, dimension(:), allocatable [orig_pndorgn](#)
- real *8, dimension(:), allocatable [orig_pndorgp](#)
- real *8, dimension(:), allocatable [orig_wetvol](#)
- real *8, dimension(:), allocatable [orig_wetsds](#)
- real *8, dimension(:), allocatable [orig_wetno3](#)
- real *8, dimension(:), allocatable [orig_wetsols](#)
- real *8, dimension(:), allocatable [orig_wetorgn](#)
- real *8, dimension(:), allocatable [orig_wetorgp](#)

- real *8, dimension(:), allocatable **orig_solcov**
- real *8, dimension(:), allocatable **orig_solsw**
- real *8, dimension(:), allocatable **orig_potno3**
- real *8, dimension(:), allocatable **orig_potsed**
- real *8, dimension(:), allocatable **wtab**
water table based on 30 day antecedent climate (precip,et) (mm)
- real *8, dimension(:), allocatable **wtab_mn**
- real *8, dimension(:), allocatable **wtab_mx**
- real *8, dimension(:), allocatable **shallst_n**
nitrate concentration in shallow aquifer converted to kg/ha (ppm NO3-N)
- real *8, dimension(:), allocatable **gw_nloss**
- real *8, dimension(:), allocatable **rchrg_n**
- real *8, dimension(:), allocatable **det_san**
- real *8, dimension(:), allocatable **det_sil**
- real *8, dimension(:), allocatable **det_cla**
- real *8, dimension(:), allocatable **det_sag**
- real *8, dimension(:), allocatable **det_lag**
- real *8, dimension(:), allocatable **afrt_surface**
fraction of fertilizer which is applied to top 10 mm of soil (the remaining fraction is applied to first soil layer) (none)
- real *8, dimension(:), allocatable **tnylda**
- real *8 **frt_surface**
fraction of fertilizer which is applied to the top 10 mm of soil (the remaining fraction is applied to the first soil layer) (none)
- real *8, dimension(:), allocatable **auto_nyr**
maximum NO3-N content allowed to be applied in one year (kg NO3-N/ha)
- real *8, dimension(:), allocatable **auto_napp**
maximum NO3-N content allowed in one fertilizer application (kg NO3-N/ha)
- real *8, dimension(:), allocatable **auto_nstrs**
nitrogen stress factor which triggers auto fertilization (none)
- real *8, dimension(:), allocatable **manure_kg**
dry weight of manure deposited on HRU daily ((kg/ha)/day)
- real *8, dimension(:,:), allocatable **rcn_mo**
- real *8, dimension(:,:), allocatable **rammo_mo**
- real *8, dimension(:,:), allocatable **drydep_no3_mo**
- real *8, dimension(:,:), allocatable **drydep_nh4_mo**
- real *8, dimension(:), allocatable **rcn_d**
- real *8, dimension(:), allocatable **rammo_d**
- real *8, dimension(:), allocatable **drydep_no3_d**
- real *8, dimension(:), allocatable **drydep_nh4_d**
- real *8, dimension(:,:), allocatable **yldn**
- integer, dimension(:,:), allocatable **gwati**
- real *8, dimension(:,:), allocatable **gwatn**
- real *8, dimension(:,:), allocatable **gwatl**
- real *8, dimension(:,:), allocatable **gwatw**
- real *8, dimension(:,:), allocatable **gwatd**
- real *8, dimension(:,:), allocatable **gwatveg**
- real *8, dimension(:,:), allocatable **gwata**
- real *8, dimension(:,:), allocatable **gwats**
- real *8, dimension(:,:), allocatable **gwatspcon**
- real *8, dimension(:,:), allocatable **rfqeo_30d**
- real *8, dimension(:,:), allocatable **eo_30d**
- real *8, dimension(:), allocatable **psetlp1**
phosphorus settling rate for 1st season (m/day)

- real *8, dimension(:), allocatable [psetlp2](#)
phosphorus settling rate for 2nd season (m/day)
- real *8, dimension(:,:), allocatable [wgnold](#)
previous value of wgncur(:,:) (none)
- real *8, dimension(:,:), allocatable [wgncur](#)
parameter to predict the impact of precip on other weather attributes (none)
wgncur(1,:) parameter which predicts impact of precip on daily maximum air temperature
wgncur(2,:) parameter which predicts impact of precip on daily minimum air temperature
wgncur(3,:) parameter which predicts impact of precip on daily solar radiation
- real *8, dimension(:), allocatable [wrt1](#)
1st shape parameter for calculation of water retention (none)
- real *8, dimension(:), allocatable [wrt2](#)
2nd shape parameter for calculation of water retention (none)
- real *8, dimension(:,:), allocatable [pst_enr](#)
pesticide enrichment ratio (none)
- real *8, dimension(:,:), allocatable [zdb](#)
- real *8, dimension(:,:), allocatable [pst_surq](#)
- real *8, dimension(:,:), allocatable [plt_pst](#)
pesticide on plant foliage (kg/ha)
- real *8, dimension(:), allocatable [psetlw1](#)
phosphorus settling rate for 1st season (m/day)
- real *8, dimension(:), allocatable [psetlw2](#)
phosphorus settling rate for 2nd season (m/day)
- real *8, dimension(:,:), allocatable [pst_sed](#)
- real *8, dimension(:,:), allocatable [wupnd](#)
average daily water removal from the pond for the month ($10^4 \text{ m}^3/\text{day}$)
- real *8, dimension(:,:), allocatable [phi](#)
phi(1,:) cross-sectional area of flow at bankfull depth (m^2) phi(2,:) (none) phi(3,:) (none) phi(4,:) (none) phi(5,:) (none) phi(6,:) bottom width of main channel (m) phi(7,:) depth of water when reach is at bankfull depth (m) phi(8,:) average velocity when reach is at bankfull depth (m/s) phi(9,:) wave celerity when reach is at bankfull depth (m/s) phi(10,:) storage time constant for reach at bankfull depth (ratio of storage to discharge) (hour) phi(11,:) average velocity when reach is at 0.1 bankfull depth (low flow) (m/s) phi(12,:) wave celerity when reach is at 0.1 bankfull depth (low flow) (m/s) phi(13,:) storage time constant for reach at 0.1 bankfull depth (low flow) (ratio of storage to discharge) (hour)
- real *8, dimension(:,:), allocatable [pcpband](#)
precipitation for the day in band in HRU (mm H₂O)
- real *8, dimension(:,:), allocatable [tavband](#)
average temperature for the day in band in HRU (deg C)
- real *8, dimension(:), allocatable [wat_phi1](#)
cross-sectional area of flow at bankfull depth (m^2)
- real *8, dimension(:), allocatable [wat_phi5](#)
flow rate when reach is at bankfull depth (m^3/s)
- real *8, dimension(:), allocatable [wat_phi6](#)
bottom width of main channel (m)
- real *8, dimension(:), allocatable [wat_phi7](#)
depth of water when reach is at bankfull (m)
- real *8, dimension(:), allocatable [wat_phi8](#)
average velocity when reach is at bankfull depth (m/s)
- real *8, dimension(:), allocatable [wat_phi9](#)
wave celerity when reach is at bankfull depth (m/s)
- real *8, dimension(:), allocatable [wat_phi10](#)
storage time constant for reach at bankfull depth (ratio of storage to discharge) (hour)
- real *8, dimension(:), allocatable [wat_phi11](#)

- average velocity when reach is at 0.1 bankfull depth (low flow) (m/s)*
- real *8, dimension(:), allocatable [wat_phi12](#)
- wave celerity when reach is at 0.1 bankfull depth (low flow) (m/s)*
- real *8, dimension(:), allocatable [wat_phi13](#)
- storage time constant for reach at 0.1 bankfull depth (low flow) (ratio of storage to discharge) (hour)*
- real *8, dimension(:,:), allocatable [snoeb](#)
- snow water content in elevation band on current day (mm H2O)*
- real *8, dimension(:,:), allocatable [wudeep](#)
- average daily water removal from the deep aquifer for the month ($10^4 \text{ m}^3/\text{day}$)*
- real *8, dimension(:,:), allocatable [wushal](#)
- average daily water removal from the shallow aquifer for the month ($10^4 \text{ m}^3/\text{day}$)*
- real *8, dimension(:,:), allocatable [tmnband](#)
- minimum temperature for the day in band in HRU (deg C)*
- real *8, dimension(:), allocatable **bs1**
- real *8, dimension(:), allocatable **bs2**
- real *8, dimension(:), allocatable **bs3**
- real *8, dimension(:), allocatable **bs4**
- real *8, dimension(:), allocatable [nsetlw1](#)
- nitrogen settling rate for 1st season (m/day)*
- real *8, dimension(:), allocatable [nsetlw2](#)
- nitrogen settling rate for 2nd season (m/day)*
- real *8, dimension(:,:), allocatable [snotmpeb](#)
- temperature of snow pack in elevation band (deg C)*
- real *8, dimension(:), allocatable [surf_bs1](#)
- amount of surface runoff lagged over one day (mm H2O)*
- real *8, dimension(:), allocatable **surf_bs2**
- real *8, dimension(:), allocatable **surf_bs3**
- real *8, dimension(:), allocatable **surf_bs4**
- real *8, dimension(:), allocatable **surf_bs5**
- real *8, dimension(:), allocatable **surf_bs6**
- real *8, dimension(:), allocatable **surf_bs7**
- real *8, dimension(:), allocatable **surf_bs8**
- real *8, dimension(:), allocatable **surf_bs9**
- real *8, dimension(:), allocatable **surf_bs10**
- real *8, dimension(:), allocatable **surf_bs11**
- real *8, dimension(:), allocatable **surf_bs12**
- real *8, dimension(:), allocatable **surf_bs13**
- real *8, dimension(:), allocatable **surf_bs14**
- real *8, dimension(:), allocatable **surf_bs15**
- real *8, dimension(:), allocatable **surf_bs16**
- real *8, dimension(:), allocatable **surf_bs17**
- real *8, dimension(:), allocatable [nsetlp1](#)
- nitrogen settling rate for 1st season (m/day)*
- real *8, dimension(:), allocatable [nsetlp2](#)
- nitrogen settling rate for 2nd season (m/day)*
- real *8, dimension(:,:), allocatable [tmxband](#)
- maximum temperature for the day in band in HRU (deg C)*
- real *8, dimension(:,:), allocatable [frad](#)
- fraction of solar radiation occuring during hour in day in HRU (none)*
- real *8, dimension(:,:), allocatable [rainsub](#)
- precipitation for the time step during the day in HRU (mm H2O)*
- real *8, dimension(:), allocatable **rstpbsb**

- real *8, dimension(:,:), allocatable **orig_snoeb**
- real *8, dimension(:,:), allocatable **orig_pltpst**
- real *8, dimension(:,:), allocatable **terr_p**
- real *8, dimension(:,:), allocatable **terr_cn**
- real *8, dimension(:,:), allocatable **terr_sl**
- real *8, dimension(:,:), allocatable **drain_d**
- real *8, dimension(:,:), allocatable **drain_t**
- real *8, dimension(:,:), allocatable **drain_g**
- real *8, dimension(:,:), allocatable **drain_idep**
- real *8, dimension(:,:), allocatable **cont_cn**
- real *8, dimension(:,:), allocatable **cont_p**
- real *8, dimension(:,:), allocatable **filt_w**
- real *8, dimension(:,:), allocatable **strip_n**
- real *8, dimension(:,:), allocatable **strip_cn**
- real *8, dimension(:,:), allocatable **strip_c**
- real *8, dimension(:,:), allocatable **strip_p**
- real *8, dimension(:,:), allocatable **fire_cn**
- real *8, dimension(:,:), allocatable **cropno_upd**
- real *8, dimension(:,:), allocatable **hi_upd**
- real *8, dimension(:,:), allocatable **laimx_upd**
- real *8, dimension(:,:), allocatable **phug**
fraction of plant heat units at which grazing begins (none)
- real *8, dimension(:,:), allocatable **pst_lag**
- integer, dimension(:), allocatable **hrupest**
pesticide use flag (none)
0: no pesticides used in HRU
1: pesticides used in HRU
- integer, dimension(:), allocatable **nrelease**
sequence number of impound/release operation within the year (none)
- integer, dimension(:), allocatable **swtrg**
rainfall event flag (none):
0: no rainfall event over midnight
1: rainfall event over midnight
- integer, dimension(:), allocatable **nrot**
number of years of rotation (none)
- integer, dimension(:), allocatable **nfert**
sequence number of fertilizer application within the year (none)
- integer, dimension(:), allocatable **nro**
sequence number of year in rotation (none)
- integer, dimension(:), allocatable **igro**
land cover status code (none). This code informs the model whether or not a land cover is growing at the beginning of the simulation
0 no land cover growing
1 land cover growing
- integer, dimension(:), allocatable **ipnd1**
beginning month of nutrient settling season (none)
- integer, dimension(:), allocatable **ipnd2**
ending month of nutrient settling season (none)
- integer, dimension(:), allocatable **nair**
sequence number of auto-irrigation application within the year (none)
- integer, dimension(:), allocatable **iflod1**
beginning month of non-flood season (none)
- integer, dimension(:), allocatable **iflod2**

- ending month of non-flood season (none)*
- integer, dimension(:), allocatable **ndtarg**
 - number of days required to reach target storage from current pond storage (none)*
- integer, dimension(:), allocatable **nirr**
 - sequence number of irrigation application within the year (none)*
- integer, dimension(:), allocatable **iafrttyp**
- integer, dimension(:), allocatable **nstress**
- integer, dimension(:), allocatable **igrotree**
- integer, dimension(:), allocatable **grz_days**
 - number of days grazing will be simulated (none)*
- integer, dimension(:), allocatable **nmgt**
 - management code (for GIS output only) (none)*
- integer, dimension(:), allocatable **nafert**
 - sequence number of auto-fert application within the year (none)*
- integer, dimension(:), allocatable **nsweep**
 - sequence number of street sweeping operation within the year (none)*
- integer, dimension(:), allocatable **icr**
 - sequence number of crop grown within the current year (none)*
- integer, dimension(:), allocatable **ncut**
 - sequence number of harvest operation within a year (none)*
- integer, dimension(:), allocatable **irrno**
 - irrigation source location (none)*
 - if IRRSC=1, IRRNO is the number of the reach*
 - if IRRSC=2, IRRNO is the number of the reservoir*
 - if IRRSC=3, IRRNO is the number of the subbasin*
 - if IRRSC=4, IRRNO is the number of the subbasin*
 - if IRRSC=5, not used*
- integer, dimension(:), allocatable **sol_nly**
 - number of soil layers in HRU (none)*
- integer, dimension(:), allocatable **npcp**
 - prior day category (none)*
 - 1 dry day*
 - 2 wet day*
- integer, dimension(:), allocatable **irn**
 - average annual number of irrigation applications in HRU (none)*
- integer, dimension(:), allocatable **ncf**
 - sequence number of continuous fertilization operation within the year (none)*
- integer, dimension(:), allocatable **ngr**
 - sequence number of grazing operation within the year (none)*
- integer, dimension(:), allocatable **igrz**
 - grazing flag for HRU (none):*
 - 0 HRU currently not grazed*
 - 1 HRU currently grazed*
- integer, dimension(:), allocatable **ndeat**
 - number of days HRU has been grazed (days)*
- integer, dimension(:), allocatable **hru_sub**
 - subbasin in which HRU is located (none)*
- integer, dimension(:), allocatable **urbldu**
 - urban land type identification number from urban.dat (none)*
- integer, dimension(:), allocatable **ldrain**
 - soil layer where drainage tile is located (none)*
- integer, dimension(:), allocatable **idorm**

- dormancy status code (none):*
 - 0 land cover growing (not dormant)
 - 1 land cover dormant
- integer, dimension(:), allocatable **hru_seq**
- integer, dimension(:), allocatable **iurban**
- urban simulation code (none):*
 - 0 no urban sections in HRU
 - 1 urban sections in HRU, simulate using USGS regression equations
 - 2 urban sections in HRU, simulate using build up/wash off algorithm
- integer, dimension(:), allocatable **iday_fert**
- integer, dimension(:), allocatable **icftrt**
- integer, dimension(:), allocatable **ifld**
- number of HRU (in subbasin) that is a floodplain (none)*
- integer, dimension(:), allocatable **irip**
- number of HRU (in subbasin) that is a riparian zone (none)*
- integer, dimension(:), allocatable **hrugis**
- GIS code printed to output files (output.hru, .rch) (none)*
- integer, dimension(:), allocatable **ndcftrt**
- integer, dimension(:), allocatable **irrsc**
- irrigation source code (none):*
 - 1 divert water from reach
 - 2 divert water from reservoir
 - 3 divert water from shallow aquifer
 - 4 divert water from deep aquifer
 - 5 divert water from source outside watershed
- integer, dimension(:), allocatable **ntil**
- sequence number of tillage operation within current year (none)*
- integer, dimension(:), allocatable **orig_igro**
- integer, dimension(:), allocatable **iwatable**
- integer, dimension(:), allocatable **curyr_mat**
- integer, dimension(:), allocatable **icpst**
- icpst = 0 do not apply*
- icpst = 1 application period*
- integer, dimension(:), allocatable **ndcpst**
- current day within the application period (day)*
- integer, dimension(:), allocatable **ncpest**
- integer, dimension(:), allocatable **iday_pest**
- current day between applications (day)*
- integer, dimension(:), allocatable **irr_flag**
- integer, dimension(:), allocatable **irra_flag**
- integer, dimension(:,:), allocatable **rndseed**
- random number generator seeds array. The seeds in the array are used to generate random numbers for the following purposes (none):*
 - (1) wet/dry day probability
 - (2) solar radiation
 - (3) precipitation
 - (4) USLE rainfall erosion index
 - (5) wind speed
 - (6) 0.5 hr rainfall fraction
 - (7) relative humidity
 - (8) maximum temperature
 - (9) minimum temperature
 - (10) generate new random numbers
- integer, dimension(:,:), allocatable **iterr**
- integer, dimension(:,:), allocatable **iyterr**
- integer, dimension(:,:), allocatable **itdrain**

- integer, dimension(:,:), allocatable **iydrain**
- integer, dimension(:,:), allocatable **ncrops**
- integer, dimension(:), allocatable [manure_id](#)
manure (fertilizer) identification number from fert.dat (none)
- integer, dimension(:,:), allocatable **mgt_sdr**
- integer, dimension(:,:), allocatable **idplot**
- integer, dimension(:,:), allocatable **icont**
- integer, dimension(:,:), allocatable **iycont**
- integer, dimension(:,:), allocatable **ifilt**
- integer, dimension(:,:), allocatable **iyfilt**
- integer, dimension(:,:), allocatable **istrip**
- integer, dimension(:,:), allocatable **ystrip**
- integer, dimension(:,:), allocatable **iopday**
- integer, dimension(:,:), allocatable **iopyr**
- integer, dimension(:,:), allocatable **mgt_ops**
- real *8, dimension(:), allocatable [wshd_pstap](#)
total amount of pesticide type applied in watershed during simulation (kg/ha)
- real *8, dimension(:), allocatable **wshd_pstdg**
- integer, dimension(12) **ndmo**
- integer, dimension(:), allocatable [npno](#)
array of unique pesticides used in watershed (none)
- integer, dimension(:), allocatable **mcrhru**
- character(len=13), dimension(18) [rfile](#)
rainfall file names (.pcp)
- character(len=13), dimension(18) [tfile](#)
temperature file names (.tmp)
- character(len=4), dimension(1000) [urbyname](#)
name of urban land use
- character(len=1), dimension(:), allocatable **kirr**
irrigation in HRU
- character(len=1), dimension(:), allocatable **hydgrp**
- character(len=16), dimension(:), allocatable **snam**
soil series name
- character(len=17), dimension(300) [pname](#)
name of pesticide/toxin
- character(len=4), dimension(60) [title](#)
description lines in file.cio (1st 3 lines)
- character(len=4), dimension(5000) [cpnm](#)
four character code to represent crop name
- character(len=17), dimension(50) **fname**
- real *8, dimension(:,:,:), allocatable [flomon](#)
average daily water loading for month (m^3/day)
- real *8, dimension(:,:,:), allocatable [solpstmon](#)
average daily soluble pesticide loading for month (mg pst/day)
- real *8, dimension(:,:,:), allocatable [srbpstmon](#)
average daily sorbed pesticide loading for month (mg pst/day)
- real *8, dimension(:,:,:), allocatable [orgnmon](#)
average daily organic N loading for month (kg N/day)
- real *8, dimension(:,:,:), allocatable [orgpmon](#)
average daily organic P loading for month (kg P/day)
- real *8, dimension(:,:,:), allocatable [sedmon](#)
average daily sediment loading for month (metric tons/day)

- real *8, dimension(:,:,:), allocatable [minpmon](#)
average daily mineral P loading for month (kg P/day)
- real *8, dimension(:,:,:), allocatable [nh3mon](#)
average amount of NH3-N loaded to stream on a given day in the month (kg N/day)
- real *8, dimension(:,:,:), allocatable [no3mon](#)
average daily NO3-N loading for month (kg N/day)
- real *8, dimension(:,:,:), allocatable [bactlpmon](#)
average amount of less persistent bacteria loaded to stream on a given day in the month (# bact/day)
- real *8, dimension(:,:,:), allocatable [bactpmon](#)
average amount of persistent bacteria loaded to stream on a given day in the month (# bact/day)
- real *8, dimension(:,:,:), allocatable [no2mon](#)
average amount of NO2-N loaded to stream on a given day in the month (kg N/day)
- real *8, dimension(:,:,:), allocatable [cmtl1mon](#)
average amount of conservative metal #1 loaded to stream on a given day in the month (# bact/day)
- real *8, dimension(:,:,:), allocatable [cmtl2mon](#)
average amount of conservative metal #2 loaded to stream on a given day in the month (# bact/day)
- real *8, dimension(:,:,:), allocatable [cmtl3mon](#)
average amount of conservative metal #3 loaded to stream on a given day in the month (# bact/day)
- real *8, dimension(:,:,:), allocatable [cbodmon](#)
average daily loading of CBOD in month (kg/day)
- real *8, dimension(:,:,:), allocatable [chlamon](#)
average daily loading of chlorophyll-a in month (kg/day)
- real *8, dimension(:,:,:), allocatable [disoxmon](#)
average daily loading of dissolved O2 in month (kg/day)
- real *8, dimension(:,:), allocatable [floyr](#)
average daily water loading for year (m^3/day)
- real *8, dimension(:,:), allocatable [orgnyr](#)
average daily organic N loading for year (kg N/day)
- real *8, dimension(:,:), allocatable [orgpyr](#)
average daily organic P loading for year (kg P/day)
- real *8, dimension(:,:), allocatable [sedyr](#)
average daily sediment loading for year (metric tons/day)
- real *8, dimension(:,:), allocatable [minpyr](#)
average daily mineral P loading for year (kg P/day)
- real *8, dimension(:,:), allocatable [nh3yr](#)
average daily NH3-N loading for year (kg N/day)
- real *8, dimension(:,:), allocatable [no2yr](#)
average daily NO2-N loading for year (kg N/day)
- real *8, dimension(:,:), allocatable [no3yr](#)
average daily NO3-N loading for year (kg N/day)
- real *8, dimension(:,:), allocatable [bactlpyr](#)
average daily loading of less persistent bacteria for year (# bact/day)
- real *8, dimension(:,:), allocatable [bactpyr](#)
average daily loading of persistent bacteria for year (# bact/day)
- real *8, dimension(:,:), allocatable [cmtl1yr](#)
average daily loading of conservative metal #1 for year (kg/day)
- real *8, dimension(:,:), allocatable [chlayr](#)
average daily loading of chlorophyll-a in year (kg/day)
- real *8, dimension(:,:), allocatable [cmtl2yr](#)
average daily loading of conservative metal #2 for year (kg/day)
- real *8, dimension(:,:), allocatable [cmtl3yr](#)

- average daily loading of conservative metal #3 for year (kg/day)*
- real *8, dimension(:,:), allocatable [cbodyr](#)
 - average daily loading of CBOD in year (kg/day)*
- real *8, dimension(:,:), allocatable [disoxyr](#)
 - average daily loading of dissolved O2 in year (kg/day)*
- real *8, dimension(:,:), allocatable [solpstyr](#)
 - average daily soluble pesticide loading for year (mg pst/day)*
- real *8, dimension(:,:), allocatable [srbpstyr](#)
 - average daily sorbed pesticide loading for year (mg pst/day)*
- real *8, dimension(:,:), allocatable [sol_mc](#)
- real *8, dimension(:,:), allocatable [sol_mn](#)
- real *8, dimension(:,:), allocatable [sol_mp](#)
- real *8, dimension(:), allocatable [flocnst](#)
- real *8, dimension(:), allocatable [orgncnst](#)
 - average daily organic N loading to reach (kg N/day)*
- real *8, dimension(:), allocatable [sedcnst](#)
 - average daily sediment loading for reach (metric tons/day)*
- real *8, dimension(:), allocatable [minpcnst](#)
 - average daily soluble P loading to reach (kg P/day)*
- real *8, dimension(:), allocatable [no3cnst](#)
 - average daily nitrate loading to reach (kg N/day)*
- real *8, dimension(:), allocatable [orgpcnst](#)
 - average daily organic P loading to reach (kg P/day)*
- real *8, dimension(:), allocatable [bactpcnst](#)
 - average daily persistent bacteria loading to reach (# bact/day)*
- real *8, dimension(:), allocatable [nh3cnst](#)
 - average daily ammonia loading to reach (kg N/day)*
- real *8, dimension(:), allocatable [no2cnst](#)
 - average daily nitrite loading to reach (kg N/day)*
- real *8, dimension(:), allocatable [bactlpcnst](#)
 - average daily less persistent bacteria loading to reach (# bact/day)*
- real *8, dimension(:), allocatable [cmtl1cnst](#)
 - average daily conservative metal #1 loading (kg/day)*
- real *8, dimension(:), allocatable [cmtl2cnst](#)
 - average daily conservative metal #2 loading (kg/day)*
- real *8, dimension(:), allocatable [chlacnst](#)
 - average daily loading of chlorophyll-a (kg/day)*
- real *8, dimension(:), allocatable [cmtl3cnst](#)
 - average daily conservative metal #3 loading (kg/day)*
- real *8, dimension(:), allocatable [disoxcnst](#)
 - average daily loading of dissolved O2 (kg/day)*
- real *8, dimension(:), allocatable [cbodcnst](#)
 - average daily loading of CBOD to reach (kg/day)*
- real *8, dimension(:), allocatable [solpstcnst](#)
 - average daily soluble pesticide loading (mg/day)*
- real *8, dimension(:), allocatable [srbpstcnst](#)
 - average daily sorbed pesticide loading (mg/day)*
- integer [nstep](#)
 - max number of time steps per day or number of lines of rainfall data for each day (none)*
- integer [idt](#)
 - length of time step used to report precipitation data for sub-daily modeling (minutes)*

- real *8, dimension(:), allocatable **hrtwtr**
- real *8, dimension(:), allocatable **hhstor**
- real *8, dimension(:), allocatable **hdepth**
- real *8, dimension(:), allocatable **hsdti**
- real *8, dimension(:), allocatable **hrchwtr**
- real *8, dimension(:), allocatable **halgae**
- real *8, dimension(:), allocatable **horgn**
- real *8, dimension(:), allocatable **hnh4**
- real *8, dimension(:), allocatable **hno2**
- real *8, dimension(:), allocatable **hno3**
- real *8, dimension(:), allocatable **horgp**
- real *8, dimension(:), allocatable **hsolp**
- real *8, dimension(:), allocatable **hbod**
- real *8, dimension(:), allocatable **hdisox**
- real *8, dimension(:), allocatable **hchla**
- real *8, dimension(:), allocatable **hsedyld**
- real *8, dimension(:), allocatable **hsedst**
- real *8, dimension(:), allocatable **hharea**
- real *8, dimension(:), allocatable **hsolpst**
- real *8, dimension(:), allocatable **hsorpst**
- real *8, dimension(:), allocatable **hhqday**
surface runoff generated each timestep of day in HRU (mm H2O)
- real *8, dimension(:), allocatable **precipdt**
precipitation, or effective precipitation reaching soil surface, in time step for HRU (mm H2O)
- real *8, dimension(:), allocatable **hhtime**
- real *8, dimension(:), allocatable **hbactp**
- real *8, dimension(:), allocatable **hbactlp**
- integer, dimension(10) **ivar_orig**
- real *8, dimension(10) **rvar_orig**
- integer **nsave**
number of save commands in .fig file
- integer **nauto**
- integer **iatmodep**
- real *8, dimension(:), allocatable **wattemp**
- real *8, dimension(:), allocatable **lkpst_mass**
- real *8, dimension(:), allocatable **lkspst_mass**
- real *8, dimension(:), allocatable **vel_chan**
- real *8, dimension(:), allocatable **vfscon**
fraction of the total runoff from the entire field entering the most concentrated 10% of the VFS (none)
- real *8, dimension(:), allocatable **vfsratio**
field area/VFS area ratio (none)
- real *8, dimension(:), allocatable **vfsch**
fraction of flow entering the most concentrated 10% of the VFS which is fully channelized (none)
- real *8, dimension(:), allocatable **vfsi**
- real *8, dimension(:, :), allocatable **filter_i**
- real *8, dimension(:, :), allocatable **filter_ratio**
- real *8, dimension(:, :), allocatable **filter_con**
- real *8, dimension(:, :), allocatable **filter_ch**
- real *8, dimension(:, :), allocatable **sol_n**
- integer **cswat**
= 0 Static soil carbon (old mineralization routines)
= 1 C-FARM one carbon pool model
= 2 Century model
- real *8, dimension(:, :), allocatable **sol_bdp**

- real *8, dimension(:,:), allocatable **tillagef**
- real *8, dimension(:), allocatable **rtfr**
- real *8, dimension(:), allocatable **stsol_rd**
- integer **urban_flag**
- integer **dorm_flag**
- real *8 **bf_flg**
- real *8 **iabstr**
- real *8, dimension(:), allocatable **ubnrunoff**
- real *8, dimension(:), allocatable **ubntss**
- real *8, dimension(:,:), allocatable **sub_ubnrunoff**
- real *8, dimension(:,:), allocatable **sub_ubntss**
- real *8, dimension(:,:), allocatable **ovrlnd_dt**
- real *8, dimension(:,:), allocatable **hhsurf_bs1**
- real *8, dimension(:,:), allocatable **hhsurf_bs2**
- integer **iuh**
unit hydrograph method: 1=triangular UH; 2=gamma funtion UH;
- integer **sed_ch**
channel routing for HOURLY; 0=Bagnold; 2=Brownlie; 3=Yang;
- real *8 **eros_expo**
an exponent in the overland flow erosion equation ranges 1.5-3.0
- real *8 **eros_spl**
coefficient of splash erosion varing 0.9-3.1
- real *8 **rill_mult**
Multiplier to USLE_K for soil susceptible to rill erosion, range 0.5-2.0.
- real *8 **sedprev**
- real *8 **c_factor**
- real *8 **ch_d50**
median particle diameter of channel bed (mm)
- real *8 **sig_g**
geometric standard deviation of particle sizes for the main channel. Mean air temperature at which precipitation is equally likely to be rain as snow/freezing rain.
- real *8 **uhalpha**
alpha coefficient for estimating unit hydrograph using a gamma function (.bsn)*
- real *8 **abstinit**
- real *8 **abstmax**
- real *8, dimension(:,:), allocatable **hhsedy**
sediment yield from HRU drung a time step applied to HRU (tons)
- real *8, dimension(:,:), allocatable **sub_subp_dt**
- real *8, dimension(:,:), allocatable **sub_hhsedy**
- real *8, dimension(:,:), allocatable **sub_atmp**
- real *8, dimension(:), allocatable **rhy**
- real *8, dimension(:), allocatable **init_abstrc**
- real *8, dimension(:), allocatable **dratio**
- real *8, dimension(:), allocatable **hrtevp**
- real *8, dimension(:), allocatable **hrttic**
- real *8, dimension(:,:), allocatable **rchhr**
- real *8, dimension(:), allocatable **hhresflwi**
- real *8, dimension(:), allocatable **hhresflwo**
- real *8, dimension(:), allocatable **hhressedi**
- real *8, dimension(:), allocatable **hhressedo**
- character(len=4), dimension(:), allocatable **lu_nodrain**
- integer, dimension(:), allocatable **bmpdrain**
- real *8, dimension(:), allocatable **sub_cn2**

- real *8, dimension(:), allocatable **sub_ha_urb**
- real *8, dimension(:), allocatable **bmp_recharge**
- real *8, dimension(:), allocatable **sub_ha_imp**
- real *8, dimension(:), allocatable **subdr_km**
- real *8, dimension(:), allocatable **subdr_ickm**
- real *8, dimension(:, :), allocatable **sf_im**
- real *8, dimension(:, :), allocatable **sf_iy**
- real *8, dimension(:, :), allocatable **sp_sa**
- real *8, dimension(:, :), allocatable **sp_pvol**
- real *8, dimension(:, :), allocatable **sp_pd**
- real *8, dimension(:, :), allocatable **sp_sedi**
- real *8, dimension(:, :), allocatable **sp_sede**
- real *8, dimension(:, :), allocatable **ft_sa**
- real *8, dimension(:, :), allocatable **ft_fsa**
- real *8, dimension(:, :), allocatable **ft_dep**
- real *8, dimension(:, :), allocatable **ft_h**
- real *8, dimension(:, :), allocatable **ft_pd**
- real *8, dimension(:, :), allocatable **ft_k**
- real *8, dimension(:, :), allocatable **ft_dp**
- real *8, dimension(:, :), allocatable **ft_dc**
- real *8, dimension(:, :), allocatable **ft_por**
- real *8, dimension(:, :), allocatable **tss_den**
- real *8, dimension(:, :), allocatable **ft_alp**
- real *8, dimension(:, :), allocatable **sf_fr**
- real *8, dimension(:, :), allocatable **sp_qi**
- real *8, dimension(:, :), allocatable **sp_k**
- real *8, dimension(:, :), allocatable **ft_qpnd**
- real *8, dimension(:, :), allocatable **sp_dp**
- real *8, dimension(:, :), allocatable **ft_qsw**
- real *8, dimension(:, :), allocatable **ft_qin**
- real *8, dimension(:, :), allocatable **ft_qout**
- real *8, dimension(:, :), allocatable **ft_sedpnd**
- real *8, dimension(:, :), allocatable **sp_bpw**
- real *8, dimension(:, :), allocatable **ft_bpw**
- real *8, dimension(:, :), allocatable **ft_sed_cumul**
- real *8, dimension(:, :), allocatable **sp_sed_cumul**
- integer, dimension(:), allocatable **num_sf**
- integer, dimension(:, :), allocatable **sf_typ**
- integer, dimension(:, :), allocatable **sf_dim**
- integer, dimension(:, :), allocatable **ft_qfg**
- integer, dimension(:, :), allocatable **sp_qfg**
- integer, dimension(:, :), allocatable **sf_ptp**
- integer, dimension(:, :), allocatable **ft_fc**
- real *8 **sfsedmean**
- real *8 **sfsedstdev**
- integer, dimension(:), allocatable **dtp_imo**
month the reservoir becomes operational (none)
- integer, dimension(:), allocatable **dtp_iyr**
year of the simulation that the reservoir becomes operational (none)
- integer, dimension(:), allocatable **dtp_numstage**
total number of stages in the weir (none)
- integer, dimension(:), allocatable **dtp_numweir**
total number of weirs in the BMP (none)
- integer, dimension(:), allocatable **dtp_onoff**

- sub-basin detention pond is associated with (none)*
- integer, dimension(:), allocatable [dtp_reltype](#)
 - equations for stage-discharge relationship (none):*
 - 1=exponential function,*
 - 2=linear,*
 - 3=logarithmic,*
 - 4=cubic,*
 - 5=power*
- integer, dimension(:), allocatable [dtp_stagdis](#)
 - (none):*
 - 0=use weir/orifice discharge equation to calculate outflow,*
 - 1=use stage-dicharge relationship*
- integer, dimension(:), allocatable **dtp_subnum**
- real *8, dimension(:), allocatable [cf](#)
 - this parameter controls the response of decomposition to the combined effect of soil temperature and moisture.*
- real *8, dimension(:), allocatable [cfh](#)
 - maximum humification rate*
- real *8, dimension(:), allocatable [cfdec](#)
 - the undisturbed soil turnover rate under optimum soil water and temperature. Increasing it will increase carbon and organic N decomp.*
- real *8, dimension(:), allocatable **lat_orgn**
- real *8, dimension(:), allocatable **lat_orgp**
- integer, dimension(:, :), allocatable [dtp_weirdim](#)
 - weir dimensions (none),*
 - 1=read user input,*
 - 0=use model calculation*
- integer, dimension(:, :), allocatable [dtp_weirtype](#)
 - type of weir (none):*
 - 1=rectangular and*
 - 2=circular*
- real *8, dimension(:), allocatable [dtp_coef1](#)
 - coefficient of 3rd degree in the polynomial equation (none)*
- real *8, dimension(:), allocatable [dtp_coef2](#)
 - coefficient of 2nd degree in the polynomial equation (none)*
- real *8, dimension(:), allocatable [dtp_coef3](#)
 - coefficient of 1st degree in the polynomial equation (none)*
- real *8, dimension(:), allocatable [dtp_evrsv](#)
 - detention pond evaporation coefficient (none)*
- real *8, dimension(:), allocatable [dtp_expont](#)
 - exponent used in the exponential equation (none)*
- real *8, dimension(:), allocatable [dtp_intcept](#)
 - intercept used in regression equations (none)*
- real *8, dimension(:), allocatable [dtp_lwratio](#)
 - ratio of length to width of water back up (none)*
- real *8, dimension(:), allocatable [dtp_totwrwid](#)
 - total constructed width of the detention wall across the creek (m)*
- real *8, dimension(:), allocatable **dtp_inflvol**
- real *8, dimension(:), allocatable **dtp_wdep**
- real *8, dimension(:), allocatable **dtp_totdep**
- real *8, dimension(:), allocatable **dtp_watdepact**
- real *8, dimension(:), allocatable **dtp_outflow**
- real *8, dimension(:), allocatable **dtp_totrel**
- real *8, dimension(:), allocatable **dtp_backoff**
- real *8, dimension(:), allocatable **dtp_seep_sa**

- real *8, dimension(:), allocatable **dtb_evap_sa**
- real *8, dimension(:), allocatable **dtb_pet_day**
- real *8, dimension(:), allocatable **dtb_pcpvol**
- real *8, dimension(:), allocatable **dtb_seepvol**
- real *8, dimension(:), allocatable **dtb_evapvol**
- real *8, dimension(:), allocatable **dtb_flowin**
- real *8, dimension(:), allocatable **dtb_backup_length**
- real *8, dimension(:), allocatable **dtb_ivol**
- real *8, dimension(:), allocatable **dtb_ised**
- integer, dimension(:,:), allocatable **so_res_flag**
- integer, dimension(:,:), allocatable **ro_bmp_flag**
- real *8, dimension(:,:), allocatable **sol_watp**
- real *8, dimension(:,:), allocatable **sol_solp_pre**
- real *8, dimension(:,:), allocatable **psp_store**
- real *8, dimension(:,:), allocatable **ssp_store**
- real *8, dimension(:,:), allocatable **so_res**
- real *8, dimension(:,:), allocatable **sol_cal**
- real *8, dimension(:,:), allocatable **sol_ph**
- integer **sol_p_model**
- integer, dimension(:,:), allocatable **a_days**
- integer, dimension(:,:), allocatable **b_days**
- real *8, dimension(:), allocatable **min_res**

minimum residue allowed due to implementation of residue managment in the OPS file (kg/ha)

- real *8, dimension(:), allocatable **harv_min**
- real *8, dimension(:), allocatable **fstap**
- real *8, dimension(:,:), allocatable **ro_bmp_flo**
- real *8, dimension(:,:), allocatable **ro_bmp_sed**
- real *8, dimension(:,:), allocatable **ro_bmp_bac**
- real *8, dimension(:,:), allocatable **ro_bmp_pp**
- real *8, dimension(:,:), allocatable **ro_bmp_sp**
- real *8, dimension(:,:), allocatable **ro_bmp_pn**
- real *8, dimension(:,:), allocatable **ro_bmp_sn**
- real *8, dimension(:,:), allocatable **ro_bmp_flos**
- real *8, dimension(:,:), allocatable **ro_bmp_seds**
- real *8, dimension(:,:), allocatable **ro_bmp_bacs**
- real *8, dimension(:,:), allocatable **ro_bmp_pps**
- real *8, dimension(:,:), allocatable **ro_bmp_sps**
- real *8, dimension(:,:), allocatable **ro_bmp_pns**
- real *8, dimension(:,:), allocatable **ro_bmp_sns**
- real *8, dimension(:,:), allocatable **ro_bmp_flot**
- real *8, dimension(:,:), allocatable **ro_bmp_sedt**
- real *8, dimension(:,:), allocatable **ro_bmp_bact**
- real *8, dimension(:,:), allocatable **ro_bmp_ppt**
- real *8, dimension(:,:), allocatable **ro_bmp_spt**
- real *8, dimension(:,:), allocatable **ro_bmp_pnt**
- real *8, dimension(:,:), allocatable **ro_bmp_snt**
- real *8, dimension(:), allocatable **bmp_flo**
- real *8, dimension(:), allocatable **bmp_sed**
- real *8, dimension(:), allocatable **bmp_bac**
- real *8, dimension(:), allocatable **bmp_pp**
- real *8, dimension(:), allocatable **bmp_sp**
- real *8, dimension(:), allocatable **bmp_pn**
- real *8, dimension(:), allocatable **bmp_sn**
- real *8, dimension(:), allocatable **bmp_flag**

- real *8, dimension(:), allocatable **bmp_flos**
- real *8, dimension(:), allocatable **bmp_seds**
- real *8, dimension(:), allocatable **bmp_bacs**
- real *8, dimension(:), allocatable **bmp_pps**
- real *8, dimension(:), allocatable **bmp_sps**
- real *8, dimension(:), allocatable **bmp_pns**
- real *8, dimension(:), allocatable **bmp_sns**
- real *8, dimension(:), allocatable **bmp_flot**
- real *8, dimension(:), allocatable **bmp_sedt**
- real *8, dimension(:), allocatable **bmp_bact**
- real *8, dimension(:), allocatable **bmp_ppt**
- real *8, dimension(:), allocatable **bmp_spt**
- real *8, dimension(:), allocatable **bmp_pnt**
- real *8, dimension(:), allocatable **bmp_snt**
- real *8, dimension(:, :), allocatable **ntp_addon**
the distance between spillway levels (m)
- real *8, dimension(:, :), allocatable **ntp_cdis**
discharge coefficient for weir/orifice flow (none)
- real *8, dimension(:, :), allocatable **ntp_depweir**
depth of rectangular weir at different stages (m)
- real *8, dimension(:, :), allocatable **ntp_diaweir**
diameter of orifice hole at different stages (m)
- real *8, dimension(:, :), allocatable **ntp_flowrate**
maximum discharge from each stage of the weir/hole (m^3/s)
- real *8, dimension(:, :), allocatable **ntp_pcp**
precipitation for different return periods (not used) (mm)
- real *8, dimension(:, :), allocatable **ntp_retperd**
return period at different stages (years)
- real *8, dimension(:, :), allocatable **ntp_wdratio**
width depth ratio of rectangular weirs (none)
- real *8, dimension(:, :), allocatable **ntp_wrwid**
- real *8, dimension(:), allocatable **ri_subkm**
- real *8, dimension(:), allocatable **ri_totpv**
- real *8, dimension(:), allocatable **irmmdt**
- real *8, dimension(:, :), allocatable **ri_sed**
- real *8, dimension(:, :), allocatable **ri_fr**
- real *8, dimension(:, :), allocatable **ri_dim**
- real *8, dimension(:, :), allocatable **ri_im**
- real *8, dimension(:, :), allocatable **ri_iz**
- real *8, dimension(:, :), allocatable **ri_sa**
- real *8, dimension(:, :), allocatable **ri_vol**
- real *8, dimension(:, :), allocatable **ri_qi**
- real *8, dimension(:, :), allocatable **ri_k**
- real *8, dimension(:, :), allocatable **ri_dd**
- real *8, dimension(:, :), allocatable **ri_evrsv**
- real *8, dimension(:, :), allocatable **ri_dep**
- real *8, dimension(:, :), allocatable **ri_ndt**
- real *8, dimension(:, :), allocatable **ri_pmpvol**
- real *8, dimension(:, :), allocatable **ri_sed_cumul**
- real *8, dimension(:, :), allocatable **hrnopcp**
- real *8, dimension(:, :), allocatable **ri_qloss**
- real *8, dimension(:, :), allocatable **ri_pumpv**
- real *8, dimension(:, :), allocatable **ri_sedi**

- character(len=4), dimension(:,:), allocatable **ri_nirr**
- integer, dimension(:), allocatable **num_ri**
- integer, dimension(:), allocatable **ri_luflg**
- integer, dimension(:), allocatable **num_noirr**
- integer, dimension(:), allocatable **wtp_subnum**
- integer, dimension(:), allocatable **wtp_onoff**
- integer, dimension(:), allocatable **wtp_imo**
- integer, dimension(:), allocatable **wtp_iyr**
- integer, dimension(:), allocatable **wtp_dim**
- integer, dimension(:), allocatable **wtp_stagdis**
- integer, dimension(:), allocatable **wtp_sdtype**
- real *8, dimension(:), allocatable **wtp_pvol**
- real *8, dimension(:), allocatable **wtp_pdepth**
- real *8, dimension(:), allocatable **wtp_sdslope**
- real *8, dimension(:), allocatable **wtp_lenwidth**
- real *8, dimension(:), allocatable **wtp_extdepth**
- real *8, dimension(:), allocatable **wtp_hydeff**
- real *8, dimension(:), allocatable **wtp_evrsv**
- real *8, dimension(:), allocatable **wtp_sdintc**
- real *8, dimension(:), allocatable **wtp_sdexp**
- real *8, dimension(:), allocatable **wtp_sdc1**
- real *8, dimension(:), allocatable **wtp_sdc2**
- real *8, dimension(:), allocatable **wtp_sdc3**
- real *8, dimension(:), allocatable **wtp_pdia**
- real *8, dimension(:), allocatable **wtp_plen**
- real *8, dimension(:), allocatable **wtp_pmann**
- real *8, dimension(:), allocatable **wtp_ploss**
- real *8, dimension(:), allocatable **wtp_k**
- real *8, dimension(:), allocatable **wtp_dp**
- real *8, dimension(:), allocatable **wtp_sedi**
- real *8, dimension(:), allocatable **wtp_sede**
- real *8, dimension(:), allocatable **wtp_qi**
- real *8 **lai_init**
initial leaf area index of transplants
- real *8 **bio_init**
initial biomass of transplants (kg/ha)
- real *8 **cnop**
SCS runoff curve number for moisture condition II (none)
- real *8 **harveff**
harvest efficiency: fraction of harvested yield that is removed from HRU; the remainder becomes residue on the soil surface(none)
- real *8 **hi_ovr**
harvest index target specified at harvest ((kg/ha)/(kg/ha))
- real *8 **frac_harvk**
- real *8 **lid_vgcl**
van Genuchten equation's coefficient, l (none)
- real *8 **lid_vgcm**
van Genuchten equation's coefficient, m (none)
- real *8 **lid_qsurf_total**
- real *8 **lid_farea_sum**
- real *8, dimension(:,:), allocatable **lid_cuminf_last**
cumulative amount of water infiltrated into the amended soil layer at the last time step in a day (mm H2O)
- real *8, dimension(:,:), allocatable **lid_cumr_last**

- cumulative amount of rainfall at the last time step in a day (mm H2O)*
- real *8, dimension(:,:), allocatable [lid_excum_last](#)
- cumulative amount of excess rainfall at the last time step in a day (mm H2O)*
- real *8, dimension(:,:), allocatable [lid_f_last](#)
- potential infiltration rate of the amended soil layer at the last time step in a day (mm/mm H2O)*
- real *8, dimension(:,:), allocatable [lid_sw_last](#)
- soil water content of the amended soil layer at the last time step in a day (mm/mm H2O)*
- real *8, dimension(:,:), allocatable [lid_qsurf](#)
- depth of runoff generated on a LID in a given time interval (mm H2O)*
- real *8, dimension(:,:), allocatable [interval_last](#)
- real *8, dimension(:,:), allocatable [lid_str_last](#)
- real *8, dimension(:,:), allocatable [lid_farea](#)
- real *8, dimension(:,:), allocatable [lid_sw_add](#)
- real *8, dimension(:,:), allocatable [lid_cumqperc_last](#)
- real *8, dimension(:,:), allocatable [lid_cumirr_last](#)
- integer, dimension(:,:), allocatable [gr_onoff](#)
- integer, dimension(:,:), allocatable [gr_imo](#)
- integer, dimension(:,:), allocatable [gr_iyr](#)
- real *8, dimension(:,:), allocatable [gr_farea](#)
- fractional area of a green roof to the HRU (none)*
- real *8, dimension(:,:), allocatable [gr_solop](#)
- real *8, dimension(:,:), allocatable [gr_etcoef](#)
- real *8, dimension(:,:), allocatable [gr_fc](#)
- real *8, dimension(:,:), allocatable [gr_wp](#)
- real *8, dimension(:,:), allocatable [gr_ksat](#)
- real *8, dimension(:,:), allocatable [gr_por](#)
- real *8, dimension(:,:), allocatable [gr_hydeff](#)
- real *8, dimension(:,:), allocatable [gr_soldpt](#)
- integer, dimension(:,:), allocatable [rg_onoff](#)
- integer, dimension(:,:), allocatable [rg_imo](#)
- integer, dimension(:,:), allocatable [rg_iyr](#)
- real *8, dimension(:,:), allocatable [rg_farea](#)
- real *8, dimension(:,:), allocatable [rg_solop](#)
- real *8, dimension(:,:), allocatable [rg_etcoef](#)
- real *8, dimension(:,:), allocatable [rg_fc](#)
- real *8, dimension(:,:), allocatable [rg_wp](#)
- real *8, dimension(:,:), allocatable [rg_ksat](#)
- real *8, dimension(:,:), allocatable [rg_por](#)
- real *8, dimension(:,:), allocatable [rg_hydeff](#)
- real *8, dimension(:,:), allocatable [rg_soldpt](#)
- real *8, dimension(:,:), allocatable [rg_dimop](#)
- real *8, dimension(:,:), allocatable [rg_sarea](#)
- real *8, dimension(:,:), allocatable [rg_vol](#)
- real *8, dimension(:,:), allocatable [rg_sth](#)
- real *8, dimension(:,:), allocatable [rg_sdia](#)
- real *8, dimension(:,:), allocatable [rg_bdia](#)
- real *8, dimension(:,:), allocatable [rg_sts](#)
- real *8, dimension(:,:), allocatable [rg_orifice](#)
- real *8, dimension(:,:), allocatable [rg_oheight](#)
- real *8, dimension(:,:), allocatable [rg_odia](#)
- integer, dimension(:,:), allocatable [cs_onoff](#)
- integer, dimension(:,:), allocatable [cs_imo](#)
- integer, dimension(:,:), allocatable [cs_iyr](#)
- integer, dimension(:,:), allocatable [cs_grcon](#)

- real *8, dimension(:,:), allocatable **cs_farea**
- real *8, dimension(:,:), allocatable **cs_vol**
- real *8, dimension(:,:), allocatable **cs_rdepth**
- integer, dimension(:,:), allocatable **pv_onoff**
- integer, dimension(:,:), allocatable **pv_imo**
- integer, dimension(:,:), allocatable **pv_iyr**
- integer, dimension(:,:), allocatable **pv_solop**
- real *8, dimension(:,:), allocatable **pv_grvdep**
- real *8, dimension(:,:), allocatable **pv_grvpor**
- real *8, dimension(:,:), allocatable **pv_farea**
- real *8, dimension(:,:), allocatable **pv_drcoef**
- real *8, dimension(:,:), allocatable **pv_fc**
- real *8, dimension(:,:), allocatable **pv_wp**
- real *8, dimension(:,:), allocatable **pv_ksat**
- real *8, dimension(:,:), allocatable **pv_por**
- real *8, dimension(:,:), allocatable **pv_hydeff**
- real *8, dimension(:,:), allocatable **pv_soldpt**
- integer, dimension(:,:), allocatable **lid_onoff**
- real *8, dimension(:,:), allocatable **sol_bmc**
- real *8, dimension(:,:), allocatable **sol_bmn**
- real *8, dimension(:,:), allocatable **sol_hsc**
- real *8, dimension(:,:), allocatable **sol_hsn**
- real *8, dimension(:,:), allocatable **sol_hpc**
- real *8, dimension(:,:), allocatable **sol_hpn**
- real *8, dimension(:,:), allocatable **sol_lm**
- real *8, dimension(:,:), allocatable **sol_lmc**
- real *8, dimension(:,:), allocatable **sol_lmn**
- real *8, dimension(:,:), allocatable **sol_ls**
- real *8, dimension(:,:), allocatable **sol_lsl**
- real *8, dimension(:,:), allocatable **sol_lsc**
- real *8, dimension(:,:), allocatable **sol_lsn**
- real *8, dimension(:,:), allocatable **sol_rnmn**
- real *8, dimension(:,:), allocatable **sol_lslc**
- real *8, dimension(:,:), allocatable **sol_lslnc**
- real *8, dimension(:,:), allocatable **sol_rspc**
- real *8, dimension(:,:), allocatable **sol_woc**
- real *8, dimension(:,:), allocatable **sol_won**
- real *8, dimension(:,:), allocatable **sol_hp**
- real *8, dimension(:,:), allocatable **sol_hs**
- real *8, dimension(:,:), allocatable **sol_bm**
- real *8, dimension(:,:), allocatable **sol_cac**
- real *8, dimension(:,:), allocatable **sol_cec**
- real *8, dimension(:,:), allocatable **sol_percc**
- real *8, dimension(:,:), allocatable **sol_latc**
- real *8, dimension(:), allocatable **sedc_d**
- real *8, dimension(:), allocatable **surfqc_d**
- real *8, dimension(:), allocatable **latc_d**
- real *8, dimension(:), allocatable **percc_d**
- real *8, dimension(:), allocatable **foc_d**
- real *8, dimension(:), allocatable **nppc_d**
- real *8, dimension(:), allocatable **rsdc_d**
- real *8, dimension(:), allocatable **grainc_d**
- real *8, dimension(:), allocatable **stoverc_d**
- real *8, dimension(:), allocatable **soc_d**
- real *8, dimension(:), allocatable **rspc_d**

- real *8, dimension(:), allocatable **emitc_d**
- real *8, dimension(:), allocatable **sub_sedc_d**
- real *8, dimension(:), allocatable **sub_surfqc_d**
- real *8, dimension(:), allocatable **sub_latc_d**
- real *8, dimension(:), allocatable **sub_percc_d**
- real *8, dimension(:), allocatable **sub_foc_d**
- real *8, dimension(:), allocatable **sub_nppc_d**
- real *8, dimension(:), allocatable **sub_rsd_c_d**
- real *8, dimension(:), allocatable **sub_grainc_d**
- real *8, dimension(:), allocatable **sub_stoverc_d**
- real *8, dimension(:), allocatable **sub_emitc_d**
- real *8, dimension(:), allocatable **sub_soc_d**
- real *8, dimension(:), allocatable **sub_rspc_d**
- real *8, dimension(:), allocatable **sedc_m**
- real *8, dimension(:), allocatable **surfqc_m**
- real *8, dimension(:), allocatable **latc_m**
- real *8, dimension(:), allocatable **percc_m**
- real *8, dimension(:), allocatable **foc_m**
- real *8, dimension(:), allocatable **nppc_m**
- real *8, dimension(:), allocatable **rsdc_m**
- real *8, dimension(:), allocatable **grainc_m**
- real *8, dimension(:), allocatable **stoverc_m**
- real *8, dimension(:), allocatable **emitc_m**
- real *8, dimension(:), allocatable **soc_m**
- real *8, dimension(:), allocatable **rspc_m**
- real *8, dimension(:), allocatable **sedc_a**
- real *8, dimension(:), allocatable **surfqc_a**
- real *8, dimension(:), allocatable **latc_a**
- real *8, dimension(:), allocatable **percc_a**
- real *8, dimension(:), allocatable **foc_a**
- real *8, dimension(:), allocatable **nppc_a**
- real *8, dimension(:), allocatable **rsdc_a**
- real *8, dimension(:), allocatable **grainc_a**
- real *8, dimension(:), allocatable **stoverc_a**
- real *8, dimension(:), allocatable **emitc_a**
- real *8, dimension(:), allocatable **soc_a**
- real *8, dimension(:), allocatable **rspc_a**
- integer, dimension(:), allocatable **tillage_switch**
- real *8, dimension(:), allocatable **tillage_depth**
- integer, dimension(:), allocatable **tillage_days**
- real *8, dimension(:), allocatable **tillage_factor**
- real *8 **dthy**
time interval for subdaily routing
- integer, dimension(4) **ihx**
- integer, dimension(:), allocatable **nhy**
- real *8, dimension(:), allocatable **rchx**
- real *8, dimension(:), allocatable **rcss**
- real *8, dimension(:), allocatable **qcap**
- real *8, dimension(:), allocatable **chxa**
- real *8, dimension(:), allocatable **chxp**
- real *8, dimension(:, :, :), allocatable **qhy**
- real *8 **ff1**
- real *8 **ff2**

5.1.1 Detailed Description

main module containing the global variables

5.1.2 Variable Documentation

5.1.2.1 igropt

`integer parm::igropt`

Qual2E option for calculating the local specific growth rate of algae
1: multiplicative.

$$u = \text{mumax} \text{ fll } f_{nn} \text{ fpp}$$

2: limiting nutrient

$$u = \text{mumax} \text{ fll } \min(f_{nn}, f_{pp})$$

3: harmonic mean

$$u = \text{mumax} \text{ fll } \frac{2}{\frac{1}{f_{nn}} + \frac{1}{f_{pp}}}$$

Chapter 6

Data Type Documentation

6.1 parm::fcgd Interface Reference

Public Member Functions

- real *8 function **fcgd** (xx)

The documentation for this interface was generated from the following file:

- [modparm.f90](#)

6.2 parm::ndenit Interface Reference

Public Member Functions

- subroutine **ndenit** (k, j, cdg, wdn, void)

The documentation for this interface was generated from the following file:

- [modparm.f90](#)

6.3 parm::regres Interface Reference

Public Member Functions

- real *8 function **regres** (k, j)

The documentation for this interface was generated from the following file:

- [modparm.f90](#)

6.4 parm::tair Interface Reference

Public Member Functions

- real *8 function **tair** (hr, jj)

The documentation for this interface was generated from the following file:

- [modparm.f90](#)

6.5 parm::theta Interface Reference

Public Member Functions

- real *8 function **theta** (r20, thk, tmp)

The documentation for this interface was generated from the following file:

- [modparm.f90](#)

Chapter 7

File Documentation

7.1 albedo.f90 File Reference

Functions/Subroutines

- subroutine [albedo](#)
this subroutine calculates albedo in the HRU for the day

7.1.1 Detailed Description

file containing the subroutine albedo

Author

modified by Javier Burguete

7.2 allocate_parms.f90 File Reference

Functions/Subroutines

- subroutine [allocate_parms](#)
this subroutine allocates array sizes

7.2.1 Detailed Description

file containing the subroutine allocate_parms

Author

modified by Javier Burguete

7.3 alph.f90 File Reference

Functions/Subroutines

- subroutine [alph](#) (iwave, j)

*this subroutine computes alpha, a dimensionless parameter that expresses the fraction of total rainfall that occurs during 0.5h @parm[in] iwave flag to differentiate calculation of HRU and subbasin sediment calculation (none)
iwave = 0 for HRU MUSLE(sedyld) each hru is calculated independently using hru area and adjusted channel length
iwave = 1 subbasin # for subbasin MUSLE is computed for entire subbasin using hru weighted KLSCP*

7.3.1 Detailed Description

file containing the subroutine alph

Author

modified by Javier Burguete

7.3.2 Function/Subroutine Documentation

7.3.2.1 alph()

```
subroutine alph (
    integer, intent(in) iwave,
    integer, intent(in) j )
```

this subroutine computes alpha, a dimensionless parameter that expresses the fraction of total rainfall that occurs during 0.5h @parm[in] iwave flag to differentiate calculation of HRU and subbasin sediment calculation (none)
iwave = 0 for HRU MUSLE(sedyld) each hru is calculated independently using hru area and adjusted channel length
iwave = 1 subbasin # for subbasin MUSLE is computed for entire subbasin using hru weighted KLSCP

Parameters

in	j	HRU number
----	---	------------

7.4 apply.f90 File Reference

Functions/Subroutines

- subroutine [apply](#) (j)

this subroutine applies pesticide

7.4.1 Detailed Description

file containing the subroutine apply

Author

modified by Javier Burguete

7.4.2 Function/Subroutine Documentation

7.4.2.1 apply()

```
subroutine apply (  
    integer, intent(in) j )
```

this subroutine applies pesticide

Parameters

in	j	HRU number
----	---	------------

7.5 ascrv.f90 File Reference

Functions/Subroutines

- subroutine [ascrv](#) (x1, x2, x3, x4, x5, x6)
this subroutine computes shape parameters x5 and x6 for the S curve equation

7.5.1 Detailed Description

file containing the subroutine ascrv

Author

modified by Javier Burguete

7.5.2 Function/Subroutine Documentation

7.5.2.1 ascrv()

```
subroutine ascrv (
    real*8, intent(in) x1,
    real*8, intent(in) x2,
    real*8, intent(in) x3,
    real*8, intent(in) x4,
    real*8, intent(out) x5,
    real*8, intent(out) x6 )
```

this subroutine computes shape parameters x5 and x6 for the S curve equation

$$x = \frac{y}{y + \exp(x5 + x6 y)}$$

given 2 (x,y) points along the curve. x5 is determined by solving the equation with x and y values measured around the midpoint of the curve (approx. 50% of the maximum value for x) and x6 is determined by solving the equation with x and y values measured close to one of the endpoints of the curve (100% of the maximum value for x). This subroutine is called from [readbsn.f90](#) and [readplant.f90](#)

Parameters

in	x1	value for x in the above equation for first datapoint, x1 should be close to 0.5 (the midpoint of the curve)
in	x2	value for x in the above equation for second datapoint, x2 should be close to 0.0 or 1.0
in	x3	value for y in the above equation corresponding to x1
in	x4	value for y in the above equation corresponding to x2
out	x5	1st shape parameter for S curve equation characterizing the midpoint of the curve
out	x6	2nd shape parameter for S curve equation characterizing the regions close to the endpoints of the curve

7.6 atri.f90 File Reference

Functions/Subroutines

- real *8 function [atri](#) (at1, at2, at3, at4i)

this function generates a random number from a triangular distribution given X axis points at start, end, and peak Y value

7.6.1 Detailed Description

file containing the function atri

Author

modified by Javier Burguete

7.6.2 Function/Subroutine Documentation

7.6.2.1 atri()

```
real*8 function atri (
    real*8, intent(in) at1,
    real*8, intent(in) at2,
    real*8, intent(in) at3,
    integer, intent(inout) at4i )
```

this function generates a random number from a triangular distribution given X axis points at start, end, and peak Y value

Parameters

in	at1	lower limit for distribution (none)
in	at2	monthly mean for distribution (none)
in	at3	upper limit for distribution (none)
in, out	at4i	random number seed (none)

Returns

daily value generated for distribution (none)

7.7 aunif.f90 File Reference

Functions/Subroutines

- real *8 function [aunif](#) (x1)

This function generates random numbers ranging from 0.0 to 1.0. In the process of calculating the random number, the seed (x1) is set to a new value. This function implements the prime-modulus generator.

7.7.1 Detailed Description

file containing the function aunif

Author

modified by Javier Burguete

7.7.2 Function/Subroutine Documentation

7.7.2.1 aunif()

```
real*8 function aunif (
    integer, intent(inout) x1 )
```

This function generates random numbers ranging from 0.0 to 1.0. In the process of calculating the random number, the seed (x1) is set to a new value. This function implements the prime-modulus generator.

$$xi = 16807 xi \bmod (2^{31} - 1)$$

using code which ensures that no intermediate result uses more than 31 bits. The theory behind the code is summarized in [1]

Parameters

in, out	<i>x1</i>	random number generator seed (integer) where $0 < x1 < 2147483647$
---------	-----------	--

Returns

random number ranging from 0.0 to 1.0

7.8 bmpinit.f90 File Reference

Functions/Subroutines

- subroutine [bmpinit](#) (*ii*)
this subroutine sets default values for urban bmp parameters

7.8.1 Detailed Description

file containing the subroutine bmpinit

Author

modified by Javier Burguete

7.8.2 Function/Subroutine Documentation

7.8.2.1 bmpinit()

```
subroutine bmpinit (
    integer, intent(in) ii )
```

this subroutine sets default values for urban bmp parameters

Parameters

in	<i>ii</i>	subbasin number
----	-----------	-----------------

7.9 burnop.f90 File Reference

Functions/Subroutines

- subroutine [burnop](#) (*j*)
this subroutine performs burning

7.9.1 Detailed Description

file containing the subroutine burnop

Author

modified by Javier Burquete

7.9.2 Function/Subroutine Documentation

7.9.2.1 burnop()

```
subroutine burnop (  
    integer, intent(in) j )
```

this subroutine performs burning

Parameters

in	j	HRU number
----	---	------------

7.10 canopyint.f90 File Reference

Functions/Subroutines

- subroutine [canopyint](#)
this subroutine computes canopy interception of rainfall used for methods other than curve number

7.10.1 Detailed Description

file containing the subroutine canopyint

Author

modified by Javier Burquete

7.11 caps.f90 File Reference

Functions/Subroutines

- subroutine [caps](#) (file_name)
this subroutine reads the input and output names given in file.cio and converts all capital letters to lowercase letters.

7.11.1 Detailed Description

file containing the subroutine caps

Author

modified by Javier Burguete

7.11.2 Function/Subroutine Documentation

7.11.2.1 caps()

```
subroutine caps (
    character (len=*) file_name )
```

this subroutine reads the input and output names given in file.cio and converts all capital letters to lowercase letters.

Parameters

<i>file_name</i>	dummy argument, file name character string
------------------	--

7.12 cfactor.f90 File Reference

Functions/Subroutines

- subroutine [cfactor](#)
this subroutine predicts daily soil loss caused by water erosion using the modified universal soil loss equation

7.12.1 Detailed Description

file containing the subroutine cfactor

Author

modified by Javier Burguete

7.13 clgen.f90 File Reference

Functions/Subroutines

- subroutine [clgen](#) (j)
this subroutine calculates the daylength, distribution of radiation throughout the day and maximum radiation for day

7.13.1 Detailed Description

file containing the subroutine clgen

Author

modified by Javier Burguete

7.13.2 Function/Subroutine Documentation

7.13.2.1 clgen()

```
subroutine clgen (
    integer, intent(in) j )
```

this subroutine calculates the daylength, distribution of radiation throughout the day and maximum radiation for day

Parameters

in	j	HRU number
----	---	------------

7.14 clicon.f90 File Reference

Functions/Subroutines

- subroutine [clicon](#) (i)

this subroutine controls weather inputs to SWAT. Precipitation and temperature data is read in and the weather generator is called to fill in radiation, wind speed and relative humidity as well as missing precipitation and temperatures. Adjustments for climate changes studies are also made in this subroutine.

7.14.1 Detailed Description

file containing the subroutine clicon

Author

modified by Javier Burguete

7.14.2 Function/Subroutine Documentation

7.14.2.1 clicon()

```
subroutine clicon (
    integer, intent(in) i )
```

this subroutine controls weather inputs to SWAT. Precipitation and temperature data is read in and the weather generator is called to fill in radiation, wind speed and relative humidity as well as missing precipitation and temperatures. Adjustments for climate changes studies are also made in this subroutine.

Parameters

in	i	current day of simulation (julian date)
----	---	---

7.15 command.f90 File Reference

Functions/Subroutines

- subroutine [command](#) (i)

for every day of simulation, this subroutine steps through the command lines in the watershed configuration (.fig) file. Depending on the command code on the .fig file line, a command loop is accessed

7.15.1 Detailed Description

file containing the subroutine command

Author

modified by Javier Burguete

7.15.2 Function/Subroutine Documentation

7.15.2.1 command()

```
subroutine command (
    integer, intent(in) i )
```

for every day of simulation, this subroutine steps through the command lines in the watershed configuration (.fig) file. Depending on the command code on the .fig file line, a command loop is accessed

Parameters

in	i	current day in simulation-loop counter (julian date)
----	---	--

7.16 crackflow.f90 File Reference

Functions/Subroutines

- subroutine [crackflow](#)
this subroutine modifies surface runoff to account for crack flow

7.16.1 Detailed Description

file containing the subroutine crackflow

Author

modified by Javier Burguete

7.17 crackvol.f90 File Reference

Functions/Subroutines

- subroutine [crackvol](#)
this subroutine computes total crack volume for the soil profile and modifies surface runoff to account for crack flow

7.17.1 Detailed Description

file containing the subroutine crackvol

Author

modified by Javier Burguete

7.18 curno.f90 File Reference

Functions/Subroutines

- subroutine [curno](#) (cnn, h)
*this subroutine determines the curve numbers for moisture conditions I and III and calculates coefficients and shape parameters for the water retention curve. The coefficients and shape parameters are calculated by one of two methods:
the default method is to make them a function of soil water,
the alternative method (labeled new) is to make them a function of accumulated PET, precipitation and surface runoff*

7.18.1 Detailed Description

file containing the subroutine curno

Author

modified by Javier Burguete

7.18.2 Function/Subroutine Documentation

7.18.2.1 curno()

```
subroutine curno (
    real*8, intent(in) cnn,
    integer, intent(in) h )
```

this subroutine determines the curve numbers for moisture conditions I and III and calculates coefficients and shape parameters for the water retention curve. The coefficients and shape parameters are calculated by one of two methods:

the default method is to make them a function of soil water,

the alternative method (labeled new) is to make them a function of accumulated PET, precipitation and surface runoff

Parameters

in	<i>cnn</i>	SCS runoff curve number for moisture condition II
in	<i>h</i>	HRU number

7.19 dailycn.f90 File Reference

Functions/Subroutines

- subroutine [dailycn](#)
calculates curve number for the day in the HRU

7.19.1 Detailed Description

file containing the subroutine dailycn

Author

modified by Javier Burguete

7.20 dstn1.f90 File Reference

Functions/Subroutines

- real *8 function [dstn1](#) (rn1, rn2)
this function computes the distance from the mean of a normal distribution with mean = 0 and standard deviation = 1, given two random numbers

7.20.1 Detailed Description

file containing the function dstn1

Author

modified by Javier Burguete

7.20.2 Function/Subroutine Documentation

7.20.2.1 dstn1()

```
real*8 function dstn1 (  
    real*8, intent(in) rn1,  
    real*8, intent(in) rn2 )
```

this function computes the distance from the mean of a normal distribution with mean = 0 and standard deviation = 1, given two random numbers

Parameters

in	<i>rn1</i>	first random number
in	<i>rn2</i>	second random number

Returns

distance from the mean

7.21 ee.f90 File Reference

Functions/Subroutines

- real *8 function ee (tk)
this function calculates saturation vapor pressure at a given air temperature

7.21.1 Detailed Description

file containing the function ee

Author

modified by Javier Burguete

7.21.2 Function/Subroutine Documentation

7.21.2.1 ee()

```
real*8 function ee (
    real*8, intent(in) tk )
```

this function calculates saturation vapor pressure at a given air temperature

Parameters

in	tk	mean air temperature (deg C)
----	----	------------------------------

Returns

saturation vapor pressure (kPa)

7.22 eiusle.f90 File Reference

Functions/Subroutines

- subroutine [eiusle](#) (j)
this subroutine computes the USLE erosion index (EI)

7.22.1 Detailed Description

file containing the subroutine eiusle

Author

modified by Javier Burguete

7.23 estimate_ksat.f90 File Reference

Functions/Subroutines

- subroutine [estimate_ksat](#) (perc_clay, esti_ksat)
This subroutine calculates ksat value for a soil layer given the % of clay in the soil layer.

7.23.1 Detailed Description

file containing the subroutine estimate_ksat

Author

modified by Javier Burguete

7.23.2 Function/Subroutine Documentation

7.23.2.1 estimate_ksat()

```
subroutine estimate_ksat (
    real*8, intent(in) perc_clay,
    real*8, intent(out) esti_ksat )
```

This subroutine calculates ksats value for a soil layer given the % of clay in the soil layer.

Background: published work of Walter Rawls. Calculated ksats values based on soil texture (sand, silt and clay).
Idea: there exists a relationship between % clay and Ksat. Equations used in this subroutine are based on the above idea (Jimmy Willimas)

Parameters

in	<i>perc_clay</i>	clay percentage (%)
out	<i>esti_ksat</i>	estimated ksats

7.24 etpot.f90 File Reference

Functions/Subroutines

- subroutine [etpot](#) (j)
this subroutine calculates potential evapotranspiration using one of three methods. If Penman-Monteith is being used, potential plant transpiration is also calculated.

7.24.1 Detailed Description

file containing the subroutine etpot

Author

modified by Javier Burguete

7.24.2 Function/Subroutine Documentation

7.24.2.1 etpot()

```
subroutine etpot (
    integer, intent(in) j )
```

this subroutine calculates potential evapotranspiration using one of three methods. If Penman-Monteith is being used, potential plant transpiration is also calculated.

Parameters

in	j	HRU number
----	---	------------

7.25 expo.f90 File Reference

Functions/Subroutines

- real *8 function `expo` (xx)

this function checks the argument against upper and lower boundary values prior to taking the Exponential

7.25.1 Detailed Description

file containing the function expo

Author

modified by Javier Burguete

7.25.2 Function/Subroutine Documentation

7.25.2.1 `expo()`

```
real*8 function expo (
    real*8, intent(in) xx )
```

this function checks the argument against upper and lower boundary values prior to taking the Exponential

Parameters

in	xx	exponential argument (none)
----	----	-----------------------------

Returns

$\exp(xx)$

7.26 fert.f90 File Reference

Functions/Subroutines

- subroutine `fert` (j)

this subroutine applies N and P specified by date and amount in the management file (.mgt)

7.26.1 Detailed Description

file containing the subroutine fert

Author

modified by Javier Burguete

7.26.2 Function/Subroutine Documentation

7.26.2.1 fert()

```
subroutine fert (
    integer, intent(in) j )
```

this subroutine applies N and P specified by date and amount in the management file (.mgt)

Parameters

in	j	HRU number
----	---	------------

7.27 finalbal.f90 File Reference

Functions/Subroutines

- subroutine [finalbal](#)
this subroutine calculates final water balance for watershed

7.27.1 Detailed Description

file containing the subroutine finalbal

Author

modified by Javier Burguete

7.28 gcycl.f90 File Reference

Functions/Subroutines

- subroutine [gcycl](#)
This subroutine initializes the random number seeds. If the user desires a different set of random numbers for each simulation run, the random number generator is used to reset the values of the seeds.

7.28.1 Detailed Description

file containing the subroutine gcycl

Author

modified by Javier Burguete

7.29 getallo.f90 File Reference

Functions/Subroutines

- subroutine [getallo](#)

This subroutine calculates the number of HRUs, subbasins, etc. in the simulation. These values are used to allocate array sizes.

7.29.1 Detailed Description

file containing the subroutine getallo

Author

modified by Javier Burguete

7.30 graze.f90 File Reference

Functions/Subroutines

- subroutine [graze](#) (j)

this subroutine simulates biomass lost to grazing

7.30.1 Detailed Description

file containing the subroutine graze

Author

modified by Javier Burguete

7.30.2 Function/Subroutine Documentation

7.30.2.1 graze()

```
subroutine graze (  
    integer, intent(in) j )
```

this subroutine simulates biomass lost to grazing

Parameters

in	j	HRU number
----	---	------------

7.31 h2omgt_init.f90 File Reference

Functions/Subroutines

- subroutine [h2omgt_init](#)

This subroutine initializes variables related to water management (irrigation, consumptive water use, etc.)

7.31.1 Detailed Description

file containing the subroutine h2omgt_init

Author

modified by Javier Burguete

7.32 harvestop.f90 File Reference

Functions/Subroutines

- subroutine [harvestop](#) (j)

this subroutine performs the harvest operation (no kill)

7.32.1 Detailed Description

file containing the subroutine harvestop

Author

modified by Javier Burguete

7.32.2 Function/Subroutine Documentation

7.32.2.1 harvestop()

```
subroutine harvestop (
    integer, intent(in) j )
```

this subroutine performs the harvest operation (no kill)

Parameters

in	j	HRU number
----	---	------------

7.33 harvkillop.f90 File Reference

Functions/Subroutines

- subroutine [harvkillop](#) (j)
this subroutine performs the harvest and kill operation

7.33.1 Detailed Description

file containing the subroutine harvkillop

Author

modified by Javier Burguete

7.33.2 Function/Subroutine Documentation

7.33.2.1 harvkillop()

```
subroutine harvkillop (
    integer, intent(in) j )
```

this subroutine performs the harvest and kill operation

Parameters

in	j	HRU number
----	---	------------

7.34 headout.f90 File Reference

Functions/Subroutines

- subroutine [headout](#)
this subroutine writes the headings to the major output files

7.34.1 Detailed Description

file containing the subroutine headout

Author

modified by Javier Burguete

7.35 hmeas.f90 File Reference

Functions/Subroutines

- subroutine [hmeas](#)

this subroutine reads in relative humidity data from file and assigns the data to the HRUs

7.35.1 Detailed Description

file containing the subroutine hmeas

Author

modified by Javier Burguete

7.36 hruaa.f90 File Reference

Functions/Subroutines

- subroutine [hruaa](#) (years)

this subroutine writes average annual HRU output to the output.hru file

7.36.1 Detailed Description

file containing the subroutine hruaa

Author

modified by Javier Burguete

7.36.2 Function/Subroutine Documentation

7.36.2.1 hruaa()

```
subroutine hruaa (  
    real*8, intent(in) years )
```

this subroutine writes average annual HRU output to the output.hru file

Parameters

in	years	length of simulation (years)
----	-------	------------------------------

7.37 hruallo.f90 File Reference

Functions/Subroutines

- subroutine [hruallo](#)

This subroutine calculates the number of management operation types, etc. used in the simulation. These values are used to allocate array sizes for processes occurring in the HRU.

7.37.1 Detailed Description

file containing the subroutine hruallo

Author

modified by Javier Burguete

7.38 hrumon.f90 File Reference

Functions/Subroutines

- subroutine [hrumon](#)

this subroutine writes monthly HRU output to the output.hru file

7.38.1 Detailed Description

file containing the subroutine hrumon

Author

modified by Javier Burguete

7.39 hruyr.f90 File Reference

Functions/Subroutines

- subroutine [hruyr](#)

this subroutine writes annual HRU output to the output.hru file

7.39.1 Detailed Description

file containing the subroutine hrubr

Author

modified by Javier Burguete

7.40 hydroinit.f90 File Reference

Functions/Subroutines

- subroutine [hydroinit](#)

This subroutine computes variables related to the watershed hydrology: the time of concentration for the subbasins, lagged surface runoff, the coefficient for the peak runoff rate equation, and lateral flow travel time.

7.40.1 Detailed Description

file containing the subroutine hydroinit

Author

modified by Javier Burguete

7.41 icl.f90 File Reference

Functions/Subroutines

- integer function [icl](#) (id)

this function determines the month and day, given the julian date

7.41.1 Detailed Description

file containing the function icl

Author

modified by Javier Burguete

7.41.2 Function/Subroutine Documentation

7.41.2.1 icl()

```
integer function icl (  
    integer, intent(in) id )
```

this function determines the month and day, given the julian date

Parameters

in	id	julian date
----	----	-------------

7.42 impnd_init.f90 File Reference

Functions/Subroutines

- subroutine [impnd_init](#)
this subroutine initializes variables related to impoundments (ponds, wetlands, reservoirs and potholes)

7.42.1 Detailed Description

file containing the subroutine impnd_init

Author

modified by Javier Burguete

7.43 impndmon.f90 File Reference

Functions/Subroutines

- subroutine [impndmon](#)
this subroutine writes monthly HRU impoundment output to the output.wtr file

7.43.1 Detailed Description

file containing the subroutine impndmon

Author

modified by Javier Burguete

7.44 impndyr.f90 File Reference

Functions/Subroutines

- subroutine [impndyr](#)
this subroutine writes annual HRU impoundment output to the output.wtr file

7.44.1 Detailed Description

file containing the subroutine `impndyr`

Author

modified by Javier Burguete

7.45 irrigate.f90 File Reference

Functions/Subroutines

- subroutine `irrigate` (`j`, `volmm`)
this subroutine applies irrigation water to HRU

7.45.1 Detailed Description

file containing the subroutine `irrigate`

Author

modified by Javier Burguete

7.45.2 Function/Subroutine Documentation

7.45.2.1 irrigate()

```
subroutine irrigate (
    integer, intent(in) j,
    real*8, intent(in) volmm )
```

this subroutine applies irrigation water to HRU

Parameters

in	<i>j</i>	HRU number (none)
in	<i>volmm</i>	depth irrigation water applied to HRU (mm H2O)

7.46 irrsb.f90 File Reference

Functions/Subroutines

- subroutine `irrsb` (`j`)

this subroutine performs the irrigation operation when the source is the shallow or deep aquifer or a source outside the watershed

7.46.1 Detailed Description

file containing the subroutine irrsb

Author

modified by Javier Burguete

7.46.2 Function/Subroutine Documentation

7.46.2.1 irrsb()

```
subroutine irrsb (
    integer, intent(in) j )
```

this subroutine performs the irrigation operation when the source is the shallow or deep aquifer or a source outside the watershed

Parameters

in	j	HRU number (none)
----	---	-------------------

7.47 jdt.f90 File Reference

Functions/Subroutines

- integer function [jdt](#) (numdays, i, m)
this function computes the julian date given the month and the day of the month

7.47.1 Detailed Description

file containing the function jdt

Author

modified by Javier Burguete

7.47.2 Function/Subroutine Documentation

7.47.2.1 jdt()

```
integer function jdt (
    integer, dimension (13), intent(in) numdays,
    integer, intent(in) i,
    integer, intent(in) m )
```

this function computes the julian date given the month and the day of the month

Parameters

in	<i>numdays</i>	julian date for last day of preceding month (where the array location is the number of the month). The dates are for leap years (numdays=ndays) (julian date)
in	<i>i</i>	day
in	<i>m</i>	month

7.48 killop.f90 File Reference**Functions/Subroutines**

- subroutine [killop](#) (j)
this subroutine performs the kill operation

7.48.1 Detailed Description

file containing the subroutine killop

Author

modified by Javier Burguete

7.48.2 Function/Subroutine Documentation**7.48.2.1 killop()**

```
subroutine killop (
    integer, intent(in) j )
```

this subroutine performs the kill operation

Parameters

in	<i>j</i>	HRU number
----	----------	------------

7.49 lid_cistern.f90 File Reference

Functions/Subroutines

- subroutine [lid_cistern](#) (sb, j, k, lid_prec)
simulate cistern processes

7.49.1 Detailed Description

file containing the subroutine lid_cistern

Author

modified by Javier Burguete

7.49.2 Function/Subroutine Documentation

7.49.2.1 lid_cistern()

```
subroutine lid_cistern (
    integer, intent(in) sb,
    integer, intent(in) j,
    integer, intent(in) k,
    real*8, intent(in) lid_prec )
```

simulate cistern processes

Parameters

in	<i>sb</i>	subbasin number (none)
in	<i>j</i>	HRU number (none)
in	<i>k</i>	subdaily time index (none)
in	<i>lid_prec</i>	precipitation depth a LID receives in a simulation time interval (mm)

7.50 lid_greenroof.f90 File Reference

Functions/Subroutines

- subroutine [lid_greenroof](#) (sb, j, k, lid_prec)
simulate green roof processes

7.50.1 Detailed Description

file containing the subroutine lid_greenroof

Author

modified by Javier Burguete

7.50.2 Function/Subroutine Documentation

7.50.2.1 lid_greenroof()

```
subroutine lid_greenroof (
    integer, intent(in) sb,
    integer, intent(in) j,
    integer, intent(in) k,
    real*8, intent(in) lid_prec )
```

simulate green roof processes

Parameters

in	<i>sb</i>	subbasin number (none)
in	<i>j</i>	HRU number (none)
in	<i>k</i>	subdaily time index (none)
in	<i>lid_prec</i>	precipitation depth a LID receives in a simulation time interval (mm)

7.51 lid_porpavement.f90 File Reference

Functions/Subroutines

- subroutine [lid_porpavement](#) (sb, j, k, lid_prec)
simulate porous pavement processes

7.51.1 Detailed Description

file containing the subroutine lid_porpavement

Author

modified by Javier Burguete

7.51.2 Function/Subroutine Documentation

7.51.2.1 lid_porpavement()

```
subroutine lid_porpavement (
    integer, intent(in) sb,
    integer, intent(in) j,
    integer, intent(in) k,
    real*8, intent(in) lid_prec )
```

simulate porous pavement processes

Parameters

in	<i>sb</i>	subbasin number (none)
in	<i>j</i>	HRU number (none)
in	<i>k</i>	subdaily time index (none)
in	<i>lid_prec</i>	precipitation depth a LID receives in a simulation time interval (mm)

7.52 lid_raingarden.f90 File Reference

Functions/Subroutines

- subroutine [lid_raingarden](#) (sb, j, k, lid_prec)
simulate rain garden processes

7.52.1 Detailed Description

file containing the subroutine lid_raingarden

Author

modified by Javier Burguete

7.52.2 Function/Subroutine Documentation

7.52.2.1 lid_raingarden()

```
subroutine lid_raingarden (
    integer, intent(in) sb,
    integer, intent(in) j,
    integer, intent(in) k,
    real*8, intent(in) lid_prec )
```

simulate rain garden processes

Parameters

in	<i>sb</i>	subbasin number (none)
in	<i>j</i>	HRU number (none)
in	<i>k</i>	subdaily time index (none)
in	<i>lid_prec</i>	precipitation depth a LID receives in a simulation time interval (mm)

7.53 lidinit.f90 File Reference

Functions/Subroutines

- subroutine [lidinit](#) (i)
this subroutine sets default values for LID parameters

7.53.1 Detailed Description

file containing the subroutine lidinit

Author

modified by Javier Burguete

7.53.2 Function/Subroutine Documentation

7.53.2.1 lidinit()

```
subroutine lidinit (
    integer, intent(in) i )
```

this subroutine sets default values for LID parameters

Parameters

in	<i>i</i>	subbasin number
----	----------	-----------------

7.54 lids.f90 File Reference

Functions/Subroutines

- subroutine [lids](#) (sb, j, k, lid_prec)
call subroutines to simulate green roof, rain garden, cistern and porous pavement processes

7.54.1 Detailed Description

file containing the subroutine lids

Author

modified by Javier Burguete

7.54.2 Function/Subroutine Documentation

7.54.2.1 lids()

```
subroutine lids (
    integer, intent(in) sb,
    integer, intent(in) j,
    integer, intent(in) k,
    real*8, intent(in) lid_prec )
```

call subroutines to simulate green roof, rain garden, cistern and porous pavement processes

Parameters

in	<i>sb</i>	subbasin number (none)
in	<i>j</i>	HRU number (none)
in	<i>k</i>	subdaily time index (none)
in	<i>lid_prec</i>	precipitation depth a LID receives in a simulation time interval (mm)

7.55 lwqdef.f90 File Reference

Functions/Subroutines

- subroutine [lwqdef](#) (ii)
this subroutine assigns default values for the lake water quality (.lwq) when the lake water quality file does not exists

7.55.1 Detailed Description

file containing the subroutine lwqdef

Author

modified by Javier Burguete

7.55.2 Function/Subroutine Documentation

7.55.2.1 lwqdef()

```
subroutine lwqdef (
    integer, intent(in) ii )
```

this subroutine assigns default values for the lake water quality (.lwq) when the lake water quality file does not exists

Parameters

in	ii	reservoir number (none)
----	----	-------------------------

7.56 main.f90 File Reference

Functions/Subroutines

- program [main](#)
this is the main program that reads input, calls the main simulation model, and writes output

7.56.1 Detailed Description

file containing the main program that reads input, calls the main simulation model, and writes output.

Author

modified by Javier Burguete Tolosa

7.57 modparm.f90 File Reference

Data Types

- interface [parm::fcgd](#)
- interface [parm::regres](#)
- interface [parm::tair](#)
- interface [parm::theta](#)
- interface [parm::ndenit](#)

Modules

- module [parm](#)
main module containing the global variables

- character(len=13), dimension(40), parameter **parm::hedwtr** = (/ " PNDPCPmm", " PND_INmm", "PSED_↵
lt/ha", " PNDEVPmm", " PNDSEPmm", " PND_OUTmm", "PSED_Ot/ha", " PNDVOLm^3", "PNDORGNppm",
PNDNO3ppm", "PNDORGPPpm", "PNDMINPppm", "PNDCHLAppm", " PNDSECIIm", " WETPCPmm", " W↵
ET_INmm", "WSED_l/ha", " WETVPmm", " WETSEPmm", " WET_OUTmm", "WSED_Ot/ha", " WETVO↵
Lm^3", "WETORGNppm", " WETNO3ppm", "WETORGPPpm", "WETMINPppm", "WETCHLAppm", " WETSE↵
CIIm", " POTPCPmm", " POT_INmm", "OSED_l/ha", " POTEVPmm", " POTSEPmm", " POT_OUTmm", "OSE↵
D_Ot/ha", " POTVOLm^3", " POT_SAha", "HRU_SURQmm", "PLANT_ETmm", " SOIL_ETmm"/)
- *column headers for HRU impoundment output file*
- integer, dimension(mhruo), parameter **parm::icols** = (/43,53,63,73,83,93,103,113,123,133,143,153,163,173,183,193,203,213,2↵
space number for beginning of column in HRU output file (none)
- integer, dimension(msubo), parameter **parm::icolb** = (/35,45,55,65,75,85,95,105,115,125,135,145,155,165,175,185,195,205,2↵
space number for beginning of column in subbasin output file (none)
- integer, dimension(mrcho), parameter **parm::icolr** = (/38,50,62,74,86,98,110,122,134,146,158,170,182,194,206,218,230,242,2↵
space number for beginning of column in reach output file (none)
- integer, dimension(41), parameter **parm::icolsv** = (/38,50,62,74,86,98,110,122,134,146,158,170,182,194,206,218,230,242,25↵
space number for beginning of column in reservoir output file (none)
- real *8, parameter **parm::ab** = 0.02083
lowest value a15 can have (mm H2O)
- integer, dimension(13), parameter **parm::ndays_leap** = (/0,31,60,91,121,152,182,213,244,274,305,335,366/)
- integer, dimension(13), parameter **parm::ndays_noleap** = (/0,31,59,90,120,151,181,212,243,273,304,334,365/)
- integer **parm::icalen**
- real *8 **parm::prf_bsn**
*Basinwide peak rate adjustment factor for sediment routing in the channel. Allows impact of peak flow rate on
sediment routing and channel reshaping to be taken into account.*
- real *8 **parm::co2_x2**
- real *8 **parm::co2_x**
- real *8, dimension(:), allocatable **parm::alph_e**
- real *8, dimension(:), allocatable **parm::cdn**
denitrification exponential rate coefficient
- real *8, dimension(:), allocatable **parm::nperco**
nitrate percolation coefficient (0-1)
0:concentration of nitrate in surface runoff is zero
1:percolate has same concentration of nitrate as surface runoff
- real *8, dimension(:), allocatable **parm::surlag**
*Surface runoff lag time. This parameter is needed in subbasins where the time of concentration is greater than 1 day.
SURLAG is used to create a "storage" for surface runoff to allow the runoff to take longer than 1 day to reach the
subbasin outlet (days)*
- real *8, dimension(:), allocatable **parm::co_p**
- real *8, dimension(:), allocatable **parm::cmn**
rate factor for humus mineralization on active organic N
- real *8, dimension(:), allocatable **parm::phoskd**
Phosphorus soil partitioning coefficient. Ratio of soluble phosphorus in surface layer to soluble phosphorus in runoff.
- real *8, dimension(:), allocatable **parm::psp**
*Phosphorus availability index. The fraction of fertilizer P remaining in labile pool after initial rapid phase of P sorption
(none)*
- real *8, dimension(:), allocatable **parm::sdnco**
denitrification threshold: fraction of field capacity triggering denitrification
- real *8 **parm::r2adj_bsn**
basinwide retention parameter adjustment factor (greater than 1)
- real *8 **parm::pst_kg**
amount of pesticide applied to HRU (kg/ha)
- real *8 **parm::yield**
yield (dry weight) (kg)

- real *8 **parm::burn_frlb**
fraction of biomass and residue that burn(input in management file) range (0 - 1.0) (none)
- real *8 **parm::yieldgrn**
- real *8 **parm::yieldbms**
- real *8 **parm::yieldtbr**
- real *8 **parm::yieldn**
- real *8 **parm::yieldp**
- real *8 **parm::hi_bms**
- real *8 **parm::hi_rsd**
- real *8 **parm::yieldrsd**
- real *8, dimension(:), allocatable **parm::l_k1**
- real *8, dimension(:), allocatable **parm::l_k2**
- real *8, dimension(:), allocatable **parm::l_lambda**
- real *8, dimension(:), allocatable **parm::l_beta**
- real *8, dimension(:), allocatable **parm::l_gama**
- real *8, dimension(:), allocatable **parm::l_harea**
- real *8, dimension(:), allocatable **parm::l_vleng**
- real *8, dimension(:), allocatable **parm::l_vslope**
- real *8, dimension(:), allocatable **parm::l_ktc**
- real *8, dimension(:), allocatable **parm::biofilm_mumax**
- real *8, dimension(:), allocatable **parm::biofilm_kinv**
- real *8, dimension(:), allocatable **parm::biofilm_klw**
- real *8, dimension(:), allocatable **parm::biofilm_kla**
- real *8, dimension(:), allocatable **parm::biofilm_cdet**
- real *8, dimension(:), allocatable **parm::biofilm_bm**
- real *8, dimension(:, :), allocatable **parm::hru_rufr**
- real *8, dimension(:, :), allocatable **parm::daru_km**
- real *8, dimension(:, :), allocatable **parm::ru_k**
- real *8, dimension(:, :), allocatable **parm::ru_c**
- real *8, dimension(:, :), allocatable **parm::ru_eiq**
- real *8, dimension(:, :), allocatable **parm::ru_ovsl**
- real *8, dimension(:, :), allocatable **parm::ru_a**
- real *8, dimension(:, :), allocatable **parm::ru_ovs**
- real *8, dimension(:, :), allocatable **parm::ru_ktc**
- real *8, dimension(:), allocatable **parm::gwq_ru**
- real *8, dimension(:), allocatable **parm::qdayout**
- integer, dimension(:), allocatable **parm::ils2**
- integer, dimension(:), allocatable **parm::ils2flag**
- integer **parm::ipest**
pesticide identification number from pest.dat (none)
- integer **parm::iru**
- integer **parm::mru**
- integer **parm::irch**
- integer **parm::isub**
- integer **parm::mhyd_bsn**
- integer **parm::ils_nofig**
- integer **parm::mhru1**
- integer, dimension(:), allocatable **parm::mhyd1**
- integer, dimension(:), allocatable **parm::irtun**
- real *8 **parm::wshd_sepno3**
- real *8 **parm::wshd_sepnh3**
- real *8 **parm::wshd_seporgn**
- real *8 **parm::wshd_sepfon**
- real *8 **parm::wshd_seporgp**

- real *8 **parm::wshd_sepfop**
- real *8 **parm::wshd_sepsolp**
- real *8 **parm::wshd_sepbod**
- real *8 **parm::wshd_sepmm**
- integer, dimension(:), allocatable **parm::isep_hru**
- real *8 **parm::fixco**
nitrogen fixation coefficient
- real *8 **parm::nfixmx**
maximum daily n-fixation (kg/ha)
- real *8 **parm::res_stlr_co**
reservoir sediment settling coefficient
- real *8 **parm::rsd_covco**
residue cover factor for computing fraction of cover
- real *8 **parm::vcrit**
critical velocity
- real *8 **parm::wshd_snob**
average amount of water stored in snow at the beginning of the simulation for the entire watershed (mm H2O)
- real *8 **parm::wshd_sw**
water in soil at beginning of simulation, or average amount of water stored in soil for the entire watershed, or difference between mass balance calculated from watershed averages and actual value for water in soil at end of simulation (goal is to have wshd_sw = 0.) (mm H2O)
- real *8 **parm::wshd_pndfr**
fraction of watershed area which drains into ponds (none)
- real *8 **parm::wshd_pndsed**
total amount of suspended sediment in ponds in the watershed (metric tons), or mass balance discrepancy for pond sediment expressed as loading per unit hectare of drainage area (metric tons/ha)
- real *8 **parm::wshd_pndv**
total volume of water in ponds in the watershed (m^3), or mass balance discrepancy for pond water volume expressed as depth over drainage area (mm H2O)
- real *8 **parm::percop**
pesticide percolation coefficient (0-1)
0: concentration of pesticide in surface runoff is zero
1: percolate has same concentration of pesticide as surface runoff
- real *8 **parm::wshd_resfr**
fraction of watershed area that drains into reservoirs (none)
- real *8 **parm::wshd_pndha**
watershed area in hectares which drains into ponds (ha)
- real *8 **parm::wshd_resha**
watershed area in hectares which drains into reservoirs (ha)
- real *8 **parm::wshd_wetfr**
fraction of watershed area which drains into wetlands (none)
- real *8 **parm::wshd_fmimp**
average annual amount of mineral P applied in watershed (kg P/ha)
- real *8 **parm::wshd_fnh3**
average annual amount of NH3-N applied in watershed (kg N/ha)
- real *8 **parm::wshd_fno3**
average annual amount of NO3-N applied in watershed (kg N/ha)
- real *8 **parm::wshd_forgn**
average annual amount of organic N applied in watershed (kg N/ha)
- real *8 **parm::wshd_ftotn**
average annual amount of N (mineral & organic) applied in watershed (kg N/ha)

- real *8 **parm::wshd_forgp**
average annual amount of organic P applied in watershed (kg P/ha)
- real *8 **parm::wshd_ftotp**
average annual amount of P (mineral & organic) applied in watershed (kg P/ha)
- real *8 **parm::wshd_yldn**
amount of nitrogen removed from soil in watershed in the yield (kg N/ha)
- real *8 **parm::wshd_yldp**
amount of phosphorus removed from soil in watershed in the yield (kg P/ha)
- real *8 **parm::wshd_fixn**
- real *8 **parm::wshd_pup**
- real *8 **parm::wshd_wstrs**
- real *8 **parm::wshd_nstrs**
- real *8 **parm::wshd_pstrs**
- real *8 **parm::wshd_tstrs**
- real *8 **parm::wshd_astrs**
- real *8 **parm::ffcb**
initial soil water content expressed as a fraction of field capacity
- real *8 **parm::wshd_hmn**
- real *8 **parm::wshd_rwn**
- real *8 **parm::wshd_hmp**
- real *8 **parm::wshd_rmn**
- real *8 **parm::wshd_dnit**
- real *8 **parm::wdpq**
die-off factor for persistent bacteria in soil solution (1/day)
- real *8 **parm::wshd_rmp**
- real *8 **parm::wshd_voln**
- real *8 **parm::wshd_nitn**
- real *8 **parm::wshd_pas**
- real *8 **parm::wshd_pal**
- real *8 **parm::wof_p**
wash off fraction for persistent bacteria on foliage during a rainfall event
- real *8 **parm::wshd_plch**
- real *8 **parm::wshd_raino3**
- real *8 **parm::ressedc**
- real *8 **parm::basno3f**
- real *8 **parm::basorgnf**
- real *8 **parm::wshd_pinlet**
- real *8 **parm::wshd_ptile**
- real *8 **parm::sftmp**
Snowfall temperature (deg C)
- real *8 **parm::smfmn**
Minimum melt rate for snow during year (Dec. 21) where deg C refers to the air temperature. (mm/deg C/day)
- real *8 **parm::smfmx**
Maximum melt rate for snow during year (June 21) where deg C refers to the air temperature. SMFMX and SMFMN allow the rate of snow melt to vary through the year. These parameters are accounting for the impact of soil temperature on snow melt. (mm/deg C/day)
- real *8 **parm::smtmp**
Snow melt base temperature. Mean air temperature at which snow melt will occur. (deg C)
- real *8 **parm::wgpq**
growth factor for persistent bacteria in soil solution (1/day)
- real *8 **parm::basminpf**
- real *8 **parm::basorgpf**
- real *8 **parm::wdlpq**

- die-off factor for less persistent bacteria in soil solution (1/day)*
- real *8 **parm::wshd_ressed**
 - total amount of suspended sediment in reservoirs in the watershed (metric tons),
or mass balance discrepancy for reservoir sediment expressed as loading per unit hectare of drainage area (metric tons/ha)*
- real *8 **parm::wshd_resv**
 - total volume of water in all reservoirs in the watershed (m^3),
or mass balance discrepancy for reservoir water volume expressed as depth over drainage area (mm H₂O)*
- real *8 **parm::basminpi**
 - average amount of phosphorus initially in the mineral P pool in watershed soil (kg P/ha)*
- real *8 **parm::basno3i**
 - average amount of nitrogen initially in the nitrate pool in watershed soil (kg N/ha)*
- real *8 **parm::basorgni**
 - average amount of nitrogen initially in the organic N pool in watershed soil (kg N/ha)*
- real *8 **parm::wdps**
 - die-off factor for persistent bacteria adsorbed to soil particles (1/day)*
- real *8 **parm::wglpq**
 - growth factor for less persistent bacteria in soil solution (1/day)*
- real *8 **parm::basorgpi**
 - average amount of phosphorus initially in the organic P pool in watershed soil (kg P/ha)*
- real *8 **parm::peakr**
 - peak runoff rate for the day in HRU (m^3/s)*
- real *8 **parm::albdlay**
 - albedo of ground for the day in HRU, the fraction of the solar radiation reflected at the soil surface back into space (none)*
- real *8 **parm::pndsedin**
 - sediment inflow to the pond from HRU (metric tons)*
- real *8 **parm::sw_excess**
 - amount of water stored in soil layer on the current day that exceeds field capacity (gravity drained water) (mm H₂O)*
- real *8 **parm::timp**
 - Snow pack temperature lag factor (0-1)
1 = no lag (snow pack temp=current day air temp) as the lag factor goes to zero, the snow pack's temperature will be less influenced by the current day's air temperature.*
- real *8 **parm::wtabelo**
- real *8 **parm::tilep**
- real *8 **parm::wt_shall**
- real *8 **parm::sq_rto**
- real *8 **parm::qtile**
 - drainage tile flow in HRU soil layer for the day (mm H₂O)*
- real *8 **parm::inflpcp**
 - amount of precipitation that infiltrates into soil (enters soil) (mm H₂O)*
- real *8 **parm::crk**
 - percolation due to crack flow (mm H₂O)*
- real *8 **parm::fixn**
 - amount of nitrogen added to plant biomass via fixation on the day in HRU (kg N/ha)*
- real *8 **parm::latlyr**
 - amount of water in lateral flow in layer in HRU for the day (mm H₂O)*
- real *8 **parm::snofall**
 - amount of precipitation falling as freezing rain/snow on day in HRU (mm H₂O)*
- real *8 **parm::snomlt**
 - amount of water in snow melt for the day in HRU (mm H₂O)*
- real *8 **parm::tloss**

- amount of water removed from surface runoff via transmission losses on day in HRU (mm H2O)*

 - real *8 **parm::pndloss**
 - real *8 **parm::wetloss**
 - real *8 **parm::potloss**
 - real *8 **parm::lpndloss**
 - real *8 **parm::lwetloss**
 - real *8 **parm::bioday**
- biomass generated on current day in HRU (kg)*

 - real *8 **parm::cfertn**
- amount of nitrogen added to soil in continuous fertilizer operation on day (kg N/ha)*

 - real *8 **parm::cfertp**
- amount of phosphorus added to soil in continuous fertilizer operation on day (kg P/ha)*

 - real *8 **parm::fertn**
- total amount of nitrogen applied to soil in HRU on day (kg N/ha)*

 - real *8 **parm::sepday**
- micropore percolation from bottom of the soil layer on day in HRU (mm H2O)*

 - real *8 **parm::sol_rd**
- current rooting depth (mm)*

 - real *8 **parm::sedrch**
 - real *8 **parm::sepcrk**
 - real *8 **parm::sepcrktot**
 - real *8 **parm::fertno3**
 - real *8 **parm::fertnh3**
 - real *8 **parm::fertorgn**
 - real *8 **parm::fertsolp**
 - real *8 **parm::fertorgp**
 - real *8 **parm::wgps**
- growth factor for persistent bacteria adsorbed to soil particles (1/day)*

 - real *8 **parm::qdfr**
- fraction of water yield that is surface runoff (none)*

 - real *8 **parm::fertp**
- total amount of phosphorus applied to soil in HRU on day (kg P/ha)*

 - real *8 **parm::grazn**
- amount of nitrogen added to soil in grazing on the day in HRU (kg N/ha)*

 - real *8 **parm::grazp**
- amount of phosphorus added to soil in grazing on the day in HRU (kg P/ha)*

 - real *8 **parm::soxy**
- saturation dissolved oxygen concentration (mg/L)*

 - real *8 **parm::sdti**
 - real *8 **parm::rtwtr**
 - real *8 **parm::ressa**
 - real *8 **parm::wdlps**
- die-off factor for less persistent bacteria absorbed to soil particles (1/day)*

 - real *8 **parm::wglps**
- growth factor for less persistent bacteria adsorbed to soil particles (1/day)*

 - real *8 **parm::da_km**
- area of the watershed in square kilometers (km²)*

 - real *8 **parm::rttime**
 - real *8 **parm::rchdep**
 - real *8 **parm::rtevp**
 - real *8 **parm::rttlc**
 - real *8 **parm::resflwi**

- real *8 **parm::wdprch**
die-off factor for persistent bacteria in streams (1/day)
- real *8 **parm::resflwo**
- real *8 **parm::respcp**
- real *8 **parm::resev**
- real *8 **parm::ressep**
- real *8 **parm::ressedi**
- real *8 **parm::ressedo**
- real *8 **parm::dtot**
- real *8 **parm::pperco_bsn**
phosphorus percolation coefficient. Ratio of soluble phosphorus in surface to soluble phosphorus in percolate
- real *8 **parm::nperco_bsn**
basin nitrate percolation coefficient (0-1)
0:concentration of nitrate in surface runoff is zero
1:percolate has same concentration of nitrate as surface runoff
- real *8 **parm::rsdco**
residue decomposition coefficient. The fraction of residue which will decompose in a day assuming optimal moisture, temperature, C:N ratio, and C:P ratio
- real *8 **parm::voltot**
total volume of cracks expressed as depth per unit area (mm)
- real *8 **parm::phoskd_bsn**
- real *8 **parm::msk_x**
weighting factor controlling relative importance of inflow rate and outflow rate in determining storage on reach
- real *8 **parm::volcrmin**
minimum crack volume allowed in any soil layer (mm)
- real *8 **parm::bactkdq**
bacteria soil partitioning coefficient. Ratio of solution bacteria in surface layer to solution bacteria in runoff soluble and sorbed phase in surface runoff.
- real *8 **parm::wdpf**
die-off factor for persistent bacteria on foliage (1/day)
- real *8 **parm::canev**
amount of water evaporated from canopy storage (mm H2O)
- real *8 **parm::precipday**
precipitation, or effective precipitation reaching soil surface, for the current day in HRU (mm H2O)
- real *8 **parm::uno3d**
plant nitrogen deficiency for day in HRU (kg N/ha)
- real *8 **parm::usle**
daily soil loss predicted with USLE equation (metric tons/ha)
- real *8 **parm::rcn**
- real *8 **parm::surlag_bsn**
- real *8 **parm::thbact**
temperature adjustment factor for bacteria die-off/growth
- real *8 **parm::wlpq20**
overall rate change for less persistent bacteria in soil solution (1/day)
- real *8 **parm::wlps20**
overall rate change for less persistent bacteria adsorbed to soil particles (1/day)
- real *8 **parm::wpq20**
overall rate change for persistent bacteria in soil solution (1/day)
- real *8 **parm::wps20**
overall rate change for persistent bacteria adsorbed to soil particles (1/day)
- real *8 **parm::bactrop**
persistent bacteria transported to main channel with surface runoff (# colonies/ha)

- real *8 `parm::bactsedp`
persistent bacteria transported with sediment in surface runoff (# colonies/ha)
- real *8 `parm::wgpf`
growth factor for persistent bacteria on foliage (1/day)
- real *8 `parm::bactlchlp`
less persistent bacteria removed from soil surface layer by percolation (# colonies/ha)
- real *8 `parm::bactlchp`
persistent bacteria removed from soil surface layer by percolation (# colonies/ha)
- real *8 `parm::enratio`
enrichment ratio calculated for day in HRU (none)
- real *8 `parm::pndpcp`
precipitation on pond during day ($m^3 H_2O$)
- real *8 `parm::wetpcp`
- real *8 `parm::wetsep`
seepage from wetland bottom for day ($m^3 H_2O$)
- real *8 `parm::pndev`
evaporation from pond on day ($m^3 H_2O$)
- real *8 `parm::pndflwi`
volume of water flowing into pond on day ($m^3 H_2O$)
- real *8 `parm::pndsedo`
sediment leaving pond during day (metric tons)
- real *8 `parm::pndsep`
seepage from pond on day ($m^3 H_2O$)
- real *8 `parm::wetev`
evaporation from wetland for day ($m^3 H_2O$)
- real *8 `parm::wetflwi`
volume of water flowing in wetland on day ($m^3 H_2O$)
- real *8 `parm::wetsedo`
sediment loading from wetland for day (metric tons)
- real *8 `parm::da_ha`
drainage area of watershed in hectares (ha)
- real *8 `parm::pndflwo`
volume of water flowing out of pond on day ($m^3 H_2O$)
- real *8 `parm::vpd`
vapor pressure deficit (kPa)
- real *8 `parm::wetflwo`
volume of water flowing out wetland on day ($m^3 H_2O$)
- real *8 `parm::wetsedi`
sediment loading to wetland for day (metric tons)
- real *8 `parm::evlai`
leaf area index at which no evaporation occurs. This variable is used in ponded HRUs where evaporation from the water surface is restricted by the plant canopy cover. Evaporation from the water surface equals potential ET when LAI = 0 and decreased linearly to 0 when LAI = EVLAI
- real *8 `parm::evrch`
Reach evaporation adjustment factor. Evaporation from the reach is multiplied by EVRCH. This variable was created to limit the evaporation predicted in arid regions.
- real *8 `parm::wdlpf`
die-off factor for less persistent bacteria on foliage (1/day)
- real *8 `parm::ep_day`
actual amount of transpiration that occurs on day in HRU (mm H_2O)
- real *8 `parm::pet_day`

- potential evapotranspiration on current day in HRU (mm H2O)*

 - real *8 **parm::bactrolp**

less persistent bacteria transported to main channel with surface runoff (# colonies/ha)
 - real *8 **parm::bactsedlp**

less persistent bacteria transported with sediment in surface runoff (# colonies/ha)
 - real *8 **parm::adj_pkr**

peak rate adjustment factor in the subbasin. Used in the MUSLE equation to account for impact of peak flow on erosion (none)
 - real *8 **parm::n_updis**

nitrogen uptake distribution parameter. This parameter controls the amount of nitrogen removed from the different soil layer layers by the plant. In particular, this parameter allows the amount of nitrogen removed from the surface layer via plant uptake to be controlled. While the relationship between UBN and N removed from the surface layer is affected by the depth of the soil profile, in general, as UBN increases the amount of N removed from the surface layer relative to the amount removed from the entire profile increases
 - real *8 **parm::nactfr**

nitrogen active pool fraction. The fraction of organic nitrogen in the active pool (none)
 - real *8 **parm::p_updis**

phosphorus uptake distribution parameter This parameter controls the amount of phosphorus removed from the different soil layers by the plant. In particular, this parameter allows the amount of phosphorus removed from the surface layer via plant uptake to be controlled. While the relationship between UBP and P uptake from the surface layer is affected by the depth of the soil profile, in general, as UBP increases the amount of P removed from the surface layer relative to the amount removed from the entire profile increases
 - real *8 **parm::snoev**

amount of water in snow lost through sublimation on current day in HRU (mm H2O)
 - real *8 **parm::sno3up**
 - real *8 **parm::reactw**
 - real *8 **parm::es_day**

actual amount of evaporation (soil et) that occurs on day in HRU (mm H2O)
 - real *8 **parm::sdiégropq**
 - real *8 **parm::sdiégrolpq**
 - real *8 **parm::sdiégrops**
 - real *8 **parm::sdiégrolps**
 - real *8 **parm::wof_lp**

wash off fraction for less persistent bacteria on foliage during a rainfall event
 - real *8 **parm::ep_max**

maximum amount of transpiration (plant et) that can occur on day in HRU (mm H2O)
 - real *8 **parm::sbactrop**
 - real *8 **parm::sbactrolp**
 - real *8 **parm::sbactsedp**
 - real *8 **parm::sbactsedlp**
 - real *8 **parm::sbactlchp**
 - real *8 **parm::sbactlchlp**
 - real *8 **parm::psp_bsn**
 - real *8 **parm::rchwtr**
 - real *8 **parm::resuspst**
 - real *8 **parm::setlpst**
 - real *8 **parm::bsprev**

surface runoff lagged from prior day of simulation (mm H2O)
 - real *8 **parm::bssprev**

lateral flow lagged from prior day of simulation (mm H2O)
 - real *8 **parm::spadyo**
 - real *8 **parm::spadyev**
 - real *8 **parm::spadyisp**
 - real *8 **parm::spadyrfv**

- real *8 **parm::spadyosp**
- real *8 **parm::qday**
amount of surface runoff loading to main channel from HRU on current day (mm H2O)
- real *8 **parm::al5**
fraction of total rainfall that occurs during 0.5h of highest intensity rain (none)
- real *8 **parm::no3pcp**
nitrate added to the soil in rainfall (kg N/ha)
- real *8 **parm::pndsedc**
net change in sediment in pond during day (metric tons)
- real *8 **parm::usle_ei**
USLE rainfall erosion index on day for HRU (100(ft·tn in)/(acre·hr))
- real *8 **parm::rcharea**
- real *8 **parm::volatpst**
- real *8 **parm::ubw**
water uptake distribution parameter. This parameter controls the amount of water removed from the different soil layers by the plant. In particular, this parameter allows the amount of water removed from the surface layer via plant uptake to be controlled. While the relationship between UBW and H2O removed from the surface layer is affected by the depth of the soil profile, in general, as UBW increases the amount of water removed from the surface layer relative to the amount removed from the entire profile increases
- real *8 **parm::uobn**
nitrogen uptake normalization parameter. This variable normalizes the nitrogen uptake so that the model can easily verify that upake from the different soil layers sums to 1.0
- real *8 **parm::uobp**
phosphorus uptake normalization parameter. This variable normalizes the phosphorus uptake so that the model can easily verify that uptake from the different soil layers sums to 1.0
- real *8 **parm::uobw**
water uptake normalization parameter. This variable normalizes the water uptake so that the model can easily verify that uptake from the different soil layers sums to 1.0
- real *8 **parm::wglpf**
growth factor for less persistent bacteria on foliage (1/day)
- real *8 **parm::wetsedc**
net change in sediment in wetland during day (metric tons)
- real *8 **parm::respesti**
- real *8 **parm::rcor**
correction coefficient for generated rainfall to ensure that the annual means for generated and observed values are comparable (needed only if IDIST=1)
- real *8 **parm::rexp**
value of exponent for mixed exponential rainfall distribution (needed only if IDIST=1)
- real *8 **parm::snocov1**
1st shape parameter for snow cover equation. This parameter is determined by solving the equation for 50% snow cover
- real *8 **parm::snocov2**
2nd shape parameter for snow cover equation. This parameter is determined by solving the equation for 95% snow cover
- real *8 **parm::snocovmx**
Minimum snow water content that corresponds to 100% snow cover. If the snow water content is less than SNOC← OVMX, then a certain percentage of the ground will be bare (mm H2O)
- real *8 **parm::lyrtile**
drainage tile flow in soil layer for day (mm H2O)
- real *8 **parm::lyrtilex**
- real *8 **parm::sno50cov**
Fraction of SNOCOVMX that corresponds to 50% snow cover. SWAT assumes a nonlinear relationship between snow water and snow cover.

- real *8 `parm::ai0`
ratio of chlorophyll-a to algal biomass (ug chla/mg alg)
- real *8 `parm::ai1`
fraction of algal biomass that is nitrogen (mg N/mg alg)
- real *8 `parm::ai2`
fraction of algal biomass that is phosphorus (mg P/mg alg)
- real *8 `parm::ai3`
the rate of oxygen production per unit of algal photosynthesis (mg O2/mg alg)
- real *8 `parm::ai4`
the rate of oxygen uptake per unit of algae respiration (mg O2/mg alg)
- real *8 `parm::ai5`
the rate of oxygen uptake per unit of NH3 nitrogen oxidation (mg O2/mg N)
- real *8 `parm::ai6`
the rate of oxygen uptake per unit of NO2 nitrogen oxidation (mg O2/mg N)
- real *8 `parm::rhoq`
algal respiration rate (1/day or 1/hr)
- real *8 `parm::tfact`
fraction of solar radiation computed in the temperature heat balance that is photosynthetically active
- real *8 `parm::k_l`
*half-saturation coefficient for light (MJ/(m2*hr))*
- real *8 `parm::k_n`
michaelis-menton half-saturation constant for nitrogen (mg N/L)
- real *8 `parm::k_p`
michaelis-menton half saturation constant for phosphorus (mg P/L)
- real *8 `parm::lambda0`
non-algal portion of the light extinction coefficient (1/m)
- real *8 `parm::lambda1`
*linear algal self-shading coefficient (1/(m*ug chla/L))*
- real *8 `parm::lambda2`
*nonlinear algal self-shading coefficient ((1/m)(ug chla/L)**(-2/3))*
- real *8 `parm::mumax`
maximum specific algal growth rate (1/day or 1/hr)
- real *8 `parm::p_n`
algal preference factor for ammonia
- real *8 `parm::rnum1`
variable to hold value for rnum1s(:) (none)
- real *8 `parm::etday`
actual evapotranspiration occurring on day in HRU (mm H2O)
- real *8 `parm::auton`
amount of nitrogen applied in auto-fert application (kg N/ha)
- real *8 `parm::autop`
amount of phosphorus applied in auto-fert application (kg P/ha)
- real *8 `parm::hmntl`
amount of nitrogen moving from active organic to nitrate pool in soil profile on current day in HRU (kg N/ha)
- real *8 `parm::hmptl`
amount of phosphorus moving from active organic to nitrate pool in soil profile on current day in HRU (kg P/ha)
- real *8 `parm::rmn2tl`
amount of nitrogen moving from the fresh organic (residue) to the nitrate(80%) and active organic(20%) pools in soil profile on current day in HRU (kg N/ha)
- real *8 `parm::rwntl`
amount of nitrogen moving from active organic to stable organic pool in soil profile on current day in HRU (kg N/ha)

- real *8 **parm::gwseep**
amount of water recharging deep aquifer on current day (mm H2O)
- real *8 **parm::revapday**
amount of water moving from the shallow aquifer into the soil profile or being taken up by plant roots in the shallow aquifer (mm H2O)
- real *8 **parm::rmp1tl**
amount of phosphorus moving from the labile mineral pool to the active mineral pool in the soil profile on the current day in the HRU (kg P/ha)
- real *8 **parm::rmp1tl**
amount of phosphorus moving from the fresh organic (residue) to the labile(80%) and organic(20%) pools in soil profile on current day in HRU (kg P/ha)
- real *8 **parm::roct1**
amount of phosphorus moving from the active mineral pool to the stable mineral pool in the soil profile on the current day in the HRU (kg P/ha)
- real *8 **parm::wdntl**
amount of nitrogen lost from nitrate pool by denitrification in soil profile on current day in HRU (kg N/ha)
- real *8 **parm::cmn_bsn**
- real *8 **parm::reswtr**
- real *8 **parm::wdlprch**
die-off factor for less persistent bacteria in streams (1/day)
- real *8 **parm::wdpres**
die-off factor for persistent bacteria in reservoirs (1/day)
- real *8 **parm::petmeas**
potential ET value read in for day (mm H2O)
- real *8 **parm::bury**
- real *8 **parm::difus**
- real *8 **parm::reactb**
- real *8 **parm::solpesto**
- real *8 **parm::wdlpres**
die-off factor for less persistent bacteria in reservoirs (1/day)
- real *8 **parm::sorpesto**
- real *8 **parm::spcon_bsn**
- real *8 **parm::spexp_bsn**
- real *8 **parm::solpesti**
- real *8 **parm::sorpesti**
- real *8 **parm::msk_co1**
calibration coefficient to control impact of the storage time constant for the reach at bankfull depth ($\phi(10, \cdot)$) upon the storage time constant for the reach used in the Muskingum flow method
- real *8 **parm::msk_co2**
calibration coefficient to control impact of the storage time constant for the reach at 0.1 bankfull depth ($\phi(13, \cdot)$) upon the storage time constant for the reach used in the Muskingum flow method
- real *8 **parm::deepstp**
depth of water in deep aquifer in HRU (mm H2O)
- real *8 **parm::shallstp**
depth of water in shallow aquifer in HRU on previous day (mm H2O)
- real *8 **parm::snoprev**
amount of water stored as snow on previous day (mm H2O)
- real *8 **parm::swprev**
amount of water stored in soil profile in the HRU on the previous day (mm H2O)
- real *8 **parm::ressolpo**
- real *8 **parm::resorgno**
- real *8 **parm::resorgpo**
- real *8 **parm::resno3o**

- real *8 **parm::reschlao**
- real *8 **parm::resno2o**
- real *8 **parm::potevmm**
volume of water evaporated from pothole expressed as depth over HRU (mm H2O)
- real *8 **parm::potflwo**
volume of water released to main channel from pothole expressed as depth over HRU (mm H2O)
- real *8 **parm::potpcpmm**
precipitation falling on pothole water body expressed as depth over HRU (mm H2O)
- real *8 **parm::potsepmmm**
seepage from pothole expressed as depth over HRU (mm H2O)
- real *8 **parm::resnh3o**
- real *8 **parm::qdbank**
- real *8 **parm::bactminlp**
Threshold detection level for less persistent bacteria. When bacteria levels drop to this amount the model considers bacteria in the soil to be insignificant and sets the levels to zero (cfu/m²)
- real *8 **parm::bactminp**
Threshold detection level for persistent bacteria. When bacteria levels drop to this amount the model considers bacteria in the soil to be insignificant and sets the levels to zero (cfu/m²)
- real *8 **parm::trnsrch**
fraction of transmission losses from main channel that enter deep aquifer
- real *8 **parm::wp20p_plt**
overall rate change for persistent bacteria on foliage (1/day)
- real *8 **parm::potsedo**
sediment released to main channel from HRU (metric tons/ha)
- real *8 **parm::pest_sol**
- real *8 **parm::bact_swf**
fraction of manure containing active colony forming units (cfu)
- real *8 **parm::bactmx**
bacteria percolation coefficient. Ratio of solution bacteria in surface layer to solution bacteria in percolate
- real *8 **parm::cncoef**
plant ET curve number coefficient
- real *8 **parm::wp20lp_plt**
overall rate change for less persistent bacteria on foliage (1/day)
- real *8 **parm::cdn_bsn**
- real *8 **parm::sdnco_bsn**
- real *8 **parm::bactmin**
- real *8 **parm::cn_froz**
drainage coefficient (mm day⁻¹)
- real *8 **parm::dorm_hr**
time threshold used to define dormant (hours)
- real *8 **parm::smxco**
adjustment factor for max curve number s factor (0-1)
- real *8 **parm::tb_adj**
adjustment factor for subdaily unit hydrograph basetime
- real *8 **parm::chla_subco**
regional adjustment on sub chla_a loading (fraction)
- real *8 **parm::depimp_bsn**
depth to impervious layer. Used to model perched water tables in all HRUs in watershed (mm)
- real *8 **parm::ddrain_bsn**
depth to the sub-surface drain (mm)
- real *8 **parm::tdrain_bsn**
time to drain soil to field capacity (hours)

- real *8 **parm::gdrain_bsn**
- real *8 **parm::rch_san**
- real *8 **parm::rch_sil**
- real *8 **parm::rch_cla**
- real *8 **parm::rch_sag**
- real *8 **parm::rch_lag**
- real *8 **parm::rch_gra**
- real *8 **parm::hlife_ngw_bsn**
Half-life of nitrogen in groundwater? (days)
- real *8 **parm::ch_opco_bsn**
- real *8 **parm::ch_onco_bsn**
- real *8 **parm::decr_min**
Minimum daily residue decay.
- real *8 **parm::rcn_sub_bsn**
Concentration of nitrogen in the rainfall (mg/kg)
- real *8 **parm::bc1_bsn**
- real *8 **parm::bc2_bsn**
- real *8 **parm::bc3_bsn**
- real *8 **parm::bc4_bsn**
- real *8 **parm::anion_excl_bsn**
- real *8, dimension(:), allocatable **parm::wat_tbl**
- real *8, dimension(:), allocatable **parm::sol_swpwt**
- real *8, dimension(:,:), allocatable **parm::vwt**
- real *8 **parm::re_bsn**
Effective radius of drains (range 3.0 - 40.0) (mm)
- real *8 **parm::sdrain_bsn**
Distance bewtween two drain or tile tubes (range 7600.0 - 30000.0) (mm)
- real *8 **parm::sstmaxd_bsn**
- real *8 **parm::drain_co_bsn**
Drainage coefficient (range 10.0 - 51.0) (mm-day-1)
- real *8 **parm::latksatf_bsn**
Multiplication factor to determine lateral ksat from SWAT ksat input value for HRU (range 0.01 - 4.0)
- real *8 **parm::pc_bsn**
Pump capacity (def val = 1.042 mm h-1 or 25 mm day-1) (mm h-1)
- integer **parm::idlast**
number of days simulated in month (none)
- integer **parm::i_subhw**
- integer **parm::imgt**
- integer **parm::iwtr**
- integer **parm::ifrttyp**
- integer **parm::mo_atmo**
- integer **parm::mo_atmo1**
- integer **parm::ifirstatmo**
- integer **parm::iyr_atmo**
- integer **parm::iyr_atmo1**
- integer **parm::matmo**
- integer **parm::mch**
maximum number of channels
- integer **parm::mcr**
maximum number of crops grown per year
- integer **parm::mcrdb**
maximum number of crops/landcover in database file (crop.dat)
- integer **parm::mfcst**

- maximum number of forecast stations*
- integer **parm::mfdb**
 - maximum number of fertilizers in fert.dat*
- integer **parm::mhru**
 - maximum number of HRUs in watershed*
- integer **parm::mhyd**
 - maximum number of hydrograph nodes*
- integer **parm::mpdb**
 - maximum number of pesticides in pest.dat*
- integer **parm::mrg**
 - maximum number of rainfall/temp gages (none)*
- integer **parm::mcut**
 - maximum number of cuttings per year*
- integer **parm::mgr**
 - maximum number of grazings per year*
- integer **parm::mnr**
 - maximum number of years of rotation*
- integer **parm::myr**
 - maximum number of years of simulation*
- integer **parm::isubwq**
 - subbasin water quality code*
 - 0 do not calculate algae/CBOD 1 calculate algae/CBOD drainmod tile equations*
- integer **parm::fcst**
- integer **parm::isproj**
 - special project code (none):*
 - 1 test rewind (run simulation twice)*
- integer **parm::nbyr**
 - number of calendar years simulated (none)*
- integer **parm::irte**
 - water routing method (none):*
 - 0 variable storage method*
 - 1 Muskingum method*
- integer **parm::nrch**
 - number of reaches in watershed (none)*
- integer **parm::nres**
 - total number of reservoirs in watershed (none)*
- integer **parm::nhru**
 - number of last HRU in previous subbasin or*
 - number of HRUs in watershed (none)*
- integer **parm::i_mo**
 - current month being simulated or month of next day of simulation (none)*
- integer **parm::immo**
 - current cumulative month of simulation (none)*
- integer **parm::mo**
- integer **parm::wndsim**
 - wind speed input code (noen)*
 - 1 measured data read for each subbasin*
 - 2 data simulated for each subbasin*
- integer **parm::ihru**
 - HRU number (none)*
- integer **parm::icode**

- variable to hold value for icodes(:) (none)*
- integer **parm::ihout**
 - variable to hold value for ihouts(:) (none)*
- integer **parm::inum1**
 - variable to hold value for inum1s(:) (subbasin number) (none)*
- integer **parm::inum2**
 - variable to hold value for inum2s(:) (none)*
- integer **parm::inum3**
 - variable to hold value for inum3s(:) (none)*
- integer **parm::inum4**
 - variable to hold value for inum4s(:) (none)*
- integer **parm::icfac**
 - icfac = 0 for C-factor calculation using Cmin (as described in manual)*
 - = 1 for new C-factor calculation from RUSLE (no minimum needed)*
- integer **parm::inum5**
- integer **parm::inum6**
- integer **parm::inum7**
- integer **parm::inum8**
- integer **parm::mrech**
 - maximum number of rechour files*
- integer **parm::nrgage**
 - number of raingage files (none)*
- integer **parm::nrgfil**
 - number of rain gages per file (none)*
- integer **parm::nrtot**
 - total number of rain gages (none)*
- integer **parm::ntgage**
 - number of temperature gage files (none)*
- integer **parm::ntgfil**
 - number of temperature gages per file (none)*
- integer **parm::nttot**
 - total number of temperature gages (none)*
- integer **parm::tmpsim**
 - temperature input code (none)*
 - 1 measured data read for each subbasin*
 - 2 data simulated for each subbasin*
- integer **parm::icrk**
 - crack flow code*
 - 1: simulate crack flow in watershed*
- integer **parm::irtpest**
 - number of pesticide to be routed through the watershed. Redefined to the sequence number of pesticide in NPNO(:) which is to be routed through the watershed (none)*
- integer **parm::igropt**
 - Qual2E option for calculating the local specific growth rate of algae*
 - 1: multiplicative.*
- integer **parm::lao**
 - Qual2E light averaging option. Qual2E defines four light averaging options. The only option currently available in SWAT is #2.*
- integer **parm::npmx**
 - number of different pesticides used in the simulation (none)*
- integer **parm::curyr**
 - current year in simulation (sequence) (none)*

- integer **parm::itdrn**
tile drainage equations flag/code
1 simulate tile flow using subroutine drains(wt_shall)
0 simulate tile flow using subroutine origtile(wt_shall,d)
- integer **parm::iwtdn**
water table depth algorithms flag/code
1 simulate wt_shall using subroutine new water table depth routine
0 simulate wt_shall using subroutine original water table depth routine
- integer **parm::ismax**
maximum depressional storage selection flag/code
0 = static depressional storage
1 = dynamic storage based on tillage and cumulative rainfall
- integer **parm::iroutunit**
not being implemented in this version drainmod tile equations
- integer **parm::ires_nut**
- integer **parm::iclb**
auto-calibration flag
- integer **parm::mrecc**
maximum number of reccnst files
- integer **parm::mrecd**
maximum number of recday files
- integer **parm::mrecm**
maximum number of recmon files
- integer **parm::mtil**
max number of tillage types in till.dat
- integer **parm::mudb**
maximum number of urban land types in urban.dat
- integer **parm::idist**
rainfall distribution code
0 for skewed normal dist
1 for mixed exponential distribution
- integer **parm::mrecy**
maximum number of recyear files
- integer **parm::nyskip**
number of years to skip output summarization and printing (none)
- integer **parm::slrsim**
solar radiation input code (none)
1 measured data read for each subbasin
2 data simulated for each subbasin
- integer **parm::ideg**
channel degradation code
0: do not compute channel degradation
1: compute channel degradation (downcutting and widening)
- integer **parm::ievent**
rainfall/runoff code (none)
0 daily rainfall/curve number technique 1 sub-daily rainfall/Green&Ampt/hourly routing 3 sub-daily rainfall/↔
Green&Ampt/hourly routing
- integer **parm::ipet**
code for potential ET method (none)
0 Priestley-Taylor method
1 Penman/Monteith method
2 Hargreaves method
3 read in daily potential ET data
- integer **parm::iopera**

- integer `parm::idaf`
beginning day of simulation (julian date)
- integer `parm::idal`
ending day of simulation (julian date)
- integer `parm::rhsim`
relative humidity input code (none)
1 measured data read for each subbasin
2 data simulated for each subbasin
- integer `parm::leapyr`
leap year flag (none)
0 leap year
1 regular year
- integer `parm::id1`
first day of simulation in current year (julian date)
- integer `parm::mo_chk`
current month of simulation (none)
- integer `parm::nhtot`
total number of relative humidity records in file
- integer `parm::nstot`
total number of solar radiation records in file (none)
- integer `parm::nwtot`
total number of wind speed records in file
- integer `parm::ifirsts`
solar radiation data search code (none)
0 first day of solar radiation data located in file
1 first day of solar radiation data not located in file
- integer `parm::ifirsth`
relative humidity data search code (none)
0 first day of relative humidity data located in file
1 first day of relative humidity data not located in file
- integer `parm::ifirstw`
wind speed data search code (none)
0 first day of wind speed data located in file
1 first day of wind speed data not located in file
- integer `parm::icst`
- integer `parm::ilog`
streamflow print code (none)
0 print streamflow in reach
1 print Log10 streamflow in reach
- integer `parm::itotr`
number of output variables printed (output.rch)
- integer `parm::iyr`
current year of simulation (year)
- integer `parm::iwq`
stream water quality code
0 do not model stream water quality
1 model stream water quality (QUAL2E & pesticide transformations)
- integer `parm::iskip`
flag for calculations performed only for the first year of simulation (none)
- integer `parm::ifirstpet`
potential ET data search code (none)
0 first day of potential ET data located in file
1 first day of potential ET data not located in file
- integer `parm::iprp`

- print code for output.pst file*
- 0 do not print pesticide output*
- 1 print pesticide output*
- integer **parm::itotb**
 - number of output variables printed (output.sub)*
- integer **parm::itots**
 - number of output variables printed (output.hru)*
- integer **parm::ith**
 - number of HRUs printed (output.hru/output.wtr)*
- integer **parm::pcpsim**
 - rainfall input code (none)*
 - 1 measured data read for each subbasin*
 - 2 data simulated for each subbasin*
- integer **parm::nd_30**
- integer **parm::iops**
- integer **parm::iphr**
- integer **parm::isto**
- integer **parm::isol**
- integer **parm::fcstcycles**
 - number of times forecast period is simulated (using different weather generator seeds each time)*
- integer **parm::fcstday**
 - beginning date of forecast period (julian date)*
- integer **parm::fcstyr**
 - beginning year of forecast period*
- integer **parm::iscen**
 - scenarios counter*
- integer **parm::subtot**
 - number of subbasins in watershed (none)*
- integer **parm::ogen**
- integer **parm::mapp**
 - maximum number of applications*
- integer **parm::mlyr**
 - maximum number of soil layers*
- integer **parm::mpst**
 - max number of pesticides used in wshed*
- integer **parm::mres**
 - maximum number of reservoirs*
- integer **parm::msub**
 - maximum number of subbasins*
- integer **parm::igen**
 - random number generator seed code (none):*
 - 0: use default numbers*
 - 1: generate new numbers in every simulation*
- integer **parm::iprint**
 - print code (none): 0=monthly, 1=daily, 2=annually*
- integer **parm::iida**
 - day being simulated (current julian date) (julian date)*
- integer **parm::icn**
 - CN method flag (for testing alternative method):*
 - 0 use traditional SWAT method which bases CN on soil moisture*
 - 1 use alternative method which bases CN on plant ET*
 - 2 use traditional SWAT method which bases CN on soil moisture but retention is adjusted for mildly-sloped tiled-drained watersheds.*

- integer **parm::ised_det**
max half-hour rainfall fraction calc option:
0 generate max half-hour rainfall fraction from triangular distribution
1 use monthly mean max half-hour rainfall fraction
- integer **parm::fcstcnt**
- integer **parm::mtran**
- integer **parm::idtill**
- integer, dimension(100) **parm::ida_lup**
- integer, dimension(100) **parm::iyr_lup**
- integer **parm::no_lup**
- integer **parm::no_up**
- integer **parm::nostep**
- character(len=8) **parm::date**
date simulation is performed where leftmost eight characters are set to a value of yyyyymmdd, where yyyy is the year, mm is the month and dd is the day
- character(len=10) **parm::time**
time simulation is performed where leftmost ten characters are set to a value of hhmmss.sss, where hh is the hour, mm is the minutes and ss.sss is the seconds and milliseconds
- character(len=5) **parm::zone**
time difference with respect to Coordinated Universal Time (ie Greenwich Mean Time)
- character(len=13) **parm::calfile**
name of file containing calibration parameters
- character(len=13) **parm::rhfile**
relative humidity file name (.hmd)
- character(len=13) **parm::slrfile**
solar radiation file name (.slr)
- character(len=13) **parm::wndfile**
wind speed file name (.wnd)
- character(len=13) **parm::petfile**
potential ET file name (.pet)
- character(len=13) **parm::atmofile**
- character(len=13) **parm::lucfile**
- character(len=13) **parm::septdb**
name of septic tank database file (septwq1.dat)
- character(len=13) **parm::dpd_file**
- character(len=13) **parm::wpd_file**
- character(len=13) **parm::rib_file**
- character(len=13) **parm::sfb_file**
- character(len=13) **parm::lid_file**
- integer, dimension(9) **parm::idg**
array location of random number seed used for a given process
- integer, dimension(:), allocatable **parm::iffirstr**
- integer, dimension(:), allocatable **parm::iffirsthr**
- integer, dimension(8) **parm::values**
values(1): year simulation is performed
values(2): month simulation is performed
values(3): day in month simulation is performed
values(4): time difference with respect to Coordinated Universal Time (ie Greenwich Mean Time)
values(5): hour simulation is performed
values(6): minute simulation is performed
values(7): second simulation is performed
values(8): millisecond simulation is performed
- integer, dimension(13) **parm::ndays**

julian date for last day of preceding month (where the array location is the number of the month). The dates are for leap years (julian date)

- integer **parm::mapex**
- real *8, dimension(:), allocatable **parm::flodaya**
- real *8, dimension(:), allocatable **parm::seddaya**
- real *8, dimension(:), allocatable **parm::orgndaya**
- real *8, dimension(:), allocatable **parm::orgpdaya**
- real *8, dimension(:), allocatable **parm::no3daya**
- real *8, dimension(:), allocatable **parm::minpdaya**
- real *8, dimension(:), allocatable **parm::hi_targ**
harvest index target of cover defined at planting ((kg/ha)/(kg/ha))
- real *8, dimension(:), allocatable **parm::bio_targ**
biomass target (kg/ha)
- real *8, dimension(:), allocatable **parm::tnyld**
modifier for autofertilization target nitrogen content for plant (kg N/kg yield)
- integer, dimension(:), allocatable **parm::idapa**
- integer, dimension(:), allocatable **parm::iypa**
- integer, dimension(:), allocatable **parm::ifirsta**
- integer, dimension(100) **parm::mo_transb**
- integer, dimension(100) **parm::mo_transe**
- integer, dimension(100) **parm::ih_tran**
- integer **parm::msdb**
maximum number of sept wq data database (none)
- integer **parm::iseptic**
- real *8, dimension(:), allocatable **parm::sptqs**
flow rate of the septic tank effluent per capita (m3/d)
- real *8, dimension(:), allocatable **parm::percp**
- real *8, dimension(:), allocatable **parm::sptbodconcs**
Biological Oxygen Demand of the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **parm::spttssconcs**
concentration of total suspended solid in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **parm::sptnconcs**
concentration of total nitrogen in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **parm::sptnh4concs**
concentration of total phosphorus of the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **parm::sptno3concs**
concentration of nitrate in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **parm::sptno2concs**
concentration of nitrite in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **parm::sptorgnconcs**
concentration of organic nitrogen in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **parm::spttpconcs**
concentration of total phosphorus in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **parm::sptminps**
concentration of mineral phosphorus in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **parm::sptorgps**
concentration of organic phosphorus in the septic tank effluent (mg/l)
- real *8, dimension(:), allocatable **parm::sptfcolis**
concentration of the facel caliform in the septic tank effluent (cfu/100ml)
- real *8, dimension(:), allocatable **parm::failyr**
- real *8, dimension(:), allocatable **parm::qstemm**
- real *8, dimension(:), allocatable **parm::bio_amn**

- real *8, dimension(:), allocatable **parm::bio_bod**
- real *8, dimension(:), allocatable **parm::biom**
- real *8, dimension(:), allocatable **parm::rbiom**
- real *8, dimension(:), allocatable **parm::fcoli**
- real *8, dimension(:), allocatable **parm::bio_ntr**
- real *8, dimension(:), allocatable **parm::bz_perc**
- real *8, dimension(:), allocatable **parm::sep_cap**
number of permanent residents in the hourse (none)
- real *8, dimension(:), allocatable **parm::plqm**
- real *8, dimension(:), allocatable **parm::bz_area**
- real *8, dimension(:), allocatable **parm::bz_z**
Depth of biozone layer(mm)
- real *8, dimension(:), allocatable **parm::bz_thk**
thickness of biozone (mm)
- real *8, dimension(:), allocatable **parm::bio_bd**
density of biomass (kg/m³) carbon outputs for .hru file
- real *8, dimension(:), allocatable **parm::cmup_kgh**
- real *8, dimension(:), allocatable **parm::cmtot_kgh**
- real *8, dimension(:), allocatable **parm::coeff_denitr**
denitrification rate coefficient (none)
- real *8, dimension(:), allocatable **parm::coeff_bod_dc**
BOD decay rate coefficient (m³/day)
- real *8, dimension(:), allocatable **parm::coeff_bod_conv**
BOD to live bacteria biomass conversion factor (none)
- real *8, dimension(:), allocatable **parm::coeff_fc1**
field capacity calibration parameter 1 (none)
- real *8, dimension(:), allocatable **parm::coeff_fc2**
field capacity calibration parameter 2 (none)
- real *8, dimension(:), allocatable **parm::coeff_fecal**
fecal coliform bacteria decay rate coefficient (m³/day)
- real *8, dimension(:), allocatable **parm::coeff_mrt**
mortality rate coefficient (none)
- real *8, dimension(:), allocatable **parm::coeff_nitr**
nitrification rate coefficient (none)
- real *8, dimension(:), allocatable **parm::coeff_plq**
conversion factor for plaque from TDS (none)
- real *8, dimension(:), allocatable **parm::coeff_rsp**
respiration rate coefficient (none)
- real *8, dimension(:), allocatable **parm::coeff_slg1**
slough-off calibration parameter (none)
- real *8, dimension(:), allocatable **parm::coeff_slg2**
slough-off calibration parameter (none)
- real *8, dimension(:), allocatable **parm::coeff_pdistrb**
- real *8, dimension(:), allocatable **parm::coeff_solpslp**
- real *8, dimension(:), allocatable **parm::coeff_solpintc**
- real *8, dimension(:), allocatable **parm::coeff_psorpmax**
- integer, dimension(:), allocatable **parm::isep_typ**
septic system type (none)
- integer, dimension(:), allocatable **parm::i_sep**
- integer, dimension(:), allocatable **parm::isep_opt**
septic system operation flag (1=active, 2=failing, 3=not operated) (none)
- integer, dimension(:), allocatable **parm::sep_tsincefail**

- integer, dimension(:), allocatable **parm::isep_tfail**
- integer, dimension(:), allocatable **parm::isep_iyr**
- integer, dimension(:), allocatable **parm::sep_strm_dist**
- integer, dimension(:), allocatable **parm::sep_den**
- real *8, dimension(:), allocatable **parm::sol_sumno3**
- real *8, dimension(:), allocatable **parm::sol_sumsolp**
- real *8, dimension(:), allocatable **parm::strsw_sum**
- real *8, dimension(:), allocatable **parm::strstmp_sum**
- real *8, dimension(:), allocatable **parm::strsn_sum**
- real *8, dimension(:), allocatable **parm::strsp_sum**
- real *8, dimension(:), allocatable **parm::strsa_sum**
- real *8, dimension(:), allocatable **parm::spill_hru**
- real *8, dimension(:), allocatable **parm::tile_out**
- real *8, dimension(:), allocatable **parm::hru_in**
- real *8, dimension(:), allocatable **parm::spill_precip**
- real *8, dimension(:), allocatable **parm::pot_seep**
- real *8, dimension(:), allocatable **parm::pot_evap**
- real *8, dimension(:), allocatable **parm::pot_sedin**
- real *8, dimension(:), allocatable **parm::pot_solp**
soluble P loss rate in the pothole (.01 - 0.5) (1/d)
- real *8, dimension(:), allocatable **parm::pot_solpi**
- real *8, dimension(:), allocatable **parm::pot_orgp**
- real *8, dimension(:), allocatable **parm::pot_orgpi**
- real *8, dimension(:), allocatable **parm::pot_orgn**
- real *8, dimension(:), allocatable **parm::pot_orgni**
- real *8, dimension(:), allocatable **parm::pot_mps**
- real *8, dimension(:), allocatable **parm::pot_mpsi**
- real *8, dimension(:), allocatable **parm::pot_mpa**
- real *8, dimension(:), allocatable **parm::pot_mpai**
- real *8, dimension(:), allocatable **parm::pot_no3i**
- real *8, dimension(:), allocatable **parm::precip_in**
- real *8, dimension(:), allocatable **parm::tile_sedo**
- real *8, dimension(:), allocatable **parm::tile_no3o**
- real *8, dimension(:), allocatable **parm::tile_solpo**
- real *8, dimension(:), allocatable **parm::tile_orgno**
- real *8, dimension(:), allocatable **parm::tile_orgpo**
- real *8, dimension(:), allocatable **parm::tile_minpso**
- real *8, dimension(:), allocatable **parm::tile_minpao**
- integer **parm::ia_b**
- integer **parm::ihumus**
- integer **parm::itemp**
- integer **parm::isnow**
- integer, dimension(46) **parm::ipdvar**
output variable codes for output.rch file (none)
- integer, dimension(mhruo) **parm::ipdvas**
output variable codes for output.hru file (none)
- integer, dimension(msubo) **parm::ipdvab**
output variable codes for output.sub file (none)
- integer, dimension(:), allocatable **parm::ipdhru**
HRUs whose output information will be printed to the output.hru and output.wtr files.
- real *8, dimension(mstdo) **parm::wshddayo**

wshddayo(1) average amount of precipitation in watershed for the day (mm H₂O)
wshddayo(3) surface runoff in watershed for day (mm H₂O)
wshddayo(4) lateral flow contribution to streamflow in watershed for day (mm H₂O)
wshddayo(5) water percolation past bottom of soil profile in watershed for day (mm H₂O)
wshddayo(6) water yield to streamflow from HRUs in watershed for day (mm H₂O)
wshddayo(7) actual evapotranspiration in watershed for day (mm H₂O)
wshddayo(12) sediment yield from HRUs in watershed for day (metric tons or metric tons/ha)
wshddayo(35) amount of water stored in soil profile in watershed at end of day (mm H₂O)
wshddayo(41) organic P loading to stream in watershed for day (kg P/ha)
wshddayo(42) nitrate loading to stream in surface runoff in watershed for day (kg N/ha)
wshddayo(43) soluble P loading to stream in watershed for day (kg P/ha)
wshddayo(44) plant uptake of N in watershed for day (kg N/ha)
wshddayo(45) nitrate loading to stream in lateral flow in watershed for day (kg N/ha)
wshddayo(46) nitrate percolation past bottom of soil profile in watershed for day (kg N/ha)
wshddayo(104) groundwater contribution to stream in watershed on day (mm H₂O)
wshddayo(108) potential evapotranspiration in watershed on day (mm H₂O)
wshddayo(109) drainage tile flow contribution to stream in watershed on day (mm H₂O)

- real *8, dimension(mstdo) [parm::wshdmono](#)

watershed monthly output array (see definitions for wshddayo array elements) (varies)
wshdmono(1) average amount of precipitation in watershed for the month (mm H₂O)
wshdmono(3) surface runoff in watershed for month (mm H₂O)
wshdmono(4) lateral flow contribution to streamflow in watershed for month (mm H₂O)
wshdmono(5) water percolation past bottom of soil profile in watershed for month (mm H₂O)
wshdmono(6) water yield to streamflow from HRUs in watershed for month (mm H₂O)
wshdmono(7) actual evapotranspiration in watershed for month (mm H₂O)
wshdmono(8) average maximum temperature in watershed for the month (deg C)
wshdmono(9) average minimum temperature in watershed for the month (deg C)
wshdmono(12) sediment yield from HRUs in watershed for the month (metric tons)
wshdmono(39) freezing rain/snow fall in watershed for the month (mm H₂O)
wshdmono(40) organic N loading to stream in watershed for the month (kg N/ha)
wshdmono(41) organic P loading to stream in watershed for the month (kg P/ha)
wshdmono(42) nitrate loading to stream in surface runoff in watershed for the month (kg N/ha)
wshdmono(43) soluble P loading to stream in watershed for the month (kg P/ha)
wshdmono(44) plant uptake of N in watershed for the month (kg N/ha)
wshdmono(45) nitrate loading to stream in lateral flow in watershed for the month (kg N/ha)
wshdmono(46) nitrate percolation past bottom of soil profile in watershed for the month (kg N/ha)
wshdmono(104) groundwater contribution to stream in watershed for the month (mm H₂O)
wshdmono(108) potential evapotranspiration in watershed for the month (mm H₂O)
wshdmono(109) drainage tile flow contribution to stream in watershed for the month (mm H₂O)

- real *8, dimension(mstdo) [parm::wshdyro](#)

watershed annual output array (varies)
wshdyro(1) average amount of precipitation in watershed for the year (mm H₂O)
wshdyro(3) surface runoff in watershed for year (mm H₂O)
wshdyro(4) lateral flow contribution to streamflow in watershed for year (mm H₂O)
wshdyro(5) water percolation past bottom of soil profile in watershed for year (mm H₂O)
wshdyro(6) water yield to streamflow from HRUs in watershed for year (mm H₂O)
wshdyro(7) actual evapotranspiration in watershed for year (mm H₂O)
wshdyro(8) average maximum temperature in watershed for the year (deg C)
wshdyro(9) average minimum temperature in watershed for the year (deg C)
wshdyro(12) sediment yield from HRUs in watershed for the year (metric tons)
wshdyro(40) organic N loading to stream in watershed for the year (kg N/ha)
wshdyro(41) organic P loading to stream in watershed for the year (kg P/ha)
wshdyro(42) nitrate loading to stream in surface runoff in watershed for the year (kg N/ha)
wshdyro(43) soluble P loading to stream in watershed for the year (kg P/ha)
wshdyro(44) plant uptake of N in watershed for the year
wshdyro(45) nitrate loading to stream in lateral flow in watershed for the year (kg N/ha)
wshdyro(46) nitrate percolation past bottom of soil profile in watershed for the year (kg N/ha)
wshdyro(104) groundwater contribution to stream in watershed for the year (mm H₂O)
wshdyro(108) potential evapotranspiration in watershed for the year (mm H₂O)
wshdyro(109) drainage tile flow contribution to stream in watershed for the year (mm H₂O)

- real *8, dimension(16) [parm::fcstaa](#)
- real *8, dimension(mstdo) [parm::wshdaao](#)

- watershed average annual output array (varies)*
wshdaao(1) precipitation in watershed (mm H2O)
wshdaao(3) surface runoff loading to main channel in watershed (mm H2O)
wshdaao(4) lateral flow loading to main channel in watershed (mm H2O)
wshdaao(5) percolation of water out of root zone in watershed (mm H2O)
wshdaao(7) actual evapotranspiration in watershed (mm H2O)
wshdaao(13) sediment loading to ponds in watershed (metric tons)
wshdaao(14) sediment loading from ponds in watershed (metric tons)
wshdaao(15) net change in sediment level in ponds in watershed (metric tons)
wshdaao(19) evaporation from ponds in watershed (m^3 H2O)
wshdaao(20) seepage from ponds in watershed (m^3 H2O)
wshdaao(21) precipitation on ponds in watershed (m^3 H2O)
wshdaao(22) volume of water entering ponds in watershed (m^3 H2O)
wshdaao(23) volume of water leaving ponds in watershed (m^3 H2O)
wshdaao(38) transmission losses in watershed (mm H2O)
- real *8, dimension(:,:), allocatable **parm::wpstdayo**
 - real *8, dimension(:,:), allocatable **parm::wpstmono**
 - real *8, dimension(:,:), allocatable **parm::wpstyro**
 - real *8, dimension(:,:), allocatable **parm::bio_hv**
harvested biomass (dry weight) (kg/ha)
 - real *8, dimension(:,:), allocatable **parm::yldkg**
yield (dry weight) by crop type in the HRU (kg/ha)
 - real *8, dimension(:,:), allocatable **parm::rchmono**
reach monthly output array (varies)
rchmono(1,:) flow into reach during month (m^3/s)
rchmono(2,:) flow out of reach during month (m^3/s)
rchmono(3,:) sediment transported into reach during month (metric tons)
rchmono(4,:) sediment transported out of reach during month (metric tons)
rchmono(5,:) sediment concentration in outflow during month (mg/L)
rchmono(6,:) organic N transported into reach during month (kg N)
rchmono(7,:) organic N transported out of reach during month (kg N)
rchmono(8,:) organic P transported into reach during month (kg P)
rchmono(9,:) organic P transported out of reach during month (kg P)
rchmono(10,:) evaporation from reach during month (m^3/s)
rchmono(11,:) transmission losses from reach during month (m^3/s)
rchmono(12,:) conservative metal #1 transported out of reach during month (kg)
rchmono(13,:) conservative metal #2 transported out of reach during month (kg)
rchmono(14,:) conservative metal #3 transported out of reach during month (kg)
rchmono(15,:) nitrate transported into reach during month (kg N)
rchmono(16,:) nitrate transported out of reach during month (kg N)
rchmono(17,:) soluble P transported into reach during month (kg P)
rchmono(18,:) soluble P transported out of reach during month (kg P)
rchmono(19,:) soluble pesticide transported into reach during month (mg pst)
rchmono(20,:) soluble pesticide transported out of reach during month (mg pst)
rchmono(21,:) sorbed pesticide transported into reach during month (mg pst)
rchmono(22,:) sorbed pesticide transported out of reach during month (mg pst)
rchmono(23,:) amount of pesticide lost through reactions in reach during month (mg pst)
rchmono(24,:) amount of pesticide lost through volatilization from reach during month (mg pst)
rchmono(25,:) amount of pesticide settling out of reach to bed sediment during month (mg pst)
rchmono(26,:) amount of pesticide resuspended from bed sediment to reach during month (mg pst)
rchmono(27,:) amount of pesticide diffusing from reach to bed sediment during month (mg pst)
rchmono(28,:) amount of pesticide in sediment layer lost through reactions during month (mg pst)
rchmono(29,:) amount of pesticide in sediment layer lost through burial during month (mg pst)
rchmono(30,:) chlorophyll-a transported into reach during month (kg chla)
rchmono(31,:) chlorophyll-a transported out of reach during month (kg chla)
rchmono(32,:) ammonia transported into reach during month (kg N)
rchmono(33,:) ammonia transported out of reach during month (kg N)
rchmono(34,:) nitrite transported into reach during month (kg N)
rchmono(35,:) nitrite transported out of reach during month (kg N)
rchmono(36,:) CBOD transported into reach during month (kg O2)
rchmono(37,:) CBOD transported out of reach during month (kg O2)
rchmono(38,:) dissolved oxygen transported into reach during month (kg O2)

- rchmono(39,:)* dissolved oxygen transported out of reach during month (kg O2)
- rchmono(40,:)* persistent bacteria transported out of reach during month (kg bact)
- rchmono(41,:)* less persistent bacteria transported out of reach during month (kg bact)
- rchmono(43,:)* total N (org N + no3 + no2 + nh4 outs) (kg)
- rchmono(44,:)* total P (org P + sol p outs) (kg)
- real *8, dimension(:,:), allocatable **parm::rchyro**
 - reach* annual output array (varies)
 - rchyro(1,:)* flow into reach during year (m^3/s)
 - rchyro(2,:)* flow out of reach during year (m^3/s)
 - rchyro(3,:)* sediment transported into reach during year (metric tons)
 - rchyro(4,:)* sediment transported out of reach during year (metric tons)
 - rchyro(5,:)* sediment concentration in outflow during year (mg/L)
 - rchyro(6,:)* organic N transported into reach during year (kg N)
 - rchyro(7,:)* organic N transported out of reach during year (kg N)
 - rchyro(8,:)* organic P transported into reach during year (kg P)
 - rchyro(9,:)* organic P transported out of reach during year (kg P)
 - rchyro(10,:)* evaporation from reach during year (m^3/s)
 - rchyro(11,:)* transmission losses from reach during year (m^3/s)
 - rchyro(12,:)* conservative metal #1 transported out of reach during year (kg)
 - rchyro(13,:)* conservative metal #2 transported out of reach during year (kg)
 - rchyro(14,:)* conservative metal #3 transported out of reach during year (kg)
 - rchyro(15,:)* nitrate transported into reach during year (kg N)
 - rchyro(16,:)* nitrate transported out of reach during year (kg N)
 - rchyro(17,:)* soluble P transported into reach during year (kg P)
 - rchyro(18,:)* soluble P transported out of reach during year (kg P)
 - rchyro(19,:)* soluble pesticide transported into reach during year (mg pst)
 - rchyro(20,:)* soluble pesticide transported out of reach during year (mg pst)
 - rchyro(21,:)* sorbed pesticide transported into reach during year (mg pst)
 - rchyro(22,:)* sorbed pesticide transported out of reach during year (mg pst)
 - rchyro(23,:)* amount of pesticide lost through reactions in reach during year!> (mg pst)
 - rchyro(24,:)* amount of pesticide lost through volatilization from reach during year (mg pst)
 - rchyro(25,:)* amount of pesticide settling out of reach to bed sediment during year (mg pst)
 - rchyro(26,:)* amount of pesticide resuspended from bed sediment to reach during year (mg pst)
 - rchyro(27,:)* amount of pesticide diffusing from reach to bed sediment during year (mg pst)
 - rchyro(28,:)* amount of pesticide in sediment layer lost through reactions during year (mg pst)
 - rchyro(29,:)* amount of pesticide in sediment layer lost through burial during year (mg pst)
 - rchyro(30,:)* chlorophyll-a transported into reach during year (kg chla)
 - rchyro(31,:)* chlorophyll-a transported out of reach during year (kg chla)
 - rchyro(32,:)* ammonia transported into reach during year (kg N)
 - rchyro(33,:)* ammonia transported out of reach during year (kg N)
 - rchyro(34,:)* nitrite transported into reach during year (kg N)
 - rchyro(35,:)* nitrite transported out of reach during year (kg N)
 - rchyro(36,:)* CBOD transported into reach during year (kg O2)
 - rchyro(37,:)* CBOD transported out of reach during year (kg O2)
 - rchyro(38,:)* dissolved oxygen transported into reach during year (kg O2)
 - rchyro(39,:)* dissolved oxygen transported out of reach during year (kg O2)
 - rchyro(40,:)* persistent bacteria transported out of reach during year (kg bact)
 - rchyro(41,:)* less persistent bacteria transported out of reach during year (kg bact)
- real *8, dimension(:,:), allocatable **parm::wpstaao**
- real *8, dimension(:,:), allocatable **parm::hrumono**
 - HRU* monthly output data array (varies)
 - hrumono(1,:)* precipitation in HRU during month (mm H2O)
 - hrumono(2,:)* amount of precipitation falling as freezing rain/snow in HRU during month (mm H2O)
 - hrumono(3,:)* amount of snow melt in HRU during month (mm H2O)
 - hrumono(4,:)* amount of surface runoff to main channel from HRU during month (ignores impact of transmission losses) (mm H2O)
 - hrumono(5,:)* amount of lateral flow contribution to main channel from HRU during month (mm H2O)
 - hrumono(6,:)* amount of groundwater flow contribution to main channel from HRU during month (mm H2O)
 - hrumono(7,:)* amount of water moving from shallow aquifer to plants or soil profile in HRU during mont (mm H2O)h
 - hrumono(8,:)* amount of water recharging deep aquifer in HRU during month (mm H2O)
 - hrumono(9,:)* total amount of water entering both aquifers from HRU during month (mm H2O)
 - hrumono(10,:)* water yield (total amount of water entering main channel) from HRU during month (mm H2O)
 - hrumono(11,:)* amount of water percolating out of the soil profile and into the vadose zone in HRU during month (mm

H2O)

humono(12,:) actual evapotranspiration in HRU during month (mm H2O)
 humono(13,:) amount of transmission losses from tributary channels in HRU for month (mm H2O)
 humono(14,:) sediment yield from HRU for month (metric tons/ha)
 humono(15,:) actual amount of transpiration that occurs during month in HRU (mm H2O)
 humono(16,:) actual amount of evaporation (from soil) that occurs during month in HRU (mm H2O)
 humono(17,:) amount of nitrogen applied in continuous fertilizer operation during month in HRU (kg N/ha)
 humono(18,:) amount of phosphorus applied in continuous fertilizer operation during month in HRU (kg P/ha)
 humono(22,:) amount of irrigation water applied to HRU during month (mm H2O)
 humono(23,:) amount of water removed from shallow aquifer in HRU for irrigation during month (mm H2O)
 humono(24,:) amount of water removed from deep aquifer in HRU for irrigation during month (mm H2O)
 humono(25,:) potential evapotranspiration in HRU during month (mm H2O)
 humono(26,:) monthly amount of N (organic & mineral) applied in HRU during grazing (kg N/ha)
 humono(27,:) monthly amount of P (organic & mineral) applied in HRU during grazing (kg P/ha)
 humono(28,:) monthly amount of N (organic & mineral) auto-applied in HRU (kg N/ha)
 humono(29,:) monthly amount of P (organic & mineral) auto-applied in HRU (kg P/ha)
 humono(31,:) water stress days in HRU during month (stress days)
 humono(32,:) temperature stress days in HRU during month (stress days)
 humono(33,:) nitrogen stress days in HRU during month (stress days)
 humono(34,:) phosphorus stress days in HRU during month (stress days)
 humono(35,:) organic nitrogen in surface runoff in HRU during month (kg N/ha)
 humono(36,:) organic phosphorus in surface runoff in HRU during month (kg P/ha)
 humono(37,:) nitrate in surface runoff in HRU during month (kg N/ha)
 humono(38,:) nitrate in lateral flow in HRU during month (kg N/ha)
 humono(39,:) soluble phosphorus in surface runoff in HRU during month (kg P/ha)
 humono(40,:) amount of nitrogen removed from soil by plant uptake in HRU during month (kg N/ha)
 humono(41,:) nitrate percolating past bottom of soil profile in HRU during month (kg N/ha)
 humono(42,:) amount of phosphorus removed from soil by plant uptake in HRU during month (kg P/ha)
 humono(43,:) amount of phosphorus moving from labile mineral to active mineral pool in HRU during month (kg P/ha)
 humono(44,:) amount of phosphorus moving from active mineral to stable mineral pool in HRU during month (kg P/ha)
 humono(45,:) amount of nitrogen applied to HRU in fertilizer and grazing operations during month (kg N/ha)
 humono(46,:) amount of phosphorus applied to HRU in fertilizer and grazing operations during month (kg P/ha)
 humono(47,:) amount of nitrogen added to soil by fixation in HRU during month (kg N/ha)
 humono(48,:) amount of nitrogen lost by denitrification in HRU during month (kg N/ha)
 humono(49,:) amount of nitrogen moving from active organic to nitrate pool in HRU during month (kg N/ha)
 humono(50,:) amount of nitrogen moving from active organic to stable organic pool in HRU during month (kg N/ha)
 humono(51,:) amount of phosphorus moving from organic to labile mineral pool in HRU during month (kg P/ha)
 humono(52,:) amount of nitrogen moving from fresh organic to nitrate and active organic pools in HRU during month (kg N/ha)
 humono(53,:) amount of phosphorus moving from fresh organic to the labile mineral and organic pools in HRU during month (kg P/ha)
 humono(54,:) amount of nitrogen added to soil in rain (kg N/ha)
 humono(61,:) daily soil loss predicted with USLE equation (metric tons/ha)
 humono(63,:) less persistent bacteria transported to main channel from HRU during month (#bacteria/ha)
 humono(64,:) persistent bacteria transported to main channel from HRU during month (#bacteria/ha)
 humono(65,:) nitrate loading from groundwater in HRU to main channel during month (kg N/ha)
 humono(66,:) soluble P loading from groundwater in HRU to main channel during month (kg P/ha)
 humono(67,:) loading of mineral P attached to sediment in HRU to main channel during month (kg P/ha)

- real *8, dimension(:,:), allocatable [parm::rchdy](#)

rchdy(1,:) flow into reach on day (m^3/s)
 rchdy(2,:) flow out of reach on day (m^3/s)
 rchdy(3,:) evaporation from reach on day (m^3/s)
 rchdy(4,:) transmission losses from reach on day (m^3/s)
 rchdy(5,:) sediment transported into reach on day (metric tons)
 rchdy(6,:) sediment transported out of reach on day (metric tons)
 rchdy(7,:) sediment concentration in outflow (mg/L)
 rchdy(8,:) organic N transported into reach on day (kg N)
 rchdy(9,:) organic N transported out of reach on day (kg N)
 rchdy(10,:) organic P transported into reach on day (kg P)
 rchdy(11,:) organic P transported out of reach on day (kg P)
 rchdy(12,:) nitrate transported into reach on day (kg N)

rchdy(13,:) nitrate transported out of reach on day (kg N)
rchdy(14,:) ammonia transported into reach on day (kg N)
rchdy(15,:) ammonia transported out of reach on day (kg N)
rchdy(16,:) nitrite transported into reach on day (kg N)
rchdy(17,:) nitrite transported out of reach on day (kg N)
rchdy(18,:) soluble P transported into reach on day (kg P)
rchdy(19,:) soluble P transported out of reach on day (kg P)
rchdy(20,:) chlorophyll-a transported into reach on day (kg chl-a)
rchdy(21,:) chlorophyll-a transported out of reach on day (kg chl-a)
rchdy(22,:) CBOD transported into reach on day (kg O₂)
rchdy(23,:) CBOD transported out of reach on day (kg O₂)
rchdy(24,:) dissolved oxygen transported into reach on day (kg O₂)
rchdy(25,:) dissolved oxygen transported out of reach on day (kg O₂)
rchdy(26,:) soluble pesticide transported into reach on day (mg pst)
rchdy(27,:) soluble pesticide transported out of reach on day (mg pst)
rchdy(28,:) sorbed pesticide transported into reach on day (mg pst)
rchdy(29,:) sorbed pesticide transported out of reach on day (mg pst)
rchdy(30,:) amount of pesticide lost through reactions in reach on day (mg pst)
rchdy(31,:) amount of pesticide lost through volatilization from reach on day (mg pst)
rchdy(32,:) amount of pesticide settling out of reach to bed sediment on day (mg pst)
rchdy(33,:) amount of pesticide resuspended from bed sediment to reach on day (mg pst)
rchdy(34,:) amount of pesticide diffusing from reach to bed sediment on day (mg pst)
rchdy(35,:) amount of pesticide in sediment layer lost through reactions on day (mg pst)
rchdy(36,:) amount of pesticide in sediment layer lost through burial on day (mg pst)
rchdy(37,:) amount of pesticide stored in river bed sediments (mg pst)
rchdy(38,:) persistent bacteria transported out of reach on day (kg bact)
rchdy(39,:) less persistent bacteria transported out of reach on day (kg bact)
rchdy(40,:) amount of conservative metal #1 transported out of reach on day (kg)
rchdy(41,:) amount of conservative metal #2 transported out of reach on day (kg)
rchdy(42,:) amount of conservative metal #3 transported out of reach on day (kg)
rchdy(43,:) total N (org N + no3 + no2 + nh4 outs) (kg)
rchdy(44,:) total P (org P + sol p outs) (kg)

- real *8, dimension(:,:), allocatable [parm::hruyro](#)

HRU annual output array (varies) *hruyro(1,:)* precipitation in HRU during year (mm H₂O)
hruyro(2,:) amount of precipitation falling as freezing rain/snow in HRU during year (mm H₂O)
hruyro(3,:) amount of snow melt in HRU during year (mm H₂O)
hruyro(4,:) amount of surface runoff to main channel from HRU during year (ignores impact of transmission losses) (mm H₂O)
hruyro(5,:) amount of lateral flow contribution to main channel from HRU during year (mm H₂O)
hruyro(6,:) amount of groundwater flow contribution to main channel from HRU during year (mm H₂O)
hruyro(7,:) amount of water moving from shallow aquifer to plants or soil profile in HRU during year (mm H₂O)
hruyro(8,:) amount of water recharging deep aquifer in HRU during year (mm H₂O)
hruyro(9,:) total amount of water entering both aquifers from HRU during year (mm H₂O)
hruyro(10,:) water yield (total amount of water entering main channel) from HRU during year (mm H₂O)
hruyro(11,:) amount of water percolating out of the soil profile and into the vadose zone in HRU during year (mm H₂O)
hruyro(12,:) actual evapotranspiration in HRU during year (mm H₂O)
hruyro(13,:) amount of transmission losses from tributary channels in HRU for year (mm H₂O)
hruyro(14,:) sediment yield from HRU for year (metric tons/ha)
hruyro(15,:) actual amount of transpiration that occurs during year in HRU (mm H₂O)
hruyro(16,:) actual amount of evaporation (from soil) that occurs during year in HRU (mm H₂O)
hruyro(17,:) amount of nitrogen applied in continuous fertilizer operation during year in HRU (kg N/ha)
hruyro(18,:) amount of phosphorus applied in continuous fertilizer operation during year in HRU (kg P/ha)
hruyro(23,:) amount of water removed from shallow aquifer in HRU for irrigation during year (mm H₂O)
hruyro(24,:) amount of water removed from deep aquifer in HRU for irrigation during year (mm H₂O)
hruyro(25,:) potential evapotranspiration in HRU during year (mm H₂O)
hruyro(26,:) annual amount of N (organic & mineral) applied in HRU during grazing (kg N/ha)
hruyro(27,:) annual amount of P (organic & mineral) applied in HRU during grazing (kg P/ha)
hruyro(28,:) annual amount of N (organic & mineral) auto-applied in HRU (kg N/ha)
hruyro(29,:) annual amount of P (organic & mineral) auto-applied in HRU (kg P/ha)
hruyro(31,:) water stress days in HRU during year (stress days)
hruyro(32,:) temperature stress days in HRU during year (stress days)
hruyro(33,:) nitrogen stress days in HRU during year (stress days)

- hruyro(34,:) phosphorus stress days in HRU during year (stress days)*
 - hruyro(35,:) organic nitrogen in surface runoff in HRU during year (kg N/ha)*
 - hruyro(36,:) organic phosphorus in surface runoff in HRU during year (kg P/ha)*
 - hruyro(37,:) nitrate in surface runoff in HRU during year (kg N/ha)*
 - hruyro(38,:) nitrate in lateral flow in HRU during year (kg N/ha)*
 - hruyro(39,:) soluble phosphorus in surface runoff in HRU during year (kg P/ha)*
 - hruyro(40,:) amount of nitrogen removed from soil by plant uptake in HRU during year (kg N/ha)*
 - hruyro(41,:) nitrate percolating past bottom of soil profile in HRU during year (kg N/ha)*
 - hruyro(42,:) amount of phosphorus removed from soil by plant uptake in HRU during year (kg P/ha)*
 - hruyro(43,:) amount of phosphorus moving from labile mineral to active mineral pool in HRU during year (kg P/ha)*
 - hruyro(44,:) amount of phosphorus moving from active mineral to stable mineral pool in HRU during year (kg P/ha)*
 - hruyro(45,:) amount of nitrogen applied to HRU in fertilizer and grazing operations during year (kg N/ha)*
 - hruyro(46,:) amount of phosphorus applied to HRU in fertilizer and grazing operations during year (kg P/ha)*
 - hruyro(47,:) amount of nitrogen added to soil by fixation in HRU during year (kg N/ha)*
 - hruyro(48,:) amount of nitrogen lost by denitrification in HRU during year (kg N/ha)*
 - hruyro(49,:) amount of nitrogen moving from active organic to nitrate pool in HRU during year (kg N/ha)*
 - hruyro(50,:) amount of nitrogen moving from active organic to stable organic pool in HRU during year (kg N/ha)*
 - hruyro(51,:) amount of phosphorus moving from organic to labile mineral pool in HRU during year (kg P/ha)*
 - hruyro(52,:) amount of nitrogen moving from fresh organic to nitrate and active organic pools in HRU during year (kg N/ha)*
 - hruyro(53,:) amount of phosphorus moving from fresh organic to the labile mineral and organic pools in HRU during year (kg P/ha)*
 - hruyro(54,:) amount of nitrogen added to soil in rain during year (kg N/ha)*
 - hruyro(61,:) daily soil loss predicted with USLE equation (metric tons/ha)*
 - hruyro(63,:) less persistent bacteria transported to main channel from HRU during year (# bacteria/ha)*
 - hruyro(64,:) persistent bacteria transported to main channel from HRU during year (# bacteria/ha)*
 - hruyro(65,:) nitrate loading from groundwater in HRU to main channel during year (kg N/ha)*
 - hruyro(66,:) soluble P loading from groundwater in HRU to main channel during year (kg P/ha)*
 - hruyro(67,:) loading of mineral P attached to sediment in HRU to main channel during year (kg P/ha)*
- **real *8, dimension(:,:), allocatable [parm::rchaa0](#)**
 - reach average annual output array (varies)*
 - rchaa0(1,:) flow into reach during simulation (m^3/s)*
 - rchaa0(2,:) flow out of reach during simulation (m^3/s)*
 - rchaa0(3,:) sediment transported into reach during simulation (metric tons)*
 - rchaa0(4,:) sediment transported out of reach during simulation (metric tons)*
 - rchaa0(5,:) sediment concentration in outflow during simulation (mg/L)*
 - rchaa0(6,:) organic N transported into reach during simulation (kg N)*
 - rchaa0(7,:) organic N transported out of reach during simulation (kg N)*
 - rchaa0(8,:) organic P transported into reach during simulation (kg P)*
 - rchaa0(9,:) organic P transported out of reach during simulation (kg P)*
 - rchaa0(10,:) evaporation from reach during simulation (m^3/s)*
 - rchaa0(11,:) transmission losses from reach during simulation (m^3/s)*
 - rchaa0(12,:) conservative metal #1 transported out of reach during simulation (kg)*
 - rchaa0(13,:) conservative metal #2 transported out of reach during simulation (kg)*
 - rchaa0(14,:) conservative metal #3 transported out of reach during simulation (kg)*
 - rchaa0(15,:) nitrate transported into reach during simulation (kg N)*
 - rchaa0(16,:) nitrate transported out of reach during simulation (kg N)*
 - rchaa0(17,:) soluble P transported into reach during simulation (kg P)*
 - rchaa0(18,:) soluble P transported out of reach during simulation (kg P)*
 - rchaa0(19,:) soluble pesticide transported into reach during simulation*
 - rchaa0(20,:) soluble pesticide transported out of reach during simulation*
 - rchaa0(21,:) sorbed pesticide transported into reach during simulation*
 - rchaa0(22,:) sorbed pesticide transported out of reach during simulation*
 - rchaa0(23,:) amount of pesticide lost through reactions in reach during simulation*
 - rchaa0(24,:) amount of pesticide lost through volatilization from reach during simulation*
 - rchaa0(25,:) amount of pesticide settling out of reach to bed sediment during simulation*
 - rchaa0(26,:) amount of pesticide resuspended from bed sediment to reach during simulation*
 - rchaa0(27,:) amount of pesticide diffusing from reach to bed sediment during simulation*
 - rchaa0(28,:) amount of pesticide in sediment layer lost through reactions during simulation*
 - rchaa0(29,:) amount of pesticide in sediment layer lost through burial during simulation*
 - rchaa0(30,:) chlorophyll-a transported into reach during simulation (kg chl-a)*
 - rchaa0(31,:) chlorophyll-a transported out of reach during simulation (kg chl-a)*
 - rchaa0(32,:) ammonia transported into reach during simulation (kg N)*

- rchaao(33,:) ammonia transported out of reach during simulation (kg N)*
- rchaao(34,:) nitrite transported into reach during simulation (kg N)*
- rchaao(35,:) nitrite transported out of reach during simulation (kg N)*
- rchaao(36,:) CBOD transported into reach during simulation (kg O2)*
- rchaao(37,:) CBOD transported out of reach during simulation (kg O2)*
- rchaao(38,:) dissolved oxygen transported into reach during simulation (kg O2)*
- rchaao(39,:) dissolved oxygen transported out of reach during simulation (kg O2)*
- rchaao(40,:) persistent bacteria transported out of reach during simulation (kg bact)*
- rchaao(41,:) less persistent bacteria transported out of reach during simulation (kg bact)*
- rchaao(43,:) Total N (org N + no3 + no2 + nh4 outs) (kg)*
- rchaao(44,:) Total P (org P + sol p outs) (kg)*
- real *8, dimension(:,:), allocatable [parm::submono](#)
 - subbasin monthly output array (varies)*
 - submono(1,:) precipitation in subbasin for month (mm H2O)*
 - submono(2,:) snow melt in subbasin for month (mm H2O)*
 - submono(3,:) surface runoff loading in subbasin for month (mm H2O)*
 - submono(4,:) water yield from subbasin for month (mm H2O)*
 - submono(5,:) potential evapotranspiration in subbasin for month (mm H2O)*
 - submono(6,:) actual evapotranspiration in subbasin for month (mm H2O)*
 - submono(7,:) sediment yield from subbasin for month (metric tons/ha)*
 - submono(8,:) organic N loading from subbasin for month (kg N/ha)*
 - submono(9,:) organic P loading from subbasin for month (kg P/ha)*
 - submono(10,:) NO3 loading from surface runoff in subbasin for month (kg N/ha)*
 - submono(11,:) soluble P loading from subbasin for month (kg P/ha)*
 - submono(12,:) groundwater loading from subbasin for month (mm H2O)*
 - submono(13,:) percolation out of soil profile in subbasin for month (mm H2O)*
 - submono(14,:) loading to reach of mineral P attached to sediment from subbasin for month (kg P/ha)*
- real *8, dimension(:,:), allocatable [parm::subyro](#)
 - subbasin annual output array (varies)*
 - subyro(1,:) precipitation in subbasin for year (mm H2O)*
 - subyro(2,:) snow melt in subbasin for year (mm H2O)*
 - subyro(3,:) surface runoff loading in subbasin for year (mm H2O)*
 - subyro(4,:) water yield from subbasin for year (mm H2O)*
 - subyro(5,:) potential evapotranspiration in subbasin for year (mm H2O)*
 - subyro(6,:) actual evapotranspiration in subbasin for year (mm H2O)*
 - subyro(7,:) sediment yield from subbasin for year (metric tons/ha)*
 - subyro(8,:) organic N loading from subbasin for year (kg N/ha)*
 - subyro(9,:) organic P loading from subbasin for year (kg P/ha)*
 - subyro(10,:) NO3 loading from surface runoff in subbasin for year (kg N/ha)*
 - subyro(11,:) soluble P loading from subbasin for year (kg P/ha)*
 - subyro(12,:) groundwater loading from subbasin for year (mm H2O)*
 - subyro(13,:) percolation out of soil profile in subbasin for year (mm H2O)*
 - subyro(14,:) loading to reach of mineral P attached to sediment from subbasin for year (kg P/ha)*
- real *8, dimension(:,:), allocatable [parm::hruaao](#)
 - HRU average annual output array (varies)*
 - hruaao(1,:) precipitation in HRU during simulation (mm H2O)*
 - hruaao(2,:) amount of precipitation falling as freezing rain/snow in HRU during simulation (mm H2O)*
 - hruaao(3,:) amount of snow melt in HRU during simulation (mm H2O)*
 - hruaao(4,:) amount of surface runoff to main channel from HRU during simulation (ignores impact of transmission losses) (mm H2O)*
 - hruaao(5,:) amount of lateral flow contribution to main channel from HRU during simulation (mm H2O)*
 - hruaao(6,:) amount of groundwater flow contribution to main channel from HRU during simulation (mm H2O)*
 - hruaao(7,:) amount of water moving from shallow aquifer to plants or soil profile in HRU during simulation (mm H2O)*
 - hruaao(8,:) amount of water recharging deep aquifer in HRU during simulation (mm H2O)*
 - hruaao(9,:) total amount of water entering both aquifers from HRU during simulation (mm H2O)*
 - hruaao(10,:) water yield (total amount of water entering main channel) from HRU during simulation (mm H2O)*
 - hruaao(11,:) amount of water percolating out of the soil profile and into the vadose zone in HRU during simulation (mm H2O)*
 - hruaao(12,:) actual evapotranspiration in HRU during simulation*
 - hruaao(13,:) amount of transmission losses from tributary channels in HRU for simulation (mm H2O)*
 - hruaao(14,:) sediment yield from HRU for simulation (metric tons/ha)*
 - hruaao(17,:) amount of nitrogen applied in continuous fertilizer operation in HRU for simulation (kg N/ha)*

- hruaao(18,:)* amount of phosphorus applied in continuous fertilizer operation in HRU for simulation (kg P/ha)
hruaao(23,:) amount of water removed from shallow aquifer in HRU for irrigation during simulation (mm H₂O)
hruaao(24,:) amount of water removed from deep aquifer in HRU for irrigation during simulation (mm H₂O)
hruaao(25,:) potential evapotranspiration in HRU during simulation (mm H₂O)
hruaao(26,:) annual amount of N (organic & mineral) applied in HRU during grazing (kg N/ha)
hruaao(27,:) annual amount of P (organic & mineral) applied in HRU during grazing (kg P/ha)
hruaao(28,:) average annual amount of N (organic & mineral) auto-applied in HRU (kg N/ha)
hruaao(29,:) average annual amount of P (organic & mineral) auto-applied in HRU (kg P/ha)
hruaao(31,:) water stress days in HRU during simulation (stress days)
hruaao(32,:) temperature stress days in HRU during simulation (stress days)
hruaao(33,:) nitrogen stress days in HRU during simulation (stress days)
hruaao(34,:) phosphorus stress days in HRU during simulation (stress days)
hruaao(35,:) organic nitrogen in surface runoff in HRU during simulation (kg N/ha)
hruaao(36,:) organic phosphorus in surface runoff in HRU during simulation (kg P/ha)
hruaao(37,:) nitrate in surface runoff in HRU during simulation (kg N/ha)
hruaao(38,:) nitrate in lateral flow in HRU during simulation (kg N/ha)
hruaao(39,:) soluble phosphorus in surface runoff in HRU during simulation (kg P/ha)
hruaao(40,:) amount of nitrogen removed from soil by plant uptake in HRU during simulation (kg N/ha)
hruaao(41,:) nitrate percolating past bottom of soil profile in HRU during simulation (kg N/ha)
hruaao(42,:) amount of phosphorus removed from soil by plant uptake in HRU during simulation (kg P/ha)
hruaao(43,:) amount of phosphorus moving from labile mineral to active mineral pool in HRU during simulation (kg P/ha)
hruaao(44,:) amount of phosphorus moving from active mineral to stable mineral pool in HRU during simulation (kg P/ha)
hruaao(45,:) amount of nitrogen applied to HRU in fertilizer and grazing operations during simulation (kg N/ha)
hruaao(46,:) amount of phosphorus applied to HRU in fertilizer and grazing operations during simulation (kg P/ha)
hruaao(47,:) amount of nitrogen added to soil by fixation in HRU during simulation (kg N/ha)
hruaao(48,:) amount of nitrogen lost by denitrification in HRU during simulation (kg N/ha)
hruaao(49,:) amount of nitrogen moving from active organic to nitrate pool in HRU during simulation (kg N/ha)
hruaao(50,:) amount of nitrogen moving from active organic to stable organic pool in HRU during simulation (kg N/ha)
hruaao(51,:) amount of phosphorus moving from organic to labile mineral pool in HRU during simulation (kg P/ha)
hruaao(52,:) amount of nitrogen moving from fresh organic to nitrate and active organic pools in HRU during simulation (kg N/ha)
hruaao(53,:) amount of phosphorus moving from fresh organic to the labile mineral and organic pools in HRU during simulation (kg P/ha)
hruaao(54,:) amount of nitrogen added to soil in rain during simulation (kg N/ha)
hruaao(61,:) daily soil loss predicted with USLE equation (metric tons/ha)
hruaao(63,:) less persistent bacteria transported to main channel from HRU during simulation (# bacteria/ha)
hruaao(64,:) persistent bacteria transported to main channel from HRU during simulation (# bacteria/ha)
hruaao(65,:) nitrate loading from groundwater in HRU to main channel during simulation (kg N/ha)
hruaao(66,:) soluble P loading from groundwater in HRU to main channel during simulation (kg P/ha)
hruaao(67,:) loading of mineral P attached to sediment in HRU to main channel during simulation (kg P/ha)
- real *8, dimension(:,:), allocatable [parm::subaao](#)
 subbasin average annual output array (varies)
 - real *8, dimension(:,:), allocatable [parm::resoutm](#)
 reservoir monthly output array (varies)
resoutm(1,:) flow into reservoir during month (m^3/s)
resoutm(2,:) flow out of reservoir during month (m^3/s)
resoutm(3,:) sediment entering reservoir during month (metric tons)
resoutm(4,:) sediment leaving reservoir during month (metric tons)
resoutm(5,:) sediment concentration in reservoir during month (mg/L)
resoutm(6,:) pesticide entering reservoir during month (mg pst)
resoutm(7,:) pesticide lost from reservoir through reactions during month (mg pst)
resoutm(8,:) pesticide lost from reservoir through volatilization during month (mg pst)
resoutm(9,:) pesticide moving from water to sediment through settling during month (mg pst)
resoutm(10,:) pesticide moving from sediment to water through resuspension during month (mg pst)
resoutm(11,:) pesticide moving from water to sediment through diffusion during month (mg pst)
resoutm(12,:) pesticide lost from reservoir sediment layer through reactions during month (mg pst)
resoutm(13,:) pesticide lost from reservoir sediment layer through burial during month (mg pst)
resoutm(14,:) pesticide transported out of reservoir during month (mg pst)
resoutm(15,:) pesticide concentration in reservoir water during month (mg pst/ m^3)
resoutm(16,:) pesticide concentration in reservoir sediment layer during month (mg pst/ m^3)
resoutm(17,:) evaporation from reservoir during month ($m^3 H_2O$)

resoutm(18,:) seepage from reservoir during month ($m^3 H_2O$)
resoutm(19,:) precipitation on reservoir during month ($m^3 H_2O$)
resoutm(22,:) organic N entering reservoir during month (kg N)
resoutm(23,:) organic N leaving reservoir during month (kg N)
resoutm(24,:) organic P entering reservoir during month (kg P)
resoutm(25,:) organic P leaving reservoir during month (kg P)
resoutm(26,:) nitrate entering reservoir during month (kg N)
resoutm(27,:) nitrate leaving reservoir during month (kg N)
resoutm(28,:) nitrite entering reservoir during month (kg N)
resoutm(29,:) nitrite leaving reservoir during month (kg N)
resoutm(30,:) ammonia entering reservoir during month (kg N)
resoutm(31,:) ammonia leaving reservoir during month (kg N)
resoutm(32,:) mineral P entering reservoir during month (kg P)
resoutm(33,:) mineral P leaving reservoir during month (kg P)
resoutm(34,:) chlorophyll-a entering reservoir during month (kg chl_a)
resoutm(35,:) chlorophyll-a leaving reservoir during month (kg chl_a)
resoutm(36,:) organic P concentration in reservoir water during month (mg P/L)
resoutm(37,:) mineral P concentration in reservoir water during month (mg P/L)
resoutm(38,:) organic N concentration in reservoir water during month (mg N/L)
resoutm(39,:) nitrate concentration in reservoir water during month (mg N/L)
resoutm(40,:) nitrite concentration in reservoir water during month (mg N/L)
resoutm(41,:) ammonia concentration in reservoir water during month (mg N/L)

- real *8, dimension(:,:), allocatable [parm::resouty](#)

reservoir annual output array (varies)
resouty(1,:) flow into reservoir during year (m^3/s)
resouty(2,:) flow out of reservoir during year (m^3/s)
resouty(3,:) sediment entering reservoir during year (metric tons)
resouty(4,:) sediment leaving reservoir during year (metric tons)
resouty(5,:) sediment concentration in reservoir during year (mg/L)
resouty(6,:) pesticide entering reservoir during year (mg pst)
resouty(7,:) pesticide lost from reservoir through reactions during year (mg pst)
resouty(8,:) pesticide lost from reservoir through volatilization during year (mg pst)
resouty(9,:) pesticide moving from water to sediment through settling during year (mg pst)
resouty(10,:) pesticide moving from sediment to water through resuspension during year (mg pst)
resouty(11,:) pesticide moving from water to sediment through diffusion during year (mg pst)
resouty(12,:) pesticide lost from reservoir sediment layer through reactions during year (mg pst)
resouty(13,:) pesticide lost from reservoir sediment layer through burial during year (mg pst)
resouty(14,:) pesticide transported out of reservoir during year (mg pst)
resouty(15,:) pesticide concentration in reservoir water during year (mg pst/ m^3)
resouty(16,:) pesticide concentration in reservoir sediment layer during year (mg pst/ m^3)
resouty(17,:) evaporation from reservoir during year ($m^3 H_2O$)
resouty(18,:) seepage from reservoir during year ($m^3 H_2O$)
resouty(19,:) precipitation on reservoir during year ($m^3 H_2O$)
resouty(22,:) organic N entering reservoir during year (kg N)
resouty(23,:) organic N leaving reservoir during year (kg N)
resouty(24,:) organic P entering reservoir during year (kg P)
resouty(25,:) organic P leaving reservoir during year (kg P)
resouty(26,:) nitrate entering reservoir during year (kg N)
resouty(27,:) nitrate leaving reservoir during year (kg N)
resouty(28,:) nitrite entering reservoir during year (kg N)
resouty(29,:) nitrite leaving reservoir during year (kg N)
resouty(30,:) ammonia entering reservoir during year (kg N)
resouty(31,:) ammonia leaving reservoir during year (kg N)
resouty(32,:) mineral P entering reservoir during year (kg P)
resouty(33,:) mineral P leaving reservoir during year (kg P)
resouty(34,:) chlorophyll-a entering reservoir during year (kg chl_a)
resouty(35,:) chlorophyll-a leaving reservoir during year (kg chl_a)
resouty(36,:) organic P concentration in reservoir water during year (mg P/L)
resouty(37,:) mineral P concentration in reservoir water during year (mg P/L)
resouty(38,:) organic N concentration in reservoir water during year (mg N/L)
resouty(39,:) nitrate concentration in reservoir water during year (mg N/L)
resouty(40,:) nitrite concentration in reservoir water during year (mg N/L)
resouty(41,:) ammonia concentration in reservoir water during year (mg N/L)

- real *8, dimension(:,:), allocatable [parm::resouta](#)
reservoir average annual output array (varies)
resouta(3,:) sediment entering reservoir during simulation (metric tons)
resouta(4,:) sediment leaving reservoir during simulation (metric tons)
resouta(17,:) evaporation from reservoir during simulation ($m^3 H_2O$)
resouta(18,:) seepage from reservoir during simulation ($m^3 H_2O$)
resouta(19,:) precipitation on reservoir during simulation ($m^3 H_2O$)
resouta(20,:) water entering reservoir during simulation ($m^3 H_2O$)
resouta(21,:) water leaving reservoir during simulation ($m^3 H_2O$)
- real *8, dimension(12, 8) [parm::wshd_aamon](#)
wshd_aamon(:,1) average annual precipitation in watershed falling during month (mm H2O)
wshd_aamon(:,2) average annual freezing rain in watershed falling during month (mm H2O)
wshd_aamon(:,3) average annual surface runoff in watershed during month (mm H2O)
wshd_aamon(:,4) average annual lateral flow in watershed during month (mm H2O)
wshd_aamon(:,5) average annual water yield in watershed during month (mm H2O)
wshd_aamon(:,6) average annual actual evapotranspiration in watershed during month (mm H2O)
wshd_aamon(:,7) average annual sediment yield in watershed during month (metric tons)
wshd_aamon(:,8) average annual potential evapotranspiration in watershed during month (mm H2O)
- real *8, dimension(:,:), allocatable [parm::wtrmon](#)
HRU monthly output data array for impoundments (varies)
wtrmon(1,:) evaporation from ponds in HRU for month (mm H2O)
wtrmon(2,:) seepage from ponds in HRU for month (mm H2O)
wtrmon(3,:) precipitation on ponds in HRU for month (mm H2O)
wtrmon(4,:) amount of water entering ponds in HRU for month (mm H2O)
wtrmon(5,:) sediment entering ponds in HRU for month (metric tons/ha)
wtrmon(6,:) amount of water leaving ponds in HRU for month (mm H2O)
wtrmon(7,:) sediment leaving ponds in HRU for month (metric tons/ha)
wtrmon(8,:) precipitation on wetlands in HRU for month (mm H2O)
wtrmon(9,:) volume of water entering wetlands from HRU for month (mm H2O)
wtrmon(10,:) sediment loading to wetlands for month from HRU (metric tons/ha)
wtrmon(11,:) evaporation from wetlands in HRU for month (mm H2O)
wtrmon(12,:) seepage from wetlands in HRU for month (mm H2O)
wtrmon(13,:) volume of water leaving wetlands in HRU for month (mm H2O)
wtrmon(14,:) sediment loading from wetlands in HRU to main channel during month (metric tons/ha)
wtrmon(15,:) precipitation on potholes in HRU for month (mm H2O)
wtrmon(16,:) evaporation from potholes in HRU for month (mm H2O)
wtrmon(17,:) seepage from potholes in HRU for month (mm H2O)
wtrmon(18,:) water leaving potholes in HRU for month (mm H2O)
wtrmon(19,:) water entering potholes in HRU for month (mm H2O)
wtrmon(20,:) sediment entering potholes in HRU for month (metric tons/ha)
wtrmon(21,:) sediment leaving potholes in HRU for month (metric tons/ha)
- real *8, dimension(:,:), allocatable [parm::wtryr](#)
HRU impoundment annual output array (varies)
wtryr(1,:) evaporation from ponds in HRU for year (mm H2O)
wtryr(2,:) seepage from ponds in HRU for year (mm H2O)
wtryr(3,:) precipitation on ponds in HRU for year (mm H2O)
wtryr(4,:) amount of water entering ponds in HRU for year (mm H2O)
wtryr(5,:) sediment entering ponds in HRU for year (metric tons/ha)
wtryr(6,:) amount of water leaving ponds in HRU for year (mm H2O)
wtryr(7,:) sediment leaving ponds in HRU for year (metric tons/ha)
wtryr(8,:) precipitation on wetlands in HRU for year (mm H2O)
wtryr(9,:) volume of water entering wetlands from HRU for year (mm H2O)
wtryr(10,:) sediment loading to wetlands for year from HRU (metric tons/ha)
wtryr(11,:) evaporation from wetlands in HRU for year (mm H2O)
wtryr(12,:) seepage from wetlands in HRU for year (mm H2O)
wtryr(13,:) volume of water leaving wetlands in HRU for year (mm H2O)
wtryr(14,:) sediment loading from wetlands in HRU to main channel during year (metric tons/ha)
wtryr(15,:) precipitation on potholes in HRU during year (mm H2O)
wtryr(16,:) evaporation from potholes in HRU during year (mm H2O)
wtryr(17,:) seepage from potholes in HRU during year (mm H2O)
wtryr(18,:) water leaving potholes in HRU during year (mm H2O)
wtryr(19,:) water entering potholes in HRU during year (mm H2O)

- wtryr(20,:)* sediment entering potholes in HRU during year (metric tons/ha)
- wtryr(21,:)* sediment leaving potholes in HRU during year (metric tons/ha)
- real *8, dimension(:,:), allocatable **parm::wtraa**
HRU impoundment average annual output array (varies)
- real *8, dimension(:,:), allocatable **parm::sub_smfmx**
max melt rate for snow during year (June 21) for subbasin(:) where deg C refers to the air temperature. SUB_SMFmx and SMFMN allow the rate of snow melt to vary through the year. These parameters are accounting for the impact of soil temperature on snow melt (range: -5.0/5.0) (mm/deg C/day)
- real *8, dimension(:,:), allocatable **parm::sub_smfmn**
min melt rate for snow during year (Dec 21) for subbasin(:) (range: -5.0/5.0) where deg C refers to the air temperature (mm/deg C/day)
- real *8, dimension(:,:), allocatable **parm::hrupstd**
hrupstd(:,1,:) amount of pesticide type in surface runoff contribution to stream from HRU on day (in solution) (mg pst)
hrupstd(:,2,:) amount of pesticide type in surface runoff contribution to stream from HRU on day (sorbed to sediment) (mg pst)
- real *8, dimension(:,:), allocatable **parm::hrupstm**
hrupstm(:, :, :) HRU monthly pesticide output array (varies)
hrupstm(:,1,:) amount of pesticide type in surface runoff contribution to stream from HRU during month (in solution) (mg pst)
hrupstm(:,2,:) amount of pesticide type in surface runoff contribution to stream from HRU during month (sorbed to sediment) (mg pst)
hrupstm(:,3,:) total pesticide loading to stream in surface runoff from HRU during month (mg pst)
- real *8, dimension(:,:), allocatable **parm::hrupsta**
HRU average annual pesticide output array (varies)
- real *8, dimension(:,:), allocatable **parm::hrupsty**
hrupsty(:, :, :) HRU annual pesticide output array (varies)
hrupsty(:,1,:) amount of pesticide type in surface runoff contribution to stream from HRU during year (in solution) (mg pst)
hrupsty(:,2,:) amount of pesticide type in surface runoff contribution to stream from HRU during year (sorbed to sediment) (mg pst)
- integer, dimension(:), allocatable **parm::ifirstt**
temperature data search code (none)
0 first day of temperature data located in file
1 first day of temperature data not located in file
- integer, dimension(:), allocatable **parm::ifirstpcp**
- integer, dimension(:), allocatable **parm::elevp**
elevation of precipitation gage station (m)
- integer, dimension(:), allocatable **parm::elevt**
elevation of temperature gage station (m)
- real *8, dimension(:,:), allocatable **parm::ftmpmn**
avg monthly minimum air temperature (deg C)
- real *8, dimension(:,:), allocatable **parm::ftmpmx**
avg monthly maximum air temperature (deg C)
- real *8, dimension(:,:), allocatable **parm::ftmpstdmn**
standard deviation for avg monthly minimum air temperature (deg C)
- real *8, dimension(:,:), allocatable **parm::ftmpstdmx**
standard deviation for avg monthly maximum air temperature (deg C)
- real *8, dimension(:,:), allocatable **parm::fpcp_stat**
fpcp_stat(:,1,:): average amount of precipitation falling in one day for the month (mm/day)
fpcp_stat(:,2,:): standard deviation for the average daily precipitation (mm/day)
fpcp_stat(:,3,:): skew coefficient for the average daily precipitation (none)
- real *8, dimension(:,:), allocatable **parm::fpr_w1**
probability of wet day after dry day in month (none)
- real *8, dimension(:,:), allocatable **parm::fpr_w2**
probability of wet day after wet day in month (none)

- real *8, dimension(:,:), allocatable **parm::fpr_w3**
proportion of wet days in the month (none)
- real *8, dimension(:), allocatable **parm::ch_d**
average depth of main channel (m)
- real *8, dimension(:), allocatable **parm::flwin**
- real *8, dimension(:), allocatable **parm::flwout**
- real *8, dimension(:), allocatable **parm::bankst**
- real *8, dimension(:), allocatable **parm::ch_wi**
- real *8, dimension(:), allocatable **parm::ch_onco**
channel organic n concentration (ppm)
- real *8, dimension(:), allocatable **parm::ch_opco**
channel organic p concentration (ppm)
- real *8, dimension(:), allocatable **parm::ch_orgn**
- real *8, dimension(:), allocatable **parm::ch_orgp**
- real *8, dimension(:), allocatable **parm::drift**
amount of pesticide drifting onto main channel in subbasin (kg)
- real *8, dimension(:), allocatable **parm::rch_dox**
- real *8, dimension(:), allocatable **parm::rch_bactp**
- real *8, dimension(:), allocatable **parm::alpha_bnk**
alpha factor for bank storage recession curve (days)
- real *8, dimension(:), allocatable **parm::alpha_bnke**
 $\exp(-\alpha_{bnk})$ (none)
- real *8, dimension(:), allocatable **parm::disolvp**
- real *8, dimension(:), allocatable **parm::algae**
- real *8, dimension(:), allocatable **parm::sedst**
- real *8, dimension(:), allocatable **parm::rchstor**
- real *8, dimension(:), allocatable **parm::organicn**
- real *8, dimension(:), allocatable **parm::organicp**
- real *8, dimension(:), allocatable **parm::chlora**
- real *8, dimension(:), allocatable **parm::ch_li**
initial length of main channel (km)
- real *8, dimension(:), allocatable **parm::ch_si**
initial slope of main channel (m/m)
- real *8, dimension(:), allocatable **parm::nitraten**
- real *8, dimension(:), allocatable **parm::nitriten**
- real *8, dimension(:), allocatable **parm::ch_bnk_san**
- real *8, dimension(:), allocatable **parm::ch_bnk_sil**
- real *8, dimension(:), allocatable **parm::ch_bnk_cla**
- real *8, dimension(:), allocatable **parm::ch_bnk_gra**
- real *8, dimension(:), allocatable **parm::ch_bed_san**
- real *8, dimension(:), allocatable **parm::ch_bed_sil**
- real *8, dimension(:), allocatable **parm::ch_bed_cla**
- real *8, dimension(:), allocatable **parm::ch_bed_gra**
- real *8, dimension(:), allocatable **parm::depfp**
- real *8, dimension(:), allocatable **parm::depsanfp**
- real *8, dimension(:), allocatable **parm::depsilfp**
- real *8, dimension(:), allocatable **parm::depclafp**
- real *8, dimension(:), allocatable **parm::depsagfp**
- real *8, dimension(:), allocatable **parm::deplagfp**
- real *8, dimension(:), allocatable **parm::depch**
- real *8, dimension(:), allocatable **parm::depsanch**
- real *8, dimension(:), allocatable **parm::depsilch**
- real *8, dimension(:), allocatable **parm::depclach**

- real *8, dimension(:), allocatable **parm::depsagch**
- real *8, dimension(:), allocatable **parm::deplagch**
- real *8, dimension(:), allocatable **parm::depgrach**
- real *8, dimension(:), allocatable **parm::depgrafp**
- real *8, dimension(:), allocatable **parm::grast**
- real *8, dimension(:), allocatable **parm::r2adj**
curve number retention parameter adjustment factor to adjust surface runoff for flat slopes (0.5 - 3.0) (dimensionless)
- real *8, dimension(:), allocatable **parm::prf**
Reach peak rate adjustment factor for sediment routing in the channel. Allows impact of peak flow rate on sediment routing and channel reshaping to be taken into account (none)
- real *8, dimension(:), allocatable **parm::depprch**
- real *8, dimension(:), allocatable **parm::depprfp**
- real *8, dimension(:), allocatable **parm::spcon**
linear parameter for calculating sediment reentrained in channel sediment routing
- real *8, dimension(:), allocatable **parm::spexp**
exponent parameter for calculating sediment reentrained in channel sediment routing
- real *8, dimension(:), allocatable **parm::sanst**
- real *8, dimension(:), allocatable **parm::silst**
- real *8, dimension(:), allocatable **parm::clast**
- real *8, dimension(:), allocatable **parm::sagst**
- real *8, dimension(:), allocatable **parm::lagst**
- real *8, dimension(:), allocatable **parm::pot_san**
- real *8, dimension(:), allocatable **parm::pot_sil**
- real *8, dimension(:), allocatable **parm::pot_cla**
- real *8, dimension(:), allocatable **parm::pot_sag**
- real *8, dimension(:), allocatable **parm::pot_lag**
- real *8, dimension(:), allocatable **parm::potsani**
- real *8, dimension(:), allocatable **parm::potsili**
- real *8, dimension(:), allocatable **parm::potclai**
- real *8, dimension(:), allocatable **parm::potsagi**
- real *8, dimension(:), allocatable **parm::potlagi**
- real *8, dimension(:), allocatable **parm::sanyld**
- real *8, dimension(:), allocatable **parm::silyld**
- real *8, dimension(:), allocatable **parm::clayld**
- real *8, dimension(:), allocatable **parm::sagyld**
- real *8, dimension(:), allocatable **parm::lagyld**
- real *8, dimension(:), allocatable **parm::grayld**
- real *8, dimension(:), allocatable **parm::res_san**
- real *8, dimension(:), allocatable **parm::res_sil**
- real *8, dimension(:), allocatable **parm::res_cla**
- real *8, dimension(:), allocatable **parm::res_sag**
- real *8, dimension(:), allocatable **parm::res_lag**
- real *8, dimension(:), allocatable **parm::res_gra**
- real *8, dimension(:), allocatable **parm::pnd_san**
- real *8, dimension(:), allocatable **parm::pnd_sil**
- real *8, dimension(:), allocatable **parm::pnd_cla**
- real *8, dimension(:), allocatable **parm::pnd_sag**
- real *8, dimension(:), allocatable **parm::pnd_lag**
- real *8, dimension(:), allocatable **parm::wet_san**
- real *8, dimension(:), allocatable **parm::wet_sil**
- real *8, dimension(:), allocatable **parm::wet_cla**
- real *8, dimension(:), allocatable **parm::wet_lag**
- real *8, dimension(:), allocatable **parm::wet_sag**
- real *8 **parm::ressano**

- real *8 **parm::ressilo**
- real *8 **parm::resclao**
- real *8 **parm::ressago**
- real *8 **parm::reslago**
- real *8 **parm::resgrao**
- real *8 **parm::ressani**
- real *8 **parm::ressili**
- real *8 **parm::resclai**
- real *8 **parm::ressagi**
- real *8 **parm::reslagi**
- real *8 **parm::resgrai**
- real *8 **parm::potsano**
- real *8 **parm::potsilo**
- real *8 **parm::potclao**
- real *8 **parm::potsago**
- real *8 **parm::potlago**
- real *8 **parm::pndsarin**
- real *8 **parm::pndsilin**
- real *8 **parm::pndclain**
- real *8 **parm::pndsagin**
- real *8 **parm::pndlugin**
- real *8 **parm::pndsano**
- real *8 **parm::pndsilo**
- real *8 **parm::pndclao**
- real *8 **parm::pndsago**
- real *8 **parm::pndlago**
- real *8, dimension(:), allocatable **parm::ch_di**
initial depth of main channel (m)
- real *8, dimension(:), allocatable **parm::ch_erod**
channel erodibility factor (0.0-1.0) (none)
0 non-erosive channel
1 no resistance to erosion
- real *8, dimension(:), allocatable **parm::ch_l2**
length of main channel (km)
- real *8, dimension(:), allocatable **parm::ch_cov**
- real *8, dimension(:), allocatable **parm::ch_bnk_bd**
bulk density of channel bank sediment (1.1-1.9) (g/cc)
- real *8, dimension(:), allocatable **parm::ch_bed_bd**
bulk density of channel bed sediment (1.1-1.9) (g/cc)
- real *8, dimension(:), allocatable **parm::ch_bnk_kd**
erodibility of channel bank sediment by jet test (Peter Allen needs to give more info on this)
- real *8, dimension(:), allocatable **parm::ch_bed_kd**
erodibility of channel bed sediment by jet test (Peter Allen needs to give more info on this)
- real *8, dimension(:), allocatable **parm::ch_bnk_d50**
D50(median) particle size diameter of channel bank sediment (0.001 - 20)
- real *8, dimension(:), allocatable **parm::ch_bed_d50**
D50(median) particle size diameter of channel bed sediment (micrometers) (0.001 - 20)
- real *8, dimension(:), allocatable **parm::ch_cov1**
channel erodibility factor (0.0-1.0) (none)
0 non-erosive channel
1 no resistance to erosion
- real *8, dimension(:), allocatable **parm::ch_cov2**

- channel cover factor (0.0-1.0) (none)*
0 channel is completely protected from erosion by cover
1 no vegetative cover on channel
- real *8, dimension(:), allocatable [parm::tc_bed](#)
critical shear stress of channel bed (N/m2)
- real *8, dimension(:), allocatable [parm::tc_bnk](#)
critical shear stress of channel bank (N/m2)
- integer, dimension(:), allocatable [parm::ch_eqn](#)
sediment routine methods (DAILY):
0 = original SWAT method
1 = Bagnold's
2 = Kodatie
3 = Molinas WU
4 = Yang
- real *8, dimension(:), allocatable [parm::chpst_rea](#)
pesticide reaction coefficient in reach (1/day)
- real *8, dimension(:), allocatable [parm::chpst_vol](#)
pesticide volatilization coefficient in reach (m/day)
- real *8, dimension(:), allocatable [parm::chpst_conc](#)
- real *8, dimension(:), allocatable [parm::chpst_koc](#)
pesticide partition coefficient between water and sediment in reach (m^3/g)
- real *8, dimension(:), allocatable [parm::chpst_rsp](#)
resuspension velocity in reach for pesticide sorbed to sediment (m/day)
- real *8, dimension(:), allocatable [parm::chpst_stl](#)
settling velocity in reach for pesticide sorbed to sediment (m/day)
- real *8, dimension(:), allocatable [parm::ch_wdr](#)
channel width to depth ratio (m/m)
- real *8, dimension(:), allocatable [parm::chpst_mix](#)
mixing velocity (diffusion/dispersion) for pesticide in reach (m/day)
- real *8, dimension(:), allocatable [parm::sedpst_conc](#)
initil pesticide concentration in river bed sediment (mg/m^3)
- real *8, dimension(:), allocatable [parm::sedpst_bry](#)
pesticide burial velocity in river bed sediment (m/day)
- real *8, dimension(:), allocatable [parm::sedpst_rea](#)
pesticide reaction coefficient in river bed sediment (1/day)
- real *8, dimension(:), allocatable [parm::sedpst_act](#)
depth of active sediment layer in reach for pesticide (m)
- real *8, dimension(:), allocatable [parm::rch_cbod](#)
- real *8, dimension(:), allocatable [parm::rch_bactlp](#)
- real *8, dimension(:), allocatable [parm::chside](#)
change in horizontal distance per unit vertical distance (0.0 - 5)
0 = for vertical channel bank
5 = for channel bank with gentl side slope
- real *8, dimension(:), allocatable [parm::rs1](#)
local algal settling rate in reach at 20 deg C (m/day or m/hour)
- real *8, dimension(:), allocatable [parm::rs2](#)
*benthos source rate for dissolved phosphorus in reach at 20 deg C ((mg disP-P)/(m²*day) or (mg disP-P)/(m²*hour))*
- real *8, dimension(:), allocatable [parm::rs3](#)
*benthos source rate for ammonia nitrogen in reach at 20 deg C ((mg NH4-N)/(m²*day) or (mg NH4-N)/(m²*hour))*
- real *8, dimension(:), allocatable [parm::rs4](#)
rate coefficient for organic nitrogen settling in reach at 20 deg C (1/day or 1/hour)
- real *8, dimension(:), allocatable [parm::rs5](#)

- organic phosphorus settling rate in reach at 20 deg C (1/day or 1/hour)*
- real *8, dimension(:), allocatable **parm::rk1**
- CBOD deoxygenation rate coefficient in reach at 20 deg C (1/day or 1/hour)*
- real *8, dimension(:), allocatable **parm::rk2**
- reaeration rate in accordance with Fickian diffusion in reach at 20 deg C (1/day or 1/hour)*
- real *8, dimension(:), allocatable **parm::rk3**
- rate of loss of CBOD due to settling in reach at 20 deg C (1/day or 1/hour)*
- real *8, dimension(:), allocatable **parm::rk4**
- sediment oxygen demand rate in reach at 20 deg C (mg O2/(m²*day) or mg O2/(m²*hour))*
- real *8, dimension(:), allocatable **parm::rk5**
- coliform die-off rate in reach (1/day)*
- real *8, dimension(:), allocatable **parm::rs6**
- rate coefficient for settling of arbitrary non-conservative constituent in reach (1/day)*
- real *8, dimension(:), allocatable **parm::rs7**
- benthic source rate for arbitrary non-conservative constituent in reach ((mg ANC)/(m²*day))*
- real *8, dimension(:), allocatable **parm::bc1**
- rate constant for biological oxidation of NH3 to NO2 in reach at 20 deg C (1/day or 1/hour)*
- real *8, dimension(:), allocatable **parm::bc2**
- rate constant for biological oxidation of NO2 to NO3 in reach at 20 deg C (1/day or 1/hour)*
- real *8, dimension(:), allocatable **parm::bc3**
- rate constant for hydrolysis of organic N to ammonia in reach at 20 deg C (1/day or 1/hour)*
- real *8, dimension(:), allocatable **parm::bc4**
- rate constant for the decay of organic P to dissolved P in reach at 20 deg C (1/day or 1/hour)*
- real *8, dimension(:), allocatable **parm::rk6**
- decay rate for arbitrary non-conservative constituent in reach (1/day)*
- real *8, dimension(:), allocatable **parm::ammonian**
- real *8, dimension(:), allocatable **parm::orig_sedpstconc**
- real *8, dimension(:,:), allocatable **parm::wurch**
- average daily water removal from the reach for the month (10⁴ m³/day)*
- integer, dimension(:), allocatable **parm::icanal**
- integer, dimension(:), allocatable **parm::itb**
- real *8, dimension(:), allocatable **parm::ch_revap**
- revap coeff: this variable controls the amount of water moving from bank storage to the root zone as a result of soil moisture depletion(none)*
- real *8, dimension(:), allocatable **parm::dep_chan**
- real *8, dimension(:), allocatable **parm::harg_petco**
- coefficient related to radiation used in hargreaves eq (range: 0.0019 - 0.0032)*
- real *8, dimension(:), allocatable **parm::subfr_nowtr**
- real *8, dimension(:), allocatable **parm::cncoef_sub**
- soil water depletion coefficient used in the new (modified curve number method) same as soil index coeff used in APEX range: 0.5 - 2.0*
- real *8, dimension(:), allocatable **parm::dr_sub**
- real *8, dimension(:), allocatable **parm::sub_fr**
- fraction of total watershed area contained in subbasin (km2/km2)*
- real *8, dimension(:), allocatable **parm::sub_sw**
- water in soil profile in subbasin (mm H2O)*
- real *8, dimension(:), allocatable **parm::wcklsp**
- real *8, dimension(:), allocatable **parm::sub_minp**
- real *8, dimension(:), allocatable **parm::sub_sumfc**
- real *8, dimension(:), allocatable **parm::sub_gwno3**
- real *8, dimension(:), allocatable **parm::sub_gwsolp**
- real *8, dimension(:), allocatable **parm::co2**

- *CO2 concentration (ppmv)*
- real *8, dimension(:), allocatable [parm::sub_km](#)
area of subbasin in square kilometers (km²)
- real *8, dimension(:), allocatable [parm::wlat](#)
latitude of weather station used to compile data (degrees)
- real *8, dimension(:), allocatable [parm::sub_tc](#)
time of concentration for subbasin (hour)
- real *8, dimension(:), allocatable [parm::sub_pet](#)
- real *8, dimension(:), allocatable [parm::welev](#)
elevation of weather station used to compile weather generator data (m)
- real *8, dimension(:), allocatable [parm::sub_orgn](#)
- real *8, dimension(:), allocatable [parm::sub_orgp](#)
- real *8, dimension(:), allocatable [parm::sub_bd](#)
- real *8, dimension(:), allocatable [parm::sub_wtmp](#)
- real *8, dimension(:), allocatable [parm::sub_sedpa](#)
- real *8, dimension(:), allocatable [parm::sub_sedps](#)
- real *8, dimension(:), allocatable [parm::daylmn](#)
shortest daylength occurring during the year (hour)
- real *8, dimension(:), allocatable [parm::sub_minpa](#)
- real *8, dimension(:), allocatable [parm::sub_minps](#)
- real *8, dimension(:), allocatable [parm::latcos](#)
cos(latitude) (none)
- real *8, dimension(:), allocatable [parm::latsin](#)
sin(latitude) (none)
- real *8, dimension(:), allocatable [parm::phutot](#)
total potential heat units for year (used when no crop is growing) (heat unit)
- real *8, dimension(:), allocatable [parm::plaps](#)
precipitation lapse rate: precipitation change due to change in elevation (mm H2O/km)
- real *8, dimension(:), allocatable [parm::tlaps](#)
temperature lapse rate: temperature change due to change in elevation (deg C/km)
- real *8, dimension(:), allocatable [parm::tmp_an](#)
average annual air temperature (deg C)
- real *8, dimension(:), allocatable [parm::sub_precip](#)
amount of water reaching soil surface in subbasin (mm H2O)
- real *8, dimension(:), allocatable [parm::rammo_sub](#)
atmospheric deposition of ammonium values for entire watershed (mg/l)
- real *8, dimension(:), allocatable [parm::rcn_sub](#)
atmospheric deposition of nitrate for entire watershed (mg/l)
- real *8, dimension(:), allocatable [parm::pcpdays](#)
- real *8, dimension(:), allocatable [parm::atmo_day](#)
- real *8, dimension(:), allocatable [parm::sub_snom](#)
amount of snow melt in subbasin on day (mm H2O)
- real *8, dimension(:), allocatable [parm::sub_qd](#)
surface runoff that reaches main channel during day in subbasin (mm H2O)
- real *8, dimension(:), allocatable [parm::sub_sedy](#)
- real *8, dimension(:), allocatable [parm::sub_tran](#)
transmission losses on day in subbasin (mm H2O)
- real *8, dimension(:), allocatable [parm::sub_no3](#)
- real *8, dimension(:), allocatable [parm::sub_latno3](#)
- real *8, dimension(:, :), allocatable [parm::sub_sftmp](#)
snowfall temperature for subbasin(:). Mean air temperature at which precip is equally likely to be rain as snow/freezing rain (range: -5.0/5.0) (deg C)

- real *8, dimension(:,:), allocatable [parm::sub_smtmp](#)
snow melt base temperature for subbasin(:) mean air temperature at which snow melt will occur (range: -5.0/5.0) (deg C)
- real *8, dimension(:,:), allocatable [parm::sub_timp](#)
snow pack temperature lag factor (0-1) (none)
1 = no lag (snow pack temp=current day air temp) as the lag factor goes to zero, the snow pack's temperature will be less influenced by the current day's air temperature
- real *8, dimension(:), allocatable **parm::sub_tileno3**
- real *8, dimension(:), allocatable **parm::sub_solp**
- real *8, dimension(:), allocatable **parm::sub_subp**
- real *8, dimension(:), allocatable **parm::sub_etday**
- real *8, dimension(:), allocatable [parm::sub_elev](#)
average elevation of HRU (m)
- real *8, dimension(:), allocatable **parm::sub_wyld**
- real *8, dimension(:), allocatable **parm::sub_surfq**
- real *8, dimension(:), allocatable **parm::qird**
- real *8, dimension(:), allocatable **parm::sub_gwq**
- real *8, dimension(:), allocatable **parm::sub_sep**
- real *8, dimension(:), allocatable **parm::sub_chl**
- real *8, dimension(:), allocatable **parm::sub_cbod**
- real *8, dimension(:), allocatable **parm::sub_dox**
- real *8, dimension(:), allocatable **parm::sub_solpst**
- real *8, dimension(:), allocatable **parm::sub_sorpst**
- real *8, dimension(:), allocatable **parm::sub_yorgn**
- real *8, dimension(:), allocatable **parm::sub_yorgp**
- real *8, dimension(:), allocatable [parm::sub_lat](#)
latitude of HRU/subbasin (degrees)
- real *8, dimension(:), allocatable **parm::sub_bactp**
- real *8, dimension(:), allocatable **parm::sub_bactlp**
- real *8, dimension(:), allocatable **parm::sub_latq**
- real *8, dimension(:), allocatable **parm::sub_gwq_d**
- real *8, dimension(:), allocatable **parm::sub_tileq**
- real *8, dimension(:), allocatable **parm::sub_vaptile**
- real *8, dimension(:), allocatable **parm::sub_dsan**
- real *8, dimension(:), allocatable **parm::sub_dsil**
- real *8, dimension(:), allocatable **parm::sub_dcla**
- real *8, dimension(:), allocatable **parm::sub_dsag**
- real *8, dimension(:), allocatable **parm::sub_dlag**
- real *8 **parm::vap_tile**
- real *8, dimension(:), allocatable **parm::wnan**
- real *8, dimension(:,:), allocatable **parm::sol_stpwt**
- real *8, dimension(:,:), allocatable **parm::sub_pst**
- real *8, dimension(:,:), allocatable **parm::sub_hhqd**
- real *8, dimension(:,:), allocatable **parm::sub_hhwtmp**
- real *8, dimension(:,:), allocatable [parm::huminc](#)
monthly humidity adjustment. Daily values for relative humidity within the month are rasied or lowered by the specified amount (used in climate change studies) (none)
- real *8, dimension(:,:), allocatable [parm::radinc](#)
monthly solar radiation adjustment. Daily radiation within the month is raised or lowered by the specified amount (used in climate change studies) (MJ/m²)
- real *8, dimension(:,:), allocatable [parm::rfinc](#)
monthly rainfall adjustment. Daily rainfall within the month is adjusted to the specified percentage of the original value (used in climate change studies)(%)
- real *8, dimension(:,:), allocatable [parm::tmpinc](#)

- monthly temperature adjustment. Daily maximum and minimum temperatures within the month are raised or lowered by the specified amount (used in climate change studies) (deg C)*
- real *8, dimension(:), allocatable [parm::ch_k1](#)
effective hydraulic conductivity of tributary channel alluvium (mm/hr)
 - real *8, dimension(:), allocatable [parm::ch_k2](#)
effective hydraulic conductivity of main channel alluvium (mm/hr)
 - real *8, dimension(:,:), allocatable [parm::elevb](#)
elevation at the center of the band (m)
 - real *8, dimension(:,:), allocatable [parm::elevb_fr](#)
fraction of subbasin area within elevation band (the same fractions should be listed for all HRUs within the subbasin) (none)
 - real *8, dimension(:,:), allocatable [parm::wndav](#)
average wind speed for the month (m/s)
 - real *8, dimension(:), allocatable [parm::ch_n1](#)
Manning's "n" value for the tributary channels (none)
 - real *8, dimension(:), allocatable [parm::ch_n2](#)
Manning's "n" value for the main channel (none)
 - real *8, dimension(:), allocatable [parm::ch_s1](#)
average slope of tributary channels (m/m)
 - real *8, dimension(:), allocatable [parm::ch_s2](#)
average slope of main channel (m/m)
 - real *8, dimension(:), allocatable [parm::ch_w1](#)
average width of tributary channels (m)
 - real *8, dimension(:), allocatable [parm::ch_w2](#)
average width of main channel (m)
 - real *8, dimension(:,:), allocatable [parm::dewpt](#)
average dew point temperature for the month (deg C)
 - real *8, dimension(:,:), allocatable [parm::amp_r](#)
average fraction of total daily rainfall occurring in maximum half-hour period for month (none)
 - real *8, dimension(:,:), allocatable [parm::solarav](#)
average daily solar radiation for the month (MJ/m²/day)
 - real *8, dimension(:,:), allocatable [parm::tmpstdmx](#)
standard deviation for avg monthly maximum air temperature (deg C)
 - real *8, dimension(:,:), allocatable [parm::pcf](#)
normalization coefficient for precipitation generated from skewed distribution (none)
 - real *8, dimension(:,:), allocatable [parm::tmpmn](#)
avg monthly minimum air temperature (deg C)
 - real *8, dimension(:,:), allocatable [parm::tmpmx](#)
avg monthly maximum air temperature (deg C)
 - real *8, dimension(:,:), allocatable [parm::tmpstdmn](#)
standard deviation for avg monthly minimum air temperature (deg C)
 - real *8, dimension(:,:), allocatable [parm::otmpstdmn](#)
 - real *8, dimension(:,:), allocatable [parm::otmpmn](#)
 - real *8, dimension(:,:), allocatable [parm::otmpmx](#)
 - real *8, dimension(:,:), allocatable [parm::otmpstdmx](#)
 - real *8, dimension(:,:), allocatable [parm::ch_erodmo](#)
 - real *8, dimension(:,:), allocatable [parm::uh](#)
 - real *8, dimension(:,:), allocatable [parm::hqdsave](#)
 - real *8, dimension(:,:), allocatable [parm::hsdsave](#)
 - real *8, dimension(:,:), allocatable [parm::pr_w1](#)
probability of wet day after dry day in month (none)
 - real *8, dimension(:,:), allocatable [parm::pr_w2](#)

- probability of wet day after wet day in month (none)*
- real *8, dimension(:,:), allocatable [parm::pr_w3](#)
- proportion of wet days in the month (none)*
- real *8, dimension(:,:), allocatable [parm::pcp_stat](#)
- real *8, dimension(:,:), allocatable [parm::opr_w1](#)
- real *8, dimension(:,:), allocatable [parm::opr_w2](#)
- real *8, dimension(:,:), allocatable [parm::opr_w3](#)
- real *8, dimension(:,:), allocatable [parm::opcp_stat](#)
- integer, dimension(:), allocatable [parm::ireg](#)
- precipitation category (none):*
 - 1 precipitation <= 508 mm/yr*
 - 2 precipitation > 508 and <= 1016 mm/yr*
 - 3 precipitation > 1016 mm/yr*
- integer, dimension(:), allocatable [parm::hrutot](#)
- number of HRUs in subbasin (none)*
- integer, dimension(:), allocatable [parm::hru1](#)
- integer, dimension(:), allocatable [parm::ihgage](#)
- HRU relative humidity data code (gage # for relative humidity data used in as HRU) (none)*
- integer, dimension(:), allocatable [parm::isgage](#)
- HRU solar radiation data code (record # for solar radiation used in HRU) (none)*
- integer, dimension(:), allocatable [parm::iwgage](#)
- HRU wind speed gage data code (gage # for wind speed data used in HRU) (none)*
- integer, dimension(:), allocatable [parm::subgis](#)
- GIS code printed to output files (output.sub, .rch) (none)*
- integer, dimension(:), allocatable [parm::irgage](#)
- subbasin rain gage data code (gage # for rainfall data used in HRU) (none)*
- integer, dimension(:), allocatable [parm::itgage](#)
- subbasin temp gage data code (gage # for temperature data used in HRU) (none)*
- integer, dimension(:), allocatable [parm::irelh](#)
- (none) irelh = 0 (dewpoint)*
 - irelh = 1 (relative humidity)*
 - note: inputs > 1.0 (dewpoint)*
 - inputs < 1.0 (relative hum)*
- integer, dimension(:), allocatable [parm::fcst_reg](#)
- real *8, dimension(:,:), allocatable [parm::sol_aorgn](#)
- amount of nitrogen stored in the active organic (humic) nitrogen pool (kg N/ha)*
- real *8, dimension(:,:), allocatable [parm::sol_fon](#)
- amount of nitrogen stored in the fresh organic (residue) pool (kg N/ha)*
- real *8, dimension(:,:), allocatable [parm::sol_tmp](#)
- average temperature of soil layer on previous day or*
 - daily average temperature of soil layer (deg C)*
- real *8, dimension(:,:), allocatable [parm::sol_awc](#)
- available water capacity of soil layer (mm H2O/mm soil)*
- real *8, dimension(:,:), allocatable [parm::volcr](#)
- crack volume for soil layer (mm)*
- real *8, dimension(:,:), allocatable [parm::sol_prk](#)
- percolation storage array (mm H2O)*
- real *8, dimension(:,:), allocatable [parm::pperco_sub](#)
- subbasin phosphorus percolation coefficient. Ratio of soluble phosphorus in surface to soluble phosphorus in percolate*
- real *8, dimension(:,:), allocatable [parm::sol_stap](#)
- amount of phosphorus in the soil layer stored in the stable mineral phosphorus pool (kg P/ha)*
- real *8, dimension(:,:), allocatable [parm::conv_wt](#)

- factor which converts kg/kg soil to kg/ha (none)*

 - real *8, dimension(:,:), allocatable [parm::sol_actp](#)
- amount of phosphorus stored in the active mineral phosphorus pool (kg P/ha)*

 - real *8, dimension(:,:), allocatable [parm::sol_solp](#)
- soluble P concentration in top soil layer (mg P/kg soil) or
amount of inorganic phosphorus stored in solution. NOTE UNIT CHANGE! (kg P/ha)*

 - real *8, dimension(:,:), allocatable [parm::crdep](#)
- maximum or potential crack volume (mm)*

 - real *8, dimension(:,:), allocatable [parm::sol_fc](#)
- amount of water available to plants in soil layer at field capacity (fc - wp) (mm H2O)*

 - real *8, dimension(:,:), allocatable [parm::sol_ul](#)
- amount of water held in the soil layer at saturation (sat - wp water) (mm H2O)*

 - real *8, dimension(:,:), allocatable [parm::sol_bd](#)
- bulk density of the soil layer in HRU (Mg/m³)*

 - real *8, dimension(:,:), allocatable [parm::sol_z](#)
- depth to bottom of soil layer (mm)*

 - real *8, dimension(:,:), allocatable [parm::sol_st](#)
- amount of water stored in the soil layer on any given day (less wp water) (mm H2O)*

 - real *8, dimension(:,:), allocatable [parm::sol_up](#)
- water content of soil at -0.033 MPa (field capacity) (mm H2O/mm soil)*

 - real *8, dimension(:,:), allocatable [parm::sol_clay](#)
- percent clay content in soil material (UNIT CHANGE!) (% or none)*

 - real *8, dimension(:,:), allocatable [parm::sol_hk](#)
- beta coefficient to calculate hydraulic conductivity (none)*

 - real *8, dimension(:,:), allocatable [parm::flat](#)
- lateral flow storage array (mm H2O)*

 - real *8, dimension(:,:), allocatable [parm::sol_nh3](#)
- amount of nitrogen stored in the ammonium pool in soil layer (kg N/ha)*

 - real *8, dimension(:,:), allocatable [parm::sol_ec](#)
- electrical conductivity of soil layer (dS/m)*

 - real *8, dimension(:,:), allocatable [parm::sol_orgn](#)
- amount of nitrogen stored in the stable organic N pool. NOTE UNIT CHANGE! (mg N/kg soil or kg N/ha)*

 - real *8, dimension(:,:), allocatable [parm::sol_por](#)
- total porosity of soil layer expressed as a fraction of the total volume (none)*

 - real *8, dimension(:,:), allocatable [parm::sol_wp](#)
- water content of soil at -1.5 MPa (wilting point) (mm H2O/mm soil)*

 - real *8, dimension(:,:), allocatable [parm::sol_orgp](#)
- amount of phosphorus stored in the organic P pool. NOTE UNIT CHANGE! (mg P/kg soil or kg P/ha)*

 - real *8, dimension(:,:), allocatable [parm::sol_hum](#)
- amount of organic matter in the soil layer classified as humic substances (kg humus/ha)*

 - real *8, dimension(:,:), allocatable [parm::sol_wpmm](#)
- water content of soil at -1.5 MPa (wilting point) (mm H2O)*

 - real *8, dimension(:,:), allocatable [parm::sol_no3](#)
- amount of nitrogen stored in the nitrate pool. This variable is read in as a concentration and converted to kg/ha (this value is read from the .sol file in units of mg/kg) (kg N/ha)*

 - real *8, dimension(:,:), allocatable [parm::sol_cbn](#)
- percent organic carbon in soil layer (%)*

 - real *8, dimension(:,:), allocatable [parm::sol_k](#)
- saturated hydraulic conductivity of soil layer (mm/hour)*

 - real *8, dimension(:,:), allocatable [parm::sol_rsd](#)
- amount of organic matter in the soil layer classified as residue (kg/ha)*

- real *8, dimension(:,:), allocatable [parm::sol_fop](#)
amount of phosphorus stored in the fresh organic (residue) pool (kg P/ha)
- real *8, dimension(:,:), allocatable [parm::sol_rock](#)
percent of rock fragments in soil layer (%)
- real *8, dimension(:,:), allocatable [parm::sol_silt](#)
percent silt content in soil material (UNIT CHANGE!) (% or none)
- real *8, dimension(:,:), allocatable [parm::sol_sand](#)
percent sand content of soil material (%)
- real *8, dimension(:,:), allocatable [parm::orig_solno3](#)
- real *8, dimension(:,:), allocatable [parm::orig_solorgn](#)
- real *8, dimension(:,:), allocatable [parm::orig_solsofp](#)
- real *8, dimension(:,:), allocatable [parm::orig_solorgp](#)
- real *8, dimension(:,:), allocatable [parm::orig_soltmp](#)
- real *8, dimension(:,:), allocatable [parm::orig_solrsd](#)
- real *8, dimension(:,:), allocatable [parm::orig_solfop](#)
- real *8, dimension(:,:), allocatable [parm::orig_solfon](#)
- real *8, dimension(:,:), allocatable [parm::orig_solaorgn](#)
- real *8, dimension(:,:), allocatable [parm::orig_solst](#)
- real *8, dimension(:,:), allocatable [parm::orig_solactp](#)
- real *8, dimension(:,:), allocatable [parm::orig_solstap](#)
- real *8, dimension(:,:), allocatable [parm::orig_volcr](#)
- real *8, dimension(:,:), allocatable [parm::conk](#)
- real *8, dimension(:,:), allocatable [parm::sol_pst](#)
sol_pst(:,1) initial amount of pesticide in first layer read in from .chm file (mg/kg)
sol_pst(:,2) amount of pesticide in layer. NOTE UNIT CHANGE! (kg/ha)
- real *8, dimension(:,:), allocatable [parm::sol_kp](#)
pesticide sorption coefficient, Kp; the ratio of the concentration in the solid phase to the concentration in solution ((mg/kg)/(mg/L))
- real *8, dimension(:,:), allocatable [parm::orig_solpst](#)
- real *8, dimension(:), allocatable [parm::velsetlr](#)
- real *8, dimension(:), allocatable [parm::velsetlp](#)
- real *8, dimension(:), allocatable [parm::br1](#)
1st shape parameter for reservoir surface area equation (none)
- real *8, dimension(:), allocatable [parm::evrsv](#)
lake evaporation coefficient (none)
- real *8, dimension(:), allocatable [parm::res_k](#)
hydraulic conductivity of the reservoir bottom (mm/hr)
- real *8, dimension(:), allocatable [parm::lkpst_conc](#)
pesticide concentration in lake water (mg/m³)
- real *8, dimension(:), allocatable [parm::res_evol](#)
volume of water needed to fill the reservoir to the emergency spillway (read in as 10⁴ m³ and converted to m³) (m³)
- real *8, dimension(:), allocatable [parm::res_pvol](#)
volume of water needed to fill the reservoir to the principal spillway (read in as 10⁴ m³ and converted to m³) (m³)
- real *8, dimension(:), allocatable [parm::res_vol](#)
reservoir volume (read in as 10⁴ m³ and converted to m³) (m³)
- real *8, dimension(:), allocatable [parm::res_psa](#)
reservoir surface area when reservoir is filled to principal spillway (ha)
- real *8, dimension(:), allocatable [parm::lkpst_rea](#)
pesticide reaction coefficient in lake water (1/day)
- real *8, dimension(:), allocatable [parm::lkpst_vol](#)
pesticide volatilization coefficient in lake water (m/day)

- real *8, dimension(:), allocatable [parm::br2](#)
2nd shape parameter for reservoir surface area equation (none)
- real *8, dimension(:), allocatable [parm::res_rr](#)
average daily principal spillway release volume (read in as a release rate in m^3/s and converted to m^3/day)
- real *8, dimension(:), allocatable [parm::res_sed](#)
amount of sediment in reservoir (read in as mg/L and converted to kg/L) (kg/L)
- real *8, dimension(:), allocatable [parm::lkpst_koc](#)
pesticide partition coefficient between water and sediment in lake water (m^3/g)
- real *8, dimension(:), allocatable [parm::lkpst_mix](#)
mixing velocity (diffusion/dispersion) in lake water for pesticide (m/day)
- real *8, dimension(:), allocatable [parm::lkpst_rsp](#)
resuspension velocity in lake water for pesticide sorbed to sediment (m/day)
- real *8, dimension(:), allocatable [parm::lkpst_stl](#)
settling velocity in lake water for pesticide sorbed to sediment (m/day)
- real *8, dimension(:), allocatable [parm::lkspst_conc](#)
pesticide concentration in lake bed sediment (mg/m^3)
- real *8, dimension(:), allocatable [parm::lkspst_rea](#)
pesticide reaction coefficient in lake bed sediment ($1/day$)
- real *8, dimension(:), allocatable [parm::theta_n](#)
- real *8, dimension(:), allocatable [parm::theta_p](#)
- real *8, dimension(:), allocatable [parm::con_nirr](#)
- real *8, dimension(:), allocatable [parm::con_pirr](#)
- real *8, dimension(:), allocatable [parm::lkspst_act](#)
depth of active sediment layer in lake for for pesticide (m)
- real *8, dimension(:), allocatable [parm::lkspst_bry](#)
pesticide burial velocity in lake bed sediment (m/day)
- real *8, dimension(:), allocatable [parm::sed_stlr](#)
- real *8, dimension(7) [parm::resdata](#)
- real *8, dimension(:), allocatable [parm::res_nsed](#)
normal amount of sediment in reservoir (read in as mg/L and convert to kg/L) (kg/L)
- real *8, dimension(:), allocatable [parm::wurtnf](#)
fraction of water removed from the reservoir via WURES which is returned and becomes flow from the reservoir outlet (none)
- real *8, dimension(:), allocatable [parm::chlar](#)
chlorophyll-a production coefficient for reservoir (none)
- real *8, dimension(:), allocatable [parm::res_no3](#)
amount of nitrate in reservoir ($kg N$)
- real *8, dimension(:), allocatable [parm::res_orgn](#)
amount of organic N in reservoir ($kg N$)
- real *8, dimension(:), allocatable [parm::res_orgp](#)
amount of organic P in reservoir ($kg P$)
- real *8, dimension(:), allocatable [parm::res_solp](#)
amount of soluble P in reservoir ($kg P$)
- real *8, dimension(:), allocatable [parm::res_seci](#)
secchi-disk depth (m)
- real *8, dimension(:), allocatable [parm::res_chla](#)
- real *8, dimension(:), allocatable [parm::res_esa](#)
reservoir surface area when reservoir is filled to emergency spillway (ha)
- real *8, dimension(:), allocatable [parm::res_nh3](#)
amount of ammonia in reservoir ($kg N$)
- real *8, dimension(:), allocatable [parm::res_no2](#)

- amount of nitrite in reservoir (kg N)*
- real *8, dimension(:), allocatable [parm::seccir](#)
- water clarity coefficient for reservoir (none)*
- real *8, dimension(:), allocatable [parm::res_bactp](#)
- real *8, dimension(:), allocatable [parm::res_bactlp](#)
- real *8, dimension(:), allocatable [parm::oflowmn_fps](#)
- minimum reservoir outflow as a fraction of the principal spillway volume (0-1) (fraction)*
- real *8, dimension(:), allocatable [parm::starg_fps](#)
- target volume as a fraction of the principal spillway volume (.1-5) (fraction)*
- real *8, dimension(:), allocatable [parm::weirc](#)
- real *8, dimension(:), allocatable [parm::weirk](#)
- real *8, dimension(:), allocatable [parm::weirw](#)
- real *8, dimension(:), allocatable [parm::acoef](#)
- real *8, dimension(:), allocatable [parm::bcoef](#)
- real *8, dimension(:), allocatable [parm::ccoef](#)
- real *8, dimension(:), allocatable [parm::orig_resvol](#)
- real *8, dimension(:), allocatable [parm::orig_ressed](#)
- real *8, dimension(:), allocatable [parm::orig_lkpstconc](#)
- real *8, dimension(:), allocatable [parm::orig_lkpststconc](#)
- real *8, dimension(:), allocatable [parm::orig_ressolp](#)
- real *8, dimension(:), allocatable [parm::orig_resorgp](#)
- real *8, dimension(:), allocatable [parm::orig_resno3](#)
- real *8, dimension(:), allocatable [parm::orig_resno2](#)
- real *8, dimension(:), allocatable [parm::orig_resnh3](#)
- real *8, dimension(:), allocatable [parm::orig_resorgn](#)
- real *8, dimension(:,:), allocatable [parm::oflowmn](#)
- minimum daily outflow for the month (read in as m^3/s and converted to m^3/day) (m^3/day)*
- real *8, dimension(:,:), allocatable [parm::oflowmx](#)
- maximum daily outflow for the month (read in as m^3/s and converted to m^3/day) (m^3/day)*
- real *8, dimension(:,:), allocatable [parm::starg](#)
- monthly target reservoir storage (needed if IRESCO=2) (read in as $10^4 m^3$ and converted to m^3) (m^3)*
- real *8, dimension(:), allocatable [parm::psetlr1](#)
- phosphorus settling rate for mid-year period (read in as $m/year$ and converted to m/day) (m/day)*
- real *8, dimension(:), allocatable [parm::psetlr2](#)
- phosphorus settling rate for remainder of year (read in as $m/year$ and converted to m/day) (m/day)*
- real *8, dimension(:), allocatable [parm::nsetlr1](#)
- nitrogen settling rate for mid-year period (read in as $m/year$ and converted to m/day) (m/day)*
- real *8, dimension(:), allocatable [parm::nsetlr2](#)
- nitrogen settling rate for remainder of year (read in as $m/year$ and converted to m/day) (m/day)*
- real *8, dimension(:,:), allocatable [parm::wuresn](#)
- average amount of water withdrawn from reservoir each month for consumptive water use (read in as $10^4 m^3$ and converted to m^3) (m^3)*
- real *8, dimension(:,:), allocatable [parm::res_out](#)
- measured average daily outflow from the reservoir for the month (needed if IRESCO=1) (read in as m^3/s and converted to m^3/day) (m^3/day)*
- integer, dimension(:), allocatable [parm::res_sub](#)
- number of subbasin reservoir is in (weather for the subbasin is used for the reservoir) (none)*
- integer, dimension(:), allocatable [parm::ires1](#)
- beginning of mid-year nutrient settling "season" (none)*
- integer, dimension(:), allocatable [parm::ires2](#)
- end of mid-year nutrient settling "season" (none)*
- integer, dimension(:), allocatable [parm::iresco](#)

- outflow simulation code (none):
 0 compute outflow for uncontrolled reservoir with average annual release rate
 1 measured monthly outflow
 2 simulated controlled outflow-target release
 3 measured daily outflow
 4 stage/volume/outflow relationship
- integer, dimension(:), allocatable **parm::iyres**
 year of the simulation that the reservoir becomes operational (none)
 - integer, dimension(:), allocatable **parm::mores**
 month the reservoir becomes operational (none)
 - integer, dimension(:), allocatable **parm::iflod1r**
 beginning month of non-flood season (needed if IRESCO=2) (none)
 - integer, dimension(:), allocatable **parm::iflod2r**
 ending month of non-flood season (needed if IRESCO=2) (none)
 - integer, dimension(:), allocatable **parm::ndtargr**
 number of days to reach target storage from current reservoir storage (needed if IRESCO=2) (days)
 - real *8, dimension(:), allocatable **parm::ap_ef**
 application efficiency (0-1) (none)
 - real *8, dimension(:), allocatable **parm::decay_f**
 exponential of the rate constant for degradation of the pesticide on foliage (none)
 - real *8, dimension(:), allocatable **parm::skoc**
 soil adsorption coefficient normalized for soil organic carbon content ((mg/kg)/(mg/L))
 - real *8, dimension(:), allocatable **parm::decay_s**
 exponential of the rate constant for degradation of the pesticide in soil (none)
 - real *8, dimension(:), allocatable **parm::hlife_f**
 half-life of pesticide on foliage (days)
 - real *8, dimension(:), allocatable **parm::hlife_s**
 half-life of pesticide in soil (days)
 - real *8, dimension(:), allocatable **parm::pst_wof**
 fraction of pesticide on foliage which is washed-off by a rainfall event (none)
 - real *8, dimension(:), allocatable **parm::pst_wsol**
 solubility of chemical in water (mg/L (ppm))
 - real *8, dimension(:), allocatable **parm::irramt**
 depth of irrigation water applied to HRU (mm H2O)
 - real *8, dimension(:), allocatable **parm::phusw**
 - real *8, dimension(:), allocatable **parm::phusw_nocrop**
 - integer, dimension(:), allocatable **parm::pstflg**
 flag for types of pesticide used in watershed. Array location is pesticide ID number
 0: pesticide not used
 1: pesticide used
 - integer, dimension(:), allocatable **parm::nope**
 sequence number of pesticide in NPNO(:) (none)
 - integer, dimension(:), allocatable **parm::nop**
 - integer, dimension(:), allocatable **parm::yr_skip**
 - integer, dimension(:), allocatable **parm::isweep**
 - integer, dimension(:), allocatable **parm::icrmx**
 - integer, dimension(:), allocatable **parm::nopmx**
 - integer, dimension(:,:), allocatable **parm::mgtop**
 - integer, dimension(:,:), allocatable **parm::idop**
 - integer, dimension(:,:), allocatable **parm::mgt1iop**
 - integer, dimension(:,:), allocatable **parm::mgt2iop**
 - integer, dimension(:,:), allocatable **parm::mgt3iop**
 - real *8, dimension(:,:), allocatable **parm::mgt4op**

- real *8, dimension(:,:), allocatable **parm::mgt5op**
- real *8, dimension(:,:), allocatable **parm::mgt6op**
- real *8, dimension(:,:), allocatable **parm::mgt7op**
- real *8, dimension(:,:), allocatable **parm::mgt8op**
- real *8, dimension(:,:), allocatable **parm::mgt9op**
- real *8, dimension(:,:), allocatable **parm::mgt10iop**
- real *8, dimension(:,:), allocatable **parm::phu_op**
- real *8, dimension(:), allocatable **parm::cnyld**
fraction of nitrogen in yield (kg N/kg yield)
- real *8, dimension(:), allocatable **parm::rsdco_pl**
plant residue decomposition coefficient. The fraction of residue which will decompose in a day assuming optimal moisture, temperature, C:N ratio, and C:P ratio (none)
- real *8, dimension(:), allocatable **parm::wac21**
1st shape parameter for radiation use efficiency equation (none)
- real *8, dimension(:), allocatable **parm::wac22**
2nd shape parameter for radiation use efficiency equation (none)
- real *8, dimension(:), allocatable **parm::alai_min**
minimum LAI during winter dormant period (m^2/m^2)
- real *8, dimension(:), allocatable **parm::leaf1**
1st shape parameter for leaf area development equation (none)
- real *8, dimension(:), allocatable **parm::leaf2**
2nd shape parameter for leaf area development equation (none)
- real *8, dimension(:), allocatable **parm::wsyf**
Value of harvest index between 0 and HVSTI which represents the lowest value expected due to water stress ((kg/ha)/(kg/ha))
- real *8, dimension(:), allocatable **parm::bio_e**
*biomass-energy ratio. The potential (unstressed) growth rate per unit of intercepted photosynthetically active radiation. ((kg/ha)/(MJ/m**2))*
- real *8, dimension(:), allocatable **parm::hvsti**
harvest index: crop yield/aboveground biomass ((kg/ha)/(kg/ha))
- real *8, dimension(:), allocatable **parm::t_base**
minimum temperature for plant growth (deg C)
- real *8, dimension(:), allocatable **parm::t_opt**
optimal temperature for plant growth (deg C)
- real *8, dimension(:), allocatable **parm::chtmx**
maximum canopy height (m)
- real *8, dimension(:), allocatable **parm::cvm**
natural log of USLE_C (the minimum value of the USLE C factor for the land cover) (none)
- real *8, dimension(:), allocatable **parm::gsi**
maximum stomatal conductance (m/s)
- real *8, dimension(:), allocatable **parm::vpd2**
rate of decline in stomatal conductance per unit increase in vapor pressure deficit ((m/s)(1/kPa))*
- real *8, dimension(:), allocatable **parm::wavp**
rate of decline in radiation use efficiency as a function of vapor pressure deficit (none)
- real *8, dimension(:), allocatable **parm::bio_leaf**
fraction of leaf/needle biomass that drops during dormancy (for trees only) (none)
- real *8, dimension(:), allocatable **parm::blai**
maximum (potential) leaf area index (none)
- real *8, dimension(:), allocatable **parm::cpyld**
fraction of phosphorus in yield (kg P/kg yield)
- real *8, dimension(:), allocatable **parm::dlai**
fraction of growing season when leaf area declines (none)

- real *8, dimension(:), allocatable [parm::rdmx](#)
maximum root depth of plant (m)
- real *8, dimension(:), allocatable [parm::bio_n1](#)
1st shape parameter for plant N uptake equation (none)
- real *8, dimension(:), allocatable [parm::bio_n2](#)
2nd shape parameter for plant N uptake equation (none)
- real *8, dimension(:), allocatable [parm::bio_p1](#)
1st shape parameter for plant P uptake equation (none)
- real *8, dimension(:), allocatable [parm::bio_p2](#)
2st shape parameter for plant P uptake equation (none)
- real *8, dimension(:), allocatable [parm::bm_dieoff](#)
fraction above ground biomass that dies off at dormancy (fraction)
- real *8, dimension(:), allocatable [parm::bmx_trees](#)
- real *8, dimension(:), allocatable [parm::ext_coef](#)
- real *8, dimension(:), allocatable [parm::rsr1](#)
initial root to shoot ratio at the beg of growing season
- real *8, dimension(:), allocatable [parm::rsr2](#)
root to shoot ratio at the end of the growing season
- real *8, dimension(:), allocatable [parm::pltnfr1](#)
nitrogen uptake parameter #1: normal fraction of N in crop biomass at emergence (kg N/kg biomass)
- real *8, dimension(:), allocatable [parm::pltnfr2](#)
nitrogen uptake parameter #2: normal fraction of N in crop biomass at 0.5 maturity (kg N/kg biomass)
- real *8, dimension(:), allocatable [parm::pltnfr3](#)
nitrogen uptake parameter #3: normal fraction of N in crop biomass at maturity (kg N/kg biomass)
- real *8, dimension(:), allocatable [parm::pltpfr1](#)
phosphorus uptake parameter #1: normal fraction of P in crop biomass at emergence (kg P/kg biomass)
- real *8, dimension(:), allocatable [parm::pltpfr2](#)
phosphorus uptake parameter #2: normal fraction of P in crop biomass at 0.5 maturity (kg P/kg biomass)
- real *8, dimension(:), allocatable [parm::pltpfr3](#)
phosphorus uptake parameter #3: normal fraction of P in crop biomass at maturity (kg P/kg biomass)
- integer, dimension(:), allocatable [parm::idc](#)
crop/landcover category:
1 warm season annual legume
2 cold season annual legume
3 perennial legume
4 warm season annual
5 cold season annual
6 perennial
7 trees
- integer, dimension(:), allocatable [parm::mat_yrs](#)
- real *8, dimension(:), allocatable [parm::bactpdb](#)
concentration of persistent bacteria in manure (fertilizer) (cfu/g manure)
- real *8, dimension(:), allocatable [parm::fminn](#)
fraction of fertilize/manure that is mineral N (NO₃ + NH₃) (kg minN/kg fert)
- real *8, dimension(:), allocatable [parm::forgn](#)
fraction of organic N in fertilizer/manure (kg orgN/kg fert)
- real *8, dimension(:), allocatable [parm::forgp](#)
fraction of fertilizer/manure that is organic P (kg orgP/kg fert)
- real *8, dimension(:), allocatable [parm::bactkddb](#)
fraction of bacteria in solution (the remaining fraction is sorbed to soil particles) (none):
1: all bacteria in solution
0: all bacteria sorbed to soil particles

- real *8, dimension(:), allocatable [parm::bactlpdb](#)
concentration of less persistent bacteria in manure (fertilizer) (cfu/g manure)
- real *8, dimension(:), allocatable [parm::fminp](#)
fraction of fertilizer that is mineral P in fertilizer/manure (kg minP/kg fert)
- real *8, dimension(:), allocatable [parm::fnh3n](#)
fraction of mineral N in fertilizer that is NH3-N in fertilizer/manure (kgNH3-N/kgminN)
- character(len=8), dimension(200) [parm::fertnm](#)
name of fertilizer
- real *8, dimension(:), allocatable [parm::curbden](#)
curb length density in HRU (km/ha)
- real *8, dimension(:), allocatable [parm::dirtmx](#)
maximum amount of solids allowed to build up on impervious surfaces (kg/curb km)
- real *8, dimension(:), allocatable [parm::fimp](#)
fraction of HRU area that is impervious (both directly and indirectly connected)(fraction)
- real *8, dimension(:), allocatable [parm::urbcoef](#)
wash-off coefficient for removal of constituents from an impervious surface (1/mm)
- real *8, dimension(:), allocatable [parm::thalf](#)
time for the amount of solids on impervious areas to build up to 1/2 the maximum level (days)
- real *8, dimension(:), allocatable [parm::tnconc](#)
concentration of total nitrogen in suspended solid load from impervious areas (mg N/kg sed)
- real *8, dimension(:), allocatable [parm::tno3conc](#)
concentration of NO3-N in suspended solid load from impervious areas (mg NO3-N/kg sed)
- real *8, dimension(:), allocatable [parm::tpconc](#)
concentration of total phosphorus in suspended solid load from impervious areas (mg P/kg sed)
- real *8, dimension(:), allocatable [parm::fcimp](#)
fraction of HRU area that is classified as directly connected impervious (fraction)
- real *8, dimension(:), allocatable [parm::urbcn2](#)
SCS curve number for moisture condition II in impervious areas (none)
- real *8 [parm::fr_curb](#)
availability factor, the fraction of the curb length that is sweepable (none)
- real *8 [parm::frt_kg](#)
amount of fertilizer applied to HRU (kg/ha)
- real *8 [parm::pst_dep](#)
depth of pesticide in the soil (mm)
- real *8 [parm::swepeff](#)
- real *8, dimension(:), allocatable [parm::ranrns_hru](#)
- integer, dimension(:), allocatable [parm::itill](#)
- real *8, dimension(:), allocatable [parm::deptil](#)
depth of mixing caused by tillage operation (mm)
- real *8, dimension(:), allocatable [parm::effmix](#)
mixing efficiency of tillage operation (none)
- real *8, dimension(:), allocatable [parm::ranrns](#)
random roughness of a given tillage operation (mm)
- character(len=8), dimension(550) [parm::tillnm](#)
8-character name for the tillage operation
- real *8, dimension(:), allocatable [parm::rnum1s](#)
*For ICODES equal to (none)
0,1,3,5,9: not used
2: Fraction of flow in channel
4: amount of water transferred (as defined by INUM4S)
7,8,10,11: drainage area in square kilometers associated with the record file
12: reparation coefficient.*

- real *8, dimension(:), allocatable [parm::hyd_dakm](#)
total drainage area of hydrograph in square kilometers (km²)
- real *8, dimension(:,:), allocatable **parm::varoute**
- real *8, dimension(:,:), allocatable **parm::shyd**
- real *8, dimension(:,:), allocatable **parm::vartran**
- real *8, dimension(:,:), allocatable **parm::hhvaroute**
- integer, dimension(:), allocatable [parm::icodes](#)
routing command code (none):
0 = finish
1 = subbasin
2 = route
3 = routres
4 = transfer
5 = add
6 = rechour
7 = recmon
8 = recyear
9 = save
10 = recday
11 = recnst
12 = structure
13 = apex
14 = saveconc
15 =
- integer, dimension(:), allocatable [parm::ihouts](#)
For ICODES equal to (none)
0: not used
1,2,3,5,7,8,10,11: hydrograph storage location number
4: departure type (1=reach, 2=reservoir)
9: hydrograph storage location of data to be printed to event file
14: hydrograph storage location of data to be printed to saveconc file.
- integer, dimension(:), allocatable [parm::inum1s](#)
For ICODES equal to (none)
0: not used
1: subbasin number
2: reach number
3: reservoir number
4: reach or res # flow is diverted from
5: hydrograph storage location of 1st dataset to be added
7,8,9,10,11,14: file number.
- integer, dimension(:), allocatable [parm::inum2s](#)
For ICODES equal to (none)
0,1,7,8,10,11: not used
2,3: inflow hydrograph storage location
4: destination type (1=reach, 2=reservoir)
5: hydrograph storage location of 2nd dataset to be added
9,14: print frequency (0=daily, 1=hourly)
- integer, dimension(:), allocatable [parm::inum3s](#)
For ICODES equal to (none)
0,1,5,7,8,10,11: not used
2,3: subbasin number 4: destination number. Reach or reservoir receiving water
9: print format (0=normal, fixed format; 1=txt format for AV interface, recday)
- integer, dimension(:), allocatable [parm::inum4s](#)
For ICODES equal to (none)
0,2,3,5,7,8,9,10,11: not used
1: GIS code printed to output file (optional)
4: rule code governing transfer of water (1=fraction transferred out, 2=min volume or flow left, 3=exact amount transferred)
- integer, dimension(:), allocatable **parm::inum5s**

- integer, dimension(:), allocatable **parm::inum6s**
- integer, dimension(:), allocatable **parm::inum7s**
- integer, dimension(:), allocatable **parm::inum8s**
- integer, dimension(:), allocatable **parm::subed**
- character(len=10), dimension(:), allocatable **parm::recmonps**
- character(len=10), dimension(:), allocatable **parm::reccnsteps**
- character(len=5), dimension(:), allocatable **parm::subnum**
- character(len=4), dimension(:), allocatable **parm::hruno**
- real *8, dimension(:), allocatable **parm::grwat_n**
Mannings's n for grassed waterway (none)
- integer, dimension(:), allocatable **parm::grwat_i**
flag for the simulation of grass waterways (none)
= 0 inactive
= 1 active
- real *8, dimension(:), allocatable **parm::grwat_l**
length of grass waterway (km)
- real *8, dimension(:), allocatable **parm::grwat_w**
average width of grassed waterway (m)
- real *8, dimension(:), allocatable **parm::grwat_d**
depth of grassed waterway from top of bank to bottom (m)
- real *8, dimension(:), allocatable **parm::grwat_s**
average slope of grassed waterway channel (m)
- real *8, dimension(:), allocatable **parm::grwat_spon**
linear parameter for calculating sediment in grassed waterways (none)
- real *8, dimension(:), allocatable **parm::tc_gwat**
- real *8, dimension(:), allocatable **parm::pot_volmm**
- real *8, dimension(:), allocatable **parm::pot_tilemm**
- real *8, dimension(:), allocatable **parm::pot_volxmm**
- real *8, dimension(:), allocatable **parm::pot_fr**
fraction of HRU area that drains into pothole (km^2/km^2)
- real *8, dimension(:), allocatable **parm::pot_tile**
average daily outflow to main channel from tile flow if drainage tiles are installed in pothole (needed only if current HRU is IPOT) (m^3/s)
- real *8, dimension(:), allocatable **parm::pot_vol**
initial or current volume of water stored in the depression/impounded area (read in as mm and converted to m^3) (needed only if current HRU is IPOT) (mm or $\text{m}^3 \text{H}_2\text{O}$)
- real *8, dimension(:), allocatable **parm::potsa**
surface area of impounded water body (ha)
- real *8, dimension(:), allocatable **parm::pot_volx**
maximum volume of water stored in the depression/impounded area (read in as mm and converted to m^3) (needed only if current HRU is IPOT) (mm)
- real *8, dimension(:), allocatable **parm::wfsh**
wetting front matric potential (average capillary suction at wetting front) (mm)
- real *8, dimension(:), allocatable **parm::potflwi**
- real *8, dimension(:), allocatable **parm::potsedi**
- real *8, dimension(:), allocatable **parm::pot_no3l**
nitrate decay rate in impounded area (1/day)
- real *8, dimension(:), allocatable **parm::pot_nsed**
normal sediment concentration in impounded water (needed only if current HRU is IPOT)(mg/L)
- real *8, dimension(:), allocatable **parm::gwno3**
nitrate-N concentration in groundwater loading to reach (mg N/L)
- real *8, dimension(:), allocatable **parm::newrti**
infiltration rate for last time step from the previous day (mm/hr)

- real *8, dimension(:), allocatable [parm::fsred](#)
reduction in bacteria loading from filter strip (none)
- real *8, dimension(:), allocatable [parm::pot_sed](#)
- real *8, dimension(:), allocatable [parm::pot_no3](#)
- real *8, dimension(:), allocatable [parm::tmpavp](#)
- real *8, dimension(:), allocatable [parm::dis_stream](#)
average distance to stream (m)
- real *8, dimension(:), allocatable [parm::evpot](#)
pothole evaporation coefficient (none)
- real *8, dimension(:), allocatable [parm::pot_solpl](#)
- real *8, dimension(:), allocatable [parm::sed_con](#)
- real *8, dimension(:), allocatable [parm::orgn_con](#)
- real *8, dimension(:), allocatable [parm::orgp_con](#)
- real *8, dimension(:), allocatable [parm::pot_k](#)
*hydraulic conductivity of soil surface of pothole defaults to conductivity of upper soil (0.↵
01-10.) layer*
- real *8, dimension(:), allocatable [parm::soln_con](#)
- real *8, dimension(:), allocatable [parm::solp_con](#)
- real *8, dimension(:), allocatable [parm::n_reduc](#)
nitrogen uptake reduction factor (not currently used; defaulted 300.)
- real *8, dimension(:), allocatable [parm::n_lag](#)
lag coefficient for calculating nitrate concentration in subsurface drains (0.001 - 1.0) (dimensionless)
- real *8, dimension(:), allocatable [parm::n_ln](#)
power function exponent for calculating nitrate concentration in subsurface drains (1.0 - 3.0) (dimensionless)
- real *8, dimension(:), allocatable [parm::n_inco](#)
coefficient for power function for calculating nitrate concentration in subsurface drains (0.5 - 4.0) (dimensionless)
- integer, dimension(:), allocatable [parm::ioper](#)
- integer, dimension(:), allocatable [parm::ngrwat](#)
- real *8, dimension(:), allocatable [parm::usle_ls](#)
USLE equation length slope (LS) factor (none)
- real *8, dimension(:), allocatable [parm::filterw](#)
filter strip width for bacteria transport (m)
- real *8, dimension(:), allocatable [parm::phuacc](#)
fraction of plant heat units accumulated (none)
- real *8, dimension(:), allocatable [parm::sumix](#)
sum of all tillage mixing efficiencies for HRU operation (none)
- real *8, dimension(:), allocatable [parm::epco](#)
plant water uptake compensation factor (0-1) (none)
- real *8, dimension(:), allocatable [parm::esco](#)
soil evaporation compensation factor (0-1) (none)
- real *8, dimension(:), allocatable [parm::hru_slp](#)
average slope steepness (m/m)
- real *8, dimension(:), allocatable [parm::slsubbsn](#)
average slope length for subbasin (m)
- real *8, dimension(:), allocatable [parm::erorgn](#)
organic N enrichment ratio, if left blank the model will calculate for every event (none)
- real *8, dimension(:), allocatable [parm::erorgp](#)
organic P enrichment ratio, if left blank the model will calculate for every event (none)
- real *8, dimension(:), allocatable [parm::biomix](#)
biological mixing efficiency. Mixing of soil due to activity of earthworms and other soil biota. Mixing is performed at the end of every calendar year (none)
- real *8, dimension(:), allocatable [parm::pnd_seci](#)

- secchi-disk depth of pond (m)*
- real *8, dimension(:), allocatable [parm::canmx](#)
maximum canopy storage (mm H2O)
- real *8, dimension(:), allocatable [parm::divmax](#)
maximum daily irrigation diversion from the reach (when IRRSC=1 or IRR=3): when value is positive the units are mm H2O; when the value is negative, the units are (10⁴ m³ H2O) (mm H2O or 10⁴ m³ H2O)
- real *8, dimension(:), allocatable [parm::flowmin](#)
minimum instream flow for irrigation diversions when IRRSC=1, irrigation water will be diverted only when streamflow is at or above FLOWMIN (m³/s)
- real *8, dimension(:), allocatable [parm::usle_p](#)
USLE equation support practice (P) factor (none)
- real *8, dimension(:), allocatable [parm::lat_sed](#)
sediment concentration in lateral flow (g/L)
- real *8, dimension(:), allocatable [parm::rch_dakm](#)
total drainage area contributing to flow at the outlet (pour point) of the reach in square kilometers (km²)
- real *8, dimension(:), allocatable [parm::cn1](#)
SCS runoff curve number for moisture condition I (none)
- real *8, dimension(:), allocatable [parm::pnd_no3s](#)
amount of nitrate originating from lateral flow in pond at end of day (kg N)
- real *8, dimension(:), allocatable [parm::lat_time](#)
lateral flow travel time or exponential of the lateral flow travel time (days or none)
- real *8, dimension(:), allocatable [parm::cn2](#)
SCS runoff curve number for moisture condition II (none)
- real *8, dimension(:), allocatable [parm::flowfr](#)
fraction of available flow in reach that is allowed to be applied to the HRU (none)
- real *8, dimension(:), allocatable [parm::sol_zmx](#)
maximum rooting depth (mm)
- real *8, dimension(:), allocatable [parm::tile_time](#)
exponential of the tile flow travel time (none)
- real *8, dimension(:), allocatable [parm::slsoil](#)
slope length for lateral subsurface flow (m)
- real *8, dimension(:), allocatable [parm::gwminp](#)
soluble P concentration in groundwater loading to reach (mg P/L)
- real *8, dimension(:), allocatable [parm::sol_cov](#)
amount of residue on soil surface (kg/ha)
- real *8, dimension(:), allocatable [parm::sed_stl](#)
fraction of sediment remaining suspended in impoundment after settling for one day (kg/kg)
- real *8, dimension(:), allocatable [parm::ov_n](#)
Manning's "n" value for overland flow (none)
- real *8, dimension(:), allocatable [parm::pnd_no3](#)
amount of nitrate originating from surface runoff in pond at end of day (kg N)
- real *8, dimension(:), allocatable [parm::pnd_solp](#)
amount of soluble P originating from surface runoff in pond at end of day (kg P)
- real *8, dimension(:), allocatable [parm::yldanu](#)
annual yield (dry weight) in the HRU (metric tons/ha)
- real *8, dimension(:), allocatable [parm::driftco](#)
coefficient for pesticide drift directly onto stream (none)
- real *8, dimension(:), allocatable [parm::pnd_orgn](#)
amount of organic N originating from surface runoff in pond at end of day (kg N)
- real *8, dimension(:), allocatable [parm::pnd_orgp](#)
amount of organic P originating from surface runoff in pond at end of day (kg P)

- real *8, dimension(:), allocatable [parm::cn3](#)
SCS runoff curve number for moisture condition III (none)
- real *8, dimension(:), allocatable [parm::twlpnd](#)
water lost through seepage from ponds on day in HRU (mm H2O)
- real *8, dimension(:), allocatable [parm::twlwet](#)
water lost through seepage from wetlands on day in HRU (mm H2O)
- real *8, dimension(:), allocatable [parm::hru_fr](#)
fraction of subbasin area contained in HRU (km^2/km^2)
- real *8, dimension(:), allocatable [parm::sol_sumul](#)
amount of water held in soil profile at saturation (mm H2O)
- real *8, dimension(:), allocatable [parm::pnd_chla](#)
amount of chlorophyll-a in pond at end of day (kg chl_a)
- real *8, dimension(:), allocatable [parm::hru_km](#)
area of HRU in square kilometers (km^2)
- real *8, dimension(:), allocatable [parm::bio_ms](#)
land cover/crop biomass (dry weight) (kg/ha)
- real *8, dimension(:), allocatable [parm::sol_alb](#)
albedo when soil is moist (none)
- real *8, dimension(:), allocatable [parm::strsw](#)
fraction of potential plant growth achieved on the day where the reduction is caused by water stress (none)
- real *8, dimension(:), allocatable [parm::pnd_fr](#)
fraction of HRU/subbasin area that drains into ponds (none)
- real *8, dimension(:), allocatable [parm::pnd_k](#)
hydraulic conductivity through bottom of ponds (mm/hr)
- real *8, dimension(:), allocatable [parm::pnd_psa](#)
surface area of ponds when filled to principal spillway (ha)
- real *8, dimension(:), allocatable [parm::pnd_pvol](#)
runoff volume from catchment area needed to fill the ponds to the principal spillway (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)
- real *8, dimension(:), allocatable [parm::pnd_esa](#)
surface area of ponds when filled to emergency spillway (ha)
- real *8, dimension(:), allocatable [parm::pnd_evol](#)
runoff volume from catchment area needed to fill the ponds to the emergency spillway (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)
- real *8, dimension(:), allocatable [parm::pnd_vol](#)
volume of water in ponds (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)
- real *8, dimension(:), allocatable [parm::yladaa](#)
average annual yield (dry weight) in the HRU (metric tons)
- real *8, dimension(:), allocatable [parm::pnd_nsed](#)
normal sediment concentration in pond water (UNIT CHANGE!) (mg/kg or kg/kg)
- real *8, dimension(:), allocatable [parm::pnd_sed](#)
sediment concentration in pond water (UNIT CHANGE!) (mg/kg or kg/kg)
- real *8, dimension(:), allocatable [parm::strsa](#)
- real *8, dimension(:), allocatable [parm::dep_imp](#)
- real *8, dimension(:), allocatable [parm::evpnd](#)
- real *8, dimension(:), allocatable [parm::evwet](#)
- real *8, dimension(:), allocatable [parm::wet_fr](#)
fraction of HRU/subbasin area that drains into wetlands (none)
- real *8, dimension(:), allocatable [parm::wet_k](#)
hydraulic conductivity of bottom of wetlands (mm/hr)
- real *8, dimension(:), allocatable [parm::wet_nsa](#)

- surface area of wetlands in subbasin at normal water level (ha)*

 - real *8, dimension(:), allocatable [parm::wet_nvol](#)

runoff volume from catchment area needed to fill wetlands to normal water level (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)

 - integer, dimension(:), allocatable [parm::iwetgw](#)
 - integer, dimension(:), allocatable [parm::iwetile](#)
 - real *8, dimension(:), allocatable [parm::wet_mxsa](#)

surface area of wetlands at maximum water level (ha)

 - real *8, dimension(:), allocatable [parm::wet_mxvol](#)

runoff volume from catchment area needed to fill wetlands to maximum water level (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)

 - real *8, dimension(:), allocatable [parm::wet_vol](#)

volume of water in wetlands (UNIT CHANGE!) ($10^4 \text{ m}^3 \text{ H}_2\text{O}$ or $\text{m}^3 \text{ H}_2\text{O}$)

 - real *8, dimension(:), allocatable [parm::wet_nsed](#)

normal sediment concentration in wetland water (UNIT CHANGE!) (mg/kg or kg/kg)

 - real *8, dimension(:), allocatable [parm::wet_sed](#)

sediment concentration in wetland water (UNIT CHANGE!) (mg/L or kg/L)

 - real *8, dimension(:), allocatable [parm::bp1](#)

1st shape parameter for pond surface area equation (none)

 - real *8, dimension(:), allocatable [parm::bp2](#)

2nd shape parameter for the pond surface area equation (none)

 - real *8, dimension(:), allocatable [parm::sci](#)

retention coefficient for CN method based on plant ET (none)

 - real *8, dimension(:), allocatable [parm::smx](#)

retention coefficient for CN method based on soil moisture (none)

 - real *8, dimension(:), allocatable [parm::bw1](#)

1st shape parameter for the wetland surface area equation (none)

 - real *8, dimension(:), allocatable [parm::bw2](#)

2nd shape parameter for the wetland surface area equation (none)

 - real *8, dimension(:), allocatable [parm::bactpq](#)

persistent bacteria in soil solution (# cfu/m²)

 - real *8, dimension(:), allocatable [parm::cnday](#)

curve number for current day, HRU and at current soil moisture (none)

 - real *8, dimension(:), allocatable [parm::bactlp_plt](#)

less persistent bacteria on foliage (# cfu/m²)

 - real *8, dimension(:), allocatable [parm::bactp_plt](#)

persistent bacteria on foliage (# cfu/m²)

 - real *8, dimension(:), allocatable [parm::auto_eff](#)

fertilizer application efficiency calculated as the amount of N applied divided by the amount of N removed at harvest (none)

 - real *8, dimension(:), allocatable [parm::secciw](#)

water clarity coefficient for wetland (none)

 - real *8, dimension(:), allocatable [parm::sol_sw](#)

amount of water stored in soil profile on current day (mm H₂O)

 - real *8, dimension(:), allocatable [parm::bactlpq](#)

less persistent bacteria in soil solution (# cfu/m²)

 - real *8, dimension(:), allocatable [parm::chlaw](#)

chlorophyll-a production coefficient for wetland (none)

 - real *8, dimension(:), allocatable [parm::tmpav](#)

average air temperature on current day in HRU (deg C)

 - real *8, dimension(:), allocatable [parm::bactlps](#)

- less persistent bacteria attached to soil particles (# cfu/m²)*
- real *8, dimension(:), allocatable [parm::bactps](#)
- persistent bacteria attached to soil particles (# cfu/m²)*
- real *8, dimension(:), allocatable [parm::sno_hru](#)
- amount of water stored as snow in HRU on current day (mm H2O)*
- real *8, dimension(:), allocatable [parm::wet_orgn](#)
- amount of organic N originating from surface runoff in wetland at end of day (kg N)*
- real *8, dimension(:), allocatable [parm::hru_ra](#)
- solar radiation for the day in HRU (MJ/m²)*
- real *8, dimension(:), allocatable [parm::subp](#)
- precipitation for the day in HRU (mm H2O)*
- real *8, dimension(:), allocatable [parm::rsdin](#)
- initial residue cover (kg/ha)*
- real *8, dimension(:), allocatable [parm::tmn](#)
- minimum air temperature on current day in HRU (deg C)*
- real *8, dimension(:), allocatable [parm::tmx](#)
- maximum air temperature on current day in HRU (deg C)*
- real *8, dimension(:), allocatable [parm::tmp_hi](#)
- real *8, dimension(:), allocatable [parm::tmp_lo](#)
- real *8, dimension(:), allocatable [parm::usle_k](#)
- USLE equation soil erodibility (K) factor (none)*
- real *8, dimension(:), allocatable [parm::tconc](#)
- time of concentration for HRU (hour)*
- real *8, dimension(:), allocatable [parm::hru_rmx](#)
- maximum possible solar radiation for the day in HRU (MJ/m²)*
- real *8, dimension(:), allocatable [parm::rwt](#)
- fraction of total plant biomass that is in roots (none)*
- real *8, dimension(:), allocatable [parm::olai](#)
- real *8, dimension(:), allocatable [parm::usle_cfac](#)
- real *8, dimension(:), allocatable [parm::usle_eifac](#)
- real *8, dimension(:), allocatable [parm::sol_sumfc](#)
- amount of water held in soil profile at field capacity (mm H2O)*
- real *8, dimension(:), allocatable [parm::t_ov](#)
- time for flow from farthest point in subbasin to enter a channel (hour)*
- real *8, dimension(:), allocatable [parm::anano3](#)
- total amount of NO3 applied during the year in auto-fertilization (kg N/ha)*
- real *8, dimension(:), allocatable [parm::aird](#)
- amount of water applied to HRU on current day (mm H2O)*
- real *8, dimension(:), allocatable [parm::wet_orgp](#)
- amount of organic P originating from surface runoff in wetland at end of day (kg P)*
- real *8, dimension(:), allocatable [parm::sol_avpor](#)
- average porosity for entire soil profile (none)*
- real *8, dimension(:), allocatable [parm::usle_mult](#)
- product of USLE K,P,LS,exp(rock) (none)*
- real *8, dimension(:), allocatable [parm::rhd](#)
- relative humidity for the day in HRU (none)*
- real *8, dimension(:), allocatable [parm::u10](#)
- wind speed (measured at 10 meters above surface) for the day in HRU (m/s)*
- real *8, dimension(:), allocatable [parm::cht](#)
- canopy height (m)*
- real *8, dimension(:), allocatable [parm::airr](#)

- average annual amount of irrigation water applied to HRU (mm H2O)*
- real *8, dimension(:), allocatable **parm::lai_aamx**
- maximum leaf area index for the entire period of simulation in the HRU (none)*
- real *8, dimension(:), allocatable **parm::deepirr**
- amount of water removed from deep aquifer for irrigation (mm H2O)*
- real *8, dimension(:), allocatable **parm::shallirr**
- amount of water removed from shallow aquifer for irrigation (mm H2O)*
- real *8, dimension(:), allocatable **parm::ch_l1**
- longest tributary channel length in subbasin (km)*
- real *8, dimension(:), allocatable **parm::wet_no3**
- amount of nitrate originating from surface runoff in wetland at end of day (kg N)*
- real *8, dimension(:), allocatable **parm::ovrlnd**
- overland flow onto HRU from upstream routing unit (mm H2O)*
- real *8, dimension(:), allocatable **parm::canstor**
- amount of water held in canopy storage (mm H2O)*
- real *8, dimension(:), allocatable **parm::irr_mx**
- maximum irrigation amount per auto application (mm)*
- real *8, dimension(:), allocatable **parm::auto_wstr**
- water stress factor which triggers auto irrigation (none or mm)*
- real *8, dimension(:), allocatable **parm::cfrt_id**
- fertilizer/manure id number from database (none)*
- real *8, dimension(:), allocatable **parm::cfrt_kg**
- amount of fertilzier applied to HRU on a given day (kg/ha)*
- real *8, dimension(:), allocatable **parm::cpst_id**
- real *8, dimension(:), allocatable **parm::cpst_kg**
- real *8, dimension(:), allocatable **parm::irr_asq**
- surface runoff ratio*
- real *8, dimension(:), allocatable **parm::irr_eff**
- real *8, dimension(:), allocatable **parm::irrsq**
- surface runoff ratio (0-1) .1 is 10% surface runoff (frac)*
- real *8, dimension(:), allocatable **parm::irrsalt**
- concentration of salt in irrigation water (mg/kg)*
- real *8, dimension(:), allocatable **parm::irrefm**
- real *8, dimension(:), allocatable **parm::bio_eat**
- dry weight of biomass removed by grazing daily ((kg/ha)/day)*
- real *8, dimension(:), allocatable **parm::bio_trmp**
- dry weight of biomass removed by trampling daily ((kg/ha)/day)*
- integer, dimension(:), allocatable **parm::ipst_freq**
- number of days between applications (days)*
- integer, dimension(:), allocatable **parm::ifrt_freq**
- integer, dimension(:), allocatable **parm::irr_noa**
- integer, dimension(:), allocatable **parm::irr_sc**
- integer, dimension(:), allocatable **parm::irr_no**
- integer, dimension(:), allocatable **parm::imp_trig**
- release/impound action code (none):*
- 0 begin impounding water*
- 1 release impounded water*
- integer, dimension(:), allocatable **parm::fert_days**
- integer, dimension(:), allocatable **parm::irr_sca**
- integer, dimension(:), allocatable **parm::idplt**
- land cover/crop identification code for first crop grown in HRU (the only crop if there is no rotation) (from crop.dat)*
- (none)*

- integer, dimension(:), allocatable **parm::pest_days**
- integer, dimension(:), allocatable **parm::wstrs_id**
- real *8, dimension(:, :), allocatable **parm::bio_aahv**
- real *8, dimension(:), allocatable **parm::cumei**
- real *8, dimension(:), allocatable **parm::cumeira**
- real *8, dimension(:), allocatable **parm::cumrt**
- real *8, dimension(:), allocatable **parm::cumrai**
- real *8, dimension(:), allocatable **parm::wet_solp**
amount of soluble P originating from surface runoff in wetland at end of day (kg P)
- real *8, dimension(:), allocatable **parm::wet_chla**
amount of chlorophyll-a in wetland at end of day (kg chla)
- real *8, dimension(:), allocatable **parm::wet_no3s**
amount of nitrate originating from lateral flow in wetland at end of day (kg N)
- real *8, dimension(:), allocatable **parm::pstsol**
soluble pesticide leached from bottom of soil profile (kg pst/ha)
- real *8, dimension(:), allocatable **parm::pnd_no3g**
amount of nitrate originating from groundwater in pond at end of day (kg N)
- real *8, dimension(:), allocatable **parm::wet_seci**
secchi-disk depth in wetland at end of day (m)
- real *8, dimension(:), allocatable **parm::delay**
groundwater delay: time required for water leaving the bottom of the root zone to reach the shallow aquifer (days)
- real *8, dimension(:), allocatable **parm::gwht**
groundwater height (m)
- real *8, dimension(:), allocatable **parm::gw_q**
groundwater contribution to streamflow from HRU on current day (mm H2O)
- real *8, dimension(:), allocatable **parm::pnd_solpg**
amount of soluble P originating from groundwater in pond at end of day (kg P)
- real *8, dimension(:), allocatable **parm::alpha_bf**
alpha factor for groundwater recession curve (1/days)
- real *8, dimension(:), allocatable **parm::alpha_bfe**
 $\exp(-\alpha_{bf}f)$ (none)
- real *8, dimension(:), allocatable **parm::gw_spyld**
specific yield for shallow aquifer (m^3/m^3)
- real *8, dimension(:), allocatable **parm::alpha_bf_d**
alpha factor for groundwater recession curve of the deep aquifer (1/days)
- real *8, dimension(:), allocatable **parm::alpha_bfe_d**
 $\exp(-\alpha_{bf_d}f_d)$ for deep aquifer (none)
- real *8, dimension(:), allocatable **parm::gw_qdeep**
- real *8, dimension(:), allocatable **parm::gw_delaye**
 $\exp(-1/delay)$ (none)
- real *8, dimension(:), allocatable **parm::gw_revap**
revap coeff: this variable controls the amount of water moving from the shallow aquifer to the root zone as a result of soil moisture depletion (none)
- real *8, dimension(:), allocatable **parm::rchrg_dp**
recharge to deep aquifer: the fraction of root zone percolation that reaches the deep aquifer (none)
- real *8, dimension(:), allocatable **parm::anion_excl**
fraction of porosity from which anions are excluded
- real *8, dimension(:), allocatable **parm::revapmn**
threshold depth of water in shallow aquifer required to allow revap to occur (mm H2O)
- real *8, dimension(:), allocatable **parm::rchrg**
- real *8, dimension(:), allocatable **parm::bio_min**

- minimum plant biomass for grazing (kg/ha)*
- real *8, dimension(:), allocatable [parm::ffc](#)
- initial HRU soil water content expressed as fraction of field capacity (none)*
- real *8, dimension(:), allocatable [parm::surqsolp](#)
- real *8, dimension(:), allocatable [parm::deepst](#)
- depth of water in deep aquifer (mm H2O)*
- real *8, dimension(:), allocatable [parm::shallst](#)
- depth of water in shallow aquifer in HRU (mm H2O)*
- real *8, dimension(:), allocatable [parm::wet_solpg](#)
- amount of soluble P originating from groundwater in wetland at end of day (kg P)*
- real *8, dimension(:), allocatable [parm::cklsp](#)
- real *8, dimension(:), allocatable [parm::rchrg_src](#)
- real *8, dimension(:), allocatable [parm::trapeff](#)
- filter strip trapping efficiency (used for everything but bacteria) (none)*
- real *8, dimension(:), allocatable [parm::sol_avbd](#)
- average bulk density for soil profile (Mg/m³)*
- real *8, dimension(:), allocatable [parm::wet_no3g](#)
- amount of nitrate originating from groundwater in wetland at end of day (kg N)*
- real *8, dimension(:), allocatable [parm::tdrain](#)
- time to drain soil to field capacity yield used in autofertilization (hours)*
- real *8, dimension(:), allocatable [parm::gwqmn](#)
- threshold depth of water in shallow aquifer required before groundwater flow will occur (mm H2O)*
- real *8, dimension(:), allocatable [parm::snotmp](#)
- temperature of snow pack in HRU (deg C)*
- real *8, dimension(:), allocatable [parm::pplnt](#)
- real *8, dimension(:), allocatable [parm::gdrain](#)
- drain tile lag time: the amount of time between the transfer of water from the soil to the drain tile and the release of the water from the drain tile to the reach (hours)*
- real *8, dimension(:), allocatable [parm::ddrain](#)
- depth to the sub-surface drain (mm)*
- real *8, dimension(:), allocatable [parm::sol_crk](#)
- crack volume potential of soil (none)*
- real *8, dimension(:), allocatable [parm::brt](#)
- fraction of surface runoff within the subbasin which takes 1 day or less to reach the subbasin outlet (none)*
- real *8, dimension(:), allocatable [parm::dayl](#)
- day length (hours)*
- real *8, dimension(:), allocatable [parm::sstmaxd](#)
- static maximum depressional storage; read from .sdr (mm)*
- real *8, dimension(:), allocatable [parm::re](#)
- effective radius of drains (mm)*
- real *8, dimension(:), allocatable [parm::sdrain](#)
- distance between two drain tubes or tiles (mm)*
- real *8, dimension(:), allocatable [parm::ddrain_hru](#)
- real *8, dimension(:), allocatable [parm::drain_co](#)
- drainage coefficient (mm/day)*
- real *8, dimension(:), allocatable [parm::latksatf](#)
- multiplication factor to determine conk(j1,j) from sol_k(j1,j) for HRU (none)*
- real *8, dimension(:), allocatable [parm::pc](#)
- pump capacity (default pump capacity = 1.042mm/hr or 25mm/day) (mm/hr)*
- real *8, dimension(:), allocatable [parm::stmaxd](#)
- real *8, dimension(:), allocatable [parm::rnd3](#)

- random number between 0.0 and 1.0 (none)*

 - real *8, dimension(:), allocatable **parm::rnd2**
- random number between 0.0 and 1.0 (none)*

 - real *8, dimension(:), allocatable **parm::twash**
 - real *8, dimension(:), allocatable **parm::sol_cnsw**
 - real *8, dimension(:), allocatable **parm::doxq**
 - real *8, dimension(:), allocatable **parm::rnd8**
- random number between 0.0 and 1.0 (none)*

 - real *8, dimension(:), allocatable **parm::rnd9**
- random number between 0.0 and 1.0 (none)*

 - real *8, dimension(:), allocatable **parm::perc**
 - real *8, dimension(:), allocatable **parm::sol_sumwp**
 - real *8, dimension(:), allocatable **parm::qdr**
- total amount of water entering main channel for day from HRU (mm H2O)*

 - real *8, dimension(:), allocatable **parm::tauton**
 - real *8, dimension(:), allocatable **parm::tautop**
 - real *8, dimension(:), allocatable **parm::cbodu**
 - real *8, dimension(:), allocatable **parm::chl_a**
 - real *8, dimension(:), allocatable **parm::tfertn**
 - real *8, dimension(:), allocatable **parm::tfertp**
 - real *8, dimension(:), allocatable **parm::tgrazn**
 - real *8, dimension(:), allocatable **parm::tgrazp**
 - real *8, dimension(:), allocatable **parm::latq**
- total lateral flow in soil profile for the day in HRU (mm H2O)*

 - real *8, dimension(:), allocatable **parm::latno3**
 - real *8, dimension(:), allocatable **parm::minpgw**
 - real *8, dimension(:), allocatable **parm::no3gw**
 - real *8, dimension(:), allocatable **parm::nplnt**
 - real *8, dimension(:), allocatable **parm::tileq**
 - real *8, dimension(:), allocatable **parm::tileno3**
 - real *8, dimension(:), allocatable **parm::sedminpa**
 - real *8, dimension(:), allocatable **parm::sedminps**
 - real *8, dimension(:), allocatable **parm::sedorgn**
 - real *8, dimension(:), allocatable **parm::sedyld**
- soil loss caused by water erosion for day in HRU (metric tons)*

 - real *8, dimension(:), allocatable **parm::sepbtm**
- percolation from bottom of soil profile for the day in HRU (mm H2O)*

 - real *8, dimension(:), allocatable **parm::sedorgp**
 - real *8, dimension(:), allocatable **parm::strsn**
 - real *8, dimension(:), allocatable **parm::surfq**
- surface runoff generated in HRU on the current day (mm H2O)*

 - real *8, dimension(:), allocatable **parm::strsp**
 - real *8, dimension(:), allocatable **parm::strstmp**
 - real *8, dimension(:), allocatable **parm::surqno3**
 - real *8, dimension(:), allocatable **parm::hru_ha**
- area of HRU in hectares (ha)*

 - real *8, dimension(:), allocatable **parm::hru_dafr**
- fraction of total watershed area contained in HRU (km2/km2)*

 - real *8, dimension(:), allocatable **parm::tcfrtn**
 - real *8, dimension(:), allocatable **parm::tcfrtp**
 - real *8, dimension(:), allocatable **parm::drydep_no3**
- atmospheric dry deposition of nitrates (kg/ha/yr)*

 - real *8, dimension(:), allocatable **parm::drydep_nh4**

- atmospheric dry deposition of ammonia (kg/ha/yr)*
- real *8, dimension(:), allocatable [parm::bio_yrms](#)
- annual biomass (dry weight) in the HRU (metric tons/ha)*
- real *8, dimension(:), allocatable [parm::phubase](#)
- base zero total heat units (used when no land cover is growing) (heat units)*
- real *8, dimension(:), allocatable [parm::hvstiadj](#)
- optimal harvest index for current time during growing season ((kg/ha)/(kg/ha))*
- real *8, dimension(:), allocatable [parm::laiday](#)
- leaf area index for HRU (m^2/m^2)*
- real *8, dimension(:), allocatable [parm::chlap](#)
- chlorophyll-a production coefficient for pond (none)*
- real *8, dimension(:), allocatable [parm::pnd_psed](#)
- amount of mineral P attached to sediment originating from surface runoff in pond at end of day (kg P)*
- real *8, dimension(:), allocatable [parm::laimxfr](#)
- real *8, dimension(:), allocatable [parm::seccip](#)
- water clarity coefficient for pond (none)*
- real *8, dimension(:), allocatable [parm::plantn](#)
- amount of nitrogen in plant biomass (kg N/ha)*
- real *8, dimension(:), allocatable [parm::plt_et](#)
- actual ET simulated during life of plant (mm H2O)*
- real *8, dimension(:), allocatable [parm::wet_psed](#)
- amount of mineral P attached to sediment originating from surface runoff in wetland at end of day (kg P)*
- real *8, dimension(:), allocatable [parm::bio_aams](#)
- average annual biomass (dry weight) in the HRU (metric tons)*
- real *8, dimension(:), allocatable [parm::plantp](#)
- amount of phosphorus in plant biomass (kg P/ha)*
- real *8, dimension(:), allocatable [parm::plt_pet](#)
- potential ET simulated during life of plant (mm H2O)*
- real *8, dimension(:), allocatable [parm::dormhr](#)
- time threshold used to define dormant period for plant (when daylength is within the time specified by dl from the minimum daylength for the area, the plant will go dormant) (hour)*
- real *8, dimension(:), allocatable [parm::lai_yrmx](#)
- maximum leaf area index for the year in the HRU (none)*
- real *8, dimension(:), allocatable [parm::bio_aamx](#)
- real *8, dimension(:), allocatable [parm::lat_pst](#)
- amount of pesticide in lateral flow in HRU for the day (kg pst/ha)*
- real *8, dimension(:), allocatable [parm::fld_fr](#)
- fraction of HRU area that drains into floodplain (km^2/km^2)*
- real *8, dimension(:), allocatable [parm::orig_snohru](#)
- real *8, dimension(:), allocatable [parm::orig_potvol](#)
- real *8, dimension(:), allocatable [parm::pltfr_n](#)
- fraction of plant biomass that is nitrogen (none)*
- real *8, dimension(:), allocatable [parm::orig_alai](#)
- real *8, dimension(:), allocatable [parm::orig_bioms](#)
- real *8, dimension(:), allocatable [parm::pltfr_p](#)
- fraction of plant biomass that is phosphorus (none)*
- real *8, dimension(:), allocatable [parm::orig_phuacc](#)
- real *8, dimension(:), allocatable [parm::orig_sumix](#)
- real *8, dimension(:), allocatable [parm::phu_plt](#)
- total number of heat units to bring plant to maturity (heat units)*
- real *8, dimension(:), allocatable [parm::orig_phu](#)
- real *8, dimension(:), allocatable [parm::orig_shallst](#)

- real *8, dimension(:), allocatable **parm::orig_deepst**
- real *8, dimension(:), allocatable **parm::rip_fr**
fraction of HRU area that drains into riparian zone (km^2/km^2)
- real *8, dimension(:), allocatable **parm::orig_pndvol**
- real *8, dimension(:), allocatable **parm::orig_pndsed**
- real *8, dimension(:), allocatable **parm::orig_pndno3**
- real *8, dimension(:), allocatable **parm::orig_pndsolv**
- real *8, dimension(:), allocatable **parm::orig_pndorgn**
- real *8, dimension(:), allocatable **parm::orig_pndorgp**
- real *8, dimension(:), allocatable **parm::orig_wetvol**
- real *8, dimension(:), allocatable **parm::orig_wetsed**
- real *8, dimension(:), allocatable **parm::orig_wetno3**
- real *8, dimension(:), allocatable **parm::orig_wetsolv**
- real *8, dimension(:), allocatable **parm::orig_wetorgn**
- real *8, dimension(:), allocatable **parm::orig_wetorgp**
- real *8, dimension(:), allocatable **parm::orig_solcov**
- real *8, dimension(:), allocatable **parm::orig_solsw**
- real *8, dimension(:), allocatable **parm::orig_potno3**
- real *8, dimension(:), allocatable **parm::orig_potsed**
- real *8, dimension(:), allocatable **parm::wtab**
water table based on 30 day antecedent climate (precip,et) (mm)
- real *8, dimension(:), allocatable **parm::wtab_mn**
- real *8, dimension(:), allocatable **parm::wtab_mx**
- real *8, dimension(:), allocatable **parm::shallst_n**
nitrate concentration in shallow aquifer converted to kg/ha (ppm NO₃-N)
- real *8, dimension(:), allocatable **parm::gw_nloss**
- real *8, dimension(:), allocatable **parm::rchrg_n**
- real *8, dimension(:), allocatable **parm::det_san**
- real *8, dimension(:), allocatable **parm::det_sil**
- real *8, dimension(:), allocatable **parm::det_cla**
- real *8, dimension(:), allocatable **parm::det_sag**
- real *8, dimension(:), allocatable **parm::det_lag**
- real *8, dimension(:), allocatable **parm::afrt_surface**
fraction of fertilizer which is applied to top 10 mm of soil (the remaining fraction is applied to first soil layer) (none)
- real *8, dimension(:), allocatable **parm::tnylida**
- real *8 **parm::frit_surface**
fraction of fertilizer which is applied to the top 10 mm of soil (the remaining fraction is applied to the first soil layer) (none)
- real *8, dimension(:), allocatable **parm::auto_nyr**
maximum NO₃-N content allowed to be applied in one year (kg NO₃-N/ha)
- real *8, dimension(:), allocatable **parm::auto_napp**
maximum NO₃-N content allowed in one fertilizer application (kg NO₃-N/ha)
- real *8, dimension(:), allocatable **parm::auto_nstrs**
nitrogen stress factor which triggers auto fertilization (none)
- real *8, dimension(:), allocatable **parm::manure_kg**
dry weight of manure deposited on HRU daily ((kg/ha)/day)
- real *8, dimension(:,:), allocatable **parm::rcn_mo**
- real *8, dimension(:,:), allocatable **parm::rammo_mo**
- real *8, dimension(:,:), allocatable **parm::drydep_no3_mo**
- real *8, dimension(:,:), allocatable **parm::drydep_nh4_mo**
- real *8, dimension(:), allocatable **parm::rcn_d**
- real *8, dimension(:), allocatable **parm::rammo_d**
- real *8, dimension(:), allocatable **parm::drydep_no3_d**

- real *8, dimension(:), allocatable **parm::drydep_nh4_d**
- real *8, dimension(:,:), allocatable **parm::yldn**
- integer, dimension(:,:), allocatable **parm::gwati**
- real *8, dimension(:,:), allocatable **parm::gwatn**
- real *8, dimension(:,:), allocatable **parm::gwatl**
- real *8, dimension(:,:), allocatable **parm::gwatw**
- real *8, dimension(:,:), allocatable **parm::gwatd**
- real *8, dimension(:,:), allocatable **parm::gwatveg**
- real *8, dimension(:,:), allocatable **parm::gwata**
- real *8, dimension(:,:), allocatable **parm::gwats**
- real *8, dimension(:,:), allocatable **parm::gwatspcon**
- real *8, dimension(:,:), allocatable **parm::rfqeo_30d**
- real *8, dimension(:,:), allocatable **parm::eo_30d**
- real *8, dimension(:), allocatable **parm::psetlp1**
phosphorus settling rate for 1st season (m/day)
- real *8, dimension(:), allocatable **parm::psetlp2**
phosphorus settling rate for 2nd season (m/day)
- real *8, dimension(:,:), allocatable **parm::wgnold**
previous value of wgncur(:,:) (none)
- real *8, dimension(:,:), allocatable **parm::wgncur**
parameter to predict the impact of precip on other weather attributes (none)
wgncur(1,:) parameter which predicts impact of precip on daily maximum air temperature
wgncur(2,:) parameter which predicts impact of precip on daily minimum air temperature
wgncur(3,:) parameter which predicts impact of precip on daily solar radiation
- real *8, dimension(:), allocatable **parm::wrt1**
1st shape parameter for calculation of water retention (none)
- real *8, dimension(:), allocatable **parm::wrt2**
2nd shape parameter for calculation of water retention (none)
- real *8, dimension(:,:), allocatable **parm::pst_enr**
pesticide enrichment ratio (none)
- real *8, dimension(:,:), allocatable **parm::zdb**
- real *8, dimension(:,:), allocatable **parm::pst_surq**
- real *8, dimension(:,:), allocatable **parm::plt_pst**
pesticide on plant foliage (kg/ha)
- real *8, dimension(:), allocatable **parm::psetlw1**
phosphorus settling rate for 1st season (m/day)
- real *8, dimension(:), allocatable **parm::psetlw2**
phosphorus settling rate for 2nd season (m/day)
- real *8, dimension(:,:), allocatable **parm::pst_sed**
- real *8, dimension(:,:), allocatable **parm::wupnd**
average daily water removal from the pond for the month ($10^4 \text{ m}^3/\text{day}$)
- real *8, dimension(:,:), allocatable **parm::phi**
phi(1,:) cross-sectional area of flow at bankfull depth (m^2) phi(2,:) (none) phi(3,:) (none) phi(4,:) (none) phi(5,:) (none) phi(6,:) bottom width of main channel (m) phi(7,:) depth of water when reach is at bankfull depth (m) phi(8,:) average velocity when reach is at bankfull depth (m/s) phi(9,:) wave celerity when reach is at bankfull depth (m/s) phi(10,:) storage time constant for reach at bankfull depth (ratio of storage to discharge) (hour) phi(11,:) average velocity when reach is at 0.1 bankfull depth (low flow) (m/s) phi(12,:) wave celerity when reach is at 0.1 bankfull depth (low flow) (m/s) phi(13,:) storage time constant for reach at 0.1 bankfull depth (low flow) (ratio of storage to discharge) (hour)
- real *8, dimension(:,:), allocatable **parm::pcpband**
precipitation for the day in band in HRU (mm H₂O)
- real *8, dimension(:,:), allocatable **parm::tavband**
average temperature for the day in band in HRU (deg C)
- real *8, dimension(:), allocatable **parm::wat_phi1**

- cross-sectional area of flow at bankfull depth (m^2)*
 - real *8, dimension(:), allocatable [parm::wat_phi5](#)
- flow rate when reach is at bankfull depth (m^3/s)*
 - real *8, dimension(:), allocatable [parm::wat_phi6](#)
- bottom width of main channel (m)*
 - real *8, dimension(:), allocatable [parm::wat_phi7](#)
- depth of water when reach is at bankfull (m)*
 - real *8, dimension(:), allocatable [parm::wat_phi8](#)
- average velocity when reach is at bankfull depth (m/s)*
 - real *8, dimension(:), allocatable [parm::wat_phi9](#)
- wave celerity when reach is at bankfull depth (m/s)*
 - real *8, dimension(:), allocatable [parm::wat_phi10](#)
- storage time constant for reach at bankfull depth (ratio of storage to discharge) (hour)*
 - real *8, dimension(:), allocatable [parm::wat_phi11](#)
- average velocity when reach is at 0.1 bankfull depth (low flow) (m/s)*
 - real *8, dimension(:), allocatable [parm::wat_phi12](#)
- wave celerity when reach is at 0.1 bankfull depth (low flow) (m/s)*
 - real *8, dimension(:), allocatable [parm::wat_phi13](#)
- storage time constant for reach at 0.1 bankfull depth (low flow) (ratio of storage to discharge) (hour)*
 - real *8, dimension(:,:), allocatable [parm::snoeb](#)
- snow water content in elevation band on current day (mm H2O)*
 - real *8, dimension(:,:), allocatable [parm::wudeep](#)
- average daily water removal from the deep aquifer for the month ($10^4 m^3/day$)*
 - real *8, dimension(:,:), allocatable [parm::wushal](#)
- average daily water removal from the shallow aquifer for the month ($10^4 m^3/day$)*
 - real *8, dimension(:,:), allocatable [parm::tmnband](#)
- minimum temperature for the day in band in HRU (deg C)*
 - real *8, dimension(:), allocatable [parm::bss1](#)
 - real *8, dimension(:), allocatable [parm::bss2](#)
 - real *8, dimension(:), allocatable [parm::bss3](#)
 - real *8, dimension(:), allocatable [parm::bss4](#)
 - real *8, dimension(:), allocatable [parm::nsetlw1](#)
- nitrogen settling rate for 1st season (m/day)*
 - real *8, dimension(:), allocatable [parm::nsetlw2](#)
- nitrogen settling rate for 2nd season (m/day)*
 - real *8, dimension(:,:), allocatable [parm::snotmpeb](#)
- temperature of snow pack in elevation band (deg C)*
 - real *8, dimension(:), allocatable [parm::surf_bs1](#)
- amount of surface runoff lagged over one day (mm H2O)*
 - real *8, dimension(:), allocatable [parm::surf_bs2](#)
 - real *8, dimension(:), allocatable [parm::surf_bs3](#)
 - real *8, dimension(:), allocatable [parm::surf_bs4](#)
 - real *8, dimension(:), allocatable [parm::surf_bs5](#)
 - real *8, dimension(:), allocatable [parm::surf_bs6](#)
 - real *8, dimension(:), allocatable [parm::surf_bs7](#)
 - real *8, dimension(:), allocatable [parm::surf_bs8](#)
 - real *8, dimension(:), allocatable [parm::surf_bs9](#)
 - real *8, dimension(:), allocatable [parm::surf_bs10](#)
 - real *8, dimension(:), allocatable [parm::surf_bs11](#)
 - real *8, dimension(:), allocatable [parm::surf_bs12](#)
 - real *8, dimension(:), allocatable [parm::surf_bs13](#)
 - real *8, dimension(:), allocatable [parm::surf_bs14](#)

- real *8, dimension(:), allocatable **parm::surf_bs15**
- real *8, dimension(:), allocatable **parm::surf_bs16**
- real *8, dimension(:), allocatable **parm::surf_bs17**
- real *8, dimension(:), allocatable **parm::nsetlp1**
nitrogen settling rate for 1st season (m/day)
- real *8, dimension(:), allocatable **parm::nsetlp2**
nitrogen settling rate for 2nd season (m/day)
- real *8, dimension(:, :), allocatable **parm::tmxband**
maximum temperature for the day in band in HRU (deg C)
- real *8, dimension(:, :), allocatable **parm::fracd**
fraction of solar radiation occurring during hour in day in HRU (none)
- real *8, dimension(:, :), allocatable **parm::rainsub**
precipitation for the time step during the day in HRU (mm H2O)
- real *8, dimension(:), allocatable **parm::rstpbsb**
- real *8, dimension(:, :), allocatable **parm::orig_snoeb**
- real *8, dimension(:, :), allocatable **parm::orig_pltpst**
- real *8, dimension(:, :), allocatable **parm::terr_p**
- real *8, dimension(:, :), allocatable **parm::terr_cn**
- real *8, dimension(:, :), allocatable **parm::terr_sl**
- real *8, dimension(:, :), allocatable **parm::drain_d**
- real *8, dimension(:, :), allocatable **parm::drain_t**
- real *8, dimension(:, :), allocatable **parm::drain_g**
- real *8, dimension(:, :), allocatable **parm::drain_idep**
- real *8, dimension(:, :), allocatable **parm::cont_cn**
- real *8, dimension(:, :), allocatable **parm::cont_p**
- real *8, dimension(:, :), allocatable **parm::filt_w**
- real *8, dimension(:, :), allocatable **parm::strip_n**
- real *8, dimension(:, :), allocatable **parm::strip_cn**
- real *8, dimension(:, :), allocatable **parm::strip_c**
- real *8, dimension(:, :), allocatable **parm::strip_p**
- real *8, dimension(:, :), allocatable **parm::fire_cn**
- real *8, dimension(:, :), allocatable **parm::cropno_upd**
- real *8, dimension(:, :), allocatable **parm::hi_upd**
- real *8, dimension(:, :), allocatable **parm::laimx_upd**
- real *8, dimension(:, :, :), allocatable **parm::phug**
fraction of plant heat units at which grazing begins (none)
- real *8, dimension(:, :, :), allocatable **parm::pst_lag**
- integer, dimension(:), allocatable **parm::hrupest**
pesticide use flag (none)
0: no pesticides used in HRU
1: pesticides used in HRU
- integer, dimension(:), allocatable **parm::nrelease**
sequence number of impound/release operation within the year (none)
- integer, dimension(:), allocatable **parm::swtrg**
rainfall event flag (none):
0: no rainfall event over midnight
1: rainfall event over midnight
- integer, dimension(:), allocatable **parm::nrot**
number of years of rotation (none)
- integer, dimension(:), allocatable **parm::nfert**
sequence number of fertilizer application within the year (none)
- integer, dimension(:), allocatable **parm::nro**
sequence number of year in rotation (none)

- integer, dimension(:), allocatable [parm::igro](#)
land cover status code (none). This code informs the model whether or not a land cover is growing at the beginning of the simulation
0 no land cover growing
1 land cover growing
- integer, dimension(:), allocatable [parm::ipnd1](#)
beginning month of nutrient settling season (none)
- integer, dimension(:), allocatable [parm::ipnd2](#)
ending month of nutrient settling season (none)
- integer, dimension(:), allocatable [parm::nair](#)
sequence number of auto-irrigation application within the year (none)
- integer, dimension(:), allocatable [parm::iflod1](#)
beginning month of non-flood season (none)
- integer, dimension(:), allocatable [parm::iflod2](#)
ending month of non-flood season (none)
- integer, dimension(:), allocatable [parm::ndtarg](#)
number of days required to reach target storage from current pond storage (none)
- integer, dimension(:), allocatable [parm::nirr](#)
sequence number of irrigation application within the year (none)
- integer, dimension(:), allocatable [parm::iafrttyp](#)
- integer, dimension(:), allocatable [parm::nstress](#)
- integer, dimension(:), allocatable [parm::igrotree](#)
- integer, dimension(:), allocatable [parm::grz_days](#)
number of days grazing will be simulated (none)
- integer, dimension(:), allocatable [parm::nmgt](#)
management code (for GIS output only) (none)
- integer, dimension(:), allocatable [parm::nafert](#)
sequence number of auto-fert application within the year (none)
- integer, dimension(:), allocatable [parm::nsweep](#)
sequence number of street sweeping operation within the year (none)
- integer, dimension(:), allocatable [parm::icr](#)
sequence number of crop grown within the current year (none)
- integer, dimension(:), allocatable [parm::ncut](#)
sequence number of harvest operation within a year (none)
- integer, dimension(:), allocatable [parm::irrno](#)
irrigation source location (none)
if IRRSC=1, IRRNO is the number of the reach
if IRRSC=2, IRRNO is the number of the reservoir
if IRRSC=3, IRRNO is the number of the subbasin
if IRRSC=4, IRRNO is the number of the subbasin
if IRRSC=5, not used
- integer, dimension(:), allocatable [parm::sol_nly](#)
number of soil layers in HRU (none)
- integer, dimension(:), allocatable [parm::npcp](#)
prior day category (none)
1 dry day
2 wet day
- integer, dimension(:), allocatable [parm::irn](#)
average annual number of irrigation applications in HRU (none)
- integer, dimension(:), allocatable [parm::ncf](#)
sequence number of continuous fertilization operation within the year (none)
- integer, dimension(:), allocatable [parm::ngr](#)
sequence number of grazing operation within the year (none)

- integer, dimension(:), allocatable **parm::igrz**
grazing flag for HRU (none):
0 HRU currently not grazed
1 HRU currently grazed
- integer, dimension(:), allocatable **parm::ndeat**
number of days HRU has been grazed (days)
- integer, dimension(:), allocatable **parm::hru_sub**
subbasin in which HRU is located (none)
- integer, dimension(:), allocatable **parm::urblu**
urban land type identification number from urban.dat (none)
- integer, dimension(:), allocatable **parm::ldrain**
soil layer where drainage tile is located (none)
- integer, dimension(:), allocatable **parm::idorm**
dormancy status code (none):
0 land cover growing (not dormant)
1 land cover dormant
- integer, dimension(:), allocatable **parm::hru_seq**
- integer, dimension(:), allocatable **parm::iurban**
urban simulation code (none):
0 no urban sections in HRU
1 urban sections in HRU, simulate using USGS regression equations
2 urban sections in HRU, simulate using build up/wash off algorithm
- integer, dimension(:), allocatable **parm::iday_fert**
- integer, dimension(:), allocatable **parm::icfrt**
- integer, dimension(:), allocatable **parm::ifld**
number of HRU (in subbasin) that is a floodplain (none)
- integer, dimension(:), allocatable **parm::irip**
number of HRU (in subbasin) that is a riparian zone (none)
- integer, dimension(:), allocatable **parm::hrugis**
GIS code printed to output files (output.hru, .rch) (none)
- integer, dimension(:), allocatable **parm::ndcfrt**
- integer, dimension(:), allocatable **parm::irrsc**
irrigation source code (none):
1 divert water from reach
2 divert water from reservoir
3 divert water from shallow aquifer
4 divert water from deep aquifer
5 divert water from source outside watershed
- integer, dimension(:), allocatable **parm::ntil**
sequence number of tillage operation within current year (none)
- integer, dimension(:), allocatable **parm::orig_igro**
- integer, dimension(:), allocatable **parm::iwatable**
- integer, dimension(:), allocatable **parm::curyr_mat**
- integer, dimension(:), allocatable **parm::icpst**
icpst = 0 do not apply
icpst = 1 application period
- integer, dimension(:), allocatable **parm::ndcpst**
current day within the application period (day)
- integer, dimension(:), allocatable **parm::ncpest**
- integer, dimension(:), allocatable **parm::iday_pest**
current day between applications (day)
- integer, dimension(:), allocatable **parm::irr_flag**
- integer, dimension(:), allocatable **parm::irra_flag**
- integer, dimension(:,:), allocatable **parm::rndseed**

random number generator seeds array. The seeds in the array are used to generate random numbers for the following purposes (none):

- (1) wet/dry day probability*
- (2) solar radiation*
- (3) precipitation*
- (4) USLE rainfall erosion index*
- (5) wind speed*
- (6) 0.5 hr rainfall fraction*
- (7) relative humidity*
- (8) maximum temperature*
- (9) minimum temperature*
- (10) generate new random numbers*

- integer, dimension(:,:), allocatable **parm::iterr**
- integer, dimension(:,:), allocatable **parm::iyterr**
- integer, dimension(:,:), allocatable **parm::itdrain**
- integer, dimension(:,:), allocatable **parm::iydrain**
- integer, dimension(:,:), allocatable **parm::ncrops**
- integer, dimension(:), allocatable **parm::manure_id**

manure (fertilizer) identification number from fert.dat (none)

- integer, dimension(:,:), allocatable **parm::mgt_sdr**
- integer, dimension(:,:), allocatable **parm::idplot**
- integer, dimension(:,:), allocatable **parm::icont**
- integer, dimension(:,:), allocatable **parm::iycont**
- integer, dimension(:,:), allocatable **parm::ifilt**
- integer, dimension(:,:), allocatable **parm::iyfilt**
- integer, dimension(:,:), allocatable **parm::istrip**
- integer, dimension(:,:), allocatable **parm::iystrip**
- integer, dimension(:,:), allocatable **parm::iopday**
- integer, dimension(:,:), allocatable **parm::iopyr**
- integer, dimension(:,:), allocatable **parm::mgt_ops**
- real *8, dimension(:), allocatable **parm::wshd_pstap**

total amount of pesticide type applied in watershed during simulation (kg/ha)

- real *8, dimension(:), allocatable **parm::wshd_pstdg**
- integer, dimension(12) **parm::ndmo**
- integer, dimension(:), allocatable **parm::npno**

array of unique pesticides used in watershed (none)

- integer, dimension(:), allocatable **parm::mcrhru**
- character(len=13), dimension(18) **parm::rfile**

rainfall file names (.pcp)

- character(len=13), dimension(18) **parm::tfile**

temperature file names (.tmp)

- character(len=4), dimension(1000) **parm::urbname**

name of urban land use

- character(len=1), dimension(:), allocatable **parm::kirr**

irrigation in HRU

- character(len=1), dimension(:), allocatable **parm::hydgrp**
- character(len=16), dimension(:), allocatable **parm::snam**

soil series name

- character(len=17), dimension(300) **parm::pname**

name of pesticide/toxin

- character(len=4), dimension(60) **parm::title**

description lines in file.cio (1st 3 lines)

- character(len=4), dimension(5000) **parm::cpnm**

four character code to represent crop name

- character(len=17), dimension(50) **parm::fname**
- real *8, dimension(:, :, :), allocatable **parm::flomon**
average daily water loading for month (m^3/day)
- real *8, dimension(:, :, :), allocatable **parm::solpstmon**
average daily soluble pesticide loading for month (mg pst/day)
- real *8, dimension(:, :, :), allocatable **parm::srbspstmon**
average daily sorbed pesticide loading for month (mg pst/day)
- real *8, dimension(:, :, :), allocatable **parm::orgnmon**
average daily organic N loading for month (kg N/day)
- real *8, dimension(:, :, :), allocatable **parm::orgpmon**
average daily organic P loading for month (kg P/day)
- real *8, dimension(:, :, :), allocatable **parm::sedmon**
average daily sediment loading for month (metric tons/day)
- real *8, dimension(:, :, :), allocatable **parm::minpmon**
average daily mineral P loading for month (kg P/day)
- real *8, dimension(:, :, :), allocatable **parm::nh3mon**
average amount of NH3-N loaded to stream on a given day in the month (kg N/day)
- real *8, dimension(:, :, :), allocatable **parm::no3mon**
average daily NO3-N loading for month (kg N/day)
- real *8, dimension(:, :, :), allocatable **parm::bactlpmon**
average amount of less persistent bacteria loaded to stream on a given day in the month (# bact/day)
- real *8, dimension(:, :, :), allocatable **parm::bactpmon**
average amount of persistent bacteria loaded to stream on a given day in the month (# bact/day)
- real *8, dimension(:, :, :), allocatable **parm::no2mon**
average amount of NO2-N loaded to stream on a given day in the month (kg N/day)
- real *8, dimension(:, :, :), allocatable **parm::cmtl1mon**
average amount of conservative metal #1 loaded to stream on a given day in the month (# bact/day)
- real *8, dimension(:, :, :), allocatable **parm::cmtl2mon**
average amount of conservative metal #2 loaded to stream on a given day in the month (# bact/day)
- real *8, dimension(:, :, :), allocatable **parm::cmtl3mon**
average amount of conservative metal #3 loaded to stream on a given day in the month (# bact/day)
- real *8, dimension(:, :, :), allocatable **parm::cbodmon**
average daily loading of CBOD in month (kg/day)
- real *8, dimension(:, :, :), allocatable **parm::chlamon**
average daily loading of chlorophyll-a in month (kg/day)
- real *8, dimension(:, :, :), allocatable **parm::disoxmon**
average daily loading of dissolved O2 in month (kg/day)
- real *8, dimension(:, :), allocatable **parm::floyr**
average daily water loading for year (m^3/day)
- real *8, dimension(:, :), allocatable **parm::orgnyr**
average daily organic N loading for year (kg N/day)
- real *8, dimension(:, :), allocatable **parm::orgpyr**
average daily organic P loading for year (kg P/day)
- real *8, dimension(:, :), allocatable **parm::sedyr**
average daily sediment loading for year (metric tons/day)
- real *8, dimension(:, :), allocatable **parm::minpyr**
average daily mineral P loading for year (kg P/day)
- real *8, dimension(:, :), allocatable **parm::nh3yr**
average daily NH3-N loading for year (kg N/day)
- real *8, dimension(:, :), allocatable **parm::no2yr**
average daily NO2-N loading for year (kg N/day)

- real *8, dimension(:,:), allocatable [parm::no3yr](#)
average daily NO3-N loading for year (kg N/day)
- real *8, dimension(:,:), allocatable [parm::bactlpyr](#)
average daily loading of less persistent bacteria for year (# bact/day)
- real *8, dimension(:,:), allocatable [parm::bactpyr](#)
average daily loading of persistent bacteria for year (# bact/day)
- real *8, dimension(:,:), allocatable [parm::cmtl1yr](#)
average daily loading of conservative metal #1 for year (kg/day)
- real *8, dimension(:,:), allocatable [parm::chlayr](#)
average daily loading of chlorophyll-a in year (kg/day)
- real *8, dimension(:,:), allocatable [parm::cmtl2yr](#)
average daily loading of conservative metal #2 for year (kg/day)
- real *8, dimension(:,:), allocatable [parm::cmtl3yr](#)
average daily loading of conservative metal #3 for year (kg/day)
- real *8, dimension(:,:), allocatable [parm::cbodyr](#)
average daily loading of CBOD in year (kg/day)
- real *8, dimension(:,:), allocatable [parm::disoxyr](#)
average daily loading of dissolved O2 in year (kg/day)
- real *8, dimension(:,:), allocatable [parm::solpstyr](#)
average daily soluble pesticide loading for year (mg pst/day)
- real *8, dimension(:,:), allocatable [parm::srbpstyr](#)
average daily sorbed pesticide loading for year (mg pst/day)
- real *8, dimension(:,:), allocatable **parm::sol_mc**
- real *8, dimension(:,:), allocatable **parm::sol_mn**
- real *8, dimension(:,:), allocatable **parm::sol_mp**
- real *8, dimension(:), allocatable **parm::flocnst**
- real *8, dimension(:), allocatable [parm::orgncnst](#)
average daily organic N loading to reach (kg N/day)
- real *8, dimension(:), allocatable [parm::sedcnst](#)
average daily sediment loading for reach (metric tons/day)
- real *8, dimension(:), allocatable [parm::minpcnst](#)
average daily soluble P loading to reach (kg P/day)
- real *8, dimension(:), allocatable [parm::no3cnst](#)
average daily nitrate loading to reach (kg N/day)
- real *8, dimension(:), allocatable [parm::orgpcnst](#)
average daily organic P loading to reach (kg P/day)
- real *8, dimension(:), allocatable [parm::bactpcnst](#)
average daily persistent bacteria loading to reach (# bact/day)
- real *8, dimension(:), allocatable [parm::nh3cnst](#)
average daily ammonia loading to reach (kg N/day)
- real *8, dimension(:), allocatable [parm::no2cnst](#)
average daily nitrite loading to reach (kg N/day)
- real *8, dimension(:), allocatable [parm::bactlpcnst](#)
average daily less persistent bacteria loading to reach (# bact/day)
- real *8, dimension(:), allocatable [parm::cmtl1cnst](#)
average daily conservative metal #1 loading (kg/day)
- real *8, dimension(:), allocatable [parm::cmtl2cnst](#)
average daily conservative metal #2 loading (kg/day)
- real *8, dimension(:), allocatable [parm::chlacnst](#)
average daily loading of chlorophyll-a (kg/day)
- real *8, dimension(:), allocatable [parm::cmtl3cnst](#)

- average daily conservative metal #3 loading (kg/day)*
 - real *8, dimension(:), allocatable [parm::disoxcnst](#)
 - average daily loading of dissolved O2 (kg/day)*
 - real *8, dimension(:), allocatable [parm::cbodcnst](#)
 - average daily loading of CBOD to reach (kg/day)*
 - real *8, dimension(:), allocatable [parm::solpstcnst](#)
 - average daily soluble pesticide loading (mg/day)*
 - real *8, dimension(:), allocatable [parm::srbspstcnst](#)
 - average daily sorbed pesticide loading (mg/day)*
 - integer [parm::nstep](#)
 - max number of time steps per day or number of lines of rainfall data for each day (none)*
 - integer [parm::idt](#)
 - length of time step used to report precipitation data for sub-daily modeling (minutes)*
 - real *8, dimension(:), allocatable [parm::hrtwtr](#)
 - real *8, dimension(:), allocatable [parm::hhstor](#)
 - real *8, dimension(:), allocatable [parm::hdepth](#)
 - real *8, dimension(:), allocatable [parm::hsdti](#)
 - real *8, dimension(:), allocatable [parm::hrchwtr](#)
 - real *8, dimension(:), allocatable [parm::halgae](#)
 - real *8, dimension(:), allocatable [parm::horgn](#)
 - real *8, dimension(:), allocatable [parm::hnh4](#)
 - real *8, dimension(:), allocatable [parm::hno2](#)
 - real *8, dimension(:), allocatable [parm::hno3](#)
 - real *8, dimension(:), allocatable [parm::horgp](#)
 - real *8, dimension(:), allocatable [parm::hsolp](#)
 - real *8, dimension(:), allocatable [parm::hbod](#)
 - real *8, dimension(:), allocatable [parm::hdisox](#)
 - real *8, dimension(:), allocatable [parm::hchla](#)
 - real *8, dimension(:), allocatable [parm::hsedyld](#)
 - real *8, dimension(:), allocatable [parm::hsedst](#)
 - real *8, dimension(:), allocatable [parm::hharea](#)
 - real *8, dimension(:), allocatable [parm::hsolpst](#)
 - real *8, dimension(:), allocatable [parm::hsorpst](#)
 - real *8, dimension(:), allocatable [parm::hhqday](#)
 - surface runoff generated each timestep of day in HRU (mm H2O)*
 - real *8, dimension(:), allocatable [parm::precipdt](#)
 - precipitation, or effective precipitation reaching soil surface, in time step for HRU (mm H2O)*
 - real *8, dimension(:), allocatable [parm::hhtime](#)
 - real *8, dimension(:), allocatable [parm::hbactp](#)
 - real *8, dimension(:), allocatable [parm::hbactlp](#)
 - integer, dimension(10) [parm::ivar_orig](#)
 - real *8, dimension(10) [parm::rvar_orig](#)
 - integer [parm::nsave](#)
 - number of save commands in .fig file*
 - integer [parm::nauto](#)
 - integer [parm::iatmodep](#)
 - real *8, dimension(:), allocatable [parm::wattemp](#)
 - real *8, dimension(:), allocatable [parm::lkpst_mass](#)
 - real *8, dimension(:), allocatable [parm::lkspst_mass](#)
 - real *8, dimension(:), allocatable [parm::vel_chan](#)
 - real *8, dimension(:), allocatable [parm::vfscon](#)
 - fraction of the total runoff from the entire field entering the most concentrated 10% of the VFS (none)*
 - real *8, dimension(:), allocatable [parm::vfsratio](#)

- field area/VFS area ratio (none)*
- real *8, dimension(:), allocatable [parm::vfsch](#)
- fraction of flow entering the most concentrated 10% of the VFS which is fully channelized (none)*
- real *8, dimension(:), allocatable **parm::vfsi**
- real *8, dimension(:,:), allocatable **parm::filter_i**
- real *8, dimension(:,:), allocatable **parm::filter_ratio**
- real *8, dimension(:,:), allocatable **parm::filter_con**
- real *8, dimension(:,:), allocatable **parm::filter_ch**
- real *8, dimension(:,:), allocatable **parm::sol_n**
- integer [parm::cswat](#)
 - = 0 Static soil carbon (old mineralization routines)
 - = 1 C-FARM one carbon pool model
 - = 2 Century model
- real *8, dimension(:,:), allocatable **parm::sol_bdp**
- real *8, dimension(:,:), allocatable **parm::tillagef**
- real *8, dimension(:), allocatable **parm::rtfr**
- real *8, dimension(:), allocatable **parm::stsol_rd**
- integer **parm::urban_flag**
- integer **parm::dorm_flag**
- real *8 **parm::bf_flg**
- real *8 **parm::iabstr**
- real *8, dimension(:), allocatable **parm::ubnrunoff**
- real *8, dimension(:), allocatable **parm::ubntss**
- real *8, dimension(:,:), allocatable **parm::sub_ubnrunoff**
- real *8, dimension(:,:), allocatable **parm::sub_ubntss**
- real *8, dimension(:,:), allocatable **parm::ovrlnd_dt**
- real *8, dimension(:,:), allocatable **parm::hhsurf_bs1**
- real *8, dimension(:,:), allocatable **parm::hhsurf_bs2**
- integer [parm::iuh](#)
 - unit hydrograph method: 1=triangular UH; 2=gamma funtion UH;*
- integer [parm::sed_ch](#)
 - channel routing for HOURLY; 0=Bagnold; 2=Brownlie; 3=Yang;*
- real *8 [parm::eros_expo](#)
 - an exponent in the overland flow erosion equation ranges 1.5-3.0*
- real *8 [parm::eros_spl](#)
 - coefficient of splash erosion varing 0.9-3.1*
- real *8 [parm::rill_mult](#)
 - Multiplier to USLE_K for soil susceptible to rill erosion, range 0.5-2.0.*
- real *8 **parm::sedprev**
- real *8 **parm::c_factor**
- real *8 [parm::ch_d50](#)
 - median particle diameter of channel bed (mm)*
- real *8 [parm::sig_g](#)
 - geometric standard deviation of particle sizes for the main channel. Mean air temperature at which precipitation is equally likely to be rain as snow/freezing rain.*
- real *8 [parm::uhalpha](#)
 - alpha coefficient for estimating unit hydrograph using a gamma function (*.bsn)*
- real *8 **parm::abstinit**
- real *8 **parm::abstmax**
- real *8, dimension(:,:), allocatable [parm::hhsedy](#)
 - sediment yield from HRU drung a time step applied to HRU (tons)*
- real *8, dimension(:,:), allocatable **parm::sub_subp_dt**
- real *8, dimension(:,:), allocatable **parm::sub_hhsedy**

- real *8, dimension(:,:), allocatable **parm::sub_atmp**
- real *8, dimension(:), allocatable **parm::rhy**
- real *8, dimension(:), allocatable **parm::init_abstrc**
- real *8, dimension(:), allocatable **parm::dratio**
- real *8, dimension(:), allocatable **parm::hrtevp**
- real *8, dimension(:), allocatable **parm::hrttlc**
- real *8, dimension(:,:), allocatable **parm::rchhr**
- real *8, dimension(:), allocatable **parm::hhresflwi**
- real *8, dimension(:), allocatable **parm::hhresflwo**
- real *8, dimension(:), allocatable **parm::hhressedi**
- real *8, dimension(:), allocatable **parm::hhressedo**
- character(len=4), dimension(:), allocatable **parm::lu_nodrain**
- integer, dimension(:), allocatable **parm::bmpdrain**
- real *8, dimension(:), allocatable **parm::sub_cn2**
- real *8, dimension(:), allocatable **parm::sub_ha_urb**
- real *8, dimension(:), allocatable **parm::bmp_recharge**
- real *8, dimension(:), allocatable **parm::sub_ha_imp**
- real *8, dimension(:), allocatable **parm::subdr_km**
- real *8, dimension(:), allocatable **parm::subdr_ickm**
- real *8, dimension(:,:), allocatable **parm::sf_im**
- real *8, dimension(:,:), allocatable **parm::sf_iy**
- real *8, dimension(:,:), allocatable **parm::sp_sa**
- real *8, dimension(:,:), allocatable **parm::sp_pvol**
- real *8, dimension(:,:), allocatable **parm::sp_pd**
- real *8, dimension(:,:), allocatable **parm::sp_sedi**
- real *8, dimension(:,:), allocatable **parm::sp_sede**
- real *8, dimension(:,:), allocatable **parm::ft_sa**
- real *8, dimension(:,:), allocatable **parm::ft_fsa**
- real *8, dimension(:,:), allocatable **parm::ft_dep**
- real *8, dimension(:,:), allocatable **parm::ft_h**
- real *8, dimension(:,:), allocatable **parm::ft_pd**
- real *8, dimension(:,:), allocatable **parm::ft_k**
- real *8, dimension(:,:), allocatable **parm::ft_dp**
- real *8, dimension(:,:), allocatable **parm::ft_dc**
- real *8, dimension(:,:), allocatable **parm::ft_por**
- real *8, dimension(:,:), allocatable **parm::tss_den**
- real *8, dimension(:,:), allocatable **parm::ft_alp**
- real *8, dimension(:,:), allocatable **parm::sf_fr**
- real *8, dimension(:,:), allocatable **parm::sp_qi**
- real *8, dimension(:,:), allocatable **parm::sp_k**
- real *8, dimension(:,:), allocatable **parm::ft_qpnd**
- real *8, dimension(:,:), allocatable **parm::sp_dp**
- real *8, dimension(:,:), allocatable **parm::ft_qsw**
- real *8, dimension(:,:), allocatable **parm::ft_qin**
- real *8, dimension(:,:), allocatable **parm::ft_qout**
- real *8, dimension(:,:), allocatable **parm::ft_sedpnd**
- real *8, dimension(:,:), allocatable **parm::sp_bpw**
- real *8, dimension(:,:), allocatable **parm::ft_bpw**
- real *8, dimension(:,:), allocatable **parm::ft_sed_cumul**
- real *8, dimension(:,:), allocatable **parm::sp_sed_cumul**
- integer, dimension(:), allocatable **parm::num_sf**
- integer, dimension(:,:), allocatable **parm::sf_typ**
- integer, dimension(:,:), allocatable **parm::sf_dim**
- integer, dimension(:,:), allocatable **parm::ft_qfg**
- integer, dimension(:,:), allocatable **parm::sp_qfg**

- integer, dimension(:,:), allocatable **parm::sf_ptp**
- integer, dimension(:,:), allocatable **parm::ft_fc**
- real *8 **parm::sfsedmean**
- real *8 **parm::sfsedstdev**
- integer, dimension(:), allocatable **parm::dtp_imo**
month the reservoir becomes operational (none)
- integer, dimension(:), allocatable **parm::dtp_jyr**
year of the simulation that the reservoir becomes operational (none)
- integer, dimension(:), allocatable **parm::dtp_numstage**
total number of stages in the weir (none)
- integer, dimension(:), allocatable **parm::dtp_numweir**
total number of weirs in the BMP (none)
- integer, dimension(:), allocatable **parm::dtp_onoff**
sub-basin detention pond is associated with (none)
- integer, dimension(:), allocatable **parm::dtp_reltype**
equations for stage-discharge relationship (none):
1=exponential function,
2=linear,
3=logarithmic,
4=cubic,
5=power
- integer, dimension(:), allocatable **parm::dtp_stagdis**
(none):
0=use weir/orifice discharge equation to calculate outflow,
1=use stage-dicharge relationship
- integer, dimension(:), allocatable **parm::dtp_subnum**
- real *8, dimension(:), allocatable **parm::cf**
this parameter controls the response of decomposition to the combined effect of soil temperature and moisture.
- real *8, dimension(:), allocatable **parm::cfh**
maximum humification rate
- real *8, dimension(:), allocatable **parm::cfdec**
the undisturbed soil turnover rate under optimum soil water and temperature. Increasing it will increase carbon and organic N decomp.
- real *8, dimension(:), allocatable **parm::lat_orgn**
- real *8, dimension(:), allocatable **parm::lat_orgp**
- integer, dimension(:,:), allocatable **parm::dtp_weirdim**
weir dimensions (none),
1=read user input,
0=use model calculation
- integer, dimension(:,:), allocatable **parm::dtp_weirtype**
type of weir (none):
1=rectangular and
2=circular
- real *8, dimension(:), allocatable **parm::dtp_coef1**
coefficient of 3rd degree in the polynomial equation (none)
- real *8, dimension(:), allocatable **parm::dtp_coef2**
coefficient of 2nd degree in the polynomial equation (none)
- real *8, dimension(:), allocatable **parm::dtp_coef3**
coefficient of 1st degree in the polynomial equation (none)
- real *8, dimension(:), allocatable **parm::dtp_evrsv**
detention pond evaporation coefficient (none)
- real *8, dimension(:), allocatable **parm::dtp_expont**
exponent used in the exponential equation (none)
- real *8, dimension(:), allocatable **parm::dtp_intcept**

- intercept used in regression equations (none)*
- real *8, dimension(:), allocatable [parm::dtp_lwratio](#)
- ratio of length to width of water back up (none)*
- real *8, dimension(:), allocatable [parm::dtp_totwrwid](#)
- total constructed width of the detention wall across the creek (m)*
- real *8, dimension(:), allocatable **parm::dtp_inflvol**
- real *8, dimension(:), allocatable **parm::dtp_wdep**
- real *8, dimension(:), allocatable **parm::dtp_totdep**
- real *8, dimension(:), allocatable **parm::dtp_watdepact**
- real *8, dimension(:), allocatable **parm::dtp_outflow**
- real *8, dimension(:), allocatable **parm::dtp_totrel**
- real *8, dimension(:), allocatable **parm::dtp_backoff**
- real *8, dimension(:), allocatable **parm::dtp_seep_sa**
- real *8, dimension(:), allocatable **parm::dtp_evap_sa**
- real *8, dimension(:), allocatable **parm::dtp_pet_day**
- real *8, dimension(:), allocatable **parm::dtp_pcpvol**
- real *8, dimension(:), allocatable **parm::dtp_seepvol**
- real *8, dimension(:), allocatable **parm::dtp_evapvol**
- real *8, dimension(:), allocatable **parm::dtp_flowin**
- real *8, dimension(:), allocatable **parm::dtp_backup_length**
- real *8, dimension(:), allocatable **parm::dtp_ivol**
- real *8, dimension(:), allocatable **parm::dtp_ised**
- integer, dimension(:,:), allocatable **parm::so_res_flag**
- integer, dimension(:,:), allocatable **parm::ro_bmp_flag**
- real *8, dimension(:,:), allocatable **parm::sol_watp**
- real *8, dimension(:,:), allocatable **parm::sol_solp_pre**
- real *8, dimension(:,:), allocatable **parm::psp_store**
- real *8, dimension(:,:), allocatable **parm::ssp_store**
- real *8, dimension(:,:), allocatable **parm::so_res**
- real *8, dimension(:,:), allocatable **parm::sol_cal**
- real *8, dimension(:,:), allocatable **parm::sol_ph**
- integer **parm::sol_p_model**
- integer, dimension(:,:), allocatable **parm::a_days**
- integer, dimension(:,:), allocatable **parm::b_days**
- real *8, dimension(:), allocatable [parm::min_res](#)
- minimum residue allowed due to implementation of residue managment in the OPS file (kg/ha)*
- real *8, dimension(:), allocatable **parm::harv_min**
- real *8, dimension(:), allocatable **parm::fstap**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_flo**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_sed**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_bac**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_pp**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_sp**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_pn**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_sn**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_flos**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_seds**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_bacs**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_pps**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_sps**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_pns**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_sns**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_flot**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_sedt**

- real *8, dimension(:,:), allocatable **parm::ro_bmp_bact**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_ppt**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_spt**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_pnt**
- real *8, dimension(:,:), allocatable **parm::ro_bmp_snt**
- real *8, dimension(:), allocatable **parm::bmp_flo**
- real *8, dimension(:), allocatable **parm::bmp_sed**
- real *8, dimension(:), allocatable **parm::bmp_bac**
- real *8, dimension(:), allocatable **parm::bmp_pp**
- real *8, dimension(:), allocatable **parm::bmp_sp**
- real *8, dimension(:), allocatable **parm::bmp_pn**
- real *8, dimension(:), allocatable **parm::bmp_sn**
- real *8, dimension(:), allocatable **parm::bmp_flag**
- real *8, dimension(:), allocatable **parm::bmp_flos**
- real *8, dimension(:), allocatable **parm::bmp_seds**
- real *8, dimension(:), allocatable **parm::bmp_bacs**
- real *8, dimension(:), allocatable **parm::bmp_pps**
- real *8, dimension(:), allocatable **parm::bmp_sps**
- real *8, dimension(:), allocatable **parm::bmp_pns**
- real *8, dimension(:), allocatable **parm::bmp_sns**
- real *8, dimension(:), allocatable **parm::bmp_flot**
- real *8, dimension(:), allocatable **parm::bmp_sedt**
- real *8, dimension(:), allocatable **parm::bmp_bact**
- real *8, dimension(:), allocatable **parm::bmp_ppt**
- real *8, dimension(:), allocatable **parm::bmp_spt**
- real *8, dimension(:), allocatable **parm::bmp_pnt**
- real *8, dimension(:), allocatable **parm::bmp_snt**
- real *8, dimension(:,:), allocatable **parm::dtp_addon**
the distance between spillway levels (m)
- real *8, dimension(:,:), allocatable **parm::dtp_cdis**
discharge coefficient for weir/orifice flow (none)
- real *8, dimension(:,:), allocatable **parm::dtp_depweir**
depth of rectangular weir at different stages (m)
- real *8, dimension(:,:), allocatable **parm::dtp_diaweir**
diameter of orifice hole at different stages (m)
- real *8, dimension(:,:), allocatable **parm::dtp_flowrate**
maximum discharge from each stage of the weir/hole (m^3/s)
- real *8, dimension(:,:), allocatable **parm::dtp_pcpret**
precipitation for different return periods (not used) (mm)
- real *8, dimension(:,:), allocatable **parm::dtp_retperd**
return period at different stages (years)
- real *8, dimension(:,:), allocatable **parm::dtp_wdratio**
width depth ratio of rectangular weirs (none)
- real *8, dimension(:,:), allocatable **parm::dtp_wrwid**
- real *8, dimension(:), allocatable **parm::ri_subkm**
- real *8, dimension(:), allocatable **parm::ri_totpvol**
- real *8, dimension(:), allocatable **parm::irmmdt**
- real *8, dimension(:,:), allocatable **parm::ri_sed**
- real *8, dimension(:,:), allocatable **parm::ri_fr**
- real *8, dimension(:,:), allocatable **parm::ri_dim**
- real *8, dimension(:,:), allocatable **parm::ri_im**
- real *8, dimension(:,:), allocatable **parm::ri_iy**
- real *8, dimension(:,:), allocatable **parm::ri_sa**

- real *8, dimension(:,:), allocatable **parm::ri_vol**
- real *8, dimension(:,:), allocatable **parm::ri_qi**
- real *8, dimension(:,:), allocatable **parm::ri_k**
- real *8, dimension(:,:), allocatable **parm::ri_dd**
- real *8, dimension(:,:), allocatable **parm::ri_evrsv**
- real *8, dimension(:,:), allocatable **parm::ri_dep**
- real *8, dimension(:,:), allocatable **parm::ri_ndt**
- real *8, dimension(:,:), allocatable **parm::ri_pmpvol**
- real *8, dimension(:,:), allocatable **parm::ri_sed_cumul**
- real *8, dimension(:,:), allocatable **parm::hrnopcp**
- real *8, dimension(:,:), allocatable **parm::ri_qloss**
- real *8, dimension(:,:), allocatable **parm::ri_pumpv**
- real *8, dimension(:,:), allocatable **parm::ri_sedi**
- character(len=4), dimension(:,:), allocatable **parm::ri_nirr**
- integer, dimension(:), allocatable **parm::num_ri**
- integer, dimension(:), allocatable **parm::ri_luflg**
- integer, dimension(:), allocatable **parm::num_noirr**
- integer, dimension(:), allocatable **parm::wtp_subnum**
- integer, dimension(:), allocatable **parm::wtp_onoff**
- integer, dimension(:), allocatable **parm::wtp_imo**
- integer, dimension(:), allocatable **parm::wtp_iyr**
- integer, dimension(:), allocatable **parm::wtp_dim**
- integer, dimension(:), allocatable **parm::wtp_stagdis**
- integer, dimension(:), allocatable **parm::wtp_sdtype**
- real *8, dimension(:), allocatable **parm::wtp_pvol**
- real *8, dimension(:), allocatable **parm::wtp_pdepth**
- real *8, dimension(:), allocatable **parm::wtp_sdslope**
- real *8, dimension(:), allocatable **parm::wtp_lenwidth**
- real *8, dimension(:), allocatable **parm::wtp_extdepth**
- real *8, dimension(:), allocatable **parm::wtp_hydeff**
- real *8, dimension(:), allocatable **parm::wtp_evrsv**
- real *8, dimension(:), allocatable **parm::wtp_sdintc**
- real *8, dimension(:), allocatable **parm::wtp_sdexp**
- real *8, dimension(:), allocatable **parm::wtp_sdc1**
- real *8, dimension(:), allocatable **parm::wtp_sdc2**
- real *8, dimension(:), allocatable **parm::wtp_sdc3**
- real *8, dimension(:), allocatable **parm::wtp_pdia**
- real *8, dimension(:), allocatable **parm::wtp_plen**
- real *8, dimension(:), allocatable **parm::wtp_pmann**
- real *8, dimension(:), allocatable **parm::wtp_ploss**
- real *8, dimension(:), allocatable **parm::wtp_k**
- real *8, dimension(:), allocatable **parm::wtp_dp**
- real *8, dimension(:), allocatable **parm::wtp_sedi**
- real *8, dimension(:), allocatable **parm::wtp_sede**
- real *8, dimension(:), allocatable **parm::wtp_qi**
- real *8 **parm::lai_init**
initial leaf area index of transplants
- real *8 **parm::bio_init**
initial biomass of transplants (kg/ha)
- real *8 **parm::cnop**
SCS runoff curve number for moisture condition II (none)
- real *8 **parm::harveff**
harvest efficiency: fraction of harvested yield that is removed from HRU; the remainder becomes residue on the soil surface(none)

- real *8 **parm::hi_ovr**
harvest index target specified at harvest ((kg/ha)/(kg/ha))
- real *8 **parm::frac_harvk**
- real *8 **parm::lid_vgcl**
van Genuchten equation's coefficient, l (none)
- real *8 **parm::lid_vgcm**
van Genuchten equation's coefficient, m (none)
- real *8 **parm::lid_qsurf_total**
- real *8 **parm::lid_farea_sum**
- real *8, dimension(:,:), allocatable **parm::lid_cuminf_last**
cumulative amount of water infiltrated into the amended soil layer at the last time step in a day (mm H2O)
- real *8, dimension(:,:), allocatable **parm::lid_cumr_last**
cumulative amount of rainfall at the last time step in a day (mm H2O)
- real *8, dimension(:,:), allocatable **parm::lid_excum_last**
cumulative amount of excess rainfall at the last time step in a day (mm H2O)
- real *8, dimension(:,:), allocatable **parm::lid_f_last**
potential infiltration rate of the amended soil layer at the last time step in a day (mm/mm H2O)
- real *8, dimension(:,:), allocatable **parm::lid_sw_last**
soil water content of the amended soil layer at the last time step in a day (mm/mm H2O)
- real *8, dimension(:,:), allocatable **parm::lid_qsurf**
depth of runoff generated on a LID in a given time interval (mm H2O)
- real *8, dimension(:,:), allocatable **parm::interval_last**
- real *8, dimension(:,:), allocatable **parm::lid_str_last**
- real *8, dimension(:,:), allocatable **parm::lid_farea**
- real *8, dimension(:,:), allocatable **parm::lid_sw_add**
- real *8, dimension(:,:), allocatable **parm::lid_cumqperc_last**
- real *8, dimension(:,:), allocatable **parm::lid_cumirr_last**
- integer, dimension(:,:), allocatable **parm::gr_onoff**
- integer, dimension(:,:), allocatable **parm::gr_imo**
- integer, dimension(:,:), allocatable **parm::gr_iyr**
- real *8, dimension(:,:), allocatable **parm::gr_farea**
fractional area of a green roof to the HRU (none)
- real *8, dimension(:,:), allocatable **parm::gr_solop**
- real *8, dimension(:,:), allocatable **parm::gr_etcoef**
- real *8, dimension(:,:), allocatable **parm::gr_fc**
- real *8, dimension(:,:), allocatable **parm::gr_wp**
- real *8, dimension(:,:), allocatable **parm::gr_ksat**
- real *8, dimension(:,:), allocatable **parm::gr_por**
- real *8, dimension(:,:), allocatable **parm::gr_hydeff**
- real *8, dimension(:,:), allocatable **parm::gr_soldpt**
- integer, dimension(:,:), allocatable **parm::rg_onoff**
- integer, dimension(:,:), allocatable **parm::rg_imo**
- integer, dimension(:,:), allocatable **parm::rg_iyr**
- real *8, dimension(:,:), allocatable **parm::rg_farea**
- real *8, dimension(:,:), allocatable **parm::rg_solop**
- real *8, dimension(:,:), allocatable **parm::rg_etcoef**
- real *8, dimension(:,:), allocatable **parm::rg_fc**
- real *8, dimension(:,:), allocatable **parm::rg_wp**
- real *8, dimension(:,:), allocatable **parm::rg_ksat**
- real *8, dimension(:,:), allocatable **parm::rg_por**
- real *8, dimension(:,:), allocatable **parm::rg_hydeff**
- real *8, dimension(:,:), allocatable **parm::rg_soldpt**
- real *8, dimension(:,:), allocatable **parm::rg_dimop**

- real *8, dimension(:,:), allocatable **parm::rg_sarea**
- real *8, dimension(:,:), allocatable **parm::rg_vol**
- real *8, dimension(:,:), allocatable **parm::rg_sth**
- real *8, dimension(:,:), allocatable **parm::rg_sdia**
- real *8, dimension(:,:), allocatable **parm::rg_bdia**
- real *8, dimension(:,:), allocatable **parm::rg_sts**
- real *8, dimension(:,:), allocatable **parm::rg_orifice**
- real *8, dimension(:,:), allocatable **parm::rg_oheight**
- real *8, dimension(:,:), allocatable **parm::rg_odia**
- integer, dimension(:,:), allocatable **parm::cs_onoff**
- integer, dimension(:,:), allocatable **parm::cs_imo**
- integer, dimension(:,:), allocatable **parm::cs_iyr**
- integer, dimension(:,:), allocatable **parm::cs_grcon**
- real *8, dimension(:,:), allocatable **parm::cs_farea**
- real *8, dimension(:,:), allocatable **parm::cs_vol**
- real *8, dimension(:,:), allocatable **parm::cs_rdepth**
- integer, dimension(:,:), allocatable **parm::pv_onoff**
- integer, dimension(:,:), allocatable **parm::pv_imo**
- integer, dimension(:,:), allocatable **parm::pv_iyr**
- integer, dimension(:,:), allocatable **parm::pv_solop**
- real *8, dimension(:,:), allocatable **parm::pv_grvdep**
- real *8, dimension(:,:), allocatable **parm::pv_grvpor**
- real *8, dimension(:,:), allocatable **parm::pv_farea**
- real *8, dimension(:,:), allocatable **parm::pv_drcoef**
- real *8, dimension(:,:), allocatable **parm::pv_fc**
- real *8, dimension(:,:), allocatable **parm::pv_wp**
- real *8, dimension(:,:), allocatable **parm::pv_ksat**
- real *8, dimension(:,:), allocatable **parm::pv_por**
- real *8, dimension(:,:), allocatable **parm::pv_hydeff**
- real *8, dimension(:,:), allocatable **parm::pv_soldpt**
- integer, dimension(:,:), allocatable **parm::lid_onoff**
- real *8, dimension(:,:), allocatable **parm::sol_bmc**
- real *8, dimension(:,:), allocatable **parm::sol_bmn**
- real *8, dimension(:,:), allocatable **parm::sol_hsc**
- real *8, dimension(:,:), allocatable **parm::sol_hsn**
- real *8, dimension(:,:), allocatable **parm::sol_hpc**
- real *8, dimension(:,:), allocatable **parm::sol_hpn**
- real *8, dimension(:,:), allocatable **parm::sol_lm**
- real *8, dimension(:,:), allocatable **parm::sol_lmc**
- real *8, dimension(:,:), allocatable **parm::sol_lmn**
- real *8, dimension(:,:), allocatable **parm::sol_ls**
- real *8, dimension(:,:), allocatable **parm::sol_lsl**
- real *8, dimension(:,:), allocatable **parm::sol_lsc**
- real *8, dimension(:,:), allocatable **parm::sol_lsn**
- real *8, dimension(:,:), allocatable **parm::sol_rnmn**
- real *8, dimension(:,:), allocatable **parm::sol_lslc**
- real *8, dimension(:,:), allocatable **parm::sol_lslnc**
- real *8, dimension(:,:), allocatable **parm::sol_rspc**
- real *8, dimension(:,:), allocatable **parm::sol_woc**
- real *8, dimension(:,:), allocatable **parm::sol_won**
- real *8, dimension(:,:), allocatable **parm::sol_hp**
- real *8, dimension(:,:), allocatable **parm::sol_hs**
- real *8, dimension(:,:), allocatable **parm::sol_bm**
- real *8, dimension(:,:), allocatable **parm::sol_cac**
- real *8, dimension(:,:), allocatable **parm::sol_cec**

- real *8, dimension(:,:), allocatable **parm::sol_percc**
- real *8, dimension(:,:), allocatable **parm::sol_latc**
- real *8, dimension(:), allocatable **parm::sedc_d**
- real *8, dimension(:), allocatable **parm::surfqc_d**
- real *8, dimension(:), allocatable **parm::latc_d**
- real *8, dimension(:), allocatable **parm::percc_d**
- real *8, dimension(:), allocatable **parm::foc_d**
- real *8, dimension(:), allocatable **parm::nppc_d**
- real *8, dimension(:), allocatable **parm::rsdc_d**
- real *8, dimension(:), allocatable **parm::grainc_d**
- real *8, dimension(:), allocatable **parm::stoverc_d**
- real *8, dimension(:), allocatable **parm::soc_d**
- real *8, dimension(:), allocatable **parm::rspc_d**
- real *8, dimension(:), allocatable **parm::emitc_d**
- real *8, dimension(:), allocatable **parm::sub_sedc_d**
- real *8, dimension(:), allocatable **parm::sub_surfqc_d**
- real *8, dimension(:), allocatable **parm::sub_latc_d**
- real *8, dimension(:), allocatable **parm::sub_percc_d**
- real *8, dimension(:), allocatable **parm::sub_foc_d**
- real *8, dimension(:), allocatable **parm::sub_nppc_d**
- real *8, dimension(:), allocatable **parm::sub_rsdcd**
- real *8, dimension(:), allocatable **parm::sub_grainc_d**
- real *8, dimension(:), allocatable **parm::sub_stoverc_d**
- real *8, dimension(:), allocatable **parm::sub_emitc_d**
- real *8, dimension(:), allocatable **parm::sub_soc_d**
- real *8, dimension(:), allocatable **parm::sub_rspc_d**
- real *8, dimension(:), allocatable **parm::sedc_m**
- real *8, dimension(:), allocatable **parm::surfqc_m**
- real *8, dimension(:), allocatable **parm::latc_m**
- real *8, dimension(:), allocatable **parm::percc_m**
- real *8, dimension(:), allocatable **parm::foc_m**
- real *8, dimension(:), allocatable **parm::nppc_m**
- real *8, dimension(:), allocatable **parm::rsdc_m**
- real *8, dimension(:), allocatable **parm::grainc_m**
- real *8, dimension(:), allocatable **parm::stoverc_m**
- real *8, dimension(:), allocatable **parm::emitc_m**
- real *8, dimension(:), allocatable **parm::soc_m**
- real *8, dimension(:), allocatable **parm::rspc_m**
- real *8, dimension(:), allocatable **parm::sedc_a**
- real *8, dimension(:), allocatable **parm::surfqc_a**
- real *8, dimension(:), allocatable **parm::latc_a**
- real *8, dimension(:), allocatable **parm::percc_a**
- real *8, dimension(:), allocatable **parm::foc_a**
- real *8, dimension(:), allocatable **parm::nppc_a**
- real *8, dimension(:), allocatable **parm::rsdc_a**
- real *8, dimension(:), allocatable **parm::grainc_a**
- real *8, dimension(:), allocatable **parm::stoverc_a**
- real *8, dimension(:), allocatable **parm::emitc_a**
- real *8, dimension(:), allocatable **parm::soc_a**
- real *8, dimension(:), allocatable **parm::rspc_a**
- integer, dimension(:), allocatable **parm::tillage_switch**
- real *8, dimension(:), allocatable **parm::tillage_depth**
- integer, dimension(:), allocatable **parm::tillage_days**
- real *8, dimension(:), allocatable **parm::tillage_factor**
- real *8 **parm::dthy**

time interval for subdaily routing

- integer, dimension(4) **parm::ihx**
- integer, dimension(:), allocatable **parm::nhy**
- real *8, dimension(:), allocatable **parm::rchx**
- real *8, dimension(:), allocatable **parm::rcss**
- real *8, dimension(:), allocatable **parm::qcap**
- real *8, dimension(:), allocatable **parm::chxa**
- real *8, dimension(:), allocatable **parm::chxp**
- real *8, dimension(:, :, :), allocatable **parm::qhy**
- real *8 **parm::ff1**
- real *8 **parm::ff2**

7.57.1 Detailed Description

file containing the module parm

Author

modified by Javier Burguete Tolosa

7.58 newtillmix.f90 File Reference

Functions/Subroutines

- subroutine [newtillmix](#) (j, bmix)

this subroutine mixes residue and nutrients during tillage and biological mixing. Mixing was extended to all layers. A subroutine to simulate stimulation of organic matter decomposition was added. March 2009: testing has been minimal and further adjustments are expected. Use with caution!

7.58.1 Detailed Description

file containing the subroutine newtillmix

Author

Armen R. Kemanian,
Stefan Julich,
Cole Rossi
modified by Javier Burguete

7.58.2 Function/Subroutine Documentation

7.58.2.1 newtillmix()

```
subroutine newtillmix (
    integer, intent(in) j,
    real*8, intent(in) bmix )
```

this subroutine mixes residue and nutrients during tillage and biological mixing. Mixing was extended to all layers. A subroutine to simulate stimulation of organic matter decomposition was added. March 2009: testing has been minimal and further adjustments are expected. Use with caution!

Parameters

| | | |
|----|-------------|---|
| in | <i>j</i> | HRU number (none) |
| in | <i>bmix</i> | biological mixing efficiency: this number is zero for tillage operations (none) |

7.59 openwth.f90 File Reference

Functions/Subroutines

- subroutine [openwth](#)
this subroutine opens the precipitation, temperature, solar radiation, relative humidity and wind speed files for simulations using measured weather data

7.59.1 Detailed Description

file containing the subroutine openwth

Author

modified by Javier Burguete

7.60 ovr_sed.f90 File Reference

Functions/Subroutines

- subroutine [ovr_sed](#) ()
this subroutine computes splash erosion by raindrop impact and flow erosion by overland flow

7.60.1 Detailed Description

file containing the subroutine ovr_sed

Author

modified by Javier Burguete

7.61 pgen.f90 File Reference

Functions/Subroutines

- subroutine [pgen](#) (j)
this subroutine generates precipitation data when the user chooses to simulate or when data is missing for particular days in the weather file

7.61.1 Detailed Description

file containing the subroutine pgen

Author

modified by Javier Burguete

7.61.2 Function/Subroutine Documentation

7.61.2.1 pgen()

```
subroutine pgen (
    integer, intent(in) j )
```

this subroutine generates precipitation data when the user chooses to simulate or when data is missing for particular days in the weather file

Parameters

| | | |
|----|---|------------|
| in | j | HRU number |
|----|---|------------|

7.62 pgenhr.f90 File Reference

Functions/Subroutines

- subroutine [pgenhr](#) (jj)
this subroutine distributes daily rainfall exponentially within the day @parameter[in] jj HRU number

7.62.1 Detailed Description

file containing the subroutine pgenhr

Author

modified by Javier Burguete

7.63 pkq.f90 File Reference

Functions/Subroutines

- subroutine [pkq](#) (iwave, j)
*this subroutine computes the peak runoff rate for each HRU and the entire subbasin using a modification of the rational formula @parm[in] iwave flag to differentiate calculation of HRU and subbasin sediment calculation (none)
iwave = 0 for HRU MUSLE(sedyld) each hru is calculated independently using hru area and adjusted channel length
iwave = 1 subbasin # for subbasin MUSLE is computed for entire subbasin using hru weighted KLSCP*

7.63.1 Detailed Description

file containing the subroutine pkq

Author

modified by Javier Burguete

7.63.2 Function/Subroutine Documentation

7.63.2.1 pkq()

```
subroutine pkq (
    integer, intent(in) iwave,
    integer, intent(in) j )
```

this subroutine computes the peak runoff rate for each HRU and the entire subbasin using a modification of the rational formula @parm[in] iwave flag to differentiate calculation of HRU and subbasin sediment calculation (none) iwave = 0 for HRU MUSLE(sedyld) each hru is calculated independently using hru area and adjusted channel length iwave = 1 subbasin # for subbasin MUSLE is computed for entire subbasin using hru weighted KLSCP

Parameters

| | | |
|----|---|-------------------|
| in | j | HRU number (none) |
|----|---|-------------------|

7.64 plantop.f90 File Reference

Functions/Subroutines

- subroutine [plantop](#) (j)
this subroutine performs the plant operation

7.64.1 Detailed Description

file containing the subroutine plantop

Author

modified by Javier Burguete

7.64.2 Function/Subroutine Documentation

7.64.2.1 plantop()

```
subroutine plantop (
    integer, intent(in) j )
```

this subroutine performs the plant operation

Parameters

| | | |
|----|----------|------------|
| in | <i>j</i> | HRU number |
|----|----------|------------|

7.65 pmeas.f90 File Reference**Functions/Subroutines**

- subroutine [pmeas](#) (i)
this subroutine reads in precipitation data and assigns it to the proper subbasins

7.65.1 Detailed Description

file containing the subroutine pmeas

Author

modified by Javier Burguete

7.65.2 Function/Subroutine Documentation**7.65.2.1 pmeas()**

```
subroutine pmeas (
    integer, intent(in) i )
```

this subroutine reads in precipitation data and assigns it to the proper subbasins

Parameters

| | | |
|----|----------|---|
| in | <i>i</i> | current day of simulation (julian date) |
|----|----------|---|

7.66 qman.f90 File Reference

Functions/Subroutines

- real*8 function `qman` (`x1`, `x2`, `x3`, `x4`)

this subroutine calculates flow rate or flow velocity using Manning's equation. If `x1` is set to 1, the velocity is calculated. If `x1` is set to cross-sectional area of flow, the flow rate is calculated.

7.66.1 Detailed Description

file containing the function `qman`

Author

modified by Javier Burguete

7.66.2 Function/Subroutine Documentation

7.66.2.1 `qman()`

```
real*8 function qman (
    real*8, intent(in) x1,
    real*8, intent(in) x2,
    real*8, intent(in) x3,
    real*8, intent(in) x4 )
```

this subroutine calculates flow rate or flow velocity using Manning's equation. If `x1` is set to 1, the velocity is calculated. If `x1` is set to cross-sectional area of flow, the flow rate is calculated.

Parameters

| | | |
|----|-----------------|---|
| in | <code>x1</code> | cross-sectional flow area or 1 (m ² or none) |
| in | <code>x2</code> | hydraulic radius (m) |
| in | <code>x3</code> | Manning's "n" value for channel (none) |
| in | <code>x4</code> | average slope of channel (m/m) |

Returns

flow rate or flow velocity (m³/s or m/s)

7.67 rchaa.f90 File Reference

Functions/Subroutines

- subroutine `rchaa` (years)

this subroutine writes the average annual reach output to the .rch file

7.67.1 Detailed Description

file containing the subroutine rchaa

Author

modified by Javier Burguete

7.67.2 Function/Subroutine Documentation

7.67.2.1 rchaa()

```
subroutine rchaa (
    real*8, intent(in) years )
```

this subroutine writes the average annual reach output to the .rch file

Parameters

| | | |
|----|-------|------------------------------|
| in | years | length of simulation (years) |
|----|-------|------------------------------|

7.68 rchday.f90 File Reference

Functions/Subroutines

- subroutine [rchday](#)
this subroutine writes the daily reach output to the .rch file

7.68.1 Detailed Description

file containing the subroutine rchday

Author

modified by Javier Burguete

7.69 rchmon.f90 File Reference

Functions/Subroutines

- subroutine [rchmon](#) (mdays)
this subroutine writes the monthly reach output to the .rch file

7.69.1 Detailed Description

file containing the subroutine `rchmon`

Author

modified by Javier Burguete

7.69.2 Function/Subroutine Documentation

7.69.2.1 `rchmon()`

```
subroutine rchmon (
    integer, intent(in) mdays )
```

this subroutine writes the monthly reach output to the `.rch` file

Parameters

| | | |
|-----------------|--------------------|-----------------------------------|
| <code>in</code> | <code>mdays</code> | number of days simulated in month |
|-----------------|--------------------|-----------------------------------|

7.70 `rchyr.f90` File Reference

Functions/Subroutines

- subroutine `rchyr` (i)
this subroutine writes the annual reach output to the `.rch` file

7.70.1 Detailed Description

file containing the subroutine `rchyr`

Author

modified by Javier Burguete

7.70.2 Function/Subroutine Documentation

7.70.2.1 `rchyr()`

```
subroutine rchyr (
    integer, intent(in) i )
```

this subroutine writes the annual reach output to the `.rch` file

Parameters

| | | |
|----|---------|---------------------------------|
| in | current | day of simulation (julian date) |
|----|---------|---------------------------------|

7.71 readatmodep.f90 File Reference

Functions/Subroutines

- subroutine [readatmodep](#)
this subroutine reads the atmospheric deposition values

7.71.1 Detailed Description

file containing the subroutine readatmodep

Author

modified by Javier Burguete

7.72 readbsn.f90 File Reference

Functions/Subroutines

- subroutine [readbsn](#)
this subroutine reads data from the basin input file (.bsn). This file contains information related to processes modeled or defined at the watershed level

7.72.1 Detailed Description

file containing the suborutine readbsn

Author

modified by Javier Burguete

7.73 readchm.f90 File Reference

Functions/Subroutines

- subroutine [readchm](#) (l)
This subroutine reads data from the HRU/subbasin soil chemical input file (.chm). This file contains initial amounts of pesticides/nutrients in the first soil layer. (Specifics about the first soil layer are given in the .sol file.) All data in the .chm file is optional input.

7.73.1 Detailed Description

file containing the subroutine readchm

Author

modified by Javier Burguete

7.73.2 Function/Subroutine Documentation

7.73.2.1 readchm()

```
subroutine readchm (
    integer, intent(in) l )
```

This subroutine reads data from the HRU/subbasin soil chemical input file (.chm). This file contains initial amounts of pesticides/nutrients in the first soil layer. (Specifics about the first soil layer are given in the .sol file.) All data in the .chm file is optional input.

Parameters

| | | |
|----|---|-------------------|
| in | / | HRU number (none) |
|----|---|-------------------|

7.74 readcnst.f90 File Reference

Functions/Subroutines

- subroutine [readcnst](#) (jj)
reads in the loading information for the reccnst command

7.74.1 Detailed Description

file containing the subroutine [readcnst.f90](#)

Author

modified by Javier Burguete

7.74.2 Function/Subroutine Documentation

7.74.2.1 readcnst()

```
subroutine readcnst (
    integer, intent(in) jj )
```

reads in the loading information for the reccnst command

Parameters

| | | |
|----|----|---|
| in | jj | file number associated with recnst command (none) |
|----|----|---|

7.75 readfcst.f90 File Reference

Functions/Subroutines

- subroutine [readfcst](#)
this subroutine reads the HRU forecast weather generator parameters from the .cst file

7.75.1 Detailed Description

file containing the subroutine readfcst

Author

modified by Javier Burguete

7.76 readfert.f90 File Reference

Functions/Subroutines

- subroutine [readfert](#)
this subroutine reads input parameters from the fertilizer/manure (i.e. nutrient) database (fert.dat)

7.76.1 Detailed Description

file containing the subroutine readfert

Author

modified by Javier Burguete

7.77 readfig.f90 File Reference

Functions/Subroutines

- subroutine [readfig](#)
reads in the routing information from the watershed configuration input file (.fig) and calculates the number of sub-basins, reaches, and reservoirs

7.77.1 Detailed Description

file containing the subroutine readfig

Author

modified by Javier Burguete

7.78 readfile.f90 File Reference

Functions/Subroutines

- subroutine [readfile](#)

this subroutine opens the main input and output files and reads watershed information from the file.cio

7.78.1 Detailed Description

file containing the subroutine readfile

Author

modified by Javier Burguete

7.79 readgw.f90 File Reference

Functions/Subroutines

- subroutine [readgw](#) (i, j)

this subroutine reads the parameters from the HRU/subbasin groundwater input file (.gw)

7.79.1 Detailed Description

file containing the subroutine readgw

Author

modified by Javier Burguete

7.79.2 Function/Subroutine Documentation

7.79.2.1 readgw()

```
subroutine readgw (  
    integer, intent(in) i,  
    integer, intent(in) j )
```

this subroutine reads the parameters from the HRU/subbasin groundwater input file (.gw)

Parameters

| | | |
|----|----------|------------------------|
| in | <i>i</i> | subbasin number (none) |
| in | <i>j</i> | HRU number (none) |

7.80 readhru.f90 File Reference

Functions/Subroutines

- subroutine [readhru](#) (*i*, *j*)

this subroutine reads data from the HRU general input file (.hru). This file contains data related to general processes modeled at the HRU level.

7.80.1 Detailed Description

file containing the subroutine readhru

Author

modified by Javier Burguete

7.80.2 Function/Subroutine Documentation

7.80.2.1 readhru()

```
subroutine readhru (
    integer, intent(in) i,
    integer, intent(in) j )
```

this subroutine reads data from the HRU general input file (.hru). This file contains data related to general processes modeled at the HRU level.

Parameters

| | | |
|----|----------|------------------------|
| in | <i>i</i> | subbasin number (none) |
| in | <i>j</i> | HRU number (none) |

7.81 readinpt.f90 File Reference

Functions/Subroutines

- subroutine [readinpt](#)

this subroutine calls subroutines which read input data for the databases and the HRUs

7.81.1 Detailed Description

file containing the subroutine readinpt

Author

modified by Javier Burguete

7.82 readlup.f90 File Reference

Functions/Subroutines

- subroutine [readlup](#)
this subroutine reads data from the HRU/subbasin management input file (.mgt). This file contains data related to management practices used in the HRU/subbasin.

7.82.1 Detailed Description

file containing the subroutine readlup

Author

modified by Javier Burguete

7.83 readlwq.f90 File Reference

Functions/Subroutines

- subroutine [readlwq](#) (ii)
this subroutine reads data from the lake water quality input file (.lwq). This file contains data related to initial pesticide and nutrient levels in the lake/reservoir and transformation processes occurring within the lake/reservoir. Data in the lake water quality input file is assumed to apply to all reservoirs in the watershed.

7.83.1 Detailed Description

file containing the subroutine readlwq

Author

modified by Javier Burguete

7.83.2 Function/Subroutine Documentation

7.83.2.1 readlwq()

```
subroutine readlwq (
    integer, intent(in) ii )
```

this subroutine reads data from the lake water quality input file (.lwq). This file contains data related to initial pesticide and nutrient levels in the lake/reservoir and transformation processes occurring within the lake/reservoir. Data in the lake water quality input file is assumed to apply to all reservoirs in the watershed.

Parameters

| | | |
|----|----|-------------------------|
| in | ii | reservoir number (none) |
|----|----|-------------------------|

7.84 readmgt.f90 File Reference

Functions/Subroutines

- subroutine [readmgt](#) (k)

this subroutine reads data from the HRU/subbasin management input file (.mgt). This file contains data related to management practices used in the HRU/subbasin.

7.84.1 Detailed Description

file containing the subroutine readmgt

Author

modified by Javier Burguete

7.84.2 Function/Subroutine Documentation

7.84.2.1 readmgt()

```
subroutine readmgt (
    integer, intent(in) k )
```

this subroutine reads data from the HRU/subbasin management input file (.mgt). This file contains data related to management practices used in the HRU/subbasin.

Parameters

| | | |
|----|---|-------------------|
| in | k | HRU number (none) |
|----|---|-------------------|

7.85 readmon.f90 File Reference

Functions/Subroutines

- subroutine [readmon](#) (i)

reads in the input data for the recmon command

7.85.1 Detailed Description

file containing the subroutine readmon

Author

modified by Javier Burguete

7.86 readops.f90 File Reference

Functions/Subroutines

- subroutine [readops](#) (k)

this subroutine reads data from the HRU/subbasin management input file (.mgt). This file contains data related to management practices used in the HRU/subbasin.

7.86.1 Detailed Description

file containing the subroutine readops

Author

modified by Javier Burguete

7.86.2 Function/Subroutine Documentation

7.86.2.1 readops()

```
subroutine readops (
    integer, intent(in) k )
```

this subroutine reads data from the HRU/subbasin management input file (.mgt). This file contains data related to management practices used in the HRU/subbasin.

Parameters

| | | |
|----|---|-------------------|
| in | k | HRU number (none) |
|----|---|-------------------|

7.87 readpest.f90 File Reference

Functions/Subroutines

- subroutine [readpest](#)

this subroutine reads parameters from the toxin/pesticide database (pest.dat)

7.87.1 Detailed Description

file containing the subroutine readpest

Author

modified by Javier Burguete

7.88 readplant.f90 File Reference

Functions/Subroutines

- subroutine [readplant](#)

this subroutine reads input parameters from the landuse/landcover database (plant.dat)

7.88.1 Detailed Description

file containing the subroutine readplant

Author

modified by Javier Burguete

7.89 readpnd.f90 File Reference

Functions/Subroutines

- subroutine [readpnd](#) (i)

This subroutine reads data from the HRU/subbasin pond input file (.pnd). This file contains data related to ponds and wetlands in the HRUs/subbasins.

7.89.1 Detailed Description

file containing the subroutine readpnd

Author

modified by Javier Burguete

7.89.2 Function/Subroutine Documentation

7.89.2.1 readpnd()

```
subroutine readpnd (
    integer, intent(in) i )
```

This subroutine reads data from the HRU/subbasin pond input file (.pnd). This file contains data related to ponds and wetlands in the HRUs/subbasins.

Parameters

| | | |
|----|---|------------------------|
| in | i | subbasin number (none) |
|----|---|------------------------|

7.90 readres.f90 File Reference

Functions/Subroutines

- subroutine [readres](#) (i)
the purpose of this subroutine is to read in data from the reservoir input file (.res)

7.90.1 Detailed Description

file containing the subroutine readres

Author

modified by Javier Burguete

7.90.2 Function/Subroutine Documentation

7.90.2.1 readres()

```
subroutine readres (
    integer, intent(in) i )
```

the purpose of this subroutine is to read in data from the reservoir input file (.res)

Parameters

| | | |
|----|---|-------------------------|
| in | i | reservoir number (none) |
|----|---|-------------------------|

7.91 readrte.f90 File Reference

Functions/Subroutines

- subroutine [readrte](#)
this subroutine reads data from the reach (main channel) input file (.rte). This file contains data related to channel attributes. Only one reach file should be made for each subbasin. If multiple HRUs are modeled within a subbasin, the same .rte file should be listed for all HRUs in file.cio

7.91.1 Detailed Description

file containing the subroutine readrte

Author

modified by Javier Burguete

7.92 readru.f90 File Reference

Functions/Subroutines

- subroutine [readru](#) (i)
this subroutine reads data from the sub input file (.sub). This file contains data related to routing

7.92.1 Detailed Description

file containing the subroutine readru

Author

modified by Javier Burguete

7.92.2 Function/Subroutine Documentation

7.92.2.1 readru()

```
subroutine readru (
    integer, intent(in) i )
```

this subroutine reads data from the sub input file (.sub). This file contains data related to routing

Parameters

| | | |
|----|---|-----------------|
| in | i | subbasin number |
|----|---|-----------------|

7.93 readsdr.f90 File Reference

Functions/Subroutines

- subroutine [readsdr](#) (j)
this subroutine reads data from the HRU/subbasin management input file (.mgt). This file contains data related to management practices used in the HRU/subbasin.

7.93.1 Detailed Description

file containing the subroutine readsdr

Author

modified by Javier Burguete

7.93.2 Function/Subroutine Documentation

7.93.2.1 readsdr()

```
subroutine readsdr (
    integer j )
```

this subroutine reads data from the HRU/subbasin management input file (.mgt). This file contains data related to management practices used in the HRU/subbasin.

Parameters

| | | |
|----|---|-------------------|
| in | j | HRU number (none) |
|----|---|-------------------|

7.94 readsepticbz.f90 File Reference

Functions/Subroutines

- subroutine [readsepticbz](#) (j)

this subroutine reads data from the septic input file (.sep). This file contains information related to septic tanks modeled or defined at the watershed level

7.94.1 Detailed Description

file containing the subroutine readsepticbz

Author

modified by Javier Burguete

7.94.2 Function/Subroutine Documentation

7.94.2.1 readsepticbz()

```
subroutine readsepticbz (  
    integer, intent(in) j )
```

this subroutine reads data from the septic input file (.sep). This file contains information related to septic tanks modeled or defined at the watershed level

Parameters

| | | |
|-----------------|----------------|-------------------|
| <code>in</code> | <code>j</code> | HRU number (none) |
|-----------------|----------------|-------------------|

7.95 readseptwq.f90 File Reference

Functions/Subroutines

- subroutine [readseptwq](#)

this subroutine reads input parameters from the sept wq database (septwq.dat). Information is used when a hru has septic tank.

7.95.1 Detailed Description

file containing the subroutine readseptwq

Author

C. Santhi, modified by Javier Burguete

7.95.2 Function/Subroutine Documentation

7.95.2.1 readseptwq()

```
subroutine readseptwq ( )
```

this subroutine reads input parameters from the sept wq database (septwq.dat). Information is used when a hru has septic tank.

This routine was developed by C. Santhi. Inputs for this routine are provided in septwq.dat of septic documentation. Data were compiled from [3] and [2].

7.96 readsno.f90 File Reference

Functions/Subroutines

- subroutine [readsno](#) (i)

this subroutine reads snow data from the HRU/subbasin soil chemical input

7.96.1 Detailed Description

file containing the subroutine readsno

Author

modified by Javier Burguete

7.96.2 Function/Subroutine Documentation

7.96.2.1 readsno()

```
subroutine readsno (  
    integer, intent(in) i )
```

this subroutine reads snow data from the HRU/subbasin soil chemical input

Parameters

| | | |
|----|---|------------------------|
| in | i | subbasin number (none) |
|----|---|------------------------|

7.97 readsol.f90 File Reference

Functions/Subroutines

- subroutine [readsol](#) (k)

this subroutine reads data from the HRU/subbasin soil properties file (.sol). This file contains data related to soil physical properties and general chemical properties

7.97.1 Detailed Description

file containing the subroutine readsol

Author

modified by Javier Burguete

7.97.2 Function/Subroutine Documentation

7.97.2.1 readsol()

```
subroutine readsol (  
    integer, intent(in) k )
```

this subroutine reads data from the HRU/subbasin soil properties file (.sol). This file contains data related to soil physical properties and general chemical properties

Parameters

| | | |
|----|---|------------|
| in | k | HRU number |
|----|---|------------|

7.98 readsub.f90 File Reference

Functions/Subroutines

- subroutine [readsub](#) (i)

this subroutine reads data from the HRU/subbasin general input file (.sub). This file contains data related to general processes modeled at the HRU/subbasin level.

7.98.1 Detailed Description

file containing the subroutine readsub

Author

modified by Javier Burguete

7.98.2 Function/Subroutine Documentation

7.98.2.1 readsub()

```
subroutine readsub (
    integer, intent(in) i )
```

this subroutine reads data from the HRU/subbasin general input file (.sub). This file contains data related to general processes modeled at the HRU/subbasin level.

Parameters

| | | |
|----|---|------------------------|
| in | i | subbasin number (none) |
|----|---|------------------------|

7.99 readswq.f90 File Reference

Functions/Subroutines

- subroutine [readswq](#)

this subroutine reads parameters from the subbasin instream water quality file (.swq) and initializes the QUAL2E variables which apply to the individual subbasins

7.99.1 Detailed Description

file containing the subroutine readswq

Author

modified by Javier Burguete

7.100 readtill.f90 File Reference

Functions/Subroutines

- subroutine [readtill](#)
this subroutine reads input data from tillage database (till.dat)

7.100.1 Detailed Description

file containing the subroutine readtill

Author

modified by Javier Burguete

7.101 readurban.f90 File Reference

Functions/Subroutines

- subroutine [readurban](#)
this subroutine reads input parameters from the urban database (urban.dat). Information from this database is used only if the urban buildup/washoff routines are selected for the modeling of urban areas

7.101.1 Detailed Description

file containing the subroutine readurban

Author

modified by Javier Burguete

7.102 readwgn.f90 File Reference

Functions/Subroutines

- subroutine [readwgn](#) (ii)
this subroutine reads the HRU weather generator parameters from the .wgn file

7.102.1 Detailed Description

file containing the subroutine readwgn

Author

modified by Javier Burguete

7.102.2 Function/Subroutine Documentation

7.102.2.1 readwgn()

```
subroutine readwgn (
    integer, intent(in) ii )
```

this subroutine reads the HRU weather generator parameters from the .wgn file

Parameters

| | | |
|----|----|------------------------|
| in | ii | subbasin number (none) |
|----|----|------------------------|

7.103 readwus.f90 File Reference

Functions/Subroutines

- subroutine [readwus](#) (i)

This subroutine reads data from the HRU/subbasin water use input file (.wus). The water use file extracts water from the subbasin and it is considered to be lost from the watershed. These variables should be used to remove water transported outside the watershed.

7.103.1 Detailed Description

file containing the subroutine readwus

Author

modified by Javier Burguete

7.103.2 Function/Subroutine Documentation

7.103.2.1 readwus()

```
subroutine readwus (
    integer, intent(in) i )
```

This subroutine reads data from the HRU/subbasin water use input file (.wus). The water use file extracts water from the subbasin and it is considered to be lost from the watershed. These variables should be used to remove water transported outside the watershed.

Parameters

| | | |
|----|---|-----------------|
| in | i | subbasin number |
|----|---|-----------------|

7.104 readwwq.f90 File Reference**Functions/Subroutines**

- subroutine [readwwq](#)
this subroutine reads the watershed stream water quality input data (.wwq file) and initializes the QUAL2E variables which apply to the entire watershed

7.104.1 Detailed Description

file containing the subroutine readwwq

Author

modified by Javier Burguete

7.105 readyr.f90 File Reference**Functions/Subroutines**

- subroutine [readyr](#) (i)
reads in the input data for the recyear command

7.105.1 Detailed Description

file containing the subroutine readyr

Author

modified by Javier Burguete

7.105.2 Function/Subroutine Documentation**7.105.2.1 readyr()**

```
subroutine readyr (
    integer, intent(in) i )
```

reads in the input data for the recyear command

Parameters

| | | |
|----|---|-------------------------|
| in | i | reservoir number (none) |
|----|---|-------------------------|

7.106 resetlu.f90 File Reference

Functions/Subroutines

- subroutine [resetlu](#)

this subroutine reads data from the HRU/subbasin management input file (.mgt). This file contains data related to management practices used in the HRU/subbasin.

7.106.1 Detailed Description

file containing the subroutine resetlu

Author

modified by Javier Burguete

7.107 rhgen.f90 File Reference

Functions/Subroutines

- subroutine [rhgen](#) (j)

this subroutine generates weather relative humidity, solar radiation, and wind speed.

7.107.1 Detailed Description

file containing the subroutine rhgen

Author

modified by Javier Burguete

7.108 rootfr.f90 File Reference

Functions/Subroutines

- subroutine [rootfr](#) (j)

this subroutine distributes dead root mass through the soil profile

7.108.1 Detailed Description

file containing the subroutine rootfr

Author

Armen R. Kemanian,
modified by Javier Burguete

7.108.2 Function/Subroutine Documentation

7.108.2.1 rootfr()

```
subroutine rootfr (
    integer, intent(in) j )
```

this subroutine distributes dead root mass through the soil profile

Parameters

| | | |
|----|---|------------|
| in | j | HRU number |
|----|---|------------|

7.109 rsedaa.f90 File Reference

Functions/Subroutines

- subroutine [rsedaa](#) (years)
this subroutine writes the annual reach output to the .sed file

7.109.1 Detailed Description

file containing the subroutine rsedaa

Author

modified by Javier Burguete

7.109.2 Function/Subroutine Documentation

7.109.2.1 rsedaa()

```
subroutine rsedaa (
    real*8, intent(in) years )
```

this subroutine writes the annual reach output to the .sed file

Parameters

| | |
|--------------|------------------------------|
| <i>years</i> | length of simulation (years) |
|--------------|------------------------------|

7.110 **rseday.f90** File Reference

Functions/Subroutines

- subroutine **rseday**

7.110.1 Detailed Description

file containing the subroutine `rseday`

Author

modified by Javier Burguete

7.111 **rsedmon.f90** File Reference

Functions/Subroutines

- subroutine `rsedmon` (*m*days)
this subroutine writes the monthly reach output to the .sed file

7.111.1 Detailed Description

file containing the subroutine `rsedmon`

Author

modified by Javier Burguete

7.111.2 Function/Subroutine Documentation

7.111.2.1 `rsedmon()`

```
subroutine rsedmon (  
    integer, intent(in) mdays )
```

this subroutine writes the monthly reach output to the .sed file

Parameters

| | | |
|----|-------|-----------------------------------|
| in | mdays | number of days simulated in month |
|----|-------|-----------------------------------|

7.112 rsedyr.f90 File Reference

Functions/Subroutines

- subroutine [rsedyr](#)
this subroutine writes the yearly reach output to the .sed file

7.112.1 Detailed Description

file containing the subroutine rsedyr

Author

modified by Javier Burguete

7.113 rteinit.f90 File Reference

Functions/Subroutines

- subroutine [rteinit](#)
This subroutine reads in the areas associated with files processed with the recday, recepic, recmon and recyear commands, calculates subbasin areas, calculates reach and hydrograph node drainage areas.

7.113.1 Detailed Description

file containing the subroutine rteinit

Author

modified by Javier Burguete

7.114 sched_mgt.f90 File Reference

Functions/Subroutines

- subroutine [sched_mgt](#) (j)
this subroutine performs all management operations

7.114.1 Detailed Description

file containing the subroutine sched_mgt

Author

modified by Javier Burguete

7.114.2 Function/Subroutine Documentation

7.114.2.1 sched_mgt()

```
subroutine sched_mgt (
    integer, intent(in) j )
```

this subroutine performs all management operations

Parameters

| | | |
|----|----------|------------|
| in | <i>j</i> | HRU number |
|----|----------|------------|

7.115 schedule_ops.f90 File Reference

Functions/Subroutines

- subroutine [schedule_ops](#) (*j*)
this subroutine controls the simulation of the land phase of the hydrologic cycle

7.115.1 Detailed Description

file containing the subroutine schedule_ops

Author

modified by Javier Burguete

7.115.2 Function/Subroutine Documentation

7.115.2.1 schedule_ops()

```
subroutine schedule_ops (
    integer, intent(in) j )
```

this subroutine controls the simulation of the land phase of the hydrologic cycle

Parameters

| | | |
|----|---|------------|
| in | j | HRU number |
|----|---|------------|

7.116 sim_inityr.f90 File Reference

Functions/Subroutines

- subroutine [sim_inityr](#)
this subroutine initializes variables at the beginning of the year

7.116.1 Detailed Description

file containing the subroutine sim_inityr

Author

modified by Javier Burguete

7.117 simulate.f90 File Reference

Functions/Subroutines

- subroutine [simulate](#)
this subroutine contains the loops governing the modeling of processes in the watershed

7.117.1 Detailed Description

file containing the subroutine simulate

Author

modified by Javier Burguete

7.118 slrgen.f90 File Reference

Functions/Subroutines

- subroutine [slrgen](#) (j)
this subroutine generates solar radiation

7.118.1 Detailed Description

file containing the subroutine slrgen

Author

modified by Javier Burguete

7.118.2 Function/Subroutine Documentation

7.118.2.1 slrgen()

```
subroutine slrgen (
    integer, intent(in) j )
```

this subroutine generates solar radiation

Parameters

| | | |
|----|---|------------|
| in | j | HRU number |
|----|---|------------|

7.119 smeas.f90 File Reference

Functions/Subroutines

- subroutine [smeas](#)
this subroutine reads in daily solar radiation data and assigns the values to the proper HRUs

7.119.1 Detailed Description

file containing the subroutine smeas

Author

modified by Javier Burguete

7.120 snom.f90 File Reference

Functions/Subroutines

- subroutine [snom](#) (j)
this subroutine predicts daily snom melt when the average air temperature exceeds 0 degrees Celsius

7.120.1 Detailed Description

file containing the subroutine snom

Author

modified by Javier Burguete

7.120.2 Function/Subroutine Documentation

7.120.2.1 snom()

```
subroutine snom (
    integer, intent(in) j )
```

this subroutine predicts daily snom melt when the average air temperature exceeds 0 degrees Celsius

Parameters

| | | |
|----|---|------------|
| in | j | HRU number |
|----|---|------------|

7.121 soil_chem.f90 File Reference

Functions/Subroutines

- subroutine [soil_chem](#) (ii)
this subroutine initializes soil chemical properties

7.121.1 Detailed Description

file containing the subroutine soil_chem

Author

modified by Javier Burguete

7.121.2 Function/Subroutine Documentation

7.121.2.1 soil_chem()

```
subroutine soil_chem (
    integer, intent(in) ii )
```

this subroutine initializes soil chemical properties

Parameters

| | | |
|----|----|------------|
| in | ii | HRU number |
|----|----|------------|

7.122 soil_phys.f90 File Reference

Functions/Subroutines

- subroutine [soil_phys](#) (ii)
this subroutine initializes soil physical properties

7.122.1 Detailed Description

file containing the subroutine soil_phys

Author

modified by Javier Burguete

7.122.2 Function/Subroutine Documentation

7.122.2.1 soil_phys()

```
subroutine soil_phys (  
    integer, intent(in) ii )
```

this subroutine initializes soil physical properties

Parameters

| | | |
|----|----|------------|
| in | ii | HRU number |
|----|----|------------|

7.123 soil_write.f90 File Reference

Functions/Subroutines

- subroutine [soil_write](#) (i)
this subroutine writes output to the output.sol file

7.123.1 Detailed Description

file containing the subroutine soil_write

Author

modified by Javier Burguete

7.123.2 Function/Subroutine Documentation

7.123.2.1 soil_write()

```
subroutine soil_write (
    integer, intent(in) i )
```

this subroutine writes output to the output.sol file

Parameters

| | | |
|----|---|--|
| in | i | current day in simulation - loop counter (julian date) |
|----|---|--|

7.124 solt.f90 File Reference

Functions/Subroutines

- subroutine [solt](#) (j)
this subroutine estimates daily average temperature at the bottom of each soil layer @parameter[in] j HRU number

7.124.1 Detailed Description

file containing the subroutine solt

Author

modified by Javier Burguete

7.125 std1.f90 File Reference

Functions/Subroutines

- subroutine [std1](#)
this subroutine writes general information to the standard output file and header lines to miscellaneous output files

7.125.1 Detailed Description

file containing the subroutine std1

Author

modified by Javier Burguete

7.126 std2.f90 File Reference

Functions/Subroutines

- subroutine [std2](#)
this subroutine writes general information to the standard output file and to miscellaneous output files

7.126.1 Detailed Description

file containing the subroutine std2

Author

modified by Javier Burguete

7.127 std3.f90 File Reference

Functions/Subroutines

- subroutine [std3](#)
this subroutine writes the annual table header to the standard output file

7.127.1 Detailed Description

file containing the subroutine std3

Author

modified by Javier Burguete

7.128 storeinitial.f90 File Reference

Functions/Subroutines

- subroutine [storeinitial](#)
this subroutine saves initial values for variables that must be reset to rerun the simulation for different real time weather scenarios

7.128.1 Detailed Description

file containing the subroutine storeinitial

Author

modified by Javier Burguete

7.129 subbasin.f90 File Reference

Functions/Subroutines

- subroutine [subbasin](#) (i)
this subroutine controls the simulation of the land phase of the hydrologic cycle

7.129.1 Detailed Description

file containing the subroutine subbasin

Author

modified by Javier Burguete

7.129.2 Function/Subroutine Documentation

7.129.2.1 subbasin()

```
subroutine subbasin (
    integer, intent(in) i )
```

this subroutine controls the simulation of the land phase of the hydrologic cycle

Parameters

| | | |
|----|---|--|
| in | i | current day in simulation—loop counter (julian date) |
|----|---|--|

7.130 submon.f90 File Reference

Functions/Subroutines

- subroutine [submon](#)
this subroutine writes monthly subbasin output to the output.sub file

7.130.1 Detailed Description

file containing the subroutine submon

Author

modified by Javier Burguete

7.131 subyr.f90 File Reference

Functions/Subroutines

- subroutine [subyr](#)
this subroutine writes annual subbasin output to the output.sub file

7.131.1 Detailed Description

file containing the subroutine subyr

Author

modified by Javier Burguete

7.132 surface.f90 File Reference

Functions/Subroutines

- subroutine [surface](#) (i, j)
this subroutine models surface hydrology at any desired time step

7.132.1 Detailed Description

file containing the subroutine surface

Author

modified by Javier Burguete

7.132.2 Function/Subroutine Documentation

7.132.2.1 surface()

```
subroutine surface (  
    integer, intent(in) i,  
    integer, intent(in) j )
```

this subroutine models surface hydrology at any desired time step

Parameters

| | | |
|----|----------|--|
| in | <i>i</i> | current day in simulation-loop counter (julian date) |
| in | <i>j</i> | HRU number (none) |

7.133 surfst_h2o.f90 File Reference

Functions/Subroutines

- subroutine [surfst_h2o](#) (*j*)

this subroutine determines the net surface runoff reaching the main channel on a given day. The net amount of water reaching the main channel can include water in surface runoff from the previous day and will exclude surface runoff generated on the current day which takes longer than one day to reach the main channel

7.133.1 Detailed Description

file containing the subroutine `surfst_h2o`

Author

modified by Javier Burguete

7.133.2 Function/Subroutine Documentation

7.133.2.1 `surfst_h2o()`

```
subroutine surfst_h2o (
    integer, intent(in) j )
```

this subroutine determines the net surface runoff reaching the main channel on a given day. The net amount of water reaching the main channel can include water in surface runoff from the previous day and will exclude surface runoff generated on the current day which takes longer than one day to reach the main channel

Parameters

| | | |
|----|----------|------------|
| in | <i>j</i> | HRU number |
|----|----------|------------|

7.134 surq_daycn.f90 File Reference

Functions/Subroutines

- subroutine [surq_daycn](#) (*j*)

predicts daily runoff given daily precipitation and snow melt using a modified SCS curve number approach

7.134.1 Detailed Description

file containing the subroutine `surq_daycn`

Author

modified by Javier Burguete

7.134.2 Function/Subroutine Documentation

7.134.2.1 `surq_daycn()`

```
subroutine surq_daycn (
    integer, intent(in) j )
```

predicts daily runoff given daily precipitation and snow melt using a modified SCS curve number approach

Parameters

| | | |
|-----------------|----------------|-------------------|
| <code>in</code> | <code>j</code> | HRU number (none) |
|-----------------|----------------|-------------------|

7.135 `surq_greenampt.f90` File Reference

Functions/Subroutines

- subroutine [surq_greenampt](#) (j)

predicts daily runoff given breakpoint precipitation and snow melt using the Green & Ampt technique

7.135.1 Detailed Description

file containing the subroutine `surq_greenampt`

Author

modified by Javier Burguete

7.135.2 Function/Subroutine Documentation

7.135.2.1 surq_greenampt()

```
subroutine surq_greenampt (
    integer, intent(in) j )
```

predicts daily runoff given breakpoint precipitation and snow melt using the Green & Ampt technique

Parameters

| | | |
|----|---|-------------------|
| in | j | HRU number (none) |
|----|---|-------------------|

7.136 swbl.f90 File Reference**Functions/Subroutines**

- subroutine [swbl](#) (snow, irrg)

this subroutine checks the soil water balance at the end of the simulation

7.136.1 Detailed Description

file containing the subroutine swbl

Author

modified by Javier Burguete

7.136.2 Function/Subroutine Documentation**7.136.2.1 swbl()**

```
subroutine swbl (
    real*8, intent(in) snow,
    real*8, intent(in) irrg )
```

this subroutine checks the soil water balance at the end of the simulation

Parameters

| | | |
|----|------|--|
| in | snow | snow in watershed at end of simulation |
| in | irrg | irrigation water applied to watershed |

7.137 tgen.f90 File Reference

Functions/Subroutines

- subroutine [tgen](#) (j)

this subroutine generates temperature data when the user chooses to simulate or when data is missing for particular days in the weather file

7.137.1 Detailed Description

file containing the subroutine tgen

Author

modified by Javier Burguete

7.137.2 Function/Subroutine Documentation

7.137.2.1 tgen()

```
subroutine tgen (
    integer, intent(in) j )
```

this subroutine generates temperature data when the user chooses to simulate or when data is missing for particular days in the weather file

Parameters

| | | |
|----|---|------------|
| in | j | HRU number |
|----|---|------------|

7.138 tillfactor.f90 File Reference

Functions/Subroutines

- subroutine [tillfactor](#) (j, bmix, emix, dtil, sol_thick)

this procedure increases tillage factor (tillagef(l,j)) per layer for each operation. The tillage factor settling will depend of soil moisture (tentatively) and must be called every day. For simplicity the settling is calculated now at the soil carbon sub because soil water content is available.

7.138.1 Detailed Description

file containing the subroutine tillfactor

Author

modified by Javier Burguete

7.138.2 Function/Subroutine Documentation

7.138.2.1 tillfactor()

```
subroutine tillfactor (
    integer, intent(in) j,
    real*8, intent(in) bmix,
    real*8, intent(inout) emix,
    real*8, intent(in) dtil,
    real*8, dimension(sol_nly(j)), intent(in) sol_thick )
```

this procedure increases tillage factor (tillagef(l,j) per layer for each operation. The tillage factor settling will depend of soil moisture (tentatively) and must be called every day. For simplicity the settling is calculated now at the soil carbon sub because soil water content is available.

Parameters

| | | |
|---------|------------------|--|
| in | <i>j</i> | HRU number (none) |
| in | <i>bmix</i> | biological mixing efficiency: this number is zero for tillage operations (none) |
| in, out | <i>emix</i> | mixing efficiency (none) |
| in | <i>dtil</i> | depth of mixing (mm) |
| in | <i>sol_thick</i> | <p>The tillage factor depends on the cumulative soil disturbance rating = csdr For simplicity, csdr is a function of emix. First step is to calculate "current" csdr by inverting tillage factor function. The effect of texture on tillage factor (ZZ) is removed first (and recovered at the end of the procedure).</p> $YY = \text{tillagef}(l, j) / ZZ$ <p>Since the tillage factor function is non linear, iterations are needed. $XX = 0.5$ is the initial value that works OK for the range of values observed. If a layer is only partially tilled then emix is corrected accordingly</p> |

7.139 tmeas.f90 File Reference

Functions/Subroutines

- subroutine [tmeas](#)

this subroutine reads in temperature data and assigns it to the HRUs

7.139.1 Detailed Description

file containing the subroutine tmeas

Author

modified by Javier Burguete

7.140 tran.f90 File Reference

Functions/Subroutines

- subroutine `tran` (j)
this subroutine computes tributary channel transmission losses

7.140.1 Detailed Description

file containing the subroutine tran

Author

modified by Javier Burguete

7.140.2 Function/Subroutine Documentation

7.140.2.1 tran()

```
subroutine tran (
    integer, intent(in) j )
```

this subroutine computes tributary channel transmission losses

Parameters

| | | |
|----|---|-------------------|
| in | j | HRU number (none) |
|----|---|-------------------|

7.141 ttcoef.f90 File Reference

Functions/Subroutines

- subroutine `ttcoef` (k)
this subroutine computes travel time coefficients for routing along the main channel

7.141.1 Detailed Description

file containing the subroutine ttcoef

Author

modified by Javier Burguete

7.141.2 Function/Subroutine Documentation**7.141.2.1 ttcoef()**

```
subroutine ttcoef (
    integer, intent(in) k )
```

this subroutine computes travel time coefficients for routing along the main channel

Parameters

| | | |
|----|---|------------|
| in | k | HRU number |
|----|---|------------|

7.142 ttcoef_wway.f90 File Reference**Functions/Subroutines**

- subroutine [ttcoef_wway](#) (j)
this subroutine computes travel time coefficients for routing along the main channel - grassed waterways

7.142.1 Detailed Description

file containing the subroutine ttcoef_wway

Author

modified by Javier Burguete

7.143 varinit.f90 File Reference**Functions/Subroutines**

- subroutine [varinit](#) (j)
this subroutine initializes variables for the daily simulation of the land phase of the hydrologic cycle (the subbasin command loop)

7.143.1 Detailed Description

file containing the subroutine varinit

Author

modified by Javier Burguete

7.143.2 Function/Subroutine Documentation

7.143.2.1 varinit()

```
subroutine varinit (
    integer, intent(in) j )
```

this subroutine initializes variables for the daily simulation of the land phase of the hydrologic cycle (the subbasin command loop)

Parameters

| | | |
|----|---|------------|
| in | j | HRU number |
|----|---|------------|

7.144 vbl.f90 File Reference

Functions/Subroutines

- subroutine [vbl](#) (evx, spx, pp, qin, ox, vx1, vy, yi, yo, ysx, vf, vyf, aha)
this subroutine checks the water and sediment balance for ponds and reservoirs at the end of a simulation

7.144.1 Detailed Description

file containing the subroutine vbl

Author

modified by Javier Burguete

7.144.2 Function/Subroutine Documentation

7.144.2.1 vbl()

```

subroutine vbl (
    real*8, intent(in) evx,
    real*8, intent(in) spx,
    real*8, intent(in) pp,
    real*8, intent(in) qin,
    real*8, intent(in) ox,
    real*8, intent(inout) vx1,
    real*8, intent(inout) vy,
    real*8, intent(in) yi,
    real*8, intent(in) yo,
    real*8, intent(in) ysx,
    real*8, intent(in) vf,
    real*8, intent(in) vyf,
    real*8, intent(in) aha )

```

this subroutine checks the water and sediment balance for ponds and reservoirs at the end of a simulation

Parameters

| | | |
|---------|-----|--|
| in | evx | evaporation from water body |
| in | spx | seepage from water body |
| in | pp | precipitation on water body |
| in | qin | water entering water body |
| in | ox | water leaving water body |
| in, out | vx1 | (in) volume of water in water body at beginning of simulation
(out) dfw expressed as depth over drainage area |
| in, out | vy | (in) sediment in water body at beginning of simulation
(out) dfy expressed as loading per unit area for drainage area |
| in | yi | sediment entering water body |
| in | yo | sediment leaving water body |
| in | ysx | change in sediment level in water body |
| in | vf | volume of water in water body at end of simulation |
| in | vyf | sediment in water body at end of simulation |
| in | aha | area draining into water body |

7.145 volq.f90 File Reference

Functions/Subroutines

- subroutine [volq](#) (j)

call subroutines to calculate the current day's CN for the HRU and to calculate surface runoff

7.145.1 Detailed Description

file containing the subroutine volq

Author

modified by Javier Burguete

7.145.2 Function/Subroutine Documentation

7.145.2.1 volq()

```
subroutine volq (
    integer, intent(in) j )
```

call subroutines to calculate the current day's CN for the HRU and to calculate surface runoff

Parameters

| | | |
|----|---|-------------------|
| in | j | HRU number (none) |
|----|---|-------------------|

7.146 water_hru.f90 File Reference

Functions/Subroutines

- subroutine [water_hru](#) (j)
this subroutine compute pet and et using Priestly-Taylor and a coefficient

7.146.1 Detailed Description

file containing the subroutine water_hru

Author

modified by Javier Burguete

7.147 watable.f90 File Reference

Functions/Subroutines

- subroutine [watable](#) (j)
this subroutine is the master soil percolation component. param[in] j HRU number

7.147.1 Detailed Description

file containing the subroutine watable

Author

modified by Javier Burguete

7.148 weatgn.f90 File Reference

Functions/Subroutines

- subroutine [weatgn](#) (*j*)

this subroutine generates weather parameters used to simulate the impact of precipitation on the other climatic processes

7.148.1 Detailed Description

file containing the subroutine weatgn

Author

modified by Javier Burguete

7.148.2 Function/Subroutine Documentation

7.148.2.1 weatgn()

```
subroutine weatgn (
    integer, intent(in) j )
```

this subroutine generates weather parameters used to simulate the impact of precipitation on the other climatic processes

Parameters

| | | |
|----|----------|------------|
| in | <i>j</i> | HRU number |
|----|----------|------------|

7.149 wmeas.f90 File Reference

Functions/Subroutines

- subroutine [wmeas](#)

this subroutine reads in wind speed data from file and assigns the data to HRUs

7.149.1 Detailed Description

file containing the subroutine wmeas

Author

modified by Javier Burguete

7.150 wndgen.f90 File Reference

Functions/Subroutines

- subroutine [wndgen](#) (j)
this subroutine generates wind speed

7.150.1 Detailed Description

file containing the subroutine wndgen

Author

modified by Javier Burguete

7.150.2 Function/Subroutine Documentation

7.150.2.1 wndgen()

```
subroutine wndgen (  
    integer, intent(in) j )
```

this subroutine generates wind speed

Parameters

| | | |
|----|----------|------------|
| in | <i>j</i> | HRU number |
|----|----------|------------|

7.151 writea.f90 File Reference

Functions/Subroutines

- subroutine [writea](#) (i)
this subroutine writes annual output

7.151.1 Detailed Description

file containing the subroutine writea

Author

modified by Javier Burguete

7.151.2 Function/Subroutine Documentation

7.151.2.1 writea()

```
subroutine writea (
    integer, intent(in) i )
```

this subroutine writes annual output

Parameters

| | | |
|-----------------|----------------|---|
| <code>in</code> | <code>i</code> | current day of simulation (julian date) |
|-----------------|----------------|---|

7.152 writed.f90 File Reference

Functions/Subroutines

- subroutine [writed](#)
this subroutine contains the daily output writes

7.152.1 Detailed Description

file containing the subroutine writed

Author

modified by Javier Burguete

7.153 writem.f90 File Reference

Functions/Subroutines

- subroutine [writem](#) (i)
this subroutine writes monthly output

7.153.1 Detailed Description

file containing the subroutine writem

Author

modified by Javier Burguete

7.153.2 Function/Subroutine Documentation

7.153.2.1 writem()

```
subroutine writem (
    integer, intent(in) i )
```

this subroutine writes monthly output

Parameters

| | | |
|----|---|---|
| in | i | current day of simulation (julian date) |
|----|---|---|

7.154 xmon.f90 File Reference

Functions/Subroutines

- subroutine [xmon](#)
this subroutine determines the month, given the julian date and leap year flag

7.154.1 Detailed Description

file containing the subroutine xmon

Author

modified by Javier Burguete

7.155 ysed.f90 File Reference

Functions/Subroutines

- subroutine [ysed](#) (iwave, j)
this subroutine predicts daily soil loss caused by water erosion using the modified universal soil loss equation

7.155.1 Detailed Description

file containing the subroutine ysed

Author

modified by Javier Burguete

7.155.2 Function/Subroutine Documentation

7.155.2.1 ysed()

```
subroutine ysed (
    integer, intent(in) iwave,
    integer, intent(in) j )
```

this subroutine predicts daily soil loss caused by water erosion using the modified universal soil loss equation

Parameters

| | | |
|----|--------------|---|
| in | <i>iwave</i> | flag to differentiate calculation of HRU and subbasin sediment calculation (none)
iwave = 0 for HRU
iwave = subbasin # for subbasin |
| in | <i>j</i> | HRU number |

7.156 zero0.f90 File Reference

Functions/Subroutines

- subroutine [zero0](#)
this subroutine initializes the values for some of the arrays

7.156.1 Detailed Description

file containing the subroutine zero0

Author

modified by Javier Burguete

7.157 zero1.f90 File Reference

Functions/Subroutines

- subroutine [zero1](#)
this subroutine initializes the values for some of the arrays

7.157.1 Detailed Description

file containing the subroutine zero1

Author

modified by Javier Burguete

7.158 zero2.f90 File Reference

Functions/Subroutines

- subroutine [zero2](#)
this subroutine zeros all array values

7.158.1 Detailed Description

file containing the subroutine zero2

Author

modified by Javier Burguete

7.159 zero_urbn.f90 File Reference

Functions/Subroutines

- subroutine [zero_urbn](#)
this subroutine zeros all array values used in urban modeling

7.159.1 Detailed Description

file containing the subroutine zero_urbn

Author

modified by Javier Burguete

7.160 zeroini.f90 File Reference

Functions/Subroutines

- subroutine [zeroini](#)
this subroutine zeros values for single array variables

7.160.1 Detailed Description

file containing the subroutine zeroini

Author

modified by Javier Burguete

Bibliography

- [1] P Bratley, B L Fox, and L E Schrage. *A Guide to Simulation*. Springer-Verlag, New York, USA, 1983. [107](#)
- [2] J. E. McCray, S. L. Kirkland, R. L. Siegrist, and G. D. Thyne. Model parameters for simulating fate and transport of on-site wastewater nutrients. *Ground Water*, 43(4):628–639, 2005. [240](#)
- [3] R. L. Siegrist, J. McCray, L. Weintraub, C. Chen, J. Bagdol, P. Lemonds, S. Van Cuyk, K. Lowe, R. Goldstein, and J. Rada. Quantifying site-scale processes and watershed-scale cumulative effects of decentralized wastewater systems, project no. wu-ht-00-27. Prepared for the National Decentralized Water Resources Capacity Development Project, Washington University, St. Louis, MO, by the Colorado School of Mines, 2005. [240](#)

Index

albedo.f90, [103](#)
allocate_parms.f90, [103](#)
alph
 alph.f90, [104](#)
alph.f90, [104](#)
 alph, [104](#)
apply
 apply.f90, [105](#)
apply.f90, [104](#)
 apply, [105](#)
ascrv
 ascrv.f90, [105](#)
ascrv.f90, [105](#)
 ascrv, [105](#)
atri
 atri.f90, [106](#)
atri.f90, [106](#)
 atri, [106](#)
aunif
 aunif.f90, [107](#)
aunif.f90, [107](#)
 aunif, [107](#)

bmpinit
 bmpinit.f90, [108](#)
bmpinit.f90, [108](#)
 bmpinit, [108](#)
burnop
 burnop.f90, [109](#)
burnop.f90, [108](#)
 burnop, [109](#)

canopyint.f90, [109](#)
caps
 caps.f90, [110](#)
caps.f90, [109](#)
 caps, [110](#)
cfactor.f90, [110](#)
clgen
 clgen.f90, [111](#)
clgen.f90, [110](#)
 clgen, [111](#)
clicon
 clicon.f90, [111](#)
clicon.f90, [111](#)
 clicon, [111](#)
command
 command.f90, [112](#)
command.f90, [112](#)
 command, [112](#)

crackflow.f90, [113](#)
crackvol.f90, [113](#)
curno
 curno.f90, [114](#)
curno.f90, [113](#)
 curno, [114](#)

dailycn.f90, [114](#)
dstn1
 dstn1.f90, [115](#)
dstn1.f90, [114](#)
 dstn1, [115](#)

ee
 ee.f90, [116](#)
ee.f90, [115](#)
 ee, [116](#)
eiusle.f90, [116](#)
estimate_ksat
 estimate_ksat.f90, [117](#)
estimate_ksat.f90, [116](#)
 estimate_ksat, [117](#)
etpot
 etpot.f90, [117](#)
etpot.f90, [117](#)
 etpot, [117](#)
expo
 expo.f90, [118](#)
expo.f90, [118](#)
 expo, [118](#)

fert
 fert.f90, [119](#)
fert.f90, [118](#)
 fert, [119](#)
finalbal.f90, [119](#)

gcycl.f90, [119](#)
getallo.f90, [120](#)
graze
 graze.f90, [120](#)
graze.f90, [120](#)
 graze, [120](#)

h2omgt_init.f90, [121](#)
harvestop
 harvestop.f90, [121](#)
harvestop.f90, [121](#)
 harvestop, [121](#)
harvkillop
 harvkillop.f90, [122](#)

- harvkillop.f90, 122
 - harvkillop, 122
- headout.f90, 122
- hmeas.f90, 123
- hruaa
 - hruaa.f90, 123
- hruaa.f90, 123
 - hruaa, 123
- hruallo.f90, 124
- hrumon.f90, 124
- hruyr.f90, 124
- hydroinit.f90, 125
- icl
 - icl.f90, 125
- icl.f90, 125
 - icl, 125
- igropt
 - parm, 99
- impnd_init.f90, 126
- impndmon.f90, 126
- impndyr.f90, 126
- irrigate
 - irrigate.f90, 127
- irrigate.f90, 127
 - irrigate, 127
- irrsb
 - irrsb.f90, 128
- irrsb.f90, 127
 - irrsb, 128
- jdt
 - jdt.f90, 128
- jdt.f90, 128
 - jdt, 128
- killop
 - killop.f90, 129
- killop.f90, 129
 - killop, 129
- lid_cistern
 - lid_cistern.f90, 130
- lid_cistern.f90, 130
 - lid_cistern, 130
- lid_greenroof
 - lid_greenroof.f90, 131
- lid_greenroof.f90, 130
 - lid_greenroof, 131
- lid_porpavement
 - lid_porpavement.f90, 132
- lid_porpavement.f90, 131
 - lid_porpavement, 132
- lid_raingarden
 - lid_raingarden.f90, 132
- lid_raingarden.f90, 132
 - lid_raingarden, 132
- lidinit
 - lidinit.f90, 133
- lidinit.f90, 133
 - lidinit, 133
- lids
 - lids.f90, 134
- lids.f90, 133
 - lids, 134
- lwqdef
 - lwqdef.f90, 135
- lwqdef.f90, 134
 - lwqdef, 135
- main.f90, 135
- modparm.f90, 135
- newtillmix
 - newtillmix.f90, 219
- newtillmix.f90, 219
 - newtillmix, 219
- openwth.f90, 220
- ovr_sed.f90, 220
- parm, 15
 - igropt, 99
- parm::fcgd, 101
- parm::ndenit, 101
- parm::regres, 101
- parm::tair, 102
- parm::theta, 102
- pgen
 - pgen.f90, 221
- pgen.f90, 220
 - pgen, 221
- pgenhr.f90, 221
- pkq
 - pkq.f90, 222
- pkq.f90, 221
 - pkq, 222
- plantop
 - plantop.f90, 222
- plantop.f90, 222
 - plantop, 222
- pmeas
 - pmeas.f90, 223
- pmeas.f90, 223
 - pmeas, 223
- qman
 - qman.f90, 224
- qman.f90, 224
 - qman, 224
- rchaa
 - rchaa.f90, 225
- rchaa.f90, 224
 - rchaa, 225
- rchday.f90, 225
- rchmon
 - rchmon.f90, 226
- rchmon.f90, 225

- rchmon, 226
- rchyr
 - rchyr.f90, 226
- rchyr.f90, 226
- rchyr, 226
- readatmodep.f90, 227
- readbsn.f90, 227
- readchm
 - readchm.f90, 228
- readchm.f90, 227
- readchm, 228
- readcnst
 - readcnst.f90, 228
- readcnst.f90, 228
- readcnst, 228
- readfcst.f90, 229
- readfert.f90, 229
- readfig.f90, 229
- readfile.f90, 230
- readgw
 - readgw.f90, 230
- readgw.f90, 230
- readgw, 230
- readhru
 - readhru.f90, 231
- readhru.f90, 231
- readhru, 231
- readinpt.f90, 231
- readlup.f90, 232
- readlwq
 - readlwq.f90, 232
- readlwq.f90, 232
- readlwq, 232
- readmgt
 - readmgt.f90, 233
- readmgt.f90, 233
- readmgt, 233
- readmon.f90, 233
- readops
 - readops.f90, 234
- readops.f90, 234
- readops, 234
- readpest.f90, 234
- readplant.f90, 235
- readpnd
 - readpnd.f90, 235
- readpnd.f90, 235
- readpnd, 235
- readres
 - readres.f90, 236
- readres.f90, 236
- readres, 236
- readrte.f90, 236
- readru
 - readru.f90, 237
- readru.f90, 237
- readru, 237
- readsdr
 - readsdr.f90, 238
- readsdr.f90, 237
- readsdr, 238
- readsepticbz
 - readsepticbz.f90, 238
- readsepticbz.f90, 238
- readsepticbz, 238
- readseptwq
 - readseptwq.f90, 240
- readseptwq.f90, 240
- readseptwq, 240
- readsno
 - readsno.f90, 241
- readsno.f90, 240
- readsno, 241
- readsol
 - readsol.f90, 241
- readsol.f90, 241
- readsol, 241
- readsub
 - readsub.f90, 242
- readsub.f90, 242
- readsub, 242
- readswq.f90, 242
- readtill.f90, 243
- readurban.f90, 243
- readwgn
 - readwgn.f90, 244
- readwgn.f90, 243
- readwgn, 244
- readwus
 - readwus.f90, 244
- readwus.f90, 244
- readwus, 244
- readwwq.f90, 245
- readyr
 - readyr.f90, 245
- readyr.f90, 245
- readyr, 245
- resetlu.f90, 246
- rhgen.f90, 246
- rootfr
 - rootfr.f90, 247
- rootfr.f90, 246
- rootfr, 247
- rsedaa
 - rsedaa.f90, 247
- rsedaa.f90, 247
- rsedaa, 247
- rseday.f90, 248
- rsedmon
 - rsedmon.f90, 248
- rsedmon.f90, 248
- rsedmon, 248
- rsedyr.f90, 249
- rteinit.f90, 249
- sched_mgt
 - sched_mgt.f90, 250

sched_mgt.f90, 249
 sched_mgt, 250
 schedule_ops
 schedule_ops.f90, 250
 schedule_ops.f90, 250
 schedule_ops, 250
 sim_inityr.f90, 251
 simulate.f90, 251
 slrgen
 slrgen.f90, 252
 slrgen.f90, 251
 slrgen, 252
 smeas.f90, 252
 snom
 snom.f90, 253
 snom.f90, 252
 snom, 253
 soil_chem
 soil_chem.f90, 253
 soil_chem.f90, 253
 soil_chem, 253
 soil_phys
 soil_phys.f90, 254
 soil_phys.f90, 254
 soil_phys, 254
 soil_write
 soil_write.f90, 255
 soil_write.f90, 254
 soil_write, 255
 solt.f90, 255
 std1.f90, 255
 std2.f90, 256
 std3.f90, 256
 storeinitial.f90, 256
 subbasin
 subbasin.f90, 257
 subbasin.f90, 257
 subbasin, 257
 submon.f90, 257
 subyr.f90, 258
 surface
 surface.f90, 258
 surface.f90, 258
 surface, 258
 surfst_h2o
 surfst_h2o.f90, 259
 surfst_h2o.f90, 259
 surfst_h2o, 259
 surq_daycn
 surq_daycn.f90, 260
 surq_daycn.f90, 259
 surq_daycn, 260
 surq_greenampt
 surq_greenampt.f90, 260
 surq_greenampt.f90, 260
 surq_greenampt, 260
 swbl
 swbl.f90, 261
 swbl.f90, 261
 swbl, 261
 tgen
 tgen.f90, 262
 tgen.f90, 262
 tgen, 262
 tillfactor
 tillfactor.f90, 263
 tillfactor.f90, 262
 tillfactor, 263
 tmeas.f90, 263
 tran
 tran.f90, 264
 tran.f90, 264
 tran, 264
 ttcoef
 ttcoef.f90, 265
 ttcoef.f90, 264
 ttcoef, 265
 ttcoef_wway.f90, 265
 varinit
 varinit.f90, 266
 varinit.f90, 265
 varinit, 266
 vbl
 vbl.f90, 266
 vbl.f90, 266
 vbl, 266
 volq
 volq.f90, 268
 volq.f90, 267
 volq, 268
 water_hru.f90, 268
 wattable.f90, 268
 weatgn
 weatgn.f90, 269
 weatgn.f90, 269
 weatgn, 269
 wmeas.f90, 269
 wndgen
 wndgen.f90, 270
 wndgen.f90, 270
 wndgen, 270
 writea
 writea.f90, 271
 writea.f90, 270
 writea, 271
 writed.f90, 271
 writem
 writem.f90, 272
 writem.f90, 271
 writem, 272
 xmon.f90, 272
 ysed

ysed.f90, [273](#)
ysed.f90, [272](#)
ysed, [273](#)

zero0.f90, [273](#)
zero1.f90, [273](#)
zero2.f90, [274](#)
zero_urban.f90, [274](#)
zeroini.f90, [274](#)