

PHY64 Experiment 2: The Millikan Oil-drop Experiment

Matthew S. E. Peterson and Jackson Burzynski
Department of Physics and Astronomy, Tufts University

I. INTRODUCTION

In 1910, American physicist Robert Millikan performed the first experiment to determine the elusive value of the electron charge. Using a microscope, Millikan observed small oil droplets as they fell through two horizontal metal electrodes. He first calculated their terminal velocity with no voltage applied to determine the radius of the droplets. Then, using the known density of the oil, Millikan was able to calculate their mass and therefore the gravitational force on each droplet. Next Millikan applied a voltage between the plates to induce an electric field. By observing the motion of a the droplets in several charge states, Millikan showed that the charges were all small integer multiples of a certain base value. Millikan calculated this value to be 1.5924×10^{-19} C, within 1% of the currently accepted value of e .

In this experiment, we measure the electron charge using the same method that Millikan used in 1910. The main device used is a cylindrical chamber manufactured by Pasco Scientific. The bottom half of the device consists of a capacitor with plate separation set by a 7.6 mm thick spacer that serves as a droplet viewing chamber. A small hole at the top of this chamber allows for oil droplets to enter the viewing region, where they are illuminated by an LED and viewed through a microscope. Using an atomizer, oil droplets are deposited into the upper chamber. In the process, the droplets acquire an electric charge. The droplets diffuse from the upper chamber through the small hole at the top of the lower chamber and enter the viewing region. By applying a voltage and observing the velocities of the oil droplets, we may deduce the concentration of charge on each droplet. Finally, we may calculate the difference between these charge states and conclude that they are integer multiples of some constant value. By calculating this value, we will obtain the charge of a single electron.

II. THEORY

Consider an oil droplet of radius a and density ρ falling through the air. The droplet will experience both a gravitational force downward as well as a viscous drag force. The gravitational force is simply $\mathbf{F}_G = -mg\hat{\mathbf{y}}$, where $m = \frac{4}{3}a^3\rho$ and g is the acceleration due to gravity. The viscous drag force is given by Stokes' law as

$$\mathbf{F}_D = 6\pi a\eta v, \quad (1)$$

where η is the viscosity of air. The viscosity of dry air at a temperature T can be approximated by the Sutherland

formula:

$$\eta \approx (1.827 \times 10^{-5}) \left(\frac{T_0 + C}{T + C} \right) \left(\frac{T}{T_0} \right)^{3/2} \text{ N s/m}^2, \quad (2)$$

where $T_0 = 291.15$ K and $C = 120$ K.

Assuming the droplet falls at its terminal velocity, v_f , the net force must be zero. This means that

$$\mathbf{F}_D + \mathbf{F}_G = \left(6\pi a\eta v_f - \frac{4}{3}a^3\rho g \right) \hat{\mathbf{y}} = 0. \quad (3)$$

Using this, we find the radius of the droplet to be

$$a = \sqrt{\frac{9\eta v_f}{2\rho g}}. \quad (4)$$

Now suppose the droplet contains some amount of charge q . If we turn on an electric field $\mathbf{E} = E\hat{\mathbf{y}}$, then the droplet will experience an additional force $\mathbf{F}_E = q\mathbf{E}$. We will assume that field is strong enough that the droplet will begin to rise at some terminal velocity v_r . The viscous force will oppose this motion, and so balancing the forces gives

$$qE - mg - 6\pi a\eta v_r = 0. \quad (5)$$

Using (3), we can see that

$$mg = 6\pi a\eta v_f.$$

Thus, we can rewrite (5) as

$$qE = 6\pi a\eta(v_f + v_r). \quad (6)$$

Solving for q and substituting for a with (4) yields

$$q = \frac{6\pi}{E}(v_f + v_r)\sqrt{v_f} \sqrt{\frac{9\eta^3}{2\rho g}}. \quad (7)$$

Millikan postulated that the terminal velocities are slightly greater than theoretically predicted by a factor of $(1 + b/pa)$, where p is the air pressure and $b = 8.2$ Pa m is an experimentally determined constant. In other words, we experimentally observe a terminal velocity v' such that

$$v' = (1 + b/pa)v,$$

where v is the theoretically predicted value. Since the derivation of (7) uses the predicted values, we must factor in this correction, giving

$$q = \frac{6\pi}{E}(v_f + v_r)\sqrt{v_f} \left(\frac{1}{1 + b/pa} \right)^{3/2} \sqrt{\frac{9\eta^3}{2\rho g}}. \quad (8)$$

This is the equation that will be used to calculate the charge on the droplets.

III. RESULTS

IV. ERROR

V. CONCLUSION