

Optimization of integration time and distance cut in the CUPID array

Joe Camilleri


DNP October 2021

Mini-Symposium: Neutrinos and Nuclei XII: Double Beta Decay Analysis Techniques

October 9, 2021

CUPID experiment

- Proposed $0\nu\beta\beta$ search using bolometric array of 1596 lithium molybdate crystals, deployed in the CUORE¹ cryostat.
- Aims to eliminate dominant background of alpha particles present in CUORE.
- **Are new backgrounds introduced with a using a new element for the bolometers?**



Rendering of proposed CUPID array of Li_2MoO_4 crystals

¹Clarke and Braginski 2004.

lithium molybdate

- Li_2MoO_4 crystals allow for discrimination of α backgrounds from $\beta\beta$ events ($Q=3034\text{keV}$) via high-light yield scintillation signals.
- relatively high isotopic abundance of ^{100}Mo (10%)
- enrichment above 95% already demonstrated in CUPID-Mo





Figure: CUPID bolometer and rejection scheme

2ν events and muons

- With respect to coincidences, the rate of $2\nu\beta\beta$ events is not negligible in CUPID array²
- Minimizing the distance cut helps avoid mis-labelling random $2\nu\beta\beta$ coincidences as multiplicity 2.
- Assuming a simple muon veto geometry, increasing the distance cut rejects more muon events.


$$T_{1/2} = 7.1 * 10^8 \text{ yr} \rightarrow A \sim 3 \text{ mHz}$$




²chernyak.

distance cut in the CUPID array


- monte-carlo simulation of 1 million 2.6MeV gamma rays in the crystal volume.
- With this energy, we expect multiple scattering events in the crystals (cite scattering length)
- multiplicity 2 > events are discarded



integration time in the CUPID array




- 100000 uranium 238 events (full chain)



$2\nu\beta\beta$ efficiency simulation

- sensitivity studies
expect on the order of 90
- large distance cut, random coincidences (more variation with integration time)
- is there a paper i can cite. $0\nu\beta\beta$ expect closer to 80
- find operating point relative to muon background, parameter of interest $-j$ at which distance cut do random coincidences play role in integration time



muon background

- muon flux LNGS
- planned muon veto , 90
- muon track + showers induced by the muon. muon background suppression vs efficiency




Figure: