PITCH PERCEPTION OF TIME-VARYING SUNG TONES

Johanna Devaney and David Wessel

SCHOOL OF MUSIC


THE OHIO STATE UNIVERSITY



Task: To study the influence of sloped F₀ trajectories in sung tones on pitch perception Goal: To develop a pitch perception model that accounts for the variation that occurs in sung tones

MOTIVATION

- The trajectory of fundamental frequencies (Fos) in sung notes often exhibit a degree of sloping
- For example, in the image below the moving average (solid line) of the F₀ trajectory (dotted line) shows the trend over the duration of the note is an upwards slope



- · This study helps to assess whether perceived pitch models based on tones with stable F₀ trajectories are appropriate to use for tones with sloping F₀ trajectories
- This project builds on earlier work that described the use of discrete cosine coefficients to describe F₀ trajectories in sung tones (Devaney, Mandel, and Fujinaga 2011)

RELATED WORK

- d'Alessandro and Castellengo (1994, 1995)
 - used short synthesized sung tones with vibrato
 - found that the end of the note was more influential on the pitch perception than the beginning of the note
- d'Alessandro, Rosset, and Piot (1995)
 - used synthesized tones with F₀ glides of various extents and durations
- findings were the opposite of D'Alessandro and Castellengo's finding that the end of the note was more influential on pitch perception than the beginning
- Gockel, Moore, and Carlyon (2001)
 - used sinusoids
 - modeled perceived pitch as a weighted average favoring the more slowly moving portions of a note

METHOD

- Paradigm: Adaptive transformed up-down two-alternative forced-choice (Levitt 1971)
 - · Versus the method-of-adjustment paradigm used in earlier studies
 - Following from Levitt (1971), six reversals were required before the measurement phase began
- Implementation: In MATLAB with psylab (Hansen 2006)
- Stimuli: 750 ms synthesized tones with either no slope or an upward or downward slope of 0.5 semitones/second
 - Vibrato and noise were added to F₀ trajectory
- Initializations: The starting mean pitch of the stimuli was 33 cents higher than mean pitch of reference and the starting step size for the stimuli adjustment was 8 cents, with a minimum step size of 1 cent
- **Trials**: Five trails were run simultaneously and interleaved
 - Flat tone versus flat tone
 - •Slope up versus flat tone
 - Flat tone versus slope up
 - •Slope down versus flat tone
- •Flat tone versus slope down Subjects: 3 subjects

•Ages: [29, 31, 36]

•Years of musical training: [20, 16, 16]

RESULTS Subject 1 Subject 2 Subject 3

REFERENCES

- d'Alessandro, C., and M. Castellengo, 1994. The pitch of short-duration vibrato tones, Journal of the Acoustical Society of America 95 (3):1617-30.
- d'Alessandro, C., and M. Castellengo. 1995. The pitch of short-duration vibrato tones: Experimental data and numerical model. In Vibrato, edited by P. H. Dejonckere, M. Hirano and J. Sundberg. San Diego, CA: Singular Publishing Group.
- d'Alessandro, C., S. Rosset, and O. Piot. Measurement of pitch perception for Fo glides. In Proceedings of the European Conference on Speech Technology, EuroSpeech, pages 957-60, Madrid, September
- Devaney, J., M. Mandel, and I. Fujinaga. 2011. Characterizing singing voice fundamental frequency trajectories. In Proceedings of the IEEE Workshop on Applications of Signal Processing to Acoustics and Audio, 73-6.
- Gockel, H., B. Moore, and R. Carlyon. 2001. Influence of rate of change of frequency on the overall pitch of frequency-modulated tones. Journal of the Acoustical Society of America 109 (2):701-12 Hansen, M. 2006. psylab MATLAB toolkit. http://www.hoertechnik-audiologie.de/psylab/
- Levitt, H. 1971. Transformed up-down procedures in psychoacoustics. Journal of the Acoustical Society of America 49:467-77.

