

Neurooncol. Author manuscript; available in PMC 2006 November 7.

Published in final edited form as:

J Neurooncol. 2006 January; 76(2): 131–138.

# Local Intracerebral Administration of Paclitaxel with the Paclimer® Delivery System: Toxicity Study in a Canine Model

Gustavo Pradilla, M.D.<sup>1</sup>, Paul P. Wang, M.D.<sup>1</sup>, Patrik Gabikian, M.D.<sup>1</sup>, Khan Li, M.D.<sup>1</sup>, Carolyn A. Magee, B.S.<sup>2</sup>, Kevin A. Walter, M.D.<sup>3</sup>, and Henry Brem, M.D.<sup>1</sup>,\*

1 Department of Neurosurgery, The Johns Hopkins University, School of Medicine, Baltimore

2 Department of Radiology MD; The Johns Hopkins University, School of Medicine, Baltimore

3Department of Neurological Surgery, University of Pittsburgh Pittsburgh, PA

#### **Abstract**

**Introduction**: Paclitaxel, a microtubule binding agent with potent anti-glioma activity *in vitro*, exhibits poor penetrance to the CNS when delivered systemically. To minimize toxicity and reach therapeutic concentrations in the CNS, paclitaxel was previously incorporated into biodegradable microspheres (Paclimer®), and the efficacy of Paclimer® was determined in a rat model of malignant glioma. In this study we report the safety of intracranial Paclimer® in a canine dose escalation toxicity study to prepare its translation into clinical scenarios.

**Methods**: Twelve normal beagle dogs underwent a right parieto-occipital craniectomy and were randomized to receive either Paclimer® at 2-mg/kg (n=5), empty microspheres at 2-mg/kg (n=1), Paclimer® at 20-mg/kg (n=5), or empty microspheres at 20-mg/kg (n=1). Post-operatively, dogs were observed daily for signs of neurotoxicity. Complete blood counts and plasma levels of paclitaxel were obtained weekly. CSF levels and MRI scans were obtained on days14-120. Paclitaxel concentrations were quantified by LC-MS.

**Results**: Animals treated with 20-mg/kg Paclimer® had minimal paclitaxel levels in plasma (range 0-7.84 ng/ml) and CSF (range 0-1.16 ng/ml). Animals treated with 2mg/kg Paclimer® had undetectable levels of paclitaxel in plasma, CSF was not obtained to minimize animal suffering. All animals exhibited normal behavior and weight gain, and were alive post-operatively through the last day of the study (day 60-120) without signs of neurological toxicity. There was no evidence of systemic toxicity or myelosuppression. MR imaging was comparable between Paclimer® animals and controls. Adverse effects included wound infections and a brain abscess, all of which responded to antibiotic therapy, and one ventriculomegaly due to communicating hydrocephalus.

**Conclusions**: Paclimer®-based delivery of paclitaxel is safe for intraparenchymal delivery at the tested doses in normal dogs.

#### **Keywords**

Paclimer; paclitaxel; taxol, microspheres; brain tumor; glioma; dogs; toxicity; safety

Disclosures:

<sup>\*</sup>Corresponding author:, Henry Brem, M.D., F.A.C.S., Professor of Neurosurgery, Oncology & Ophthalmology, Chairman, Department of Neurosurgery, Johns Hopkins University School of Medicine, 725 N. Wolfe Street, Hunterian 817, Baltimore, MD 21205, Office: 410-614-0477, 410-614-0478 (fax), E-mail: hbrem@jhmi.edu

Under separate licensing agreements between the Johns Hopkins University and Guilford Pharmaceuticals, Inc. and the Johns Hopkins University and Angiotech Pharmaceuticals, Inc., Dr. Brem is entitled to a share of royalty received by the University on sales of products described in this article. Dr. Brem owns Guilford Pharmaceuticals, Inc. stock, which is subject to certain restrictions under University policy. Dr. Brem also is a paid consultant to Guilford Pharmaceuticals, Inc. The terms of this arrangement are being managed by the Johns Hopkins University in accordance with its conflict of interest policies.

# Introduction:

Despite significant advances in diagnostic modalities and therapeutic interventions, the prognosis for patients with malignant gliomas remains poor (1,2). The restricted permeability of the blood-brain barrier (BBB) permits only the entry of small hydrophobic molecules, a limited set of actively transported molecules such as glucose and particular amino-acids, and some macromolecules (3). Ideal drug candidates tend to be smaller molecules with simple structures, increased plasma availability, and high lipid solubility (4,5). These characteristics limit the spectrum of effective antineoplastic agents that can achieve interstitial therapeutic levels in the central nervous system (CNS) with acceptable systemic toxicity (3,6,7).

The development of controlled-release polymers for local delivery of drugs to the brain bypasses the limitations of the BBB, minimizes systemic toxicity, and enhances local concentrations of the agents delivered (3,6). Biodegradable controlled-release polymers loaded with carmustine have been established as a safe and effective therapy for patients with malignant gliomas (8-12). Given the inherent advantages of locally-delivered chemotherapy, there has been continued interest in the development of additional controlled-release drug delivery systems suitable for intraparenchymal administration of novel antineoplastic compounds.

Paclitaxel is a microtubule-binding agent that has proven to be clinically efficacious for the treatment of various human neoplasias including ovarian and breast cancers when administered systemically (13-18). Paclitaxel exhibits potent cytotoxic activity against malignant gliomas *in vitro*, but has failed to prolong survival of brain tumor patients in clinical trials when administered systemically (19-21). Systemic administration of paclitaxel also induces a dose-dependent toxicity that ranges from sensory neuropathy and gastrointestinal disturbances to severe myelosuppression (22-25). The lack of efficacy observed with systemic paclitaxel in patients with brain tumors could be attributed to the low permeability of the BBB to paclitaxel and the resultant inadequate bioavailability of paclitaxel in the CNS at maximally tolerated systemic doses of the drug (26-28).

The results of a phase II study of locally-delivered paclitaxel using convection-enhanced drug delivery (CEDD) for treatment of patients with recurrent glioblastoma multiforme were recently published and showed that paclitaxel administered via CEDD had marked anti-tumor responses but generated a significant incidence of treatment-associated complications (29). The delivery of paclitaxel via CEDD and its effect within the tumor can be monitored clinically using diffusion-weighted magnetic resonance imaging (DW-MRI)(30). In this study paclitaxel was administered to 15 patients, 3 patients received a dose of 7.2 mg/day and the remaining 12 patients received a dose of 3.6 mg/day. Complications reported by the authors included chemical and bacterial meningitis, ventriculomegaly, subdural empyema, poor wound healing, and increased peri-tumoral edema. These studies are encouraging and validate the use of local delivery for intratumoral paclitaxel administration.

In order to achieve local sustained administration of paclitaxel into the CNS, we had previously incorporated paclitaxel into a biodegradable polymer matrix consisting of a polyphosphoester polymer (polilactofate) in a microsphere form with a loading concentration of 10% paclitaxel by weight (Paclimer® delivery system)(27). In that study we found that Paclimer® was capable of delivering paclitaxel in a sustained fashion for up to 90 days *in vitro*. There was no local or systemic toxicity in the rat brain and Paclimer®-delivered paclitaxel prolonged the survival of rats challenged intracranially with 9L gliosarcoma (27).

The results of these preclinical studies further support the role of local-delivery of paclitaxel in brain tumor therapy and confirm the need for a safer and more effective system for

intracranial paclitaxel delivery. Therefore we conducted the present study to establish the safety of intracerebral Paclimer® when administered at doses of 2 and 20 mg/kg in the dog brain.

#### Materials and methods:

# Formulation of paclitaxel microspheres

The Paclimer® delivery system, a controlled release formulation of paclitaxel, was obtained from Guilford Pharmaceuticals, Inc. (Baltimore, MD). We have previously described the technique for preparation of Paclimer®(27). Briefly, Paclimer® was synthesized by incorporating paclitaxel into the bioerodable polyphosphoester (PPE) polymer, p(DAPG-EOP), in the form of microspheres at 10% loading by weight. Both substances were dissolved in ethyl acetate and the resulting solution was pumped through an in-line homogenizer with 0.5% polyvinyl acetate solution in a container with an overhead stirrer. The microspheres were allowed to harden, and the product obtained was filtered and lyophilized. Control empty PPE microspheres were synthesized in a similar fashion.

#### **Animals**

Twelve normal adult male beagle dogs weighing 12-14 kg were obtained from Harlan Inc. (Indianapolis, IN) and housed individually with free access to Baltimore city water and dog chow. All animal protocols were approved by the Animal Care and Use Committee of The Johns Hopkins University School of Medicine.

# **Experimental design**

Animals (n=12) were assigned to two experimental groups. Dogs in the first group (n=6) were surgically implanted with either intracerebral Paclimer® at a dose of 20 mg/kg (n=5) or with the equivalent amount of empty microspheres (n=1). Dogs in the second group (n=6) were surgically implanted with either intracerebral Paclimer® at a dose of 2 mg/kg (n=5), or with the equivalent amount of empty microspheres (n=1). All animals were evaluated daily for signs of neurological or systemic toxicity, and sacrificed at pre-determined time points for complete pathological analysis. Animals in the 20 mg/kg Paclimer® group were sacrificed in a staggered fashion on post-operative days (POD) 59 (n=1), 62 (n=1), 99 (n=1), 101 (n=1), and 125 (n=2). All animals in the 2 mg/kg Paclimer® group were sacrificed on POD 120-123 (Table 1).

### Surgical technique

The dogs were sedated with an intramuscular (IM) injection of acepromazine and transferred to the operating suite. An intravenous (I.V.) line was started for continuous infusion of Lactated Ringer's solution and general anesthesia was induced with I.V. sodium thiopental. The animals were intubated endotracheally for mechanical ventilation. General anesthesia was maintained using inhalation isofluorane with continuous cardiac and respiratory monitoring. Each dog was positioned prone and the scalp over the operative site was shaved, disinfected, and draped in sterile fashion. A small parasagittal incision was made over the right parieto-occipital bone and scalp bleeding was controlled with electrocautery. The underlying connective tissue and muscle were divided with electrocautery, and a high-speed air drill was used to create a 2.0cm parieto-occipital craniectomy. A 1.5-cm dural incision was made to expose the underlying parieto-occipital cortex, and a 1.5-cm corticotomy was made using bipolar cautery and gentle suction. Once hemostasis was achieved, sterile Paclimer® or empty microspheres were placed intraparenchymally and secured by placing a rectangular piece of Surgicel® over the corticotomy site. The surgical field was irrigated and a rectangular piece of Gelfoam® was placed to fit the craniectomy defect. The incision was closed in a layered fashion using interrupted 3.0 Vicryl and running 3.0 Prolene® sutures. After discontinuation of anesthesia, the animals were extubated uneventfully and transferred to the housing facility for observation.

#### **Blood and CSF Sampling and Analysis**

Blood was collected from peripheral veins pre-operatively, as well as weekly during the post-operative period in all animals. The blood samples were analyzed by an automated cell counter for complete blood count and white blood cell (WBC) differential. Animals implanted with 20 mg/kg Paclimer® underwent lumbar or cisternal punctures for collection of cerebrospinal fluid (CSF) on POD 14, 44, 59, 61, 99, 101, and 125 and Gram staining, cell counts, protein concentration, and glucose levels were performed. The blood and CSF samples were also assayed for paclitaxel concentration by liquid chromatography mass spectrometry (LC-MS). In animals implanted with 2 mg/kg Paclimer® CSF was not collected and Paclimer® levels were not established to minimize animal suffering since the levels of paclitaxel obtained from the animals treated with the 20 mg/kg dose were minimal.

#### Radiographic studies

In weeks 2, 6, 8, and 18 post-implantation, the animals were sedated, anesthetized, and intubated as described previously, and magnetic resonance imaging (MRI) with and without gadolinium contrast enhancement was conducted.

#### **Necropsy and Histology**

All animals were sacrificed at the scheduled time points. Before sacrifice, blood and CSF were obtained as described above. Complete necropsies were performed on all animals by a veterinary pathologist and the brain, the meninges, and other organs were fixed in 10% paraformaldehyde and embedded in paraffin for histopathological examination.

#### Results:

# Survival, toxicity, and adverse events

In the 20 mg/kg Paclimer® group, the control animal and four out of five dogs that received the intracranial Paclimer® microspheres survived to the scheduled time points of 30, 60, 90, and 120 post-operative days, and exhibited no signs of neurological or systemic adverse events (Table 1). Two of the animals developed wound infections and partial incisional dehiscence that promptly responded to wound irrigation and a prophylactic seven day course of I.M. ceftriaxone. On POD 56, one of the treatment animals developed signs of increased intracranial pressure (i.e. somnolence, bradycardia) that were secondary to a communicating hydrocephalus, which was initially managed by a large-volume CSF drainage; however, because of ethical considerations, the animal was euthanized on POD 59, one day before his scheduled sacrifice date of POD 60. This complication was likely caused by accidental intraventricular administration of Paclimer®.

In the 2 mg/kg Paclimer® group, the control animal and all five dogs that received the intracranial Paclimer® microspheres survived to their scheduled time points of 120 days (Table 1). One animal implanted with Paclimer® developed transient gait disturbances and signs of meningismus on POD 37. MRI showed a deep brain abscess with significant edema and mass effect that was successfully treated with 4 weeks of I.M. ceftriaxone (Figure 1). The neurological exam of this animal returned to baseline after treatment. Two treatment animals developed wound infections or partial incisional dehiscence that responded to wound irrigation and a prophylactic seven day course of IM ceftriaxone.

Throughout the study period, animals in both groups were weighed at weekly intervals. Animals receiving blank microspheres and those receiving Paclimer® displayed appropriate weight gain regardless of the dose administered.

Myelosuppression following systemic paclitaxel administration is a serious dose-limiting toxicity (25). When compared to the control animal, the animals receiving intracranial Paclimer® at 20 mg/kg did not develop leukopenia (Figure 2-A), anemia (Figure 2-B), or <a href="mailto:thrombocytopenia">thrombocytopenia</a> (Figure 2-C) regardless of the dose administered. Animals treated with 2 mg/kg Paclimer® or control microsphere also had unremarkable hematological values throughout the study.

#### Systemic and CSF concentrations of paclitaxel

Animals in the 20 mg/kg Paclimer® group had minimal amounts of paclitaxel in plasma (range 0-4.08 ng/ml) (Figure 3-A). Likewise, the concentration of paclitaxel in the CSF was either minimal or below the level of quantification (<0.5 ng/ml) (Figure 3-B). These measurements were not obtained in the 2 mg/kg Paclimer® group to minimize the number of invasive procedures in the animals.

#### Biocompatibility of Paclimer® in the CNS

The response of the brain parenchyma to Paclimer® and blank microspheres at both doses was followed over time using MRI with and without gadolinium enhancement. In the 20 mg/kg Paclimer® group images showed mild edema surrounding the site of implantation that was comparable between control and Paclimer® animals, which resolved by the last day of follow-up (Figure 4). In the 2 mg/kg Paclimer® mild edema surrounding the implantation site was also observed, which resolved by the last day of follow-up.

Detailed necropsy examination of the animals at 120 days for the 2 mg/kg group, and at 60, 100, and 120 days for the 20 mg/kg group, did not reveal any overt signs of organ toxicity attributable to polilactofate microspheres or paclitaxel. A closer inspection of brains of the control and treatment animals showed an area of fibrous adhesion present between the inner table of the calvarium and the implant site (Figure 5).

Histological examination of meningeal and brain specimens obtained after necropsy near the implant site revealed chronic inflammatory reaction and superficial gliosis that was limited to the parenchyma immediately surrounding the implant site. The extent of this cuff of gliosis and chronic inflammatory response was similar in the control and treatment animals (Figure 5). The brain of the Paclimer®-treated dog that developed communicating hydrocephalus displayed marked ventriculomegaly and mild meningeal hyperplasia.

#### **Discussion:**

In this study we report the safety of intracranial administration of paclitaxel delivered via the Paclimer® system at 2 and 20 mg/kg in dogs. We found that at a dose of 2 mg/kg of Paclimer®, paclitaxel generates no signs of systemic or neurologic toxicity. When Paclimer® was administered at a dose of 20 mg/kg no signs of systemic toxicity were observed, minimal levels of paclitaxel were detected in plasma and CSF, but a neurologic complication occurred in one of the treated animals. This complication was secondary to accidental intraventricular administration of Paclimer®. Even though this type of complication would be amenable to treatment in clinical settings the animal was euthanized to minimize suffering. All other animals treated with 20 mg/kg had no neurologic complications. Clinical studies using CEDD of paclitaxel showed a similar complication, which was attributed to leakage of paclitaxel into the subarachnoid space (29). Accidental intrathecal administration of other anti-microtubule agents has been shown to produce a wide range of complications and should be avoided (31, 32).

Other treatment-related complications found in the animals included wound dehiscence and wound infection. It has been proposed that this complication is caused by subcutaneous leaking

of paclitaxel, which interferes with adequate wound healing (29). In our study one animal in every group experienced suture dehiscence, but a complete response was obtained after wound irrigation and antibiotic therapy. Wound infections were present in one animal of the 2 mg/kg group and in two animals in the 20 mg/kg group (including 1 animal treated with blank microspheres), all of these animals responded to antibiotic therapy. One animal in the 2 mg/kg group presented with a deep brain abscess that was successfully treated with antibiotics, this was most likely a complication resulting from surgical contamination.

Radiological evaluation of the animals showed minimal edema surrounding the site of implantation that resolved by the day of sacrifice in both treatment and control animals for both dose groups. Pathological evaluation of the implanted areas 120 days after implantation of Paclimer® in the 2 mg/kg group and 60, 100, and 120 days in the 20 mg/kg showed changes consistent with the surgical manipulation that were comparable between animals treated with Paclimer® at both doses and animals treated with blank microspheres. No abnormalities were found in the surrounding brain parenchyma in any of the animals in the study. The chronic inflammatory response seen around the implantation area tends to decrease over time as the polymer is degraded. By day 125 most of the polymer has been degraded and absorbed and changes compatible with parenchymal healing are present with minimal chronic inflammatory changes.

The toxicity of systemic administration of paclitaxel has been previously determined by Poirier et al in dogs (33). In that study severe myelosuppression was consistently observed in the animals at clinically relevant doses. Intraparenchymal implantation of Paclimer® microspheres did not induce significant myelosuppression in dogs at the doses administered. As shown in Figure 2, a descending trend in WBC counts and platelets was observed in animals treated with 20 mg/kg of Paclimer® towards the end of the study. This decline, however, was not clinically significant. In order to fully determine additional toxicities evaluation of other parameters such as activity of natural-killer lymphocytes, mitogen responses, and phagocytic activity, is warranted and should be considered during clinical testing. Although complications observed in the animals including wound infections, sutures dehiscence, and brain abscess could be related to immune system dysfunction, at the time of presentation, WBC and platelet counts were within normal limits.

In the present study we selected normal canines over lower species in order to evaluate their neurological status in a more detailed fashion. We acknowledge that the effects from implantation of the Paclimer® system in a normal brain could differ from the effects of implantation in tumor-bearing tissue where tumorigenic edema and hypoxia are present. Furthermore, Paclimer® will be tested clinically in patients that are likely to have received both radiation therapy and cytotoxic chemotherapy, which could also alter the diffusion and absorption of paclitaxel.

The pharmacokinetics of Paclimer® have been previously described *in vitro* (27). The polilactofate microspheres used in this study where selected based on the safety and efficacy studies in the rat brain (27). In previous studies by our group, paclitaxel was initially incorporated into a biodegradable polymer matrix consisting of paclitaxel and a polyanhydride poly[1,3-bis (carboxyphenoxy) propane-co-sebacic-acid] (PCPP:SA) polymer. The safety and efficacy of this formulation were also evaluated in a rodent model of malignant glioma (34). In this study a bi-phasic release profile of paclitaxel generated sporadic toxicity, for this reason polilactofate was preferred over PCPP:SA for paclitaxel delivery. The efficacy studies showed that locally delivered Paclimer® doubled the median survival of F344 rats bearing intracranial 9L gliosarcoma (35 days vs. 16 days; n=10 animals/group; P < 0.001) compared to controls, and that systemic administration of an equivalent dose of Paclitaxel had no effect on animal survival (27).

Paclimer® has also demonstrated beneficial effects against human prostate tumors implanted in nude mice (35). Treatment of the xenografts with intratumoral Paclimer® alone significantly decreased tumor size compared to controls, and showed synergistic activity when combined with radiation therapy (35). This effect has also been observed by our group in intracranial 9L gliosarcomas in rats using a combination of Paclimer® and radiation therapy (36). This effect appears to be related to the radiosensitizing properties of paclitaxel (20,37), which depends on stabilization of tubulin formation that creates a block in the G2-M phase of the cell cycle (the most radiosensitive phases)(35).

Paclimer® is currently undergoing clinical testing for recurrent ovarian cancer and the results of a phase I study are being prepared for publication (Gynecologic Oncology Group, Dr. Deborah Armstrong, personal communication).

In summary, Paclimer® has demonstrated potent anti-glioma effects in animal models and can be administered intracranially at doses ranging between 2 and 20mg/kg with acceptable margins of safety within the limitations of this toxicity study.

#### Acknowledgements:

This work was supported in part by National Cancer Institute grants U19 CA52857 (H.B.), NCI NSRA T32 CA09574 (P.P.W.), and F32 CA83287 (K.A.W.). We would like to thank John Ziewackz and Jonathan Winter for their assistance in this project.

#### References

- 1. Gurney JG, Kadan-Lottick N. Brain and other central nervous system tumors: rates, trends, and epidemiology. Curr Opin Oncol 2001;13:160–166. [PubMed: 11307058]
- Legler JM, Gloeckler Ries LA, Smith MA, Warren JL, Heineman EF, Kaplan RS, Linet MS. RESPONSE: Re: Brain and Other Central Nervous System Cancers: Recent Trends in Incidence and Mortality. J Natl Cancer Inst 1999;91:2050A–22051. [PubMed: 10580036]
- 3. Rautioa J, Chikhale PJ. Drug delivery systems for brain tumor therapy. Curr Pharm Des 2004;10:1341–1353. [PubMed: 15134485]
- Tamargo RJ, Brem H. Drug delivery to the central nervous system: a review. Neurosurgery Quarterly 1992;2:259–279.
- 5. Fung LK, Ewend MG, Sills A, Sipos EP, Thompson R, Watts M, Colvin OM, Brem H, Saltzman WM. Pharmacokinetics of interstitial delivery of carmustine, 4-hydroperoxycyclophosphamide, and paclitaxel from a biodegradable polymer implant in the monkey brain. Cancer Res 1998;58:672–684. [PubMed: 9485020]
- Lesniak MS, Brem H. Targeted therapy for brain tumours. Nat Rev Drug Discov 2004;3:499–508.
   [PubMed: 15173839]
- Moses MA, Brem H, Langer R. Advancing the field of drug delivery: taking aim at cancer. Cancer Cell 2003;4:337–341. [PubMed: 14667500]
- 8. Walter KA, Tamargo RJ, Olivi A, Burger PC, Brem H. Intratumoral chemotherapy. Neurosurgery 1995;37:1128–1145. [PubMed: 8584154]
- 9. Brem H, Piantadosi S, Burger PC, Walker M, Selker R, Vick NA, Black K, Sisti M, Brem S, Mohr G, et al. Placebo-controlled trial of safety and efficacy of intraoperative controlled delivery by biodegradable polymers of chemotherapy for recurrent gliomas. The Polymer-brain Tumor Treatment Group. Lancet 1995;345:1008–1012. [PubMed: 7723496]
- 10. Westphal M, Hilt DC, Bortey E, Delavault P, Olivares R, Warnke PC, Whittle IR, Jaaskelainen J, Ram Z. A phase 3 trial of local chemotherapy with biodegradable carmustine (BCNU) wafers (Gliadel wafers) in patients with primary malignant glioma. Neuro-oncol 2003;5:79–88. [PubMed: 12672279]
- 11. Valtonen S, Timonen U, Toivanen P, Kalimo H, Kivipelto L, Heiskanen O, Unsgaard G, Kuurne T. Interstitial chemotherapy with carmustine-loaded polymers for high-grade gliomas: a randomized double-blind study. Neurosurgery 1997;41:44–48. [PubMed: 9218294]


12. Brem H, Ewend MG, Piantadosi S, Greenhoot J, Burger PC, Sisti M. The safety of interstitial chemotherapy with BCNU-loaded polymer followed by radiation therapy in the treatment of newly diagnosed malignant gliomas: phase I trial. J Neurooncol 1995;26:111–123. [PubMed: 8787853]

- 13. Buda A, Floriani I, Rossi R, Colombo N, Torri V, Conte PF, Fossati R, Ravaioli A, Mangioni C. Randomised controlled trial comparing single agent paclitaxel vs epidoxorubicin plus paclitaxel in patients with advanced ovarian cancer in early progression after platinum-based chemotherapy: an Italian Collaborative Study from the Mario Negri Institute, Milan, G.O.N.O. (Gruppo Oncologico Nord Ovest) group and I.O.R. (Istituto Oncologico Romagnolo) group. Br J Cancer 2004;90:2112–2117. [PubMed: 15150623]
- Garcia AA, O'Meara A, Bahador A, Facio G, Jeffers S, Kim DY, Roman L. Phase II study of gemcitabine and weekly paclitaxel in recurrent platinum-resistant ovarian cancer. Gynecol Oncol 2004;93:493–498. [PubMed: 15099968]
- 15. Garcia AA, Keren-Rosenberg S, Parimoo D, Rogers M, Jeffers S, Koda R, Muggia FM. Phase I and pharmacologic study of estramustine phosphate and short infusions of paclitaxel in women with solid tumors. J Clin Oncol 1998;16:2959–2963. [PubMed: 9738564]
- Guppy AE, Nelstrop AE, Foster T, Agarwal R, Seckl MJ, Rustin GJ. A phase II study of sequential carboplatin, paclitaxel and topotecan in patients with previously untreated advanced ovarian cancer. Br J Cancer 2004;90:810–814. [PubMed: 14970858]
- 17. Nishi T, Yamanaka E, Yamasaki M, Tanaka H, Takeda C, Ikeda N, Hoashi T, Higaki J, Matsunami T, Nakano K, Kogire M, Furukawa J, Inui N, Yamauchi E, Takagaki K, Ogino N, Nomura T, Yamada S, Fujimoto M, Uenishi M, Yayoi E. [Efficacy and safety of weekly taxol (TXL) for advanced recurrent breast cancer evaluated in a multi-center cooperative clinical trial]. Gan To Kagaku Ryoho 2004;31:897–901. [PubMed: 15222108]
- 18. Delfino C, Caccia G, Gonzales LR, Mickiewicz E, Rodger J, Balbiani L, Morales DF, Comba AZ, Brosio C. Gemcitabine plus paclitaxel as first-line chemotherapy for patients with advanced breast cancer. Oncology 2004;66:18–23. [PubMed: 15031594]
- 19. Cahan MA, Walter KA, Colvin OM, Brem H. Cytotoxicity of taxol in vitro against human and rat malignant brain tumors. Cancer Chemother Pharmacol 1994;33:441–444. [PubMed: 7905792]
- 20. Tishler RB, Geard CR, Hall EJ, Schiff PB. Taxol sensitizes human astrocytoma cells to radiation. Cancer Res 1992;52:3495–3497. [PubMed: 1350755]
- 21. Chang SM, Kuhn JG, Robins HI, Schold SC Jr. Spence AM, Berger MS, Mehta M, Pollack IF, Rankin C, Prados MD. A Phase II study of paclitaxel in patients with recurrent malignant glioma using different doses depending upon the concomitant use of anticonvulsants: a North American Brain Tumor Consortium report. Cancer 2001;91:417–422. [PubMed: 11180089]
- 22. Winer EP, Berry DA, Woolf S, Duggan D, Kornblith A, Harris LN, Michaelson RA, Kirshner JA, Fleming GF, Perry MC, Graham ML, Sharp SA, Keresztes R, Henderson IC, Hudis C, Muss H, Norton L. Failure of higher-dose paclitaxel to improve outcome in patients with metastatic breast cancer: cancer and leukemia group B trial 9342. J Clin Oncol 2004;22:2061–2068. [PubMed: 15169793]
- 23. Wist EA, Sommer HH, Ostenstad B, Risberg T, Fjaestad K. Weekly one-hour paclitaxel as first-line chemotherapy for metastatic breast cancer. Acta Oncol 2004;43:11–14. [PubMed: 15068314]
- 24. Hagiwara H, Sunada Y. Mechanism of taxane neurotoxicity. Breast Cancer 2004;11:82–85. [PubMed: 14718798]
- 25. Nieto Y, Vaughan WP. Pharmacokinetics of high-dose chemotherapy. Bone Marrow Transplant 2004;33:259–269. [PubMed: 14647243]
- 26. Fetell MR, Grossman SA, Fisher JD, Erlanger B, Rowinsky E, Stockel J, Piantadosi S. Preirradiation paclitaxel in glioblastoma multiforme: efficacy, pharmacology, and drug interactions. New Approaches to Brain Tumor Therapy Central Nervous System Consortium. J Clin Oncol 1997;15:3121–3128. [PubMed: 9294475]
- 27. Li KW, Dang W, Tyler BM, Troiano G, Tihan T, Brem H, Walter KA. Polilactofate microspheres for Paclitaxel delivery to central nervous system malignancies. Clin Cancer Res 2003;9:3441–3447. [PubMed: 12960135]


28. Glantz MJ, Choy H, Kearns CM, Mills PC, Wahlberg LU, Zuhowski EG, Calabresi P, Egorin MJ. Paclitaxel disposition in plasma and central nervous systems of humans and rats with brain tumors. J Natl Cancer Inst 1995;87:1077–1081. [PubMed: 7616600]


- Lidar Z, Mardor Y, Jonas T, Pfeffer R, Faibel M, Nass D, Hadani M, Ram Z. Convection-enhanced delivery of paclitaxel for the treatment of recurrent malignant glioma: a phase I/II clinical study. J Neurosurg 2004;100:472–479. [PubMed: 15035283]
- 30. Mardor Y, Roth Y, Lidar Z, Jonas T, Pfeffer R, Maier SE, Faibel M, Nass D, Hadani M, Orenstein A, Cohen JS, Ram Z. Monitoring response to convection-enhanced taxol delivery in brain tumor patients using diffusion-weighted magnetic resonance imaging. Cancer Res 2001;61:4971–4973. [PubMed: 11431326]
- Al Ferayan A, Russell NA, Al Wohaibi M, Awada A, Scherman B. Cerebrospinal fluid lavage in the treatment of inadvertent intrathecal vincristine injection. Childs Nerv Syst 1999;15:87–89. [PubMed: 10230662]
- Raney B, Tefft M, Heyn R, Newton W, Jones PM, Haeberlen V, Maurer H. Ascending myelitis after intensive chemotherapy and radiation therapy in children with cranial parameningeal sarcoma. Cancer 1992;69:1498–1506. [PubMed: 1540886]
- 33. Poirier VJ, Hershey AE, Burgess KE, Phillips B, Turek MM, Forrest LJ, Beaver L, Vail DM. Efficacy and toxicity of paclitaxel (Taxol) for the treatment of canine malignant tumors. J Vet Intern Med 2004;18:219–222. [PubMed: 15058774]
- 34. Walter KA, Cahan MA, Gur A, Tyler B, Hilton J, Colvin OM, Burger PC, Domb A, Brem H. Interstitial taxol delivered from a biodegradable polymer implant against experimental malignant glioma. Cancer Res 1994;54:2207–2212. [PubMed: 7909720]
- 35. Lapidus RG, Dang W, Rosen DM, Gady AM, Zabelinka Y, O'Meally R, DeWeese TL, Denmeade SR. Anti-tumor effect of combination therapy with intratumoral controlled-release paclitaxel (PACLIMER microspheres) and radiation. Prostate 2004;58:291–298. [PubMed: 14743469]
- 36. Gabikian, P.; Li, KW.; Tyler, BM.; Brem, H.; Walter, KA. Intracranial delivery of taxol by biodegradable polymer microspheres radiosensitizes malignant gliomas. 2004. In preparation
- 37. Tishler RB, Schiff PB, Geard CR, Hall EJ. Taxol: a novel radiation sensitizer. Int J Radiat Oncol Biol Phys 1992;22:613–617. [PubMed: 1346533]







**Figure 1.** MRI of an animal treated with 2 mg/kg Paclimer® that presented with a deep brain abscess.









**Figure 2.**Panel shows the progression of hematological values in animals treated with 20 mg/kg Paclimer®. Line graphs show the WBC counts (A), hematocrit (B) and platelets counts (C).






**A.** Levels of Paclimer® in plasma from animals treated with 20 mg/kg. **B.** Levels of Paclimer® in CSF from animals treated with 20 mg/kg.

# MRI Paclimer 20 mg/kg animals



**Figure 4.**MRI showing coronal sections of the brain of animals treated with Paclimer® 20 mg/kg (bottom) and blank microspheres (top).



**Figure 5.** Photographs of the brains of animal treated with: Paclimer® 20 mg/kg, Gross specimen (bottom left) and microphotograph of an H&E section (bottom right) and Blank microspheres, gross specimen (top left) and microphotograph of an H&E section (top right).

Table 1

2 mg/kg Paclimer®				
Animal #	Treatment	Length of Survival (days post- implant)	Complications/Adverse Events	Outcome
1	Blank microspheres	120	None	No neurological deficit; scheduled sacrifice
2	Paclimer®	120	None	No neurological deficit; scheduled sacrifice
3	Paclimer®	123	Wound Infection	No neurological deficit; scheduled sacrifice
4	Paclimer®	123	Incisional Dehiscence	No neurological deficit; scheduled sacrifice
5	Paclimer®	123	Brain Abscess	No neurological deficit; scheduled sacrifice
6	Paclimer®	120	None	No neurological deficit; scheduled sacrifice
		20	) mg/kg Paclimer®	
Animal #	Treatment	Length of Survival (days post- implant)	Complications/ Adverse Events	Outcome
1	Blank microspheres	125	Wound Infection	No neurological deficit; scheduled sacrifice
2	Paclimer®	125	None	No neurological deficit; scheduled sacrifice
3	Paclimer®	62	Wound Infection	No neurological deficit; scheduled sacrifice
4	Paclimer®	101	Incisional Dehiscence	No neurological deficit; scheduled sacrifice
5	Paclimer®	59	Communicating Hydrocephalus	Lethargy; scheduled sacrifice
6	Paclimer®	99	None	No neurological deficit; scheduled sacrifice