

NEW DELIVERY SYSTEMS FOR BRAIN TUMOR THERAPY

Eric P. Sipos, MD, and Henry Brem, MD

Therapy of brain tumors has been limited by lack of effective methods of drug delivery. Systemic administration often fails to achieve therapeutic concentrations of drugs within intracranial tumors despite toxic systemic levels. This limitation of drug delivery results from abnormalities in the tumor microvasculature as well as restrictions imposed by the normal physiologic blood-brain barrier.

TUMOR MICROVASCULAR ARCHITECTURE

The microvasculature of solid tumors differs structurally from normal capillary beds in several important ways that restrict drug delivery to neoplastic cells. Tumor microvessels are frequently dilated, saccular, and tortuous. Their organization is highly variable from one region to another within the same tumor, and microvascular branching patterns are often abnormal. Architectural features such as arteriovenous shunts or anastomoses between venules may also contribute to highly variable patterns of blood flow within solid tumors.^{34,35,41} Interstitial pressures are elevated and microvascular pressures are abnormally low, so that many of the tumor microvessels may be partially or completely collapsed.^{33,34} These factors contribute to diminished blood flow measured in brain tumor vessels compared with that in adjacent normal pial vessels and limit the tissue penetration of pharmacologic agents from the bloodstream to the tumor parenchyma.⁸¹

This work was supported by National Cooperative Drug Discovery Group grant no. U01CA5 52857 and by the P20-NS31081 grant from the National Institutes of Health. Dr. Sipos is a recipient of NIH National Research Service Award CA-09574.

From the Departments of Neurosurgical Oncology (EPS) and Neurological Surgery, Ophthalmology, and Oncology (HB), The Johns Hopkins University School of Medicine, Baltimore, Maryland

THE BLOOD-BRAIN BARRIER

The tight junctions of the cerebral capillary endothelium, which form the anatomic basis for the blood-brain barrier, restrict the influx of molecules from the bloodstream into the brain. Although the blood-brain barrier is critical for homeostatic control of the biochemical milieu of the central nervous system, it also restricts access of many potentially beneficial therapeutic agents to the brain parenchyma. Attempts to improve drug delivery by biochemical or osmotic modification of the barrier^{1,15,26,29-31,50} or by modification of pharmacological agents to make them more lipophilic have been moderately successful;⁵⁵ however, these strategies have not yet significantly increased the number of drugs available for treatment of brain tumors.

NEW DELIVERY METHODS

Conventional strategies for improving intravenous drug delivery to brain tumors usually require increasing the dose, frequency, or duration of drug administration. High systemic drug levels are often required to achieve adequate drug concentrations within the central nervous system. This approach is usually limited by systemic toxicity. A variety of alternative techniques have been designed to improve drug delivery to brain tumors and limit systemic toxicity by reducing the volume of distribution of the drug. These techniques include intra-arterial, intrathecal, intraventricular, and interstitial drug administration.⁶⁴

Intra-arterial Therapy

Catheterization of the carotid or vertebral arteries for intra-arterial administration of chemotherapeutic agents such as the nitrosoureas has been used to permit more selective drug delivery to brain tumors. In rat brain tumor models, this technique was shown to permit higher drug concentrations in the brain on the side of the infusion than is possible with intravenous drug delivery.⁷⁸ This permitted the use of significantly lower doses than were necessary for equivalent therapeutic effects using intravenous injections.¹³

Unfortunately, the therapeutic promise of intra-arterial drug delivery has not been supported by clinical trials in patients with metastatic⁴⁴ or primary brain tumors.^{38,60,62,76} In these studies, complications associated with intra-arterial therapy included transient orbital or head pain, periorbital and scleral erythema, retinal toxicity and blindness, and significant leukoencephalopathy. In a phase III study, survival rates in patients treated with intra-arterial carmustine (BCNU) injections were unchanged or worse than those for patients treated with conventional intravenous therapy. Thus, it was concluded that intra-arterial therapy with BCNU is neither safe nor effective.⁶⁰ The toxicity and lack of efficacy of intra-arterial therapy may be related to inhomogeneous mixing of the infused drug in the arterial bloodstream, a phenomenon known as "streaming" that has been demonstrated *in vitro* and *in vivo*.^{2,16,43} The use of infusion techniques that minimize streaming may be of clinical value in the future and warrant further investigation.

Intraventricular and Intrathecal Therapy

Injection of drugs into the lumbar thecal sac, the cerebral ventricles, or the basal cisterns of the brain has been an important method of treatment for patients

with malignancies affecting the cerebrospinal fluid (CSF) spaces or adjacent surfaces of the brain and spinal cord. The CSF-brain barrier, however, prevents significant drug penetration from the CSF spaces into the brain parenchyma; consequently, intrathecal and intraventricular drug infusions have been of limited benefit for treatment of intraparenchymal brain tumors.⁶⁴

In appropriately selected patients, lumbar puncture for intrathecal administration of antineoplastic agents has been safely used for treatment of carcinomatous or leukemic meningitis.³⁷

The Ommaya reservoir, introduced in 1963,⁵⁴ has facilitated the use of intermittent bolus injections of drug into the ventricular CSF, a process that previously required an externalized catheter or repeated direct needle punctures of the lateral ventricles through a frontal burr hole. More recently, continuous sustained intraventricular or intrathecal drug infusions have been made possible by a variety of implantable mechanical or electronic pumps.^{27,45,57}

Interstitial Therapy

The most direct method of by-passing the barriers to drug delivery imposed by the abnormal tumor microvasculature, elevated interstitial pressure, and the normal brain capillary endothelial tight junctions is to administer therapeutic agents directly into the tumor parenchyma. Theroretically, interstitial therapy should maximize drug delivery to the tumor while minimizing systemic toxicity. With this approach to drug delivery, interstitial pressure gradients and drug concentration gradients favor drug redistribution from the center to the periphery of the tumor by a combination of convection and diffusion.³⁴ Interstitial drug delivery is particularly appealing for the treatment of primary brain tumors such as the malignant gliomas because these tumors generally recur locally and systemic metastases are exceedingly rare.²⁸ Approaches to local drug delivery include the use of implantable catheters and implantable controlled-release polymer systems.

Catheter and Pump Systems

Soon after its introduction, the Ommaya reservoir was first used to treat malignant gliomas with methotrexate.⁵⁹ Since that time, catheters with or without pump systems have been used to administer methotrexate,^{23,51,75} nitrosoureas,⁷⁹ bleomycin,^{4,48} adriamycin,³² cisplatin,⁵ and lymphokine-activated killer cells⁴⁹ directly into tumor parenchyma or resection cavities.

Although these studies demonstrate that intratumoral infusion is feasible and may be well tolerated, this has not been proven more effective than less-invasive methods of treatment. The major risks of CNS infusion techniques include infection, catheter obstruction, and neurotoxicity. Intermittent drug infusions may also be limited by the failure to generate an adequate sustained concentration gradient to promote drug penetration into the tumor parenchyma.⁴² To overcome this drug penetration problem, Bouvier and coworkers⁵ reported the implantation of 68 catheters for the delivery of cisplatin into the brain of a patient with a malignant glioma. Although this system halted progression of the disease, the tumor recurred and the patient died 6 months after therapy.⁵ Another strategy for improving drug penetration through tumor and brain parenchyma by using continuous drug infusions to maintain pressure gradients favoring drug convection is being developed and tested at the National Institutes of Health.³

Polymer Systems

A variety of biocompatible controlled release polymers have been developed as alternative systems for interstitial drug delivery. These devices are available in biodegradable and nonbiodegradable forms.

The best example of a nonbiodegradable controlled release polymer is the ethylene-vinyl acetate copolymer (EVA), a biologically inert polymer that releases incorporated drugs by simple diffusion. Although this system has been studied extensively and is used clinically, it has not yet been approved for clinical investigation as a drug delivery device for the brain. The primary limitation of EVA and other nonbiodegradable controlled release polymers is that they remain as foreign bodies in the brain long after drug release has ceased. Nevertheless, EVA has proven invaluable as a research tool permitting proof of the principle that use of controlled release polymers for interstitial drug delivery is feasible, well-tolerated, and effective against experimental brain tumors. Because it is easy to work with, EVA has also been useful in the laboratory as a screening tool to test new pharmacological agents for safety and efficacy against animal brain tumor models.

Biodegradable polymers such as the polyanhydrides release drugs by a combination of drug diffusion and polymer degradation. Because these polymers degrade completely in the presence of water, they do not remain *in vivo* as foreign bodies but rather are hydrolyzed to yield their component monomers. These degradation products are noncytotoxic, nonmutagenic, and nonteratogenic.⁴⁰

Two polyanhydride polymer systems have been studied extensively as vehicles for interstitial drug delivery in the brain. The first of these devices is poly[bis(*p*-carboxyphenoxy) propane-sebacic acid] or *p*(CPP-SA). Incorporation of drugs into this polymer is a simple process that is performed at room temperature. The polymer can be manufactured in a variety of physical forms, such as sheets, rods, microspheres, nanospheres, or wafers and may therefore be implanted by a variety of surgical techniques, including stereotaxy. By modifying the ratio of CPP to sebacic acid (SA), the polymer degradation time can be adjusted according to the desired period of drug release. *p*(CPP-SA) is particularly well-suited for delivery of hydrophobic compounds. In preclinical animal studies and human clinical trials, it has been demonstrated to be biocompatible and safe for use in the brain.^{7-10,65}

A second polyanhydride polymer, fatty acid dimer-sebacic acid (FAD-SA), shares many of the advantages listed for *p*(CPP-SA) but was designed to optimize interstitial delivery of hydrophilic and hydrolytically unstable compounds such as methotrexate, carboplatin, and 4-hydroperoxycyclophosphamide (4-HC).¹⁷ This device has also been extensively tested in animal models^{6,12} and appears to be a promising candidate for human clinical trials.

POLYMER-MEDIATED INTERSTITIAL DELIVERY OF BCNU

The nitrosoureas are a class of alkylating agents that are highly lipid-soluble and essentially nonionized; therefore, they cross the blood-brain barrier and have often been used for the treatment of brain tumors. The single most commonly used and most-effective chemotherapeutic agent for brain tumors has been BCNU. Systemic administration of this drug has provided modest improvements in patient survival, but its efficacy has been limited by myelosuppression, hepatic toxicity, and pulmonary fibrosis.³⁹

Because delivery of BCNU directly into the parenchyma of brain tumors could result in very high local drug concentrations while by-passing the systemic circulation, it was postulated that interstitial BCNU therapy via implantable polymers

would maximize this drug's proven efficacy by eliminating or minimizing the constraints imposed by its systemic toxicity. This hypothesis has been rigorously tested in a series of preclinical investigations and clinical trials.

Once the biocompatibility of EVAc and *p*(CPP-SA) was established,^{8,65} BCNU was incorporated into these polymers and the distribution and pharmacokinetics of active drug release were determined in vitro and in vivo. These studies demonstrated that controlled, sustained release of intact BCNU is possible by use of these polymer systems, and that high local concentrations of the drug in brain tissue can be achieved with these implants.^{25,77,80} Subsequent preclinical studies have shown that interstitial delivery of BCNU from polymers is well-tolerated in the brains of rodents⁶⁷ and primates¹¹ and that concomitant radiation therapy did not lead to any adverse effects.¹¹ These implants effectively reduce the rate of tumor growth and prolong survival in the rat 9L-gliosarcoma model.⁶⁷

On the basis of the preclinical studies, phase I-II clinical trials of the effect of interstitial BCNU incorporated into the *p*(CPP-SA) polymer were conducted in 21 patients with recurrent malignant gliomas. In these patients, up to eight polymer wafers containing BCNU were implanted into the surgical resection cavity at the time of tumor debulking (Fig. 1 through 3). Three increasing doses of BCNU were studied and all were found to be well-tolerated and safe in the brain.⁹

A phase III randomized, double-blind, placebo-controlled clinical trial of *p*(CPP-SA) containing 3.8% BCNU by weight was conducted at 27 medical centers in 222 patients with recurrent malignant gliomas. In this study, the BCNU-containing polymers significantly prolonged patient survival. Furthermore, in marked contrast to the many toxic effects commonly experienced by patients treated with systemic or intraarterial BCNU therapy, no significant adverse effects were associated with these implants.¹⁰

On the basis of the safety and significant efficacy demonstrated in the clinical trials of 3.8% BCNU in *p*(CPPA-SA), future studies to evaluate higher doses of BCNU in the polymer are planned. The role of these polymers as part of the initial treatment of gliomas is also being evaluated.




Figure 1. A dime-sized polyanhydride polymer disc containing BCNU is inserted into the surgical resection cavity after removal of a malignant glioma.




Figure 2. After completion of surgical resection of a malignant glioma, the walls of the tumor cavity are seen lined with polymer discs containing BCNU.

OTHER DRUG-POLYMER COMBINATIONS

The safety and efficacy with which the polyanhydride polymer permits intracranial BCNU administration suggest that this drug delivery method could be used for a wide array of therapeutic agents that have not previously been successfully used for treatment of brain tumors or other central nervous system diseases.



Figure 3. Nonenhanced T1-weighted MR imaging of a patient after resection of a malignant glioma and implantation of polymers containing BCNU. The polymer discs appear as linear low signal intensity objects on this imaging sequence (*arrows*).

Chemotherapeutic Drugs

A variety of other chemotherapeutic agents, including 4-HC,^{12,36} carboplatin,^{52,53} camptothecin (National Cancer Institute, Bethesda, MD),⁷³ and paclitaxel (Taxol)⁷² have been safely delivered intracranially in implantable polymers and effectively improve treatment of brain tumors in rats. 4-HC is a hydrophilic derivative of cyclophosphamide that does not cross the blood-brain barrier. Unlike its parent compound, 4-HC is appropriate for intracranial delivery because it does not require hepatic activation. Carboplatin is a derivative of cisplatin that was selected for intracranial use because of reduced neurotoxicity and adverse systemic effects compared with its parent compound.⁵³ Camptothecin is a naturally occurring inhibitor of topoisomerase I that has potent antitumor effects *in vitro*. Unexpected toxicity and lack of efficacy against melanoma²⁴ and gastrointestinal carcinoma⁴⁷ in early clinical trials, however, have prevented systemic use of this drug in patients with malignant brain tumors and justify consideration of local delivery. Paclitaxel (Taxol) is a microtubule binding agent with demonstrated clinical efficacy against breast, ovarian, and non-small cell lung carcinoma.⁵⁸ Because it is effective *in vitro* against glioma cell lines¹⁴ but does not cross the blood-brain barrier, paclitaxel (Taxol) is another promising agent for use interstitially.

Each of these agents prolongs survival in rodent brain tumor models when implanted interstitially using biodegradable polymers.^{12,36,52,53,72,73} Preclinical safety and drug distribution studies are underway in primates in preparation for possible clinical trials.

Anticonvulsants

The use of phenytoin as an anticonvulsant is associated with a variety of adverse systemic effects, including cutaneous rashes, hepatitis, and megaloblastic anemia. Because more than 90% of the circulating drug is bound to plasma proteins, it crosses the blood-brain barrier poorly. Because of fluctuating hepatic clearance and erratic drug absorption, it is often difficult to maintain proper therapeutic drug levels in plasma. For these reasons, phenytoin is a good candidate for local drug delivery in patients who undergo craniotomy.

Chronic seizure disorders in rats may be produced by applying cobalt chloride to the cerebral cortex. In this model, interstitial delivery of phenytoin safely and effectively reduced the incidence of seizures as measured by the overall EEG grade, the total number of seizure spikes on the EEG, and the Racine behavioral scale. The polymers released phenytoin in a sustained fashion with no apparent behavioral toxicity over the course of 11 months.⁶⁸

Steroids for Vasogenic Edema

Administration of high systemic doses of corticosteroids is widely used to control the vasogenic edema associated with brain tumors. The prolonged use of therapeutic doses of steroids, however, is associated with serious adverse effects, including hemorrhagic gastric ulcers, atrophic skin changes, diabetes, myopathy, and osteoporosis.⁴⁶ It was hypothesized that interstitial delivery of dexamethasone would more safely control peritumoral edema. To test this hypothesis, *in vitro* and *in vivo* pharmacokinetic studies of dexamethasone release from polymer implants were first conducted. These studies demonstrated that, in contrast to systemic injections, interstitial therapy resulted in higher steroid concentrations in the brain and

lower concentrations in plasma.⁵⁶ Comparable results were obtained in tumor-bearing rats:⁶⁹ controlled-release polymers deliver biologically active dexamethasone in a sustained fashion and vasogenic edema is effectively reduced by using these implants in the rat 9L-gliosarcoma model.⁶⁹

Inhibitors of Tumor Angiogenesis

The growth of solid tumors in the brain, as in other locations throughout the body, is dependent on the ability of neoplastic cells to induce new vessel growth from the host vasculature, a process known as tumor angiogenesis.²¹ By inhibiting vascular proliferation, it is therefore possible to modulate the growth of these neoplasms.¹⁸⁻²⁰

Cortisone, when administered in combination with heparin, has potent antiantiogenic effects.²² This property is preserved when heparin and cortisone are loaded into polymers for interstitial delivery, and when implanted into rat 9L-gliosarcoma, the polymers inhibit tumor growth.⁶⁶ These experiments established that polymer-mediated interstitial antiangiogenic therapy is feasible and effective in controlling tumor growth.

Minocycline, a semisynthetic derivative of tetracycline, is another agent that has been shown to inhibit tumor angiogenesis.⁶³ When delivered systemically, minocycline effectively prolongs the survival of mice with subcutaneously implanted Lewis lung carcinoma.^{61,70,71} For efficacy against established intracranial tumors in the rat 9L-gliosarcoma, however, minocycline must be delivered interstitially.⁷⁴ This effect is potentiated by the concurrent administration of systemic BCNU.⁷⁴

FUTURE APPLICATIONS

As with other malignant neoplasms, a human glioma is a heterogeneous population of cells that is diffusely infiltrative and tends to recur locally. No single therapeutic approach is likely to be curative in the foreseeable future. Instead, the most successful strategies will use rational combinations of synergistic or complementary treatment modalities. Drug delivery polymers are particularly important because they permit treatment of brain tumors and other central nervous system pathology with a diverse array of agents that previously could not be efficiently, safely, or effectively delivered to the brain. This new route of administration will allow use of combinations of new and established chemotherapeutic agents along with modulators of drug resistance, antiangiogenic agents, cytokines, immunotoxins, anticonvulsants, or inhibitors of vasogenic edema. Interstitial chemotherapy may soon be used to enhance the efficacy of systemically administered drugs by locally disrupting the blood-brain barrier. Radiosensitizing agents, implanted locally at the time of tumor resection, could potentiate the antineoplastic effects of postoperative external beam irradiation.

We anticipate that in the future, patients with brain tumors will first undergo biopsy. The tumor specimen will then be studied in the laboratory to characterize the lesion fully, including its patterns of drug resistance. At the time of tumor resection, the specific best therapeutic combinations for that patient's tumor will be administered by polymer implantation. Adjuvant therapy will be administered postoperatively. If the tumor recurs, the same cycle of diagnostic and therapeutic interventions will be repeated. The new biopsy sample will be studied to determine whether the tumor has acquired new patterns of drug resistance since it was first characterized. This will permit selection of the optimal interstitial agents for treatment of the recurrence. With the use of this treatment approach (Fig. 4), im-




Figure 4. Proposed future clinical approach to management of patients with malignant brain tumors.

plantable polymer drug delivery systems will permit highly individualized therapy for patients with malignant brain tumors.

ACKNOWLEDGMENTS

The authors thank Dr. Pamela Talalay for reviewing the manuscript.

References

1. Black KL, Chio CC: Increased opening of blood-tumour barrier by leukotriene C4 is dependent on size of molecules. *Neurol Res* 14:402-404, 1992
2. Blacklock JB, Wright DC, Dedrick RL, et al: Drug streaming during intra-arterial chemotherapy. *J Neurosurg* 64:284-291, 1986
3. Bobo RH, Laske DW, Akbasak A, et al: Convection-enhanced delivery of macromolecules in the brain. *Proc Natl Acad Sci USA* 91:2076-2080, 1994
4. Bosch DA, Hindmarch T, Larsson S, et al: Intraneoplastic administration of bleomycin in intracerebral gliomas: A pilot study. *Acta Neurochir (Wien)* 30 (suppl):441-444, 1980
5. Bouvier G, Penn RD, Kroin JS, et al: Direct delivery of medication into a brain tumor through multiple chronically implanted catheters. *Neurosurgery* 20:286-291, 1987
6. Brem H, Domb A, Lenartz D, et al: Brain biocompatibility of a biodegradable controlled release polymer consisting of anhydride copolymer of fatty acid and sebacic acid. *J Control Rel* 19:325-330, 1992
7. Brem H, Ewend MG, Greenhoot J, et al: The safety of interstitial chemotherapy with BCNU-loaded polymer in the treatment of newly diagnosed malignant gliomas: A multi-institutional study with three year follow-up. *In Annual Meeting of the American Association of Neurological Surgeons*, Orlando, FL 1995
8. Brem H, Kader A, Epstein JJ, et al: Biocompatibility of a biodegradable, controlled-release polymer in the rabbit brain. *Sel Cancer Ther* 5:55-65, 1989
9. Brem H, Mahaley MSJ, Vick NA, et al: Interstitial chemotherapy with drug polymer implants for the treatment of recurrent gliomas. *J Neurosurg* 74:441-446, 1991
10. Brem H, Piantadosi S, Burger PC, et al: Intraoperative controlled delivery of chemotherapy by biodegradable polymers: Safety and effectiveness for recurrent gliomas evaluated by a prospective, multi-institutional placebo-controlled clinical trial. *Lancet* 345:1008-1012, 1995
11. Brem H, Tamargo RJ, Olivi A, et al: Biodegradable polymers for controlled delivery of chemotherapy with and without radiation therapy in the monkey brain. *J Neurosurg* 80:283-290, 1994
12. Buahin KG, Judy KD, Hartke C, et al: Controlled release of 4-hydroperoxycyclophosphamide from the fatty acid dimer sebacic acid copolymer. *Polymers for Advanced Technologies* 3:331, 1992
13. Bullard DE, Bigner SH, Bigner DD: Comparison of intravenous versus intracarotid therapy with 1,3-bis(2-chloroethyl)-1-nitrosourea in a rat brain tumor model. *Cancer Res* 45:5240-5245, 1985
14. Cahan MA, Walter KA, Colvin OM, et al: Cytotoxicity of taxol in vitro against human and rat malignant brain tumors. *Cancer Chemother Pharmacol* 33:441-444, 1994
15. Chio CC, Baba T, Black KL: Selective blood-tumor barrier disruption by leukotrienes. *J Neurosurg* 77:407-410, 1992
16. Dedrick RL: Arterial drug infusion: pharmacokinetic problems and pitfalls. *J Natl Cancer Inst* 80:84-89, 1988
17. Domb A, Bogdanský S, Olivi A, et al: Controlled delivery of water soluble and hydrolytically unstable anti-cancer drugs for polymeric implants. *Polymer Preprints* 32:219-220, 1991
18. Folkman J: Tumor angiogenesis: Therapeutic implications. *N Engl J Med* 285:1182-1186, 1971
19. Folkman J: Anti-angiogenesis: New concept for therapy of solid tumors. *Ann Surg* 175:409-416, 1972
20. Folkman J: Tumor angiogenesis: A possible control point in tumor growth. *Ann Intern Med* 82:96-100, 1975
21. Folkman J: What is the evidence that tumors are angiogenesis dependent? *J Natl Cancer Inst* 82:4-6, 1990
22. Folkman J, Langer R, Linhardt R, et al: Angiogenesis inhibition and tumor regression caused by heparin or a heparin fragment in the presence of cortisone. *Science* 221:719-725, 1983
23. Garfield J, Dayan AD: Postoperative intracavitary chemotherapy of malignant gliomas: A preliminary study using methotrexate. *J Neurosurg* 39:315-322, 1973
24. Gottlieb J, Luce J: Treatment of malignant melanoma with camptothecin (NCS-100880). *Cancer Chemotherapy Reports* 56:103-105, 1972

25. Grossman SA, Reinhard C, Colvin OM, et al: The intracerebral distribution of BCNU delivered by surgically implanted biodegradable polymers. *J Neurosurg* 76:640–647, 1992
26. Gumerlock MK, Belshe BD, Madsen R, et al: Osmotic blood-brain barrier disruption and chemotherapy in the treatment of high grade malignant glioma: Patient series and literature review. *J Neurooncol* 12:33–46, 1992
27. Harbaugh RE, Saunders RL, Reeder RF: Use of implantable pumps for central nervous system drug infusions to treat neurological disease. *Neurosurgery* 23:693–698, 1988
28. Hochberg FH, Pruitt A: Assumptions in the radiotherapy of glioblastoma. *Neurology* 30:907–911, 1980
29. Inamura T, Black KL: Bradykinin selectively opens blood-tumor barrier in experimental brain tumors. *J Cereb Blood Flow Metab* 14:862–870, 1994
30. Inamura T, Nomura T, Bartus RT, et al: Intracarotid infusion of RMP-7, a bradykinin analog: A method for selective drug delivery to brain tumors. *J Neurosurg* 81:752–758, 1994
31. Inamura T, Nomura T, Ikezaki K, et al: Intracarotid histamine infusion increases blood tumour permeability in RG2 glioma. *Neurol Res* 16:125–128, 1994
32. Itho Y, Nakazawa S, Ohwaki K, et al: Treatment of malignant brain tumor by local injection of adriamycin. *Neurochirurgia (suppl)*:363, 1981
33. Jain RK: Determinants of tumor blood flow: A review. *Cancer Res* 48:2641–2658, 1988
34. Jain RK: Delivery of novel therapeutic agents in tumors: Physiological barriers and strategies. *J Natl Cancer Inst* 81:570–576, 1989
35. Jain RK: Tumor blood Flow - II: characterization and modulation. *In* Critical Issues in Tumor Microcirculation, Angiogenesis and Metastasis. Harvard Medical School Conference, Cambridge, MA, 1994
36. Judy K, Olivi A, Buahin K, et al: Effectiveness of controlled release of a cyclophosphamide derivative with polymers against rat gliomas. *J Neurosurg*, 82:481–486, 1995
37. Keizer HJ, Pinedo HM: Cancer chemotherapy: Alternative routes of drug administration. A review. *Cancer Drug Del* 2:147–169, 1985
38. Kleinschmidt-DeMasters BK, Geier JM: Pathology of high-dose intraarterial BCNU. *Surg Neurol* 31:435–443, 1989
39. Kornblith P, Walker M: Chemotherapy for malignant gliomas. *J Neurosurg* 68:1–17, 1988
40. Leong KW, D'Amore P, Marletta M, et al: Bioerodible polyanhydrides as drug-carrier matrices. II: Biocompatibility and chemical reactivity. *J Biomed Mater Res* 20:51–64, 1986
41. Less JR, Skalak TC, Sevic EM, et al: Microvascular architecture in a mammary carcinoma: Branching patterns and vessel dimensions. *Cancer Res* 51:265–273, 1991
42. Levin VA, Patlak CS, Landahl HD: Heuristic modeling of drug delivery to malignant brain tumors. *J Pharmacokinet Biopharm* 8:257–296, 1980
43. Lutz RJ, Dedrick RL, Boretos JW, et al: Mixing studies during intracarotid artery infusions in an in vitro model. *J Neurosurg* 64:277–283, 1986
44. Madajewicz S, West CR, Park HC, et al: Phase II study—Intraarterial BCNU therapy for metastatic brain tumors. *Cancer* 47:653–657, 1981
45. Madrid Y, Langer LF, Brem H, et al: New directions in the delivery of drugs and other substances to the central nervous system. *Adv Pharmacol* 22:299–324, 1991
46. Melby JC: Systemic corticosteroid therapy: Pharmacology and endocrinologic considerations. *Ann Intern Med* 81:505–512, 1974
47. Moertel CG, Schutt AJ, Reitemeier RJ, et al: Phase II study of camptothecin (NSC-100880) in the treatment of advanced gastrointestinal cancer. *Cancer Chemotherapy Reports* 65:95–101, 1972
48. Morantz RA, Kimler BF, Vats TS, et al: Bleomycin and brain tumors: A review. *J Neurooncol* 1:249–255, 1983
49. Naganuma H, Kimurat R, Sasaki A, et al: Complete remission of recurrent glioblastoma multiforme following local infusions of lymphokine activated killer cells. *Acta Neurochir (Wien)* 99:157–160, 1989
50. Neuwelt EA, Howieson J, Frenkel EP, et al: Therapeutic efficacy of multiagent chemotherapy with drug delivery enhancement by blood-brain barrier modification in glioblastoma. *Neurosurgery* 19:573–582, 1986
51. Nierenberg D, Harbaugh R, Maurer LH, et al: Continuous intratumoral infusion of methotrexate for recurrent glioblastoma: A pilot study. *Neurosurgery* 28:752–761, 1991
52. Olivi A, Domb A, Dureza C, et al: Treatment of F98 rat glioma with local controlled-release of carboplatin from biodegradable polymer. *In* Annual Meeting of the American Association of Neurological Surgeons, New Orleans, LA, 1991

53. Olivi A, Gilbert M, Duncan KL, et al: Direct delivery of platinum-based antineoplastics to the central nervous system: A toxicity and ultrastructural study. *Cancer Chemother Pharmacol* 31:449–454, 1993
54. Ommaya AK: Subcutaneous reservoir and pump for sterile access to ventricular cerebrospinal fluid. *Lancet* 2:983–984, 1963
55. Pardridge WM, Boado RJ, Black KL, et al: Blood-brain barrier and new approaches to brain drug delivery. *West J Med* 156:281–286, 1992
56. Reinhard CS, Radomsky ML, Saltzman WM, et al: Polymeric controlled release of dexamethasone in normal rat brain. *Journal of Controlled Release* 16:331–340, 1991
57. Rohde T, Buchwald H, Blackshear P: Implantable infusion pumps. In Tyle P (ed): *Drug Delivery Devices, Fundamentals and Applications*. New York, Dekker, 1988, pp 235–260
58. Rowinsky EK, McGuire WP, Donehower RC: The current status of taxol. In Huskins WJ, Perez CA, Young RC (eds): *Principles and Practice of Gynecologic Oncology Updates*. Philadelphia, JB Lippincott, 1993, pp 1–16
59. Rubin RC, Ommaya AK, Henderson ES, et al: Cerebrospinal fluid perfusion for central nervous system neoplasms. *Neurology* 16:680–692, 1966
60. Shapiro WR, Green SB, Burger PC, et al: A randomized comparison of intra-arterial versus intravenous BCNU, with or without intravenous 5-fluorouracil, for newly diagnosed patients with malignant glioma. *J Neurosurg* 76:772–781, 1992
61. Sotomayor EA, Teicher BA, Schwartz GN, et al: Minocycline in combination with chemotherapy or radiation therapy in vitro and in vivo. *Cancer Chemother Pharmacol* 30:377–384, 1992
62. Stewart DJ, Grahovac Z, Hugenholtz H, et al: Combined intraarterial and systemic chemotherapy for intracerebral tumors. *Neurosurgery* 21:207–213, 1987
63. Tamargo RJ, Bok RA, Brem H: Angiogenesis inhibition by minocycline. *Cancer Res* 51:672–675, 1991
64. Tamargo RJ, Brem H: Drug delivery to the central nervous system: a review. *Neurosurgery Quarterly* 2:259–279, 1992
65. Tamargo RJ, Epstein JI, Reinhard CS, et al: Brain Biocompatibility of a biodegradable, controlled-release polymer in rats. *J Biomed Mater Res* 23:253–266, 1989
66. Tamargo RJ, Leong RW, Brem H: Growth inhibition of the 9L glioma using polymers to release heparin and cortisone acetate. *J Neurooncol* 9:131–138, 1990
67. Tamargo RJ, Myseros JS, Epstein JI, et al: Interstitial chemotherapy of the 9L gliosarcoma: Controlled release polymers for drug delivery in the brain. *Cancer Res* 53:329–333, 1993
68. Tamargo RJ, Rossell LA, Tyler BM, et al: Interstitial delivery of diphenylhydantoin in the brain for the treatment of seizures in a rat model. *J Neurosurg* 80:372, 1994
69. Tamargo RJ, Sills AKJ, Reinhard CS, et al: Interstitial delivery of dexamethasone in the brain for the reduction of peritumoral edema. *J Neurosurg* 74:956–961, 1991
70. Teicher BA, Holden SA, Ara G, et al: Potentiation of cytotoxic cancer therapies by TNP-470 alone and with other antiangiogenic agents. *Int J Cancer* 57:920–925, 1994
71. Teicher BA, Sotomayor EA, Huang ZD: Antiangiogenic agents potentiate cytotoxic cancer therapies against primary and metastatic disease. *Cancer Res* 52:6702–6704, 1992
72. Walter KA, Cahan MA, Gur A, et al: Interstitial taxol delivered from a biodegradable polymer implant against experimental malignant glioma. *Cancer Res* 54:2207–2212, 1994
73. Weingart J, Tyler B, Colvin OM, et al: Local delivery of camptothecin as a new treatment for malignant brain tumors. In *Third NIH Drug Discovery and Development Symposium*, San Diego, CA, 1993
74. Weingart JD, Sipos EP, Brem H: The role of minocycline in the treatment of 9L glioma. *J Neurosurg* 82:635–640, 1995
75. Weiss SR, Raskind R: Treatment of malignant brain tumors by local methotrexate. *Int Surg* 51:149–155, 1969
76. West CR, Avellanosa AM, Barua NR, et al: Intraarterial 1,3-bis(2-chloroethyl)-1-nitrosourea (BCNU) and systemic chemotherapy for malignant gliomas: A follow-up study. *Neurosurgery* 13:420–426, 1983
77. Wu MP, Tamanda JA, Brem H, et al: In vivo versus in vitro degradation of controlled release polymers for intracranial surgical therapy. *J Biomed Mater Res* 28:387–395, 1994
78. Yamada K, Ushio Y, Hayakawa T, et al: Distribution of radiolabeled 1-(4-amino-2-methyl-5-pyrimidinyl)methyl-3-(2-chloroethyl)-3-nitrosourea hydrochloride in rat brain tumor: intraarterial versus intravenous administration. *Cancer Res* 47:2123–2128, 1987

79. Yamashima T, Tamashita J, Shoin K: Neurotoxicity of local administration of two nitrosoureas in malignant gliomas. *Neurosurgery* 26:794–800, 1990
80. Yang MB, Tamargo RJ, Brem H: Controlled delivery of 1,3-bis(2-chloroethyl)-1-nitrosourea from ethylene-vinyl acetate copolymer. *Cancer Res* 49:5103–5107, 1989
81. Yuan F, Salehi HA, Boucher Y, et al: Vascular permeability and microcirculation of gliomas and mammary carcinomas transplanted in rat and mouse cranial windows. *Cancer Res* 54:4564–4568, 1994

Address reprint requests to

Henry Brem, MD
Department of Neurological Surgery and Oncology
Hunterian 817
The Johns Hopkins University School of Medicine
725 North Wolfe Street
Baltimore, MD 21205