

## Local anti-angiogenic brain tumor therapies

Eric P. Sipsos<sup>1,2,3</sup> and Henry Brem<sup>3,4,5</sup>

<sup>1</sup>*Division of Neurosurgery, Walter Reed Army Medical Center, Washington D.C., USA;* <sup>2</sup>*Department of Surgery, F. Edward Hébert School of Medicine, Uniformed Services University of the Health Sciences, Bethesda, Maryland, USA;* <sup>3</sup>*Department of Neurosurgery;* <sup>4</sup>*Department of Oncology, and* <sup>5</sup>*Department of Ophthalmology, The Johns Hopkins Medical Institutions, Baltimore, Maryland, USA*

**Key words:** brain tumor therapy, angiogenesis, drug delivery, minocycline, amiloride

### Summary

The critical role of angiogenesis in the growth of solid tumors, including neoplasms of the central nervous system, has provided the impetus for research leading to the discovery of inhibitors of tumor neovascularization. The therapeutic potential of systemically administered antiangiogenic drugs for brain tumors, however, is limited by a variety of anatomic and physiologic barriers to drug delivery. Implantable controlled-release polymers for local drug administration directly into the tumor parenchyma have therefore been developed to achieve therapeutic concentrations of these drugs within the brain while minimizing systemic toxicity. With use of these polymers, successful antiangiogenic therapy for treatment of experimental intracranial malignancies has been achieved. This has been demonstrated with a variety of otherwise unrelated drugs – including the angiostatic steroids, tetracycline derivatives, and amiloride – which modulate collagenase activity, and thus, basement membrane and interstitial matrix metabolism. Controlled-release polymers provide a clinically practicable method of achieving sustained antiangiogenic therapy which can be readily used in combination with other treatment modalities such as cytoreductive surgery, radiation, and cytotoxic chemotherapy.

### Introduction

The growth of solid tumors in three dimensions is critically dependent upon the presence of a blood supply. During the initial, avascular phase of growth, tumors exist as small aggregates of neoplastic cells sustained by simple diffusion for delivery of nutrients and removal of cellular waste products [1]. Avascular tumors grow at a slow, linear pace until they reach the limits of diffusion and an equilibrium is achieved between the rate of cellular proliferation at the periphery and cell death at the center [2]. Within the anterior chamber of the rabbit eye, the upper limit of avascular tumor growth is 1–2 mm in diameter [3]. Similarly, the growth of tumor spheroids *in vitro* plateaus once aggregates of fewer than 10<sup>6</sup> cells reach diameters less than 4 mm [2]. These avascular neoplasms are dormant from a growth perspective and have little metastatic potential; however, they may remain viable for prolonged periods.

Tumors produce diffusable angiogenic factors which activate endothelial cells in nearby established capillaries or venules and induce endothelial proliferation, migration, and new vessel formation through a series of sequential but partially overlapping steps [4–9]. Once vascularized, previously dormant tumors begin to grow rapidly and their volumes increase exponentially [1,2,10]. Mitotic activity within such tumors occurs predominantly within cylindrical clusters around new microvessels [11]. For malignant neoplasms, it is during this phase of tumor growth that metastatic colonies are most commonly detected [4].

The potential therapeutic value of inhibition of tumor angiogenesis, first recognized nearly three decades ago [12], is also supported by a variety of indirect clinical and experimental evidence in neuro-oncology. The angiogenic capacity of brain tumor cells correlates with tumor malignancy [13] and a histologic grading system developed to quantify tumor angiogenesis predicts survival of patients with astrocytomas [14]. On the basis of

this system, the glioblastoma multiforme is among the most 'endothelial-rich' tumors studied [15]. Similarly, microvascular density correlates with tumor recurrence and patient mortality in children with brain tumors [16]. Other less infiltrative intracranial neoplasms such as meningiomas and hemangiopericytomas are also dependent upon angiogenesis for growth [17]. Cerebrospinal fluid (CSF) from brain tumor patients is up to 10 times as potent in inducing capillary endothelial cell migration than CSF from patients without tumors [18]. Furthermore, in a study of 26 pediatric patients with intracranial tumors, basic fibroblast growth factor, a potent angiogenic factor, was detected in the CSF of 62% of the children studied and this CSF was mitogenic for cultured endothelial cells [16]. Observations such as these provide the rationale for development of anti-angiogenic therapies for treatment of patients with brain tumors.

### **Barriers to systemic drug delivery for intracranial neoplasms**

The normal blood–brain barrier is disrupted in newly-formed tumor vessels. Altered expression of biochemical barrier proteins such as molecular transporters has been shown to correlate with the histologic grade of central nervous system tumors [19]. The permeability of newly induced capillary endothelial cells is also increased as a direct result of their location in the peritumoral environment of malignant glial neoplasms [20]. The contrast enhancement and cerebral edema seen clinically on computerized tomography or magnetic resonance imaging scans of patients with malignant gliomas provides a radiographic correlate of the disruption in the normal blood–brain barrier. Despite the permeability of the tumor microvasculature, however, a variety of structural and physiologic anomalies provide barriers to effective systemic drug delivery for treatment of brain tumor patients.

The microvasculature of malignant brain tumors such as the glioblastoma multiforme differs structurally and physiologically from normal cerebral capillaries in several important ways. Histologically, a normal brain capillary has one or two endothelial cells per lumen when viewed in cross-section. In malignant tumors, however, there may be dramatic endothelial hyperplasia with up to 10 endothelial cells per lumen [15]. These vessels are often dilated, saccular, and tortuous. The microvascular architecture often varies from one intratumoral location to another and abnormal branching

patterns are common. Tumors may contain giant capillaries, arteriovenous shunts, or anastomotic channels between venules [21–23].

Because of the elevated interstitial pressures resulting from increased vascular permeability and the decreased intravascular pressures resulting from the anatomic peculiarities of the tumor microvasculature, many of the small tumor vessels are partially or totally collapsed [24]. These factors probably contribute to the diminished blood flow measured in brain tumor vessels compared to that in adjacent normal pial vessels [25] and may explain the areas of necrosis seen histologically and radiographically in glioblastoma multiforme. Furthermore, these observations predict that systemic administration of drugs for patients with malignant brain tumors would yield inhomogeneous and problematic penetration within the tumor and the surrounding brain.

High systemic drug levels are often required to achieve therapeutic concentrations within the central nervous system. Conventional strategies for improving systemic drug delivery to brain tumors have typically relied upon increased doses, frequency, or duration of drug administration. This approach is generally limited by systemic toxicity. Several alternative methods have therefore been developed in attempts to improve drug delivery to brain tumors [26]. These methods have included chemical modification of pharmacological agents to make them more lipophilic, and biochemical or osmotic modification of the blood–brain barrier. Alternative routes of drug delivery include intraarterial, intrathecal, intraventricular, and interstitial drug administration [27].

### **Controlled-release polymers for interstitial drug delivery**

The most direct method of bypassing the many barriers to drug delivery is to deliver drugs directly into the tumor parenchyma, thus maximizing drug concentrations within the tumor while minimizing systemic toxicity. Theoretically, with local delivery, interstitial pressure gradients and drug concentration gradients would favor redistribution from the center to the periphery of the tumor by a combination of diffusion and convection [28]. Additionally, local drug delivery is appealing for the treatment of malignant gliomas because these tumors usually recur locally and systemic metastases are rare [29]. Examples of local drug delivery strategies include implantable

catheters and implantable controlled-release polymer systems [26].

One of the first applications of controlled-release polymers for *in vivo* drug delivery was to provide a method for testing small quantities of putative inhibitors of angiogenesis in simple experimental models such as the rabbit cornea assay and the chick chorioallantoic membrane assay [30,31]. These models continue to serve as valuable tools in the search for new anti-angiogenic compounds; however, the role of drug delivery polymers has since been translated to therapeutic setting, providing a new method for local delivery of antiangiogenic agents directly within the substance of malignant brain tumors [32,33].

The biodegradable polyanhydride polymer, polybis(p-carboxyphenoxy)propane-sebacic acid (pCPP:SA) copolymer, undergoes slow but complete hydrolytic degradation in the presence of water, thus releasing biologically active drugs incorporated within the polymer matrix gradually over time [27]. Biocompatibility studies of p(CPP:SA) have demonstrated that the parent compound and its degradation products are noncytotoxic, nonmutagenic, and nonteratogenic [34]. Since clinical testing (including prospective, double-blind, placebo-controlled multicenter trials in North America and Europe) demonstrated its safety and efficacy [35–37], this drug delivery polymer has become widely used clinically for the local delivery of carmustine in patients with malignant gliomas. It is currently being studied in the laboratory for local delivery of a variety of antiangiogenic agents [38].

### Local anti-angiogenic therapy

The therapeutic potential of angiogenesis inhibition for treatment of tumors and other diseases has provided a strong impetus for research and has led to the discovery of many new antiangiogenic compounds which have been summarized elsewhere [38]. The antiangiogenic agents reviewed below have been effective when delivered locally using controlled-release polymers.

#### *Cartilage-derived inhibitor*

Cartilage is a normally avascular tissue generally resistant to tumor invasion [39]. The vascularized embryonic precursor of cartilage loses its blood vessels in the early neonatal period [40,41]. Observations such as these led Brem and Folkman to hypothesize

and demonstrate that cartilage produces a potent diffusible inhibitor of tumor angiogenesis [42]. Once isolated and purified, the cartilage extract was incorporated into the nonbiodegradable ethylene-vinyl acetate copolymer and was tested in the rabbit cornea angiogenesis assay [30]. This was the first reported use of a polymer for local delivery of an antiangiogenic compound. Subsequently, the purified cartilage-derived inhibitor was found to be a protein with potent anticollagenase properties and a high sequence homology to a collagenase inhibitor isolated from cultured human skin fibroblasts [43,44]. More recently, a potent and specific cartilage-derived angiogenesis inhibitor was isolated, cloned, purified, sequenced, and identified as troponin I, a subunit of the troponin complex [45].

#### *Heparin and the angiostatic steroids*

The observation that cortisone administered in combination with heparin is antiangiogenic, causes tumor regression, and inhibits tumor metastases [46] led to the discovery of a new class of compounds referred to as the 'angiostatic steroids'. The potency of the antiangiogenic effects of these steroids in the presence of heparin is independent of their relative mineralocorticoid or glucocorticoid activities [47]. Tetrahydrocortisol, for example, is the most potent of the known angiostatic steroids and is completely lacking in glucocorticoid or mineralocorticoid activity [48]. The mechanism of action for the angiostatic steroids appears related to modulation of collagen metabolism [49,50], induction of basement membrane dissolution [51], and promotion of plasminogen activator inhibitor synthesis which in turn reduces synthesis of plasminogen activator, an enzyme involved in remodeling the extracellular matrix and the capillary basement membrane [52,53].

The antitumor effects of angiostatic steroids in the presence of heparin have been confirmed in a series of *in vivo* studies. In a nude mouse model, orally administered heparin and hydrocortisone inhibited angiogenesis and growth of human neurofibrosarcoma xenografts [54]. When incorporated into an implantable controlled-release polymer for local delivery, the combination of heparin and cortisone inhibited angiogenesis in the rabbit cornea assay and inhibited tumor growth in the rat 9L-gliosarcoma model [55]. This study established that local delivery of antiangiogenic agents using implantable biodegradable

polymers was technically feasible and effective. It also demonstrated that these polymers could be used to deliver a combination of drugs simultaneously.

### *Tetracycline derivatives*

As the mechanisms of action for many of the seemingly unrelated inhibitors of angiogenesis become better understood, a recurrent theme involves collagenase inhibition and alteration of basement membrane and extracellular matrix metabolism, critical components of the series of events leading to tumor angiogenesis, cancer invasion, and metastasis. This observation stimulated interest in other commercially available drugs known to have similar effects and led to study of the antitumor effects of the tetracycline derivatives.

The tetracyclines are inhibitors of mitochondrial protein synthesis which have been used for many years as broad-spectrum antibiotics, and more recently have been found to inhibit tumor growth [56]. Although the antineoplastic activity of these drugs were initially attributed to inhibition of mitochondrial protein synthesis within tumor cells, the tetracyclines also inhibit extracellular matrix metalloproteinases [57–59]. Thus, the antitumor effect of the tetracyclines may result in part or entirely from modulation of collagen synthesis and consequently inhibition of angiogenesis.

Of the tetracycline derivatives in commercial use as antibiotics, minocycline is the most lipid-soluble and therefore has the best tissue-penetration [60,61]. It is also a potent inhibitor of collagenase [62] and a potent inhibitor of tumor angiogenesis in the rabbit cornea [63]. Minocycline inhibits the growth of early-passage endothelial cells at concentrations that have minimal effects on the growth of pericytes, smooth-muscle cells, and C6 glioma cells. Deoxyribonucleic acid and protein synthesis in endothelial cells are also selectively inhibited by minocycline. These effects on endothelial cell growth *in vitro* correlate with the anticollagenase potency of minocycline and other tetracycline derivatives rather than their antimicrobial activity [64]. Others have demonstrated that minocycline has minimal cytotoxic effects against tumor cells *in vitro* [65]. These studies lend further support to the hypothesis that the antineoplastic effects of minocycline are mediated by inhibition of angiogenesis.

When delivered systemically, minocycline prolongs survival of mice with subcutaneously implanted Lewis lung carcinoma [65–67]. In the intracranial 9L-glioma model, however, systemic administration

of minocycline was ineffective. In this tumor model, local delivery of minocycline using controlled release polymers significantly improved survival [68].

Col-3 (6-dimethyl-6-deoxy-4-dedimethylaminotetracycline, Metastat), is another non-antimicrobial tetracycline derivative with antineoplastic and antiangiogenic properties. In addition to its potent anticollagenase activity, col-3 is as effective as minocycline in inhibition of brain endothelial cell proliferation [69]. Col-3 is one of several newly developed synthetic matrix metalloproteinase inhibitors currently in clinical trials [70] and is now being studied in a New Approaches to Brain Tumor Therapy (NABTT) clinical consortium trial for patients with malignant gliomas [71]. Although oral or parenteral administration of col-3 may prove effective for extracranial malignancies, the barriers to effective drug delivery summarized above may warrant trials of local delivery for this drug.

### *Amiloride*

Amiloride is another drug which was available commercially for many years before its antiangiogenic properties were recognized. Originally developed as a potassium-sparing diuretic, amiloride was subsequently found to inhibit urokinase-type plasminogen activator (u-PA) [72], a serine protease which has an important role in the proteolytic degradation of the extracellular matrix. In central nervous system gliomas, immunohistochemical staining for u-PA correlates with tumor grade [73,74] and shorter patient survival [74,75] and the genes for both u-PA and its receptor are upregulated [76]. Perhaps through its effects on u-PA, amiloride inhibits primary tumor growth [77] and metastasis [78] and has antiangiogenic effects [79,80]. When administered systemically, amiloride inhibits the growth of rat 9L-gliomas implanted subcutaneously but has no effect on the survival of rats with intracranial 9L-gliomas [81], as one might expect given the poor central nervous system penetration of this drug [82,83]. When treated with drug delivery polymers containing amiloride, however, the median survival of rats with intracranial 9L-gliomas was increased 50% [81].

### **Future applications for antiangiogenic brain tumor therapy**

As a treatment strategy for intracranial or extracranial neoplasms, successful antiangiogenic therapy will

provide tumoristatic rather than tumoricidal effects. Once the antiangiogenic effect is eliminated, tumor growth suppression might no longer be expected. Thus, long-term therapy could be required for any antiangiogenic drug. Recent experimental evidence, however, provides cause for greater optimism [84]. These experiments demonstrated no development of drug resistance to an inhibitor of angiogenesis in three tumor types tested. Furthermore, repeated cycles of treatment resulted in prolonged tumor dormancy after cessation of antiangiogenic therapy [84].

The recent discovery that thalidomide is an orally effective antiangiogenic agent [85] provides an obvious potential solution for clinically achievable long-term therapy of extracranial malignancies and this drug is now being tested clinically. Similarly, clinical trials of col-3 administered as a single daily oral dose are underway as noted above. Development of an effective oral agent for antiangiogenic therapy of intracranial malignancies remains a challenge. It is, however, technically feasible to alter the duration of local drug delivery by changing the composition of implantable controlled-release polymers [86,87]. Thus, long-term local antiangiogenic brain tumor therapy should be possible for these and other promising new antiangiogenic drugs such as Squalamine [88], Angiostatin [89], and Endostatin [90], which are currently in clinical trials for extracranial malignancies. The efficacy of these new drugs against malignant brain tumors has yet to be determined clinically; however, by providing a therapeutic alternative to systemic drug administration, local drug delivery polymers may allow us to use drugs that otherwise would be effective only against extracranial malignancies.

In order to maximize the therapeutic benefits of antiangiogenic therapy, it will be necessary to use these drugs in combination with cytoreductive treatment modalities such as radiation, chemotherapy, and surgical resection. Minocycline, for example, synergistically potentiates the survival benefits achieved with surgical tumor resection or systemic carmustine (BCNU) in the rat intracranial 9L-glioma model [68]. Similarly, minocycline potentiates the therapeutic benefits of radiation and a variety of cytotoxic drugs in the murine Lewis lung carcinoma model [65,66]. Greater than additive effects have also been achieved when different antiangiogenic drugs have been used in combination [91–93]. As we learn more about the inhibitors of angiogenesis, additional synergistic drug combinations may become obvious on the basis of complementary mechanisms of action.

## Acknowledgements

This work was partially supported by a grant from the N.I.H., NCDDG CA52857. Dr. Brem is a consultant to Guilford Pharmaceuticals, Inc. and to Rhone-Poulenc Rorer. The Johns Hopkins University and Dr. Brem own Guilford stock, the sale of which is subject to certain restrictions under University policy. The terms of this arrangement are being managed by the University in accordance with its conflict of interest policies.

## References

1. Knighton D, Ausprunk D, Tapper D, Folkman J: Avascular and vascular phases of tumour growth in the chick embryo. *Br J Cancer* 35: 347–356, 1977
2. Gimbrone M, Leapman S, Cotran R, Folkman J: Tumor dormancy *in vivo* by prevention of neovascularization. *J Exp Med* 136: 261–276, 1972
3. Folkman J, Hochberg M: Self-regulation of growth in three dimensions. *J Exp Med* 138: 745–753, 1973
4. Blood CH, Zetter BR: Tumor interactions with the vasculature: angiogenesis and tumor metastasis. *Biochim Biophys Acta* 1032: 89–118, 1990
5. Folkman J, Brem H: Angiogenesis and inflammation. In: Gallin JI, Goldstein IM, Snyderman R (eds) *Inflammation: Basic Principles and Clinical Correlates*. Raven Press, Ltd., New York, 1992, pp 821–839
6. Folkman J: How is blood vessel growth regulated in normal and neoplastic tissue? G.H.A. Clowes Memorial Award lecture. *Cancer Res* 46: 467–473, 1986
7. Furcht LT: Critical factors controlling angiogenesis: cell products, cell matrix, and growth factors. *Lab Invest* 55: 505–509, 1986
8. D'Amore PA, Thompson RW: Mechanisms of angiogenesis. *Ann Rev Physiol* 49: 453–464, 1987
9. Liotta LA, Steeg PS, Stetler-Stevenson WG: Cancer metastasis and angiogenesis: an imbalance of positive and negative regulation. *Cell* 64: 327–336, 1991
10. Gimbrone M, Cotran R, Leapman S, Folkman J: Tumor growth and neovascularization: an experimental model using the rabbit cornea. *J Natl Cancer Inst* 52: 413–419, 1974
11. Folkman J: Tumor angiogenesis: a possible control point in tumor growth. *Ann Int Med* 82: 96–100, 1975
12. Folkman J: Tumor angiogenesis: therapeutic implications. *N Engl J Med* 285: 1182–1186, 1971
13. Brem H, Thompson D, Long DM, Patz A: Human brain tumors: differences in ability to stimulate angiogenesis. *Surg Forum* 31: 471–473, 1980
14. Brem S: The role of vascular proliferation in the growth of brain tumors. *Clin Neurosurg* 23: 440–453, 1976
15. Brem S, Cotran R, Folkman J: Tumor angiogenesis: a quantitative method for histologic grading. *J Natl Cancer Inst* 48: 347–356, 1972

16. Li VW, Folkerth RD, Watanabe H, Yu C, Rupnick M, Barnes P, Scott RM, Black PM, Sallan SE, Folkman J: Microvessel count and cerebrospinal fluid basic fibroblast growth factor in children with brain tumours. *Lancet* 344: 82–86, 1994
17. Wesseling P, Vandersteenhoven JJ, Downey BT, Ruiter DJ, Burger PC: Cellular components of microvascular proliferation in human glial and metastatic brain neoplasms. A light microscopic and immunohistochemical study of formalin-fixed, routinely processed material. *Acta Neuropathol (Berl)* 85: 508–514, 1993
18. Brem H, Patz J, Tapper D: Detection of human central nervous system tumors: use of migration-stimulating activity of the cerebrospinal fluid. *Surg Forum* 34: 532–534, 1983
19. Guerin C, Wolff JE, Laterra J, Drewes LR, Brem H, Goldstein GW: Vascular differentiation and glucose transporter expression in rat gliomas: effects of steroids. *Ann Neurol* 31: 481–487, 1992
20. Grabb PA, Gilbert MR: Neoplastic and pharmacological influence on the permeability of an *in vitro* blood–brain barrier. *J Neurosurg* 82: 1053–1058, 1995
21. Less JR, Skalak TC, Sevick EM, Jain RK: Microvascular architecture in a mammary carcinoma: branching patterns and vessel dimensions. *Cancer Res* 51: 265–273, 1991
22. Less JR, Skalak TC, Sevick EM, Jain RK: Microvascular network architecture in a mammary carcinoma. *Exs* 61: 74–80, 1992
23. Jain RK: Tumor blood Flow – II: characterization and modulation. *Critical Issues in Tumor Microcirculation, Angiogenesis and Metastasis (conference proceedings)* 1994. Cambridge, Massachusetts. Harvard Medical School and the Massachusetts General Hospital
24. Jain RK: Determinants of tumor blood flow: a review. *Cancer Res* 48: 2641–2658, 1988
25. Yuan F, Salehi HA, Boucher Y, Vasthare US, Tuma RF, Jain RK: Vascular permeability and microcirculation of gliomas and mammary carcinomas transplanted in rat and mouse cranial windows. *Cancer Res* 54: 4564–4568, 1994
26. Sipsos EP, Brem H: New delivery systems for brain tumor therapy. *Neurologic Clinics* 13: 813–825, 1995
27. Tamargo RJ, Brem H: Drug delivery to the central nervous system: a review. *Neurosurg Quarterly* 2: 259–279, 1992
28. Jain RK: Delivery of novel therapeutic agents in tumors: physiological barriers and strategies. *J Natl Cancer Inst* 81: 570–576, 1989
29. Hochberg FH, Pruitt A: Assumptions in the radiotherapy of glioblastoma. *Neurology* 30: 907–911, 1980
30. Langer R, Brem H, Faltermann K, Klein M, Folkman J: Isolation of a cartilage factor that inhibits tumor neovascularization. *Science* 193: 70–72, 1976
31. Langer R, Folkman J: Polymers for the sustained release of proteins and other macromolecules. *Nature* 263: 797–800, 1976
32. Langer R, Murray J: Angiogenesis inhibitors and their delivery systems. *Appl Biochem Biotech* 8: 9–24, 1983
33. Guerin C, Tamargo RJ, Olivi A, Brem H: Brain tumor angiogenesis: drug delivery and new inhibitors. In: Maragoudakis ME, Gullino P, Lelkes PI (eds) *Angiogenesis in Health and Disease*. Plenum Press, New York, 1992, pp 265–274
34. Leong KW, D'Amore P, Marletta M, Langer R: Bioerodible polyanhydrides as drug-carrier matrices. II: Biocompatibility and chemical reactivity. *J Biomed Mater Res* 20: 51–64, 1986
35. Brem H, Mahaley MSJ, Vick NA, Black KL, Schold SCJ, Burger PC, Friedman AH, Ciric IS, Eller TW, Cozzens JW, Kenealy JN: Interstitial chemotherapy with drug polymer implants for the treatment of recurrent gliomas. *J Neurosurg* 74: 441–446, 1991
36. Brem H, Piantadosi S, Burger P, Walker M, Selker R, Vick N, Black K, Sisti M, Brem S, Mohr G, Muller P, Morawetz R, Schold S, Group P-BTT: Placebo-controlled trial of safety and efficacy of intraoperative controlled delivery by biodegradable polymers of chemotherapy for recurrent gliomas. *Lancet* 345: 1008–1012, 1995
37. Valtonen S, Timonen U, Toivanen P, Kalimo H, Kivipelto L, Heiskanen O, Unsgaard G, Kuurne T: Interstitial chemotherapy with carmustine-loaded polymers for high-grade gliomas: A randomized double-blind study. *Neurosurgery* 41: 44–49, 1997
38. Sipsos EP, Brem H: Angiogenesis in nervous system tumors. In: Black PM, Loeffler JS (eds) *Cancer of the Nervous System*. Blackwell Scientific Publications, Inc., Cambridge, Massachusetts, 1997, pp 858–873
39. Eisenstein R, Sorgente N, Soble LW, Miller A, Kuettner KE: The resistance of certain tissues to invasion: penetrability of explanted tissues by vascularized mesenchyme. *Am J Pathol* 73: 765–774, 1973
40. Haraldsson S: The vascular pattern of a growing and full-grown human epiphysis. *Acta Anat* 48: 156, 1962
41. Blackwood HJJ: Vascularization of the condylar cartilage of the human mandible. *J Anat* 99: 551, 1965
42. Brem H, Folkman J: Inhibition of tumor angiogenesis mediated by cartilage. *J Exp Med* 141: 427–439, 1975
43. Moses MA, Sudhalter J, Langer R: Identification of an inhibitor of neovascularization from cartilage. *Science* 248: 1408–1410, 1990
44. Murray JB, Allison K, Sudhalter J, Langer R: Purification and partial amino acid sequence of a bovine cartilage-derived collagenase inhibitor. *J Biol Chem* 261: 4154–4159, 1986
45. Moses MA, Wiederschain D, Wu I, Fernandez CA, Ghazizadeh V, Lane WS, Flynn E, Sytkowski A, Tao T, Langer R: Troponin I is present in human cartilage and inhibits angiogenesis. *Proc Natl Acad Sci USA* 96: 2645–2650, 1999
46. Folkman J, Langer R, Linhardt R, Haudenschild C, Taylor S: Angiogenesis inhibition and tumor regression caused by heparin or a heparin fragment in the presence of cortisone. *Science* 221: 719–725, 1983
47. Crum R, Szabo S, Folkman J: A new class of steroids inhibits angiogenesis in the presence of heparin or a heparin fragment. *Science* 230: 1375–1378, 1985
48. Folkman J, Ingber DE: Angiostatic steroids. Method of discovery and mechanism of action. *Ann Surg* 206: 374–383, 1987
49. Maragoudakis M, Sarmonika M, Panoutsacopoulou M: Antiangiogenic action of heparin plus cortisone is associated with decreased collagenous protein synthesis in the chick

- chorioallantoic membrane system. *J Pharmacol Exp Ther* 251: 679–682, 1989
50. Harada I, Kikuchi T, Shimomura Y, Yamamoto M, Ohno H, Sato N: The mode of action of anti-angiogenic steroid and heparin. *Exs* 61: 445–448, 1992
  51. Ingber D, Madri J, Folkman J: A possible mechanism for inhibition of angiogenesis by angiostatic steroids: induction of capillary basement membrane dissolution. *Endocrinology* 119: 1768–1775, 1986
  52. Ashino-Fuse H, Takano Y, Oikawa T, Shimamura M, Iwaguchi T: Medroxyprogesterone acetate, an anti-cancer and anti-angiogenic steroid, inhibits the plasminogen activator in bovine endothelial cells. *Int J Cancer* 44: 859–864, 1989
  53. Blei F, Wilson EL, Mignatti P, Rifkin DB: Mechanism of action of angiostatic steroids: suppression of plasminogen activator activity via stimulation of plasminogen activator inhibitor synthesis. *J Cell Physiol* 155: 568–578, 1993
  54. Lee JK, Choi B, Sobel RA, Chiocca EA, Martuza RL: Inhibition of growth and angiogenesis of human neurofibrosarcoma by heparin and hydrocortisone. *J Neurosurg* 73: 429–435, 1990
  55. Tamargo RJ, Leong KW, Brem H: Growth inhibition of the 9L glioma using polymers to release heparin and cortisone acetate. *J Neuro-Oncol* 9: 131–138, 1990
  56. Kroon AM, Dontje BHJ, Holtrop M, Van den Bogert C: The mitochondrial genetic system as a target for chemotherapy: tetracyclines as cytostatics. *Cancer Lett* 25: 33–40, 1984
  57. Golub LM, Ramamurthy N, McNamara TF, Gomes B, Wolff M, Casino A, Kapoor A, Zambon J, Ciancio S, Schneir M: Tetracyclines inhibit tissue collagenase activity. A new mechanism in the treatment of periodontal disease. *J Periodontal Res* 19: 651–655, 1984
  58. Golub LM, Wolff M, Lee HM, McNamara TF, Ramamurthy NS, Zambon J, Ciancio S: Further evidence that tetracyclines inhibit collagenase activity in human crevicular fluid and from other mammalian sources. *J Periodontal Res* 20: 12–23, 1985
  59. Golub LM, Ramamurthy NS, McNamara TF, Greenwald RA, Rifkin BR: Tetracyclines inhibit connective tissue breakdown: new therapeutic implications for an old family of drugs. *Crit Rev Oral Biol Med* 2: 297–321, 1991
  60. Zbinovsky V, Chrekian GP: Minocycline. In: Florey K (eds) *Analytical Profiles of Drug Substances*. Academic Press, New York, 1977, pp 323–339
  61. Sande MA, Mandell GL: Antimicrobial agents: tetracyclines, chloramphenicol, erythromycin, and miscellaneous antibacterial agents. In: Gilman AG, Rall TW, Nies AS, Taylor P (eds) *Goodman and Gilman's The Pharmacological Basis of Therapeutics*. Pergamon Press, New York, 1990, pp 1117–1125
  62. Golub LM, Lee HM, Lehrer G, Nemiroff A, McNamara TF, Kaplan R, Ramamurthy NS: Minocycline reduces gingival collagenolytic activity during diabetes. Preliminary observations and a proposed new mechanism of action. *J Periodontal Res* 18: 516–526, 1983
  63. Tamargo RJ, Bok RA, Brem H: Angiogenesis inhibition by minocycline. *Cancer Res* 51: 672–675, 1991
  64. Guerin C, Laterra J, Masnyk T, Golub LM, Brem H: Selective endothelial growth inhibition by tetracyclines that inhibit collagenase. *Biochem Biophys Res Commun* 188: 740–745, 1992
  65. Sotomayor EA, Teicher BA, Schwartz GN, Holden SA, Menon K, Herman TS, Frei E: Minocycline in combination with chemotherapy or radiation therapy *in vitro* and *in vivo*. *Cancer Chemother Pharmacol* 30: 377–384, 1992
  66. Teicher BA, Sotomayor EA, Huang ZD: Antiangiogenic agents potentiate cytotoxic cancer therapies against primary and metastatic disease. *Cancer Res* 52: 6702–6704, 1992
  67. Teicher BA, Holden SA, Ara G, Sotomayor EA, Huang ZD, Chen YN, Brem H: Potentiation of cytotoxic cancer therapies by TNP-470 alone and with other anti-angiogenic agents. *Int J Cancer* 57: 920–925, 1994
  68. Weingart JD, Sipos EP, Brem H: The role of minocycline in the treatment of intracranial 9L glioma. *J Neurosurg* 82: 635–640, 1995
  69. Haroun RI, Bettegowda C, Li K, Brem H: Effect of Col-3, an anti-angiogenesis agent, on rat brain endothelial cell proliferation. Submitted for the Annual Meeting of the American Association of Neurological Surgeons 2000. San Francisco, CA, AANS
  70. Wojtowicz-Praga S: Clinical potential of matrix metalloprotease inhibitors. *Drugs R D* 1: 117–129, 1999
  71. Grossman S, Fisher J, Piantadosi S, Brem H: The new approaches to brain tumor therapy (NABTT) CNS consortium: organization, objectives, and activities. *Cancer Control* 5: 107–114, 1998
  72. Vassalli JD, Belin D: Amiloride selectively inhibits the urokinase-type plasminogen activator. *FEBS Lett* 214: 187–191, 1987
  73. Caccamo DV, Keohane ME, McKeever PE: Plasminogen activators and inhibitors in gliomas: an immunohistochemical study. *Mod Pathol* 7: 99–104, 1994
  74. Hsu DW, Efird JT, Hedley-Whyte ET: Prognostic role of urokinase-type plasminogen activator in human gliomas. *Am J Pathol* 147: 114–123, 1995
  75. Landau BJ, Kwaan HC, Verrusio EN, Brem SS: Elevated levels of urokinase-type plasminogen activator and plasminogen activator inhibitor type-1 in malignant human brain tumors. *Cancer Res* 54: 1105–1108, 1994
  76. Gladson DL, Pijuan-Thompson V, Olman MA, Gillespie GY, Yacoub IZ: Upregulation of urokinase and urokinase receptor genes in malignant astrocytoma. *Am J Pathol* 146: 1150–1160, 1995
  77. Sparks RL, Pool TB, Smith NK, Cameron IL: Effects of amiloride on tumor growth and intracellular element content of tumor cells *in vivo*. *Cancer Res* 43: 73–77, 1983
  78. Kellen JA, Mirakian A, Kolin A: Antimetastatic effect of amiloride in an animal tumour model. *Anticancer Res* 8: 1373–1376, 1988
  79. Alliegro MC, Alliegro MA, Cragoe EJJ, Glaser BM: Amiloride inhibition of angiogenesis *in vitro*. *J Exp Zool* 267: 245–252, 1993
  80. Avery RL, Connor TBJ, Farazdaghi M: Systemic amiloride inhibits experimentally induced neovascularization. *Arch Ophthalmol* 108: 1474–1476, 1990

81. Gibbons MC, Sipos EP, Tyler B, Pinn M, Brem H: The anti-angiogenic agent amiloride inhibits tumor growth and prolongs survival in the rat 9L-glioma model (abstract). Annual Meeting of the Congress of Neurological Surgeons 1995. San Francisco, California
82. Pruitt AW, McNay JL, Dayton PG: Transfer characteristics of triamterene and its analogues: central nervous system, placenta, and kidney. *Drug Metab Dispos* 3: 30–41, 1975
83. Garritsen A, Ijzerman AP, Beukers MW, Cragoe EJJ, Soudijn W: Interaction of amiloride and its analogues with adenosine A1 receptors in calf brain. *Biochem Pharmacol* 40: 827–834, 1990
84. Boehm T, Folkman J, Browder T, O'Reilly MS: Antiangiogenic therapy of experimental cancer does not induce acquired drug resistance. *Nature* 390: 404–407, 1997
85. D'Amato RJ, Loughnan MS, Flynn E, Folkman J: Thalidomide is an inhibitor of angiogenesis. *Proc Natl Acad Sci USA* 91: 4082–4085, 1994
86. Burke M, Langer R, Brem H: Drug delivery to the central nervous system. In: Mathiowitz E (eds) *The Encyclopedia of Controlled Drug Delivery*. John Wiley and Sons, Inc., New York, 1999, pp 184–212
87. Sipos E, Tyler B, Piantadosi S, Burger P, Brem H: Optimizing interstitial delivery of BCNU from controlled release polymers for the treatment of brain tumors. *Cancer Chemother Pharmacol* 39: 383–389, 1997
88. Sills AK, Williams JI, Tyler BM, Epstein DS, Sipos EP, Davis JD, McLane MP, Pitchford S, Cheshire K, Gannon FH, Kinney WA, Chao TL, Donowitz M, Laterra J, Zasloff M, Brem H: Squalamine inhibits angiogenesis and solid tumor growth *in vivo* and perturbs embryonic vasculature. *Cancer Res* 58: 2784–2792, 1998
89. O'Reilly MS, Holmgren L, Shing Y, Chen C, Rosenthal RA, Moses M, Lane WS, Cao Y, Sage EH, Folkman J: Angiostatin: a novel angiogenesis inhibitor that mediates the suppression of metastases by a Lewis lung carcinoma (see comments). *Cell* 79: 315–328, 1994
90. O'Reilly MS, Boehm T, Shing Y, Fukai N, Vasios G, Lane WS, Flynn E, Birkhead JR, Olsen BR, Folkman J: Endostatin: an endogenous inhibitor of angiogenesis and tumor growth. *Cell* 88: 277–285, 1997
91. Brem H, Gresser I, Grosfeld J, Folkman J: The combination of antiangiogenic agents to inhibit primary tumor growth and metastasis. *J Pediatr Surg* 28: 1253–1257, 1993
92. Gagliardi A, Hadd H, Collins DC: Inhibition of angiogenesis by suramin. *Cancer Res* 52: 5073–5075, 1992
93. Wilks JW, Scott PS, Vrba LK, Cocuzza JM: Inhibition of angiogenesis with combination treatments of angiostatic steroids and suramin. *Int J Radiat Biol* 60: 73–77, 1991

*Address for offprints:* Henry Brem, Department of Neurosurgery, The Johns Hopkins School of Medicine, Hunterian 817, 725 North Wolfe Street, Baltimore, MD 21205, USA; Tel.: 410-614-0477; Fax: 410-614-0478; E-mail: hbrem@jhmi.edu