


Chapter 3 The Sun

Content

- 3.1 solar structure and energy
- 3.2 solar radiation
- 3.3 steady and transit condition



The Sun: Energy source of Space

Age = 4.5×10^9 yr

Mass = 1.99×10^{30} kg

Radius = 696,000 km (696 Mm)

Mean density = 1.4×10^3 kg · m⁻³ (1.4 g · cm⁻³)

Mean distance from earth (1 AU) = 150×10^6 km (215 solar radii)

Surface gravity = 274 m · s⁻²

Escape velocity at surface = 618 km · s⁻¹

Radiation emitted (luminosity) = 3.86×10^{26} W (3.86×10^{33} erg · s⁻¹)

Equatorial rotation period = 26 days


Mass loss rate = 10^9 kg · s⁻¹

Effective blackbody temperature = 5,785K

Inclination of sun's equator to plane of earth's orbit = 7°

Composition: approximately 90% H, 10% He, 0.1% other elements
(C, N, O, . . .)

The Sun



- Age: 5×10^9 years
- Mass: $3.3 \times 10^5 M_E$
- Radius: $100 R_E$
- Density: 1.4 g/cm^3
- Gravitation: $2.7 \times 10^2 \text{ m/s}^2$
- Rotation Period: 27 days

$$\bullet M_E = 5.97 \times 10^{27} \text{ Kg}$$

$$\bullet R_E = 6400 \text{ Km}$$

TABLE 3.1. Solar-system, Coronal, and Solar-Wind Compositions



Element ^a	Solar System ^b	Corona ^c	Solar Wind ^d
H	1,350	—	1,900
He	108	72	75
C	0.60	0.41	0.43
N	0.12	0.12	0.15
O	1.00	1.00	1.00
Ne	0.14	0.14	0.17
Na	0.003	0.012	—
Mg	0.053	0.192	—
Al	0.004	0.147	—
Si	0.050	0.176	0.22
S	0.026	0.043	—
Ar	0.005	0.004	0.004
Ca	0.003	0.014	—
Fe	0.045	0.223	0.190
Ni	0.002	0.008	—



^aAbundances referenced to that of oxygen. Only elements with abundances greater than 10^{-3} are listed.

^bValues from Anders and Ebihara (1982).

^cValues derived from measurements of solar-energetic particles (Crook et al., 1984; Breneman and Stone, 1985).

^dValues from Bochsler and Geiss (1989).





- **Core:** energy source (nuclear fusion)
 - $0.3R_{\odot}$, $1.5 \times 10^7 K$
- **Radiative Zone:** scattering
 - $0.5R_{\odot}$, $5.0 \times 10^5 K$
- **Convective Zone:** convection
 - $0.2R_{\odot}$, $6.6 \times 10^5 K$
- **Photosphere:** visible solar disc
 - $0.001R_{\odot}$, $5750 K$
- **Chromosphere:** thin, low temp
 - $10000 K M$, $4500-500000 K$
- **Corona:** Sun's atmosphere
 - $>R_{\odot}$, $1.5 \times 10^6 K$







FIG. 3.2. Overall structure of the solar interior (including the core, the radiative zone, and the convection zone) and the solar atmosphere (i.e., the photosphere, chromosphere, and corona).

TABLE 3.2. Fusion Reactions in the Interior of the Sun



$$m_{^4\text{H}} = 6.693 \times 10^{-27} \text{ kg}$$

$$m_{^4\text{He}} = 6.645 \times 10^{-27} \text{ kg}$$

$$\Delta m = 0.048 \times 10^{-27} \text{ kg}$$

$$6.2 \times 10^{11}$$

The source of the Sun's energy is nuclear fusion

Proton-proton cycle, four protons merges to one α -particle, the mass difference between the four protons and the α -particle corresponds to an energy of $4.3 \cdot 10^{-12} \text{ J}$ or 26.2 MeV.

The first step:

Two protons merge to a deuteron, emitting a positron and a neutrino.

The second step:



A proton collides and merges with the deuteron, forming a ^3He -nuclei under emission of a γ -quant.


The third step:

Two ^3He -nuclei collide, they merge to an α -particle, emitting two protons and a γ -quant .

The time scale of the proton-proton cycle basically is by the first step, half of the hydrogen initially present is converted into deuteron within 10^{10} year.

Although a photon travels with the speed of light, it needs about 100000 years to diffuse from the Sun's core to its surface.











FIG. 3.1. Spectral radiance of the surface of the sun as a function of wavelength.

Solar under various temperatures



EIT195 1.5 million	EIT284 2 million	EIT304 80,000
MDI 6000	EIT171 1million	

The Sun's composition:


hydrogen 92% in terms of particle number or 72% in terms of mass
helium 8%
oxygen 0.06%
carbon 0.03%
the rest <0.01%.


The Sun radiates as a black body at about 6000 K.


The flux of visible solar energy varies very little and, however, the emissions at shorter wavelengths, the EUV and X-ray, vary by orders of magnitude depending on sunspot number and solar activity.

Table 6.1. Properties of the Sun

Radius	$r_{\odot} = 696\,000 \text{ km}$
Mass	$M_{\odot} = 1.99 \times 10^{31} \text{ kg}$
Average density	$\rho_{\odot} = 1.91 \text{ g/cm}^3$
Gravity at the surface	$g_{\odot} = 274 \text{ m/s}^2$
Escape velocity at the surface	$v_{\text{esc}} = 618 \text{ km/s}$
Luminosity	$L_{\odot} = 3.86 \times 10^{23} \text{ kW}$
Magnetic field	
polar	1 G
general	some G
protuberance	10–100 G
sunspot	3 000 G
Temperature	
core	15 million K
photosphere	5780 K
sunspot (typical)	4200 K
chromosphere	4400–10 000 K
transition region	10 000–800 000 K
corona	2 million K
Sidereal rotation	
equator	26.8 d
30° latitude	Differential rotation
60° latitude	28.2 d
75° latitude	30.8 d
	31.8 d








2000/09/22 09:36 UT

Sunspot characteristics:

- (i) Low temperature: 3900°K
- (ii) Intense magnetic field: 200gauss
- (iii) Pair
- (iv) Cycle~11.25 years



Wolf Sunspot number (Solar activity index, **SSN, R**)

Wolf number: $R=k(10^*g+f)$


f: number of individual spot g: number of group

k: comparison constant ~1 Solar min R~0 Solar max R~200

R_{12} : Sum 12 monthly mean of R and then divide by 12

$$R_{12} = (0.5R_{-6} + R_{-5} + R_{-4} + R_{-3} + R_{-2} + R_{-1} + R_0 + R_1 + R_2 + R_3 + R_4 + R_5 + 0.5R_6)/12$$

Near Solar Max - March 2001



2001/03/29 09:36 UT

Near Solar Min - January 2005




2005/01/07 09:50


The 1st cycle: 1755~1766

Now: 24th cycle, 2008.01.04 ~




The Maunder Minimum: 1645~1715




DAILY SUNSPOT AREA AVERAGED OVER INDIVIDUAL SOLAR ROTATIONS



The Sun's rotation – differential rotation



*A sketch of the formation of sunspots and the 22-years sunspot cycle
due to the differential rotation of plasma in the photosphere*



Sunlight and Solar wind




Fig. 2.2 Solar features (observed at various times on different wavelengths) that can have important consequences on the ionosphere (NASA photograph MSFC-5/80-STE 3826)

The solar-terrestrial system

- The Earth's upper atmosphere is ionized by solar radiations
 - (a) electromagnetic: radio - - X-ray; speed 300000 km/s, traveling time 8.3 min.
 - (b) particle (corpuscular): solar wind (H^+ and e^- mainly); speed 300-800 km/s
- Solar terrestrial system: The Sun, the interplanetary medium, and the Earth's magnetosphere, Ionosphere and neutral atmosphere.





Fig. 1.1 The various forms of solar energy and the space environment

- Sunspots: low temp \sim 3000 K, strong magnetic field 0.4 T (4000 G), cycle \sim 11 yrs.
- The period between about 1640 and 1710 is called the Maunder minimum. Solar activity vs. climate.
- The Wolf (or Zurich) sunspot number $R=k(10g+s)$, where g and s are the group and individual spot number, and k denotes the correction factor.
- The spots, after a sunspot min., occur at the latitudes of 20° to 30° north and south whereas at sunspot max. spots occur at $\pm 15^\circ$ and, as the sunspot number declines, spots occur in the latitudes of 5° and 10° .
- The 12-month smoothed relative sunspot number

$$R_{12} = (0.5R_{-6} + R_{-5} + \dots + R_5 + 0.5 R_6)/12$$
- Solar activity index: sunspots, F10.7




The solar wind


The solar wind is a continuous flow of charged particle

Slow solar wind 250-400 km/s, 8 ions/cm³ at 1AU, Streamer belt


Strongly overexpanding flux tubes


Fast solar wind 400-800 km/s, 3 ions/cm³ at 1AU, Coronal hole





Laval's nozzle





$$\frac{dA}{A} = \left(\frac{v^2}{c_s^2} - 1 \right) \frac{dv}{v}$$

$$c_s = \sqrt{\frac{p}{\rho}} = \sqrt{\frac{2k_B T}{m}}$$

Equation of motion:

$$\rho(\vec{u} \cdot \nabla) \vec{u} = -\nabla p - \frac{\rho M_{\Theta} G}{r^2} \quad (4.16)$$

$$u_r \frac{du_r}{dr} = \frac{1}{nm} \frac{d}{dr} (2nk_B T) - \frac{GM_{\Theta}}{r^2} \quad (4.17)$$

Continuity Equation:

$$n(r) \cdot u_r(r) \cdot r^2 = n_0 \cdot u_{r0} \cdot r_0^2 \quad (4.18)$$

$$\frac{du_r}{dr} \left[u_r - \frac{2k_B T}{mv_r} \right] = \frac{2k_B r^2}{m} \frac{d}{dr} \frac{T}{r^2} - \frac{GM_{\Theta}}{r^2} \quad (4.19)$$

$$\left[\frac{v}{v_c} - 1 \right] \frac{dv}{v} = 2 \left[1 - \frac{r_c}{r} \right] \frac{dr}{r}$$

$$r_c = \frac{GM_\Theta m}{4k_B T} = \frac{GM_\Theta}{2c_s}$$

$$u_c = \sqrt{\frac{2k_B T_0}{m}}$$




TABLE 4.2. Solar-Wind Flux Densities and Fluxes near the Orbit of the Earth

	Flux Density	Flux Through Sphere at 1 AU
Protons	$3.0 \times 10^8 \text{ cm}^{-2} \cdot \text{s}^{-1}$	$8.4 \times 10^{35} \text{ s}^{-1}$
Mass	$5.8 \times 10^{-16} \text{ g} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$	$1.6 \times 10^{12} \text{ g} \cdot \text{s}^{-1}$
Radial momentum	$2.6 \times 10^{-9} \text{ pascal (Pa)}$	$7.3 \times 10^{14} \text{ newton (N)}$
Kinetic energy	$0.6 \text{ erg} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$	$1.7 \times 10^{27} \text{ erg} \cdot \text{s}^{-1}$
Thermal energy	$0.02 \text{ erg} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$	$0.05 \times 10^{27} \text{ erg} \cdot \text{s}^{-1}$
Magnetic energy	$0.01 \text{ erg} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$	$0.025 \times 10^{27} \text{ erg} \cdot \text{s}^{-1}$
Radial magnetic flux	$5 \times 10^{-9} \text{ T}$	$1.4 \times 10^{15} \text{ weber (Wb)}$

TABLE 4.3. Some Derived Properties of the Solar Wind near the Orbit of the Earth

Gas pressure	30 pPa
Sound speed	$60 \text{ km}\cdot\text{s}^{-1}$
Magnetic pressure	19 pPa
Alfvén speed	$40 \text{ km}\cdot\text{s}^{-1}$
Proton gyroradius	80 km
Proton–proton collision time	$4 \times 10^6 \text{ s}$
Electron–electron collision time	$3 \times 10^5 \text{ s}$
Time for wind to flow from corona to 1 AU	$\sim 4 \text{ days} = 3.5 \times 10^5 \text{ s}$




FIG. 4.1. Coronal temperature measurements at various heliocentric distances.
(Adapted from Newkirk, 1967.)




FIG. 4.2. Equatorial electron density for 1954 corona. Number density of electrons (cm⁻³) inferred by different observers from data obtained at a single eclipse of the sun. (Adapted from Billings, 1966.)

The Fluid Theory of Solar Wind Formation

$$\rho \frac{\partial \mathbf{u}}{\partial t} + \rho \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla p + \mathbf{j} \times \mathbf{B} + \rho \mathbf{F}_g \quad (\text{momentum})$$

For a plasma flow that is *steady* or independent of time, all time derivatives are zero, and we need deal only with simpler equations:

$$\nabla \cdot \rho \mathbf{u} = 0 \quad (4.1)$$

$$\rho \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla p + \mathbf{j} \times \mathbf{B} + \rho \mathbf{F}_g \quad (4.2)$$

$$\nabla \cdot \rho \mathbf{u} = \frac{1}{r^2} \frac{d}{dr} \rho u r^2$$

$$\rho \mathbf{u} \cdot \nabla \mathbf{u} = \rho u \frac{du}{dr} \hat{\mathbf{e}}_r$$

we can write the flow speed as

$$\mathbf{u} = u(r) \hat{\mathbf{e}}_r$$

and express the sun's gravitational field as

$$\mathbf{F}_g = -\frac{GM_{\odot}}{r^2} \hat{\mathbf{e}}_r$$

where G is the gravitational constant, and M_{\odot} is the mass of the sun.
The pressure gradient is

$$\nabla p = \frac{dp}{dr} \hat{\mathbf{e}}_r$$

$$\frac{1}{r^2} \frac{d}{dr} \rho u r^2 = 0 \quad (4.3)$$

$$\rho u \frac{du}{dr} = -\frac{dp}{dr} - \rho \frac{GM_{\odot}}{r^2} \quad (4.4)$$

$$-\frac{dp}{dr} - \rho \frac{GM_{\odot}}{r^2} = 0 \quad (4.5)$$

$$p = nk(T_e + T_i) = 2nkT$$

Solving for the mass density $\rho = n(m_e + m_p) = nm$, where m is the sum of the proton and electron masses, yields

$$\rho = m \cdot \frac{p}{2kT}$$

Substitution into (4.5), assuming that the temperature T is constant, finally gives the equation for pressure in a *static, isothermal* atmosphere:

$$\frac{1}{p} \frac{dp}{dr} = -\frac{GM_{\odot}m}{2kT} \frac{1}{r^2}$$

The solution to this differential equation for $p(r)$ is

$$\ln p = -\frac{GM_{\odot}m}{2kT} \frac{1}{r} + K$$

where K is an arbitrary constant. Setting $p(r)=p_0$ at the base of the corona, $r=R$, gives the value of the constant

$$K = \ln p_0 - \frac{GM_{\odot}m}{2kT} \frac{1}{R}$$

and allows us to write the solution as

$$\ln \frac{p}{p_0} = -\frac{GM_{\odot}m}{2kT} \left(\frac{1}{r} - \frac{1}{R} \right)$$

or

$$p(r) = p_0 \exp \left\{ -\frac{GM_{\odot}m}{2kT} \left(\frac{1}{r} - \frac{1}{R} \right) \right\} \quad (4.6)$$

For $r > R$, $1/r - 1/R$ is negative, and $p < p_0$; that is, solution (4.6) shows what we would expect, that $p(r)$ decreases with increasing distance from the sun.

that is of interest. The more familiar form can be recovered from (4.6) by measuring r from the base of the corona, in terms of the height

$$h = r - R$$

and considering only $h \ll R$. Then

$$\frac{1}{r} = \frac{1}{R+h} = \frac{1}{R(1+h/R)} \approx \frac{1}{R} \left(1 - \frac{h}{R}\right)$$

In this approximation, equation (4.6) becomes

$$p = p_0 \exp \left\{ \frac{GM_{\odot}m}{2kT} \left(\frac{1}{R} - \frac{h}{R^2} - \frac{1}{R} \right) \right\} = p_0 \exp \left\{ -\frac{GM_{\odot}m}{2kTR^2} \cdot h \right\} = p_0 e^{-h/\lambda} \quad (4.7)$$

where $\lambda = 2kTR^2/GM_{\odot}m = 2kT/mg$ is the *scale height* given in terms of the acceleration due to gravity g at the base of the atmosphere.

The deficiency of equation (4.6) as a model for the equilibrium state of the corona stems from the variation in gravity over the great heliocentric distance spanned by the corona. If we let $r \rightarrow \infty$, the pressure given by (4.6) does not continue to decay exponentially as in (4.7); rather, it approaches the value

$$p_\infty = p_0 \exp \left\{ -\frac{GM_\odot m}{2kTR} \right\}$$

For a coronal temperature of 10^6 K, this is only about e^{-8} or 3×10^{-4} of the high pressure at the base of the corona. It is many orders of magnitude higher than the pressure thought to exist in the interstellar medium ($p_{\text{int}} \approx 10^{-13} - 10^{-14}$ Pa), and thus could not represent an equilibrium state between the corona and that distant medium.

It was this problem that motivated E. N. Parker in the 1950s to reexamine the equilibrium states implied by equations (4.3) and (4.4) by considering solutions with nonzero flow speeds. The first of these equations is satisfied if

$$\rho u r^2 = C \quad (\text{a constant})$$

The meaning of this expression becomes clear if it is multiplied by 4π to give

$$4\pi r^2 \cdot \rho u = I \quad (\text{a constant}) \tag{4.8}$$

The momentum equation (4.4) involves the same two fluid properties, ρ and u , as our first integral (4.8), plus the pressure p . Under the same isothermal assumption made in our analysis of a static equilibrium,

$$p = 2nkT$$

can be differentiated and substituted into (4.4) to give

$$\rho u \frac{du}{dr} = -2kT \frac{dn}{dr} - \rho \frac{GM_{\odot}}{r^2}$$

or, using $\rho = nm$ and dividing through by ρ ,

$$u \frac{du}{dr} = -\frac{2kT}{m} \frac{1}{n} \frac{dn}{dr} - \frac{GM_{\odot}}{r^2} \quad (4.9)$$

This equation can be written with the speed $u(r)$ as the only remaining fluid property by writing (4.8) as

$$4\pi r^2 m n u = I$$

solving for

$$n = \frac{I}{4\pi m} \frac{1}{u r^2}$$

and differentiating to obtain

$$\frac{dn}{dr} = \frac{I}{4\pi m} \left\{ -\frac{1}{u} \cdot \frac{2}{r^3} - \frac{1}{r^2} \frac{1}{u^2} \frac{du}{dr} \right\} \quad (4.10)$$

$$\frac{1}{n} \frac{dn}{dr} = \frac{4\pi m}{I} \cdot u r^2 \cdot \frac{I}{4\pi m} \left\{ -\frac{2}{ur^3} - \frac{1}{r^2 u^2} \frac{du}{dr} \right\} = -\frac{2}{r} - \frac{1}{u} \frac{du}{dr}$$

Substitution into (4.9) then reduces the momentum equation to

$$u \frac{du}{dr} = \frac{4kT}{mr} + \frac{2kT}{m} \frac{1}{u} \frac{du}{dr} - \frac{GM_{\odot}}{r^2} \quad (4.11)$$

This is a differential equation for $u(r)$ and its derivative du/dr in an *expanding, isothermal* atmosphere.

Analysis of equation (4.11) is facilitated by moving all terms involving $u(r)$ to the left-hand side to give

$$u \frac{du}{dr} - \frac{2kT}{m} \frac{1}{u} \frac{du}{dr} = \frac{4kT}{mr} - \frac{GM_{\odot}}{r^2} \quad \left(u^2 - \frac{2kT}{m} \right) \frac{1}{u} \frac{du}{dr} = \frac{4kT}{mr} - \frac{GM_{\odot}}{r^2}$$

$$\left(u^2 - \frac{2kT}{m}\right) \frac{1}{u} \frac{du}{dr} = \frac{4kT}{mr} - \frac{GM_{\odot}}{r^2} \quad (4.12)$$

This form of the momentum equation was recognized by Parker (1958) as revealing the existence of a solar wind. For any realistic coronal temperature T , the second term on the right-hand side of (4.12), GM/r^2 , is larger than the first term, $4kT/mr$, at the base of the corona. This is basically a statement that despite its high temperature, the corona is gravitationally bound. Thus the right-hand side of (4.12) is negative near the base of the corona. However, GM/r^2 falls off more rapidly with r than does $4kT/mr$, so that the right-hand side grows with increasing r , passes through zero at the radius

$$r_c = \frac{GM_{\odot}m}{4kT}$$

$$\left. \frac{du}{dr} \right|_{r_c} = 0$$

Section 4.2, the sound speed in the corona is given by

$$c_s^2 = \frac{\gamma p}{\rho} = \gamma \frac{2nkT}{nm} = \gamma \frac{2kT}{m}$$

For an isothermal plasma, $\gamma = 1$. All other solutions (that start with small u) It is not difficult to write down actual solutions to equation (4.12). The reader can easily verify by differentiation that the expression

$$\frac{1}{2} u^2 - \frac{2kT}{m} \ln u = \frac{4kT}{m} \ln r + \frac{GM_{\odot}}{r} + K'$$

where K' is again an arbitrary constant, is a solution of (4.12). Imposition of the requirement that $u^2 = 2kT/m$ where $r = r_c = GM_{\odot} m / 4kT$ determines the constant K' and gives the explicit form of the special *solar-wind solution*

$$u^2 - \frac{2kT}{m} - \frac{2kT}{m} \ln \frac{mu^2}{2kT} = 8 \frac{kT}{m} \ln \left(\frac{r}{r_c} \right) + 2GM_{\odot} \left(\frac{1}{r} - \frac{1}{r_c} \right) \quad (4.13)$$




FIG. 4.3. Radial-expansion speed $u(r)$ derived from isothermal coronal-expansion models with coronal temperatures ranging from $5 \times 10^5 \text{ K}$ to $4 \times 10^6 \text{ K}$ (Adapted from Parker, 1958.)




FIG. 4.4. Geometry of a flow (or flux) tube defined by radial streamlines (or magnetic-field lines).

magnetic flux within the tube (as implied by $\nabla \cdot \mathbf{B} = 0$) then would require that if B_0 were the (radial) field at the base of the flux tube, and $B(r)$ were the field at any radius r ,

$$B(r) \left(\frac{r}{R}\right)^2 dA_0 = B_0 dA_0$$

or

$$B(r) = B_0 \left(\frac{R}{r}\right)^2$$

That is, the intensity of the purely radial magnetic field would fall off as $1/r^2$.

$$\omega = \frac{2\pi \text{ rad}}{25.4 \text{ days}} = 2.7 \times 10^{-6} \text{ rad} \cdot \text{s}^{-1}$$




FIG. 4.5. Loci of a succession of fluid parcels (eight of them in this sketch) emitted at constant speed from a source fixed on the rotating sun.

Then the magnetic field stretched out along the path of plasma flowing from the fixed source in this coordinate system has components related by

$$\frac{B_\varphi}{B_r} = \frac{U_\varphi}{U_r} = \frac{-\omega r}{u(r)} \quad (4.14)$$

This gives a differential equation for the field lines near the solar equator (solar latitude = 0) as

$$\frac{r d\varphi}{dr} = \frac{-\omega r}{u(r)}$$

If the radial-expansion speed is constant, as in the solar wind well out in interplanetary space, this equation becomes

$$\frac{dr}{d\varphi} = -\frac{u}{\omega}$$

and has the obvious solution

$$r = -\frac{u}{\omega} \varphi + K''$$

Specification of the location of the source of a field line at longitude φ_0 at $r=R$ then yields

$$r - R = -\frac{u}{\omega}(\varphi - \varphi_0)$$

This is in fact a geometric figure of Grecian antiquity and respectability known as the spiral of Archimedes.

$$B_r(r) \frac{r^2}{R^2} dA_0 = B_0 dA_0$$

or

$$B_r(r) = B_0 \left(\frac{R^2}{r^2} \right)$$

The longitude (or azimuthal) component of the field can then be derived from (4.14):

$$B_\phi(r) = -\frac{\omega r}{u} B_r = -B_0 \frac{\omega R}{u} \frac{R}{r}$$




FIG. 4.6. Spiral
interplanetary magnetic-field
lines frozen into a radial solar-
wind expansion at $400 \text{ km} \cdot \text{s}^{-1}$.

Solar Terrestrial Environment







Photo Courtesy of Prof. Yohsuke Kamide

National Geophysical Data Center



低速太陽風




高速太陽風

The Sun


Solar Sectors

Solar sectors are a result of the solar wind carrying charged particles.



$$\tan\theta = r\omega/V_{sw}$$

Solar Minimum



Solar Maximum




Fig. 6.20. Waviness of the heliospheric current sheet during solar minimum and maximum conditions, based on a sketch by R. Jokipii, University of Arizona





Fig. 1.10. Three-dimensional sketch of the solar equatorial current sheet and associated magnetic field lines. The current sheet is shown as lying near the solar equator with spiraled, outward-pointing magnetic fields lying above it and inward-pointing fields lying below it. The current sheet contains folds or flutes. When the sun rotates, an observer near the ecliptic will alternately lie above and below the current and will see a changing sector pattern. The inset above shows a meridional cross section with the earth below the current sheet. (Figure courtesy of S.-I. Akasofu.)

Solar Storm

Solar energy	Effects on space environment
<u>Steady energy radiation</u>	<p>Control of the structure and dynamics of the atmosphere.</p> <p>Formation of the ionosphere, production of ionospheric currents.</p> <p>Formation of the magnetosphere, drives magnetospheric convection.</p>
<u>Transient energy radiation</u>	<p>Anomalous ionization of the ionosphere, radio noise.</p> <p>Magnetocospheric and ionospheric storms.</p> <p>Production of high-energy particles.</p> <p>Radiation, anomalous ionization in the polar ionosphere.</p>

- Solar index, F10.7: Solar radiation intensity of wavelength 10.7cm(2800MHz) and SSN (Sunspot number)



- The active Sun

- Flare


- A flare is a burst of “light” (easily observed at red H α 6563A) occurring in the chromosphere near a sunspot. Life time 3 min up to 2 hr.
- H α -flares are ranked in the size (importance, 1-4) and brilliance (f, n, b).
- X-ray flares are classified by power flux level of 1-8A (C, M, X).


- Radio burst

- Radio bursts are generally associated with flares and originate from all levels of the solar atmosphere.

- X-Ray flare classification

Class	Peak (W/m ²) between 1 and 8 Angstroms
B	$I < 10^{-6}$
C	$10^{-6} \leq I < 10^{-5}$
M	$10^{-5} \leq I < 10^{-4}$
X	$I \geq 10^{-4}$





Type	Time to earth	Consequence
Solar flare	8.5 minute	Sudden ionospheric disturbance Short wave (1-30MHz) fadeout
CME	1-2 day	Magnetic storm Ionospheric storm (1-2 day later)
High-energy particle	1-2 hour	Electronic equipment damage

	Type	Rate
Energy radiation from sun	Electromagnetic wave	3.8×10^{26} W/sec
	Solar wind	4.1×10^{20} W/sec
	Explosive event	7×10^{18} W/sec
Rate of mass-lose from sun	Electromagnetic wave	4.2×10^9 kg/sec
	Solar wind	1.4×10^9 kg/sec

TABLE 2. *Spectral density of power in solar radiation*


A	100	500	1000	1216	2000	3000	5000	10000
B	10	50	100	121.6	200	300	500	1000
C	10^{-5}	5×10^{-5}	3×10^{-5}	(Ly α)	10^{-2}	0.6	2	0.7
D	10^{-80}	10^{-12}	10^{-4}		0.7	1.5	2	0.5
E		3×10^{-3}		5×10^{-3}			1200	

Line A Wavelength in Å.

Line B Wavelength in nm.


Line C Observed spectral density ($\text{W m}^{-2} \text{ nm}^{-1}$).Line D Spectral density from black body at 6000°K normalized to equal observed density at 500 nm.Line E Observed total power flux in part of spectrum indicated (W m^{-2}).

Solar flare




2003/11/04 19:13

FIG. 3.3. Chromosphere
in H α , showing active regions
(bright) and filaments or
prominences (thin dark
ribbons on the disk). (Courtesy B.
Schmieder, Meudon
Observatory.)



Corona Mass Ejection (CME)

18 Aug 1980: White Light



Source: High Altitude Observatory/Solar Maximum Mission Archives

HAO A-013