

Assignment 5

https://github.com/jchryssanthacopoulos/quantum_information/tree/main/assignment_5

Quantum Information and Computing AA 2022–23

James Chryssanthacopoulos
6 December 2022



UNIVERSITÀ
DEGLI STUDI
DI PADOVA

- Goal is to solve 1D time-dependent quantum harmonic oscillator

$$\hat{H}|\Psi(x, t)\rangle = i\hbar \frac{\partial}{\partial t} |\Psi(x, t)\rangle, \quad \hat{H} \equiv \hat{T} + \hat{V} = \frac{\hat{p}^2}{2m} + \frac{1}{2}m\omega^2 \left(\hat{x} - \frac{t}{T}\right)^2, \quad t \in [0, T]$$

- Problem was solved using **split operator method**, where state is evolved using

$$|\Psi(x, t + \Delta t)\rangle = e^{-\frac{i\hat{V}\Delta t}{2}} \mathcal{F}^{-1} e^{-i\hat{T}\Delta t} \mathcal{F} e^{-\frac{i\hat{V}\Delta t}{2}} |\Psi(x, t)\rangle$$

where \mathcal{F} and \mathcal{F}^{-1} are the Fourier transform and its inverse, respectively
($\hbar = \omega = m = 1$ was used)

```
! multiply by potential part of Hamiltonian
do ii = 1, Nx
  V(ii) = potential(x_grid(ii), time, tmax)
  final_state(ii) = cexp(cmplx(0.0, -0.5 * V(ii) * dt)) * init_state(ii)
end do



! normalize
call normalize(final_state, dx)

! call FFT to go from coordinate space to momentum space
call dfftw_plan_dft_1d(plan, Nx, final_state, state_transform, -1, 64)
call dfftw_execute_dft(plan, final_state, state_transform)
call dfftw_destroy_plan(plan)
```

```
$ compiled/solve_time_dep_ho --xmin -15 --xmax 15 --tmax 10 --num_x_pts 100 --num_t_pts 100
xmin = -15.000
xmax = 15.000
tmax = 10.000
num_x_pts = 100
num_t_pts = 100
debug = F
output_filename = solution.txt
```

- `fftw` library installed from source and used to compute FFT
- System simulated using discretization parameters and wavefunction saved to file, results analyzed in Jupyter notebook


- $|\Psi(x, t)\rangle$ starts in ground state, $(\frac{1}{\pi})^{1/4} e^{-\frac{x^2}{2}}$
- Position and time discretized using N_x and N_t equispaced bins of size Δx and Δt
- Discretized momentum is $\left[0, \frac{2\pi}{L}, \dots, \frac{\pi(N_x-2)}{L}, -\frac{\pi N_x}{L}, \dots, -\frac{2\pi}{L}\right]$, where L is x range
- Wavefunction moves to right, developing complex phase, as expected




Wavefunction for Different T



- Expected position $E[x](t)$ computed using $\sum_{i=1}^{N_x} x_i |\Psi(x_i, t)|^2 \Delta x$
- Result agrees well with theoretical result, $E[x](t) = \frac{t}{T} - \sin(\frac{t}{T})$
- Deviations from linear fit decrease with increasing T



- Position oscillations decrease when T increases because when T is large, potential changes slowly (i.e., **adiabatic approximation**)
- Fit of oscillations to $\alpha e^{\beta x}$ results in $\alpha = 0.60 \pm 0.03$ and $\beta = -0.88 \pm 0.01$



- **Correctness.** Results closely match theoretical result
- **Stability.** Code is numerically stable, but results sensitive to discretization
- **Accurate Discretization.** Error of split operator method is $\mathcal{O}(\Delta t^3)$. For $\Delta t \sim 1$, expected final state not reproduced
- **Flexibility.** Split operator method assumes \hat{V} is only function of \hat{x} and \hat{T} is only function of \hat{p} , enabling exact diagonal representation
- **Efficiently.** Overall time complexity is $\mathcal{O}(N_t N_x \log N_x)$. It took ~ 8 s to solve using $N_x = N_t = 10^4$