

Newton-Cotes

Math 131: Numerical Analysis

J.C. Meza

April 11, 2024

- The basic quadrature rules derived so far are generally good, but what if we wanted to have formulas with greater accuracy.
- The general approach we used still holds and leads to a family of quadrature formulas known as ***Newton-Cotes*** formulas.
- These are classified under either open or closed depending on whether the formulas include the end points or not.
 - ▶ ***Closed Newton-Cotes*** include the endpoints of closed interval $[a, b]$ as nodes.
 - ▶ ***Open Newton-Cotes*** do not include the endpoints.

In particular

To be specific, for a closed Newton-Cotes quadrature formula we would choose the node points x_i through the formula:

$$x_i = a + i \frac{b-a}{n-1}, \quad i = 0, 1, \dots, n-1. \quad (1)$$

For an open Newton-Cotes quadrature formula we would use the formula:

$$x_i = a + (i+1) \frac{b-a}{n+1}, \quad i = 0, 1, \dots, n-1. \quad (2)$$

Example

Suppose, we choose $n = 5$ on the interval $[a,b] = [0,1]$.

Then the closed Newton-Cotes formula would generate the points:

$$\begin{aligned}x_i &= a + i \cdot \frac{b-a}{n-1}, \\&= 0 + i \frac{1}{4}, \\&= \frac{i}{4}, \quad i = 0, 1, \dots, 4,\end{aligned}$$

thereby yielding the set of nodes: $\{x\} = \{ 0, .25, .5, .75, 1.0 \}$.

Example

Similarly for the open Newton-Cotes formula would generate the points:

$$\begin{aligned}x_i &= a + (i + 1) \cdot \frac{b - a}{n + 1}, \\&= 0 + (i + 1) \frac{1}{6}, \\&= \frac{i + 1}{6}, \quad i = 0, 1, \dots, 4,\end{aligned}$$

which generates the set of nodes:

$$\{x\} = \{ 1/6, 2/6, 3/6, 4/6, 5/6 \}.$$

Midpoint Rule

One example of an Open Newton-Cotes is the midpoint rule

$$\int_a^b f(x)dx = 2hf(x_0) + \frac{h^3}{3}f''(\xi) \quad \xi \in (a, b)$$

where x_0 is the midpoint between a and b .

Likewise, both Trapezoidal and Simpson's rules, which we introduced earlier can be categorized as Closed Newton-Cotes.

Other formulas

- There are many different formulas of both the Closed and Open variety all with corresponding error terms.
- All of them can be derived by the methods we've used for Trapezoid and Simpson's rule, so there is little to be gained by re-deriving them.
- Instead we will present them here because an interesting pattern arises that is worth knowing about:

Closed Newton-Cotes formulas:

$n = 2$ (Trapezoid)

$$I(f) = \frac{b-a}{2}[f(x_0) + f(x_1)] \quad (3)$$

$n = 3$ (Simpson's)

$$I(f) = \frac{b-a}{6}[f(x_0) + 4f(x_1) + f(x_2)] \quad (4)$$

$n = 4$ (Simpson's 3/8)

$$I(f) = \frac{b-a}{8}[f(x_0) + 3f(x_1) + 3f(x_2) + f(x_3)]$$

$n = 5$ (Boole's rule)

$$I(f) = \frac{b-a}{90}[7f(x_0) + 32f(x_1) + 12f(x_2) + 32f(x_3) + 7f(x_4)]$$

Notation

- The formulas here are written using $b - a$ versus h to make them easier to compare.
- However, you will see these formulas written in terms of h in many other places.
- You should be careful in understanding exactly what h represents as it often is taken to mean $h = (b - a)/(n - 1)$, $n \geq 1$, which corresponds to the number of node points used in the quadrature formula.

Higher-order formulas

- In theory, we could go as high as we wanted (and people have) in generating higher-order quadrature formulas, and of course with additional computational work.
- However, for large n the formulas can be shown to become numerically unstable ($n \geq 11$.) One can actually prove that formulas do not converge for all integrands that are analytic.
- In practice, we tend to only use low-order formulas since they can still give us good accuracy (especially over small intervals (see Exercise 1.1 below)).

Open Newton-Cotes formulas:

$n = 1$ (Midpoint)

$$I(f) = (b - a)f(x_0)$$

$n = 2$

$$I(f) = \frac{b - a}{2}[f(x_0) + f(x_1)]$$

$n = 3$

$$I(f) = \frac{b - a}{3}[2f(x_0) - f(x_1) + 2f(x_2)]$$

Similarly to the closed Newton-Cotes formulas, we could continue and derive higher-order formulas - with the same consequences.

Section 1

Error Estimates

Error Estimates

In both the closed and open Newton-Cotes cases, the formulas have error terms, which we have summarized in the table below, along with the precision of each:

Table 1: Summary of Error Terms for Newton-Cotes quadrature formulas

Name	npts	Error	Precision
Trapezoid	2	$-\frac{(b-a)^3}{12} f^{(2)}(\xi)$	1
Simpson's	3	$-\frac{(b-a)^5}{2880} f^{(4)}(\xi)$	3
Simpson's 3/8	4	$-\frac{(b-a)^5}{6480} f^{(4)}(\xi)$	3
Boole	5	$-\frac{(b-a)^7}{1935360} f^{(6)}(\xi)$	5
Midpoint	1	$\frac{(b-a)^3}{24} f^{(2)}(\xi)$	1
	2	$\frac{(b-a)^3}{36} f^{(2)}(\xi)$	1
	3	$\frac{(b-a)^5}{23040} f^{(4)}(\xi)$	3

Important

An interesting feature of the quadrature formulas is that whenever N is odd then the precision of the formula $= N$. But when N is even then the precision is only $N - 1$. We lose one order in the precision whenever N is even! Or we could also say that we gain one order of precision for N odd.

As a final look into these methods, let's compare several of the methods on a simple function to gain further insight into the behavior of the formulas.

Exercise

Exercise

Compute the value of

$$\int_0^1 e^x dx$$

using the Trapezoid and Simpson's Rule for:

- 1 $a = 0, b = 1$
- 2 $a = 0.9, b = 1$

Trapezoid Rule

Trapezoid Rule:

$$\int_a^b f(x)dx = \frac{(b-a)}{2} [f(x_0) + f(x_1)],$$

Simpson's Rule:

$$\int_a^b f(x)dx = \frac{(b-a)}{6}[f(x_0) + 4f(x_1) + f(x_2)]$$

Remarks

The lesson from this example is that all of the formulas have a rather large error when we compute the integral over a large interval, whereas when we considered a smaller interval, the error was in fact quite small.

Section 2

Summary

Summary

Let's take a step back and summarize the main results:

- We can use a simple approach towards deriving basic quadrature rules by replacing the integrand with an interpolating polynomial and integrating the polynomial on a chosen set of N points.
- Using Taylor's theorem, we can also generate corresponding error terms that can provide us with estimates on how well the quadrature formula approximated the given integral.

Summary (cont.)

- The precision of a quadrature formula is the highest degree of the polynomial for which the formula is exact. When N is odd, the precision is also N ; but when N is even, the precision is $N - 1$.
- Higher-order quadrature formulas yield greater accuracy, but at greater additional computational work as well as a fundamental assumption on the higher-order derivatives being nicely behaved (i.e. bounded).
- Basic (and low-order) formulas can be quite accurate, but usually require a small interval. This observation will prove useful in the next sections.