

NCT07561983: Tislelizumab Plus Chemotherapy and BACE for Unresectable NSCLC

Gender: ALL

Age: 18 Years - 80 Years

Criteria:

Inclusion Criteria:

- * Age 18 to 80 years
- * Histologically or cytologically confirmed non-small cell lung cancer (NSCLC)
- * Newly diagnosed, previously untreated stage IIIA-IIIB NSCLC according to the 9th edition TNM staging system
- * Initially unresectable disease as determined by multidisciplinary team (MDT) assessment
- * At least 1 measurable intrapulmonary lesion according to RECIST version 1.1
- * Eastern Cooperative Oncology Group (ECOG) performance status of 0 or 1
- * Forced expiratory volume in the first second (FEV1) $\geq$ 1.0 L and $\geq$ 40% of predicted normal value
- * Estimated life expectancy of at least 3 months
- * Adequate organ function
- * Willingness to provide tumor tissue for pathology, molecular testing, and PD-L1 assessment before enrollment
- * Women of childbearing potential must have a negative pregnancy test within 72 hours before the first dose and agree to use effective contraception during the study and for 3 months after the last dose of tislelizumab
- * Men with partners of childbearing potential must agree to use effective contraception during the study and for 3 months after the last dose of tislelizumab
- * Ability to understand and willingness to sign a written informed consent form

Exclusion Criteria:

- * Prior local therapy for NSCLC, including radiotherapy or interventional therapy
- * Known positive driver genomic alterations, including EGFR mutations, ALK rearrangements, ROS1 rearrangements, and MET exon 14 skipping alterations
- * Distant organ metastasis
- * History of another malignancy within the past 5 years
- * Active autoimmune disease or history of autoimmune disease requiring systemic treatment
- * Known allergy to any study drug or excipient
- * Interstitial lung disease, non-infectious pneumonitis, chronic obstructive pulmonary disease, or other uncontrolled systemic diseases judged to interfere with study treatment
- * Severe chronic or active infection requiring systemic antibacterial, antifungal, or antiviral therapy, including active tuberculosis
- * Major surgery requiring general anesthesia within 4 weeks before first dose
- * Any medical condition, alcohol or drug abuse, or dependence that may interfere with study treatment, interpretation of results, or increase treatment risk
- * Participation in another interventional therapeutic clinical study
- * Psychiatric illness or history of psychotropic drug abuse that may compromise study participation
- * Any condition judged by the investigator to make the patient unsuitable for the study