

NCT07562932: Assessing Catecholamine Treatment Initiation Options: Norepinephrine vs Dopamine for Cardiogenic Shock

Gender: ALL

Age: 19 Years - MISSING

Criteria:

Inclusion Criteria: 1\&2

1. Age ≥ 19
2. Cardiogenic shocka in the stage of Society for cardiovascular angiography and intervention (SCAI) C or D

* Cardiogenic shock was defined as follows, and should fulfill both 1) and 2)

1. systolic blood pressure ≤ 90 mm Hg for ≥ 30 min or need of inotropes or vasopressors to maintain systolic blood pressure ≤ 90 mm Hg And

2. Impaired cardiac function confirmed by cardiac catheterization or echocardiography

* Patients will be further classified based on the SCAI shock classification system as follows:

1. SCAI B: no signs of hypoperfusion

2. SCAI C: any signs of hypoperfusion, cardiogenic shock will be classified as SCAI C, and these include mental status change, cool and clammy skin, mottled skin appearance, decreased urine output (30ml/hour) or lactate over 2.0mmol/L

3. SCAI D: requirement of second-line vasoactive drug or mechanical circulatory support based on the predefined treatment protocol

Exclusion Criteria: any of these,

1. Administration of vasoactive drug more than 6 hours before enrollment
2. Patients already on temporary mechanical circulatory supportb before enrollment
3. Glasgow Coma Scale lower than 8 or other evidence of irreversible brain injury
4. Shock etiologies other than cardiogenic which include postcardiotomy shock or mixed shock
5. Pregnancy or lactation