

NCT07563075: Effect of High-Flow Nasal Cannula Oxygen Therapy on Hypoxemia in Pediatric

Sedated Bronchoscopy

Gender: ALL

Age: 1 Year - 6 Years

Criteria:

Inclusion Criteria:

- * 1 year $\leq$ age $\leq$ 6 years,
- * Weight $\geq$ 10 kg,
- * Undergoing elective sedated bronchoscopy,
- * Expected procedure duration $\leq$ 45 minutes,
- * Written informed consent obtained from the subject's legal guardian.

Exclusion Criteria:

- * preoperative SpO₂ $<$ 95% on room air,
- * Already sedated and tracheally intubated,
- * Known history of pneumothorax, known congenital or acquired upper airway abnormalities (e.g., nasopharyngeal structural anomalies), or history of difficult airway,
- * Coagulation disorders or predisposition to oral/nasal bleeding, mucosal injury, or space-occupying lesions,
- * Severe cardiac insufficiency ($<$ 4 METs), severe renal insufficiency (requiring dialysis), diagnosed severe hepatic insufficiency, Increased intracranial pressure, or ASA classification $\geq$ IV,
- * Allergy to propofol or sufentanil,
- * Multiple traumatic injuries,
- * Current participation in another clinical trial,
- * Other conditions deemed unsuitable by the investigator.