

NCT07563088: SPARE-1st: Surgery Alone vs Surgery Plus Postoperative Radiotherapy for Sinonasal Adenoid Cystic Carcinoma

Gender: ALL

Age: 18 Years - 75 Years

Criteria:

Inclusion Criteria:

- * Age ≥ 18 years and ≤ 75 years.
- * Histologically confirmed adenoid cystic carcinoma of cribriform or tubular type (solid type excluded) by central pathology review.
- * Completed radical surgical resection with postoperative pathology confirming R0 resection (negative microscopic margins).
- * Tumor stage T1, T2, or T3 according to AJCC 9th edition, with N0 and M0 status.
- * Tumor originating from the nasal cavity or paranasal sinuses (excluding primary salivary gland tumors metastatic to this region).
- * ECOG performance status 0 or 1.
- * Adequate bone marrow, liver, and kidney function within 14 days before enrollment:
 - * Absolute neutrophil count $\geq 1.5 \times 10^9/L$, platelet count $\geq 100 \times 10^9/L$, hemoglobin ≥ 90 g/L.
 - * Total bilirubin ≤ 1.5 upper limit of normal (ULN); AST and ALT ≤ 2.5 ULN.
 - * Serum creatinine ≤ 1.5 ULN or calculated creatinine clearance ≥ 60 mL/min (Cockcroft-Gault formula).
- * Willing and able to provide written informed consent.

Exclusion Criteria:

- * Tumor stage T4, or presence of regional lymph node metastasis (N+), or distant metastasis (M1).
- * Postoperative pathology showing positive margins (R1 or R2).
- * Prior head and neck radiotherapy.
- * Concurrent active malignancy other than adequately treated basal cell carcinoma of the skin or carcinoma in situ of the cervix.
- * Uncontrolled serious concomitant disease (e.g., unstable angina, recent myocardial infarction, uncontrolled hypertension).
- * Pregnant or breastfeeding women, or women of childbearing age not using effective contraception.
- * Psychological, social, or geographic factors that may interfere with study compliance.