

NCT07563179: Bugitinib Combined With Venetoclax and Cytarabine in Relapsed/Refractory

Acute Myeloid Leukemia

Gender: ALL

Age: 18 Years - 70 Years

Criteria:

Inclusion Criteria:

Participants must meet all of the following criteria:

- * Age 18-70 years (inclusive), regardless of sex;
- * Diagnosis of non-M3 acute myeloid leukemia (AML), and meeting at least one of the following conditions:
 - * Failure to achieve complete remission (CR) after standard induction chemotherapy;
 - * First CR duration ≤ 12 months;
 - * First CR duration > 12 months, followed by relapse and failure of at least one line of standard chemotherapy;
 - * Relapsed disease after ≥ 2 prior lines of therapy;
- * Eastern Cooperative Oncology Group (ECOG) performance status of 0-2;
- * Estimated life expectancy ≥ 3 months;
- * Adequate organ function, defined as:
 - * Cardiac: left ventricular ejection fraction (LVEF) $\geq 50\%$ by echocardiography, with no clinically significant ECG abnormalities;
 - * Renal: serum creatinine ≤ 2.0 upper limit of normal (ULN) and creatinine clearance ≥ 60 mL/min (Cockcroft-Gault);
 - * Hepatic: ALT and AST ≤ 3.0 ULN;
 - * Total bilirubin ≤ 2.0 ULN (≤ 3.0 ULN for patients with Gilbert syndrome);
 - * Oxygen saturation $\geq 92\%$ on room air;
- * Ability to take oral medications and comply with study procedures;
- * Willingness to participate and provide written informed consent.

Exclusion Criteria:

Participants meeting any of the following criteria will be excluded:

- * Known hypersensitivity to any study drug or its excipients;
- * Active central nervous system (CNS) leukemia (patients with prior CNS involvement who achieved complete remission and are clinically stable may be eligible);
- * Active, uncontrolled bacterial, viral, or systemic fungal infection;
- * Known hereditary bleeding or coagulation disorders, history of non-traumatic bleeding or thromboembolism, or other conditions that may increase bleeding risk;
- * Evidence of active infection, including:
 - * Hepatitis B virus (HBV) infection with positive HBsAg or HBcAb and HBV DNA > 100 copies/mL;
 - * Hepatitis C virus (HCV) antibody positive with detectable HCV RNA;
 - * Human immunodeficiency virus (HIV) infection;
 - * Positive test for syphilis;
- * Active bleeding or clinically significant bleeding tendency, including but not limited to:
 - * Active gastrointestinal bleeding, intracranial hemorrhage, or bleeding in other vital organs;
 - * Grade ≥ 3 bleeding event within 4 weeks prior to enrollment (per CTCAE);

- * Known hereditary bleeding disorders (e.g., hemophilia);
- * Severe platelet dysfunction not correctable;
- * Significant cardiovascular disease, including but not limited to:
 - * New York Heart Association (NYHA) class III-IV heart failure;
 - * Myocardial infarction or coronary artery bypass grafting (CABG) within 6 months prior to enrollment;
 - * Clinically significant ventricular arrhythmias or unexplained syncope (excluding vasovagal or dehydration-related causes);
 - * Severe cardiomyopathy;
- * History of or concurrent other malignancies (except for adequately treated basal cell carcinoma of the skin or carcinoma in situ of the cervix);
- * Clinically significant central nervous system disorders (e.g., epilepsy, stroke, psychiatric disorders) that may affect compliance or safety;
- * Participation in another interventional clinical trial within 30 days prior to enrollment;
- * Pregnant or breastfeeding women, or subjects planning pregnancy during the study period or within 6 months after the last dose and unwilling to use effective contraception;
- * Known or suspected drug abuse or alcohol dependence;
- * Requirement for concomitant use of strong CYP3A inhibitors or inducers that cannot be discontinued or dose-adjusted per protocol;
- * Any other condition that, in the investigator's opinion, would make the subject unsuitable for participation.