

Conceptual Test

Chapter 24

ASTRONOMY TODAY

6th edition

Chaisson

McMillan

© 2008 Pearson Addison-Wesley

This work is protected by United States copyright laws and is provided solely for the use of instructors in teaching their courses and assessing student learning. Dissemination or sale of any part of this work (including on the World Wide Web) will destroy the integrity of the work and is not permitted. The work and materials from it should never be made available to students except by instructors using the accompanying text in their classes. All recipients of this work are expected to abide by these restrictions and to honor the intended pedagogical purposes and the needs of other instructors who rely on these materials.

Question 1

Which of the following is NOT one of the basic types of galaxy?

- 1) spiral**
- 2) lenticular**
- 3) globular**
- 4) irregular**
- 5) elliptical**

Question 1

Which of the following is NOT one of the basic types of galaxy?

- 1) spiral
- 2) lenticular
- 3) globular
- 4) irregular
- 5) elliptical



Question 2

Which type of galaxy contains the most population II stars?

- 1) spiral**
- 2) globular**
- 3) irregular**
- 4) elliptical**

Question 2

Which type of galaxy contains the most population II stars?

- 1) spiral
- 2) globular
- 3) irregular
- 4) elliptical




(a) M49 Type E2



(b) M84 Type E3



(c) M110 Type E5



Question 3

Which type of galaxy most resembles our Milky Way?

- 1) spiral**
- 2) globular**
- 3) irregular**
- 4) elliptical**
- 5) lenticular**

Question 3

Which type of galaxy most resembles our Milky Way?

1) spiral

2) globular

3) irregular

4) elliptical

5) lenticular



Question 4

Which type of galaxy can grow to the greatest mass and possess the largest number of globular clusters?

- 1) spiral**
- 2) globular**
- 3) irregular**
- 4) elliptical**
- 5) lenticular**

Question 4

Which type of galaxy can grow to the greatest mass and possess the largest number of globular clusters?

- 1) spiral
- 2) globular
- 3) irregular
- 4) elliptical
- 5) lenticular




(a) M49 Type E2



(b) M84 Type E3



(c) M110 Type E5



Question 5

Hubble took spectra of galaxies in the 1930s. What did he find?


- 1) All the galaxies showed blue line shifts.**
- 2) Most galaxies showed redshifts.**
- 3) Galaxies showed about half redshifts and half blueshifts.**
- 4) Galaxies showed no line shifts at all.**
- 5) Some galaxies showed a redshift that changed into a blueshift at other times.**



Question 5

Hubble took spectra of galaxies in the 1930s. What did he find?


- 1) All the galaxies showed blue line shifts.
- 2) Most galaxies showed redshifts.
- 3) Galaxies showed about half redshifts and half blueshifts.
- 4) Galaxies showed no line shifts at all.
- 5) Some galaxies showed a redshift that changed into a blueshift at other times.



Question 6

What did the discovery suggest to cosmologists?


- 1) The universe is static.
- 2) The universe is collapsing.
- 3) The universe is expanding.
- 4) The universe is contracting.
- 5) There is no accepted interpretation.



Question 6

What did the discovery suggest to cosmologists?

- 1) The universe is static.
- 2) The universe is collapsing.
- 3) The universe is expanding.
- 4) The universe is contracting.
- 5) There is no accepted interpretation.



Question 7


**Most of the
galaxies (99%) are**

- 1) moving around randomly.**
- 2) showing Doppler blueshifts.**
- 3) moving in circles.**
- 4) rushing away from us.**
- 5) approaching us.**


Question 7

Most of the galaxies (99%) are

- 1) moving around randomly.
- 2) showing Doppler blueshifts.
- 3) moving in circles.
- 4) rushing away from us.**
- 5) approaching us.



(a)




(b)


Question 8

Galaxy distance and Doppler redshift are correlated for galaxies. What do we call that correlation?

- 1) the Hubble “law”
- 2) Wein’s relation
- 3) redshift–distance relation
- 4) period–luminosity relation
- 5) both 1 and 3



(a)



(b)

Question 8

Galaxy distance and Doppler redshift are correlated for galaxies. What do we call that correlation?


1) the Hubble “law”

2) Wein’s relation


3) redshift–distance relation

4) period–luminosity relation

5) both 1 and 3



(a)





(b)

Question 9

What is the H in $V = Hr$?

- 1) Cepheid relation
- 2) Hubble constant
- 3) radial velocity
- 4) distance in megaparsecs
- 5) Wien's constant



Question 9

What is the H in $V = Hr$?



1) Cepheid relation

2) Hubble constant

3) radial velocity

4) distance in megaparsecs


5) Wien's constant



Question 10

We estimate the value of H by getting what measure for many galaxies at many different (Doppler redshift) recession velocities?


- 1) temperature
- 2) luminosity
- 3) distance
- 4) color
- 5) size



Question 10

We estimate the value of H by getting what measure for many galaxies at many different (Doppler redshift) recession velocities?


- 1) temperature
- 2) luminosity
- 3) distance
- 4) color
- 5) size



Question 11

_____ galaxies may bridge the gap between spiral galaxies and quasars.


- 1) Hickson
- 2) Arp
- 3) Core–halo
- 4) Seyfert
- 5) Hubble



Question 11

_____ galaxies may bridge the gap between spiral galaxies and quasars.

- 1) Hickson
- 2) Arp
- 3) Core-halo
- 4) Seyfert
- 5) Hubble



Question 12


Not only does the central engine of active galaxies and quasars require a black hole, but _____ is also needed to provide the energy radiated.

- 1) a collision with another galaxy**
- 2) an accretion disk of matter**
- 3) a very strong magnetic field**
- 4) a source of very high energy electrons**
- 5) a high rate of rotation of the black hole**

Question 12

Not only does the central engine of active galaxies and quasars require a black hole, but _____ is also needed to provide the energy radiated.

- 1) a collision with another galaxy
- 2) an accretion disk of matter
- 3) a very strong magnetic field
- 4) a source of very high energy electrons
- 5) a high rate of rotation of the black hole



Question 13

**How would we get
Doppler velocity
independent distances to
the nearest galaxies (a
few megaparsecs)?**

- 1) trig parallax**
- 2) period–luminosity relation for Cepheids**
- 3) Hubble's law ($V = Hr$)**
- 4) radar reflection**
- 5) main sequence fitting**

Question 13

How would we get
Doppler velocity
independent distances to
the nearest galaxies (a
few megaparsecs)?

1) trig parallax

2) period–luminosity relation for
Cepheids

3) Hubble's law ($V = Hr$)

4) radar reflection

5) main sequence fitting

Question 14

How would we get Doppler velocity independent distances to more distant galaxies (tens of megaparsecs) without using the Hubble law?

- 1) trig parallax**
- 2) period–luminosity relation for Cepheids**
- 3) Comparing brightest stars/clusters/nebulae with similar objects in MW.**
- 4) main sequence fitting**
- 5) impossible to do it**

Question 14

How would we get Doppler velocity independent distances to more distant galaxies (tens of megaparsecs) without using the Hubble law?

- 1) trig parallax
- 2) period–luminosity relation for Cepheids
- 3) Comparing brightest stars/clusters/nebulae with similar objects in MW.
- 4) main sequence fitting
- 5) impossible to do it

Question 15

You see a distant car in the night. What two things must you know in order to estimate its distance?

- 1) how fast it is moving towards or away from me (Doppler shift)**
- 2) the apparent brightness (flux) of the distant headlights**
- 3) the intrinsic brightness of the headlights (e.g., 500 or 1000 watts)**
- 4) 1 and 2**
- 5) 2 and 3**

Question 15

You see a distant car in the night. What two things must you know in order to estimate its distance?

- 1) how fast it is moving towards or away from me (Doppler shift)
- 2) the apparent brightness (flux) of the distant headlights
- 3) the intrinsic brightness of the headlights (e.g., 500 or 1000 watts)

4) 1 and 2

5) 2 and 3

Question 16

You see a distant (main-sequence!) B-type star like Rigel in a rather distant galaxy. What two things do you need to know in order to estimate its distance?

- 1) how fast it is moving towards or away from me**
- 2) the apparent brightness (flux or apparent magnitude)**
- 3) the intrinsic brightness of main-sequence B stars like Rigel**
- 4) 1 and 2**
- 5) 2 and 3**

Question 16

You see a distant (main-sequence!) B-type star like Rigel in a rather distant galaxy. What two things do you need to know in order to estimate its distance?

- 1) how fast it is moving towards or away from me
- 2) the apparent brightness (flux or apparent magnitude)
- 3) the intrinsic brightness of main-sequence B stars like Rigel
- 4) 1 and 2
- 5) 2 and 3

Question 17

What do you need to know in order to estimate the distance of galaxies where we can no longer see individual stars / clusters / nebulae?

- 1) trig parallax**
- 2) period–luminosity relation for Cepheids**
- 3) main sequence fitting**
- 4) Supernovae give us a reasonable guess.**
- 5) We cannot do it without $V = Hr$.**

Question 17


What do you need to know in order to estimate the distance of galaxies where we can no longer see individual stars / clusters / nebulae?

- 1) trig parallax
- 2) period–luminosity relation for Cepheids
- 3) main sequence fitting
- 4) Supernovae give us a reasonable guess.**
- 5) We cannot do it without $V = Hr$.

Question 18

The spectral lines of quasars are


- 1) emission lines with large redshifts.
- 2) a continuum from the synchrotron radiation.
- 3) fuzzy absorption lines from the merged light of the billions of stars.
- 4) too complex for any interpretation.
- 5) nonexistent, the gas so hot as to be totally ionized, so no lines are seen.



Question 18

The spectral lines of quasars are

- 1) emission lines with large redshifts.
- 2) a continuum from the synchrotron radiation.
- 3) fuzzy absorption lines from the merged light of the billions of stars.
- 4) too complex for any interpretation.
- 5) nonexistent, the gas so hot as to be totally ionized, so no lines are seen.



Question 19

What is the most likely source of energy for active galaxies and quasars?

- 1) collisions of large spiral galaxies**
- 2) large clusters of very massive, luminous stars**
- 3) a single supermassive, superluminous star**
- 4) accretion onto a supermassive black hole**
- 5) numerous supernovae from rapid star formation in young galaxies**

Question 19

What is the most likely source of energy for active galaxies and quasars?

- 1) collisions of large spiral galaxies
- 2) large clusters of very massive, luminous stars
- 3) a single supermassive, superluminous star
- 4) accretion onto a supermassive black hole**
- 5) numerous supernovae from rapid star formation in young galaxies

