


Chapter 7

Energy of a System



Outline for Week 7,D1

Exam I return

Work by a constant force $W = F \Delta r \cos \theta$

Scalar dot products, $\mathbf{A} \cdot \mathbf{B} = C$

Homework

Ch. 7 Read 7.1-7.4

P. 1,2,5,8,12,18,20,22,25,37,39,41,45,55,
56,59,60,65

MiscQ: 1-13 (odd)

Do for next Mon

Notes:

“NEW STUFF”: added chapter7.pdf, two practice quizzes, and Ch 7 hwk key.

Outline for Week 7,D2

What is energy?

Work by a constant force $W = F \Delta r \cos \theta$

Scalar dot products, $\mathbf{A} \cdot \mathbf{B} = C$

Work by a variable force $W = \int \vec{F} \cdot d\vec{r}$

Homework

Ch. 7 Read 7.1-7.4

P. 1,2,5,8,12,18,20,22,25,37,39,41,45,55,
56,59,60,65

MiscQ: 1-13 (odd)

Do for next Mon (after break)

Notes:

“NEW STUFF”: updated chapter7.pdf.

Midterms will be based on Exam 1.

What is Energy?

Some problems are very difficult to solve with Newton's Laws alone, but the *Law of Conservation of Energy* makes them easier.

Energy: in physics, a quantity measured in Joules (SI), ergs, calories, BTU, N-m, ft-lbs, or kW-hrs. It's dimensions are ML^2/T^2 . It is a scalar, NOT a vector.

Energy: “the ability to do work” (Energy is not easily defined!)

Types of energy:

Energies stored in a system: thermal (or internal), kinetic (energy of motion), potential (energy of position: gravitational, elastic, nuclear, chemical)

Energies transferred into a system: **work** (W) and heat (Q). (Others: T_{ET} T_{MT} T_{ER})

It is wrong to say “the system contains a lot of work”. Instead you can say “work was done on the system”. You can say “the system contains a lot of kinetic and potential energy” or “work increased the kinetic energy of the system”.

Work

The **work**, W , done on a system by a *constant force* is given by:

$$W = F \Delta r \cos \theta$$


where F is the magnitude of the force, Δr is the magnitude of the displacement of the point of application of the force, and θ is the angle between the force and the displacement vectors.

- The meaning of the term *work* is distinctly different in physics than in everyday meaning.
- Work is not a vector, even though it involves two vectors, \mathbf{F} and $\Delta \mathbf{r}$.
- Work can be + or – because $\cos \theta$ can be + or -.
- Two geometric interpretations:
 - $F \cos \theta$ is the component of F parallel to $\Delta \mathbf{r}$, (i.e., $W = F_{\parallel} \Delta r$)
 - $\Delta r \cos \theta$ is the component of $\Delta \mathbf{r}$ parallel to F . (i.e., $W = F \Delta r_{\parallel}$)

Work, cont.


$$W = F \Delta r \cos \theta$$

- The displacement, $\Delta \mathbf{r}$, is that of the point of application of the force.
- A force does no work on the object if the force does not move through a displacement.
- The work done by a force on a moving object is zero when the force applied is perpendicular to the displacement of its point of application.



Force without motion

When a force is exerted on an object which does not move, no work is done on the object.



Force perpendicular to the motion
constant velocity

When an object is carried at constant velocity by a force which acts at right angles to the motion, no work is done on the object.

Work, cont.

$$W = F \Delta r \cos \theta$$

Q1: What work is done by $F = 10 \text{ N}$ if $\theta = 0$ degrees and $\Delta r = 2 \text{ m}$?


Q2: What work is done by $F = 10 \text{ N}$ if $\theta = 60$ degrees and $\Delta r = 2 \text{ m}$?

Q3: What work is done by $F = 10 \text{ N}$ if $\theta = 90$ degrees and $\Delta r = 2 \text{ m}$?

Q4: The box weighs 20 N . What is the work done by the normal force on the box when $\theta = 0$ degrees and $\Delta r = 2 \text{ m}$?

Q5: The box weighs 20 N . What is the work done by the force of gravity on the box when $\theta = 0$ degrees and $\Delta r = 2 \text{ m}$?

Q6: What is the work done by the force of friction on the 20 N box if $\mu_k = 0.5$ and when $\theta = 0$ degrees and $\Delta r = 2 \text{ m}$?



Work, cont.


$$W = F \Delta r \cos \theta$$

Q7: What work is done by $\mathbf{F} = 20 \text{ N } \hat{i}$ if $\Delta \mathbf{r} = 2 \text{ m } \hat{i}$?

Q8: What work is done by $\mathbf{F} = -20 \text{ N } \hat{i}$ if $\Delta \mathbf{r} = 2 \text{ m } \hat{i}$? How could this happen?

Q9: What work is done by $\mathbf{F} = 20 \hat{i} + 15 \hat{j} \text{ N}$ if $\Delta \mathbf{r} = 3 \text{ m } \hat{i}$?

Q10: What work is done by $\mathbf{F} = 20 \hat{i} + 0 \hat{j} \text{ N}$ if $\Delta \mathbf{r} = 3 \text{ m } \hat{i} - 2 \text{ m } \hat{j}$?



© Cengage Learning. All Rights Reserved.


Work, cont.

What is the work done by the tension force in 1 revolution of the mass?

What is the work by the tension force in $\frac{1}{2}$ revolution of the mass?

What is the work done by the centripetal force for any angle swept out by the string?

What is the work done by the net force on the mass for any angle swept out by the string?



Displacement in the Work Equation

The displacement is that of the point of application of the force.

If the force is applied to a rigid object that can be modeled as a particle, the displacement is the same as that of the particle.

For a deformable system, the displacement of the object generally is not the same as the displacement associated with the forces applied.

Work by friction is also hard to explain because there is no single point of application. But it can ultimately be modeled by $W = f_k \Delta r \cos \theta$ and is usually negative because $\theta=180$.

Outline for Week 7,D3

Scalar dot products, $\mathbf{A} \cdot \mathbf{B} = C$

Work by a variable force $W = \int \vec{F} \cdot d\vec{r}$

Homework

Ch. 7 Read 7.1-7.4

P. 1,2,5,8,12,18,20,22,25,37,39,41,45,55,
56,59,60,65

MiscQ: 1-13 (odd)

Do for next Mon (after break)

Notes:

“NEW STUFF”: added exam-like questions for Ch.7,
updated chapter7.pdf.

Midterms will be based on Exam 1.

Work (cont.)

The normal force and the gravitational force do no work on the object.

- $\cos \theta = \cos 90^\circ = 0$


The force \mathbf{F} is the only force that does work on the object.

Example with acceleration.

Q: How much work is done pushing a 46 kg crate to the right with $\mathbf{a}=2.5 \text{ m/s}^2\hat{i}$ over 10.3 m across a floor with $\mu_k=0.4$?

Sol: First find F_{app} , then $W=F_{\text{app}}\Delta r$
 $\mathbf{F}_{\text{net}} = m\mathbf{a} = \mathbf{F}_{\text{app}} + \mathbf{f}_k$

... $W = 3042 \text{ J}$.



© Cengage Learning. All Rights Reserved.

Work Is An Energy Transfer

If the work done on a system is positive, energy is transferred into the system.


If the work done on the system is negative, energy is transferred out of the system.

Example system: a mass on a string (a pendulum).

Energy that can be stored: gravitational potential energy and kinetic energy

Q: How can you use work to transfer energy in?

Q: How can you use work to transfer energy out?



Scalar (Dot) Product of Two Vectors

The scalar product of two vectors is written as $\vec{A} \cdot \vec{B}$.

- It is also called the dot product.


$$\vec{A} \cdot \vec{B} = AB \cos \theta$$

- θ is the angle *between* A and B

Applied to work, this means

$$W = F \Delta r \cos \theta = \vec{F} \cdot \Delta \vec{r}$$

(for the case of constant forces)



© Cengage Learning. All Rights Reserved.

Scalar Product, cont

The scalar product is commutative.

- $\vec{A} \cdot \vec{B} = \vec{B} \cdot \vec{A}$

The scalar product obeys the distributive law of multiplication.

- $\vec{A} \cdot (\vec{B} + \vec{C}) = \vec{A} \cdot \vec{B} + \vec{A} \cdot \vec{C}$

Dot Products of Unit Vectors

$$\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1$$

$$\hat{i} \cdot \hat{j} = \hat{i} \cdot \hat{k} = \hat{j} \cdot \hat{k} = 0$$

Using component form with vectors:


$$\vec{A} = A_x \hat{i} + A_y \hat{j} + A_z \hat{k}$$

$$\vec{B} = B_x \hat{i} + B_y \hat{j} + B_z \hat{k}$$

$$\vec{A} \cdot \vec{B} = A_x B_x + A_y B_y + A_z B_z$$

In the special case where

$$\vec{A} = A_x \hat{i} + A_y \hat{j} + A_z \hat{k}$$



Dot Products - example

$$\vec{A} \cdot \vec{B} = A_x B_x + A_y B_y + A_z B_z$$

$$\vec{A} \cdot \vec{B} = AB \cos \theta$$

Ex) (P. 20) Find the angle between $\mathbf{A}=5.8\mathbf{i}-3.5\mathbf{j}-6.2\mathbf{k}$
and $\mathbf{B}=8.2\mathbf{i}+4.3\mathbf{j}-7.0\mathbf{k}$

$$\vec{A} \cdot \vec{B} = 5.8 \cdot 8.2 + (-3.5) \cdot 4.3 + (-6.2) \cdot (-7.0)$$

$$\mathbf{A} \cdot \mathbf{B} = 75.91$$

Find $|\mathbf{A}| = 9.145$ and $|\mathbf{B}| = 11.61$

Then $|\mathbf{A}||\mathbf{B}| \cos \theta = 75.91$

$$\cos \theta = 75.91 / (9.145 \cdot 11.61) \rightarrow \theta = 44.4^\circ \text{ (fixed)}$$

Work Done by a Varying Force

$$W = \int \vec{F} \cdot d\vec{r}$$

To use $W = F \Delta r \cos \theta$, the force must be constant, so the equation cannot be used to calculate the work done by a varying force.


Assume that during a very small displacement, Δx , F is constant.

For that displacement, $W \sim F \Delta x$

For all of the intervals,

$$W \approx \sum_{x_i}^{x_f} F_x \Delta x$$

The total work done for the displacement from x_i to x_f is approximately equal to the sum of the areas of all the rectangles.



a

© Cengage Learning. All Rights Reserved.

Work Done by a Varying Force, cont.

Let the size of the small displacements approach zero .

Since


$$\lim_{\Delta x \rightarrow 0} \sum_{x_i}^{x_f} F_x \Delta x = \int_{x_i}^{x_f} F_x dx$$

Or, more generally,

$$W = \int \vec{F} \cdot d\vec{r}$$

The work done is equal to the area under the curve between x_i and x_f .

The work done by the component F_x of the varying force as the particle moves from x_i to x_f is *exactly* equal to the area under the curve.



b

© Cengage Learning. All Rights Reserved.

Work Done by a Varying Force, cont.

$$W = \int \vec{F} \cdot d\vec{r}$$

Example) P. 7.41. Find the work done on a particle moving according to this graph ...

a) between $x=0$ to $x=10$ m

Ans: 2800 J

b) between $x=0$ to $x=15$ m

Ans: 2100 J

Work Done by a Varying Force, cont.

$$W = \int \vec{F} \cdot d\vec{r}$$

Example) P. 7.45. $F(x) = \frac{A}{\sqrt{x}}$

With $A = 3.00 \text{ Nm}^{1/2}$ what is the work done by this force when the object moves from $x_i = 0$ to $x_f = 1.0$?

Work Done By Multiple Forces

If more than one force acts on a system *and the system can be modeled as a particle*, the total work done on the system is the work done by the net force.

$$\sum W = W_{\text{ext}} = \int_{x_i}^{x_f} (\sum F_x) dx$$

In the general case of a net force whose magnitude and direction may vary.

$$\sum W = W_{\text{ext}} = \int_{x_i}^{x_f} (\sum \mathbf{F}) d\vec{r}$$

The subscript “ext” indicates the work is done by an *external* agent on the system.

Work Done by Multiple Forces, cont.

If the system cannot be modeled as a particle, then the total work is equal to the algebraic sum of the work done by the individual forces.

$$\sum W = W_{\text{ext}} = \sum_{\text{forces}} (\mathbf{F} \cdot d\mathbf{r})$$

- Remember work is a scalar, so this is the sum of a bunch of scalars.

Work Done By A Spring

The spring force varies with position according to Hooke's Law:


$$F_s = -kx$$

Where F_s is the spring force, k is the spring constant (N/m), and x is the distance of [the block] from the equilibrium position.


The block is on a horizontal, frictionless surface.

Observe the motion of the block with various values of the spring constant.

See 7.9.swf



Spring Force (Hooke's Law)



© Cengage Learning. All Rights Reserved.

The force exerted by the spring is

$$F_s = -kx$$

- x is the position of the block with respect to the equilibrium position ($x = 0$).
- k is called the spring constant or force constant and measures the stiffness of the spring.
 - k measures the *stiffness* of the spring.

This is called Hooke's Law.

Hooke's Law, cont.


The vector form of Hooke's Law is

$$\mathbf{F}_s = F_x \hat{\mathbf{i}} = -kx\hat{\mathbf{i}}$$

When x is positive (spring is stretched),
 F is negative

When x is 0 (at the equilibrium
position), F is 0

When x is negative (spring is
compressed), F is positive



Hooke's Law, final

The force exerted by the spring is always directed opposite to the displacement from equilibrium.

The spring force is sometimes called the *restoring force*.

If the block is released it will oscillate back and forth between $-x$ and x .

Work Done by a Spring


Identify the block as the system.

Calculate the work by the spring as the block moves from $x_i = -x_{\max}$ to $x_f = 0$.

$$W_s = \int \vec{F}_s \cdot d\vec{r} = \int_{-x_{\max}}^0 (-kx \hat{i}) \cdot (dx \hat{i})$$

$$W_s = \int_{-x_{\max}}^0 -kx dx = \frac{1}{2}(-k)(0^2 - (-x_{\max})^2) = -\frac{1}{2}kx_{\max}^2$$

Notice that this is the area of the shaded triangle!



Q: What is the net work done by the spring if the block moves from $-x_{\max}$ to x_{\max} ?

Q: What is the work done by a hand pushing the block at constant speed from $x_i=0$ to $x_f=0.5$ m if $k=300$ N/m?

Work Done by a Spring, cont.

Assume the block undergoes an arbitrary displacement from $x = x_i$ to $x = x_f$.

The work done by the spring on the block is

$$W_s = \int_{x_i}^{x_f} (-kx) dx = \frac{1}{2}kx_i^2 - \frac{1}{2}kx_f^2$$

- If the motion ends where it begins, $W = 0$

Spring with an Applied Force

Suppose an external agent, F_{app} , stretches the spring.

The applied force is equal and opposite to the spring force (if block speed constant).

$$\mathbf{F}_{\text{app}} = F_{\text{app}} \hat{\mathbf{i}} = -\mathbf{F}_s = -(-kx\hat{\mathbf{i}}) = kx\hat{\mathbf{i}}$$


Work done by F_{app} as the block moves from $-x_{\text{max}}$ to $x = 0$ is equal to

$$-\frac{1}{2} kx_{\text{max}}^2$$

For any displacement, the work done by the applied force is

$$W_{\text{app}} = \int_{x_i}^{x_f} (kx) dx = \frac{1}{2} kx_f^2 - \frac{1}{2} kx_i^2$$

If the process of moving the block is carried out very slowly, then \vec{F}_{app} is equal in magnitude and opposite in direction to \vec{F}_s at all times.



© Cengage Learning. All Rights Reserved.

Kinetic Energy

One possible result of work acting as an influence on a system is that the system changes its speed.

The system could possess *kinetic energy*.

Kinetic Energy is the energy of a particle due to its motion.

- $K = \frac{1}{2} mv^2$
 - K is the kinetic energy
 - m is the mass of the particle
 - v is the speed of the particle

A change in kinetic energy is one possible result of doing work to transfer energy into a system.

Kinetic Energy, cont


Calculating the work:

$$W_{\text{ext}} = \int_{x_i}^{x_f} \sum F \, dx = \int_{x_i}^{x_f} ma \, dx$$

$$W_{\text{ext}} = \int_{v_i}^{v_f} mv \, dv$$

$$W_{\text{ext}} = \frac{1}{2}mv_f^2 - \frac{1}{2}mv_i^2$$

$$W_{\text{ext}} = K_f - K_i = \Delta K$$



© Cengage Learning. All Rights Reserved.

Work-Kinetic Energy Theorem

The Work-Kinetic Energy Theorem states $W_{\text{ext}} = K_f - K_i = \Delta K$

When work is done on a system and the only change in the system is in its speed, the net work done on the system equals the change in kinetic energy of the system.

- The speed of the system increases if the work done on it is positive.
- The speed of the system decreases if the net work is negative.
- Also valid for changes in rotational speed

The work-kinetic energy theorem is not valid if other changes (besides its speed) occur in the system or if there are other interactions with the environment besides work.

The work-kinetic energy theorem applies to the speed of the system, not its velocity.


Work-Kinetic Energy Theorem – Example

The block is the system and three external forces act on it.

The normal and gravitational forces do no work since they are perpendicular to the direction of the displacement.

$$W_{\text{ext}} = \Delta K = \frac{1}{2} m v_f^2 - 0$$

The answer could be checked by modeling the block as a particle and using the kinematic equations.



© Cengage Learning. All Rights Reserved.

Potential Energy

Potential energy is energy determined by the configuration of a system in which the components of the system interact by forces.

- The forces are internal to the system.
- Can be associated with only specific types of forces acting between members of a system

Gravitational Potential Energy

The system is the Earth and the book.


Do work on the book by lifting it slowly through a vertical displacement.

$$\Delta \mathbf{r} = (y_f - y_i) \hat{\mathbf{j}}$$

The work done on the system must appear as an increase in the energy of the system.

The energy storage mechanism is called *potential energy*.

The work done by the agent on the book–Earth system is $mgy_f - mgy_i$.



© Cengage Learning. All Rights Reserved.

Gravitational Potential Energy, cont

Assume the book in fig. 7.15 is allowed to fall.

There is no change in kinetic energy since the book starts and ends at rest.

Gravitational potential energy is the energy associated with an object at a given location above the surface of the Earth.

$$W_{\text{ext}} = (\mathbf{F}_{\text{app}}) \cdot \Delta \mathbf{r}$$

$$W_{\text{ext}} = (mg\hat{\mathbf{j}}) \cdot [(y_f - y_i)\hat{\mathbf{j}}]$$

$$W_{\text{ext}} = mgy_f - mgy_i$$

Gravitational Potential Energy, final

The quantity mgy is identified as the gravitational potential energy, U_g .

- $U_g = mgy$

Units are joules (J)

Is a scalar

Work may change the gravitational potential energy of the system.

- $W_{\text{ext}} = \Delta u_g$

Potential energy is always associated with a system of two or more interacting objects.

Gravitational Potential Energy, Problem Solving

The gravitational potential energy depends only on the vertical height of the object above Earth's surface.

In solving problems, you must choose a reference configuration for which the gravitational potential energy is set equal to some reference value, normally zero.

- The choice is arbitrary because you normally need the difference in potential energy, which is independent of the choice of reference configuration.

Often having the object on the surface of the Earth is a convenient zero gravitational potential energy configuration.

The problem may suggest a convenient configuration to use.

Elastic Potential Energy

Elastic Potential Energy is associated with a spring.

The force the spring exerts (on a block, for example) is $F_s = -kx$

The work done by an external applied force on a spring-block system is

- $W = \frac{1}{2} kx_f^2 - \frac{1}{2} kx_i^2$
- The work is equal to the difference between the initial and final values of an expression related to the configuration of the system.

Elastic Potential Energy, cont.


This expression is the elastic potential energy:

$$U_s = \frac{1}{2} kx^2$$

The elastic potential energy can be thought of as the energy stored in the deformed spring.

The stored potential energy can be converted into kinetic energy.

Observe the effects of different amounts of compression of the spring.



Elastic Potential Energy, final

The elastic potential energy stored in a spring is zero whenever the spring is not deformed ($U = 0$ when $x = 0$).


- The energy is stored in the spring only when the spring is stretched or compressed.

The elastic potential energy is a maximum when the spring has reached its maximum extension or compression.

The elastic potential energy is always positive.

- x^2 will always be positive.

Energy Bar Chart Example




An energy bar chart is an important graphical representation of information related to the energy of a system.

- The vertical axis represents the amount of energy of a given type in the system.
- The horizontal axis shows the types of energy in the system.

In a, there is no energy.

- The spring is relaxed, the block is not moving


Energy Bar Chart Example, cont.



Between b and c, the hand has done work on the system.

- The spring is compressed.
- There is elastic potential energy in the system.
- There is no kinetic energy since the block is held steady.

Energy Bar Chart Example, final



In d, the block has been released and is moving to the right while still in contact with the spring.

- The elastic potential energy of the system decreases while the kinetic energy increases.

In e, the spring has returned to its relaxed length and the system contains only kinetic energy associated with the moving block.

Internal Energy


The energy associated with an object's temperature is called its *internal energy*, E_{int} .

In this example, the surface is the system.

The friction does work and increases the internal energy of the surface.

When the book stops, all of its kinetic energy has been transformed to internal energy.

The total energy remains the same.



© Cengage Learning. All Rights Reserved.

Conservative Forces

The work done by a conservative force on a particle moving between any two points is independent of the path taken by the particle.

The work done by a conservative force on a particle moving through any closed path is zero.

- A closed path is one in which the beginning and ending points are the same.

Examples of conservative forces:

- Gravity
- Spring force

Conservative Forces, cont

We can associate a potential energy for a system with any conservative force acting between members of the system.

- This can be done only for conservative forces.
- In general: $W_{int} = - \Delta U$
 - W_{int} is used as a reminder that the work is done by one member of the system on another member and is internal to the system.
- Positive work done by an outside agent on a system causes an increase in the potential energy of the system.
- Work done on a component of a system by a conservative force internal to an isolated system causes a decrease in the potential energy of the system.

Non-conservative Forces

A non-conservative force does not satisfy the conditions of conservative forces.

Non-conservative forces acting in a system cause a *change* in the mechanical energy of the system.


$$E_{\text{mech}} = K + U$$

- K includes the kinetic energy of all moving members of the system.
- U includes all types of potential energy in the system.

Non-conservative Forces, cont.

The work done against friction is greater along the brown path than along the blue path.

Because the work done depends on the path, friction is a non-conservative force.



© Cengage Learning. All Rights Reserved.

Conservative Forces and Potential Energy

Define a potential energy function, U , such that the work done by a conservative force equals the decrease in the potential energy of the system.

The work done by such a force, F , is

$$W_{\text{int}} = \int_{x_i}^{x_f} F_x dx = -\Delta U$$

- ΔU is negative when F and x are in the same direction

Conservative Forces and Potential Energy

The conservative force is related to the potential energy function through.

$$F_x = - \frac{dU}{dx}$$

The x component of a conservative force acting on an object within a system equals the negative of the potential energy of the system with respect to x.

- Can be extended to three dimensions

Conservative Forces and Potential Energy – Check


Look at the case of a deformed spring:

$$F_s = - \frac{dU_s}{dx} = - \frac{d}{dx} \left(\frac{1}{2} kx^2 \right) = - kx$$

- This is Hooke's Law and confirms the equation for U

U is an important function because a conservative force can be derived from it.

Energy Diagrams and Equilibrium



Motion in a system can be observed in terms of a graph of its position and energy.

In a spring-mass system example, the block oscillates between the turning points, $x = \pm x_{\max}$.

The block will always accelerate back toward $x = 0$.


Energy Diagrams and Stable Equilibrium

The $x = 0$ position is one of **stable equilibrium**.

- Any movement away from this position results in a force directed back toward $x = 0$.

Configurations of stable equilibrium correspond to those for which $U(x)$ is a minimum.

$x = x_{\max}$ and $x = -x_{\max}$ are called the turning points.



a

© Cengage Learning. All Rights Reserved.


Energy Diagrams and Unstable Equilibrium

$F_x = 0$ at $x = 0$, so the particle is in equilibrium.

For any other value of x , the particle moves away from the equilibrium position.

This is an example of ***unstable equilibrium***.

Configurations of unstable equilibrium correspond to those for which $U(x)$ is a maximum.



© Cengage Learning. All Rights Reserved.

Neutral Equilibrium

Neutral equilibrium occurs in a configuration when U is constant over some region.

A small displacement from a position in this region will produce neither restoring nor disrupting forces.


Potential Energy in Molecules

There is potential energy associated with the force between two neutral atoms in a molecule which can be modeled by the *Lennard-Jones* function.


$$U(x) = 4 \epsilon \left[\left(\frac{\sigma}{x} \right)^{12} - \left(\frac{\sigma}{x} \right)^6 \right]$$

Find the minimum of the function (take the derivative and set it equal to 0) to find the separation for stable equilibrium.


The graph of the Lennard-Jones function shows the most likely separation between the atoms in the molecule (at minimum energy).




7.9 Hooke's Law and Restoring Forces



7.15 Work Done By External Agent on System of the Book




7.16 Conservation Mechanical Energy in Block-Spring System



7.18 Work When Kinetic Friction is Present



7.20 An Oscillating Block-Spring System



7.20b Conservation of Mechanical Energy for a Pendulum

