

Chapter 27

Magnetic Fields



PHYS 2321
Week 9: Magnetism




Day 2 Outline

1) Hwk: Ch. 27 P. 3-5,10,13,17,21,28, 33,39,49 -Due next Wed
   Read Ch. 27.1-7

2) Magnetic Fields and Forces (Ch 27)
   a. Bar magnets (behavior and field lines)
   b. Force on moving charged particles
      * Lorentz force: $\vec{F}_B = q\,\vec{v} \times \vec{B}$
      * Right hand rule

Notes: Quiz next Wednesday (on B-fields). Try testbank quiz. Ch. 25 &26 graded. Mean = 9.4/10. Checked P.9 MQ 5,7,9. Gone Friday. Instead, work on homework problems. Figure out whether parallel currents will attract or repel.


A Brief History of Magnetism




- 13th century BC
  - Chinese used a compass
    - Uses a magnetic needle
    - Probably an invention of Arabic or Indian origin
- 800 BC
  - Greeks
    - Discovered magnetite (Fe_3O_4) attracts pieces of iron

A Brief History of Magnetism

- The mid-oceanic ridge between N. America and Europe shows that Earth's magnetic field has switched polarity.





Figures by the USGS

A Brief History of Magnetism, 4

- 1819
  - Hans Christian Oersted
    - Discovered the relationship between electricity and magnetism
    - An electric current in a wire deflected a nearby compass needle



© 2004 Thomson - Brooks/Cole

Fundamental source of B-fields: current and current loops!

Magnetic Poles

- Every magnet, regardless of its shape, has two poles
  - Called north and south poles
  - Poles exert forces on one another
    - Similar to the way electric charges exert forces on each other
    - Like poles repel each other
      - N-N or S-S
    - Unlike poles attract each other
      - N-S

Magnetic Poles, cont.

- The poles received their names due to the way a magnet behaves in the Earth's magnetic field
- If a bar magnet is suspended so that it can move freely, it will rotate
  - The magnet's north pole points toward the Earth's north geographic pole
    - This means the Earth's north geographic pole is a magnetic south pole
    - Similarly, the Earth's south geographic pole is a magnetic north pole


Magnetic Poles, final

- The force between two poles varies as the inverse square of the distance between them
- A single magnetic pole has never been isolated ("No magnetic monopoles")
  - In other words, magnetic poles are always found in pairs
  - All attempts so far to detect an isolated magnetic pole has been unsuccessful
    - No matter how many times a permanent magnetic is cut in two, each piece always has a north and south pole

Magnetic Fields

- Reminder: an electric field surrounds any electric charge
- The region of space surrounding any *moving* electric charge also contains a magnetic field
- A magnetic field also surrounds a magnetic substance making up a permanent magnet


Magnetic Fields, cont.



- A vector quantity
- Symbolized by $\vec{B}$
- Direction is given by the direction a north pole of a compass needle points in that location
- Magnetic field lines can be traced out by a compass

Magnetic Field Lines, Bar Magnet Example

- The compass can be used to trace the field lines
- The lines outside the magnet point from the North pole to the South pole
- Inside, lines point South to North, continuing the loops




© Thomson Higher Education

PLAY ACTIVE FIGURE

Magnetic Field Lines, Bar Magnet


- Iron filings are used to show the pattern of the magnetic field lines
- The direction of the field is the direction a north pole would point



© 2003 Thomson - Brooks Cole

Magnetic Field Lines, Unlike Poles


- Iron filings are used to show the pattern of the magnetic field lines
- The direction of the field is the direction a north pole would point
  - Compare to the electric field produced by an electric dipole



© 2003 Thomson - Brooks Cole


Magnetic Field Lines, Like Poles

- Iron filings are used to show the pattern of the magnetic field lines
- The direction of the field is the direction a north pole would point
  - Compare to the electric field produced by like charges




© 2003 Thomson - Brooks Cole

Definition of Magnetic Field




- The magnetic field at some point in space can be defined in terms of the magnetic force, $\vec{F}_B$
- The magnetic force will be exerted on a charged particle moving with a velocity, $\vec{v}$
  - Assume (for now) there are no gravitational or electric fields present

Force on a Charge Moving in a Magnetic Field



- The magnitude F_B of the magnetic force exerted on the particle is proportional to the charge, q, and to the speed, v, of the particle
- When a charged particle moves parallel to the magnetic field vector, the magnetic force acting on the particle is zero
- When the particle's velocity vector makes any angle $\theta \neq 0$ with the field, the force acts in a direction perpendicular to both the velocity and the field

F_B on a Charge Moving in a Magnetic Field, final



- The magnetic force exerted on a positive charge is in the direction opposite the direction of the magnetic force exerted on a negative charge moving in the same direction
- The magnitude of the magnetic force is proportional to sin θ, where θ is the angle the particle's velocity makes with the direction of the magnetic field

PHYS 2321
Week 10: Magnetism



Day 1 Outline

1) Hwk: Ch. 27 P. 3-5,10,13,17,21,28, 33,39,49 -Due Wed
   Read Ch. 27.1-7

2) Magnetic Fields and Forces (Ch 27)

   a. Force on moving charged particles

      * Lorentz force: $\vec{F}_B = q\,\vec{v} \times \vec{B}$

      * Right hand rule


   b. Force on current-carrying wire

   c. Combining electric and magnetic forces on a charge.

$$\vec{F} = q\,(\vec{E} + \vec{v} \times \vec{B})$$

Notes: Quiz next Wednesday (on B-fields). Try test bank quiz.
Ch. 26 graded, but not ready for return.
Last Friday: Figure out whether parallel currents will attract or repel.

Force on a Charge Moving in a Magnetic Field, Formula




- The properties can be summarized in a vector equation:

$$\vec{F}_B = q\left(\vec{v} \times \vec{B}\right)$$

  - $\vec{F}_B$ is the magnetic force
  - q is the charge
  - $\vec{v}$ is the velocity of the moving charge
  - $\vec{B}$ is the magnetic field

More About Direction




© Thomson Higher Education

- $\vec{F}_B$ is perpendicular to the plane formed by $\vec{v}$ and $\vec{B}$
- Oppositely directed forces exerted on oppositely charged particles will cause the particles to move in opposite directions

Direction: Right-Hand Rule #1


- The fingers point in the direction of **v**
- Orient hand so fingers bend towards **B**
- The thumb points in the direction of $\mathbf{F}_B$ which is the direction of the force on a *positive (+)* charge.



More About Magnitude of F


- The magnitude of the magnetic force on a charged particle is $F_B = |q|\ v\ B\ \sin\theta$
  - θ is the smaller angle between v and B
  - F_B is zero when the field and velocity are parallel or antiparallel
    - $\theta = 0$ or 180°
  - F_B is a maximum when the field and velocity are perpendicular
    - $\theta = 90^o$

Differences Between Electric and Magnetic Fields




- Direction of force
  - The electric force acts along the direction of the electric field
  - The magnetic force acts perpendicular to the magnetic field
- Motion
  - The electric force acts on a charged particle regardless of whether the particle is moving
  - The magnetic force acts on a charged particle only when the particle is in motion

More Differences Between Electric and Magnetic Fields



- Work
  - The electric force does work in displacing a charged particle
  - The magnetic force associated with a steady magnetic field **does no work** when a particle is displaced
    - This is because the force is perpendicular to the displacement

Work in Fields, cont.



- The kinetic energy of a charged particle moving through a magnetic field cannot be altered by the magnetic field alone
- When a charged particle moves with a given velocity through a magnetic field, the field can alter the direction of the velocity, but not the speed or the kinetic energy

Units of Magnetic Field


- The SI unit of magnetic field is the tesla (T)

  (Technically, a tesla is a magnetic flux density. It can be multiplied by an area to get total magnetic flux through that area measured in weber, Wb.)

  - Wb is a weber
- A non-SI commonly used unit is a gauss (G)
  - $1\ T = 10^4\ G$


Notation Notes

- When vectors are perpendicular to the page, dots and crosses are used
  - The dots represent the arrows coming out of the page
  - The crosses represent the arrows going into the page



© Thomson Higher Education

PHYS 2321
Week 10: Magnetism



Day 2 Outline

1) Quiz 4 on Magnetism

2) Magnetic Fields and Forces (Ch 27)

a. Force on current-carrying conductor (cont.)

b. Torque on a current loop

c. Cyclotron radius and frequency

d. Combining electric and magnetic forces on a charge.

$$\vec{F} = q\left(\vec{E} + \vec{v} \times \vec{B}\right)$$

Hwk: Ch. 27 P. 3-5,10,13,17,21,28, 33,39,49 -Due < 2 pm.


Next: Ch. 28 P. 1,4,5,19,27,29,37,38, MQ. 1-13 odd

Notes:

Ch. 26 graded. Mean = 4.98,4.93/5.

Charged Particle in a Magnetic Field

- Consider a particle moving in an external magnetic field with its velocity perpendicular to the field
- The force is always directed toward the center of the circular path
- The magnetic force causes a centripetal acceleration, changing the direction of the velocity of the particle



© Thomson Higher Education

PLAY ACTIVE FIGURE

Force on a Charged Particle

- For q in uniform circular motion: $q\vec{v}\times\vec{B}=qvB$
- Equating the magnetic and centripetal forces:


$$qvB=\frac{mv^2}{r}$$

- Solving for r:

$$r=\frac{mv}{qB}$$

  - r is proportional to the linear momentum of the particle and inversely proportional to the magnetic field

More About Motion of Charged Particle



- The angular speed of the particle is

$$\omega = \frac{v}{r} = \frac{qB}{m}$$

  - The angular speed, ω, is also referred to as the **cyclotron frequency**
- The period of the motion is


$$P = \frac{2\pi}{\omega}$$

Motion of a Particle, General

- If a charged particle moves in a magnetic field at some arbitrary angle with respect to the field, its path is a helix
- Same equations apply, with

$$v_{\perp} = \sqrt{v_y^2 + v_z^2}$$


- Use the active figure to vary the initial velocity and observe the resulting motion



PLAY
ACTIVE FIGURE


Bending of an Electron Beam

- Electrons are accelerated from rest through a potential difference
- The electrons travel in a curved path
- Conservation of energy will give *v*
- Other parameters can be found


© 2004 Thomson - Brooks/Cole

Particle in a Nonuniform Magnetic Field


- The motion is complex
- For example, the particles can oscillate back and forth between two positions
- This configuration is known as a *magnetic bottle*



©2004 Thomson - Brooks/Cole


Van Allen Radiation Belts

- The Van Allen radiation belts consist of charged particles surrounding the Earth in doughnut-shaped regions
- The particles are trapped by the Earth's magnetic field
- The particles spiral from pole to pole
  - May result in Auroras



©2004 Thomson - Brooks/Cole

Charged Particles Moving in Electric and Magnetic Fields




- In many applications, charged particles will move in the presence of both magnetic and electric fields
- In that case, the total force is the sum of the forces due to the individual fields
- In general:

$$\vec{F} = q\left\{\vec{E} + \left(\vec{v} \times \vec{B}\right)\right\}$$

Velocity Selector

- Used when particles of a certain velocity are needed.
- A uniform electric field is perpendicular to a uniform magnetic field



PLAY ACTIVE FIGURE


Velocity Selector, cont.

- When the force due to the electric field is equal but opposite to the force due to the magnetic field, the particle moves in a straight line
- This occurs for velocities of value

  $v = E / B$

Mass Spectrometer

- A mass spectrometer separates ions according to their mass-to-charge ratio
- A beam of ions passes through a velocity selector and enters a second magnetic field



© Thomson Higher Education


PLAY ACTIVE FIGURE

Mass Spectrometer, cont.

- After entering the second magnetic field, the ions move in a semicircle of radius r before striking a detector at P
- If the ions are positively charged, they deflect to the left
- If the ions are negatively charged, they deflect to the right

Thomson's *e*/*m* Experiment

- Electrons are accelerated from the cathode
- They are deflected by electric and magnetic fields
- The beam of electrons strikes a fluorescent screen
- e/m was measured




©2004 Thomson - Brooks/Cole

Cyclotron

- A **cyclotron** is a device that can accelerate charged particles to very high speeds
- The energetic particles produced are used to bombard atomic nuclei and thereby produce reactions
- These reactions can be analyzed by researchers

Cyclotron, 2

- D_1 and D_2 are called *dees* because of their shape
- A high frequency alternating potential is applied to the dees
- A uniform magnetic field is perpendicular to them



(a)

© Thomson Higher Education



(b)

© Thomson Higher Education

Cyclotron, 3

- A positive ion is released near the center and moves in a semicircular path
- The potential difference is adjusted so that the polarity of the dees is reversed in the same time interval as the particle travels around one dee
- This ensures the kinetic energy of the particle increases each trip


Cyclotron, final

- The cyclotron's operation is based on the fact that T is independent of the speed of the particles and of the radius of their path

$$K = \frac{1}{2}mv^2 = \frac{q^2B^2R^2}{2m}$$

- When the energy of the ions in a cyclotron exceeds about 20 MeV, relativistic effects come into play


Magnetic Force on a Current Carrying Conductor



- A force is exerted on a current-carrying wire placed in a magnetic field
  - The current is a collection of many charged particles in motion
- The direction of the force is given by the right-hand rule

Force on a Wire


- In this case, there is no current, so there is no force
- Therefore, the wire remains vertical



(b)

Force on a Wire (2)


- The magnetic field is into the page
- The current is up the page
- The force is to the left



(c)

Force on a Wire, (3)

- The magnetic field is into the page
- The current is down the page
- The force is to the right




(d)

©2004 Thomson - Brooks/Cole

Force on a Wire, equation

- The magnetic force is exerted on each moving charge in the wire
  - $\vec{F} = q\ (\vec{v}_d \times \vec{B})$
- The total force is the product of the force on one charge and the number of charges
  - $\vec{F}_{tot} = q(\vec{v}_d \times \vec{B})nAL$



Force on a Wire, (4)

- In terms of the current, this becomes

$$\vec{F}_B = I\ \vec{L} \times \vec{B}$$

  - I is the current
  - $\vec{L}$ is a vector that points in the direction of the current
    - Its magnitude is the length *L* of the segment
  - $\vec{B}$ is the magnetic field


Force on a Wire, Arbitrary Shape

- Consider a small segment of the wire,
- The force exerted on this segment is

$$d\vec{F}_B = Id\vec{s} \times \vec{B}$$

- The total force is


$$\vec{F}_B = \int I\, d\vec{s} \times \vec{B}$$



© Thomson Higher Education

Torque on a Current Loop


- The rectangular loop carries a current I in a uniform magnetic field
- No magnetic force acts on sides 1 & 3
  - The wires are parallel to the field and $\mathbf{\Lambda} \cdot \mathbf{B} = 0$



(a)

Torque on a Current Loop, 2

- There is a force on sides 2 & 4 since they are perpendicular to the field
- The magnitude of the magnetic force on these sides will be:
  - $F_2 = F_4 = I\ a\ B$
- The direction of F_2 is out of the page
- The direction of F_4 is into the page




(a)

Torque on a Current Loop, 3

- The forces are equal and in opposite directions, but not along the same line of action
- The forces produce a torque around point *O*


View loop from **below**:



(b)

© Thomson Higher Education

Torque on a Current Loop, Equation




- The maximum torque is found by:

$$\tau = IaB\frac{b}{2} + IaB\frac{b}{2} = IaBb$$

- The area enclosed by the loop is *ab*, so $\tau_{max} = IAB$
  - This maximum value occurs only when the field is parallel to the plane of the loop

Torque on a Current Loop, General


- Assume the magnetic field makes an angle of
$\theta < 90^\circ$ with a line perpendicular to the plane of the loop
- The net torque about point *O* will be $\tau = IAB \sin \theta$
- Use the active figure to vary the initial settings and observe the resulting motion



© Thomson Higher Education

PLAY ACTIVE FIGURE


Torque on a Current Loop, Summary



- The torque has a maximum value when the field is perpendicular to the normal to the plane of the loop
- The torque is zero when the field is parallel to the normal to the plane of the loop
- $\tau = I\,\vec{A} \times \vec{B}$ where $\vec{A}$ is perpendicular to the plane of the loop and has a magnitude equal to the area of the loop


Direction

- A right-hand rule can be used to determine the direction of **A**
- Curl your fingers in the direction of the current in the loop
- Your thumb points in the direction of **A**




© Thomson Higher Education

Hall Effect (skip)



- When a current carrying conductor is placed in a magnetic field, a potential difference is generated in a direction perpendicular to both the current and the magnetic field
- This phenomena is known as the Hall effect
- It arises from the deflection of charge carriers to one side of the conductor as a result of the magnetic forces they experience


Hall Effect, cont.




- The Hall effect gives information regarding the sign of the charge carriers and their density
- It can also be used to measure magnetic fields

Hall Voltage

- This shows an arrangement for observing the Hall effect
- The Hall voltage is measured between points *a* and *c*



Hall Voltage, cont



© Thomson Higher Education

- When the charge carriers are negative, the upper edge of the conductor becomes negatively charged
  - *c* is at a lower potential than *a*
- When the charge carriers are positive, the upper edge becomes positively charged
  - *c* is at a higher potential than *a*

Hall Voltage, final

- $\Delta V_H = E_H d = v_d B d$
  - d is the width of the conductor
  - v_d is the drift velocity
  - If B and d are known, v_d can be found
- $$\Delta V_H = \frac{I B}{n q t} = \frac{R_H I B}{t}$$
  - $R_H = 1 / nq$ is called the Hall coefficient
  - A properly calibrated conductor can be used to measure the magnitude of an unknown magnetic field