

Quantifying shoreline modifications adjacent to eelgrass meadows in the Strait of Georgia Bioregion

John Cristiani, Katherine Bannar-Martin, Emily M. Rubidge

Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
Nanaimo, British Columbia
V9T 6N7

2022

**Canadian Technical Report of
Fisheries and Aquatic Sciences #####**



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Canada 

Canadian Technical Report of Fisheries and Aquatic Sciences

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Rapport technique canadien des sciences halieutiques et aquatiques

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques de Pêches et Océans Canada, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications à part entière. Le titre exact figure au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre.

Les numéros 1 à 456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Canadian Technical Report of
Fisheries and Aquatic Sciences nnn

2022

QUANTIFYING SHORELINE MODIFICATIONS ADJACENT TO EELGRASS MEADOWS IN THE
STRAIT OF GEORGIA BIOREGION

by

John M. Cristiani¹ Katherine H. Bannar-Martin² and Emily M. Rubidge²

¹Pacific Biological Station
Fisheries and Oceans Canada, 3190 Hammond Bay Road
Nanaimo, British Columbia, V9T 6N7, Canada
²Institute of Ocean Sciences
Fisheries and Oceans Canada, 9860 W Saanich Road
Sidney, British Columbia, V8L 4B2, Canada

16
17

© His Majesty the King in Right of Canada, 2022
Cat. No. Fs97-6/nnnE-PDF ISBN ISSN 1488-5379

18 Correct citation for this publication:

19 Cristiani, J.C., Bannar-Martin, K.H, and Rubidge, E.M. 2022. Quantifying shoreline modifications
20 adjacent to eelgrass meadows in the Strait of Georgia Bioregion. Can. Tech. Rep. Fish. Aquat.
21 Sci. nnn: v + 9p.

23	ABSTRACT	iv
24	RÉSUMÉ	v
25	1 Introduction	1
26	2 Methods	1
27	2.1 Seagrass spatial data	1
28	2.2 Shoreline area adjacent to seagrass meadows	2
29	2.3 Shoreline modifications	3
30	3 Results	4
31	4 Discussion	7
32	5 Data availability	7
33	6 References	8

ABSTRACT

35 Cristiani, J.C., Bannar-Martin, K.H, and Rubidge, E.M. 2022. Quantifying shoreline modifications
36 adjacent to eelgrass meadows in the Strait of Georgia Bioregion. Can. Tech. Rep. Fish. Aquat.
37 Sci. nnn: v + 9 p.

38 Here is the abstract text. Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do
39 eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis
40 nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure
41 dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint
42 occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

RÉSUMÉ

44 Cristiani, J.C., Bannar-Martin, K.H, and Rubidge, E.M. 2022. Quantifying shoreline modifications
45 adjacent to eelgrass meadows in the Strait of Georgia Bioregion. Can. Tech. Rep. Fish. Aquat.
46 Sci. nnn: v + 9 p.

47 Voici le résumé. Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor
48 incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation
49 ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit
50 in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat
51 non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

1 Introduction

The health and functioning of coastal marine ecosystems are under threat from a variety of human activities (Halpern et al. 2019). Coastal activities such as agriculture, industrial and residential development, forestry, and shoreline hardening can create pressures on the marine environment. A modified shoreline may alter levels of sedimentation, nutrient runoff, pollution, and wave energy (Dethier et al. 2016; Todd et al. 2019). For coastal biogenic habitat in British Columbia such as seagrass, these pressures may impact seagrass productivity and survival, and thus impact the community of species that rely on seagrass (Iacarella et al. 2018; Nahirnick et al. 2019; Murphy et al. 2021). Therefore, knowing the presence of shoreline modifications adjacent to seagrass meadows would allow us to predict ecological impacts and understand seagrass ecosystem dynamics in a broader seascape context.

Assessing human activities for an entire coastal region is generally done at broad spatial scales. For example, impact mapping and assessments for all of BC have been done with a 2 km+ spatial resolution (Clarke Murray et al. 2015), which exceeds the size of many seagrass meadows as well as the size of the shoreline region which may be locally impacting a meadow. In addition, many spatially distinct meadows may exist close together, where only a high resolution assessment of shoreline modifications could distinguish the potential impacts between them. Fine-scale assessments of impacts to seagrass exist for the BC coast, but these are typically done in detail for only a few meadows due to logistical constraints (Iacarella et al. 2018; Nagel et al. 2020).

The objective of this study is to map and quantify the shoreline modifications adjacent to all known seagrass meadows in the Strait of Georgia Bioregion of British Columbia. Eelgrass (*Zostera marina*, the dominant habitat-forming seagrass species) is a conservation priority in British Columbia (DFO 2019), and eelgrass meadows have been designated as Ecologically and Biologically Significant Areas (EBSA) due to their productivity, sensitivity, and support for biological diversity (Rubidge et al. 2020). Therefore, it is important to acquire information on human activities to predict impacts and categorize meadows by their degree of naturalness, as areas of high naturalness may be a priority for additional management and conservation efforts (UN CBD 2008). While shoreline modifications do not represent all of the human activities potentially threatening seagrass, a high resolution dataset is currently needed and can complement other existing human impact data.

2 Methods

2.1 Seagrass spatial data

Eelgrass (*Z. marina*) is the primary subtidal and intertidal meadow-forming seagrass in British Columbia. Meadows may also consist of the non-native seagrass, *Zostera japonica*, in the intertidal zone. Seagrass occurs to depths of 10 meters and can form meadows many km² in size (Murphy et al. 2021). We used a spatial dataset of seagrass in the Salish Sea compiled in Cristiani et al. (2021), which consists of surveyed and modeled data from a variety of government and non-governmental sources. Due to the combination of data with varying

collection methodologies and error, there is high uncertainty in the dataset, and it is best used for coarse estimates of extent. The dataset includes 685 spatially distinct meadows across the Strait of Georgia Bioregion as well as in the southern portions of the Northern Shelf Bioregion and Southern Shelf Bioregion (Figure 1).




Figure 1. Study area. Finish caption on seagrass and bioregions.

2.2 Shoreline area adjacent to seagrass meadows

Past guidelines on the width of marine riparian buffer zones for protecting sensitive habitat typically range from 50-150 meters (Levings and Jamieson 2001; Lemieux et al. 2004). We followed the methodology in a similar study of anthropogenic impacts (Iacarella et al. 2018) and examined shoreline modifications within 100 m of the coastline adjacent to each seagrass meadow. Quantifying modifications within a buffer zone requires generating consistent buffers from all meadows onto land. The perimeters of meadows, however, do not always exactly border

the shoreline due to variable mapping accuracies and errors, as well as some meadows only existing in the subtidal zone. This results in slightly different buffer extents on to land. Aside from a few exceptions, the majority of meadows are in close proximity to a coastline, and therefore, to create consistent width buffers on land we first adjusted the perimeter of meadows to match the coastline using digitization tools in ArcGIS.

2.3 Shoreline modifications

To quantify shoreline modification adjacent to seagrass, we identified any structures (e.g. buildings, houses) and areas de-vegetated from their natural state (e.g. lawns, logged areas, agriculture, armoured shoreline) within the 100 meter buffer. De-vegetated areas can increase nutrient run-off from agricultural areas and sewage outfalls potentially resulting in eutrophication (Hauxwell et al. 2003; Vandermeulen 2005; Quiros et al. 2017). Hardened and de-vegated areas may also increase the outflow of sedimentation resulting in decreased light levels in seagrass meadows (Vandermeulen et al. 2012; Dethier et al. 2016; Todd et al. 2019).

Shoreline modifications were digitized from satellite and aerial imagery in Google Earth Pro. The most recent imagery available was used, but the dates of the imagery varied across the study area. The majority of roads were included using an existing provincial dataset (Province of British Columbia - Digital Road Atlas). This dataset consists of linear features which were buffered using the number of lanes and standard lane widths. Modifications were classified into six categories: road, residential, cropland, industrial, greenspace, unclear. Within each category, an additional descriptor was also listed (Table 1). Overwater structures (e.g. floathomes, docks, aquaculture) were not considered to be shoreline modifications as these are associated with a different suite of impacts (e.g. shading, boat traffic), and these features were captured in a separate study.

To quantify the overall level of shoreline modification we totaled the area of all modification types per buffered area and calculated the percent of the buffered area that is modified. The percent modified was then associated back with the adjacent seagrass meadow.

Table 1. Shoreline modification classifications and descriptors

Modification	Subcategory descriptors
Road	paved; dirt; rail
Residential	house; RV; lighthouse
Industrial	logging; airport; general development; parking; storage; marina; church; electrical; ferry; hospital; train yard; shipping
Cropland	crop; fishfarm
Greenspace	cleared; campground; golf; park; recreation
Unclear	unclear

3 Results

128 The spatial distribution of shoreline modifications varied across the region. Examples of the
129 extent and diversity of modifications are shown in Figure 2, and the full dataset is accessible
130 online (see Data availability section). As expected, the seagrass meadows with the highest
131 total levels of adjacent shoreline modification occurred near population centers (Figure 3). The
132 lowest levels of modification occurred near islands and in the northern part of the study area
133 (i.e. Johnstone Strait). The dominant modification type tended to vary by latitude. For example,
134 there is more coastal agriculture in the south and more logging in the north (Figure).

135 (*TODO: make colors more bold in example figure.*) (*TODO: add cities and towns to overall figure*)
136 (*TODO: do a full page figure for the 5 types. If these come after the total one, then I could try to*
137 *fit them on one page.*)



Figure 2. Shoreline modifications within 100 meter buffered areas. The six selected areas are shown for example and do not imply any significance over other areas.




Figure 3. The percentage of the shoreline buffered area that is modified.

4 Discussion

By quantifying shoreline modifications adjacent to seagrass meadows, we've provided a novel high resolution dataset for documenting land use over a large spatial extent and predicting potential impacts to nearshore ecosystems. The variation in the distribution and patterns of modifications across the region illustrate the different threats that seagrass meadows face based on their location. If an area including a meadow is selected for management (e.g. an EBSA), then it's likely the shoreline modifications will have to be considered for possible mitigation of impacts. Knowing the spatial distribution of modifications is important because the management action will vary with the type of modification. While areas of naturalness are generally targeted for conservation over degraded areas (UN CBD 2008; Rubidge et al. 2020), other criteria such as biological diversity, productivity and connectivity may result in targeting meadows that are threatened by shoreline activities (Rubidge et al. 2020; Cristiani 2022).

This analysis could be strengthened by addressing some of its assumptions and limitations. We used a uniform buffer from the coast, but there is likely a distance-decay of the impacts from certain activities, such that subtidal meadows may experience impacts differently than intertidal meadows. In addition, a narrow buffer of vegetation on the edge of the coast could be enough to mitigate the effects of a certain modifications behind this buffer. Most importantly, it will be necessary to quantify the relative severity of each modification and the vulnerability of seagrass to different pressures. For example, runoff from agricultural areas may be more damaging than runoff from residential areas (Teck et al. 2010). Shoreline modifications are representative of only one type of threat to seagrass. However, by quantifying severity and vulnerability scores, this data could be incorporated into larger cumulative effects analyses in which the overall impact of many stressors is quantified (Clarke Murray et al. 2016, 2020; Murphy et al. 2022).

Our analysis emphasizes the importance of managing seagrass and the biodiversity it supports in the spatial context of the larger seascape and landscape. While eelgrass area is declining globally, it appears to be stable in the north Pacific (Dunic et al. 2021), and in the Washington state portion of the Salish Sea, eelgrass has been resilient despite an increase in human activities (Shelton et al. 2016). Understanding exactly which activities may be impacting eelgrass locally will likely require a more precise analysis of human activities, the stressors they generate, and the vulnerability of eelgrass to these stressors. Therefore, it will be important to continue to gather the high resolution spatial data that informs the initial stage of these analyses.

5 Data availability

Link to code in github repo
Link to dataset wherever it gets hosted

6 References

- Clarke Murray, C., Agbayani, S., Alidina, H.M., and Ban, N.C. 2015. [Advancing marine cumulative effects mapping: An update in Canada's Pacific waters](#). Marine Policy 58: 71–77.
- Clarke Murray, C., Hannah, L., and Locke, H. 2020. A Review of Cumulative Effects Research and Assessment in Fisheries and A Review of Cumulative Effects Research and Assessment in Fisheries and Oceans Canada. Can. Tech. Rep. Fish. Aquat. Sci. 3357: vii + 51 p.
- Clarke Murray, C., Mach, M.E., and O, M. 2016. Pilot ecosystem risk assessment to assess cumulative risk to species in the Pacific North Coast Integrated Management Area (PNCIMA). Canadian Science Advisory Secretariat.
- Cristiani, J. 2022. [The connectivity of coastal marine populations : Uncovering patterns of dispersal and the implications for management](#). University of British Columbia.
- Cristiani, J., Rubidge, E., Forbes, C., Moore-Maley, B., and O'Connor, M.I. 2021. [A Biophysical Model and Network Analysis of Invertebrate Community Dispersal Reveals Regional Patterns of Seagrass Habitat Connectivity](#). Frontiers in Marine Science 8: 1–19.
- Dethier, M.N., Raymond, W.W., McBride, A.N., Toft, J.D., Cordell, J.R., Ogston, A.S., Heerhartz, S.M., and Berry, H.D. 2016. [Multiscale impacts of armoring on Salish Sea shorelines: Evidence for cumulative and threshold effects](#). Estuarine, Coastal and Shelf Science 175: 106–117. Elsevier Ltd.
- DFO. 2019. Design strategies for the Northern Shelf Bioregion Marine Protected Area Network. CSAS.
- Dunic, J.C., Brown, C.J., Connolly, R.M., Turschwell, M.P., and Côt'e, I.M. 2021. [Long-term declines and recovery of meadow area across the world's seagrass bioregions](#). Global Change Biology: gcb.15684.
- Halpern, B.S., Frazier, M., Afflerbach, J., Lowndes, J.S., Micheli, F., O'NAHara, C., Scarborough, C., and Selko, K.A. 2019. [Recent pace of change in human impact on the world's ocean](#). Scientific Reports 9(1): 1–8.
- Hauxwell, J., Cebri'an, J., and Valiela, I. 2003. [Eelgrass *Zostera marina* loss in temperate estuaries: Relationship to land-derived nitrogen loads and effect of light limitation imposed by algae](#). Mar. Ecol. Prog. Ser. 247: 59–73.
- Iacarella, J.C., Adamczyk, E., Bowen, D., Chalifour, L., Eger, A., Heath, W., Helms, S., Hessing-Lewis, M., Hunt, B.P.V., MacInnis, A., O'Connor, M.I., Robinson, C.L.K., Yakimishyn, J., and Baum, J.K. 2018. [Anthropogenic disturbance homogenizes seagrass fish communities](#). Global Change Biology 24(5): 1904–1918.
- Lemieux, J.P., Brennan, J.S., Farrell, M., Levings, C.D., and Myers, D. 2004. PROCEEDINGS OF THE DFO/PSAT SPONSORED MARINE RIPARIAN EXPERTS WORKSHOP, TSAWWASSEN, BC, FEBRUARY 17-18, 2004.

- Levings, C., and Jamieson, G. 2001. Marine and estuarine riparian habitats and their role in coastal ecosystems, Pacific region. Canadian Science Advisory Secretariat: 41.
- Murphy, G.E.P., Dunic, J.C., Adamczyk, E.M., Bittick, S.J., Côté, I.M., Cristiani, J., Geissinger, E.A., Gregory, R.S., Lotze, H.K., O'Connor, M.I., Ara'ujo, C.A.S., Rubidge, E.M., Templeman, N.D., and Wong, M.C. 2021. [From coast to coast to coast : Ecology and management of seagrass ecosystems across Canada](#). Facets 6: 1–41.
- Murphy, G.E.P., Kelly, N.E., Lotze, H.K., and Wong, M.C. 2022. [Incorporating anthropogenic thresholds to improve understanding of cumulative effects on seagrass beds](#). FACETS 7: 966–987.
- Nagel, E.J., Murphy, G.E.P., Fast, J., Bittick, S.J., Adamczyk, M., Connor, M.I.O., Wong, M.C., and Lotze, H.K. 2020. Application of a coastal human impact metric and nitrogen loading model to 10 eelgrass (*Zostera marina*) meadows in British Columbia. Fisheries and Oceans Canada.
- Nahirnick, N.K., Costa, M., Schroeder, S., and Sharma, T. 2019. [Long-Term Eelgrass Habitat Change and Associated Human Impacts on the West Coast of Canada](#). Journal of Coastal Research 36(1): 30.
- Quiros, T.E.A.L., Croll, D., Tershy, B., Fortes, M.D., and Raimondi, P. 2017. [Land use is a better predictor of tropical seagrass condition than marine protection](#). Biological Conservation 209: 454–463. The Authors.
- Rubidge, E., Jeffery, S., Gregr, E.J., Gale, K.S.P., and Frid, A. 2020. Assessment of nearshore features in the Northern Shelf Bioregion against criteria for determining Ecologically and Biologically Significant Areas (EBSAs). DFO Canadian Science Advisory Secretariat. Canadian Science Advisory Secretariat.
- Shelton, A.O., Francis, T.B., Feist, B.E., Williams, G.D., Lindquist, A., and Levin, P. 2016. [Forty years of seagrass population stability and resilience in an urbanizing estuary](#). Journal of Ecology: 1–13.
- Teck, S., Halpern, B., and Kappel, C. 2010. [Using expert judgement to estimate marine ecosystem vulnerability in the California Current](#). Ecological ... 20(5): 1–16.
- Todd, P.A., Heery, E.C., Loke, L.H.L., Thurstan, R.H., Kotze, D.J., and Swan, C. 2019. [Towards an urban marine ecology: Characterizing the drivers, patterns and processes of marine ecosystems in coastal cities](#). Oikos: 1215–1242.
- UN CBD. 2008. Decision adopted by the conference of the parties to the convention on biological diversity at its ninth meeting. UNEP/CBD/COP/DEC/IX/20.
- Vandermeulen, H. 2005. Assessing Marine Habitat Sensitivity: A case study with eelgrass (*Zostera marina* L.) And kelps (*Laminaria*, *Macrocystis*). DFO Canadian Science Advisory Secretariat, Fisheries and Oceans Canada.
- Vandermeulen, H., Surette, J., and Skinner, M.A. 2012. Responses of Eelgrass (*Zostera Marina* L.) To stress. DFO Can. Sci. Advis. Sec. Res. Doc. 2011/095. vi + 43 p.