

# **Quantifying shoreline modifications adjacent to eelgrass meadows in the Strait of Georgia Bioregion**

John Cristiani, Katherine Bannar-Martin, Emily M. Rubidge

Fisheries and Oceans Canada  
Science Branch, Pacific Region  
Pacific Biological Station  
Nanaimo, British Columbia  
V9T 6N7

2022

**Canadian Technical Report of  
Fisheries and Aquatic Sciences #####**



Fisheries and Oceans  
Canada

Pêches et Océans  
Canada

**Canada** 

## **Canadian Technical Report of Fisheries and Aquatic Sciences**

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

## **Rapport technique canadien des sciences halieutiques et aquatiques**

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques de Pêches et Océans Canada, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications à part entière. Le titre exact figure au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre.

Les numéros 1 à 456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Canadian Technical Report of  
Fisheries and Aquatic Sciences nnn

2022

QUANTIFYING SHORELINE MODIFICATIONS ADJACENT TO EELGRASS MEADOWS IN THE  
STRAIT OF GEORGIA BIOREGION

by

John M. Cristiani<sup>1</sup> Katherine H. Bannar-Martin<sup>2</sup> and Emily M. Rubidge<sup>2</sup>

<sup>1</sup>Pacific Biological Station  
Fisheries and Oceans Canada, 3190 Hammond Bay Road  
Nanaimo, British Columbia, V9T 6N7, Canada  
<sup>2</sup>Institute of Ocean Sciences  
Fisheries and Oceans Canada, 9860 W Saanich Road  
Sidney, British Columbia, V8L 4B2, Canada

16  
17

© His Majesty the King in Right of Canada, 2022  
Cat. No. Fs97-6/nnnE-PDF ISBN ISSN 1488-5379

18 Correct citation for this publication:

19 Cristiani, J.C., Bannar-Martin, K.H, and Rubidge, E.M. 2022. Quantifying shoreline modifications  
20 adjacent to eelgrass meadows in the Strait of Georgia Bioregion. Can. Tech. Rep. Fish. Aquat.  
21 Sci. nnn: v + 9p.

23	<b>ABSTRACT</b>	<b>iv</b>
24	<b>RÉSUMÉ</b>	<b>v</b>
25	<b>1 Introduction</b>	<b>1</b>
26	<b>2 Methods</b>	<b>1</b>
27	2.1 Seagrass spatial data . . . . .	1
28	2.2 Shoreline area adjacent to seagrass meadows . . . . .	2
29	2.3 Shoreline modifications . . . . .	2
30	<b>3 Results</b>	<b>4</b>
31	<b>4 Discussion</b>	<b>7</b>
32	<b>5 Data availability</b>	<b>7</b>
33	<b>6 References</b>	<b>8</b>

**ABSTRACT**

35 Cristiani, J.C., Bannar-Martin, K.H, and Rubidge, E.M. 2022. Quantifying shoreline modifications  
36 adjacent to eelgrass meadows in the Strait of Georgia Bioregion. Can. Tech. Rep. Fish. Aquat.  
37 Sci. nnn: v + 9 p.

38 Here is the abstract text. Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do  
39 eiusmod tempor incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis  
40 nostrud exercitation ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure  
41 dolor in reprehenderit in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint  
42 occaecat cupidatat non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

## RÉSUMÉ

44 Cristiani, J.C., Bannar-Martin, K.H, and Rubidge, E.M. 2022. Quantifying shoreline modifications  
45 adjacent to eelgrass meadows in the Strait of Georgia Bioregion. Can. Tech. Rep. Fish. Aquat.  
46 Sci. nnn: v + 9 p.

47 Voici le résumé. Lorem ipsum dolor sit amet, consectetur adipisicing elit, sed do eiusmod tempor  
48 incididunt ut labore et dolore magna aliqua. Ut enim ad minim veniam, quis nostrud exercitation  
49 ullamco laboris nisi ut aliquip ex ea commodo consequat. Duis aute irure dolor in reprehenderit  
50 in voluptate velit esse cillum dolore eu fugiat nulla pariatur. Excepteur sint occaecat cupidatat  
51 non proident, sunt in culpa qui officia deserunt mollit anim id est laborum.

## 1 Introduction

The health and functioning of coastal marine ecosystems are under threat from a variety of human activities (Halpern et al. 2019). Coastal activities such as agriculture, industrial and residential development, forestry, and shoreline hardening can create pressures on the marine environment. A modified shoreline may alter levels of sedimentation, nutrient runoff, pollution, and wave energy (Dethier et al. 2016; Todd et al. 2019). For coastal biogenic habitat in British Columbia such as seagrass, these pressures may impact seagrass productivity and survival, and thus impact the community of species that rely on seagrass (Iacarella et al. 2018; Nahirnick et al. 2019; Murphy et al. 2021). Therefore, knowing the presence of shoreline modifications adjacent to seagrass meadows would allow us to predict ecological impacts and understand seagrass ecosystem dynamics in a broader seascape context.

Assessing human activities for an entire coastal region is generally done at broad spatial scales. For example, impact mapping and assessments for all of BC have been done with a 2 km+ spatial resolution (Clarke Murray et al. 2015), which exceeds the size of many seagrass meadows as well as the size of the shoreline region which may be locally impacting a meadow. In addition, many spatially distinct meadows may exist close together, where only a high resolution assessment of shoreline modifications could distinguish the potential impacts between them. Fine-scale assessments of impacts to seagrass exist for the BC coast, but these are typically done in detail for only a few meadows due to logistical constraints (Iacarella et al. 2018; Nagel et al. 2020).

The objective of this study is to map and quantify the shoreline modifications adjacent to all known seagrass meadows in the Strait of Georgia Bioregion of British Columbia. Eelgrass (*Zostera marina*, the dominant habitat-forming seagrass species) is a conservation priority in British Columbia (DFO 2019), and eelgrass meadows have been designated as Ecologically and Biologically Significant Areas (EBSA) due to their productivity, sensitivity, and support for biological diversity (Rubidge et al. 2020). Therefore, it is important to acquire information on human activities to predict impacts and categorize meadows by their degree of naturalness, as areas of high naturalness may be a priority for additional management and conservation efforts (UN CBD 2008). While shoreline modifications do not represent all of the human activities potentially threatening seagrass, a high resolution dataset is currently needed and can complement other existing human impact data.

## 2 Methods

### 2.1 Seagrass spatial data

Eelgrass (*Z. marina*) is the primary subtidal and intertidal meadow-forming seagrass in British Columbia. Meadows may also consist of the non-native seagrass, *Zostera japonica*, in the intertidal zone. Seagrass occurs to depths of 10 meters and can form meadows many km<sup>2</sup> in size (Murphy et al. 2021). We used a spatial dataset of seagrass in the Salish Sea compiled in Cristiani et al. (2021), which consists of surveyed and modeled data from a variety of government and non-governmental sources. Due to the combination of data with varying



collection methodologies and error, there is high uncertainty in the dataset, and it is best used for coarse estimates of extent. The dataset includes 685 spatially distinct meadows across the Strait of Georgia Bioregion as well as in the southern portions of the Northern Shelf Bioregion and Southern Shelf Bioregion (Figure @ref:studyareafig).

## 2.2 Shoreline area adjacent to seagrass meadows

Past guidelines on the width of marine riparian buffer zones for protecting sensitive habitat typically range from 50-150 meters (Levings and Jamieson 2001; Lemieux et al. 2004). We followed the methodology in a similar study of anthropogenic impacts (Iacarella et al. 2018) and examined shoreline modifications within 100 m of the coastline adjacent to each seagrass meadow. Quantifying modifications within a buffer zone requires generating consistent buffers from all meadows onto land. The perimeters of meadows, however, do not always exactly border the shoreline due to variable mapping accuracies and errors, as well as some meadows only existing in the subtidal zone. This results in slightly different buffer extents on to land. Aside from a few exceptions, the majority of meadows are in close proximity to a coastline, and therefore, to create consistent width buffers on land we first adjusted the perimeter of meadows to match the coastline using digitization tools in ArcGIS.

## 2.3 Shoreline modifications

To quantify shoreline modification adjacent to seagrass, we identified any structures (e.g. buildings, houses) and areas de-vegetated from their natural state (e.g. lawns, logged areas, agriculture, armoured shoreline) within the 100 meter buffer. De-vegetated areas can increase nutrient run-off from agricultural areas and sewage outfalls potentially resulting in eutrophication (Hauxwell et al. 2003; Vandermeulen 2005; Quiros et al. 2017). Hardened and de-vegetated areas may also increase the outflow of sedimentation resulting in decreased light levels in seagrass meadows (Vandermeulen et al. 2012; Dethier et al. 2016; Todd et al. 2019).

Shoreline modifications were digitized from satellite and aerial imagery in Google Earth Pro. The most recent imagery available was used, but the dates of the imagery varied across the study area. The majority of roads were included using an existing provincial dataset (Province of British Columbia - Digital Road Atlas). This dataset consists of linear features which were buffered using the number of lanes and standard lane widths. Modifications were classified into six categories: road, residential, cropland, industrial, greenspace, unclear. Within each category, an additional descriptor was also listed (Table 1). Overwater structures (e.g. floathomes, docks, aquaculture) were not considered to be shoreline modifications as these are associated with a different suite of impacts (e.g. shading, boat traffic), and these features were captured in a separate study.

To quantify the overall level of shoreline modification we totaled the area of all modification types per buffered area and calculated the percent of the buffered area that is modified. The percent modified was then associated back with the adjacent seagrass meadow.

Table 1. Shoreline modification classifications and descriptors

Modification	Subcategory descriptors
Road	paved; dirt; rail
Residential	house; RV; lighthouse
Industrial	logging; airport; general development; parking; storage; marina; church; electrical; ferry; hospital; train yard; shipping
Cropland	crop; fishfarm
Greenspace	cleared; campground; golf; park; recreation
Unclear	unclear




Figure 1. Study area

127

### 3 Results

128

129

- A few sentences on the spatial distribution (e.g. more modifications in the south, more ag in south, more forestry in north).



Figure 2. Shoreline modifications within 100 meter buffered areas. The six selected areas are shown for example and do not imply any significance over other areas.






Figure 3. The percentage of the shoreline buffer modified.

## 4 Discussion

- summary paragraph
- spatial distribution of activities and the significance for management
- assumptions and limitations
  - No point measurements in field to confirm impacts. Some activities are well regulated in Canada, like logging, and there may be very minimal impact.
  - Meadows that aren't right on the coast may not experience any impacts
  - If the modification exists behind some vegetation then the impact may be less (cite lacarella)
  - We use a uniform buffer, but there is a likely a distance decay for some of these activities.
- Future directions
  - incorporate vulnerability scores to specific types of modifications
  - Consider management goals of protecting the most natural meadows and how this might change given the spatial distribution of activities and other information (e.g. biodiversity in meadows, connectivity)
- some stuff from my thesis:
  - “managing seagrass habitat and associated species in a landscape context, in which patterns of distribution, dispersal, and impacts will interact to influence regional management strategies (Murphy et al. 2021b). Although eelgrass is declining globally (Dunic et al. 2021), eelgrass in nearby Puget Sound, Washington is stable and resilient overall, despite a significant increase in local human and climactic stressors (Shelton et al. 2016). Assessing the relevance of managing for human impacts in the Salish Sea will therefore require a deeper understanding of seagrass and invertebrate responses to stressors and the mechanisms (e.g., dispersal) that allow for resilience to these stressors. Ultimately, refining and validating our models will increase their utility and promote their incorporation into broader marine spatial planning efforts.”

## 5 Data availability

Link to github repo

## 6 References

- Clarke Murray, C., Agbayani, S., Alidina, H.M., and Ban, N.C. 2015. [Advancing marine cumulative effects mapping: An update in Canada's Pacific waters](#). Marine Policy 58: 71–77.
- Cristiani, J., Rubidge, E., Forbes, C., Moore-Maley, B., and O'Connor, M.I. 2021. [A Biophysical Model and Network Analysis of Invertebrate Community Dispersal Reveals Regional Patterns of Seagrass Habitat Connectivity](#). Frontiers in Marine Science 8: 1–19.
- Dethier, M.N., Raymond, W.W., McBride, A.N., Toft, J.D., Cordell, J.R., Ogston, A.S., Heerhartz, S.M., and Berry, H.D. 2016. [Multiscale impacts of armoring on Salish Sea shorelines: Evidence for cumulative and threshold effects](#). Estuarine, Coastal and Shelf Science 175: 106–117. Elsevier Ltd.
- DFO. 2019. Design strategies for the Northern Shelf Bioregion Marine Protected Area Network. CSAS.
- Halpern, B.S., Frazier, M., Afflerbach, J., Lowndes, J.S., Micheli, F., O'NAHara, C., Scarborough, C., and Selko, K.A. 2019. [Recent pace of change in human impact on the world's ocean](#). Scientific Reports 9(1): 1–8.
- Hauxwell, J., Cebrian, J., and Valiela, I. 2003. [Eelgrass \*Zostera marina\* loss in temperate estuaries: Relationship to land-derived nitrogen loads and effect of light limitation imposed by algae](#). Mar. Ecol. Prog. Ser. 247: 59–73.
- Iacarella, J.C., Adamczyk, E., Bowen, D., Chalifour, L., Eger, A., Heath, W., Helms, S., Hessing-Lewis, M., Hunt, B.P.V., MacInnis, A., O'Connor, M.I., Robinson, C.L.K., Yakimishyn, J., and Baum, J.K. 2018. [Anthropogenic disturbance homogenizes seagrass fish communities](#). Global Change Biology 24(5): 1904–1918.
- Lemieux, J.P., Brennan, J.S., Farrell, M., Levings, C.D., and Myers, D. 2004. PROCEEDINGS OF THE DFO/PSAT SPONSORED MARINE RIPARIAN EXPERTS WORKSHOP, TSAWWASSEN, BC, FEBRUARY 17-18, 2004.
- Levings, C., and Jamieson, G. 2001. Marine and estuarine riparian habitats and their role in coastal ecosystems, Pacific region. Canadian Science Advisory Secretariat: 41.
- Murphy, G.E.P., Dunic, J.C., Adamczyk, E.M., Bittick, S.J., Côté, I.M., Cristiani, J., Geissinger, E.A., Gregory, R.S., Lotze, H.K., O'Connor, M.I., Ara'ujo, C.A.S., Rubidge, E.M., Templeman, N.D., and Wong, M.C. 2021. [From coast to coast to coast : Ecology and management of seagrass ecosystems across Canada](#). Facets 6: 1–41.
- Nagel, E.J., Murphy, G.E.P., Fast, J., Bittick, S.J., Adamczyk, M., Connor, M.I.O., Wong, M.C., and Lotze, H.K. 2020. Application of a coastal human impact metric and nitrogen loading model to 10 eelgrass ( *Zostera marina* ) meadows in British Columbia. Fisheries and Oceans Canada.
- Nahirnick, N.K., Costa, M., Schroeder, S., and Sharma, T. 2019. [Long-Term Eelgrass Habitat Change and Associated Human Impacts on the West Coast](#)

199 [of Canada](#). Journal of Coastal Research 36(1): 30.

200 Quiros, T.E.A.L., Croll, D., Tershy, B., Fortes, M.D., and Raimondi, P. 2017. [Land use](#)  
201 [is a better predictor of tropical seagrass condition than marine protection](#). Biological  
202 Conservation 209: 454–463. The Authors.

203 Rubidge, E., Jeffery, S., Gregr, E.J., Gale, K.S.P., and Frid, A. 2020. Assessment of nearshore  
204 features in the Northern Shelf Bioregion against criteria for determining Ecologically and  
205 Biologically Significant Areas ( EBSAs ). DFO Canadian Science Advisory Secretariat.  
206 Canadian Science Advisory Secretariat.

207 Todd, P.A., Heery, E.C., Loke, L.H.L., Thurstan, R.H., Kotze, D.J., and Swan, C. 2019.  
208 [Towards an urban marine ecology: Characterizing the drivers, patterns and processes of](#)  
209 [marine ecosystems in coastal cities](#). Oikos: 1215–1242.

210 UN CBD. 2008. Decision adopted by the conference of the parties to the convention on  
211 biological diversity at its ninth meeting. UNEP/CBD/COP/DEC/IX/20.

212 Vandermeulen, H. 2005. Assessing Marine Habitat Sensitivity: A case study with eelgrass  
213 (*Zostera marina* L.) And kelps (*Laminaria*, *Macrocystis*). DFO Canadian Science Advisory  
214 Secretariat, Fisheries and Oceans Canada.

215 Vandermeulen, H., Surette, J., and Skinner, M.A. 2012. Responses of Eelgrass (*Zostera*  
216 *Marina* L.) To stress. DFO Can. Sci. Advis. Sec. Res. Doc. 2011/095. vi + 43 p.